



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Cohort Profile: The Alberta Pregnancy Outcomes and Nutrition Cohort Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-047503
Article Type:	Cohort profile
Date Submitted by the Author:	11-Dec-2020
Complete List of Authors:	<p>Letourneau, Nicole; University of Calgary, Faculty of Nursing; University of Calgary Cumming School of Medicine, Pediatrics, Psychiatry and Community Health Sciences</p> <p>Ali, Elena; University of Calgary</p> <p>Aghajafari, Fariba; University of Calgary, Family Medicine</p> <p>Bell, R; University of Alberta, Agricultural, Food and Nutritional Science; University of Alberta</p> <p>Deane, Andrea</p> <p>Dewey, Deborah ; University of Calgary, Paediatrics and Community Health Sciences</p> <p>Field, Catherine; University of Alberta, Department of Agricultural, Food and Nutritional Science;</p> <p>Giesbrecht, Gerald ; University of Calgary,</p> <p>Kaplan, Bonnie; University of Calgary Cumming School of Medicine, ;</p> <p>Leung, Brenda; University of Calgary,</p> <p>Ntanda, Henry</p>
Keywords:	NUTRITION & DIETETICS, MENTAL HEALTH, PAEDIATRICS, PUBLIC HEALTH

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Cohort Profile: The Alberta Pregnancy Outcomes and Nutrition Cohort Study

Authors: Nicole Letourneau^{1,2}, Elena Ali¹, Fariba Aghajafari³, Rhonda Bell⁴, Andrea Deane⁵, Deborah Dewey⁶, Catherine Field⁴, Gerald Giesbrecht⁷, Bonnie Kaplan⁷, Brenda Leung⁸, Henry Ntanda⁵, & APrON Study Team

Author affiliations: ¹ Faculty of Nursing, University of Calgary, Calgary, AB, Canada;

² Cumming School of Medicine (Paediatrics, Psychiatry & Community Health Sciences),

University of Calgary, Calgary, AB, Canada; ³ Cumming School of Medicine (Family Medicine

& Community Health Sciences), University of Calgary, Calgary, AB, Canada; ⁴ Faculty of

Agricultural, Life & Environmental Sciences, University of Alberta, Edmonton, AB, Canada;

⁵ Cumming School of Medicine, University of Calgary, Calgary, AB, Canada; ⁶ Cumming School

of Medicine (Paediatrics), University of Calgary, Calgary, AB, Canada; ⁷ Cumming School of

Medicine (Paediatrics & Community Health Sciences), University of Calgary, Calgary, AB,

Canada; ⁸ University of Lethbridge (Health Science – Public Health Program), University of

Lethbridge, Lethbridge, AB, Canada

Corresponding author:

Nicole Letourneau

Faculty of Nursing

Cumming School of Medicine (Paediatrics, Psychiatry & Community Health Sciences)

University of Calgary

2500 University Drive NW

Calgary, AB

T2N 1N4

Telephone: (403) 210-3833

E-mail: nicole.letourneau@ucalgary.ca

Cohort Profile: The Alberta Pregnancy Outcomes and Nutrition Cohort Study

ABSTRACT

Purpose: The purposes of the Alberta Pregnancy Outcomes and Nutrition (APrON) study were: to determine the relationship between maternal nutrient intake and status before, during, and after pregnancy, and (a) maternal mental health, (b) pregnancy and birth outcomes, and (c) infant neurodevelopment; to examine the relationship between maternal thyroid status and neonatal outcomes, and to establish nutrient predictors of infant behavioral patterns; and to establish a DNA biobank to examine the influence of nutritional genomics on children neurodevelopment and behavioral patterns.

Participants: APrON is an ongoing prospective transgenerational cohort study consisting of participants who reside in Calgary and Edmonton, Alberta, Canada. Data comprise of: self-completed mother, father, and child questionnaires; maternal and paternal dietary interviews; clinical assessments; linkage to hospital obstetrical records; and biological samples. DNA samples were banked for future studies of nutrient/gene and genetic links to infant development.

Findings to date: Key findings pertaining to mental health and nutrition outcomes are presented. Specifically, risk factors for depressive symptoms and the positive association between maternal exposure to adverse childhood experiences and prenatal depression and anxiety are discussed. The consumption of specific food groups that is associated with meeting choline adequate intake recommendations is discussed, along with findings pertaining to mothers' gestational weight gain and reported dietary Vitamin D intake.

Future plans: The APrON cohort offers a unique opportunity to follow a group of Canadian children to advance understanding of the fetal origins of diseases. The APrON study aspires to transition to an intergenerational study in order to investigate current child participants in

adulthood. Accordingly, the APrON study will use a detailed array of standardized validated psychological and physiological measures and biospecimens to collect detailed information on maternal nutrition during pregnancy and maternal and paternal perinatal mental health.

ARTICLE SUMMARY

- The APrON study used a detailed array of standardized validated psychological and physiological measures and biospecimens. The prospective data collection and follow-up enabled investigation of a wide range of maternal and paternal health outcomes.
- The APrON team has been successful in continuously engaging participants, as evidenced by high response rates during pregnancy, postpartum and early childhood.
- One limitation is that the sample characteristics are skewed toward a high percentage of white, well-educated and married participants. Consequently, generalizability of findings to those with different socio-demographic backgrounds should be made with caution.
- Another limitation is that selective attrition bias has likely occurred, which may lead to underestimation of the effect of parental mental health and lifestyle choices on the child's development.

INTRODUCTION

The Alberta Pregnancy Outcomes and Nutrition (APrON) study is a longitudinal cohort of pregnant women, their partners¹, and children in Alberta, Canada. The APrON study was designed to investigate the effect of nutrient intake and status during pregnancy on maternal mental health and children's neurodevelopment [1, 2]. The original grant of \$5 million provided the foundation for the APrON study, which has expanded its focus and contributed to over 50

¹ Mothers in our study are biological mothers. Fathers self-identified as fathers of their children, whether biological or not.

additional projects on the psychobiology of stress, parent-child relationships and attachment, neurotoxicant exposure, genetics, epigenetics, inflammation, and immune activity, probiotics, the microbiome and children's brain development. To date, the APrON study has received over \$14 million in additional funding. This funding enabled follow-up of APrON families to when children reach age 12 to address the impact of medical, biological, and environmental factors on children's development, behavior and mental health [3]. Data collection for the five-year follow-up questionnaire is complete, eight-year follow-up is ongoing and a 12-year follow-up will start in 2021. The objective of this paper is to describe key findings on maternal and paternal mental health outcomes and maternal nutrient intake and status of APrON participants from pregnancy to three years after delivery.

COHORT DESCRIPTION

Women were recruited into the cohort if they were 16 years of age or older, able to speak and read English, < 27 weeks gestation and willing to come for on-site visits at the University of Calgary, Calgary or University of Alberta, Edmonton, Alberta. Men were eligible if they were cohabitating with the participating woman. Participants were excluded if they planned to move out of the region during the on-site visit timeframe (pregnancy to 3 months postpartum) of the study. A total of 2189 pregnant women (aged 16 to 44) and 1325 men (aged 18 to 52), residing around Calgary or Edmonton, were enrolled in the study between May 2009 and June 2012. The APrON study was approved by the University of Calgary Health Research Ethics Board (REB14-1702) and University of Alberta Health Research Ethics Biomedical Panel (00002954). Full details on how women and their partners were recruited are provided by Kaplan et al. (2014) and in previous publications describing the APrON cohort [4, 5]. To assess the representativeness of the APrON cohort (see Table 1), socio-demographic characteristics of participants at the time of

recruitment were compared to characteristics of women with children in Calgary, Alberta, and

Category	Maternal (n / %)	Paternal (n / %)
Demographics		
Age (years)		
16 – 24	100 (4.6)	19 (1.4)
25 – 29	481 (22.0)	216 (15.4)
30 -34	980 (44.78)	493 (35.2)
35 -39	512 (23.4)	329 (23.5)
≥ 40	105 (4.8)	162 (11.6)
Missing age information	11 (0.5)	181 (12.9)
Marital status		
Married/common-law	2035 (93.0)	1275 (91.1)
Single/divorced/separated	97 (4.4)	27 (1.9)
Missing marital information	57 (2.6)	98 (7.0)
Education		
Less than a high school diploma	59 (2.7)	31 (2.2)
Completed high school diploma	202 (9.2)	147 (10.5)
Completed trade, technical	404 (18.5)	376 (26.9)
Completed university or more	1442 (66.8)	746 (53.3)
Missing education information	83 (3.7)	100 (7.1)
Ethnicity		
White	1693 (77.3)	1082 (77.3)
Non-white	485 (22.2)	206 (14.7)
Missing ethnic information	11 (0.5)	112 (8.0)
Mothers' Report on total household income		
< \$40,000	190 (8.7)	-
\$40,000 - \$69,999	279 (12.7)	-
\$70,000 - \$99,999	464(21.2)	-
>\$100,000	1146 (52.3)	-
Missing household income information	110 (5.0)	-
Born in Canada		
Yes	1618 (73.9)	1024 (73.1)
No	481 (22.0)	277 (19.8)
Missing country of birth	90 (4.1)	99 (7.1)
Gravidity		
Multigravida	1107 (51.6)	n/a
Primigravida	996 (46.4)	n/a
Missing gravidity information	42 (2.0)	n/a
Parity		
0	1222 (55.8)	n/a
1	732 (33.4)	n/a

Canada [3, 5, 6].

2	181 (8.3)	n/a
≥3	37 (1.7)	n/a
Missing parity information	17 (0.8)	n/a

Table 1. *Participant Intake Characteristics (N = 3514)*

A greater proportion of women in the cohort (28.0%) were over 34 years old compared with women in Calgary (20.0%), Alberta (16.0%), and Canada (18.0%). More APrON women were married (95%) compared with women in Calgary (73.0%), Alberta (70.0%), and Canada (60.0%). A greater proportion of women had household income of \$70,000 or more (77.0%) compared to Calgary (65.0%), Alberta (61.0%), and Canada (58.0%). More APrON women completed post-secondary education (87.0%) compared to women in Calgary (69.0%), Alberta (70.0%), and Canada (72.0%). The proportion of APrON women who were born outside of Canada (23.0%) was similar to the proportions in Calgary (25.0%), Alberta (22%), and Canada (20.0%) (Figure 1).

How Often Have They Been Followed Up?

Between 2009 and 2012 the initial cohort completed questionnaires and clinical assessments three times during pregnancy, and at 3 months postpartum. Follow-up questionnaires were provided when their child was 6 months, and 1-, 2- and 3- years of age (see Figure 2).

In total, 2189 women and 1325 men completed at least one questionnaire. Of these, 1945 (88.8%), are continuing participants, who were defined as those who did not become ineligible or lost to follow-up. Participants who discontinued the study included women who miscarried, had a stillbirth, lost custody of their child, or for whom a maternal/child death was reported ($n = 52$, 2.4%); moved out of Calgary or Edmonton before three months postpartum ($n = 43$, 2.0%); or withdrew from the study citing loss of interest ($n = 147$; 6.7%). The demographic characteristics

No	425 (22.5)	13 (32.5)	14(37.8)	28 (21.2)	0.186
Yes	1463 (77.5)	27 (67.5)	23 (62.2)	104 (78.8)	

Note. Assessing the null hypothesis that there is no difference in distributions between those participants who continued and those who did not continue in the study; chi-square test

Continuing participants were more likely older, married, and had higher education and household income (all $ps < 0.05$). The proportion of continuing participants born in Canada was similar to discontinuing participants. When mothers were lost to follow-up, so too were fathers and children.

What Has Been Measured?

Women were asked to complete questionnaires three times during pregnancy (<14, 14-26, and 27-40 weeks gestation, number of scales administered at this time was 37) and six times postnatally (3 and 6 months, 1, 2, 3, and 5 years, number of scales administered at this time was 65). Fathers were asked to complete questionnaires when the women were 14-26 weeks gestational age (number of scales administered at this time was 16), and twice postnatally (3 months and 5 years, number of scales administered at this time were 14). Questionnaires administered during pregnancy measured previous and current mental and physical health status, nutritional, medication and supplement intake, lifestyle choices, healthcare services used, social support, attitudes towards breastfeeding, and sociodemographic variables. The follow-up postpartum questionnaires collected information about labor and delivery; postpartum medical problems; maternal mental health; nutritional, medication and supplement intake; lifestyle choices; food security; and socio-demographics (see Figure 2). A detailed list of data collected, including measures used if applicable, is provided in Table 3.1 for mothers and Table 3.2 for fathers.

Table 3.1. *Maternal Variables and Biological Specimens Collected in All Questionnaires and Clinical Visits*

Variables	<14 weeks	14-26 weeks	27-40 weeks	3 months postpartum	6 months postpartum	1 year	2 years	3 years
Socio-demographics								
Household income	X							
Education	X							
Marital status	X							
Age	X							
Primary language	X							
Born in Canada	X							
Family background		X	X	X	X	X	X	X
Family history		X						
Health and lifestyle								
General health				SF-8	SF-8	SF-8	SF-8	SF-8
Current medical conditions and medications	X	X	X	X	X	X	X	X
Substance use	X	X	X	X	X	X	X	X
Pregnancy history								
Pre-pregnancy physical activity	X	X	X	X	X	X	X	X
Physical activity	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE
Dental care	X	X	X					
Dietary changes since pregnancy	X							
Food security			CCHSC		CCHSC			
Nutrition counselling information				X				
Diet history pre-pregnancy		FFQ						
Nausea and vomiting	PUQUE	PUQUE	PUQUE					
Breastfeeding information				X	X	X	X	X
24-hour recall of foods consumed	X	X	X	X	X	X	X	X
Psychosocial health								
Depression	EDS	EDS	EDS	EDS	EDS	EDS	EDS	EDS
Stress	SLEQ		SLEQ	SLEQ		SLEQ	SLEQ	SLEQ
Social support	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ

DSM-IV Axis I disorders	SCL-90-R	PDSQ	SCL-90-R	SCL-90-R	PDSQ	SCL-90-R	SCL-90-R	SCL-90-R	SCL-90-R
Anxiety	ACEs	R	R	R	R	R	R	R	R
Adverse childhood events	X	X	X	X	X	X	X	X	X
History of mental health	X	X	X	X	X	X	X	X	X
Anthropometrics (BMI/height/weight)									
Biological specimens									
Urine		X			X				
Venous blood	X	X	X	X	X	X	X	X	X
Breast milk					X				

Note. X refers to investigator-developed measures. SCL-90-R, Symptom Checklist-90-Revised; EDS, Edinburgh Depression Scale; PUQE, Pregnancy-Unique Quantification of Emesis and Nausea questionnaire; SLEQ, Stressful Life Events Questionnaire; SSQ, Social Support Questionnaire; PDSQ, Psychiatric Diagnostic Screening Questionnaire; ACEs, Adverse Childhood Experiences Questionnaire; SF-8, Short Form Health Survey-8; CCHSC, Canadian Community Health Survey Cycle 2.2, modified; GLTE, Godin Leisure-Time Exercise Questionnaire; FFQ, Food Frequency Questionnaire, adapted from the Canadian version of the Diet History Questionnaire.*Breastfeeding mothers only.

Table 3.2. *Paternal Variables and Biological Specimens Collected in All Questionnaires and Clinical Visits*

Variables	<14 weeks gestation	14-26 weeks	27-40 weeks	3 months postpartum
Socio-demographics				
Household income			X	
Education			X	
Marital status			X	
Age			X	
Primary language			X	
Born in Canada			X	
Family background			X	
Family history*			X	
Health and lifestyle		X	X	
General health				
Current medical conditions and medications		X	X	X
Substance use		X	X	X
Physical Activity Anthropometrics (BMI/height/weight)		GLTE X	GLTE X	GLTE X
Psychosocial health				
Depression		EDS	EDS	EDS
Social support		SSQ	SSQ	SSQ
Stress			SLEQ	
Biological specimens*				
Buccal cell swab			X	
DNA from blood			X	

Maternal anthropometric measurements were collected at each visit during pregnancy and three months postpartum. These measures included height, pre-pregnancy and current weight, highest weight during pregnancy; circumference measurements (i.e., mid-upper arm, waist, hip, thigh); and skinfold thickness (i.e., biceps, triceps, subscapular, suprailiac, thigh). Each measurement was taken three times and averages were reported for analyses. Paternal height and weight were collected during the prenatal period. One objective of APrON was to establish a biobank for future research that would allow investigation on genetics, epigenetics [1], nutrient exposures and neurotoxicants. Maternal non-fasting venous blood samples were collected from

mothers at <14, 14-26, and 27-40 weeks gestation, and at three months postpartum. Maternal blood collection provided whole blood, plasma, and serum, which were used for DNA, RNA, cytokines, hematocrit, hemoglobin measures, thyroid hormones and measures of heavy metal (e.g., mercury, lead) and perfluorinated chemical exposure. Buccal cell samples were also collected at 14-26 weeks gestation from fathers for DNA extraction and future genetic and epigenetic studies. Purified DNA is available at the Gene Expression Omnibus data repository (<https://www.ncbi.nlm.nih.gov/geo/>). Midstream random urine samples (10 mL) were collected from expectant mothers at 14-26 weeks gestation and three months postpartum. Samples have been used in the Neurotoxicant study ($N = 546$) to examine maternal exposure to endocrine-disrupting hormones [2]. A breast milk sample was collected from breastfeeding mothers to determine fatty acids composition.

Patient and Public Involvement

There is an APrON Patient Engagement committee, who advises on research directions, but not design. Findings have been disseminated to participants through newsletters and an online conference.

FINDINGS TO DATE

A complete list of publications from APrON can be found at <http://www.apronstudy.ca/research/publications-and-press>. Key findings on maternal and paternal mental health and maternal nutrition outcomes have been summarized in Box 1.

Box 1 Key findings

Mental health outcomes

- In partnered couples, both mothers and fathers had depressive symptoms in 2.3% (*n* = 19) of couples. Low household income and prenatal maternal depression were associated with a higher probability of depressive symptoms in both partners. In 78.5% (*n* = 664) of couples, neither father nor mother experienced depressive symptoms. Postnatal social support was found to be a protective factor associated with decreased risk of maternal and paternal postpartum depressive symptoms. Risk factors for maternal postpartum depression symptoms included low household income, and prenatal EDS score > 10. For paternal postpartum depression symptoms, risk factors included low household income, prenatal EDS score of > 9, postpartum stressful life events, and smoking.
- Women with EDS score of < 10 were more likely to take micronutrients (e.g., vitamins B6, B9, B12, and E; essential fatty acids; selenium) compared to women with EDS score of < 10.
- One hundred and twenty-two women (13.4 %) had a score of 3 or more ACEs. Maternal exposure to ACEs was positively associated with both prenatal depression and anxiety.

Nutrition outcomes

- Approximately 70% of women who enter pregnancy with a body mass index (BMI) > 25 kg/m² were likely to exceed GWG guidelines in the beginning of the second trimester. Women with excessive weight gain also gained higher amounts of body fat and retain higher amounts of fat compared to women who gain within the guidelines. Women who consumed a healthy diet pattern prior to pregnancy were less likely to develop complications such as gestational hypertension.
- The median vitamin D intake from diet and supplements was 600 IU/day during pregnancy, which was not enough to achieve a target circulation 25(OH)D concentration. A significant relationship between maternal reported dietary vitamin D intake and plasma 25(OH)D and 3-epi-25(OH)D3 concentration were identified.
- Only 23% (*n*= 138) of mothers met the adequate intake (AI) recommendations for choline; the number was even lower (10%) in the postpartum period. Consuming eggs and milk during pregnancy increased mothers' likelihood of meeting choline AI recommendations.

Mental Health Outcomes

Four key papers [7-10] reported on perinatal depression and anxiety prevalence and risk factors, and the impact of adverse childhood experiences (ACEs) on maternal prenatal depression and anxiety. Letourneau et al. [8] reported that 10.7% (*n* = 68) of mothers experienced depressive symptoms in late pregnancy, defined as a score above the Edinburgh Depression Scale (EDS) cut-off of ≥ 10 for mothers [11]. Similarly, 10.7 % (*n* = 68) reported depressive symptoms at 3 months postpartum (8). For fathers, cut-off of $EDS \geq 9$ was used [12]. Nearly 14.0% of fathers (*n* = 139) reported depressive symptoms in either first or second trimester, with rates going down to 9.1% (58) at 3 months postpartum (8). Leung et al. [9] examined 846 cohabitating (i.e., married

and common-law) mother-father couples, with 2.3% ($n = 19$) of couples having both mothers and fathers reporting perinatal depressive symptoms. Neither mother nor father reported depressive symptoms in 78.5% ($n = 664$) of couples. In 9.5% ($n = 80$) of couples, only mothers experienced depressive symptoms during postpartum; similarly, in 9.8% ($n = 83$) of couples only fathers reported depressive symptoms. Low household income and prenatal maternal depression were associated with a higher incidence of depressive postpartum symptoms in both mothers and fathers. Risk factors for maternal postpartum depressive symptoms included the low household income and prenatal depressive symptoms. Risk factors for paternal postpartum depressive symptoms included low household income, prenatal depressive symptoms, postpartum stressful life events, and smoking. Postpartum social support was found to be a protective factor associated with decreased risk of maternal and paternal postpartum depressive symptoms. Additionally, Leung et al. [7] found that mothers who reported no depressive symptoms during the perinatal period were more likely to take micronutrients (e.g., vitamins B6, B9, B12, and E; essential fatty acids; selenium) compared to mothers who experienced perinatal depressive symptoms.

Letourneau et al. [10] reported that the higher ACEs score was positively associated with both prenatal depression and anxiety. ACEs were defined as traumatic events that include physical, sexual, and emotional abuse; household dysfunction; domestic violence; or living with an adult experiencing mental illness, substance use, or incarceration during childhood [13]. Nearly 14% ($n = 122$) of women reported having experienced 3 or more ACEs (10). This finding is consistent with previous research [14] that suggest about a graded relationship between the number of ACEs and later increased risk for health issues. Routine screening for ACEs may facilitate early identification and preventative interventions; however, this has been recently debated (14). In sum, the identified risks and protective factors provide opportunities for prevention of perinatal depression and anxiety in mothers and fathers.

Nutrition Outcomes

Gestational weight gain and dietary intake, and perinatal nutrient intake and status were examined in 8 key papers (15-18, 20-22, 26).

Gestational Weight Gain and Dietary Intake

Gestational weight gain over or below Health Canada's Gestational Weight Gain (GWG) Guidelines is associated with an increased risk of adverse outcomes for mother and infant [15]. The APrON cohort was the first to provide evidence that more than half of women gain more than Health Canada's Guidelines, using prospectively collected body weights from across pregnancy [16]. Since then, several studies have been completed with body weight or body composition in pregnancy or postpartum as a main outcome. Analysis of the APrON data by Jarman et al. [17] provided a clear picture indicating that ~70% of women who enter pregnancy with a body mass index (BMI) > 25 kg/m² are likely to exceed GWG guidelines, by about 19 weeks gestation. Women with excessive weight gain also gain higher amounts of body fat and retain higher amounts of fat compared to women who gain within the guidelines [18]. The detailed dietary intake information that was collected showed that women who consumed a healthy diet pattern prior to pregnancy were less likely to develop complications such as gestational hypertension [19]. An individualized approach to supportive dietary counseling that considers pre-pregnancy BMI in addition to the woman's social and financial context is key to helping women meet national recommendations for both diet and GWG in pregnancy [20].

Nutrient Intake and Status

Vitamin D is critical for a healthy pregnancy [21]. Three studies that included APrON participants examined the role of maternal vitamin D intake during pregnancy [22-24]. Aghajafari et al. [23] reported that the median vitamin D intake from diet and supplements was 600 IU/day during pregnancy, which was not sufficient to achieve a target circulation 25(OH)D

concentration. When the 3-epimer was included in the estimation of vitamin D status, the prevalence of vitamin D insufficiency (<75 nmol/L) was lower (33.0 %) compared to when it was excluded (38.0 %). Vitamin D supplementation (2000 IU/day) was associated with 25(OH)D₃ sufficiency [22-24]. A significant relationship between maternal reported dietary vitamin D intake and plasma 25(OH)D and 3-epi-25(OH)D₃ concentration were identified.

Choline intake during pregnancy and lactation is critical to the offspring's brain function [25]. Lewis et al.[26] reported that the mean choline intake in pregnant and lactating women ranged between 340 (SD 148) in the second trimester and 346 (SD 151) mg/day at 3 months postpartum. Only 23% ($n = 138$) of mothers met the adequate intake (AI) recommendations for choline; the number was even lower (10%) in the postpartum period [26]. Consuming eggs and milk during pregnancy increased mothers' likelihood of meeting choline AI recommendations. These findings contributed to the European Food Safety Authority (EFSA) dietary reference values for choline guidelines [27]. To conclude, a significant number of women were not meeting recommendations for vitamin D and choline intake during pregnancy and postpartum. Increased consumption of these nutrients may be necessary for pregnant and lactating women to improve the health and development of their infants.

STRENGTHS AND WEAKNESSES

The major strengths of APrON are recruitment and retention of the large community-based cohort during pregnancy, postpartum and early childhood. The APrON study used a detailed array of standardized validated psychological and physiological measures and biospecimens. The prospective data collection and follow-up enabled investigation of a wide range of maternal and paternal health outcomes. The APrON team has been successful in continuously engaging participants, as evidenced by high response rates over the years (see Table 4).

Table 4. *Response Rates for Questionnaires*

Data collection time point	Targeted participants (n)	Participants responded (n)	Response rate (%)
Mothers < 14 weeks gestation	2189	2145	97.9
Mothers at 14-26 weeks gestation			
Mothers at 14-26 weeks gestation and those recruited at < 14 weeks gestation	563	475	84.3
Fathers at 14-26 weeks gestation	1325	1315	99.2
Mothers at 27-40 weeks gestation	1838	1830	99.5
Mothers at 3 months postpartum	2051	1811	88.3
Fathers at 3 months postpartum	1287	1150	89.3
Mothers at 6 months postpartum	1727	1523	88.1
Mothers at 1 year postpartum	1747	1305	74.7
Mothers at 2 year postpartum	1809	1299	71.8
Mothers at 3 year postpartum	1648	1282	77.7

Limitations of this study include the sample characteristics that are skewed toward a high percentage of white, well-educated and married participants. Consequently, generalizability of findings to those with different socio-demographic backgrounds should be made with caution. Another limitation is that selective attrition bias has likely occurred, which may lead to underestimation of the effect of parental mental health and lifestyle choices on the child's development. The study likely included participants with healthier lifestyle behaviors or with

more positive attitudes towards research, because such characteristics are associated with higher socioeconomic status [28]. The differences in the demographic characteristics of the continuing participants compared with discontinuing participants should be considered when interpreting the findings. There were also fewer fathers than mothers in the sample; thus, most developmental and behavioral outcomes were based on maternal reports.

FURTHER DETAILS

Collaboration

Information about the APrON study is available at www.apronstudy.ca. The APrON data is available through SAGE (Secondary Evidence to Generate Evidence; <https://policywise.com/sage/>). For more information contact Dr. Nicole Letourneau at Nicole.letourneau@ucalgary.ca. Collaboration or data access inquiries will be considered by the APrON study team.

Profile in a nutshell

- The APrON study is a longitudinal cohort of mothers, fathers and their children investigating maternal, parental, birth, and child development outcomes.
- A total of 2189 women (aged 16 to 44) and 1325 fathers (aged 18 to 52), residing around either Calgary or Edmonton, Alberta, enrolled and completed at least one questionnaire between May 2009 and June 2012. The majority were married, educated, had annual household incomes >\$70 000 and were born in Canada.
- Mothers completed 18 questionnaires and up to 4 clinic visits spanning pregnancy to three years postpartum and provided access to their medical records as well as three to four blood and two urine samples. To date, 1648 mothers and 1255 fathers remain eligible for follow-up.
- The APrON dataset includes comprehensive socio-demographic and psychological data from pregnancy to three years postpartum, as well as maternal blood and urine samples. The 8-year data collection follow-up is ongoing and 12-year data follow-up will start in 2021.

Requests for data and collaboration are welcomed, contact Dr. Nicole Letourneau at Nicole.letourneau@ucalgary.ca

Data Availability Statement

Data can be collected at the Secondary Analysis to General Evidence (SAGE) data repository or by contacting Dr. Nicole Letourneau (the principal investigator for the APrON Study).

Funding

This work was supported by the Canadian Institutes of Health Research Doctoral Award (CIHR, grant number 394652).

Acknowledgements

The authors acknowledge the contributions of all the families who took part in the APrON study, and the investigators, managers, research assistants, graduate and undergraduate students, volunteers and clerical staff of the APrON study team.

Conflict of Interest

None declared.

AUTHOR CONTRIBUTORSHIP STATEMENT

1. Elena Ali designed the manuscript, drafted the whole manuscript, analyzed and interpreted statistical data; created data tables; worked with co-authors to facilitate feedback and group work on the manuscripts; incorporated co-authors comments into manuscripts; edited the draft of the whole manuscript for content, grammar, format and style to create a final version for submission to the journal; revised and approved final version of manuscript; and agreed to be accountable for all aspects of the work involved in this manuscript.
2. Nicole Letourneau conceived the APrON study and the idea for manuscript, facilitated involvement with co-authors in creation of manuscript; designed the manuscript; obtained data; revised the whole draft for grammar, content, and style; provided final approval of the final version; agreed to be accountable for all aspects of the work.

3. Fariba Aghajafari analyzed and interpreted data; participated in discussions about design of the manuscript; drafted and revised a Nutrient Intake and Status paragraph in the manuscript; provided approval of final version; agreed to be accountable for all aspects of the work.
4. Rhonda Bell contributed to design of the manuscript by suggesting what data to include in the manuscript; obtained and analyzed data; drafted and revised Gestational Weight Gain paragraph in the manuscript; provided final approval; and agreed to be accountable for all aspects of the work.
5. Andrea Deane obtained and interpreted data; revised draft for statistical content accuracy description of the methods; assisted with creating data tables; provided final approval for the manuscript; and agreed to be accountable for all aspects of the work.
6. Deborah Dewey obtained and interpreted data; contributed to design of the draft by suggesting what to include; revised manuscript for brevity and intellectual content; approved final version; and agreed to be accountable for all aspects of the work.
7. Catherine Field obtained, analyzed and interpreted data; drafted the Nutrient Intake and Status paragraph in the manuscript, and provided comments for the whole draft; provided final approval of the manuscript; and agreed to be accountable for all aspects of the work.
8. Gerald Giesbrecht conceived the APrON study; obtained and analyzed data; revised the draft; provided final approval of the final version; agreed to be accountable for all aspects of the work.
9. Bonnie Kaplan conceived the idea for the APrON study; overseen the obtainment of data; revised the manuscript; provided final approval; and agreed to be accountable for all aspects of the work.
10. Brenda Mun Ying Leung conceived the idea for the APrON study; overseen the obtainment of data; revised the manuscript; provided final approval; and agreed to be accountable for all aspects of the work.
11. Henry Ntanda analyzed and interpreted data; revised the draft for statistical accuracy; approved final version; and agreed to be accountable for all aspects of the work.
12. APrON Team includes a large number of individuals who all had all met the ICMJE criteria for

authorship.

FIGURE LEGENDS

- Figure 1. Socio-demographic characteristics of women in APrON, Calgary, Alberta, and Canada
- Figure 2. Data collection procedures

REFERENCES

1. Leung, B.M., et al., *Perinatal nutrition in maternal mental health and child development: Birth of a pregnancy cohort*. Early Hum Dev, 2016. **93**: p. 1-7.
2. Kaplan, B.J., et al., *The Alberta Pregnancy Outcomes and Nutrition (APrON) cohort study: rationale and methods*. Matern Child Nutr, 2014. **10**(1): p. 44-60.
3. Tough, S.C., et al., *Cohort Profile: The All Our Babies pregnancy cohort (AOB)*. Int J Epidemiol, 2017. **46**(5): p. 1389-1390k.
4. Manca, D.P., et al., *The most effective strategy for recruiting a pregnancy cohort: a tale of two cities*. BMC Pregnancy Childb, 2013. **13**: p. 75.
5. Leung, B.M., et al., *Comparison of sample characteristics in two pregnancy cohorts: community-based versus population-based recruitment methods*. BMC Med Res Methodol, 2013. **13**: p. 149.
6. Dzakpasu, S., et al., *The Canadian maternity experiences survey: design and methods*. J Obstet Gynaecol Can, 2008. **30**(3): p. 207-216.
7. Leung, B.M., et al., *Prenatal micronutrient supplementation and postpartum depressive symptoms in a pregnancy cohort*. BMC Pregnancy Childb, 2013. **13**: p. 2.

8. Letourneau, N., et al., *Maternal and paternal perinatal depressive symptoms associate with 2- and 3-year-old children's behaviour: findings from the APrON longitudinal study*. BMC Pediatr, 2019. **19**(1): p. 435.
9. Leung, B.M., et al., *Predictors of Postpartum Depression in Partnered Mothers and Fathers from a Longitudinal Cohort*. Community Ment Health J, 2017. **53**(4): p. 420-431.
10. Letourneau, N., et al., *Intergenerational transmission of adverse childhood experiences via maternal depression and anxiety and moderation by child sex*. J Dev Orig Health Dis, 2019. **10**(1): p. 88-99.
11. Bergink, V., et al., *Validation of the Edinburgh Depression Scale during pregnancy*. J Psychosom Res, 2011. **70**: p. 385-389.
12. Paulson, J.F. and S.D. Bazemore, *Prenatal and Postpartum Depression in Fathers and Its Association With Maternal Depression*. JAMA, 2010. **303**(19): p. 1961-1969.
13. Felitti, V.J., et al., *Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study*. Am J Prev Med, 1998. **14**(4): p. 245-58.
14. Hughes, K., et al., *The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis*. Lancet Publ Heal, 2017. **2**(8): p. e356-e366.
15. Government of Canada. *Prenatal Nutrition Guidelines for Health Professionals: Gestational Weight Gain*. 2010 [cited 2020 21 Jan]; Available from: <https://www.canada.ca/en/health-canada/services/canada-food-guide/resources/prenatal-nutrition/eating-well-being-active-towards-healthy-weight-gain-pregnancy-2010.html>.
16. Begum, F., et al., *Gestational weight gain and early postpartum weight retention in a prospective cohort of Alberta women*. J Obstet Gynaecol Can, 2012. **34**(7): p. 637-47.

17. Jarman, M., et al., *Patterns and trajectories of gestational weight gain: a prospective cohort study*. CMAJ Open, 2016. **4**(2): p. E338-45.

18. Subhan, F.B., et al., *Association of pre-pregnancy BMI and gestational weight gain with fat mass distribution and accretion during pregnancy and early postpartum: a prospective study of Albertan women*. BMJ Open, 2019. **9**(7): p. e026908.

19. Jarman, M., et al., *Dietary Patterns Prior to Pregnancy and Associations with Pregnancy Complications*. Nutrients, 2018. **10**(7).

20. APrON and ENRICH Study Teams, *Adherence to Canada's Food Guide Recommendations during Pregnancy: Nutritional Epidemiology and Public Health*. Curr Dev Nutr, 2017. **1**(7): p. e000356.

21. Aghajafari, F., et al., *Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies*. Bmj, 2013. **346**: p. f1169.

22. Aghajafari, F., et al., *Plasma 3-Epi-25-Hydroxycholecalciferol Can Alter the Assessment of Vitamin D Status Using the Current Reference Ranges for Pregnant Women and Their Newborns*. J Nutr, 2016. **146**(1): p. 70-5.

23. Aghajafari, F., et al., *The Current Recommended Vitamin D Intake Guideline for Diet and Supplements During Pregnancy Is Not Adequate to Achieve Vitamin D Sufficiency for Most Pregnant Women*. PLoS One, 2016. **11**(7): p. e0157262.

24. Aghajafari, F., et al., *The High Prevalence of Vitamin D Insufficiency in Cord Blood in Calgary, Alberta (APrON-D Study)*. J Obstet Gynaecol Can, 2017. **39**(5): p. 347-353.e1.

25. Wallace, T.C., et al., *Choline: The Neurocognitive Essential Nutrient of Interest to Obstetricians and Gynecologists*. J Diet Suppl, 2019: p. 1-20.

- 1
2
3 26. Lewis, E.D., et al., *Estimation of choline intake from 24 h dietary intake recalls and*
4 *contribution of egg and milk consumption to intake among pregnant and lactating women*
5 *in Alberta*. Br J Nutr, 2014. **112**(1): p. 112-21.
6
7
8
9
10 27. European Food Safety Authority, *Dietary reference values: EFSA publishes advice on*
11 *choline*. 2016. p. 1-70.
12
13
14 28. Daniels, J.L., et al., *Attitudes toward participation in a pregnancy and child cohort study.*
15 *Paediatr Perinat Epidemiol*, 2006. **20**(3): p. 260-6.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Socio-demographic characteristics of women in APrON, Calgary, Alberta, and Canada





Figure 2. Data collection procedures



BMJ Open

The Alberta Pregnancy Outcomes and Nutrition Cohort (APrON) Study: Cohort Profile and Key Findings from the First Three Years

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-047503.R1
Article Type:	Cohort profile
Date Submitted by the Author:	23-Aug-2021
Complete List of Authors:	Letourneau, Nicole; University of Calgary, Faculty of Nursing; University of Calgary Cumming School of Medicine, Pediatrics, Psychiatry and Community Health Sciences Ali, Elena; University of Calgary, Faculty of Nursing Aghajafari, Fariba; University of Calgary, Family Medicine; University of Calgary Cumming School of Medicine, Community Health Sciences Bell, Rhonda; University of Alberta, Agricultural, Food and Nutritional Science Deane, Andrea; University of Calgary Cumming School of Medicine, Pediatrics Dewey, Deborah ; University of Calgary Cumming School of Medicine, Paediatrics; University of Calgary Cumming School of Medicine, Community Health Sciences Field, Catherine; University of Alberta, Department of Agricultural, Food and Nutritional Science Giesbrecht, Gerald ; University of Calgary, Pediatrics; University of Calgary, Community Health Sciences Kaplan, Bonnie; University of Calgary Cumming School of Medicine, Pediatrics Leung, Brenda; University of Calgary Cumming School of Medicine, Community Health Sciences; University of Lethbridge, Faculty of Health Sciences Ntanda, Henry; University of Calgary Cumming School of Medicine, Pediatrics
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Nutrition and metabolism, Paediatrics, Public health
Keywords:	NUTRITION & DIETETICS, MENTAL HEALTH, PAEDIATRICS, PUBLIC HEALTH

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Word count: 4457 words

Keywords: Nutrition & dietetics, mental health, paediatrics, public health

Acknowledgements

The authors acknowledge the contributions of all the families who took part in the APrON study, and the investigators, managers, research assistants, graduate and undergraduate students, volunteers and clerical staff of the APrON study team.

Abstract

Objectives. The objectives of the ongoing Canadian longitudinal cohort called the Alberta Pregnancy Outcomes and Nutrition (APrON) Study were to: (1) determine the relationship between maternal nutrient intake and status before, during, after pregnancy, and (a) maternal mental health, (b) pregnancy and birth outcomes, and (c) infant/child neurodevelopment and behavior; (2) identify maternal mental health and nutrient predictors of child behaviour; and (3) establish a DNA biobank to explore genomic predictors of children’s neurodevelopment and behavior. Drawn mostly from data collected during the period from pregnancy to when children are 3 years of age, the objectives of this paper are to describe the participants, measures, and key findings on maternal and paternal mental health, maternal nutrition, and child outcomes. However data have been collected to 8 years of age, with planned follow-up at 12 years of age.

Data Collection. Data comprise: questionnaires completed by pregnant women/mothers and their partners (usually fathers) on mothers’, fathers’ and children’s health; dietary interviews; clinical assessments; linkage to hospital obstetrical records; and biological samples such as DNA.

Key Findings. Mental health, nutrition and child outcomes are presented. For example, APrON women who consumed more selenium and omega-3 were less likely to develop symptoms of perinatal depression. Couples in which both mothers and fathers were affected by perinatal depression reported lower incomes and higher maternal prenatal depressive symptoms and lower support from fathers postnatally and their children presented with the most behavioural problems. Higher prenatal consumption of choline rich foods such as eggs and milk were recommended as was vitamin D supplementation for both mothers and children to meet guidelines. Maternal experiences of early adversity predicted increased likelihood of perinatal depression and anxiety and children’s behavioural problems.

Conclusion. The APrON cohort offers a unique opportunity to advance understanding of the developmental origins of health and disease.

Strengths and Limitations

1. APrON is a large, ongoing longitudinal study designed to understand the early origins of health and disease;
2. APrON has collected data to eight years of age and has planned follow up at 12 years of age;
3. APrON has retained more than 88% of participants at the 8-year data collection; and
4. A limitation is that the cohort is largely low-risk, limiting generalizability of study findings to higher-risk populations.

Who is in the APrON cohort?

Women were recruited into the cohort if they were 16 years of age or older, able to speak and read English, < 27 weeks gestation and willing to come for on-site clinic visits at the University of Calgary, Calgary or University of Alberta, Edmonton, Alberta. Partners were eligible if they were cohabitating with the participating woman. Participants were excluded if they planned to move out of the region during the clinic visit timeframe of pregnancy to 3 months postpartum. A total of 2189 pregnant women (aged 16 to 44) and 1325 men (aged 18 to 52), residing around Calgary or Edmonton, were enrolled in the study between May 2009 and June 2012. The APrON study was approved by the University of Calgary Health Research Ethics Board (REB14-1702) and University of Alberta Health Research Ethics Biomedical Panel (Pro00002954). To assess the representativeness of the APrON cohort (see Table 1), sociodemographic characteristics of participants at the time of recruitment were compared to characteristics of women with children in Calgary, Alberta, and Canada [6, 8, 9].

Patient and Public Involvement

APrON cohort participants were not involved in the development of the initial study design or in the recruitment of participants. However, APrON established a participant advisory committee (PAC) shortly after recruitment ended in 2012. Over time, 70 parents have participated on the PAC, with approximately 10 to 15 parents taking part in twice yearly meetings to advise the Principal and Co-Investigators on a variety of topics, for example, how to keep participants engaged and interested in APrON research, priorities for future APrON research, and feedback on laymen research summaries posted to APrON's website. APrON has also extensively disseminated knowledge to the public via various modalities including: multiple presentations to various stakeholders, APrON's Newsletter, website (<https://apronstudy.ca/>) and conferences. Held via Zoom due to COVID-19 on October 29-30th, 2020, APrON hosted an online conference

titled *Celebrating 10 Years: Alberta Pregnancy Outcomes and Nutrition (APrON) Achievements & Evolution Conference*, that attracted over 300 attendees, many who were APrON participants. The APrON cohort PAC members held their own panel session where they spoke to the attendees about their experiences with APrON, why they think the research is so important and how we can continue for years to come.

For peer review only

Table 1. *Participant Intake Characteristics (N = 3514)*

For peer review only

Category	Maternal (n / %)	Paternal (n / %)
Demographics		
Age (years)		
16 – 24	100 (4.6)	19 (1.4)
25 – 29	481 (22.0)	216 (15.4)
30 -34	980 (44.78)	493 (35.2)
35 -39	512 (23.4)	329 (23.5)
≥ 40	105 (4.8)	162 (11.6)
Missing age information	11 (0.5)	181 (12.9)
Marital status		
Married/common-law	2035 (93.0)	1275 (91.1)
Single/divorced/separated	97 (4.4)	27 (1.9)
Missing marital information	57 (2.6)	98 (7.0)
Education		
Less than a high school diploma	59 (2.7)	31 (2.2)
Completed high school diploma	202 (9.2)	147 (10.5)
Completed trade, technical	404 (18.5)	376 (26.9)
Completed university or more	1442 (66.8)	746 (53.3)
Missing education information	83 (3.7)	100 (7.1)
Ethnicity		
White	1693 (77.3)	1082 (77.3)
Non-white	485 (22.2)	206 (14.7)
Missing ethnic information	11 (0.5)	112 (8.0)
Mothers' Report on total household income		
< \$40,000	190 (8.7)	-
\$40,000 - \$69,999	279 (12.7)	-
\$70,000 - \$99,999	464(21.2)	-
>\$100,000	1146 (52.3)	-
Missing household income information	110 (5.0)	-
Born in Canada		
Yes	1618 (73.9)	1024 (73.1)
No	481 (22.0)	277 (19.8)
Missing country of birth	90 (4.1)	99 (7.1)
Gravidity		
Multigravida	1107 (51.6)	n/a
Primigravida	996 (46.4)	n/a
Missing gravidity information	42 (2.0)	n/a
Parity		
0	1222 (55.8)	n/a
1	732 (33.4)	n/a
2	181 (8.3)	n/a
≥3	37 (1.7)	n/a
Missing parity information	17 (0.8)	n/a

A greater proportion of women in the cohort (28.0%) were over 34 years old compared with women in Calgary (20.0%), Alberta (16.0%), and Canada (18.0%). More APrON women were married (95%) compared with women in Calgary (73.0%), Alberta (70.0%), and Canada (60.0%). A greater proportion of women had household income of \$70,000 or more (77.0%) compared to Calgary (65.0%), Alberta (61.0%), and Canada (58.0%). More APrON women completed post-secondary education (87.0%) compared to women in Calgary (69.0%), Alberta (70.0%), and Canada (72.0%). The proportion of APrON women who were born outside of Canada (23.0%) was similar to the proportions in Calgary (25.0%), Alberta (22%), and Canada (20.0%) (see Figure 1).

How often have APrON families been followed up?

Between 2009 and 2012 the initial cohort completed questionnaires and clinical assessments three times during pregnancy, and at 3 months postpartum. Follow-up questionnaires were provided when their child was 6 months, and 1, 2 3, 5 and 8 years of age (see Figure 2). We plan a 12 year follow-up beginning in 2022.

In total, 2189 women and 1325 men completed at least one questionnaire. Of the women, and their children (88.9%) are continuing participants, defined as those who did not become ineligible or were lost to follow-up. Participants who discontinued the study included women who miscarried, had a stillbirth, lost custody of their child, or for whom a maternal/child death was reported ($n = 52$, 2.4%); moved out of Calgary or Edmonton before three months postpartum ($n = 43$, 2.0%); or withdrew from the study citing loss of interest ($n = 149$; 6.8%). The demographic characteristics of participants who continued in APrON compared with those that discontinued are provided in Table 2.

Table 2. Comparison of Demographic Characteristics of Continuing and Discontinued Participants

Characteristic	Continuing participants n (%)	Discontinuing participants			P-value*
		Become ineligible n (%)	Lost to follow-up n (%)	Dropped out n (%)	
Maternal age					
16 – 24	71 (3.6)	8 (18.6)	15 (34.8)	6 (4.1)	< 0.001
25 – 29	424 (16.2)	7 (16.2)	7 (16.2)	43 (29.2)	
30 – 34	899 (46.2)	15 (34.8)	14 (32.5)	51 (34.7)	
35 – 39	454 (23.3)	9 (20.9)	7 (16.2)	41 (27.9)	
≥40	95 (4.8)	4 (9.3)	0 (0.0)	6 (4.1)	
Marital status					
Single	72 (3.8)	12 (24.5)	8 (21.1)	5 (3.8)	0.007
Married	1838 (96.2)	37 (75.5)	30 (78.9)	128 (96.2)	
Maternal education					
Less than a high school diploma	41 (2.2)	6 (16.2)	8 (20.5)	4 (3.01)	<0.001
Completed high school diploma	164 (8.6)	9 (24.3)	9 (23.1)	20 (15.0)	
Completed trade, technical diploma	362 (19.1)	5 (13.5)	10 (25.6)	26 (19.5)	
Completed university or more	1329 (70.1)	17 (45.9)	12 (30.6)	83 (62.4)	
Household income					
< \$40,000	151 (8.1)	12 (30.8)	19 (54.3)	8 (6.15)	<0.001
\$40,000-\$60,000	252 (13.4)	6 (15.4)	5 (14.3)	16 (12.31)	
\$70,000-\$99,999	428 (22.8)	7 (17.9)	4 (11.4)	24 (18.46)	
≥ \$100,000	1042 (55.6)	14 (35.9)	7 (20.0)	82 (63.08)	
Born in Canada					
No	425 (22.5)	13 (32.5)	14 (37.8)	28 (21.2)	0.186
Yes	1463 (77.5)	27 (67.5)	23 (62.2)	104 (78.8)	

Note. Assessing the null hypothesis that there is no difference in distributions between those participants who continued and those who did not continue in the study; chi-square test

Continuing participants were more likely to be older, married, and to have higher education and household incomes (all *ps* < 0.05). The proportion of continuing participants born in Canada was similar to discontinuing participants. When mothers were lost to follow-up, so too were fathers and children.

What has the APrON study measured?

Women were asked to complete questionnaires three times during pregnancy (<14, 14-26, and 27-40 weeks gestation, number of scales administered was 37) and seven times postnatally (3 and 6 months, 1, 2, 3, 5, and 8 years, number of scales administered was 77). Partners (usually self-reported fathers) were asked to complete questionnaires when their partners were 14-26 weeks gestational age (number of scales administered was 16), and twice postnatally (3 months and 5 years, number of scales administered was 14). Questionnaires administered during pregnancy measured previous and current mental and physical health status, nutritional, medication and supplement intake, lifestyle choices, healthcare services used, social support, attitudes towards breastfeeding, and sociodemographic variables. The follow-up postpartum questionnaires collected information about labor and delivery, postpartum medical problems, maternal and paternal mental health and stress, nutritional, medication and supplement intake, lifestyle choices, food security, parenting, experiences in close relationships, and sociodemographics (see Figure 2). Measures were selected based on several synergistic priorities. First, the APrON team had to judge measures to be reliable and valid indicators of the constructs of interest. Second, measures had to minimize burden to participants in terms of time and difficulty of completion, for example measures with fewer items and more accessible language were valued most. Finally, an APrON team member had to express an interest in using the measure. Full details and documentation of all APrON measures is available on request.

Clinic visits were undertaken to gather maternal anthropometric measurements during pregnancy and three months postpartum. These measures included height, pre-pregnancy and current weight, highest weight during pregnancy, circumference measurements (i.e., mid-upper arm, waist, hip, thigh), and skinfold thickness (i.e., biceps, triceps, subscapular, suprailiac, thigh). Each measurement was taken three times and averages were reported for analyses. Paternal height and weight were collected during the prenatal period. One objective of APrON was to

establish a biobank for future research that would allow investigation on genetics, epigenetics [1], nutrient exposures and neurotoxicants. Thus, biosamples were also collected from mothers, their partners and from infants during clinic visits. Midstream random urine samples (10 mL) were collected from expectant mothers at 14-26 weeks gestation and three months postpartum. A breast milk sample was collected from breastfeeding mothers to determine fatty acid composition. Maternal non-fasting venous blood samples were collected from mothers at <14, 14-26, and 27-40 weeks gestation, and at three months postpartum. Maternal blood collection provided whole blood, plasma, and serum, which were used for DNA, RNA, cytokines, hematocrit, hemoglobin measures, thyroid hormones and measures of heavy metals (e.g., mercury, lead) and perfluorinated chemical exposure. Buccal cell samples were also collected at 14-26 weeks gestation from partners for DNA extraction and future genetic and epigenetic studies. From infants, buccal and/or blood samples were collected at 3 months of age for DNA extraction. Purified DNA is available at the Gene Expression Omnibus data repository (<https://www.ncbi.nlm.nih.gov/geo/>). A detailed list of data collected up to 8 years of age on the full sample is provided in Table 3.1 for mothers, Table 3.2 for fathers and Table 3.3 for children.

Table 3.1. *Maternal Variables and Biological Samples Collected via Questionnaires and Clinic Visits*

Variables	<14 weeks	14-26 weeks	27-40 weeks	3 months postpartum	6 months postpartum	1 year	2 years	3 years	5 years	8 years
Socio-demographics										
Household income	X								X	X
Education	X									
Marital status	X								X	X
Age	X									
Primary language	X									
Born in Canada	X									
Family background		X	X	X	X	X	X	X	X	X
Family history		X								
Health and lifestyle										
General health				SF-8	SF-8	SF-8	SF-8	SF-8		
Current medical conditions and medications	X	X	X	X	X	X	X	X		
Substance use	X	X	X	X	X	X	X	X	X	
Parenting									X	X
Pregnancy history										
Pre-pregnancy physical activity	X	X	X	X	X	X	X	X		
Physical activity	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	GLTE	
Dental care	X	X	X							
Dietary changes since pregnancy	X									
Food security			CCHSC		CCHSC					
Nutrition counselling information				X						
Diet history pre-pregnancy		FFQ								
Nausea and vomiting	PUQUE	PUQUE	PUQUE							
Breastfeeding information				X	X	X	X	X		
24-hour recall of foods consumed	X	X	X	X	X	X	X	X		

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Psychosocial health										
Depression										
Stress	EDS SLEQ	EDS	EDS SLEQ	EDS SLEQ	EDS	EDS SLEQ	EDS SLEQ	EDS SLEQ PSS	CESD SLEQ, PSS	CESD SLEQ,
Social support										
Attachment	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ	SSQ, SSI, ECR	SSI, ECR
DSM-IV Axis I disorders		PDSQ		PDSQ						
Anxiety	SCL-90-R	SCL-90-R	SCL-90-R	SCL-90-R	SCL-90-R	SCL-90-R	SCL-90-R X	SCL-90-R	SCL-90-R	STAI
Adverse childhood events										
History of mental health	X	X	X	X		X	X	X		
Anthropometrics (BMI/height/weight)	X	X	X							
Personality traits										
Personality traits										TIPI
Biological specimens										
Urine	X	X	X	X						
Venous blood		X		X						
Breast milk				X						

Note. X refers to investigator-developed measures. SCL-90-R, Symptom Checklist-90-Revised; EDS, Edinburgh Depression Scale; PUQE, Pregnancy-Unique Quantification of Emesis and Nausea questionnaire; SLEQ, Stressful Life Events Questionnaire ; SSQ, Social Support Questionnaire; PDSQ, Psychiatric Diagnostic Screening Questionnaire; ACEs, Adverse Childhood Experiences Questionnaire; SF-8, Short Form Health Survey-8; CCHSC, Canadian Community Health Survey Cycle 2.2, modified; GLTE, Godin Leisure-Time Exercise Questionnaire; FFQ, Food Frequency Questionnaire, adapted from the Canadian version of the Diet History Questionnaire; SSI, Social Support Index; ECR, Experiences in Close Relationship Scale; PSS, Perceived Stress, CES-D, Centre for Epidemiologic Studies Depression Scale; OECD, Organization for Economic Cooperation and Development Guidelines on Measuring Subjective Wellbeing; STAI, State-Trait Anxiety Inventory; TIPI, Ten-Item Personality Inventory. *Breastfeeding mothers only.

136/bmjopen-2020-047503 first published February 17, 2022. Downloaded from <http://bmjopen.bmj.com/> on April 28, 2024 by guest. Protected by copyright.

Table 3.2. *Paternal Variables and Biological Specimens Collected via Questionnaires and Clinic Visits*

Variables	<14 weeks gestation	14-26 weeks	27-40 weeks	3 months postpartum	5 years postpartum
Socio-demographics					
Household income			X		
Education			X		
Marital status			X		
Age			X		
Primary language			X		
Born in Canada			X		
Family background			X		X
Family history*			X		
Health and lifestyle		X	X		
General health					
Current medical conditions and medications		X	X	X	
Substance use		X	X	X	X
Physical Activity Anthropometrics (BMI/height/weight)		GLTE X	GLTE X	GLTE X	GLTE
Psychosocial health					
Depression		EDS	EDS	EDS	EDS
Social support		SSQ	SSQ	SSQ	SSQ
Stress			SLEQ		SLEQ
Anxiety					SCL-90-
Biological specimens*					
Buccal cell swab			X		
DNA from blood			X		

Note. X refers to investigator-developed measures. EDS, Edinburgh Depression Scale; SLEQ, Stressful Life Events Questionnaire; SSQ, Social Support Questionnaire; GLTE, Godin Leisure-Time Exercise Questionnaire.

Table 3.3. *Child Variables and Biological Specimens Collected in Questionnaires and Clinic Visits*

Variables	3 months	6 months	1 year	2 years	3 years	5 years	8 years
Health and Development		X	X	X	X	X	X
Occupation	X	X	X	X	X	X	
Childcare	X	X	X	X	X	X	
Vaccines						X	
Child food and liquid intake	X	X	X	X	X	X	
Children’s eating and drinking behaviours						X	X
Children’s activities						X	X
Children’s injuries						X	
Community							X
Children’s education							X
General infant behaviours							
Infant Behavior Questionnaire (IBQ)	X	X	X	X			
Scales of Independent Behaviors (SIB-R)	X	X	X	X	X	X	
Fussing and Crying	X	X	X	X			
Sleeping		X	X	X	X	X	X
Early Childhood Behavior Questionnaire (ECBQ)				X			
Children’s Behavior Questionnaire (CBQ)					X		
Child Behaviour Checklist Ages 1.5 – 5 (CBCL)				X	X		
Behaviour Rating Inventory of Executive Function- Preschool (BRIEF-P)				X	X		
Ages and Stages Questionnaire – Third Edition (ASQ-3)						X	
Children’s Communication Checklist – Second Edition (CCC-Q)						X	
Behaviour Assessment System for Children – Second Edition (BASC-2)						X	X
Media use						X	X
Quantitative Checklist for Autism in Toddlers				X	X		
Temper Tantrum Scale – Past 2 Weeks				X	X		
Temper Tantrum Scale – Past 6 Months				X	X		
Parenting						X	X
Biological specimens							
Buccal cells	X						
Venous blood	X						

136/bmjopen-2020-047503 on 7 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 28, 2024 by guest. Protected by copyright.

Numerous sub-studies, employing portions of the APrON sample have employed the originally collected biosamples in related research, for example the Neurotoxicant Study (n=546) that examined maternal exposure to endocrine hormone disrupting chemicals (i.e. bisphenol a and phthalates) on child outcomes [e.g., 10, 11-15] and the Parenting Research on Mental Illness, Stress and Epi/genetics (PROMISE; n=276) Study of gene by environment interactions [e.g., 16, 17-20]. Other sub-studies have collected additional biosamples for related research, for example maternal and infant cortisol and other measures such heart rate variability and parent-child interaction quality observations in the Fetal Programming Study [n=276; e.g., 21, 22-29].

What has APrON found?

A complete list of publications from APrON can be found at <http://www.apronstudy.ca/research/publications-and-press>. Key findings from the full sample on maternal and paternal mental health, maternal nutrition and child outcomes have been summarized in Box 1 and below.

Box 1 Key findings

Mental health outcomes

Mental health outcomes

- In partnered couples, both mothers and fathers had depressive symptoms in 2.3% of couples. Low household income and prenatal maternal depression were associated with a higher probability of depressive symptoms in both partners. In 78.5% of couples, neither father nor mother experienced depressive symptoms. Postnatal social support was found to be a protective factor associated with decreased risk of maternal and paternal postpartum depressive symptoms. Risk factors for maternal postpartum depression symptoms included low household income, and prenatal EDS score ≥ 10 . For paternal postpartum depression symptoms, risk factors included low household income, prenatal EDS score of < 10 , postpartum stressful life events, and smoking.
- Women with EDS scores < 10 were more likely to take micronutrients (e.g., vitamins B6, B9, B12, and E; essential fatty acids; selenium); however only selenium and omega-3 were significantly different between depressed and non-depressed groups, with non-depressed mothers consuming higher amounts.
- 14% of mothers had a score of three or more adverse childhood experiences. Maternal exposure to early adversities predicted higher reported symptoms of perinatal depression and anxiety.

Nutrition outcomes

- Approximately 70% of women who entered pregnancy with a body mass index (BMI) $> 25 \text{ kg/m}^2$ were likely to exceed gestational weight gain guidelines in the beginning of the second trimester. Women with excessive weight gain also gained higher amounts of body fat and retained higher amounts of fat compared to women who gained

within the guidelines. Women who consumed a healthy diet pattern prior to pregnancy were less likely to develop complications such as gestational hypertension.

- The median vitamin D intake from diet and supplements was 600 IU/day during pregnancy, which was not enough to achieve a target plasma circulation 25(OH)D concentration. A significant relationship between maternal reported dietary vitamin D intake and plasma 25(OH)D and 3-epi-25(OH)D3 concentration were identified.
- Only 23% of mothers met the adequate intake (AI) recommendations for choline; the number was even lower (10%) in the postpartum period. Consuming eggs and milk during pregnancy increased mothers' likelihood of meeting choline AI recommendations.

Child Outcomes

- Infants with higher birth weights and at 3 months of age had mothers with a higher pre-pregnancy BMI. Gestational weight gain above recommendations was associated with higher infant weight at birth and three months of age as well as more rapid postnatal growth.
- Infants' vitamin D status increases in direct proportion to mothers' vitamin D intake. Given many mothers are below recommended AI for vitamin D, both infants and breastfeeding mothers require vitamin D supplementation.
- Maternal experience of early adversity associates with maternal symptoms of anxiety and depression during the perinatal period as well as externalizing behavioral problems in their 2-year old children. Boys demonstrate greater vulnerability to the indirect effects of maternal adversity via both depression and anxiety.
- Both mothers' perinatal depressive symptoms and mothers' and fathers' co-occurring perinatal depressive symptoms predicted more problematic emotionally reactive, withdrawn, and total internalizing behaviors in their 2- to 3-year old children. Children's aggression, attention problems and total externalizing behaviors were only predicted by mothers' perinatal depressive symptoms.

Mental health outcomes

Drawn from the full APrON cohort, three key papers [30-32] reported on perinatal depression and anxiety symptom prevalence and risk factors and the impact of adverse childhood experiences on maternal prenatal depression and anxiety. Many more papers report on parental mental health in sub-studies, see APrONstudy.ca for details.

Predictors of Depression

Maternal prenatal depression is known to have negative impacts on pregnancy, maternal postnatal mental health and neonatal outcomes [33, 34], and psychosocial predictors of maternal postpartum depression are well known [35, 36]. However, less well known is whether mothers and fathers in partnered, cohabitating couples (e.g. married, common-law) share common predictors and experiences of depressive symptoms in the perinatal period. Further, whether maternal prenatal nutrition impacts postpartum depressive symptoms is less well studied. Thus,

using APrON data, Leung and colleagues [32] reported on the prevalence of depressive symptoms in couples. For partnered mothers' depressive symptoms, using the recommended Edinburgh Depression Scale (EDS) cut-off score of ≥ 10 [37], findings revealed that 15.2% and 11.18% of mothers had significant symptoms prenatally and at 3 months postpartum, respectively. For fathers in couples, the cut-off of $\text{EDS} \geq 9$ was used as per recommendations [38] revealing that 14.0% and 12.1% of fathers had significant symptoms prenatally and at 3 months postpartum, respectively. These rates are consistent with other large samples [39, 40]. Further, 2.3% of couples experienced perinatal depressive symptoms in both mothers and fathers, 78.5% of couples did not experience depressive symptoms in either partner, 9.5% of couples experienced depressive symptoms in mothers only, and 9.8% of couples experienced depressive symptoms in fathers only. Leung and colleagues [30] also determined the predictors of depression at 3 months postpartum from second trimester data. Risk factors for maternal postpartum depressive symptoms included low household income, high prenatal depressive symptoms, low postnatal support from fathers, and high stressful life events. Risk factors for paternal postpartum depressive symptoms included low household income, prenatal depressive symptoms, postpartum stressful life events, and smoking. For couples in which both partners were depressed, risk factors included low income, high maternal depressive symptoms and low prenatal social support from fathers. Postpartum social support was found to be a protective factor associated with decreased risk of maternal and paternal postpartum depressive symptoms.

Additionally, Leung and colleagues [30] also examined nutritional predictors of maternal postpartum depression, focusing on APrON mothers' reports of their multivitamin and mineral intake in each trimester of pregnancy. Almost all women (99%) took some type of micronutrient supplement during the prenatal period, with the most commonly consumed being vitamins B6, B9 (folate), B12, and E, with more than 90% of mothers consuming above recommended dietary

allowances. With respect to predictors of postpartum depression, findings revealed that mothers who experienced postpartum depressive symptoms ($EDS \geq 10$) were statistically more likely to take the supplemental micronutrients selenium and omega-3, compared to mothers who reported no postpartum depressive symptoms ($EDS < 10$). Selenium intake above the recommended dietary allowance appeared to be protective, and other research suggests that this micronutrient has a role in normal brain function and in the pathway between dopamine and the pathophysiology of depression [41]. The supplement taken the least was omega-3; however, 68.5% of women with low depressive symptoms ($EDS < 10$) did not take omega-3, versus 78.0% of women with high depressive symptoms ($EDS \geq 10$), a finding consistent with other research [42]. The mean intakes of other nutrients were more likely to be higher in women with low depressive symptoms than those with high symptoms, although not statistically significant. The authors conclude with a call for more study into the value of selenium and omega-3 in the prevention of postpartum depression.

Associations with adverse childhood experiences

Adverse childhood experiences are defined as a set of exposures to personal abuse, neglect and household dysfunction prior 18 years of age that includes physical, sexual, and emotional abuse, domestic violence, and parental mental illness, substance use, and incarceration [43]. A graded relationship has been repeatedly observed between the number of adversities experienced in childhood and later increased risk for poor health, including depression [44, 45]; however whether such adversities affect perinatal mood more broadly including anxiety was less studied. Felitti and colleagues' [43] classic measure of adverse childhood experiences was administered to APrON mothers, revealing that while 55% reported none or one early adversity, 31% reported one or two adversities and nearly 14% reported three or more adverse childhood experiences. Employing data from each trimester of pregnancy and 3, 6 and 12 months

postpartum, more reported early adversities predicted higher reported symptoms of prenatal and postnatal depression and anxiety [31]. Routine screening for maternal childhood adversities may facilitate early identification and preventative interventions for perinatal mood disorders including depression and anxiety; however, this has been recently debated [46]. In sum, the identified risks and protective factors provide opportunities for prevention of perinatal depression and anxiety in mothers and fathers.

Nutrition outcomes

Gestational weight gain and dietary intake, especially perinatal nutrient intake and status focused on vitamin D and choline were examined in eight key papers [47-54].

Gestational weight gain and dietary intake

Gestational weight gain over or below Health Canada's Gestational Weight Gain Guidelines is associated with an increased risk of adverse outcomes for mother and infant [55]. The APrON cohort was the first to provide evidence that more than half of women gain more than Health Canada's Guidelines, using prospectively collected body weights from across pregnancy [47]. Since then, several studies have been completed with body weight or body composition in pregnancy or postpartum as a main outcome. Analysis of the APrON data by Jarman and colleagues [48] provided a clear picture indicating that approximately 70% of women who become pregnant with a body mass index (BMI) $> 25 \text{ kg/m}^2$ are likely to exceed gestational weight gain guidelines, by about 19 weeks gestation. Women with excessive weight gain also gain higher amounts of body fat and retain higher amounts of fat compared to women who gain within the guidelines [49]. The detailed dietary intake information that was collected showed that women who consumed a healthy diet pattern prior to pregnancy were less likely to develop complications such as gestational hypertension [50]. An individualized approach to supportive dietary counseling that considers pre-pregnancy BMI in addition to the woman's social and

financial context may be key to helping women meet national recommendations for both diet and gestational weigh gain in pregnancy [56].

Nutrient intake and status

Vitamin D is critical for a healthy pregnancy and the major circulating form of vitamin D is 25-hydroxyvitamin D (25(OH)D); thus, the total serum 25(OH)D level is currently considered the best indicator of vitamin D supply [57]. Three studies that included APrON participants examined the role of maternal vitamin D intake during pregnancy [51-53]. Aghajafari and colleagues [52] reported that the median vitamin D intake from diet and supplements was 600 International Units (IU)/day during pregnancy, which was not sufficient to achieve a target circulation of 25(OH)D concentration. When the 3-epimer (3-epi-25(OH)D3) was included in the estimation of vitamin D status, the prevalence of vitamin D insufficiency (<75 nmol/L) was lower (33.0 %) compared to when it was excluded (38.0 %). Vitamin D supplementation (2000 IU/day) was associated with 25(OH)D3 sufficiency [51-53]. A significant relationship between maternal reported dietary vitamin D intake and plasma 25(OH)D and 3-epi-25(OH)D3 concentration were identified.

Choline intake during pregnancy and lactation is critical to the offspring's brain function [58]. Lewis and colleagues [54] reported that the mean choline intake in pregnant and lactating women ranged between 340 (SD 148) in the second trimester and 346 (SD 151) mg/day at 3 months postpartum. Only 23% of mothers met the adequate intake (AI) recommendations for choline; the number was even lower (10%) in the postpartum period [54]. Consuming eggs and milk during pregnancy increased mothers' likelihood of meeting choline AI recommendations. These findings contributed to the European Food Safety Authority (EFSA) dietary reference values for choline guidelines [59]. To conclude, a significant number of women were not meeting recommendations for vitamin D and choline intake during pregnancy and postpartum. Increased

consumption of these nutrients may be necessary for pregnant and lactating women to improve the health and development of their infants.

Child outcomes

The impacts of maternal and infant nutrient intake, maternal BMI and gestational weight gain, and perinatal depression and maternal adverse childhood experiences on child outcomes were examined in four key papers [31, 60-62]. Many more papers will emerge as the cohort matures and data become available.

Maternal BMI, gestational weight gain and vitamin D

Subhan and colleagues [60] described the effects of maternal pre-pregnancy BMI and gestational weight gain on infant anthropometrics at birth and three months and infant growth rates between birth and three months. Other research shows that children who are heavier at birth have an increased risk of high BMI in childhood, being overweight or obese as adults [63, 64], a condition linked increased risk for hypertension, type 2 diabetes and cardiovascular diseases [63, 65]. APrON findings revealed that infants with higher weight at birth and at 3 months of age had mothers with a higher pre-pregnancy BMI. Also gestational weight gain above recommendations was associated with higher infant weight at birth and three months of age as well as more rapid postnatal growth. The authors concluded that clinicians and health care professionals should encourage women to enter pregnancy with a healthy BMI and adhere to the current gestational weight gain recommendations. Aghajafari and colleagues [61] examined the association between APrON mothers' vitamin D intake during breastfeeding with their infants' vitamin D status to determine whether infant supplementation was sufficient. Other research reveals that low vitamin D is associated with increased rates of infections, autoimmunity and allergies [66] and poor bone health in children [67]. Using plasma from a subset of breastfed infants, vitamin D status was measured by liquid chromatography-tandem mass spectrometry and maternal and infants' dietary

data were attained from dietary questionnaires. Controlling for race, season, prenatal maternal vitamin D status, infant birth weight, and daily infant vitamin D supplementation, infants vitamin D status increased in direct proportion to mothers’ vitamin D intake. Moreover, a quarter of mothers were below recommended AI for vitamin D. The authors concluded that to ensure optimal infant vitamin D status, both infants and breastfeeding mothers require vitamin D supplementation.

Perinatal depression and adverse childhood experiences

Two studies addressed impacts of perinatal depression and adverse childhood experiences on children’s behaviour. It is well established that children’s internalizing behavioural problems (e.g. emotional reactivity, anxiety, depression, somatic complaints, and social withdrawal) and externalizing behavioral problems (i.e. inattention, hyperactivity and aggression) predict adult mental health problems [68-70]. As indicated above, adverse childhood experiences increased risk for poor health over the lifespan [71-73]; however intergenerational impacts have also been noted [74, 75] with an incomplete understanding of the mechanism of transmission form one generation to the next [31]. In examining intergenerational impacts, Letourneau and colleagues³¹ found that APrON mothers’ experiences of early adversity were associated with maternal symptoms of anxiety and depression during the perinatal period (discussed earlier) as well as externalizing behavioral problems in their 2-year old children. Indirect associations were observed between maternal adverse childhood experiences and children’s internalizing and externalizing problems via maternal anxiety and depression. Sex differences were also observed with boys demonstrating greater vulnerability to the indirect effects of maternal adversity via both depression and anxiety. The authors concluded that interventions may be targeted to women who have multiple adverse childhood experiences as well as mental health problems to prevent behavioral problems in their children. Letourneau and colleagues [62] further examined perinatal

depression in mothers, as well as fathers or both parents and how symptoms in one or both parents predict children's behavioral problems. Other research has shown that fathers may buffer the impacts of maternal depression on children's outcomes and thus this protective factor is likely undermined when fathers are depressed as well as mothers [34, 76]. Mothers' and fathers' depressive symptoms were measured during pregnancy and three months postpartum, and children's behavior was measured at 24 and 36 months of age. Results revealed that both mothers' perinatal depressive symptoms and mothers' and fathers' co-occurring perinatal depressive symptoms predicted more problematic emotionally reactive, withdrawn, and total internalizing behaviors. In contrast, children's aggression, attention problems and total externalizing behaviors were only predicted by mothers' perinatal depressive symptoms. The authors urge health care providers to consider the whole family, including fathers, when treating maternal symptoms of depression, aligned with review recommendations to also consider fathers [34, 77].

What are the main strengths and weaknesses?

The major strengths of APrON are recruitment and retention of the large community-based cohort during pregnancy, postpartum and early childhood. The APrON study used a detailed array of standardized validated psychological and physiological measures and biospecimens. The prospective data collection and follow-up enabled investigation of a wide range of maternal and paternal health outcomes. The APrON team has been successful in continuously engaging participants, as evidenced by high response rates over the years (see Table 4).

Table 4. *Response Rates for Questionnaires*

Data collection time point	Targeted participants (n)	Participants responded (n)	Response rate (%)
	2189	2145	97.9

Mothers < 14 weeks gestation			
Mothers at 14-26 weeks gestation			
Mothers at 14-26 weeks gestation and those recruited at < 14 weeks gestation	563	475	84.3
Fathers at 14-26 weeks gestation	1325	1315	99.2
Mothers at 27-40 weeks gestation	1838	1830	99.5
Mothers at 3 months postpartum	2051	1811	88.3
Fathers at 3 months postpartum	1287	1150	89.3
Mothers at 6 months postpartum	1727	1523	88.1
Mothers at 1 year postpartum	1747	1305	74.7
Mothers at 2 years postpartum	1809	1299	71.8
Mothers at 3 years postpartum	1648	1282	77.7

The high response rates over time may be due to the recruitment of a relatively low-risk sample as sociodemographic characteristics are skewed toward a high percentage of well-educated, higher income, married participants. Consequently, generalizability of findings to those with different sociodemographic backgrounds should be made with caution. Further, the study likely included participants with healthier lifestyle behaviors or with more positive attitudes towards research, because such characteristics are associated with higher socioeconomic status [78]. Selective attrition bias is likely operating in the cohort, which may lead to underestimation of the effect of parental mental health and lifestyle choices on children's development. While

attrition has been relatively low over time, differences in the sociodemographic characteristics of the continuing participants compared with discontinuing participants should be considered when interpreting the findings. There were also fewer fathers than mothers in the sample; thus, most developmental and behavioral outcomes are based on maternal reports.

Where can I find out more about APrON research?

Information about the APrON study is available at www.apronstudy.ca. APrON data are available through SAGE (Secondary Evidence to Generate Evidence; <https://policywise.com/sage/>). For more information contact Dr. Nicole Letourneau at Nicole.letourneau@ucalgary.ca. Collaboration or data access inquiries will be considered by the APrON Study team.

Profile in a nutshell

- The APrON study is a longitudinal cohort of mothers, fathers and their children investigating maternal, parental, birth, and child development outcomes.
- A total of 2189 women (aged 16 to 44) and 1325 fathers (aged 18 to 52), residing around either Calgary or Edmonton, Alberta, enrolled and completed at least one questionnaire between May 2009 and June 2012. The majority were married, educated, had annual household incomes >\$70 000 and were born in Canada.
- Mothers completed 18 questionnaires and up to 4 clinic visits spanning pregnancy to eight years postpartum and provided access to their medical records as well as three to four blood and two urine samples. To date, 1648 mothers and 1255 fathers remain eligible for follow-up.
- The APrON dataset includes comprehensive maternal and child sociodemographic data, maternal nutrition and psychological data from pregnancy to eight years postpartum, and infant and child neurodevelopmental and behavioural data to eight years of age. Maternal, paternal and infant biosamples for genetics were also collected. Five and 8-year data collection is complete and 12-year data follow-up will start in 2022.

Requests for data and collaboration are welcomed, contact Dr. Nicole Letourneau at nicole.letourneau@ucalgary.ca

Figure Legends

- *Figure 1.* Socio-demographic characteristics of women in APrON, Calgary, Alberta, Canada.
- *Figure 2.* Data collection to 8 years of child age

Contributorship Statement

Nicole Letourneau (NL) conceived of the idea for the paper, oversees all aspects of the APrON cohort, including the collection, cleaning and analysis of data for reviewed studies, and wrote major sections of the paper, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Elena Ali (EA) wrote the initial draft of the paper with support from NL, GG, FA, RB, and DD, approved of the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Fariba Aghajafari (FA) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Rhonda Bell (RB) wrote a key section of the paper, contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published and agreed to be accountable for all aspects of the work.

Andrea J. Deane (AJD) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved of the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Deborah Dewey (DD) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Catherine Field (CF) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Gerald Giesbrecht (GG) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Bonnie Kaplan (BK) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Brenda Leung (BL) contributed to the collection, cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

Henry Ntanda (HN) contributed to the cleaning and analysis of data for reviewed studies, critically reviewed the drafts, approved the final version of the paper to be published, and agreed to be accountable for all aspects of the work.

APrON Study Cohort Participants volunteered time and energy to share their individual and family's experiences with us to help improve the long-term health outcomes for future newborns, mother, fathers, and families.

Competing Interest

None declared.

Funding Statement

This work was supported by the Alberta Children's Hospital Foundation (grant/award number N/A), Alberta Innovates Health Solutions Foundation (grant/award number N/A), Canadian Institutes for Health Research (Operating Grants), Canadian Institutes for Health Research (CIHR

1
2
3 Doctoral Award, grant number 394652), Kids Brain Health Network (grant/award number N/A),
4
5 and the National Centre of Excellence: Allergen (grant/award number N/A).
6

7
8 **Data Sharing Statement**

9 Data may be obtained from a third party and are not publicly available and data are available
10
11 upon reasonable request. Accordingly, data can be collected at the Secondary Analysis to General
12
13 Evidence (SAGE) data repository or by contacting Dr. Nicole Letourneau (the principal
14
15 investigator for the APrON study).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Leung, B.M., et al., *Perinatal nutrition in maternal mental health and child development: Birth of a pregnancy cohort*. Early Hum Dev, 2016. **93**: p. 1-7.
2. Kaplan, B.J., et al., *The Alberta Pregnancy Outcomes and Nutrition (APrON) cohort study: rationale and methods*. Matern Child Nutr, 2014. **10**(1): p. 44-60.
3. Manca, D.P., et al., *The most effective strategy for recruiting a pregnancy cohort: a tale of two cities*. BMC pregnancy and childbirth, 2013. **13**(1): p. 1-7.
4. Kaplan, B.J., et al., *The Alberta Pregnancy Outcomes and Nutrition (APrON) cohort study: rationale and methods*. Maternal & child nutrition, 2014. **10**(1): p. 44-60.
5. Manca, D.P., et al., *The most effective strategy for recruiting a pregnancy cohort: a tale of two cities*. BMC Pregnancy Childbirth, 2013. **13**: p. 75.
6. Leung, B.M., et al., *Comparison of sample characteristics in two pregnancy cohorts: community-based versus population-based recruitment methods*. BMC Med Res Methodol, 2013. **13**: p. 149.
7. Leung, B.M., et al., *Perinatal nutrition in maternal mental health and child development: Birth of a pregnancy cohort*. Early human development, 2016. **93**: p. 1-7.
8. Dzakpasu, S., et al., *The Canadian maternity experiences survey: design and methods*. J Obstet Gynaecol Can, 2008. **30**(3): p. 207-216.
9. Tough, S.C., et al., *Cohort Profile: The All Our Babies pregnancy cohort (AOB)*. Int J Epidemiol, 2017. **46**(5): p. 1389-1390k.
10. England-Mason, G., et al., *Postnatal BPA is associated with increasing executive function difficulties in preschool children*. Pediatric research, 2020: p. 1-8.

11. Liu, J., et al., *Exposure and dietary sources of bisphenol A (BPA) and BPA-alternatives among mothers in the APrON cohort study*. Environment international, 2018. **119**: p. 319-326.

12. Giesbrecht, G.F., et al., *Prenatal bisphenol a exposure and dysregulation of infant hypothalamic-pituitary-adrenal axis function: findings from the APrON cohort study*. Environmental Health, 2017. **16**(1): p. 1-11.

13. Giesbrecht, G.F., et al., *Urinary bisphenol A is associated with dysregulation of HPA-axis function in pregnant women: Findings from the APrON cohort study*. Environmental research, 2016. **151**: p. 689-697.

14. Liu, J., et al., *Interaction of prenatal bisphenols, maternal nutrients, and toxic metal exposures on neurodevelopment of 2-year-olds in the APrON cohort*. Environment International, 2021. **155**: p. 106601.

15. Grohs, M.N., et al., *Prenatal maternal and childhood bisphenol a exposure and brain structure and behavior of young children*. Environmental Health, 2019. **18**(1): p. 1-12.

16. Letourneau, N.L., et al., *Parenting Interacts With Plasticity Genes in Predicting Behavioral Outcomes in Preschoolers*. Canadian Journal of Nursing Research, 2019: p. 0844562119863612.

17. Merrill, S.M., et al., *Associations of peripheral blood DNA methylation and estimated monocyte proportion differences during infancy with toddler attachment style*. Attachment & Human Development, 2021: p. 1-30.

18. Merrill, S.M., et al., *Paternal adverse childhood experiences: Associations with infant DNA methylation*. Developmental Psychobiology, 2021.

19. Letourneau, N., et al., *Prenatal maternal distress and immune cell epigenetic profiles at 3-months of age*. Developmental Psychobiology, 2021.

20. Letourneau, N.L., et al., *Maternal sensitivity and social support protect against childhood atopic dermatitis*. Allergy, Asthma & Clinical Immunology, 2017. **13**(1): p. 1-14.
21. Giesbrecht, G., et al., *Advancing gestation does not attenuate biobehavioural coherence between psychological distress and cortisol*. Biological psychiatry, 2013. **93**(1): p. 45-51.
22. Giesbrecht, G., et al., *The buffering effect of social support on HPA axis function during pregnancy*. Psychosomatic medicine, 2013. **75**(9): p. 856-862.
23. Giesbrecht, G., et al., *Sexually dimorphic adaptations in basal maternal stress physiology during pregnancy and implications for fetal development*. Psychoneuroendocrinology, 2015. **56**: p. 168-178.
24. Giesbrecht, G.F., et al., *The buffering effect of social support on hypothalamic-pituitary-adrenal axis function during pregnancy*. Psychosomatic medicine, 2013. **75**(9): p. 856-862.
25. Thomas-Argyriou, J.C., et al., *The role of HPA-axis function during pregnancy in the intergenerational transmission of maternal adverse childhood experiences to child behavior problems*. Development and Psychopathology, 2020: p. 1-17.
26. Giesbrecht, G.F., et al., *Psychological distress and salivary cortisol covary within persons during pregnancy*. Psychoneuroendocrinology, 2012. **37**(2): p. 270-279.
27. Thomas, J.C., et al., *Biological embedding of perinatal social relationships in infant stress reactivity*. Developmental psychobiology, 2017. **59**(4): p. 425-435.
28. Rash, J.A., et al., *Maternal cortisol during pregnancy is related to infant cardiac vagal control*. Psychoneuroendocrinology, 2015. **54**: p. 78-89.
29. Thomas, J.C., et al., *Developmental origins of infant emotion regulation: Mediation by temperamental negativity and moderation by maternal sensitivity*. Developmental Psychology, 2017. **53**(4): p. 611.

30. Leung, B.M., et al., *Prenatal micronutrient supplementation and postpartum depressive symptoms in a pregnancy cohort*. BMC Pregnancy Childbirth, 2013. **13**: p. 2.

31. Letourneau, N., et al., *Intergenerational transmission of adverse childhood experiences via maternal depression and anxiety and moderation by child sex*. Journal of developmental origins of health and disease, 2019. **10**(1): p. 88-99.

32. Leung, B.M., et al., *Predictors of postpartum depression in partnered mothers and fathers from a longitudinal cohort*. Community mental health journal, 2017. **53**(4): p. 420-431.

33. Field, T., et al., *Comorbid depression and anxiety effects on pregnancy and neonatal outcome*. Infant Behavior and Development, 2010. **33**(1): p. 23-29.

34. Letourneau, N.L., L. Tramonte, and J.D. Willms, *Maternal depression, family functioning and children's longitudinal development*. Journal of pediatric nursing, 2013. **28**(3): p. 223-234.

35. Yim, I.S., et al., *Biological and psychosocial predictors of postpartum depression: systematic review and call for integration*. Annual review of clinical psychology, 2015. **11**: p. 99-137.

36. Guintivano, J., T. Manuck, and S. Meltzer-Brody, *Predictors of postpartum depression: a comprehensive review of the last decade of evidence*. Clinical obstetrics and gynecology, 2018. **61**(3): p. 591.

37. Bergink, V., et al., *Validation of the Edinburgh Depression Scale during pregnancy*. Journal of Psychosomatic Research, 2011. **70**: p. 385-389.

38. Paulson, J.F. and S.D. Bazemore, *Prenatal and Postpartum Depression in Fathers and Its Association With Maternal Depression*. JAMA, 2010. **303**(19): p. 1961-1969.

39. Gavin, N.I., et al., *Perinatal depression: a systematic review of prevalence and incidence*. Obstetrics & Gynecology, 2005. **106**(5 Part 1): p. 1071-1083.

40. Carlberg, M., M. Edhborg, and L. Lindberg, *Paternal perinatal depression assessed by the Edinburgh Postnatal Depression Scale and the Gotland Male Depression Scale: Prevalence and possible risk factors*. American journal of men's health, 2018. **12**(4): p. 720-729.
41. Pasco, J.A., et al., *Dietary selenium and major depression: a nested case-control study*. Complementary therapies in medicine, 2012. **20**(3): p. 119-123.
42. Markhus, M.W., et al., *Low omega-3 index in pregnancy is a possible biological risk factor for postpartum depression*. PloS one, 2013. **8**(7): p. e67617.
43. Felitti, V.J., et al., *Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study*. Am J Prev Med, 1998. **14**(4): p. 245-58.
44. Danese, A., et al., *Adverse childhood experiences and adult risk factors for age-related disease: depression, inflammation, and clustering of metabolic risk markers*. Archives of pediatrics & adolescent medicine, 2009. **163**(12): p. 1135-1143.
45. Hughes, K., et al., *The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis*. Lancet Public Health, 2017. **2**(8): p. e356-e366.
46. Hughes, K., et al., *The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis*. The Lancet Public Health, 2017. **2**(8): p. e356-e366.
47. Begum, F., et al., *Gestational weight gain and early postpartum weight retention in a prospective cohort of Alberta women*. J Obstet Gynaecol Can, 2012. **34**(7): p. 637-47.
48. Jarman, M., et al., *Patterns and trajectories of gestational weight gain: a prospective cohort study*. CMAJ Open, 2016. **4**(2): p. E338-45.

49. Subhan, F.B., et al., *Association of pre-pregnancy BMI and gestational weight gain with fat mass distribution and accretion during pregnancy and early postpartum: a prospective study of Albertan women*. BMJ Open, 2019. **9**(7): p. e026908.

50. Jarman, M., et al., *Dietary Patterns Prior to Pregnancy and Associations with Pregnancy Complications*. Nutrients, 2018. **10**(7).

51. Aghajafari, F., et al., *Plasma 3-Epi-25-Hydroxycholecalciferol Can Alter the Assessment of Vitamin D Status Using the Current Reference Ranges for Pregnant Women and Their Newborns*. J Nutr, 2016. **146**(1): p. 70-5.

52. Aghajafari, F., et al., *The Current Recommended Vitamin D Intake Guideline for Diet and Supplements During Pregnancy Is Not Adequate to Achieve Vitamin D Sufficiency for Most Pregnant Women*. PLoS One, 2016. **11**(7): p. e0157262.

53. Aghajafari, F., et al., *The High Prevalence of Vitamin D Insufficiency in Cord Blood in Calgary, Alberta (APrON-D Study)*. J Obstet Gynaecol Can, 2017. **39**(5): p. 347-353.e1.

54. Lewis, E.D., et al., *Estimation of choline intake from 24 h dietary intake recalls and contribution of egg and milk consumption to intake among pregnant and lactating women in Alberta*. Br J Nutr, 2014. **112**(1): p. 112-21.

55. Government of Canada. *Prenatal Nutrition Guidelines for Health Professionals: Gestational Weight Gain*. 2010 [cited 2020 21 Jan]; Available from: <https://www.canada.ca/en/health-canada/services/canada-food-guide/resources/prenatal-nutrition/eating-well-being-active-towards-healthy-weight-gain-pregnancy-2010.html>.

56. APrON and ENRICH Study Teams, *Adherence to Canada's Food Guide Recommendations during Pregnancy: Nutritional Epidemiology and Public Health*. Curr Dev Nutr, 2017. **1**(7): p. e000356.

57. Aghajafari, F., et al., *Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies*. *Bmj*, 2013. **346**: p. f1169.
58. Wallace, T.C., et al., *Choline: The Neurocognitive Essential Nutrient of Interest to Obstetricians and Gynecologists*. *J Diet Suppl*, 2019: p. 1-20.
59. European Food Safety Authority, *Dietary reference values: EFSA publishes advice on choline*. 2016. p. 1-70.
60. Subhan, F.B., et al., *Higher pre-pregnancy BMI and excessive gestational weight gain are risk factors for rapid weight gain in infants*. *Maternal and child health journal*, 2017. **21**(6): p. 1396-1407.
61. Aghajafari, F., et al., *Both mother and infant require a vitamin D supplement to ensure that infants' vitamin D status meets current guidelines*. *Nutrients*, 2018. **10**(4): p. 429.
62. Letourneau, N., et al., *Maternal and paternal perinatal depressive symptoms associate with 2- and 3-year-old children's behaviour: findings from the APrON longitudinal study*. *BMC Pediatr*, 2019. **19**(1): p. 435.
63. Sommer, A. and G. Twig, *The impact of childhood and adolescent obesity on cardiovascular risk in adulthood: a systematic review*. *Current diabetes reports*, 2018. **18**(10): p. 1-6.
64. Sacco, M., et al., *Birth weight, rapid weight gain in infancy and markers of overweight and obesity in childhood*. *European journal of clinical nutrition*, 2013. **67**(11): p. 1147-1153.
65. Ghoorah, K., et al., *Obesity and cardiovascular outcomes: a review*. *European Heart Journal: Acute Cardiovascular Care*, 2016. **5**(1): p. 77-85.

66. Maillhot, G. and J.H. White, *Vitamin D and immunity in infants and children*. *Nutrients*, 2020. **12**(5): p. 1233.

67. Arman, S., *What are the effects of vitamin D supplementation for term breastfed infants to prevent vitamin D deficiency and improve bone health?-A Cochrane Review summary with commentary*. *Journal of Musculoskeletal & Neuronal Interactions*, 2021. **21**(2): p. 193.

68. Aronen, E.T. and M. Soininen, *Childhood depressive symptoms predict psychiatric problems in young adults*. *The Canadian Journal of Psychiatry*, 2000. **45**(5): p. 465-470.

69. Benjet, C., et al., *Incidence and recurrence of depression from adolescence to early adulthood: A longitudinal follow-up of the mexican adolescent mental health survey*. *Journal of affective disorders*, 2020. **263**: p. 540-546.

70. Copeland, W.E., et al., *Childhood and adolescent psychiatric disorders as predictors of young adult disorders*. *Archives of general psychiatry*, 2009. **66**(7): p. 764-772.

71. Hege, A., et al., *Adverse childhood experiences among adults in North Carolina, USA: influences on risk factors for poor health across the lifespan and intergenerational implications*. *International journal of environmental research and public health*, 2020. **17**(22): p. 8548.

72. Sheffler, J.L., I. Stanley, and N. Sachs-Ericsson, *ACEs and mental health outcomes*, in *Adverse childhood experiences*. 2020, Elsevier. p. 47-69.

73. Vig, K.D., M.M. Paluszec, and G.J. Asmundson, *ACEs and physical health outcomes*, in *Adverse Childhood Experiences*. 2020, Elsevier. p. 71-90.

74. McDonald, S., et al., *Maternal adverse childhood experiences, mental health, and child behaviour at age 3: The all our families community cohort study*. *Preventive medicine*, 2019. **118**: p. 286-294.

75. Racine, N., et al., *Maternal adverse childhood experiences and infant development*. Pediatrics, 2018. **141**(4).
76. Letourneau, N., L. Duffett-Leger, and M. Salmani, *The role of paternal support in the behavioural development of children exposed to postpartum depression*. Canadian Journal of Nursing Research, 2009. **41**(3): p. 86-106.
77. Rao, W.-W., et al., *Prevalence of prenatal and postpartum depression in fathers: A comprehensive meta-analysis of observational surveys*. Journal of affective disorders, 2020. **263**: p. 491-499.
78. Daniels, J.L., et al., *Attitudes toward participation in a pregnancy and child cohort study*. Paediatr Perinat Epidemiol, 2006. **20**(3): p. 260-6.




Figure 1. Sociodemographic characteristics of women in APrON, Calgary, Alberta, and Canada.

Sociodemographic characteristics of women in APrON, Calgary, Alberta, and Canada
215x279mm (200 x 200 DPI)




Figure 2. Data collection to 8 years of child age

Data collection to 8 years of child age

215x279mm (96 x 96 DPI)