

BMJ Open

Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women's accounts of their birth choices

Journal:	BMJ Open
Manuscript ID:	bmjopen-2015-008881
Article Type:	Research
Date Submitted by the Author:	24-May-2015
Complete List of Authors:	Black, Mairead; University of Aberdeen, Division of Applied Health Sciences Entwistle, Vikki; University of Aberdeen, Health Services Research Unit Bhattacharya, Siladitya; University of Aberdeen, Division of Applied Health Sciences Gillies, Katie; University of Aberdeen, Health Services Research Unit
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Health informatics
Keywords:	OBSTETRICS, PUBLIC HEALTH, PERINATOLOGY, Maternal medicine < OBSTETRICS

SCHOLARONE™
Manuscripts

TITLE:

Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women’s accounts of their birth choices

Mairead Black, Vikki A Entwistle, Siladitya Bhattacharya, Katie Gillies

Mairead Black (corresponding author), Division of Applied Health Sciences, University of Aberdeen, Aberdeen Maternity Hospital, Aberdeen, UK, AB252ZD. Email: Mairead.black@abdn.ac.uk, Tel: 01224438420

Research fellow

Vikki Entwistle, Division of Applied Health Sciences, University of Aberdeen, Aberdeen, UK, AB252ZD, Professor

Siladitya Bhattacharya, Division of Applied Health Sciences, University of Aberdeen, Aberdeen, UK Professor

Katie Gillies, Division of Applied Health Sciences, University of Aberdeen, Health Sciences Building, Aberdeen, UK, AB252ZD, Research fellow

Keywords: caesarean section, vaginal birth after caesarean section, decision-making

Word count: 4301

ABSTRACT

Objective: To identify what women report influences their preferred mode of birth after caesarean section.

Design: Systematic review of qualitative literature using meta-ethnography.

Data sources: Medline, EMBASE, ASSIA, CINAHL and PsycINFO (1996 until April 2013). Hand-searched journals, reference lists and abstract authors.

Study selection: Primary qualitative studies reporting women's accounts of what influenced their preferred mode of birth after caesarean section.

Data extraction and synthesis: Both primary data (quotations from study participants) and authors' interpretations of these were extracted, compared and contrasted between studies, and grouped into themes to support the development of a 'line of argument' synthesis.

Results: Twenty papers reporting the views of 507 women from four countries were included. Distinctive clusters of influences were identified for each of three groups of women. Women who confidently sought vaginal birth after a caesarean section were typically driven by a longstanding anticipation of vaginal birth. Women who sought a repeat caesarean section were strongly influenced by distressing previous birth experiences, and at times, by encouragement from social contacts. Women who were more open to information and professional guidance had fewer strong preconceptions and concerns and viewed a range of considerations as potentially important.

Conclusions: Women's attitudes towards birth after CS appear to be shaped by distinct clusters of influences, suggesting that opportunities exist for clinicians to stratify and personalise decision support by addressing relevant ideas, concerns and experiences from the first CS delivery onwards.

STRENGTHS AND LIMITATIONS OF THE STUDY

- Meta-ethnographic methods ensured sensitivity to contextual factors surrounding the influences reported by women planning birth after CS
- The contextual factors considered included the circumstances under which women were recruited and interviewed, and the timing of the interventions or exposures that influenced their views
- The iterative process of reciprocal translation of study findings facilitated a higher level of understanding than previous mixed-method review methodology has allowed
- The focus on women’s perspectives is consistent with woman-centred approaches to care, but this review did not consider the views of health professionals and family
- The identification of clustering of influences was robust to ‘testing back the fit’ which confirmed that primary authors’ interpretations supported the synthesis ‘line of argument’.

INTRODUCTION

Caesarean section (CS) deliveries are described as being at epidemic levels across middle-and high-income countries.^{1,2} One in three babies in the US are delivered by CS.¹ South American CS rates exceed 50% in many areas, with over 70% of births in private healthcare settings being by CS.^{3,4} Concern to reduce overall CS rates is in tension with efforts to promote patient choice as women themselves often request this mode of birth.⁵

The greatest contribution to current high CS rates comes from repeat CS procedures.⁶ Worldwide rates of vaginal birth after CS (VBAC) have dropped dramatically in recent years. Between 1999 and 2002, US VBAC attempts fell from 48.3% in 2000 to 30.7% in 2002, with 73.4% of VBAC attempts being successful.⁷ The UK saw actual VBAC rates fall from 45.9% in 1988 to 36% between 2004 and 2011.^{8,9} Health service support for VBAC diminished after retrospective data published in 1996 favoured the maternal safety profile of repeat CS.¹⁰ Although more evidence for the relative safety of VBAC has emerged in recent years,¹¹ and efforts have been made to increase VBAC attempts, rates have never fully recovered.^{12,13}

Enthusiasm to reduce CS rates stems from policy concerns about both the relatively high financial costs and the greater maternal morbidity and mortality of CS when compared to vaginal birth.¹⁴ It can also be linked to broader concerns about unnecessary medical intervention (too much medicine).¹⁵ However, the costs and harms that are evident when CS is considered at a population level are much less apparent at the level of individual women. Absolute rates of serious morbidity from CS are low^{2,16} and there is little evidence that women themselves regret CS deliveries when they have requested this mode of birth.¹⁷ At the same time, potential benefits of CS delivery can often be identified for (and by) individual women.¹⁸ Population data suggest that increasing CS rates are not contributing to parallel improvements in neonatal outcomes.¹⁹

Broad policy consensus in high-income countries supports offering women who become pregnant after CS a choice between repeat CS and attempting VBAC unless clinical circumstances or available services preclude this (e.g. when a high risk of CS scar rupture contraindicates VBAC).^{12,16,20} UK guidance outlines which risks (including probabilities) should be discussed by women and health professionals before agreeing on the planned mode of birth by 36 weeks gestation.²⁰ Although probabilistic information about the physical health outcomes of VBAC and repeat CS might seem to support VBAC, the introduction of decision support interventions in the latter part of pregnancy after CS has made little difference to women's choices.^{21,22} There are several plausible explanations for this, including the likelihood that decision-making is influenced by a much broader range of cultural values and social and emotional considerations than are addressed through existing decision support. It is known, for example, that some women have a strong desire to experience vaginal birth,^{23,24} and that some fear dissatisfaction if they choose VBAC but their attempt fails.²⁵⁻²⁷ However, the insights that have emerged from studies to date have been somewhat fragmented. A more comprehensive and nuanced understanding of the complex range of influences on women's decisions is needed to support informed ethical judgements about efforts either to reduce CS rates or to support women's decision-making. Development of public health policy and clinical practice would benefit from as robust as possible an understanding of the diverse perspectives that women bring to decisions about modes of delivery following a previous caesarean, as would debate about what range of options, information, advice and decision support could be appropriately provided by health services. In order to address this need, we aimed to identify, contextualise and synthesise understanding of the reasons why women prefer VBAC or elective repeat CS (ERCS).

METHODS

A systematic literature search and meta-ethnography was conducted. The seven steps of meta-ethnography described by Noblit and Hare, as listed in Box 1, were followed to synthesise the available primary research studies.²⁸

A systematic search was conducted using Medline, EMBASE, ASSIA, CINAHL and PsycINFO in April 2013 using multiple subject headings and free text key words relating to modes of delivery and to exploration of women’s preferences or choices (full search strategy available from authors on request). Inclusion and exclusion criteria are outlined in table 1.

Table 1. Inclusion and exclusion criteria

Inclusion criteria	
Study population	<ul style="list-style-type: none">Comprised or included an identifiable subgroup of women who have had at least one previous caesarean section
Study design	<ul style="list-style-type: none">Primary research that included and clearly reported a qualitative element
Study findings	<ul style="list-style-type: none">Included accounts of influences on preferred mode of delivery after a previous caesarean section from a woman’s perspectivePrimary data provided relevant to the research question and target population of this synthesis
Language	<ul style="list-style-type: none">Any - no language restrictions applied
Exclusion criteria	
Date of publication	<ul style="list-style-type: none">Studies published before 1996.

Titles, abstracts and, where necessary, full papers were screened for potential eligibility. Inclusion and exclusion criteria were applied to full papers. Authors were contacted when only abstracts were published and studies appeared relevant. Three journals containing the greatest number of relevant studies (British Journal of Midwifery, International Journal of Nursing Practice, and BJOG: an International Journal of Obstetrics and Gynaecology) were hand searched to identify any further relevant papers. High quality translation of two abstracts and one full article was obtained. Quality assessment was performed using the Critical Appraisal Skills Programme checklist for qualitative

studies²⁹ to prompt reflection on study quality, but studies were not excluded on the basis of quality if they contained some qualitative data of value to our research question.

The key characteristics of included studies were extracted and summarised (see Table 2). The studies were initially read individually, in chronological order, and relevant points from both the primary data (first-order constructs) and the study authors' descriptions and interpretations (second-order constructs) were extracted. First-order constructs were all obtained from the 'results' section of each study, while second-order constructs were obtained from both 'results' and 'discussion' sections. All first and second-order constructs were tabulated in the form of primary quotes or exact author interpretations to support the identification of key themes.

Screening and identification of studies, followed by coding of constructs was conducted by two authors (one clinical, one non-clinical) independently with regular meetings to establish agreement. During these meetings, provisional third order constructs (our interpretation of both primary authors' interpretations and primary data) and key themes were identified and further developed in discussion with the remaining authors, who each reviewed a different sample of included studies.

The key interpretive aspect, step five of Noblit and Hare's approach, involved comparing and contrasting the constructs and themes that featured in the different studies in an iterative manner. The findings of each study were interpreted in light of each of the other relevant studies in turn. This allowed for detailed consideration of how study design and context could have shaped study findings (for example which women were included and when they were interviewed in relation to their original CS and/or subsequent delivery). During this process, third order constructs were confirmed and a line-of-argument synthesis developed.

The potential for the clinical background of two authors to influence the findings was recognised from the outset. Their preconceptions were continually challenged and utilised in a constructive manner throughout the synthesis process.

A further relevant study was published in 2014, after our line of argument synthesis had been developed. The findings were considered against our line of argument to test the fit.³⁰

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

The search results are outlined in figure 1. Of 2391 citations obtained, 1174 duplicates were excluded. Screening of 1217 titles and/or abstracts resulted in a further 1092 exclusions for lack of relevance. 71 full papers and two sets of conference proceedings were obtained, and attempts made to contact four authors, of which two were unsuccessful. A total of 57 titles lacked relevant primary data or were published before 1996 and were excluded. 20 papers reporting from 15 primary studies were included.

The focus and key study characteristics for the 20 included papers are outlined in table 2.

The studies were conducted in four countries (UK, USA, China and Australia) and each included between 4 and 170 women, with findings from 507 women in total reported across the papers. Six papers reported on women who planned VBAC, four reported on women who planned ERCS, nine reported on both, and one reported on women who planned ERCS but would have desired VBAC in other circumstances.

Table 2. Characteristics of included studies

ID No	Author	Year	Country	Study Aim	Data collection method	Planned birth method at time of study	Participants (n)	Timing of interview
M1	Ridley ³¹	2002	US	Discover what influences women in the decision to deliver via vaginal birth after CS	Interview (FTF)	VBAC	5	Postnatal (2-4/12)
M2	York ³²	2005	UK	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (FTF)	CS	10	Antenatal (Third trimester)
M3	Liu ²³	2006	China	Investigate the decision factors involved and experience of women who had successful VBAC	Interview (FTF), researcher diary, field notes	VBAC	10	Postnatal (1-2/7)
M4 [†]	Fenwick ¹⁸	2006	Australia	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (T), field notes	CS	49	Pre-pregnancy, antenatal and postnatal (no limits)
M5	Emmett ³³	2006	UK	Explore women's experience of decision-making regarding mode of delivery after having a previous CS	Interview (FTF)	VBAC and CS	21	Postnatal (2-8/12)
M6	Cheung ³⁴	2006	China	Understand Chinese women's perceptions and interpretations of their own CS decision-making, and to investigate how their negotiation with healthcare professionals may be improved	Interview (FTF), field notes	CS	52	Postnatal (1/52 or 8/12)
M7	Meddings ³⁵	2006	UK	Examine the lived experience of women who elected to attempt a vaginal birth following a previous CS delivery	Interview (FTF)*2	VBAC	8	Antenatal (>34/40) and postnatal (~6/52)
M8	Moffat ³⁶	2007	UK	Prospectively explore women's decision making regarding mode of delivery after a previous CS	Consultation observation, patient diaries, interview (FTF)	VBAC and CS	26	Antenatal (from 20/40) and postnatal(6/52)
M9 [†]	Fenwick ³⁷	2007	Australia	Explore childbirth expectations and knowledge of women who had experienced a CS and would prefer a vaginal birth in a subsequent pregnancy	Interview (T)	VBAC	35	Pre-pregnancy, Antenatal and Postnatal (no limits)
M10	Farnworth ³⁸	2007	UK	Identify and describe factors which influence women making a choice regarding mode of delivery after a previous CS delivery in a UK setting, and to identify the role of the obstetrician in this process	Interview (FTF)	VBAC and CS	10	Antenatal (36/40)
M11	Cox ³⁹	2007	UK	Explore issues around the choices between VBAC and elective CS based on the nature and extent of the	Interview (type not clear)	VBAC and CS	7	Postnatal (timing not clear)

				information women actually received when making a decision between elective CS and VBAC, the sources of that information and its importance in terms of the influence it had on their decision				
M12	Farnworth ⁴⁰	2008	UK	Examine the impact of a decision support intervention designed for women choosing mode of delivery after one previous CS	Interview (FTF)	VBAC and CS	18	Antenatal (37/40)
M13 [*]	McGrath ⁴¹	2009(a)	Australia	Explore, from the mother's perspective, the decision-making experience with regards to subsequent birth choice for women who had delivered previously by CS	Interview (FTF)	CS	16	Postnatal (6/52)
M14 [*]	McGrath ⁴²	2009(b)	Australia	Describe the perspective of mothers who underwent elective CS on risks associated with the delivery modes of VBAC and elective CS, and their experience discussing such risks with their health professionals	Interview (FTF)	CS	16	Postnatal (6/52)
M15	Goodall ⁴³	2009	UK	Explore women's perceptions of the role of health professionals in their decision regarding mode of delivery, following previous delivery by CS	Interview (FTF)	VBAC and CS	8	Antenatal (20-40/40)
M16	Frost ⁴⁴	2009	UK	Obtain the views of women on their experiences of decision making about the method of delivery following a previous CS section and the role of decision aids in this process	Interview (FTF)	VBAC and CS	30	Antenatal (37/40), Postnatal (6-8/52)
M17 [*]	Phillips ²⁴	2009	Australia	Explore, from a phenomenological perspective, the reasons motivating women to try for or achieve VBAC	Interview (FTF)	VBAC	4	Postnatal (6/52)
M18 [*]	McGrath ⁴⁵	2010(a)	Australia	Explore, from the mothers' perspective, the process of decision-making about mode of delivery for a subsequent birth after a previous CS	Interview (FTF)	VBAC	4	Postnatal (6/52)
M19	David ⁴⁶ Originates from same study as	2010	Australia	Provide maternity health care providers with an increased understanding of, and insight into, the different information needs of this specific group of maternity care consumers.	Telephone log and field notes	VBAC	170	Antenatal (various gestations)
M20 [*]	McGrath ⁴⁷	2010(b)	Australia	To focus on findings which recorded the frustration of women who valued a vaginal delivery but who delivered by CS	Interview (FTF)	CS	8	Postnatal (6/52)

FTF=face-to-face, T=telephone, M=Manuscript. ^{*}Originates from same study (M4 and M9). ^{*}Originates from same study (M13, M14, M17, M18 and M20)

Quality assessment of the papers is presented in Appendix 1. All papers had a clear statement of study aim which deemed qualitative methods appropriate. Common quality concerns included lack of information on: justification for the theoretical approach; lack of information about women who declined to take part; the interview guide used; and data saturation. Only one paper included a discussion of the potential for the researcher's role to influence the study's findings, although two further papers described involvement of a multi-disciplinary team to perform the data analysis, mitigating the risk of dominance of a single interpretive perspective.

Our initial grouping of first and second order constructs resulted in 40 sub-themes. These were then categorised into six key themes which formed the basis of the synthesis: longstanding anticipation of vaginal birth; previous birth experience; influential people; risk-related information from the media and health services; women's perceived balance of consequences of each birth option; and women's role in decision-making.

Key Themes

The six key themes identified as shaping birth preferences after CS are illustrated with example data in Table 3. Primary study participant quotes are displayed in bold text and primary author interpretations are presented in italics.

Table 3. Key themes of influence on birth preferences after CS with corresponding example data. Primary study participant quotes are displayed in bold text and primary author interpretations are presented in italics.

Theme	Exemplary quote
Longstanding anticipation of vaginal birth	<p>'Right from the start I wanted a natural delivery. All the women in my family just gave birth naturally and so I was very disappointed when it didn't work out that way for the first baby' (M17)</p> <p><i>'Despite their CS they still considered women's bodies were 'designed' to give birth vaginally'. (M9)</i></p> <p><i>'Some of the study cases believed, due to their own notions, that there was only one way to feel like a real mother, ie. experiencing vaginal birth and the delivery pain in person. This was why they chose VBAC'. (M3)</i></p>
Previous birth experience	<p>'If my body can't do it [vaginal birth], why put myself and bub [baby] through all the stress and heartache' (M13)</p> <p><i>'Many of these women also expressed that the CS experience had made them feel powerless and helpless; 'taking away total control' (M9)</i></p>
Influential people	'they [doctors] said you can try normally, but they didn't seem very

	<p>positive that it would work and I think they preferred me to have a caesarean'. (M11)</p> <p><i>'Horror stories' and the knowledge and/or personal experience of friends also worked to reinforce their emerging view that CS was the safest birthing option' (M4)</i></p> <p><i>'..other sources of information were noted as mothers groups and/or playgroups.[where] ..sharing of knowledge 'inspired' them' to pursue VBAC (M19)</i></p>
Risk-related information from the media and health professionals	<p>'I like to gather as much information as I can about things and then make my own decisions from that' (M17)</p> <p><i>'A persistent theme appeared to be the lack of both local written information and professional opinion..this led the women to base their knowledge on a mixture of media, professional and personal sources.' (M2)</i></p> <p><i>'Some women described feeling very sure about their preferred mode of delivery from the beginning of pregnancy and those women generally needed little in the way of decisional support' (M8).</i></p>
Perceived balance of consequences	<p>'supposed to have all that stuff squeezed out and that's not done in a CS but it's probably less risky for the baby'. (M4)</p> <p>'About the biggest thing for me was the success rate. . . . There was more positive than negative. . . . 80% of the women who tried it were able to do it'. (M1)</p> <p><i>'When deciding whether to accept the VBAC or not, in most cases patients would first evaluate the advantages and disadvantages. which included the recovery time after delivery, time of hospitalisation, potential harms to the mother and baby.' (M3)</i></p> <p><i>'women.. considered CS a physical, emotional and lifestyle disruption that was risky and had the potential to cause harm to both mother and baby; separated them from their baby; and interrupted the postnatal period' (M9)</i></p>
Role in decision-making	<p>'I was basically told they would prefer for me to try vaginal delivery but I could have a section if I really wanted' (M8)</p> <p>'I feel every time I go and see the doctor or the midwife they keep talking about elective Caesareans... they keep finding reasons why I'll probably need an elective Caesarean so yeah it feels like choice is lot more limited this time' (M15)</p> <p><i>'The important point is that the mothers who tried for a VBAC were clear and focused in their determination to own the decision-making process' (M1)</i></p>

Patterns of influence: a line of argument

Women approaching a birth after a CS generally have either a clear preference for VBAC, a clear preference for ERCS or a relatively open mind to either option. Although some studies by design included women from only one or two of these categories, looking across the studies and using the process of meta-ethnography, we were able to develop a line of argument to explain how their findings were related. In summary the line of argument is that three distinctive clusters of influences support the three attitudinal positions that women adopt towards mode of delivery after CS. These influences include expectations, experiences, exposure to information from health professionals and others, and, crucially, interpretations of each of these.

The three positions and the distinctive influences on these are summarised in Figure 2 and described below. We note that the influences could be operative from different times, and that some were significant before and around the first CS.

Preferences for vaginal birth

Preferences for vaginal birth could be shaped by influences acting over a period of time, which for some women reached several years, and for many was linked to key events or periods of their lives. Personal ambition to achieve vaginal birth, often predating a first pregnancy, drove many women to pursue VBAC. (M17 and M3) This could act synergistically with unpleasant memories of the initial CS experience, particularly where women had felt a loss of control over that birth, leading women to view VBAC as a potentially life-enriching experience that met their ambitions and avoided further negative emotions. (M1, M19, M9, M3, M8, M14) This impression was often enhanced by inter-pregnancy social interaction with other women who had positive VBAC experiences. (M19) For some, the probability of successful VBAC was pivotal. (M1, M3)

Future considerations played an important role in the shaping of preferences for VBAC. Several women believed that VBAC offered physiological benefits to physical and emotional health of both themselves and their offspring, with particular emphasis on the facilitation of bonding and breastfeeding. (M17, M3) This was a particularly dominant issue among women who experienced breastfeeding difficulties after a previous planned CS, especially in those who had successfully breast fed their babies born vaginally in prior pregnancies. (M3) The social benefits of being able to return to usual family roles and resume driving as soon as possible in the postnatal period were also cited as reasons for preferring to avoid CS particularly within UK study settings. (M7, M8 and M9)

Support, advice or direction from health professionals who favoured VBAC had been influential for some women. Although there were women, particularly in the UK and Australia, who were confident about their right to decide how to plan their birth, (M18, M7, M17, M1) others judged any personal reasons they had in favour of ERCS unimportant or unjustified when considered in the light of medical advice in favour of VBAC.(M8)

Preferences for elective repeat CS

The nature of the previous birth experience was the central theme among women who demonstrated a clear preference to have an ERCS. The previous birth and its outcome could also play a role in these women's safety concerns about VBAC, in their discussions with social contacts and on the degree to which they felt they had a choice to make in the subsequent pregnancy.

A previous emergency CS in labour appeared to lead many women to believe their bodies were incapable of vaginal birth. (M8, M10, M13) Some women sought an ERCS to actively avoid any possibility of a repeat emergency abdominal delivery, (M8, M10, M13) while others feared the possibility of a recurrence of the factors which led to the previous CS. Others opted for ERCS on the grounds that it was a familiar and positive birth experience. (M19, M5, M6)

Safety concerns were described as influential among some women in Australia who wished to avoid VBAC due to fear of the uterine scar '**splitting**', or '**rupturing**' during labour. This feeling dominated their preference for ERCS despite awareness of neonatal breathing problems being more common following this mode of birth. (M4) Some women with a strong preference for VBAC had been influenced, sometimes powerfully, by family, friends and health professionals who recommended ERCS as a safer and more predictable mode of birth than VBAC. (M13, M4)

Ownership of choice, or lack of the same, appeared instrumental in those opting for ERCS in many who perceived that their health professionals would prefer this option, and as such, that VBAC was not available to them. (M15) Others choosing ERCS felt happy to exercise their preference as they had been positively encouraged to opt for the mode of birth that felt right for them. (M5)

Open-minded approach

Women who did not have a firm preference for either VBAC or ERCS appeared to be less strongly influenced by prior expectations about childbirth or by their previous birth experience than those who were more committed to one particular mode of delivery. These women valued and often actively

sought the opinion of health professionals during their pregnancy, processed information on the options available, and put considerable effort into weighing up the attributes of the birth options available. An exception to this involved women who felt overwhelmed by the decision-making responsibility and preferred to follow health professionals' advice. (M19, M8 and M18) Obstetricians, and, at times, midwives, appeared to have particular influence over women who were open to considering either mode of birth, even when women were not actively advised how to deliver but perceived subtle signals that their health professional had a preference. (M11) Some women said their choice should be based upon information alone, rather than the input or opinions of others, recognising that other people are not necessarily impartial. (M17)

Robustness of findings

On 'testing back the fit' of our line of argument, we found that the clusters of influence we identified were consistent with the findings of each of the individual studies, but that none of these studies included a broad enough mix of participants to have enabled the development of this level of understanding. Further 'testing' of the line of argument was made possible by the publication of a new study following completion of this synthesis. Shorten et al analysed written text in which women explained their reasons for choosing either mode of birth after CS. They highlighted the significance of previous birth experience, safety concerns and speed of recovery along with health professionals' preferences in shaping eventual decisions. Although they did not describe a clear distinction between the attitudinal groups, their findings were broadly supportive of the conclusions of this synthesis, with no evidence of confliction or contradiction.³⁰

DISCUSSION

Summary of main findings

This study sought to answer the research question 'What influences women's preferred mode of delivery after previous caesarean section?' We have identified distinctive clusters of influences that tend to underpin the three main positions that pregnant women adopt towards modes of delivery. After an initial CS, women tend to approach childbirth with one of three broad attitudinal positions meaning that they: 1) seek vaginal birth 2) seek repeat caesarean or 3) are open minded to consideration of either mode of birth. These positions reflect thought processes which are likely to evolve from at least as early as the primary CS, with some influential cultural norms in operation well before that time. A strong preference for VBAC appears to be driven by a belief that vaginal birth is 'normal' and has some intrinsic

value. This belief is often accompanied by a keen desire to resume a normal life soon after vaginal delivery. In contrast, a clear preference for elective repeat CS from early in pregnancy can be driven by a previous negative experience of childbirth and a positive emphasis on the predictability of ERCS. Finally, there are women who embark upon their next pregnancy undecided about mode of birth. These women are more open to external influence: they appreciate the benefits of both written information and personalised expert advice, which they use to weigh up what they see as the advantages and disadvantages of their options. The recognition of these clusters of influences according to attitude towards birth from early in the pregnancy after CS is a novel finding made possible by looking across the range of relevant studies. Both historical and contemporary studies have highlighted influences on birth preferences after CS which resonate with those identified in this synthesis, but without identification of attitudinal groups or attention to the multiple influences and the ways these may vary over time.^{27,27,48,49,49} The importance of timing of influence has, however, been highlighted recently by prospective work which found that first-trimester preferences for either ERCS or VBAC persist by early in the third trimester in over 70% of women.⁵⁰

Benefits of a meta-ethnographic approach

Meta-ethnography enabled an interpretation of the available research that incorporated a sensitivity to the contextual factors surrounding the influences reported by specific groups of women planning birth after CS.⁵¹ Contextual factors considered included key time points at which influences took hold, fundamental study characteristics (setting; eligibility criteria; recruitment processes; timing of interviews; healthcare systems) and factors unique to individual women. These contextual considerations limit the likelihood that findings would be generalised inappropriately. The iterative process of reciprocal translation used to build upon emergent themes facilitated a higher level of understanding than previous mixed-method review methodology has allowed.²⁵ The clustering of influences identified within specific attitudinal groups provided clinically relevant insight into the nature of women’s decision-making behaviour. In addition, the identification of clustering was considered robust in light of the ‘testing back the fit’ which confirmed that primary authors’ interpretations supported specific attitudinal clusters.

Women’s perspective

The specific focus upon *women’s* perspectives on what influences birth preferences after CS complements the current focus on joint health care decision-making in which informed patients

contribute to decisions which reflect their beliefs and preferences.⁵² This therefore provides insight which has maximal clinical application in settings where every effort should be made to ensure decisions about mode of birth after CS incorporate women's values and preferences. Given that health professionals have a variable level of input into shaping the eventual mode of birth, it is possible that consideration of health professionals' perspectives may have further developed our understanding of the decision-making process.⁵³ However, women's insights were considered central to achieving the goal of informing future efforts to optimise and support woman-centred planning of birth after CS.

Clinical and research implications

Reflection on current practice

The strength of evidence supporting the first CS birth experience as a key influence upon future birth preferences demands immediate attention. Women should be effectively supported in dealing with the unexpected and potentially traumatic nature of a primary CS. Efforts to promptly address any inaccurate perceptions of their CS birth events, and to provide personally specific information about the risks and benefits of future birth options could be made following the first CS and be reiterated early in the pregnancy after CS. The findings of this synthesis suggest that women's concerns about serious maternal or offspring health risks (beyond those of CS scar rupture) are not important influences on their birth choices after CS. This is of particular interest because information currently provided by health professionals for women planning birth after CS focuses largely on these risks and clinical health considerations.²⁰ Recognition of this mismatch between what women and health professionals prioritise should prompt health professionals to engage in discussion with women which allows identification of their main concerns and places sufficient emphasis on the psychological and social as well as the physical health consequences of modes of birth after CS. The heterogeneity of influences on birth choices after CS demonstrated in this synthesis highlight why approaching all women planning birth after CS with eg. the same decision support tool in the latter part of pregnancy is unlikely to alter their prior attitudinal positions.

Implications for future research and practice

Recognition of the diverse range of influences on, and attitudes towards, birth after CS enables us to understand why decision support interventions have had limited effects on ERCS rates so far,^{21,22} and opens up the possibility of a more targeted approach. We suggest that future interventions should aim to promote positive experiences of informed and shared decision-making, while minimising maternal

and fetal morbidity and avoiding unnecessary healthcare costs. Insights from this synthesis suggest that future strategies should ensure early consideration of both women’s concerns and preferences and their likelihood of achieving good physical birth outcomes. Women may be broadly categorised in early pregnancy after CS as being in favour of either VBAC or ERCS or being open to either option. At the same time, their prognosis for successful VBAC may also be assessed based upon factors such as their age, BMI and indication for previous CS.^{8,54} In order to support high quality decision making and increase VBAC success rates, efforts could be made to ensure design of decision support which reflects women’s prognosis for VBAC success and is sensitive to any early preferences regarding birth mode after CS. The six main prognosis/preference categories are represented in figure 3.

Decision support for women may be delivered via conversations with health professionals, advice and information, including decision aids.⁵⁵ Decision aids provide women with information about options relevant to their health status, while helping them to reflect and draw on their personal values. Previous research has demonstrated that use of some such tools in supporting birth choices after CS improved decision satisfaction but had minimal impact upon VBAC rates.²² The lack of success in increasing VBAC rates may reflect that the tools that were tested were not tailored to women’s early attitudes towards each birth mode, but instead delivered advice according to outcomes which women prioritised. Faced with a choice of surgery and less invasive options, decision-aids have been shown to lead patients to choose conservative or less invasive treatments.⁵⁶

In the context of planning birth after CS, decision aids might usefully be stratified according to predicted VBAC success and responsive to individual women’s early birth mode preferences and priorities. It is likely to be particularly important to engage women who are open-minded (groups E and F on figure 3) and women with a VBAC prognosis which is at odds with their preferred mode of birth (groups B and C on figure 3) by the second trimester, in conversations with health professionals to ensure sufficient time to explore their views and discuss and allow them to consider their options. In such situations, a ‘consider a recommendation’ approach may be warranted – explaining why either ERCS or VBAC is recommended, but leaving sufficient scope and ensuring sufficient support for women to assess and discuss the recommendation before making their own mind up about it.⁵⁷ In those pursuing VBAC despite a poor prognosis for success, there could be a discussion about criteria for conversion to CS and adequate counselling in preparation for the possible psychological impact of such an outcome. Those in whom VBAC prognosis is in keeping with their preferred mode of birth (groups A and D on figure 3) might need less in the way of information, conversation and recommendations from health

professionals, but their needs for information and reassurance about their decisions should not be neglected: balanced written information regarding the risks and benefits of both birth options, and clarification/confirmation of ongoing preferences are still likely to be important. As events unfold during subsequent pregnancies, ongoing communication and decision support for all women would need to be tailored to accommodate new clinical information, concerns and preferences, but a broad pathway identified following the first CS would ensure timely and relevant intervention to address modifiable influences.

CONCLUSIONS

Forming a preference for repeat CS or VBAC is a dynamic process, shaped by many influences which appear to cluster distinctively in the development of strongly held positions. Long-standing expectations of childbirth and perceptions of previous birth experiences appear particularly influential upon VBAC and ERCS preferences respectively. This suggests that early communication to discuss women's prospects for VBAC success and explore and discuss their attitudes towards future births may be valuable, and could perhaps start from as early as the first CS. This might help increase the proportion of women who approach birth after CS with an open mind are more likely to be receptive to written information and the advice of health professionals. Our synthesis has highlighted why current care models involving information provision in pregnancy after CS may not lead to the birth choices which could contribute to a reduction in the unnecessary CS rate. It suggests a need to address women's social and psychological concerns, and not just the currently recommended information, both to support women's autonomy in decision-making and to address public health concerns about rising rates of clinically unnecessary CS.

References

1. World Health Organisation. World health statistics 2013.
http://www.who.int/gho/publications/world_health_statistics/2013/en/. Updated 2013.

2. Stavrou EP, Ford JB, Shand AW, Morris JM, Roberts CL. Epidemiology and trends for caesarean section births in new south wales, australia: A population-based study. *BMC Pregnancy & Childbirth*. 2011;11:8.

3. Almeida S, Bettiol H, Barbieri MA, Silva AA, Ribeiro VS. Significant differences in cesarean section rates between a private and a public hospital in brazil. *Cad Saude Publica*. 2008;24(12):2909-2918.

4. Lumbiganon P, Laopaiboon M, Gulmezoglu AM, et al. Method of delivery and pregnancy outcomes in asia: The WHO global survey on maternal and perinatal health 2007-08. *Lancet*. 2010;375(9713):490-499.

5. Minkoff H, Powderly KR, Chervenak F, McCullough LB. Ethical dimensions of elective primary cesarean delivery. *Obstetrics & Gynecology*. 2004;103(2):387-392.

6. Timofeev J, Reddy UM, Huang CC, Driggers RW, Landy HJ, Laughon SK. Obstetric complications, neonatal morbidity, and indications for cesarean delivery by maternal age. *Obstet Gynecol*. 2013;122(6):1184-1195.

7. Landon MB, Hauth JC, Leveno KJ, et al. Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med*. 2004;351(25):2581-2589.

8. Knight HE, Gurol-Urganci I, van der Meulen JH, et al. Vaginal birth after caesarean section: A cohort study investigating factors associated with its uptake and success. *BJOG*. 2014;121(2):183-192.

9. Paterson CM, Saunders NJ. Mode of delivery after one caesarean section: Audit of current practice in a health region. *BMJ*. 1991;303(6806):818-821.
10. McMahon MJ, Luther ER, Bowes WA, Jr, Olshan AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med*. 1996;335(10):689-695.
11. Dodd JM, Crowther CA, Huertas E, Guise JM, Horey D. Planned elective repeat caesarean section versus planned vaginal birth for women with a previous caesarean birth. *Cochrane Database of Systematic Reviews*. 2004(4). doi: 10.1002/14651858.
12. American College of Obstetricians and Gynecologists. ACOG practice bulletin no. 115: Vaginal birth after previous cesarean delivery. *Obstet Gynecol*. 2010;116(2 Pt 1):450-463.
13. Centres for Disease Control and Prevention. Trends in cesarean birth and vaginal birth after previous cesarean, 1991-99. *National vital statistics reports: from the Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System*. 2002;49(13):15.
14. Main EK, Morton CH, Melsop K, Hopkins D, Giuliani G, Gould JB. Creating a public agenda for maternity safety and quality in cesarean delivery. *Obstet Gynecol*. 2012;120(5):1194-1198.
15. Macdonald H, Loder E. Too much medicine: The challenge of finding common ground. *BMJ*. 2015;350:1163.
16. National Institute for Health and Clinical Excellence. Caesarean section; clinical guideline no. 132. . 2011.
17. Dursun P, Yanik FB, Zeyneloglu HB, Baser E, Kuscu E, Ayhan A. Why women request cesarean section without medical indication? *J Matern Fetal Neonatal Med*. 2011;24(9):1133-1137.

18. Fenwick J, Gamble J, Hauck Y. Reframing birth: A consequence of cesarean section. *J Adv Nurs*. 2006;56(2):121-130.

19. Guise J-M, Eden K, Emeis C, Denman MA, Marshall N, Fu R, Janik R, Nygren P, Walker M, McDonagh M. Vaginal birth after cesarean: New insights. evidence report/technology assessment no.191. . 2010;AHRQ Publication No. 10-E003.

20. Royal College of Obstetricians and Gynaecologists. Birth after previous caesarean birth; greentop guideline. . 2007;45.

21. Shorten A, Shorten B, Keogh J, West S, Morris J. Making choices for childbirth: A randomized controlled trial of a decision-aid for informed birth after cesarean. *Birth*. 2005;32(4):252-261.

22. Montgomery AA, Emmett CL, Fahey T, et al. Two decision aids for mode of delivery among women with previous caesarean section: Randomised controlled trial. *Br Med J*. 2007;334(7607):1305-1309.

23. Liu S, Li H, Lee S. The experiences of multipara who chose to undergo vaginal birth after cesarean [chinese]. *J EVID BASED NURS*. 2006;2(3):241-249.

24. Phillips E, McGrath P, Vaughan G. 'I wanted desperately to have a natural birth': Mothers' insights on vaginal birth after caesarean (VBAC). *Contemporary Nurse*. 2009;34(1):77-84.

25. Eden KB, Hashima JN, Osterweil P, Nygren P, Guise JM. Childbirth preferences after cesarean birth: A review of the evidence. *Birth*. 2004;31(1):49-60.

26. Shorten A, Shorten B. The importance of mode of birth after previous cesarean: Success, satisfaction, and postnatal health. *J Midwifery Womens Health*. 2012;57(2):126-132.

27. Flannagan C, Reid B. Repeat CS or VBAC? A systematic review of the factors influencing pregnant women's decision-making processes. *Evidence Based Midwifery*. 2012;10(3):80-86.
28. Noblit G. HR. *Meta-ethnography: Synthesising qualitative studies*. Sage Publications; 1988.
29. Collaboration for qualitative methodologies. Critical appraisal skills program. www.casp-uk.net. Updated 1998.
30. Shorten A, Shorten B, Kennedy HP. Complexities of choice after prior cesarean: A narrative analysis. *Birth*. 2014;41(2):178-184.
31. Ridley RT, Davis PA, Bright JH, Sinclair D. What influences a woman to choose vaginal birth after cesarean? *J Obstet Gynecol Neonatal Nurs*. 2002;31(6):665-672.
32. York S, Briscoe L, Walkinshaw S, Lavender T. Why women choose to have a repeat caesarean section. *BR J MIDWIFERY*. 2005;13(7):440-445.
33. Emmett CL, Shaw ARG, Montgomery AA, Murphy DJ. Women's experience of decision making about mode of delivery after a previous caesarean section: The role of health professionals and information about health risks. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2006;113(12):1438-1445.
34. Cheung NF, Mander R, Cheng L, Chen VY, Yang X. Caesarean decision-making: Negotiation between chinese women and healthcare professionals. *EVID BASED MIDWIFERY*. 2006;4(1):24-30.
35. Meddings F, Phipps FM, Haith-Cooper M, Haigh J. Vaginal birth after caesarean section (VBAC): Exploring women's perceptions. *J Clin Nurs*. 2007;16(1):160-167.

36. Moffat MA, Bell JS, Porter MA, et al. Decision making about mode of delivery among pregnant women who have previously had a caesarean section: A qualitative study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2007;114(1):86-93.

37. Fenwick J, Gamble J, Hauck Y. Believing in birth - choosing VBAC: The childbirth expectations of a self-selected cohort of australian women. *J Clin Nurs*. 2007;16(8):1561-1570.

38. Farnsworth A, Pearson PH. Choosing mode of delivery after previous caesarean birth. *BR J MIDWIFERY*. 2007;15(4):188.

39. Cox B. Hot topic. women's perceptions of their access to, and value of, information as part of their decision making on mode of birth following a previous caesarean section delivery. *MIDIRS MIDWIFERY DIGEST*. 2007;17(2):159-168.

40. Farnworth A, Robson SC, Thomson RG, Watson DB, Murtagh MJ. Decision support for women choosing mode of delivery after a previous caesarean section: A developmental study. *Patient Educ Couns*. 2008;71(1):116-124.

41. McGrath P, Ray-Barruel G. The easy option? australian findings on mothers' perception of elective caesarean as a birth choice after a prior caesarean section. *Int J Nurs Pract*. 2009;15(4):271-279.

42. McGrath P, Phillips E, Ray-Barruel G. Bioethics and birth: Insights on risk decision-making for an elective caesarean after a prior caesarean delivery. *Monash Bioethics Review*. 2009;28(3):22.1-2219.

43. Goodall KE, McVittie C, Magill M. Birth choice following primary caesarean section: Mothers' perceptions of the influence of health professionals on decision-making. *J REPROD INFANT PSYCHOL*. 2009;27(1):4-14.

44. Frost J, Shaw A, Ontgomery A, Murphy D. Women's views on the use of decision aids for decision making about the method of delivery following a previous caesarean section: Qualitative interview study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2009;116(7):896-905.
45. McGrath P, Phillips E, Vaughan G. Vaginal birth after caesarean risk decision-making: Australian findings on the mothers' perspective. *Int J Nurs Pract*. 2010;16(3):274-281.
46. David S, Fenwick J, Bayes S, Martin T. A qualitative analysis of the content of telephone calls made by women to a dedicated 'next birth after caesarean' antenatal clinic. *Women & Birth: Journal of the Australian College of Midwives*. 2010;23(4):166-171.
47. McGrath P, Phillips E, Vaughan G. Speaking out! qualitative insights on the experience of mothers who wanted a vaginal birth after a birth by cesarean section. *The Patient: Patient-Centered Outcomes Research*. 2010;3(1):25-32.
48. McClain CS. Patient decision making: The case of delivery method after a previous cesarean section. *Cult Med Psychiatry*. 1987;11(4):495-508.
49. Lundgren I, Begley C, Gross MM, Bondas T. 'Groping through the fog': A metasynthesis of women's experiences on VBAC (vaginal birth after caesarean section). *BMC Pregnancy and Childbirth*. 2012;12.
50. Shorten A, Shorten B. Timing the provision of a pregnancy decision-aid: Temporal patterns of preference for mode of birth during pregnancy. *Patient Educ Couns*. 2014;97(1):108-113.
51. Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: A critical review. *BMC Med Res Methodol*. 2009;9:59-2288-9-59.

52. Dy SM, Purnell TS. Key concepts relevant to quality of complex and shared decision-making in health care: A literature review. *Soc Sci Med*. 2012;74(4):582-587.

53. Kamal P, Dixon-Woods M, Kurinczuk JJ, Oppenheimer C, Squire P, Waugh J. Factors influencing repeat caesarean section: Qualitative exploratory study of obstetricians' and midwives' accounts. *BJOG : an international journal of obstetrics and gynaecology*. 2005;112(8):1054-1060.

54. Grobman WA, Lai Y, Landon MB, et al. Development of a nomogram for prediction of vaginal birth after cesarean delivery. *Obstet Gynecol*. 2007;109(4):806-812.

55. Say R, Murtagh M, Thomson R. Patients' preference for involvement in medical decision making: A narrative review. *Patient Educ Couns*. 2006;60(2):102-114.

56. Knops AM, Legemate DA, Goossens A, Bossuyt PM, Ubbink DT. Decision aids for patients facing a surgical treatment decision: A systematic review and meta-analysis. *Ann Surg*. 2013;257(5):860-866.

57. Entwistle VA, Carter SM, Trevena L, et al. Communicating about screening. *BMJ*. 2008;337:a1591.

Acknowledgements: None.

Contributorship: MB and SB conceived the idea of the study. MB, SB and KG planned the study. MB and KG conducted the literature search and analysed all data. VE and SB contributed to data analysis and interpretation. MB wrote all drafts of the manuscript and is guarantor. VE, SB and KG contributed to all drafts of the manuscript.

Funding: Mairead Black is a research training fellow funded by The Wellcome Trust. The funder played no role in the design and conduct of the study, in the collection, analysis and interpretation of the data, or in the preparation, review, or approval of the manuscript.

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, [a worldwide licence](http://creativecommons.org/licenses/by/4.0/) to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints,

1
2
3 *include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create*
4 *any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the*
5 *Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever*
6 *it may be located; and, vi) licence any third party to do any or all of the above.”*
7
8

9
10 Ethical approval: Not required.

11
12 Transparency: The lead author (the manuscript’s guarantor) affirms that the manuscript is an honest,
13 accurate, and transparent account of the study being reported; no important aspects of the study have
14 been omitted.
15

16
17 Data sharing: All data analysed is available in the published literature.

18
19 All authors had full access to all of the data in the study and can take responsibility for the integrity of
20 the data and the accuracy of the data analysis.
21

22
23
24 Figure 1. ‘Box 1. Meta-ethnography steps’
25

26
27 Figure 2. ‘Figure 1. Flow diagram of search results’
28


29
30 Figure 3. ‘Figure 2. Summary attitudinal positions of women early in the pregnancy after CS and key
31 influences acting upon their eventual birth preferences.’
32

33
34 Figure 4. ‘Figure 3. Table represents how women may be categorised according to their preferred mode
35 of birth in early pregnancy and their prognosis for VBAC success’
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1. Identify the research question
- 2. Identify relevant studies
- 3. Read the studies
- 4. Identify themes
- 5. Translate the findings of each study into those of the others
- 6. Synthesise the findings
- 7. Express the synthesis

209x297mm (300 x 300 DPI)



209x297mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sought Vaginal birth	Sought repeat CS	Open minded
<ul style="list-style-type: none">•Longstanding anticipation of vaginal birth•Previous (negative) birth experience•Perceived balance of consequences (social considerations about role resumption predominating)•Role in decision-making•Influential people	<ul style="list-style-type: none">•Previous (negative or positive) birth experience•Influential people (friends and health professionals)•Role in decision-making•Risk-related information (media)•Perceived balance of consequences	<ul style="list-style-type: none">•Influential people (health professionals)•Risk-related information (provided by health professionals)•Perceived balance of consequences (mix of health and social considerations predominating)•Longstanding anticipation of vaginal birth•Role in decision-making

209x297mm (300 x 300 DPI)

	Good VBAC prognosis	Poor VBAC prognosis
Prefers VBAC	A	B
Prefers ERCS	C	D
Open minded	E	F

209x297mm (300 x 300 DPI)

Author	Clear study aim	Qualitative methods appropriate	Design appropriate	Recruitment strategy appropriate	Data collection process appropriate	Researcher role in forming research questions discussed	Ethical issues considered	In-depth description of analysis	Adequate discussion of the evidence	Contribution to existing knowledge discussed
Ridley ³¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
York ³²	Yes	Yes	Yes	N/C	Yes	N/C	Yes	No	Yes	Yes
Liu ²³	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Fenwick ¹⁸	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Emmett ³³	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Cheung ³⁴	Yes	Yes	Yes	No – snowball*	Yes	N/C	Yes	Yes	Yes	Yes
Meddings ³⁵	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Moffat ³⁶	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Fenwick ³⁷	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Farnworth ³⁸	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Cox ³⁹	Yes	Yes	Yes	Yes	Yes	N/C	Yes	No	Yes	Yes
Farnworth ⁴⁰	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴¹	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴²	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
Goodall ⁴³	Yes	Yes	Yes	No	Yes	N/C	Yes	Yes	Yes	Yes
Frost ⁴⁴	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Phillips ²⁴	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴⁵	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
David ⁴⁶	Yes	Yes	Yes	Yes	Yes	N/C	N/C	Yes	Yes	Yes
McGrath ⁴⁷	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes

Appendix Table 1. Critical Appraisal Skills Programme Assessment of Included Studies

N/C=Not clear *judged inappropriate to meet the specific study aim

Enhancing transparency in reporting the synthesis of qualitative research: the ENTREQ statement

No	Item	Guide and description	Met
1	Aim	State the research question the synthesis addresses.	✓
2	Synthesis methodology	Identify the synthesis methodology or theoretical framework which underpins the synthesis, and describe the rationale for choice of methodology (e.g. <i>meta-ethnography, thematic synthesis, critical interpretive synthesis, grounded theory synthesis, realist synthesis, meta-aggregation, meta-study, framework synthesis</i>).	✓
3	Approach to searching	Indicate whether the search was pre-planned (<i>comprehensive search strategies to seek all available studies</i>) or iterative (<i>to seek all available concepts until they theoretical saturation is achieved</i>).	✓
4	Inclusion criteria	Specify the inclusion/exclusion criteria (e.g. <i>in terms of population, language, year limits, type of publication, study type</i>).	✓
5	Data sources	Describe the information sources used (e.g. <i>electronic databases (MEDLINE, EMBASE, CINAHL, psycINFO, Econlit), grey literature databases (digital thesis, policy reports), relevant organisational websites, experts, information specialists, generic web searches (Google Scholar) hand searching, reference lists</i>) and when the searches conducted; provide the rationale for using the data sources.	✓
6	Electronic Search strategy	Describe the literature search (e.g. <i>provide electronic search strategies with population terms, clinical or health topic terms, experiential or social phenomena related terms, filters for qualitative research, and search limits</i>).	✓
7	Study screening methods	Describe the process of study screening and sifting (e.g. <i>title, abstract and full text review, number of independent reviewers who screened studies</i>).	✓
8	Study characteristics	Present the characteristics of the included studies (e.g. <i>year of publication, country, population, number of participants, data collection, methodology, analysis, research questions</i>).	✓
9	Study selection results	Identify the number of studies screened and provide reasons for study exclusion (e.g. <i>for comprehensive searching, provide numbers of studies screened and reasons for exclusion indicated in a figure/flowchart; for iterative searching describe reasons for study exclusion and inclusion based on modifications to the research question and/or contribution to theory development</i>).	✓
10	Rationale for appraisal	Describe the rationale and approach used to appraise the included studies or selected findings (e.g. <i>assessment of conduct (validity and robustness), assessment of reporting (transparency), assessment of content and utility of the findings</i>).	✓
11	Appraisal items	State the tools, frameworks and criteria used to appraise the studies or selected findings (e.g. <i>Existing tools: CASP, QARI, COREQ, Mays and Pope [25]; reviewer developed tools; describe the domains assessed: research team, study design, data analysis and interpretations, reporting</i>).	✓
12	Appraisal process	Indicate whether the appraisal was conducted independently by more than one reviewer and if consensus was required.	-
13	Appraisal results	Present results of the quality assessment and indicate which articles, if any, were weighted/excluded based on the assessment and give the rationale.	✓
14	Data extraction	Indicate which sections of the primary studies were analysed and how were the data extracted from the primary studies? (e.g. <i>all text under the headings "results /conclusions" were extracted electronically and entered into a computer software</i>).	✓
15	Software	State the computer software used, if any.	-
16	Number of reviewers	Identify who was involved in coding and analysis.	In part
17	Coding	Describe the process for coding of data (e.g. <i>line by line coding to search for concepts</i>).	✓
18	Study comparison	Describe how were comparisons made within and across studies (e.g. <i>subsequent studies were coded into pre-existing concepts, and new concepts were created when deemed necessary</i>).	✓
19	Derivation of themes	Explain whether the process of deriving the themes or constructs was inductive or deductive.	✓

20	Quotations	Provide quotations from the primary studies to illustrate themes/constructs, and identify whether the quotations were participant quotations of the author's interpretation.	✓
21	Synthesis output	Present rich, compelling and useful results that go beyond a summary of the primary studies (e.g. <i>new interpretation, models of evidence, conceptual models, analytical framework, development of a new theory or construct</i>).	✓

For peer review only

BMJ Open

Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women's accounts of their birth choices

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-008881.R1
Article Type:	Research
Date Submitted by the Author:	01-Oct-2015
Complete List of Authors:	Black, Mairead; University of Aberdeen, Division of Applied Health Sciences Entwistle, Vikki; University of Aberdeen, Health Services Research Unit Bhattacharya, Siladitya; University of Aberdeen, Division of Applied Health Sciences Gillies, Katie; University of Aberdeen, Health Services Research Unit
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Qualitative research
Keywords:	OBSTETRICS, PUBLIC HEALTH, PERINATOLOGY, Maternal medicine < OBSTETRICS

SCHOLARONE™
Manuscripts

TITLE:

Vaginal birth after caesarean section: why is uptake so low? Insights from a meta-ethnographic synthesis of women’s accounts of their birth choices

Mairead Black, Vikki A Entwistle, Siladitya Bhattacharya, Katie Gillies

Mairead Black (corresponding author), Division of Applied Health Sciences, University of Aberdeen, Aberdeen Maternity Hospital, Aberdeen, UK, AB252ZD. Email: Mairead.black@abdn.ac.uk, Tel: 01224438420

Research fellow

Vikki Entwistle, Division of Applied Health Sciences, University of Aberdeen, Aberdeen, UK, AB252ZD, Professor

Siladitya Bhattacharya, Division of Applied Health Sciences, University of Aberdeen, Aberdeen, UK Professor

Katie Gillies, Division of Applied Health Sciences, University of Aberdeen, Health Sciences Building, Aberdeen, UK, AB252ZD, Research fellow

Keywords: caesarean section, vaginal birth after caesarean section, decision-making

Word count: 5159

ABSTRACT

Objective: To identify what women report influences their preferred mode of birth after caesarean section.

Design: Systematic review of qualitative literature using meta-ethnography.

Data sources: Medline, EMBASE, ASSIA, CINAHL and PsycINFO (1996 until April 2013 –updated September 2015). Hand-searched journals, reference lists and abstract authors.

Study selection: Primary qualitative studies reporting women's accounts of what influenced their preferred mode of birth after caesarean section.

Data extraction and synthesis: Both primary data (quotations from study participants) and authors' interpretations of these were extracted, compared and contrasted between studies, and grouped into themes to support the development of a 'line of argument' synthesis.

Results: Twenty papers reporting the views of 507 women from four countries were included. Distinctive clusters of influences were identified for each of three groups of women. Women who confidently sought vaginal birth after a caesarean section were typically driven by a longstanding anticipation of vaginal birth. Women who sought a repeat caesarean section were strongly influenced by distressing previous birth experiences, and at times, by encouragement from social contacts. Women who were more open to information and professional guidance had fewer strong preconceptions and concerns and viewed a range of considerations as potentially important.

Conclusions: Women's attitudes towards birth after CS appear to be shaped by distinct clusters of influences, suggesting that opportunities exist for clinicians to stratify and personalise decision support by addressing relevant ideas, concerns and experiences from the first CS birth onwards.

STRENGTHS AND LIMITATIONS OF THE STUDY

- Meta-ethnographic methods ensured sensitivity to contextual factors surrounding the influences reported by women planning birth after CS
- The contextual factors considered included the circumstances under which women were recruited and interviewed, and the timing of the interventions or exposures that influenced their views
- The iterative process of reciprocal translation of study findings facilitated a higher level of understanding than previous mixed-method review methodology has allowed
- The focus on women’s perspectives is consistent with woman-centred approaches to care, but this review did not consider the views of health professionals and family
- The identification of clustering of influences was robust to ‘testing back the fit’ which confirmed that primary authors’ interpretations supported the synthesis ‘line of argument’.

INTRODUCTION

Caesarean section (CS) births are described as being at epidemic levels across middle-and high-income countries.^{1,2} One in three babies in the US are born by CS.¹ South American CS rates exceed 50% in many areas, with over 70% of births in private healthcare settings being by CS.^{3,4} Concern to reduce overall CS rates is in tension with efforts to promote patient choice as women themselves often request this mode of birth.⁵

The greatest contribution to current high CS rates comes from repeat CS procedures.⁶ Worldwide rates of vaginal birth after CS (VBAC) have dropped dramatically in recent years. Between 1999 and 2002, US VBAC attempts fell from 48.3% in 2000 to 30.7% in 2002, with 73.4% of VBAC attempts being successful.⁷ The UK saw actual VBAC rates fall from 45.9% in 1988 to 36% between 2004 and 2011.^{8,9} Health service support for VBAC diminished after retrospective data published in 1996 favoured the maternal safety profile of repeat CS.¹⁰ Although more evidence for the relative safety of VBAC has emerged in recent years,¹¹ and efforts have been made to increase VBAC attempts, rates have never fully recovered.^{12,13}

Enthusiasm to reduce CS rates stems from policy concerns about both the relatively high financial costs and the greater maternal morbidity and mortality of CS when compared to vaginal birth.¹⁴ It can also be linked to broader concerns about unnecessary medical intervention (too much medicine).¹⁵ However, the costs and harms that are evident when CS is considered at a population level are much less apparent at the level of individual women. Absolute rates of serious morbidity from CS are low^{2,16} and there is little evidence that women themselves regret CS when they have requested this mode of birth.¹⁷ At the same time, potential benefits of CS can often be identified for (and by) individual women.¹⁸ Population data suggest that increasing CS rates are not contributing to parallel improvements in neonatal outcomes.¹⁹

Broad policy consensus in high-income countries supports offering women who become pregnant after CS a choice between repeat CS and attempting VBAC unless clinical circumstances or available services preclude this (e.g. when a high risk of CS scar rupture contraindicates VBAC).^{12,16,20} UK guidance outlines which risks (including probabilities) should be discussed by women and health professionals before agreeing on the planned mode of birth by 36 weeks gestation.²⁰ Although probabilistic information about the physical health outcomes of VBAC and repeat CS might seem to support VBAC, the introduction of decision support interventions in the latter part of pregnancy after CS has made little difference to women's choices.^{21,22} There are several plausible explanations for this, including the likelihood that decision-making is influenced by a much broader range of cultural values and social and emotional considerations than are addressed through existing decision support. It is known, for example, that some women have a strong desire to experience vaginal birth,^{23,24} and that some fear dissatisfaction if they choose VBAC but their attempt fails.²⁵⁻²⁷ However, the insights that have emerged from studies to date have been somewhat fragmented. A more comprehensive and nuanced understanding of the complex range of influences on women's decisions is needed to support informed ethical judgements about efforts either to reduce CS rates or to support women's decision-making. Development of public health policy and clinical practice would benefit from as robust as possible an understanding of the diverse perspectives that women bring to decisions about mode of birth following a previous caesarean, as would debate about what range of options, information, advice and decision support could be appropriately provided by health services. In order to address this need, we aimed to identify, contextualise and synthesise an understanding of the reasons why women prefer VBAC or elective repeat CS (ERCS).

METHODS

A systematic literature search and meta-ethnography was conducted. The seven steps of meta-ethnography described by Noblit and Hare, as listed in Box 1, were followed to synthesise the available primary research studies.²⁸

A systematic search was conducted using Medline, EMBASE, ASSIA, CINAHL and PsycINFO in April 2013 (updated in September 2015) using multiple subject headings and free text key words relating to modes of birth and exploration of women’s preferences or choices (the full search strategy for Medline is provided as appendix 1 and further search strategies are available from the authors on request).

Inclusion and exclusion criteria are outlined in table 1.

Table 1. Inclusion and exclusion criteria

Inclusion criteria	
Study population	<ul style="list-style-type: none">Comprised or included an identifiable subgroup of women who have had at least one previous caesarean section
Study design	<ul style="list-style-type: none">Primary research that included and clearly reported a qualitative element
Study findings	<ul style="list-style-type: none">Included accounts of influences on preferred mode of birth after a previous caesarean section from the women’s perspectivesPrimary data provided relevant to the research question and target population of this synthesis
Language	<ul style="list-style-type: none">Any - no language restrictions applied
Exclusion criteria	
Date of publication	<ul style="list-style-type: none">Studies published before 1996.

Titles, abstracts and, where necessary, full papers were screened for potential eligibility. Inclusion and exclusion criteria were applied to full papers. Authors were contacted when only abstracts were published and studies appeared relevant. Three journals containing the greatest number of relevant studies in the 2013 search (British Journal of Midwifery, International Journal of Nursing Practice, and BJOG: an International Journal of Obstetrics and Gynaecology) were hand searched to identify any further relevant papers. High quality translation of two abstracts and one full article was obtained.

Quality assessment was performed using the Critical Appraisal Skills Programme checklist for qualitative studies²⁹ to prompt reflection on study quality, but studies were not excluded on the basis of quality if they contained some qualitative data of value to our research question.

The key characteristics of included studies were extracted and summarised (see Table 2). The studies were initially read individually, in chronological order, and relevant points from both the primary data (first-order constructs) and the study authors' descriptions and interpretations (second-order constructs) were extracted. First-order constructs were obtained from quotations from women reported in the 'results' section of each study, while second-order constructs (primary authors' account and interpretation of their findings) were obtained from both 'results' and 'discussion' sections. All first and second-order constructs were tabulated in the form of primary quotes or exact author interpretations to support the identification of key themes.

Searching was conducted by one author, with input from an information specialist. Screening and identification of studies, followed by coding of constructs was conducted by two authors {one clinical (MB), one non-clinical (KG)} independently with regular meetings to establish agreement. During these meetings, provisional third order constructs (our interpretation of both primary authors' interpretations and primary data) and key themes were identified. A third and fourth author (VE and SB) were involved in further development of these themes, having each reviewed a different sample of included studies.

The key interpretive aspect, step five of Noblit and Hare's approach, involved one author comparing and contrasting the constructs and themes that featured in the different studies in an iterative manner. The findings of each study were interpreted in light of each of the other relevant studies in turn. This allowed for detailed consideration of how study design and context could have shaped study findings (for example which women were included and when they were interviewed in relation to their original CS and/or subsequent birth). During this process, third order constructs were confirmed and a line-of-argument synthesis developed. All four authors contributed to the development of the line of argument.

The potential for the clinical background of two authors (MB and SB) in particular to influence the findings was recognised from the outset. All team members' interpretations and preconceptions were continually challenged and utilised in a constructive manner during discussions throughout the synthesis process to ensure all reported perspectives were fairly considered and the line of argument developed was robust.

Following the updated search in September 2015, additional eligible papers were identified. Relevant findings were used to test the fit of the line of argument. This involved identification of first and second order constructs (primary data and authors’ interpretations respectively) in the additional papers and analysing these for relevant themes of influence on birth preferences after CS. These themes were compared and contrasted with the content of the line of argument to assess the extent to which they appeared to ‘fit’ together or conflict with one another.³⁰

RESULTS

The search results are outlined in Figure 1. Of 2391 citations obtained in the original search, 1174 duplicates were excluded. Screening of 1217 titles and/or abstracts resulted in a further 1092 exclusions for lack of relevance. 71 full papers and two sets of conference proceedings were obtained, and attempts made to contact four authors, of which two were unsuccessful. A total of 57 titles lacked relevant primary data or were published before 1996 and were excluded. 20 papers reporting from 15 primary studies were included following resolution of disagreement over eligibility of two papers.

The focus and key study characteristics for the 20 included papers are outlined in table 2.

The identified studies were conducted in four countries (UK, USA, China and Australia) and each included between 4 and 170 women, with findings from 507 women in total reported across the papers. Six papers reported on women who planned VBAC, four reported on women who planned ERCS, nine reported on both, and one reported on women who planned ERCS but would have desired VBAC in other circumstances.

Table 2. Characteristics of included studies

ID No	Author	Year	Country	Study Aim	Data collection method	Planned birth method at time of study	Participants (n)	Timing of interview
M1	Ridley ³¹	2002	US	Discover what influences women in the decision to deliver via vaginal birth after CS	Interview (FTF)	VBAC	5	Postnatal (2-4/12)

M2	York ³²	2005	UK	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (FTF)	CS	10	Antenatal (Third trimester)
M3	Liu ²³	2006	China	Investigate the decision factors involved and experience of women who had successful VBAC	Interview (FTF), researcher diary, field notes	VBAC	10	Postnatal (1-2/7)
M4 [¥]	Fenwick ¹⁸	2006	Australia	Describe childbirth expectations, influences and knowledge in women who had experienced emergency CS and planned subsequent CS	Interview (T), field notes	CS	49	Pre-pregnancy, antenatal and postnatal (no limits)
M5	Emmett ³³	2006	UK	Explore women's experience of decision-making regarding mode of delivery after having a previous CS	Interview (FTF)	VBAC and CS	21	Postnatal (2-8/12)
M6	Cheung ³⁴	2006	China	Understand Chinese women's perceptions and interpretations of their own CS decision-making, and to investigate how their negotiation with healthcare professionals may be improved	Interview (FTF), field notes	CS	52	Postnatal (1/52 or 8/12)
M7	Meddings ³⁵	2006	UK	Examine the lived experience of women who elected to attempt a vaginal birth following a previous CS delivery	Interview (FTF)*2	VBAC	8	Antenatal (>34/40) and postnatal (~6/52)
M8	Moffat ³⁶	2007	UK	Prospectively explore women's decision making regarding mode of delivery after a previous CS	Consultation observation, patient diaries, interview (FTF)	VBAC and CS	26	Antenatal (from 20/40) and postnatal(6/52)
M9 [¥]	Fenwick ³⁷	2007	Australia	Explore childbirth expectations and knowledge of women who had experienced a CS and would prefer a vaginal birth in a subsequent pregnancy	Interview (T)	VBAC	35	Pre-pregnancy, Antenatal and Postnatal (no limits)
M10	Farnworth ³⁸	2007	UK	Identify and describe factors which influence women making a choice regarding mode of delivery after a previous CS delivery in a UK setting, and to identify the role of the obstetrician in this process	Interview (FTF)	VBAC and CS	10	Antenatal (36/40)
M11	Cox ³⁹	2007	UK	Explore issues around the choices between VBAC and elective CS based on the nature and extent of the information women actually received when making a decision between elective CS and VBAC, the sources of that information and its importance in terms of the influence it had on their decision	Interview (type not clear)	VBAC and CS	7	Postnatal (timing not clear)

M12	Farnworth ⁴⁰	2008	UK	Examine the impact of a decision support intervention designed for women choosing mode of delivery after one previous CS	Interview (FTF)	VBAC and CS	18	Antenatal (37/40)
M13*	McGrath ⁴¹	2009(a)	Australia	Explore, from the mother's perspective, the decision-making experience with regards to subsequent birth choice for women who had delivered previously by CS	Interview (FTF)	CS	16	Postnatal (6/52)
M14*	McGrath ⁴²	2009(b)	Australia	Describe the perspective of mothers who underwent elective CS on risks associated with the delivery modes of VBAC and elective CS, and their experience discussing such risks with their health professionals	Interview (FTF)	CS	16	Postnatal (6/52)
M15	Goodall ⁴³	2009	UK	Explore women's perceptions of the role of health professionals in their decision regarding mode of delivery, following previous delivery by CS	Interview (FTF)	VBAC and CS	8	Antenatal (20-40/40)
M16	Frost ⁴⁴	2009	UK	Obtain the views of women on their experiences of decision making about the method of delivery following a previous CS section and the role of decision aids in this process	Interview (FTF)	VBAC and CS	30	Antenatal (37/40), Postnatal (6-8/52)
M17*	Phillips ²⁴	2009	Australia	Explore, from a phenomenological perspective, the reasons motivating women to try for or achieve VBAC	Interview (FTF)	VBAC	4	Postnatal (6/52)
M18*	McGrath ⁴⁵	2010(a)	Australia	Explore, from the mothers' perspective, the process of decision-making about mode of delivery for a subsequent birth after a previous CS	Interview (FTF)	VBAC	4	Postnatal (6/52)
M19	David ⁴⁶ Originates from same study as	2010	Australia	Provide maternity health care providers with an increased understanding of, and insight into, the different information needs of this specific group of maternity care consumers.	Telephone log and field notes	VBAC	170	Antenatal (various gestations)
M20*	McGrath ⁴⁷	2010(b)	Australia	To focus on findings which recorded the frustration of women who valued a vaginal delivery but who delivered by CS	Interview (FTF)	CS	8	Postnatal (6/52)

FTF=face-to-face, T=telephone, M=Manuscript, VBAC=vaginal birth after CS, CS=Caesarean section.

*Originates from same study (M4 and M9). *Originates from same study (M13, M14, M17, M18 and M20)

Quality assessment of the papers is presented in Appendix 2. All papers had a clear statement of study aim which deemed qualitative methods appropriate. Common quality concerns included lack of information on: justification for the theoretical approach; lack of information about women who

declined to take part; the interview guide used; and data saturation. Only one paper included a discussion of the potential for the researcher's role to influence the study's findings, although two further papers described involvement of a multi-disciplinary team to perform the data analysis, mitigating the risk of dominance of a single interpretive perspective.

Our initial grouping of first and second order constructs resulted in 40 sub-themes. These were then categorised into six key themes which characterised the main kinds of consideration and features of decision-making processes that appeared to influence preferences for mode of birth. These themes were: longstanding anticipation of vaginal birth; responses to previous birth experiences (positive and/or negative); encouragement or dissuasion from influential people for either birth mode; fear or reassurance from risk-related information on VBAC; perceived net benefit or harm of birth options; and extent and nature of involvement in decision-making. As the labels suggest, several of these themes accommodate a spectrum of views or experiences.

Key Themes

The six key themes identified as shaping birth preferences after CS are illustrated with example data in Table 3. Primary study participant quotes illustrating first order constructs are displayed in bold text and primary author interpretations illustrating second order constructs are presented in italics.

Table 3. Key themes of influence on birth preferences after CS with corresponding example data. Primary study participant quotes are displayed in bold text and primary author interpretations are presented in italics.

Theme	Exemplary quote
Longstanding anticipation of vaginal birth	<p>'Right from the start I wanted a natural delivery. All the women in my family just gave birth naturally and so I was very disappointed when it didn't work out that way for the first baby' (M17)</p> <p><i>'Despite their CS they still considered women's bodies were 'designed' to give birth vaginally'. (M9)</i></p> <p><i>'Some of the study cases believed, due to their own notions, that there was only one way to feel like a real mother, ie. experiencing vaginal birth and the delivery pain in person. This was why they chose VBAC'. (M3)</i></p>
Responses to previous birth experience (positive and/or negative)	<p>'If my body can't do it [vaginal birth], why put myself and bub [baby] through all the stress and heartache' (M13)</p> <p><i>'Many of these women also expressed that the CS experience had made them feel powerless and helpless; 'taking away total control'(M9)</i></p> <p>'In the end we said, look, we're going to go with what we know. What we did first time worked out okay' (M13)</p>

Encouragement or dissuasion from influential people for either birth mode	<p>'they [doctors] said you can try normally, but they didn't seem very positive that it would work and I think they preferred me to have a caesarean'. (M11)</p> <p><i>'Horror stories' and the knowledge and/or personal experience of friends also worked to reinforce their emerging view that CS was the safest birthing option' (M4)</i></p> <p><i>'...other sources of information were noted as mothers groups and/or playgroups.[where] ...sharing of knowledge 'inspired' them' to pursue VBAC (M19)</i></p>
Fear or reassurance from risk-related information on VBAC	<p>'I like to gather as much information as I can about things and then make my own decisions from that' (M17)</p> <p><i>'A persistent theme appeared to be the lack of both local written information and professional opinion... this led the women to base their knowledge on a mixture of media, professional and personal sources.' (M2)</i></p> <p><i>'Some women described feeling very sure about their preferred mode of delivery from the beginning of pregnancy and those women generally needed little in the way of decisional support' (M8).</i></p> <p><i>'Information and support gave women confidence in their decision, and ultimately, the power to own and justify the decision that they had made' (M12)</i></p> <p>'Oh yeah, the riskiest approach was to try a vaginal delivery. Yeah, no I wouldn't even have attempted it. And everything I read backed that up, yes.' (M14)</p>
Perceived net benefit or harm of birth options;	<p>'supposed to have all that stuff squeezed out and that's not done in a CS but it's probably less risky for the baby'. (M4)</p> <p>'About the biggest thing for me was the success rate. . . . There was more positive than negative. . . . 80% of the women who tried it were able to do it'. (M1)</p> <p><i>'When deciding whether to accept the VBAC or not, in most cases patients would first evaluate the advantages and disadvantages. which included the recovery time after delivery, time of hospitalisation, potential harms to the mother and baby.' (M3)</i></p> <p><i>'women... considered CS a physical, emotional and lifestyle disruption that was risky and had the potential to cause harm to both mother and baby; separated them from their baby; and interrupted the postnatal period' (M9)</i></p>
Extent and nature of involvement in decision-making	<p>'I was basically told they would prefer for me to try vaginal delivery but I could have a section if I really wanted' (M8)</p> <p>'I feel every time I go and see the doctor or the midwife they keep talking about elective Caesareans... they keep finding reasons why I'll probably need an elective Caesarean so yeah it feels like choice is lot</p>

more limited this time' (M15)

'The important point is that the mothers who tried for a VBAC were clear and focused in their determination to own the decision-making process'
(M1)

Patterns of influence: a line of argument

We noted that some kinds of views and experiences (specific instances of the six key themes) tended to cluster together in support of the main birth preferences. These clusterings are discussed here in the context of the line of argument we developed using the process of meta-ethnography to synthesise knowledge of influences on women's birth preferences after CS.

Women approaching a birth after a CS generally have either a clear preference for VBAC, a clear preference for ERCS or a relatively open mind to either option. Although some studies by design included women from only one or two of these categories, looking across the studies, we were able to develop a line of argument to explain how their findings were related. In summary the line of argument is that three distinctive clusters of influences support the three attitudinal positions that women adopt towards mode of birth after CS.

The three positions and the distinctive influences on these are summarised in figure 2 and described below. We note that the influences could be operative from different times, and that some were significant before and around the first CS.

Preferences for vaginal birth

Preferences for vaginal birth could be shaped by influences acting over a period of time, which for some women reached several years, and for many was linked to key events or periods of their lives. With respect to women's longstanding anticipation of vaginal birth, some women had a personal ambition to achieve vaginal birth that predated their first pregnancy and drove them to pursue VBAC. (M17 and M3) This could act synergistically with negative responses to a previous birth experience. For example, unpleasant memories of the initial CS experience, particularly where women had felt a loss of control over that birth, led some women to view VBAC as a potentially life-enriching experience that met their ambitions and avoided further negative emotions. (M1, M19, M9, M3, M8, M14) This impression was often enhanced by inter-pregnancy social interaction with influential others, including women who

provided encouragement by sharing accounts of their own positive VBAC experiences. (M19) For some, the probability of successful VBAC was pivotal. (M1, M3)

Future considerations could also play an important role in the shaping of preferences for VBAC, as women considered implications beyond the birth itself when evaluating their expected net gain from VBAC. Several women believed that VBAC offered physiological benefits to physical and emotional health of both themselves and their offspring, with particular emphasis on the facilitation of bonding and breastfeeding. (M17, M3) This was a particularly dominant issue among women who experienced breastfeeding difficulties after a previous planned CS, especially in those who had successfully breast fed their babies born vaginally in prior pregnancies. (M3) The social benefits of being able to return to usual family roles and resume driving as soon as possible in the postnatal period were also cited as reasons for preferring to avoid CS particularly within UK study settings. (M7, M8 and M9)

Further influential people included health professionals who provided support, advice or encouragement in favour of VBAC.. Women’s perception of the extent to which they themselves should make the decision regarding planned birth mode was important. Although some women, particularly in the UK and Australia, were confident about their right to decide how to plan their birth, (M18, M7, M17, M1) others judged any personal reasons they had in favour of ERCS to be unimportant or unjustified when considered in light of medical advice in favour of VBAC.(M8)

Preferences for elective repeat CS

Response to the previous birth experience was the central theme among women who demonstrated a clear preference to have an ERCS. A previous emergency CS in labour appeared to lead many women to believe their bodies were incapable of vaginal birth. (M8, M10, M13) Some women sought an ERCS to actively avoid any possibility of a repeat emergency CS, (M8, M10, M13) while others feared the possibility of a recurrence of the factors which led to the previous CS. Others opted for ERCS on the grounds that it was a familiar and positive birth experience. (M19, M5, M6)

The previous birth and its outcome could also shape women’s perceptions of the safety of VBAC (as outlined, it could lead to an assessment of net harm from planning VBAC), moderate the influence of social contacts (favouring those who encouraged ERCS and/or discouraged from planning VBAC) and limit the degree to which they felt they had a choice to make in the subsequent pregnancy (role in decision-making).

Safety concerns were described as particularly influential among some women in Australia who wished to avoid VBAC due to fear of the uterine scar 'splitting', or 'rupturing' during labour. This feeling dominated their preference for ERCS despite awareness of neonatal breathing problems being more common following this mode of birth. (M4) Some women with a strong preference for VBAC had been influenced, sometimes powerfully, by family, friends and health professionals who recommended ERCS as a safer and more predictable mode of birth than VBAC. (M13, M4)

Ownership of choice, or lack of the same, appeared crucial in determining whether or not some women opted for ERCS. Many women perceived that their health professionals would prefer this option, and as such, that VBAC was not available to them. (M15) Others choosing ERCS felt happy to exercise their preference as they had been positively encouraged to opt for the mode of birth that felt right for them. (M5)

Open-minded approach

Women who did not have a firm preference for either VBAC or ERCS appeared to be less strongly influenced by prior expectations about childbirth or by their previous birth experience than those who were more committed to one particular mode of birth. Influential others were apparently key to the decisions made in this context. These women valued and often actively sought the opinion of health professionals during their pregnancy, processed information on the options available, and put considerable effort into weighing up the attributes of the birth options available to assess net benefit. An exception to this involved women who felt overwhelmed by the decision-making responsibility and preferred to follow health professionals' advice. (M19, M8 and M18) Obstetricians, and, at times, midwives, appeared to have particular influence over women who were open to considering either mode of birth, even when women were not actively advised how to deliver but perceived subtle signals that their health professional had a preference. (M11) Some women said their choice should be based upon information alone, rather than the input or opinions of others, recognising that other people are not necessarily impartial. (M17)

Robustness of findings

On 'testing back the fit' of our line of argument, we found that the clusters of influence we identified were consistent with the findings of each of the individual included studies, but that none of these studies included a broad enough mix of participants to have enabled the development of this level of understanding in isolation.

Further ‘testing’ of the line of argument was made possible by the publication of the three new studies identified in the update of the search conducted in 2015, which are summarised in table 3. . Shorten et al analysed written text in which women explained their reasons for choosing either mode of birth after CS. They highlighted the significance of previous birth experience, safety concerns and speed of recovery along with health professionals’ preferences in shaping eventual decisions. Although they did not describe a clear distinction between the attitudinal groups, their findings were broadly supportive of the conclusions of this synthesis, with no evidence of confliction or contradiction.³⁰ Kennedy et al performed an institutional ethnography exploring the complexity of choice around elective CS.⁴⁸ This included interviews with women within English National Health Service provider settings. The authors identified that women planning birth after CS negotiated with clinicians to reach a ‘comfortable compromise’ which facilitated a plan for VBAC that included adequate assurance of early recourse to CS if labour progress was suboptimal. This supports our findings of the crucial role of health professionals in influencing VBAC decisions by providing support for this option. Further author interpretation echoed our emphasis on the importance of predicted VBAC success in influencing women to aim for this mode of birth. Finally, the authors highlighted the desire for information among some women, providing an exemplary quote which supported our impression that women with an open mind to birth mode after CS place great emphasis upon the content, and in this case, quality of information accessed: ‘When I was getting told about the 0.3% chance of a scar rupturing, you know, when I was asking people about how that statistic was arrived at no one could tell me, so I kept digging for more and more information, ‘and there’s just not enough research, there’s not enough studies that have been done, the women aren’t in the same circumstances, they’re not all in even one country, it’s international, it’s in under- developed countries, so you’re pulling together these statistics from a complete diverse set of sample set,.and how can you make judgements on what an individual’s circumstances are going to be based on that? There’s just not enough there’s not enough information out there to be able to say you’re going to be one of those statistics.(P108;woman pondering VBAC decision)

Tully et al presented findings of an interview study of 115 mothers recently delivered by CS over a three year period in England.⁴⁹ Although minimal primary or secondary constructs related to birth after CS were presented, there was evidence that predicted VBAC success was important to women aiming for a vaginal birth, and that a negative previous birth experience drove women to seek control and predictability in the form of an ERCS. These observations are consistent with our findings, and no evidence of contradictory interpretations was identified.

Table 3. Studies identified in the updated search which were used to ‘test the fit’ of the line of argument

Author	Year	Country	Study Aim	Data collection method	Planned birth method at time of	Participants (n)	Timing of data collection

					study		
Shorten ³⁰	2014	Australia	explore values and expectations that guide women during decision making about the next birth after caesarean and identify factors that influence consistency between women's choices and actual birth experiences	Written surveys and narrative accounts	VBAC and CS	187	36-37 weeks gestation and postnatal
Kennedy ⁴⁸	2013	UK	To explore the complexities of women's and clinicians' choices around elective caesarean delivery	Interview (FTF) and consultation observations.	CS and vaginal birth (sample not restricted to birth after CS)	27 women of whom three had VBAC and 19 had no history of prior CS. Previous obstetric history of 5 participants who underwent CS was not clear	Not specified - appears to span antenatal and postnatal period
Tully ⁴⁹	2013	UK	To document the circumstances in which caesarean section was deemed to be appropriate in one UK hospital through the eyes of the women and their partners experiencing the operative delivery of their infant.	Interview (FTF)	Not applicable (postnatal)	115 women	Postnatal hospital stay

FTF=face-to-face, VBAC=vaginal birth after CS, CS=Caesarean section.

DISCUSSION

Summary of main findings

This study sought to answer the research question 'What influences women's preferred mode of birth after previous caesarean section?' We have identified distinct clusters of influences that tend to underpin the three main positions that pregnant women adopt towards modes of birth. After an initial CS, women tend to approach childbirth with one of three broad attitudinal positions meaning that they: 1) seek vaginal birth 2) seek repeat caesarean or 3) are open minded to consideration of either mode of birth. These positions reflect thought processes which are likely to evolve from at least as early as the primary CS, with some influential cultural norms in operation well before that time. A strong preference for VBAC appears to be driven by a belief that vaginal birth is 'normal' and has some intrinsic value. This belief is often accompanied by a keen desire to resume a normal life soon after vaginal birth. In contrast, a clear preference for elective repeat CS from early in pregnancy can be driven by a previous negative experience of attempting but failing to achieve vaginal birth and a positive emphasis on the predictability of ERCS. Finally, there are women who embark upon their next pregnancy undecided about mode of birth. These women are more open to external influence: they appreciate the benefits of both written information and personalised expert advice, which they use to weigh up what they see as

the advantages and disadvantages of their options. The recognition of these clusters of influences according to attitude towards birth from early in the pregnancy after CS is a novel finding made possible by looking across the range of relevant studies. Both historical and contemporary studies have highlighted influences on birth preferences after CS which resonate with those identified in this synthesis, but without identification of attitudinal groups or attention to the multiple influences and the ways these may vary over time.^{27,27,50,51,51} The importance of timing of influence has, however, been highlighted recently by prospective work which found that first-trimester preferences for either ERCS or VBAC persist by early in the third trimester in over 70% of women.⁵²

Benefits of a meta-ethnographic approach

Meta-ethnography enabled an interpretation of the available research that incorporated a sensitivity to the contextual factors surrounding the influences reported by specific groups of women planning birth after CS.⁵³ Contextual factors considered included key time points at which influences took hold, fundamental study characteristics (setting; eligibility criteria; recruitment processes; timing of interviews; healthcare systems) and factors unique to individual women. These contextual considerations limit the likelihood that findings would be generalised inappropriately. The iterative process of reciprocal translation used to build upon emergent themes facilitated a higher level of understanding than previous mixed-method review methodology has allowed, particularly that of quantitative work, where presence or absence of potential influences has been the focus.²⁵ The clustering of influences identified within specific attitudinal groups provided clinically relevant insight into the nature of women’s decision-making behaviour. In addition, the identification of clustering was considered robust in light of the ‘testing back the fit’ which confirmed that primary authors’ interpretations supported specific attitudinal clusters.

Women’s perspectives

The specific focus upon *women’s* perspectives on what influences birth preferences after CS complements the current focus on joint health care decision-making in which informed patients contribute to decisions which reflect their beliefs and preferences.⁵⁴ This therefore provides insight which has maximal clinical application in settings where every effort should be made to ensure decisions about mode of birth after CS incorporate women’s values and preferences. Given that health professionals have a variable level of input into shaping the eventual mode of birth, it is possible that consideration of health professionals’ perspectives may have further developed our understanding of

the decision-making process.⁵⁵ However, women's insights were considered central to achieving the goal of informing future efforts to optimise and support woman-centred planning of birth after CS.

Clinical and research implications

Reflection on current practice

The strength of evidence supporting the first CS birth experience as a key influence upon future birth preferences demands immediate attention. Women should be effectively supported in dealing with the unexpected and potentially traumatic nature of a primary CS. Efforts to promptly address any inaccurate perceptions of their CS birth events, and to provide personally specific information about the risks and benefits of future birth options could be made following the first CS and be reiterated early in the pregnancy after CS. The findings of this synthesis suggest that women's concerns about serious maternal or offspring health risks (beyond those of CS scar rupture) are not important influences on their birth choices after CS. This is of particular interest because information currently provided by health professionals for women planning birth after CS focuses largely on these risks and clinical health considerations.²⁰ Recognition of this mismatch between what women and health professionals prioritise should prompt health professionals to engage in discussion with women which allows identification of their main concerns and places sufficient emphasis on the psychological and social as well as the physical health consequences of modes of birth after CS. The heterogeneity of influences on birth choices after CS demonstrated in this synthesis highlight why approaching all women planning birth after CS with eg. the same decision support tool in the latter part of pregnancy is unlikely to alter their prior attitudinal positions.

Implications for future research and practice

Recognition of the diverse range of influences on, and attitudes towards, birth after CS enables us to understand why decision support interventions have had limited effects on ERCS rates so far,^{21,22} and opens up the possibility of a more targeted approach. We suggest that future interventions should aim to promote positive experiences of informed and shared decision-making, while minimising maternal and fetal morbidity and avoiding unnecessary healthcare costs. Insights from this synthesis suggest that future strategies should ensure early consideration of both women's concerns and preferences and their likelihood of achieving good physical birth outcomes. Women may be broadly categorised in early pregnancy after CS as being in favour of either VBAC or ERCS or being open to either option. At the same time, their prognosis for successful VBAC may also be assessed based upon factors such as their age,

BMI and indication for previous CS.^{8,56} In order to support high quality decision making and increase VBAC success rates, efforts could be made to ensure design of decision support which reflects women’s prognosis for VBAC success and is sensitive to any early preferences regarding birth mode after CS. The six main prognosis/preference categories are represented in figure 3.

Decision support for women may be delivered via conversations with health professionals, advice and information, including decision aids.⁵⁷ Decision aids provide women with information about options relevant to their health status, while helping them to reflect and draw on their personal values. Previous research has demonstrated that use of some such tools in supporting birth choices after CS improved decision satisfaction but had minimal impact upon VBAC rates.²² The lack of success in increasing VBAC rates may reflect that the tools that were tested were not tailored to women’s early attitudes towards each birth mode, but instead delivered advice according to outcomes which women prioritised. Faced with a choice of surgery and less invasive options, decision-aids have been shown to lead patients to choose conservative or less invasive treatments.⁵⁸

In the context of planning birth after CS, decision aids might usefully be stratified according to predicted VBAC success and responsive to individual women’s early birth mode preferences and priorities. It is likely to be particularly important to engage women who are open-minded (groups E and F on figure 3) and women with a VBAC prognosis which is at odds with their preferred mode of birth (groups B and C on figure 3) by the second trimester, in conversations with health professionals to ensure sufficient time to explore their views and discuss and allow them to consider their options. In such situations, a ‘consider a recommendation’ approach may be warranted – explaining why either ERCS or VBAC is recommended, but leaving sufficient scope and ensuring sufficient support for women to assess and discuss the recommendation before making their own mind up about it.⁵⁹ In those pursuing VBAC despite a poor prognosis for success, there could be a discussion about criteria for conversion to CS and adequate counselling in preparation for the possible psychological impact of such an outcome. Those in whom VBAC prognosis is in keeping with their preferred mode of birth (groups A and D on figure 3) might need less in the way of information, conversation and recommendations from health professionals, but their needs for information and reassurance about their decisions should not be neglected: balanced written information regarding the risks and benefits of both birth options, and clarification/confirmation of ongoing preferences are still likely to be important. As events unfold during subsequent pregnancies, ongoing communication and decision support for all women would need to be tailored to accommodate new clinical information, concerns and preferences, but a broad pathway

identified following the first CS would ensure timely and relevant intervention to address modifiable influences.

CONCLUSIONS

Forming a preference for repeat CS or VBAC is a dynamic process, shaped by many influences which appear to cluster distinctively in the development of strongly held positions. Long-standing expectations of childbirth and perceptions of previous birth experiences appear particularly influential upon VBAC and ERCS preferences respectively. This suggests that early communication to discuss women's prospects for VBAC success and explore and discuss their attitudes towards future births may be valuable, and could perhaps start from as early as the first CS. This might help increase the proportion of women who approach birth after CS with an open mind being receptive to written information and the advice of health professionals. Our synthesis has highlighted why current care models involving information provision in pregnancy after CS may not lead to the birth choices which could help reduce the unnecessary CS rate. It suggests a need to address women's social and psychological concerns, and not just the currently recommended information, both to support women's autonomy in decision-making and to address public health concerns about rising rates of clinically unnecessary CS.

References

1. World Health Organisation. World health statistics 2013.

http://www.who.int/gho/publications/world_health_statistics/2013/en/. Updated 2013.

2. Stavrou EP, Ford JB, Shand AW, Morris JM, Roberts CL. Epidemiology and trends for caesarean section births in New South Wales, Australia: A population-based study. *BMC Pregnancy & Childbirth*. 2011;11:8.

3. Almeida S, Bettiol H, Barbieri MA, Silva AA, Ribeiro VS. Significant differences in cesarean section rates between a private and a public hospital in Brazil. *Cad Saude Publica*. 2008;24(12):2909-2918.

4. Lumbiganon P, Laopaiboon M, Gulmezoglu AM, et al. Method of delivery and pregnancy outcomes in Asia: The WHO global survey on maternal and perinatal health 2007-08. *Lancet*. 2010;375(9713):490-499.

5. Minkoff H, Powderly KR, Chervenak F, McCullough LB. Ethical dimensions of elective primary cesarean delivery. *Obstetrics & Gynecology*. 2004;103(2):387-392.

6. Timofeev J, Reddy UM, Huang CC, Driggers RW, Landy HJ, Laughon SK. Obstetric complications, neonatal morbidity, and indications for cesarean delivery by maternal age. *Obstet Gynecol*. 2013;122(6):1184-1195.

7. Landon MB, Hauth JC, Leveno KJ, et al. Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med*. 2004;351(25):2581-2589.

8. Knight HE, Gurol-Urganci I, van der Meulen JH, et al. Vaginal birth after caesarean section: A cohort study investigating factors associated with its uptake and success. *BJOG*. 2014;121(2):183-192.

9. Paterson CM, Saunders NJ. Mode of delivery after one caesarean section: Audit of current practice in a health region. *BMJ*. 1991;303(6806):818-821.

10. McMahon MJ, Luther ER, Bowes WA, Jr, Olshan AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med*. 1996;335(10):689-695.

11. Dodd JM, Crowther CA, Huertas E, Guise JM, Horey D. Planned elective repeat caesarean section versus planned vaginal birth for women with a previous caesarean birth. *Cochrane Database of Systematic Reviews*. 2004(4). doi: 10.1002/14651858.
12. American College of Obstetricians and Gynecologists. ACOG practice bulletin no. 115: Vaginal birth after previous cesarean delivery. *Obstet Gynecol*. 2010;116(2 Pt 1):450-463.
13. Centres for Disease Control and Prevention. Trends in cesarean birth and vaginal birth after previous cesarean, 1991-99. *National vital statistics reports: from the Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System*. 2002;49(13):15.
14. Main EK, Morton CH, Melsop K, Hopkins D, Giuliani G, Gould JB. Creating a public agenda for maternity safety and quality in cesarean delivery. *Obstet Gynecol*. 2012;120(5):1194-1198.
15. Macdonald H, Loder E. Too much medicine: The challenge of finding common ground. *BMJ*. 2015;350:1163.
16. National Institute for Health and Clinical Excellence. Caesarean section; clinical guideline no. 132. 2011.
17. Dursun P, Yanik FB, Zeyneloglu HB, Baser E, Kuscu E, Ayhan A. Why women request cesarean section without medical indication? *J Matern Fetal Neonatal Med*. 2011;24 (9):1133-1137.
18. Fenwick J, Gamble J, Hauck Y. Reframing birth: A consequence of cesarean section. *J Adv Nurs*. 2006;56 (2):121-130.

19. Guise J-M, Eden K, Emeis C, Denman MA, Marshall N, Fu R, Janik R, Nygren P, Walker M, McDonagh M. Vaginal birth after cesarean: New insights. evidence report/technology assessment no.191. . 2010;AHRQ Publication No. 10-E003.

20. Royal College of Obstetricians and Gynaecologists. Birth after previous caesarean birth; greentop guideline. 2007;45.

21. Shorten A, Shorten B, Keogh J, West S, Morris J. Making choices for childbirth: A randomized controlled trial of a decision-aid for informed birth after cesarean. *Birth*. 2005;32 (4):252-261.

22. Montgomery AA, Emmett CL, Fahey T, et al. Two decision aids for mode of delivery among women with previous caesarean section: Randomised controlled trial. *Br Med J*. 2007;334 (7607):1305-1309.

23. Liu S, Li H, Lee S. The experiences of multipara who chose to undergo vaginal birth after cesarean [chinese]. *J EVID BASED NURS*. 2006;2(3):241-249.

24. Phillips E, McGrath P, Vaughan G. 'I wanted desperately to have a natural birth': Mothers' insights on vaginal birth after caesarean (VBAC). *Contemporary Nurse*. 2009;34(1):77-84.

25. Eden KB, Hashima JN, Osterweil P, Nygren P, Guise JM. Childbirth preferences after cesarean birth: A review of the evidence. *Birth*. 2004;31(1):49-60.

26. Shorten A, Shorten B. The importance of mode of birth after previous cesarean: Success, satisfaction, and postnatal health. *J Midwifery Womens Health*. 2012;57(2):126-132.

27. Flannagan C, Reid B. Repeat CS or VBAC? A systematic review of the factors influencing pregnant women's decision-making processes. *Evidence Based Midwifery*. 2012;10(3):80-86.

28. Noblit G. HR. *Meta-ethnography: Synthesising qualitative studies*. Sage Publications; 1988.
29. Collaboration for qualitative methodologies. Critical appraisal skills program. www.casp-uk.net. Updated 1998.
30. Shorten A, Shorten B, Kennedy HP. Complexities of choice after prior cesarean: A narrative analysis. *Birth*. 2014;41(2):178-184.
31. Ridley RT, Davis PA, Bright JH, Sinclair D. What influences a woman to choose vaginal birth after cesarean? *J Obstet Gynecol Neonatal Nurs*. 2002;31(6):665-672.
32. York S, Briscoe L, Walkinshaw S, Lavender T. Why women choose to have a repeat caesarean section. *BR J MIDWIFERY*. 2005;13(7):440-445.
33. Emmett CL, Shaw ARG, Montgomery AA, Murphy DJ. Women's experience of decision making about mode of delivery after a previous caesarean section: The role of health professionals and information about health risks. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2006;113(12):1438-1445.
34. Cheung NF, Mander R, Cheng L, Chen VY, Yang X. Caesarean decision-making: Negotiation between chinese women and healthcare professionals. *EVID BASED MIDWIFERY*. 2006;4(1):24-30.
35. Meddings F, Phipps FM, Haith-Cooper M, Haigh J. Vaginal birth after caesarean section (VBAC): Exploring women's perceptions. *J Clin Nurs*. 2007;16(1):160-167.
36. Moffat MA, Bell JS, Porter MA, et al. Decision making about mode of delivery among pregnant women who have previously had a caesarean section: A qualitative study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2007;114(1):86-93.

37. Fenwick J, Gamble J, Hauck Y. Believing in birth - choosing VBAC: The childbirth expectations of a self-selected cohort of australian women. *J Clin Nurs*. 2007;16(8):1561-1570.

38. Farnsworth A, Pearson PH. Choosing mode of delivery after previous caesarean birth. *BR J MIDWIFERY*. 2007;15(4):188.

39. Cox B. Hot topic. women's perceptions of their access to, and value of, information as part of their decision making on mode of birth following a previous caesarean section delivery. *MIDIRS MIDWIFERY DIGEST*. 2007;17(2):159-168.

40. Farnworth A, Robson SC, Thomson RG, Watson DB, Murtagh MJ. Decision support for women choosing mode of delivery after a previous caesarean section: A developmental study. *Patient Educ Couns*. 2008;71(1):116-124.

41. McGrath P, Ray-Barruel G. The easy option? australian findings on mothers' perception of elective caesarean as a birth choice after a prior caesarean section. *Int J Nurs Pract*. 2009;15(4):271-279.

42. McGrath P, Phillips E, Ray-Barruel G. Bioethics and birth: Insights on risk decision-making for an elective caesarean after a prior caesarean delivery. *Monash Bioethics Review*. 2009;28(3):22.1-2219.

43. Goodall KE, McVittie C, Magill M. Birth choice following primary caesarean section: Mothers' perceptions of the influence of health professionals on decision-making. *J REPROD INFANT PSYCHOL*. 2009;27(1):4-14.

44. Frost J, Shaw A, Ontgomery A, Murphy D. Women's views on the use of decision aids for decision making about the method of delivery following a previous caesarean section: Qualitative interview study. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2009;116(7):896-905.

45. McGrath P, Phillips E, Vaughan G. Vaginal birth after caesarean risk decision-making: Australian findings on the mothers' perspective. *Int J Nurs Pract*. 2010;16(3):274-281.
46. David S, Fenwick J, Bayes S, Martin T. A qualitative analysis of the content of telephone calls made by women to a dedicated 'next birth after caesarean' antenatal clinic. *Women & Birth: Journal of the Australian College of Midwives*. 2010;23(4):166-171.
47. McGrath P, Phillips E, Vaughan G. Speaking out! qualitative insights on the experience of mothers who wanted a vaginal birth after a birth by cesarean section. *The Patient: Patient-Centered Outcomes Research*. 2010;3(1):25-32.
48. Kennedy HP, Grant J, Walton C, Sandall J. Elective caesarean delivery: A mixed method qualitative investigation. *Midwifery*. 2013;29(12):e138-44.
49. Tully K, Ball H. Misrecognition of need: Women's experiences of and explanations of cesarean delivery. *Social Science and Medicine*. 2013;85:103-111.
50. McClain CS. Patient decision making: The case of delivery method after a previous cesarean section. *Cult Med Psychiatry*. 1987;11(4):495-508.
51. Lundgren I, Begley C, Gross MM, Bondas T. 'Groping through the fog': A metasynthesis of women's experiences on VBAC (vaginal birth after caesarean section). *BMC Pregnancy and Childbirth*. 2012;12.
52. Shorten A, Shorten B. Timing the provision of a pregnancy decision-aid: Temporal patterns of preference for mode of birth during pregnancy. *Patient Educ Couns*. 2014;97(1):108-113.
53. Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: A critical review. *BMC Med Res Methodol*. 2009;9:59-2288-9-59.

54. Dy SM, Purnell TS. Key concepts relevant to quality of complex and shared decision-making in health care: A literature review. *Soc Sci Med*. 2012;74(4):582-587.

55. Kamal P, Dixon-Woods M, Kurinczuk JJ, Oppenheimer C, Squire P, Waugh J. Factors influencing repeat caesarean section: Qualitative exploratory study of obstetricians' and midwives' accounts. *BJOG : an international journal of obstetrics and gynaecology*. 2005;112(8):1054-1060.

56. Grobman WA, Lai Y, Landon MB, et al. Development of a nomogram for prediction of vaginal birth after cesarean delivery. *Obstet Gynecol*. 2007;109(4):806-812.

57. Say R, Murtagh M, Thomson R. Patients' preference for involvement in medical decision making: A narrative review. *Patient Educ Couns*. 2006;60(2):102-114.

58. Knops AM, Legemate DA, Goossens A, Bossuyt PM, Ubbink DT. Decision aids for patients facing a surgical treatment decision: A systematic review and meta-analysis. *Ann Surg*. 2013;257(5):860-866.

59. Entwistle VA, Carter SM, Trevena L, et al. Communicating about screening. *BMJ*. 2008;337:a1591.

Competing interests. No, there are no competing interests.

Acknowledgements: None.

Contributorship: MB and SB conceived the idea of the study. MB, SB and KG planned the study. MB and KG conducted the literature search and analysed all data. VE and SB contributed to data analysis and interpretation. MB wrote all drafts of the manuscript and is guarantor. VE, SB and KG contributed to all drafts of the manuscript.

Funding: Mairead Black is a research training fellow funded by The Wellcome Trust. The funder played no role in the design and conduct of the study, in the collection, analysis and interpretation of the data, or in the preparation, review, or approval of the manuscript.

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, [a worldwide licence](#) to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints,

1
2
3 *include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create*
4 *any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the*
5 *Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever*
6 *it may be located; and, vi) licence any third party to do any or all of the above."*
7
8

9
10 Ethical approval: Not required.

11
12 Transparency: The lead author (the manuscript's guarantor) affirms that the manuscript is an honest,
13 accurate, and transparent account of the study being reported; no important aspects of the study have
14 been omitted.
15

16
17 Data sharing: Additional data is available by emailing the corresponding author.

18
19 All authors had full access to all of the data in the study and can take responsibility for the integrity of
20 the data and the accuracy of the data analysis.
21

22
23
24 Box 1. 'Box 1. Meta-ethnography steps as described by Noblit and Hare, 1988.'

25
26
27 Figure 1. 'Figure 1. Flow diagram of search results' CS, caesarean section.

28
29 Figure 2. 'Figure 2. Summary attitudinal positions of women early in the pregnancy after CS and clusters
30 of key influences acting upon their eventual birth preferences.'

31
32 Figure 3. 'Figure 3. Table represents how women may be categorised according to their preferred mode
33 of birth in early pregnancy and their prognosis for VBAC success' VBAC, vaginal birth after caesarean;
34 ERCS, elective repeat caesarean section.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1. Identify the research question
- 2. Identify relevant studies
- 3. Read the studies
- 4. Identify themes
- 5. Translate the findings of each study into those of the others
- 6. Synthesise the findings
- 7. Express the synthesis

Box 1. Meta-ethnography steps as described by Noblit and Hare, 1988
297x420mm (300 x 300 DPI)




Figure 1. Flow diagram of search results' CS, caesarean section
297x420mm (300 x 300 DPI)

Sought Vaginal birth	Sought repeat CS	Open minded
<ul style="list-style-type: none">•Longstanding anticipation of vaginal birth•Responses to previous birth experiences (negative)•Perceived net benefit or harm of birth options (postnatal issues predominating)•Encouragement and dissuasion from influential people (for VBAC and CS respectively)•Extent and nature of involvement in decision-making	<ul style="list-style-type: none">•Responses to previous birth experiences (negative and positive)•Fear or reassurance from risk-related information on VBAC•Encouragement and dissuasion from influential people [for CS and VBAC respectively]•Extent and nature of involvement in decision-making•Perceived net benefit or harm of birth options	<ul style="list-style-type: none">•Encouragement and dissuasion from influential people for either birth mode•Fear or reassurance from risk-related information on VBAC•Perceived net benefit or harm of birth options (mix of health and social considerations predominating)•Extent and nature of involvement in decision-making

Figure 2. Summary attitudinal positions of women early in the pregnancy after CS and clusters of key influences acting upon their eventual birth preferences
297x420mm (300 x 300 DPI)

	Good VBAC prognosis	Poor VBAC prognosis
Prefers VBAC	A	B
Prefers ERCS	C	D
Open minded	E	F

Figure 3. Table represents how women may be categorised according to their preferred mode of birth in early pregnancy and their prognosis for VBAC success' VBAC, vaginal birth after caesarean; ERCS, elective repeat caesarean section
297x420mm (300 x 300 DPI)

Search Strategy V3 Medline(R) In-process and other non-indexed citations and Ovid MEDLINE(R)
<1946 to Present>

Date;

1 exp caesarean section/ or vaginal birth after caesarean/

2 c?esarean section?.ti.

3 1 or 2

4 qualitative research/

5 questionnaires/

6 exp interviews as topic/

7 (qualitative or interview\$ or focus group? Or questionnaire\$ or survey\$).tw.

8 (ethno\$ or grounded or thematic or interpretive or narrative).tw.

9 or/4-8

10 3 and 9

11 Choice behaviour/

12 Decision Making/

13 Patient Preference/

14 Patient Acceptance of Health Care/

15 Attitude to health/

16 intention/

17 (choice? Adj2 (birth or childbirth or delivery or vaginal or c?esarean)).tw.

18 (elective adj2 (c?esarean or CS)).tw.

19 vbac.tw.

20 (prefer\$ adj3 (birth or childbirth or delivery or vaginal or c?esarean)).tw.

21 (c?esarean adj2 (after or subsequent or prior or previous or follow\$)).tw.

22 or/11-21

23 10 and 22

Author	Clear study aim	Qualitative methods appropriate	Design appropriate	Recruitment strategy appropriate	Data collection process appropriate	Researcher role in forming research questions discussed	Ethical issues considered	In-depth description of analysis	Adequate discussion of the evidence	Contribution to existing knowledge discussed
Ridley ³¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
York ³²	Yes	Yes	Yes	N/C	Yes	N/C	Yes	No	Yes	Yes
Liu ²³	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Fenwick ¹⁸	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Emmett ³³	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Cheung ³⁴	Yes	Yes	Yes	No – snowball*	Yes	N/C	Yes	Yes	Yes	Yes
Meddings ³⁵	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Moffat ³⁶	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Fenwick ³⁷	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Farnworth ³⁸	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Cox ³⁹	Yes	Yes	Yes	Yes	Yes	N/C	Yes	No	Yes	Yes
Farnworth ⁴⁰	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴¹	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴²	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes
Goodall ⁴³	Yes	Yes	Yes	No	Yes	N/C	Yes	Yes	Yes	Yes
Frost ⁴⁴	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
Phillips ²⁴	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
McGrath ⁴⁵	Yes	Yes	Yes	Yes	Yes	N/C	Yes	Yes	Yes	Yes
David ⁴⁶	Yes	Yes	Yes	Yes	Yes	N/C	N/C	Yes	Yes	Yes
McGrath ⁴⁷	Yes	Yes	Yes	N/C	Yes	N/C	Yes	Yes	Yes	Yes

Appendix Table 1. Critical Appraisal Skills Programme Assessment of Included Studies

N/C=Not clear *judged inappropriate to meet the specific study aim

Enhancing transparency in reporting the synthesis of qualitative research: the ENTREQ statement

No	Item	Guide and description	Section heading containing relevant information
1	Aim	State the research question the synthesis addresses.	Abstract and introduction
2	Synthesis methodology	Identify the synthesis methodology or theoretical framework which underpins the synthesis, and describe the rationale for choice of methodology (e.g. meta-ethnography, thematic synthesis, critical interpretive synthesis, grounded theory synthesis, realist synthesis, meta-aggregation, meta-study, framework synthesis).	Methods
3	Approach to searching	Indicate whether the search was pre-planned (comprehensive search strategies to seek all available studies) or iterative (to seek all available concepts until they theoretical saturation is achieved).	Methods and appendix
4	Inclusion criteria	Specify the inclusion/exclusion criteria (e.g. in terms of population, language, year limits, type of publication, study type).	Methods Table 1
5	Data sources	Describe the information sources used (e.g. electronic databases (MEDLINE, EMBASE, CINAHL, psycINFO, Econlit), grey literature databases (digital thesis, policy reports), relevant organisational websites, experts, information specialists, generic web searches (Google Scholar) hand searching, reference lists) and when the searches conducted; provide the rationale for using the data sources.	Methods
6	Electronic Search strategy	Describe the literature search (e.g. provide electronic search strategies with population terms, clinical or health topic terms, experiential or social phenomena related terms, filters for qualitative research, and search limits).	Methods and appendix 1
7	Study screening methods	Describe the process of study screening and sifting (e.g. title, abstract and full text review, number of independent reviewers who screened studies).	Methods
8	Study characteristics	Present the characteristics of the included studies (e.g. year of publication, country, population, number of participants, data collection, methodology, analysis, research questions).	Results Table 2
9	Study selection results	Identify the number of studies screened and provide reasons for study exclusion (e.g. for comprehensive searching, provide numbers of studies screened and reasons for exclusion indicated in a figure/flowchart; for iterative searching describe reasons for study exclusion and inclusion based on modifications to the research question and/or contribution to theory development).	Results Figure 1
10	Rationale for appraisal	Describe the rationale and approach used to appraise the included studies or selected findings (e.g. assessment of conduct (validity and robustness), assessment of reporting (transparency), assessment of content and utility of the findings).	Methods
11	Appraisal items	State the tools, frameworks and criteria used to appraise the studies or selected findings (e.g. Existing tools: CASP, QARI, COREQ, Mays and Pope [25]; reviewer developed tools; describe the domains assessed: research team, study design, data analysis and interpretations, reporting).	Methods
12	Appraisal process	Indicate whether the appraisal was conducted independently by more than one reviewer and if consensus was required.	Methods
13	Appraisal results	Present results of the quality assessment and indicate which articles, if any, were weighted/excluded based on the assessment and give the rationale.	Methods and appendix 2
14	Data extraction	Indicate which sections of the primary studies were analysed and how were the data extracted from the primary studies? (e.g. all text under the headings "results /conclusions" were extracted electronically and entered into a computer software).	Methods
15	Software	State the computer software used, if any.	N/A
16	Number of reviewers	Identify who was involved in coding and analysis.	Methods
17	Coding	Describe the process for coding of data (e.g. line by line coding to search for concepts).	Methods
18	Study comparison	Describe how were comparisons made within and across studies (e.g. subsequent studies were coded into pre-existing concepts, and new concepts were created when deemed necessary).	Methods

19	Derivation of themes	Explain whether the process of deriving the themes or constructs was inductive or deductive.	Methods
20	Quotations	Provide quotations from the primary studies to illustrate themes/constructs, and identify whether the quotations were participant quotations of the author's interpretation.	Methods
21	Synthesis output	Present rich, compelling and useful results that go beyond a summary of the primary studies (e.g. <i>new interpretation, models of evidence, conceptual models, analytical framework, development of a new theory or construct</i>).	Results, particularly 'patterns of influence: a line of argument' and discussion 'clinical and research implications' section.