BMJ Open

Does case management for patients with heart failure reduce unplanned hospital admissions? A systematic review and meta-analysis

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-010933
Article Type:	Research
Date Submitted by the Author:	22-Dec-2015
Complete List of Authors:	Huntley, Alyson; University of Bristol, Centre of Academic Primary care Johnson, Rachel; Academic unit of primary health care, School of social and community medicine King, Anna; University of Bristol, Centre of academic Primary care Morris, Richard; University of Bristol, School of Social & Community Medicine Purdy, Sarah; University of Bristol, Centre for Academic Primary Care
Primary Subject Heading :	Cardiovascular medicine
Secondary Subject Heading:	General practice / Family practice
Keywords:	systematic review, meta-analysis, Heart failure < CARDIOLOGY, case management, hospital admission

SCHOLARONE™ Manuscripts

Version 20th December 2015

Does case management for patients with heart failure reduce unplanned hospital admissions? A systematic review and meta-analysis

Huntley AL1*

Johnson R¹

King A¹

Morris RW¹

Purdy S¹

Key words: systematic review, meta-analysis, heart failure, case management, hospital admission

Word count: 3,823

^{1.} Centre of Academic Primary Care, School of Social and Community Medicine, University of Bristol, Canynge Hall, Bristol, BS8 2PS, UK.

^{*}Dr Alyson Louise Huntley Research Fellow <u>alyson.huntley@bristol.ac.uk</u> Tel no: 0117 3314545

ABSTRACT

Objectives

This systematic review aims to extend our knowledge of the effectiveness and cost- effectiveness of case management (CM) for patients with heart failure (HF) in reducing unplanned admissions.

Setting

CM interventions initiated either whilst as an inpatient or on discharge from acute care hospitals including the emergency department, or in the community and then continuing on in the community.

Participants

Adults with a diagnosis of HF and resident in Organisation for Economic Co-operation and Development countries.

Intervention

CM based on nurse co-ordinated multi-component care of individual patients which is applicable to the primary care based health systems.

Primary and secondary outcomes

The primary outcomes of interest were unscheduled secondary care and cost effectiveness.

Secondary outcomes were primary health care resources.

Results

Twenty-two studies were included: 17 RCTs and five controlled trials. Seventeen of these studies described hospital-initiated CM and five described community-initiated CM of HF. Hospital -initiated CM showed a reduced risk of readmission (rate ratio 0.77 [0.63, 0.94] p=0.01) and length of hospital stay (mean difference -1.28 days [-2.03,-0.53] p=0.0008) in favour of CM compared to usual care. Nine trials described cost data of which six reported no difference between CM and usual care. Data from three of five trials of community-initiated CM showed no differences in risk of admission compared to the usual care group (rate ratio 1.08 [0.62, 1.87] p=0.8). Data from some individual studies suggests that family involvement and education/self-management are likely to be important in CM but this was not demonstrated in sub-group analysis.

Conclusions

Hospital-initiated CM can be successful in reducing unplanned hospital readmissions for HF and length of hospital stay for people with HF. Limited data suggest no difference in costs between usual care and CM. There were limited data for community-initiated CM which suggested it does not reduce hospital admission.

Protocol registration

No

Strengths of review

- High quality systematic review
- Interventions examine nurse-led multicomponent care of heart failure patients
- Focus on use of resources specific to heart failure

Limitations of review

- Community-initiated case management trials were limited in quantity and were mostly of low quality.
- Lack of cost data in most trials

INTRODUCTION

Applying current prevalence figures to population estimates suggests that more than 550,000 individuals (more than 308,000 men and slightly fewer than 250,000 women) in the UK are living with heart failure (HF). Quality and Outcome Framework (QOF) data supports this: in 2012/13 just over 480,000 patients were recorded as having heart failure. The average age of HF patients in general practice in the UK is 77 years.

Prior to 1990, 60–70% of patients died within 5 years of diagnosis, and admission to hospital with worsening symptoms was a regular and recurrent event. Effective treatment has improved care, with a relative reduction in hospitalization in recent years of 30–50%, and smaller but significant decreases in mortality. 4-6

More than £6.8 billion was spent on treating all cardiovascular disease within the NHS in England in 2012/13 with 63% of these costs coming from within secondary care and 21% within primary care. Within secondary care, non-elective inpatient admittance for cardiovascular disease, i.e. emergency admissions, had the greatest expenditure with £1,925 million.¹

Case management (CM) is the process of planning, coordinating and reviewing the care of an individual. We used the definition cited by the King's Fund in the UK 'A collaborative process of assessment, planning, facilitation, care coordination, evaluation, and advocacy for options and services to meet an individual's and family's comprehensive health needs through communication and available resources to promote quality cost-effective outcomes.' The NHS has used less-intensive approaches than the traditional US model, for example, through the use of nurses to support older people and those with long-term conditions at home. In this review we have focused on CM based on nurse co-ordinated multi-component care of patients which is applicable to the primary care based health systems such as that in the UK.

Previous research suggests that whilst CM is not effective in reducing unplanned hospital admissions for the general older/elderly population, the evidence for patients with HF is promising. ^{9,10} Therefore this review aimed to a) update the evidence of the effectiveness and cost-effectiveness of interventions by examining both randomised and non-randomised controlled trials and b) to better understand the potential success of CM by examining the components of tested interventions.

Version 20th December 2015

METHODS

Search

Databases and registries

A search strategy was developed using keywords for the electronic databases according to their specific subject headings or searching structure. The search strategy was run from 1985 – 2012 in the OVID databases - Medline®, Excerpta Medica Database (Embase), Cumulative Index to Nursing and Allied Health Literature (CINAHL®), PsycINFO® on the 2nd July2014. (Appendix one) The search strategy was modified to search internet sites such as the Agency for Healthcare Research and Quality (AHRQ) and the King's Fund. A pragmatic update of these searches was conducted on the 20/11/15 using the same search strategy and run in Medline and Medline in process only.

Other sources

Once the included papers were determined, both backwards (reference list of paper) and forwards citation searching (via google scholar) was performed to identify any other potentially relevant studies.

All authors of included studies in the field were contacted with data queries and to identify additional relevant studies.

Eligibility criteria:

Randomised controlled trials, and other controlled studies (controlled trials, controlled before and after studies, analytic cohorts, comparative studies) were included as determined by our eligibility criteria. CM interventions needed to be initiated either whilst as an inpatient or on discharge from acute care hospitals including the emergency department (ED), or in the community, and then continue on in the community. Only studies including adults with HF in Organisation for Economic Co-operation and Development (OECD) countries were included as the outcomes were more likely to be comparable for synthesis, and relevant to the UK situation. Studies were included as long as one of the outcomes of interest was unscheduled secondary care [unplanned hospital admissions/readmissions, ED attendance, length of hospital stay] as well as related cost and cost effectiveness of the interventions. Other outcomes of interest were primary health care resources e.g. GP visits, visits to other primary care health professionals or services and prescriptions. Studies written in any language were considered if there was an English abstract.

Reference management and study selection

EndNote and Excel were used to manage the references. Duplicates were removed from the Endnote file. References underwent a two stage process of screening using the inclusion and exclusion criteria by two reviewers independently (AH, AK, RJ). Firstly, a screen of titles and abstracts (if abstract available) and secondly screening of the full paper was conducted. Where there was continued disagreement between reviewers about including or excluding a paper, a third reviewer made the final decision (SP or RJ).

In addition to the included quantitative intervention papers, we identified relevant reviews from the search. Any potentially relevant conference proceedings were followed up, firstly by searching in Medline to see if the study had been published. If the study was not published, the authors were contacted where possible to check if the studies were likely to be published within the work frame of this review.

Data extraction

Data were extracted into a custom-designed table which included description of trial type, participants, intervention, controls, outcome measures and results. Based on the Kings Fund definition of case management we devised taxonomy of intervention components. (table 1) As part of this data extraction process, the intervention and control treatments were also described by their component parts e.g. monitoring signs and symptoms using the framework of the CM definition.

Quantitative data concerning the outcomes of interest were extracted into the Cochrane Revman software. The Cochrane risk of bias tool was used to record trial bias. This process was performed by one author and checked by a second. (AH, AK) Any disagreements were resolved by discussion and if necessary with a third author. (RJ, SP)

Version 20th December 2015

Data analysis

Trials were divided as previously described by Huntley *et al* 2013 into hospital-initiated CM and community-initiated CM.⁹ Where there were data from three or more studies, effect sizes were calculated and presented in forest plots as rate ratios (admissions) or mean differences (length of stay) using Revman software. If the heterogeneity of the combined data was greater than 50%, a random effects model was used for analysis.

We conducted sensitivity analysis in response to the risk of bias assessment of studies, removing high risk of bias studies as appropriate; the results of both analyses are presented. We conducted subgroup-analysis to explore the effects of CM duration (3,6 and 12 months plus) on hospital admission and LOS. There was insufficient detail in trials to perform sub-analysis by severity of HF or intensity of intervention.

Data were assessed narratively in respect of the components of interventions using the CM definition cited above as guidance.⁸ (table 1) In addition, where possible subgroup-analysis was conducted in Revman in which interventions with components of interest were compared with those that did not have these components.

RESULTS

The systematic review yielded 22 studies with data published over 32 papers of which 17 were RCTs and five were CTs. ¹⁴⁻⁴⁵ No relevant studies were identified in the pragmatic update search. Seventeen of these studies described hospital-initiated CM^{14,15,17,18,20-24,26-28,31,32,} and five described community-initiated CM of HF. ^{38,42-45}

Profile of patients (table 2)

The range of female participants in the trials was 1-58%, but the majority of trials had relatively even gender divide. Co and multi-morbidity were common. Eight of the 22 trials gave no detail on ethnicity of participants; in four studies the trialists used white/non-white and English speaking/non-English speaking categories. In the remaining 10 studies, a fuller profile was described. Twelve of the 22 trials were conducted in the USA and the ethnicity profile reflected that including Spanish speaking/Hispanic, American Indian, Black, African American, Asian and White participants.

The majority of trials described the severity of HF using New York Heart Association (NYHA) classification. Twelve of the trials gave a breakdown of numbers or percentages in the I-IV classes with some trials only giving numbers of participants for the III and IV class. In these trials the percentage range of III and IV class patients was 6-98%. Four trials gave mean and median values of NYHA status, one trial used the APR-DRG severity of illness scale, and five trials did not describe disease severity.

Profile of interventions (table 1)

The majority of studies (n=15) described the intervention being delivered by a case manager/ specialist nurse with no specific mention of other health professionals, and the remaining seven studies described a case manager/speciality nurse working as part of a multi-disciplinary team. All but two studies compared CM with usual care although the control group was not always described. The two remaining studies were comparative: one RCT comparing CM with specialist clinics and one RCT comparing CM with telemedicine plus CM. ^{32,42}

The duration of the CM interventions in the studies was 1-24 months with the majority having a 3 or 6 month duration. The majority of studies were conducted face to face or a combination of in-person and by phone. Four interventions were conducted purely by phone. Outcomes were measured to match the total duration of intervention in the majority of studies. For many of the studies the intensity of interventions was not stated explicitly. When intensity was described it was always a tapered approach after an initial intensive period.

Version 20th December 2015

Risk of bias (Figure 2a&b)

The degree of risk of bias was starkly different between the RCTs and CTs. All five of the CTs were rated at high risk or unknown risk for most domains. The majority of the RCTs were rated at low risk for most domains with the exception of the domain of blinding of the participants and personnel which is not applicable to this type of intervention. Three RCTs were assessed as at high risk for at least one domain: both Hancock and Wade gave no description of the randomisation process or allocation concealment, Riegel 2002 was randomised at physician level and patients were chosen by physician preference. Pour of the five community-initiated trials (2RCTs and 2CTs) were assessed to be at high risk of bias, and in some studies did not present usable data.

All the intervention studies reported unplanned hospital admissions ¹⁴⁻⁴⁵ and 17 reported number of days in hospital. ^{14, 15, 17, 18, 20-22, 24, 26-28, 35, 38, 42-45} There were few data on A and E attendance and primary care resource use. However, only some of the data could be used in meta-analysis with the main reasons being that data were presented in different formats where neither confidence intervals, standard errors nor raw data were given. Due to heterogeneity of data all analysis was conducted using a random effects model.

Unplanned HF (re)admissions data (figure 3)

Hospital-initiated CM

Thirteen of hospital-initiated CM trials had data that could be used in the meta-analysis. Overall the data showed the rate ratio of readmissions was 0.77 [95% CI 0.63, 0.94] p=0.01 I² =69% in favour of hospital- initiated CM. (figure 3a) A sensitivity analysis was conducted, removing Riegel 2000 (high risk CT) and 2002 (RCT with high risk for randomisation domain) which had a minimal effect on the rate ratio and heterogeneity 0.77 [0.61, 0.96] p=0.02 I² =68%. Sub-analysis looking at 3, 6 and 12-18 month data did not produce a clear time-related effect which is most likely due to heterogeneity within and between studies. (figure 3b,c,d). There was one hospital-initiated CM trial which compared CM with specialist clinics which reported no differences in hospital readmissions between the two groups. 32

Community initiated CM

Version 20th December 2015

Of the four community initiated trials comparing admissions between CM with usual care, two reported no significant differences^{38,43} and two reported statistically significant reductions in favour of CM. Data were available from three of these trials for meta-analysis and showed no significant differences in admissions between groups, (rate ratio 1.08 [0.62,1.87] p=0.8 I²=78% (figure 3e). One further trial compared CM, with telehealth and CM and reported no differences in admissions but data were not presented. Description

Length of hospital stay (LOS) (figure 4)

Hospital-initiated CM

Nine of the hospital-initiated CM trials had data that could be used in a meta-analysis. Overall the data showed that LOS was reduced in the CM group compared to usual care mean difference (MD) - $1.28 \text{ days} [-2.03, -0.53] \text{ p=}0.0008 \text{ l}^2 = 68\%$. (figure 4a) A sensitivity analysis was conducted removing Riegel 2000 and 2002 which had an important effect on the rate ratio and heterogeneity MD -1.76 [-2.29, -1.23] p<0.00001 $\text{l}^2 = 14\%$. Sub-analysis looking at 3, 6 and 12-18 month data suggests this effect is short-term (first 3 months) but a longer time-related effect was difficult to assess due to lack of data.(figure 4b,c,d).

The one study comparing CM with specialist clinic care, reported that CM patients accumulated 592 and clinic patients 547 *all-cause* hospitalizations (p = 0.087) associated with 3067 versus 4410 days of hospital stay (p < 0.01 for rate and duration of hospital stay).³²

Community initiated CM

Of the 4 community initiated trials comparing CM with usual care, two did not report LOS ^{38, 43} one reported median values in favour of CM⁴⁴ and the remaining one reported a mean reduction in LOS. ⁴⁵ (table 2). The one comparative trial between community-initiated CM, and telehealth and CM did not report any useful data.⁴²

Intervention components (table 2 and appendix 2)

Fourteen intervention components were identified and grouped as per the CM definition in methods and prevalence determined for both hospital and community –initiated CM studies with a usual care control group (table 2). ⁷

Version 20th December 2015

Hospital-initiated CM (appendix2)

Data from individual studies which contained components of either family involvement or education/self-management showed an overall reduction in hospital readmissions in comparison with usual care. They also showed a reduction in hospital readmissions observationally in comparison with interventions which did not contain these components. However, analysis comparing the studies in which the component was present with those in which the component was absent did not yield any statistically significant differences. (appendix 2a &b) The same calculations for medication review, referral to other services, and assessment of home environment, CM meetings and patient-directed access did not indicate any specific effect of these components of hospital- initiated CM on rates of admission. (appendix 2 c-e)

Community initiated CM

There were insufficient data to conduct any subgroup-analysis on any of the remaining components of hospital-initiated CM, community initiated studies or the length of stay data.

Outpatient health care resources

Only six of the included studies measured outpatient resource use. In some studies, outpatient resource data were all-cause and not HF-specific. In some studies primary and secondary use was combined. ^{23,24,35,38,42,45} Two of these studies also reported ED attendance. ^{23,42} All but one of these studies reported no difference between intervention and control group for these measures with the exception of Lowery which showed a statistically significant greater use of outpatient resources in the usual care group (optional primary care visits 1yr 16.75(13.62),10.43(9.6) p<0.001, 2yr 14.27(11.98),9.35(9.97) p<0.001). ⁴⁵

Costs

Nine of the seventeen hospital initiated trials described cost comparison data. Of these, six reported no statistically significant difference between CM (3 or 6 month duration) and usual care. ^{17,18,22,24,35,26} and three reported costs in favour of CM although data from Stauffer was brief. ^{15,32,37} One of these was 12-18 months ³² and two were 3 months in duration. It was difficult from the intervention

descriptions to determine their intensity. There were no cost data reported from the community-initiated trials.

DISCUSSION

 This systematic review confirms that hospital-initiated CM can be successful in reducing unplanned hospital readmissions, and reducing length of stay in hospital in the short term for people with HF. There were only five community-initiated CM studies of which four were at high risk of bias. Limited data shows no effect of community-initiated CM on hospital admissions. A minority of trials report cost comparisons with usual care and most of those show no difference. There were limited data on the effect of CM on other health care resources. Observational intervention component data suggests that care providing family involvement and education/self-management are likely to be important in case management but these observations did not stand when subjected to subgroup analysis.

Many factors are likely to modify the effect of CM on use of emergency care seen in these studies. It is generally accepted that CM is more appropriate for people with severe HF and poorer general health. However it was difficult to compare the health status of the study participants in both hospital-initiated and community-initiated trials as in some studies there was little detail, others gave median and mean figures for NYHA status, and the presentation format and detail of co-morbidities varied.

Seventeen studies described hospital-initiated CM and five described community initiated CM of heart failure although often the participants were identified via hospital clinic records. Overall the meta-analysis showed that a positive effect of CM on readmissions and hospital length of stay. This may be explained by the fact that in most of the trials the participants were identified via hospital contact and therefore were likely to have had a recent exacerbation of their HF and to be at increased risk of readmission in the post-discharge period. In addition, it is likely that interventions are acting at a time of highest risk as reflected by HF mortality in first year of diagnosis. Therefore once they were assessed and given extra support they were stable for a period of time. Previous work by Roland et al 2005 suggests that admission rates in people aged 65 with two or more emergency admissions in 12 months fall in subsequent years without any intervention and account for fewer than 10% of admissions in the following year and thus effectiveness of admission avoidance schemes cannot be judged by tracking admission rates without careful comparison with a control group. The data from

Version 20th December 2015

trials of community-initiated CM was lacking both in the number of studies, and the fact there were limited useable data that showed no effect on unplanned hospital admissions. It is likely that these patients were likely to be in more stable health.⁴¹

BMJ Open

A meta-review of a wide range of HF disease management programs by Savard 2011 reports that nine previous systematic reviews (2001-2009) identified significant reductions in HF admissions with reductions in risk ranging from 30-56%. However the authors caution that these reviews are limited by inadequate reporting in the population, setting, intervention and comparator components. They report that reviewers have not taken into account statistical, clinical and methodological heterogeneity in interventions. Our review focussed specifically on CM avoiding some of these limitations and indicates a reduction in HF admissions with hospital-initiated CM in the range of 10-30%.

Wakefield *et al* in 2013 looked at common components of a range of HF care programs focusing mainly on disease management and education investigated in RCTs, and 10/35 of the discussed studies were included in our review.⁴⁸ They described patient education, symptom management by health professionals and by patients, and medication adherence strategies as the most commonly occurring elements of care. A literature review by Jaarsma 2012 looked at 70 'home care' controlled studies (mostly RCTs) which encompassed 9 of our included CM studies covering a wide range of approaches such as telemedicine, hospital at home and health buddies for patients with HF.⁴⁹ They identified a multi-disciplinary team, continuity of care, care plans, optimising titration of medication, education/counselling of patients and caregivers and increased access as important.

In order to understand why our meta-analysis reported that the specific approach of CM was successful in reducing admissions and length of stay we also conducted component analysis. Our analysis agreed with the above reviews that patient education/self-management is significant but also flagged up the importance of family involvement. The magnitude of effect was not so convincing for medication review (appendix 2c) but it was difficult from the study description to distinguish between medication management and formal reviewing of medication by either the CM or pharmacist.

Previous systematic reviews have investigated the role of the lay caregiver in HF patient management. 50-52 These suggest that better relationship quality and communication were related to reduced mortality, increased health status and less distress and improved patient self-care outcomes.

Our review adds to this evidence base by suggesting that more family involvement in CM may also reduce unscheduled secondary care.

Education about HF and about its pharmacological and non-pharmaceutical treatment has been well-reviewed both as an individual approach and as part of complex interventions, and is considered to be essential for improving many patient outcomes. ^{49, 53,54}A recent mixed method study, suggests asking patients with HF to write down their learning needs before the education increases their chances of receiving education based on their individual needs. ⁵⁵ Qualitative interviews with health professionals caring for HF patients suggest that communication with, and education by specialist nurses facilitated by continuity of care is essential to good care of HF patients. The authors also highlight the role of the specialist nurse in multi-disciplinary team communication and functioning; essentially describing the role of the specialist nurse as a case manager. ⁵⁶

Our review of CM suggests that the evidence for its cost-effectiveness is lacking with most studies that have performed cost comparisons with usual care show no advantage. Previous work by de Bruin et al 2001 looked at cost effectiveness of disease management for a range of chronic conditions and concluded that the data is most positive for HF with 5 out of the 8 included studies showing cost-effectiveness.⁵⁷

Strengths and limitations

The contribution of our high quality systematic review to the above is that we have focused on CM which is based on nurse co-ordinated multi-component care of patients which is applicable to the primary care based health systems such as that in the UK. We have focused on HF admissions and length of stay as opposed to all-cause data which many of the previous reviews have used.

By examining the components of CM we have a profile of the components most likely to lead to the success of CM of patients with heart failure in terms of reducing admissions and hospital length of stay. Our review has high-lighted the potential importance of family involvement and education/self-management.

The limitations of this review are that majority of the community-initiated CM studies were of low quality with the exception of one low risk RCT, and provided limited data for meta-analysis. This was counteracted by the fact that the hospital-initiated studies comprised of predominantly community-

Version 20th December 2015

based case management. There is a lack of cost data and analysis in the included papers. This point needs to be emphasised for future trials. It is possible that cost effectiveness will be more likely with intervention for patients with more severe HF.

CONCLUSIONS

Hospital-initiated CM reduces unplanned hospital admissions, and length of stay for people with HF in the short term. Cost data is limited. There were limited data for community-initiated CM which suggested it does not reduce hospital admission. Further research is needed to determine the individual components of CM that contribute to reduced admissions.

Contributorship statement

Alyson Huntley. Main systematic reviewer, worked across all stages of the review from inception to completed draft.

Rachel Johnson. Cardiology and primary care expertise. Worked on screening, selection of studies, commenting on analysis, and development and checking of final document content.

Anna King. Second reviewer, involved in screening, selection, data checking and commenting on developing and final document content.

Richard Morris. Statistical expertise, advising on data analysis and commenting on the developing and final document content.

Sarah Purdy. Primary care and admission avoidance expertise. Advised throughout project, third reviewer for screening process and commenting on the developing and final document content.

Competing interests

None declared

Funding

This project was funded by the National School of Primary Health Care project no.238.

Data sharing statement

Full data extraction tables and data analysis files are available on request

REFERENCES

- Cardiovascular disease statistics 2014 British Heart Foundation accessed at https://www.bhf.org.uk/~/media/files/publications/research/bhf_cvd-statistics-2014_web_2.pdf on the 7/10/15.
- Quality and Outcomes Framework Achievement, prevalence and exceptions data 2012/13. 29
 October 2013. Accessed at http://www.hscic.gov.uk/catalogue/PUB12262/qual-outc-fram-12-13-rep.pdf on the7/10/15
- de Giuli F, Khaw KT, Cowie MR, et al. Incidence and outcome of persons with a clinical diagnosis of heart failure in a general practice population of 696,884 in the United Kingdom. Eur J Heart Fail 2005; 7(3):295–302.
- 4. Stewart S, MacIntyre K, Hole DJ *et al.* More 'malignant' than cancer? Five-year survival following a first admission for heart failure. *Eur J Heart Fail* 2001; 3:315–322.
- Stewart S, Ekman I, Ekman T et al. Population impact of heart failure and the most common forms of cancer: a study of 1 162 309 hospital cases in Sweden (1988 to 2004). Circ Cardiovasc Qual Outcomes 2010; 3:573–580.523.
- Jhund PS, Macintyre K, Simpson CR et al. Long-term trends in first hospitalization for heart failure and subsequent survival between 1986 and 2003: a population study of 5.1 million people. Circulation 2009; 119:515–523.
- Ross S, Curry N, Goodwin N. Case Management. What Is It and How It Can Best Be Implemented. London, UK: The Kings Fund, 2011. Accessed at http://www.kingsfund.org.uk/publications/case_management.html on the7/10/15.
- Department of Health. Supporting People with Long Term Conditions. Liberating the Talents
 of Nurses Who Care for People with Long Term Conditions. London, UK: Department of
 Health, 2005. Accessed at
 www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH
 _4102469 on 01 /12/11.
- Huntley AL, Thomas R, Mann M et al. Is case management effective in reducing the risk of unplanned hospital admissions for older people? A systematic review and meta-analysis. Fam Pract 2013; 30(3):266-75

- 10. Purdy S. Interventions to reduce unplanned hospital admission: a series of systematic reviews. Report accessed at http://www.bristol.ac.uk/medialibrary/sites/primaryhealthcare/migrated/documents/unplannedadmissions.pdf on 07/10/15.
- List of OECD Member countries accessed at http://www.oecd.org/about/membersandpartners/list-oecd-member-countries.htm on 07/10/15.
- 12. Chapter 8: Assessing risk of bias in included studies accessed at http://handbook.cochrane.org/chapter_8/8_assessing_risk_of_bias_in_included_studies.htm on 07/10/15.
- Chapter 9- 9.7 Sensitivity analyses accessed at http://handbook.cochrane.org/chapter_9/9_7_sensitivity_analyses.htmon07/10/15.
- 14. Rich, Vinson JM, Sperry JC *et al.* Prevention of readmission in elderly patients with congestive heart failure: results of a prospective, randomized pilot study. *J Gen Intern Med* 1993; 8(11): 585-90. **MAIN PUBLICATION**
- Rich MW, Beckham V, Wittenberg C et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. NEJM 1995; 333(18): 1190-5.

MAIN PUBLICATION

16. Rich MW, Gray DB, Beckham V *et al.* Effect of a multidisciplinary intervention on medication compliance in elderly patients with congestive heart failure. *Am J Med* 1996; 101 (3): 270-6.

EXTRA

 Stewart S, Pearson S, Luke CG et al. Effects of home-based intervention on unplanned readmissions and out-of-hospital deaths. J Am Geriatr Soc 1998; 46(2):174-80. MAIN PUBLICATION

- 18. Stewart S, Marley JE, Horowitz JD. Effects of a multidisciplinary, home-based intervention on unplanned readmissions and survival among patients with chronic congestive heart failure: a
- 19. Stewart S, Horowitz JD. Home-based intervention in congestive heart failure: long-term implications on readmission and survival. *Circulation* 2002; 105 (24): 2861-6. **EXTRA**

randomised controlled study. Lancet 1999; 354 (9184): 1077-83. MAIN PUBLICATION

20. Blue L, Lang E, McMurray *et al.* Randomised controlled trial of specialist nurse intervention in heart failure. *BMJ* 2001; 323(7315): 715-718. **MAIN PUBLICATION**

- 21. Riegel B, Carlson B, Kopp Z *et al.* Effect of a standardized nurse case-management telephone intervention on resource use in patients with chronic heart failure. *Arch Intern Med* 2002; 162(6):705-12. **MAIN PUBLICATION**
- 22. Laramee S, Levinsky SK, Sargent J et al. Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. Arch Intern Med 2003; 163(7): 809-17.
 MAIN PUBLICATION
- 23. DeBusk RF, Miller NH, Parker KM *et al.* Care management for low-risk patients with heart failure: a randomized, controlled trial. *Ann Intern Med* 2004; 141(8):606-13. **MAIN**PUBLICATION
- 24. Naylor MD, Brooten DA, Campbell RL *et al.* Transitional Care of Older Adults Hospitalized with Heart Failure: A Randomized, Controlled Trial. *J Am Geriatr Soc* 2004; 52(5):675-684.

 MAIN PUBLICATION
- McCauley KM, Bixby MB, Naylor MD. Advanced practice nurse strategies to improve outcomes and reduce cost in elders with heart failure. *Dis Manag* 2006; 9(5):302-10. EXTRA
- 26. Riegel B, Carlson B, Glaser D, Romero T.
 Randomized controlled trial of telephone case management in Hispanics of Mexican origin wit h heart failure. J Card Fail 2006; 12(3):211-9. MAIN PUBLICATION
- Thompson DR, Roebuck A, Stewart S. Effects of a nurse-led, clinic and home-based intervention on recurrent hospital use in chronic heart failure. Eur J Heart Failure 2005; 7(3): 377-384. MAIN PUBLICATION
- 28. Jaarsma T, van der Wal MH, Lesman-Leegte I *et al.* A Coordinating Study Evaluating
 Outcomes of and I. Counseling in Heart Failure Effect of moderate or intensive disease
 management program on outcome in patients with heart failure: Coordinating Study
 Evaluating Outcomes of Advising and Counseling in Heart Failure (COACH). Arch Intern Med
 2008; 168(3): 316-24. MAIN PUBLICATION
- 29. Postmus D, Pari AA, Jaarsma T *et al.* A trial-based economic evaluation of 2 nurse-led disease management programs in heart failure. *Am Heart J* 2011; 162(6): 1096-104. **EXTRA**
- 30. Inglis SC, Pearson S, Treen S *et al.* Extending the horizon in chronic heart failure: effects of multidisciplinary, home-based intervention relative to usual care. *Circulation* 2006; 114 (23): 2466-73. **EXTRA**

- 31. Brotons C, Falces C, Alegre J *et al.* Randomized clinical trial of the effectiveness of a home based intervention in patients with heart failure: the IC-DOM study. *Rev Esp Cardiol* 2009; 62(4):400-8. **MAIN PUBLICATION**
- 32. Stewart S, Carrington MJ, Marwick TH et al. Impact of home versus clinic-based management of chronic heart failure: the WHICH? (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care) multicenter, randomized trial. J Am Coll Cardiol 2012; 60(14);1239-48. MAIN PUBLICATION
- 33. Stewart S, Carrington MJ, Horowitz JD et al.
 Prolonged impact of home versus clinicbased management of chronic heart failure:
 extended follow-up of a pragmatic, multicentre randomized trial cohort. Int J Cardiol 2014;
 174(3):600-10. EXTRA
- 34. Stewart S, Carrington MJ, Chan YK. Prolonged benefits of nurse-led, home-based intervention versus a specialist heart failure clinic: Extended follow-up of the WHICH? Trial Cohort. *Eur J Heart Failure* 2014; 16(106) . **EXTRA**
- 35. Riegel B, Carlson B, Glaser D *et al.* Which patients with heart failure respond best to multidisciplinary disease management? *J Cardiac Fail* 2000; 6 (4): 290-9. **MAIN**PUBLICATION
- 36. Russell D, Rosati RJ, Sobolewski S *et al.* Implementing a transitional care program for highrisk heart failure patients: findings from a community-based partnership between a certified home healthcare agency and regional hospital. *J Healthcare Qual* 2011; 33(6):17-23. **MAIN PUBLICATION**
- 37. Stauffer BD, Fullerton C, Fleming N et al.
 Effectiveness and cost of transitional care program for heart failure:
 a prospective study with concurrent controls. Arch Intern Med 2011; 171(14):1238-43. MAIN
 PUBLICATION
- 38. Peters-Klimm F, Campbell S, Hermann K *et al.* Case management for patients with chronic systolic heart failure in primary care: The HICMan exploratory randomised controlled trial.

 Trials 2010; 11: 56. MAIN PUBLICATION
- 39. Peters-Klimm F, Muller-Tasch T, Schellberg D *et al.* Rationale, design and conduct of a randomised controlled trial evaluating a primary care-based complex intervention to improve

- the quality of life of heart failure patients: HICMan (Heidelberg Integrated Case Management).

 BMC Cardiovascular Disorders 2007: 7, (25). PROTOCOL
- 40. Freund F, Baldauf A, Muth C *et al.* Practice-based home visit and telephone monitoring of chronic heart failure patients: rationale, design and practical application of monitoring lists in the HICMan trial *Zeitschrift fur Evidenz Fortbildung und Qualitat im Gesundheitswesen* 2011; 105 (6): 434-45. **EXTRA**
- 41. Peters-Klimm F, Campbell S, Muller-Tasch T et al. Primary care-based multifaceted, interdisciplinary medical educational intervention for patients with systolic heart failure: lessons learned from a cluster randomised controlled trial. *Trials* 2009; 10:68. EXTRA
- 42. Wade MJ, Desai AS, Spettell CM *et al.* Telemonitoring with case management for seniors with heart failure. *Am J Manag Care* 2011; 17(3): e71-9. **MAIN PUBLICATION**
- 43. Hancock HC, Close H, Mason JM et al. Feasibility of evidence-based diagnosis and management of heart failure in older people in care: a pilot randomised controlled trial. BMC Geriatr 2012 Nov 14;12:70.
- 44. Bonarek-Hessamfar M, Benchimol D, Lauribe P et al.
 Multidisciplinary network in heart failure management in a community
 based population: results and benefits at 2 years. Int J Cardiol 2009 May 1;134(1):120-2.
- 45. Lowery J, Hopp F, Subramanian U et al. Evaluation of a nurse practitioner disease management model for chronic heart failure: a multi-site implementation study. Congest Heart Fail 2012; 18(1):64-71.
- 46. Roland M, Dusheiko M, Gravelle H *et al.* Follow up of people aged 65 and over with a history of emergency admissions: analysis of routine admission data. *BMJ* 2005; 330(7486):289-92.
- 47. Savard LA, Thompson DR, Clark AM.
 A metareview of evidence on heart failure disease management programs:
 the challenges of describing and synthesizing evidence on complex interventions. *Trials* 2011;
 12:194.
- 48. Wakefield BJ, Boren SA, Groves PS *et al.* Heart failure care management programs: a review of study interventions and meta-analysis of outcomes. *J Cardiovasc Nurs* 2013; 28(1):8-19
- 49. Jaarsma T, Brons M, Kraai I *et al.* Components of heart failure management in home care; a literature review. *Eur J Cardiovasc Nurs* 2013; 12(3):230-41.

Version 20th December 2015

- 50. Hooker SA, Grigsby ME, Riegel B *et al.* The Impact of Relationship Quality on Health-Related Outcomes in Heart Failure Patients and Informal Family Caregivers: An Integrative Review. *J Cardiovasc Nurs* 2015;30(4 Suppl 1):S52-63.
- 51. Buck HG, Harkness K, Wion R *et al.* Caregivers' contributions to heart failure self-care: a systematic review. *Eur J Cardiovasc Nurs* 2015; 14(1):79-89.
- 52. Clark AM, Spaling M, Harkness K *et al.* Determinants of effective heart failure self-care: a systematic review of patients' and caregivers' perceptions. *Heart* 2014;100(9):716-21.
- 53. Spatola CF, Cocchieri A, De Marinis MG *et al.* Educational interventions in patients with heart failure: a review of the literature. Article in Italian *Ig Sanita Pubbl* 2013; 69(5):557-74.
- 54. Boyde M, Turner C, Thompson DR *et al.* Educational interventions for patients with heart failure: a systematic review of randomized controlled trials. *J Cardiovasc Nurs* 2011; 26(4):E27-35.
- 55. Ross A, Ohlsson U, Blomberg K *et al.* Evaluation of an intervention to individualise patient education at a nurse-led heart failure clinic: a mixed-method study. *J Clin Nurs* 2015; 24(11-12):1594-602.
- 56. Glogowska M, Simmonds R, McLachlan S et al. Managing Patients With Heart Failure: A Qualitative Study of Multidisciplinary Teams With Specialist Heart Failure Nurses. Ann Fam Med 2015; 13(5):466-71.
- 57. de Bruin SR, Heijink R, Lemmens LC *et al.* Impact of disease management programs on healthcare expenditures for patients with diabetes, depression, heart failure or chronic obstructive pulmonary disease: a systematic review of the literature.

 Health Policy 2011; 101(2):105-21.



Table 1: Components of CM interventions

Definition & total prevalence of Components of CM interventions	Number of hospital- initiated CM vs. usual care with component present (total studies=16)	Number of community-initiated CM vs. usual care with component present (total studies=4)
Assessment /Evaluation		
Monitoring signs & symptoms (n=18) Encompasses general care of CHF patients which is likely to include establishing a relationship with patient over visits, physical and cardiac status checking, lifestyle assessment, general medication check and screening tests e.g. depression, dementia	14	4
Medication review	6	2
(n=8) Review and adjustment of medication by experienced case manager (nurse), pharmacist, GP or consultant often using a		
combination of these health professionals.		
Assessment of home environment (n=4) Assessment carried out by case manager to identify any issues or potential issues with home environment e.g. stairs	4	0
Planning		
CM meetings/feedback to other HPs	3	2
(n=5) Group meetings of health professionals involved in CHF patients care with the aim of reporting on and planning for patients care.		
Appointment organisation	2	0
(n=2) Case manager checking medical appointments, ensuring ability to go etc.		
Advance care planning (n=1)	1	
Facilitation	-	
Education/self-management	15	3
(n=18) Educating CHF patients about their condition, treatment and what to expect. The aim of this is to assist self-management (care with assistance of health professionals) and self-care (patient engaging in activities to promote their health and well-being).		
Patient-directed access (n=6) The ability of CHF patients to initiate care from the case manager or case management service.	6	0
Care co-ordination		
Referral to (n=14) When the case manager refers the patient to other health or social care professionals, this can be GP hospital consultant, social care or tests.	11	3
Advocacy for options & services		
Equipment (n=4)	3	1
Provision of items to assist patient's health care such as pill counters, weighing scales and measured water bottles. Physical therapy	1	0
(n=1) CHF patient receiving physical therapy/rehabilitation		
Support group (n=1) CHF attending or being offered the opportunity of a support group.	1	0
Other		
Family involvement (n=9) When the patient's family in terms of information, education or involvement e.g. goal setting in	8	1
patients' care or active monitoring Emotional support (n=1)	0	1
Case manager providing emotional support to CHF patient.		

Version 20th December 2015

Table 2: study characteristics of intervention studies


	•		
Study n=randomised	Baseline characteristics of participants:	Intervention	Control
Recruitment/setting	CM vs. usual care		
Hospital Initiated CM -			
RCTs			
Rich ¹⁴	Age 80(6.3),77.3(6.1) yrs p=0.04	Non-pharmacological comprehensive	Usual care (n=35)
1993 USA	% Female: 60.3, 47.1%	multidisciplinary treatment strategy (n=63)	Components of intervention: Visits by
	Ethnicity:	, ,	home nurse
n=98 randomised	White 46,57.1% Disease Status:	Components of intervention at home: Monitoring signs& symptoms	
Patients ≥70 yrs admitted	Mean NYHA status	Medication review (nurse)	
to medical wards of Jewish	2.7(1.1),3.0(1.0)	Education /Self-management support	
Hospital at Washington University Medical Centre		Assessment of home environment Patient directed access to study	
were screened for		personnel	
congestive HF.	Acc. 90 1/5 0) 79 4/6 1) yrs	Nurse directed multidisciplinary	Herrel core (p=140)
1995	Age: 80.1(5.9),78.4(6.1) yrs p=0.02	Nurse-directed multidisciplinary intervention (n=142)	Usual care (n=140)
USA	% Female:68,59%	As above for Rich 1993	As above for Rich 1993
n=285 randomised	Ethnicity: Non-white race 52.59%		
As above for Rich 1993	Disease Status:		
•	Mean NYHA class 2.4(1.0),2.4(1.1)		
Stewart 17	Age: 76 (11), 74(10) yrs	Home-based intervention (n=49)	Usual care (n=48)
1998 Australia	% Female:55,48% Ethnicity: Non-English speaking	Components of intervention at home:	Components of intervention:
	20.4,18.75%	Monitoring signs& symptoms	Disease management
n=97 randomised Patients were recruited	Disease Status: NYHA II 49,50%	Education (pharmacist)/self-management support	
whilst admitted to a large	III 47,42%	Medication review (pharmacist)	
tertiary hospital	III 4,4%	Referral to GP Family involvement	
		Equipment	
Stewart 18,19	Age: 75.2(7.1),76.1(9.3)	Multidisciplinary home base	Usual care (n=100)
1999	% Female: 35,41% Ethnicity: Primary language not	intervention (n=100)	Components of intervention:
Australia	English 32,32	Components of intervention at home:	Contact with other health and social
n=200 randomised	Disease Status: NYHA II 42,48	Monitoring signs& symptoms Referral to other health & social care	professionals Appointment with GP or cardiac clinic or
Patients admitted to a	III 46,43	Appointment organisation	both
tertiary referral hospital	IV 12,9	Assessment of home environment Family involvement	
		Education/self-management support	
		Medication review (nurse/GP/cardiologist)	
Blue ²⁰	Age: (SD)74.4(8.6),75.6(7.9)yrs	Specialist nurse intervention (n=82)	Usual care (n=75)
2001	% Female: 36,49%		
UK	Ethnicity: Not reported Disease Status: NYHA	Components of intervention at home: Monitoring signs& symptoms	Components of intervention: GP care
n=165 randomised	II 19(23), 16(20)	Education/Self-management support	GI care
Patients admitted as an	III 28(34), 33 (42) IV 36 (43), 30(38)	Referral to other health & social care Appointment organisation	
emergency to the acute	Co or Multi-morbidity:	Medication review (nurse, cardiologist)	
medical ward of the hospital	Angina 40(49), 38(45) Past MI 41 (51), 46(55)		
Поорна	Diabetes 15(19), 15(18)		
	Chronic lung disease 18(22),23(27)		
	Hypertension 42(52), 36(43)		
	AF 42(52), 29(35) Valve disease 12(15),15(18)		
Riegel ²¹	Age: 72.52(13.05), 74.63(12)	Telephonic case management (n=130)	Usual care (n=228)
2002 USA	% Female:46.2,53.9 Ethnicity:(Primary language)	Components of intervention at home:	Components of intervention: Not known
	English 91(70), 168(73.7)	Monitoring signs& symptoms	
n=281 physicians randomised	Spanish 35 (26.9), 58(25.4) Disease Status: NYHA	Self-management support Referral to other HPs (includ.GP) &	
	II 2.3,3.6	social care	
Patients admitted at 2 Southern California	III 35.9, 38.4 IV 61.7,58.0	Family involvement	
hospitals.			
Laramee ²² 2003	Age: 70.6(11.4), 70.8(12.2) yrs. % female:	Case management (n=131 data available)	Usual care (n=125 data available)
USA	42,50%	,	Components of intervention: Not
n=287 randomised	Ethnicity: Not reported	Components of intervention at home: Monitoring signs& symptoms	known
	Disease Status (SD):	Education/Self-management support	
Patients admitted to hospital for CHF were	NYHA I 10(7),35(26)	Family involvement Equipment	
screened.	II 76(55),47(36)	Patient–directed access to CM	
	III 50(36),46(35) IV 3(2),4(3)		
	Note P<0.001		

Version 20th December 2015

De Busk ²³	Age: <60yrs 15,14%,	Case management (n=228)	Usual care (n=234)
2004 USA n=462 randomised	6-70yrs 22,24%,70-80yrs 40,37%, >80yrs 21,26% % Female:52,45% Ethnicity:	Components of intervention at home: Monitoring signs& symptoms Education /Self-management support	Components of interventions: not known
Patients who were admitted with a provisional diagnosis of HF from Kaiser Permanente medical centres in California	American Indian 0,1% Asian 4,8% Black 2,2% White 5,6% Hispanic 3,3% Disease Status: NHYA	CM meetings/feedback to other health providers	
	I-II 50,50% III-IV 50,50%		
Naylor ^{24,25} 2004	Age: 76.4(6.9),75.6(6.5) % Female:60, 56%	Transitional care intervention with advanced practice nurses (APNs)	Usual care (n=121)
USA n=239 patients randomised Patients aged 65yrs+ admitted to 6 study hospitals from home with a diagnosis of HF were screened for participation.	Ethnicity: African American 34,38%, White 66, 62% Disease Status: Functional Status (Moinpur C 1992) Personal 17.1(5.8),16.9(5.8) Social 8.4(2.6),8.6(2.6) Total 25.5(8),25.4(7.8)	(n=118) Components of intervention at home: Monitoring signs& symptoms Education/Self-management support Family involvement CM meetings/feedback to other health providers	Components of intervention: Care from standard home care services Patient–directed access to home car services
		Patient-directed access to CM	
Riegel ²⁶	Age: 71.6910.8),72.7(11.2)	Telephonic case management (n=69)	Usual care (n=65)
2006 San Diego USA n=135 randomised Self-identified Hispanics were identified at 2 community hospitals close	% Female:58,49.2% Ethnicity: Hispanic patients Speak/read only Spanish 60.9,65.1% Disease Status: NYHA II 17.4,20% III 44.9.47.7% IV 37.7,32.3%	Components of intervention at home: Monitoring signs& symptoms self-management support Referral to other HPs (includ.GP) & social care Family involvement	Components of intervention: Disease management Information
to US-Mexico border			
Thompson ²⁷ 2005 UK	Age: 73(14),72(12) % Female:38,27% Ethnicity:No details	Clinic & home based intervention (n=58)	Usual care (n=48) Components of intervention at home:
Randomisation was at GP practice level	Disease Status: NYHA III &IV 76,73%	Components of intervention at home: Monitoring signs& symptoms Education/self-management support Family involvement	unknown
Patients recruited from 2 North of England general hospitals following an admission		In outpatient clinic Monitoring signs& symptoms Education/self-management support Family involvement Referral to other health and social care	
Jaarsma ^{28,29} 2008	Age: 71(11),70(12),72(11) % Female:34,39,40%	Basic nurse support (n=340)	Usual care group (n=339)
The Netherlands	Ethnicity: No detail Disease Status: NYHA II 51,48,54%	Components of intervention: Outpatients Education /Self-management support	Components of intervention: Disease management
n=1049 randomised All patients had been admitted to hospital with	III 47,48,42% IV3,4,4%	Patient directed access to HF nurse Intensive nurse support (n=344)	
symptoms of HF		Components of intervention at home: Patient directed access to HF nurse Referral to other health & social care Education/Self-management support Equipment	
Brotons ³¹	Age: 76.6(7.5),76(8.9)yrs.	Home based intervention (n=144)	Usual care (n=139)
2009 Spain n=283 randomised Patients were recruited by well-trained nurses at 2 university hospitals	% Female:54.2,56.1% Ethnicity: Not reported Disease Status: NHYA 142.4,55.4% II 52.1,37.4% III 4.9,5.8% IV 0.7,1.4%	Components of intervention at home: Monitoring signs& symptoms Education/Self-management support Medication review (nurse, physician, cardiologist) Referral to physician or cardiologist as necessary	Components of intervention: not known
Stewart 32	Home vs. Clinic	Home based intervention (n=143)	Clinic-based intervention (n=137)
2012 WHICH trial Australia	Age: 70 (15), 73(13) yrs % Female:27,28% Ethnicity: No details Disease Status: NYHA II or III	Components of intervention at home: Monitoring signs& symptoms Family involvement	Components of intervention: In clinic Disease management
n=280 randomised	83,88%	CM meetings/feedback to other health providers	Assessment of home environment
Patients admitted to participating hospitals were screened for study eligibility	Months since CHF diagnosis 34.6(55.3),44.8(71.0)	providers Referral to other health or social care Assessment of home environment Medication review (nurse, pharmacist, physician, cardiologist)	Family involvement? Referral to other health or social care CM meetings/feedback to other health providers
Hospital Initiated CM - CTS			
Riegel ³⁵	Age: 74.44 yrs. (10.65), 70.77	Multidisciplinary disease management	Usual Care (n=120)
2000 USA n=240 were randomised Patients were recruited	(11.77) % female:55, 55% Ethnicity: No details Disease Status: NYHA I 19.2,24.2%	(DM) (n=120) Components of intervention at home: Monitoring signs& symptoms Support group	Components of intervention at home Disease management
from 5 hospitals following a hospitalisation for HF.	II 26.7,18.3% III 43.3,44.2% IV 10.8,13.3%	Referral to specialist RN visits	

Russell ³⁶	Age: 79.4(10.7),79.9(10.7)	Transitional care service (n=223)	Usual home care services (n=224)
2011	% Female:55.6,57.6 (numbers)	Components of interpreting at he con	Components of inter
USA n=447	Ethnicity: White non-Hispanic 56.9,58.4	Components of intervention at home: Self-management support	Components of intervention at home: Nurse visits
11-771	African-American 17.0,16.5	Referral to other health & social care	Physical therapy (44.6)
Patients were referred from	Hispanic 14.8,14.	Assessment of home environment	Home health aide service (27.7)
a single large not-for-profit	Asian/other 11.2,10.7	CM meetings/feedback to other health	,
general medical and	Disease Status: Patients with a	providers	
surgical hospital	primary or secondary diagnosis of CHF	Advance care planning	
Stauffer ³⁷	Age: 78.9(8.3), 81.4(8.3)	Physical therapy Nurse-led transitional care	Control group (n=84)
2011	% Female:58.1, 54.8%	intervention (n=56)	Control group (ii 64)
USA	Ethnicity: Hispanic ethnicity		Components of interventions:
	7.1,3.6%	Components of intervention at home:	unknown
n=140	Disease Status:	Monitoring signs& symptoms Education/Self-management support	
Patients were screened for eligibility within 48 hours of	APR-DRG severity of illness 1 5.4,1.2%	Family involvement	
hospital admission	2 44.6,31%	Referral (Assessing availability of social	
	3 37.5,57.1%	care)	
	4 12.5,10.7%	Patient directed access to study	
		personnel	
Community-initiated CM- RCTs			
Peters-Klimm ³⁹	Baseline characteristics of	Case management (n=97)	Usual care (n=100)
2010	participants: CM,UC	- acc management (ii or)	
Germany	Age: 70.4yrs (10.0), 68.9 (9.7)	Components of intervention at home:	Components of control intervention:
•	% Female:29, 26%	Monitoring signs& symptoms	Disease management
n=100 at randiti	Ethnicity: No details Disease Status: NYHA	Education/Self-management support	Education
n=199 at randomisation	1 1(1.0), 5 (5)	Medication review (CM/GP) Referral to GP	
Recruitment was via	II 63 (64.9),67 (67)	relevanto or	
general practice by mail.	III 33 (34),27 (27)		
	IV 0, 1(1.0)		
	Mean yrs. with CHD		
Wade ⁴²	6.2(4.6) (n=79),6.8(6.3)(n=74 Age: 75.8, 77.7 yrs.	Case management (n=152)	Tele health with case management
2011	% female: No detail	ouse management (ii 102)	(THCM) (n=164)
USA	Ethnicity: Black/African	Components of intervention at home:	
n=2,200 were randomised	American	Referral to other health & social care	Components of intervention:
Aetna Medicare Advantage	24,20.4% Disease Status (SD):	Equipment	Disease management Education
members with medical &	No detail		Referral to other health & social care
pharmacy benefits were	. To dota		Troibinar to out of mountain a coolar saile
identified through analysis			
of claims	A	0	B. C. OBlatan (v. 40)
Hancock ⁴³ 2012	Age: 85.1 (6.7), 81.8 (7.1)yrs % Female:56%, 58%	Case management (n=16)	Routine GP-led care (n=12)
UK	Ethnicity: 100% white British	Components of interventions at home:	Components of intervention:
n=28 randomised	Disease Status:	Monitoring signs and symptoms	Disease management
	I:II:III:IV	Education	
Residents from 33 of 35 long-term residential &	10:1:4:1, 5:4:1:1	CM meetings/feedback to other health	
nursing homes	1	providers Medication review (CM/GP/cardiologist)	
Community –initiated CM		(Sim of road along lot)	
-CCTs			
Bonarek- Hessamfar 44	Age: Median 78, 80 yrs. % Female: No details	Co-ordinated care via	Usual care (n=233)
2008 France	% Female: No details Ethnicity: No details	multidisciplinary network (n=129)	Components of intervention: not
n=362	Disease Status: NHYA	Components of intervention at home:	known
Compared patients	Median of	Monitoring signs& symptoms	
included prospectively from	III, IV	Education (diet)	
Jan 1st 2004- Dec 31st	1	Physical therapy	
2005 from GP list	1	CM meetings/feedback to other health provider.	
Lowery ⁴⁵	Age: 65.4(0.51),67.4(0.45) yrs.	Nurse-practitioner-led disease	Usual care (n=510)
2012	% female:1, 1%	management model (n=457)	Coudi Gaio (ii G 10)
USA	Ethnicity:		Components of intervention: not
10.10	White 71.2,79.9%	Components of intervention at home:	known
n=1043		Location was lead tertiary centre, other	
	Black 24,16.1%		
Intervention implemented in	Other 4.8,4.0%	medical centres (some primary care) or	
		medical centres (some primary care) or one affiliated outpatient clinic.	
Intervention implemented in 4 Midwest VA medical	Other 4.8,4.0%	one affiliated outpatient clinic. Monitoring signs& symptoms	
Intervention implemented in 4 Midwest VA medical centres from the same region & one affiliated outpatient clinic and 2 VA	Other 4.8,4.0%	one affiliated outpatient clinic. Monitoring signs& symptoms Education /Self-management support	
Intervention implemented in 4 Midwest VA medical centres from the same region & one affiliated	Other 4.8,4.0%	one affiliated outpatient clinic. Monitoring signs& symptoms	


Key: AF atrial fibrillation, CHF Chronic heart failure, COPD chronic obstructive pulmonary disease, CM case management or case manager, HOCM/RCM Hypertrophic Obstructive/Restrictive Cardiomyopathy, GP general practitioner, UC usual care, NYHA, LV left ventricular SNF skilled nursing facility, PAD peripheral arterial disease,




From: Moher D, Liberati A, Telzielf J, Altman OG, The PRISMA Group (2009). Preferred Reporting Name for Systematic Reviews and Meta


122x146mm (96 x 96 DPI)








177x132mm (96 x 96 DPI)







306x127mm (96 x 96 DPI)

2

4

5 6

7 8 9

10

11 12

13

14 15

16

17 18

19

20 21

22

23

24

25

26 27

28

29 30

31

32 33

34

35

36 37

38 39

40

41 42

43

44

45 46

47

48 49

50

51 52

53

54 55

56

Appendix 1:

Database: Medline In-process - Current week, Medline 1950 to present

Search Strategy:

- 1 randomized controlled trial.pt. (376608)
- 2 random\$.tw. (717987)
- 3 control\$.tw. (2630023)
- 4 intervention \$.tw. (556397)
- 5 evaluat\$.tw. (2214967)
- 6 or/1-5 (5041451)
- 7 Qualitative Research/ (20094)
- 8 semi-structured questionnaire.mp. (1162)
- 9 observation methods.mp. (152)
- 10 Observation/mt [Methods] (635)
- 11 Nvivo.mp. (639)
- 12 interview/ (25018)
- 13 Personal Narratives/ (877)
- 14 Focus Groups/ (16824)
- 15 patient experience*.mp. (8525)
- 16 or/7-15 (70071)
- 17 exp Heart Failure/ (87270)
- 18 exp Heart Failure, Diastolic/ (496)
- 19 exp heart failure, systolic/ (789)
- 20 exp Ventricular Dysfunction/ (26332)
- 21 chronic heart failure.mp. (11341)
- 22 congestive heart failure.mp. (33082)
- 23 cardiac failure.mp. (10151)
- 24 LV dysfunction.mp. (2827)
- 25 left ventricular dysfunction.mp. (9373)
- 26 left ventricular impairment.mp. (188)
- 27 diastolic impairment.mp. (121)
- 28 systolic impairment.mp. (93)
- 29 or/17-28 (135885)
- 30 exp Case Management/ (8326)
- 31 exp Patient Care Planning/ (52319)
- 32 organisation of care.mp. (367)
- 33 community matron.mp. (44)

- 34 "Continuity of Patient Care"/ (14497)
- 35 Community Health Nursing/ (18371)
- 36 transit* care.mp. (580)

2

4 5

6

7 8

9

10 11

12

13 14

15

16 17

18

19 20

21

22 23

24

25 26

27

28 29

30

31 32

33

34 35

36

37

38 39

40 41

42

43

44 45

46

47 48

49

50 51

52

53 54

55

56 57

- 37 Interdisciplinary Communication/ (10602)
- 38 Patient Discharge/ (18977)
- 39 discharge plan.mp. (176)
- 40 exp Patient Care Management/ (535496)
- 41 Comprehensive Health Care/ (6078)
- 42 exp Managed Care Programs/ (38918)
- 43 Primary Health Care/ (54234)
- 44 Community Health Services/ (26923)
- 45 General Practitioners/ (1943)
- 46 Family Practice/ (60223)
- 47 Physicians, Family/ (14745)
- 48 multidisciplinary.mp. (44988)
- 49 or/30-48 (674050)
- 50 6 or 16 (5094326)
- 51 29 and 50 (47009)
- 52 49 and 51 (2590)
- 53 52 not (case report/ or case study/ or letter/ or editorial/ or expert opinion.mp.) (2489)
- 54 53 not (Algeria\$ or Egypt\$ or Liby\$ or Morocc\$ or Tunisia\$ or Western Sahara\$ or Angola\$ or Benin or Botswana\$ or Burkina Faso or Burundi or Cameroon or Cape Verde or Central African Republic or Chad or Comoros or Congo or Diibouti or Eritrea or Ethiopia\$ or Gabon or Gambia\$ or Ghana or Guinea or Keny\$ or Lesotho or Liberia or Madagasca\$ or Malawi or Mali or Mauritania or Mauritius or Mayotte or Mozambiq\$ or Namibia\$ or Niger or Nigeria\$ or Reunion or Rwand\$ or Saint Helena or Senegal or Seychelles or Sierra Leone or Somalia or South Africa\$ or Sudan or Swaziland or Tanzania or Togo or Ugand\$ or Zambia\$ or Zimbabw\$ or China or Chinese or Hong Kong or Macao or Mongolia\$ or Taiwan\$ or Belarus or Moldov\$ or Russia\$ or Ukraine or Afghanistan or Armenia\$ or Azerbaijan or Bahrain or Cyprus or Cypriot or Georgia\$ or Iran\$ or Iraq\$ or Israel\$ or Jordan\$ or Kazakhstan or Kuwait or Kyrgyzstan or Leban\$ or Oman or Pakistan\$ or Palestin\$ or Qatar or Saudi Arabia or Syria\$ or Tajikistan or Turkmenistan or United Arab Emirates or Uzbekistan or Yemen or Bangladesh\$ or Bhutan or British Indian Ocean Territory or Brunei Darussalam or Cambodia\$ or India\$ or Indonesia\$ or Lao or People's Democratic Republic or Malaysia\$ or Maldives or Myanmar or Nepal or Philippin\$ or Singapore or Sri Lanka or Thai\$ or Timor Leste or Vietnam or Albania\$ or Andorra or Bosnia\$ or Herzegovina\$ or Bulgaria\$ or Croatia\$ or Estonia or Faroe Islands or Greenland or Liechtenstein or Lithuani\$ or Macedonia or Malta or maltese or Romania or Serbia\$ or Montenegro or Slovenia or Svalbard or

Argentina\$ or Belize or Bolivia\$ or Brazil\$ or chile or Chilean or Colombia\$ or Costa Rica\$ or



Appendix 2: Subgroup-analysis by intervention component- hospital initiated studies

a) Family involvement

			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.12.1 CM & FI								
Riegel 2002	-0.675	0.219	130	228	25.9%	0.51 [0.33, 0.78]	2002	
Naylor 2004	-0.563	0.197	118	121	26.9%	0.57 [0.39, 0.84]	2004	
Thompson 2005	-1.288	0.298	58	48	22.3%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	24.9%	1.12 [0.70, 1.79]	2006	-
Subtotal (95% CI)			375	462	100.0%	0.56 [0.34, 0.92]		•
Heterogeneity: Tau2=	0.20; Chi2 = 14.0	7, df = 3	3 (P = 0.003); I ² = 79%					
Test for overall effect:	Z = 2.30 (P = 0.02)	2)						
1.12.2 CM minus FI								
Rich 1993	-0.136	0.332	63	35	6.5%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	10.2%	0.44 [0.27, 0.71]	1995	
Riegel 2000	0.054	0.164	140	140	16.6%	1.06 [0.77, 1.46]	2000	+
Blue 2001	-0.129	0.248	82	75	10.1%	0.88 [0.54, 1.43]	2001	- +
Laramee 2003	-0.201	0.321	131	125	6.8%	0.82 [0.44, 1.53]	2003	
DeBusk 2004	-0.098	0.157	228	234	17.4%	0.91 [0.67, 1.23]	2004	+
Jaarsma 2008	0.076	0.151	344	339	18.0%	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	139	14.4%		2009	*
Subtotal (95% CI)			1274	1227	100.0%	0.87 [0.72, 1.04]		•
Heterogeneity: Tau2=	0.03; Chi2 = 11.4	1, df = 7	7 (P = 0.12); I ² = 39%					
Test for overall effect:	Z = 1.54 (P = 0.12)	2)						
								0.01 0.1 1 10 100
Test for subgroup diff	ferences: Chi² = 2.	64, df=	= 1 (P = 0.10), I ^z = 62.2	%			Fa	avours [case management] Favours [usual care]

h) education & self-management

) education	& seit-mai	nage	ement					
			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.20.1 Education and	self-managemer	nt						
Rich 1993	-0.136	0.332	63	35	5.9%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	9.1%	0.44 [0.27, 0.71]	1995	
Blue 2001	-0.129	0.248	84	81	8.9%	0.88 [0.54, 1.43]	2001	-
Riegel 2002	-0.675	0.219	130	228		Not estimable	2002	
aramee 2003	-0.201	0.321	131	125	6.2%	0.82 [0.44, 1.53]	2003	
eBusk 2004	-0.098	0.157	228	234	14.6%	0.91 [0.67, 1.23]	2004	+
laylor 2004	-0.563	0.197	118	121	11.7%	0.57 [0.39, 0.84]	2004	
hompson 2005	-0.1288	0.298	58	48	6.9%	0.88 [0.49, 1.58]	2005	
Riegel 2006	0.112	0.24	69	65	9.3%	1.12 [0.70, 1.79]	2006	
aarsma 2008	0.076	0.151	344	339	15.0%	1.08 [0.80, 1.45]	2008	+
rotons 2009	-0.211	0.188	144	139	12.3%	0.81 [0.56, 1.17]	2009	-
ubtotal (95% CI)			1381	1327	100.0%	0.82 [0.68, 0.98]		♦
leterogeneity: Tau² =	0.03; Chi2 = 15.41	6, df = 9	3 (P = 0.08); I ² = 42%					
est for overall effect:	Z = 2.17 (P = 0.03)	()						
.20.2 No education	and self-manager	nent						
tewart 1999	-0.279	0.122	100	100	55.4%	0.76 [0.60, 0.96]	1999	-
tiegel 2000	0.054	0.164	140	140	44.6%	1.06 [0.77, 1.46]	2000	+
tewart 2012	0.018	0.071	143	137	0.0%	1.02 [0.89, 1.17]	2012	
ubtotal (95% CI)			240	240	100.0%	0.88 [0.63, 1.21]		•
leterogeneity: Tau ² =	0.03; Chi ² = 2.65,	df = 1	$(P = 0.10); I^2 = 62\%$					
est for overall effect:	Z = 0.79 (P = 0.43)	1)						
								0.01 0.1 1 10 100
							Fa	avours [case management] Favours [Usual care]
at for aubarous diff	foroncoo: Chiz - O	1 A AF -	4 /D = 0.74\ IZ = 00/					norman transfer and a company of the contract

Test for subgroup differences: Chi2 = 0.14, df = 1 (P = 0.71), I2 = 0%

c) medication review

			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.19.1 Medication re	view							
Rich 1993	-0.136	0.332	63	35	9.3%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	15.4%	0.44 [0.27, 0.71]	1995	
Stewart 1999	-0.279	0.122	100	100	37.5%	0.76 [0.60, 0.96]	1999	-
Blue 2001	-0.129	0.248	84	81	15.1%	0.88 [0.54, 1.43]	2001	
Brotons 2009	-0.211	0.188	144	139	22.7%	0.81 [0.56, 1.17]	2009	 +
Stewart 2012	0.018	0.071	143	137	0.0%	1.02 [0.89, 1.17]	2012	
Subtotal (95% CI)			533	495	100.0%	0.73 [0.59, 0.91]		♦
Heterogeneity: Tau ² =	0.02; Chi2 = 5.56,	df = 4	$(P = 0.23); I^2 = 28\%$					
Test for overall effect:	Z = 2.87 (P = 0.00	4)						
1.19.2 No medication	review							
Riegel 2000	0.054	0.164	140	140	13.6%	1.06 [0.77, 1.46]	2000	+
Riegel 2002	-0.675	0.219	130	228	12.5%	0.51 [0.33, 0.78]	2002	
Laramee 2003	-0.201	0.321	131	125	10.5%	0.82 [0.44, 1.53]	2003	
DeBusk 2004	-0.98	0.157	228	234	13.7%	0.38 [0.28, 0.51]	2004	-
Naylor 2004	-0.563	0.197	118	121	13.0%	0.57 [0.39, 0.84]	2004	-
Thompson 2005	-1.288	0.298	58	48	10.9%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	12.1%	1.12 [0.70, 1.79]	2006	 -
Jaarsma 2008	0.076	0.151	344	339	13.8%	1.08 [0.80, 1.45]	2008	.+
Subtotal (95% CI)			1218	1300	100.0%	0.66 [0.45, 0.95]		•
Heterogeneity: Tau ² =	0.23; Chi2 = 47.20	0, df = 7	$(P < 0.00001); I^2 = 85$	5%				
Test for overall effect:	Z = 2.26 (P = 0.02)						
								0.01 0.1 1 10 100
							Fa	vours (case management) Favours (usual care)

Test for subgroup differences: $Chi^2 = 0.26$, df = 1 (P = 0.61), $I^2 = 0\%$

d) Referral to other services

			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.16.1 Referral to oth	iers							
Stewart 1999	-0.279	0.122	100	100	16.4%	0.76 [0.60, 0.96]	1999	*
Riegel 2000	0.486	0.318	120	120	9.4%	1.63 [0.87, 3.03]	2000	
Blue 2001	-0.129	0.248	84	81	11.7%	0.88 [0.54, 1.43]	2001	
Riegel 2002	-0.445	0.268	120	120	11.0%	0.64 [0.38, 1.08]	2002	-
Thompson 2005	-1.288	0.298	58	48	10.1%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	12.0%	1.12 [0.70, 1.79]	2006	
Jaarsma 2008	0.076	0.151	344	339	15.4%	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	0	14.0%	0.81 [0.56, 1.17]	2009	-+
Russell 2011	0	0	0	0		Not estimable	2011	
Stewart 2012	0.018	0.071	143	137	0.0%	1.02 [0.89, 1.17]	2012	
Subtotal (95% CI)			1039	873	100.0%	0.82 [0.63, 1.08]		•
Heterogeneity: Tau2=	0.10; Chi2 = 24.2	7, df = 7	(P = 0.001); I2 = 71%					
Test for overall effect:	Z = 1.43 (P = 0.15	0)						
1.16.2 No referral								
Rich 1993	-0.136	0.332	63	35	13.7%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	19.6%	0.44 [0.27, 0.71]	1995	
Laramee 2003	-0.201	0.321	131	125	14.3%	0.82 [0.44, 1.53]	2003	
DeBusk 2004	-0.098	0.157	228	234	28.3%	0.91 [0.67, 1.23]	2004	*
Naylor 2004	-0.563	0.197	118	121	24.1%	0.57 [0.39, 0.84]	2004	-
Subtotal (95% CI)			682	655	100.0%	0.69 [0.51, 0.93]		•
Heterogeneity: Tau ² =	0.06; Chi2 = 8.13,	df = 4	P = 0.09); I ² = 51%					
Test for overall effect:	Z = 2.47 (P = 0.01)						
								0.01 0.1 1 10 100
Test for subgroup differences: Chi² = 0.73, df = 1 (P = 0.39), i² = 0% Favours [case management] Favours [usual care]								

e) Assessment of home environment

			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.18.1 Home assess	ment							
Rich 1993	-0.136	0.332	63	35	21.5%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	30.3%	0.44 [0.27, 0.71]	1995	-
Stewart 1999	-0.279	0.122	100	100	48.2%	0.76 [0.60, 0.96]	1999	=
Stewart 2012	0.018	0.071	143		0.0%	1.02 [0.89, 1.17]	2012	
Subtotal (95% CI)			305	275	100.0%	0.66 [0.45, 0.96]		•
Heterogeneity: Tau ² =	0.06; Chi ² = 4.50	df = 2 (P = 0.11); I ² = 56%					
Test for overall effect:	Z = 2.15 (P = 0.03)	()						
1.18.2 No home asse								
Riegel 2000		0.164	140		11.6%	1.06 [0.77, 1.46]	2000	
Blue 2001	-0.129		84		9.0%	0.88 [0.54, 1.43]		
Riegel 2002	-0.675		130		9.9%	0.51 [0.33, 0.78]	2002	-
Laramee 2003		0.321	131	125	7.1%	0.81 [0.43, 1.52]	2003	
DeBusk 2004	-0.098		228		11.9%	0.91 [0.67, 1.23]	2004	-
Naylor 2004	-0.563		118		10.6%	0.57 [0.39, 0.84]	2004	*
Thompson 2005	-1.288	0.298	58	48	7.7%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	9.3%	1.12 [0.70, 1.79]	2006	+
Jaarsma 2008	0.076	0.151	344		12.1%	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	139	10.9%	0.81 [0.56, 1.17]	2009	<u> </u>
Subtotal (95% CI)			1446	1520	100.0%	0.77 [0.61, 0.97]		•
Heterogeneity: Tau ² =	0.09; Chi ² = 29.3	7, df = 9	$(P = 0.0006); I^2 = 699$	%				
Test for overall effect:	Z = 2.21 (P = 0.03)	1)						
								0.01 0.1 1 10 100
							F	avours [case mangement] Favours [usual care]
Test for subgroup diff	erences: Chi ² = 0.	46, df=	1 (P = 0.50), P = 0%					

f) Case management/health professional meetings

•	· ·	•		_				
			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.21.1 CM/HP meetin	g							
DeBusk 2004	-0.098	0.157	228	234	53.3%	0.91 [0.67, 1.23]	2004	#
Naylor 2004	-0.563	0.197	118	121	46.7%	0.57 [0.39, 0.84]	2004	
Stewart 2012	0.018	0.071	143	137	0.0%	1.02 [0.89, 1.17]	2012	
Subtotal (95% CI)			346	355	100.0%	0.73 [0.46, 1.15]		•
Heterogeneity: Tau2=	0.08; Chi2 = 3.41,	df = 1 ($P = 0.06$); $I^2 = 71\%$					
Test for overall effect:	Z = 1.36 (P = 0.17)						
1.21.2 No CM/HP med	etings							
Rich 1993	-0.136	0.332	63	35	6.2%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	8.3%	0.44 [0.27, 0.71]	1995	
Stewart 1999	-0.279	0.122	100	100	12.0%	0.76 [0.60, 0.96]	1999	*
Riegel 2000	0.054	0.164	140	140	10.7%	1.06 [0.77, 1.46]	2000	+
Blue 2001	-0.129	0.248	84	81	8.2%	0.88 [0.54, 1.43]	2001	
Riegel 2002	-0.675	0.219	130	228	9.0%	0.51 [0.33, 0.78]	2002	-
Laramee 2003	-0.201	0.219	131	125	9.0%	0.82 [0.53, 1.26]	2003	-
Thompson 2005	-1.288	0.298	58	48	7.0%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	8.4%	1.12 [0.70, 1.79]	2006	+
Jaarsma 2008	0.076	0.151	344	339	11.1%	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	139	10.0%	0.81 [0.56, 1.17]	2009	.
Subtotal (95% CI)			1405	1440	100.0%	0.75 [0.60, 0.93]		•
Heterogeneity: Tau ² =	0.09; Chi ² = 32.20	D, df = 1	$0 (P = 0.0004); I^2 = 69$	%				
Test for overall effect:	Z = 2.59 (P = 0.01	0)						
								0.01 0.1 1 10 100
							Fa	avours [case management] Favours [usual care]
To at fav and avenue diff	foroncoo: Chiz - 0	04 46-	1 /D = 0.01\ IZ = 00/					()

Test for subgroup differences: $Chi^2 = 0.01$, df = 1 (P = 0.91), $I^2 = 0\%$

g) Patient-directed access

			Case management	Usual care		Rate Ratio	Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total		Weight	IV, Random, 95% CI	
1.22.1 Patient directe						,	
Jaarsma 2008	0.076	0.151	344	339	25.4%	1.08 [0.80, 1.45]	<u> </u>
Laramee 2003	-0.201		131	125	16.1%	0.82 [0.44, 1.53]	
Naylor 2004	-0.563		118			0.57 [0.39, 0.84]	
Rich 1993	-0.136	0.332	63	35	15.6%	0.87 [0.46, 1.67]	
Rich 1995	-0.825		100	100	20.1%	0.44 [0.27, 0.71]	
Subtotal (95% CI)			756	720	100.0%	0.72 [0.50, 1.04]	◆
Heterogeneity: Tau ² =	0.12; Chi2 = 12.8	D, df = 4	(P = 0.01); I ² = 69%				
Test for overall effect:	Z = 1.75 (P = 0.08)	1)					
4 00 0 11							
1.22.2 No patient-dire							
Blue 2001	-0.129		84		9.2%	0.88 [0.54, 1.43]	
Brotons 2009	-0.211		144			0.81 [0.56, 1.17]	
DeBusk 2004	-0.098		228			0.91 [0.67, 1.23]	
Jaarsma 2008		0.151	344			1.08 [0.80, 1.45]	
Riegel 2000		0.164	140			1.06 [0.77, 1.46]	
Riegel 2002	-0.675		130			0.51 [0.33, 0.78]	
Riegel 2006		0.24	69		9.5%	1.12 [0.70, 1.79]	
Stewart 1999	-0.279		100		14.0%	0.76 [0.60, 0.96]	
Stewart 2012	0.018		143			Not estimable	
Thompson 2005	-1.288	0.298	58		7.7%	0.28 [0.15, 0.49]	
Subtotal (95% CI)	0.00.01.3 00.0		1297		100.0%	0.80 [0.64, 0.99]	T
Heterogeneity: Tau ² =			(P = 0.0009); P = 70	%			
Test for overall effect:	Z = 2.03 (P = 0.04	•)					
							0.01 0.1 1 10 100
Test for subgroup diff	foroncos: Chi²= N	21 df=	1 (P = 0.65) P = 0%			F	avours [case management] Favours [usual carel]
restror subgroup uni	erences. On = 0	21, u1-	1 (1 - 0.05), 1 - 0.0				



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
'Rationale	3	Describe the rationale for the review in the context of what is already known.	4-5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
		Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	N/A
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5
Information sources 7 Sources 7 Sources 8		Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
		Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Appendix one
Study selection 9 State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).		6	
Data collection process	Data collection process 10 Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.		6
B Data items	Data items 11 List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.		6
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6-7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	6-7
Synthesis of results			6-7



PRISMA 2009 Checklist

	Page 1 of 2				
Section/topic	#	Checklist item	Reported on page		
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	6-7		
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7		
RESULTS					
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7 & Figure 1		
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7-8		
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	9		
esults of individual studies 20 For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.					
5 6 7 8 9		Present results of each meta-analysis done, including confidence intervals and measures of consistency.	9-11 plus figures 3 & 4 and appendix 2		
		Present results of any assessment of risk of bias across studies (see Item 15).	9-11 plus figures 3 & 4		
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9-11 plus figures 3 & 4		
DISCUSSION					
Summary of evidence 24 Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).		12			
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	12,14-15		
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	15		



PRISMA 2009 Checklist

FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	21

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. For m. 10 doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.



BMJ Open

Does case management for patients with heart failure based in the community reduce unplanned hospital admissions? A systematic review and meta-analysis

Journal:	BMJ Open
Manuscript ID	bmjopen-2015-010933.R1
Article Type:	Research
Date Submitted by the Author:	24-Mar-2016
Complete List of Authors:	Huntley, Alyson; University of Bristol , Centre of Academic Primary care Johnson, Rachel; Academic unit of primary health care, School of social and community medicine King , Anna; University of Bristol , Centre of academic Primary care Morris, Richard; University of Bristol, School of Social & Community Medicine Purdy, Sarah; University of Bristol, Centre for Academic Primary Care
Primary Subject Heading :	Cardiovascular medicine
Secondary Subject Heading:	General practice / Family practice, Emergency medicine
Keywords:	systematic review, meta-analysis, Heart failure < CARDIOLOGY, case management, hospital admission

Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.

Figure_2a.tiff

SCHOLARONE™ Manuscripts

Version 21st March 2016

Does case management for patients with heart failure based in the community reduce unplanned hospital admissions? A systematic review and meta-analysis

Huntley AL1*

Johnson R¹

King A¹

Morris RW¹

Purdy S¹

Key words: systematic review, meta-analysis, heart failure, case management, hospital admission

Word count: 3887

^{1.} Centre of Academic Primary Care, School of Social and Community Medicine, University of Bristol, Canynge Hall, Bristol, BS8 2PS, UK.

^{*}Dr Alyson Louise Huntley Research Fellow <u>alyson.huntley@bristol.ac.uk</u> Tel no: 0117 3314545

Version 21st March 2016

ABSTRACT

Objectives

The aim of this systematic review of randomised controlled trials (RCTs) and controlled trials (NRCTs) is to investigate the effectiveness and related costs of case management (CM) for patients with heart failure (HF) predominantly based in the community in reducing unplanned readmissions and length of stay (LOS).

Setting

CM initiated either whilst as an inpatient, or on discharge from acute care hospitals, or in the community and then continuing on in the community.

Participants

Adults with a diagnosis of HF and resident in Organisation for Economic Co-operation and Development countries.

Intervention

CM based on nurse co-ordinated multi-component care which is applicable to the primary care based health systems.

Primary and secondary outcomes

Primary outcomes of interest were unplanned (re)admissions, LOS and any related cost data. Secondary outcomes were primary health care resources.

Results

Twenty-two studies were included: 17 RCTs and five NRCTs. Seventeen studies described hospital-initiated CM (n=4794) and five described community-initiated CM of HF (n=3832). Hospital-initiated CM reduced readmissions (rate ratio 0.74 [95%CI 0.60, 0.92] p=0.008) and LOS (mean difference - 1.28 days [95%CI -2.04,-0.52] p=0.001) in favour of CM compared to usual care. Nine trials described cost data of which six reported no difference between CM and usual care. There were four studies of community-initiated CM versus usual care (2RCTs and 2NRCTs) with only the 2 NRCTs showing a reduction in admissions.

Version 21st March 2016

Conclusions

Hospital-initiated CM can be successful in reducing unplanned hospital readmissions for HF and length of hospital stay for people with HF. Nine trials described cost data; no clear difference emerged between CM and usual care. There was limited evidence for community-initiated CM which suggested it does not reduce admission.

Protocol registration

No

Strengths of review

- High quality systematic review
- Interventions examine nurse-led multicomponent care of heart failure patients
- Focus on use of resources specific to heart failure

Limitations of review

- Community-initiated case management trials were limited in quantity and were mostly of low quality.
- Lack of cost data in most trials

INTRODUCTION

 Applying current prevalence figures to population estimates suggests that more than 550,000 individuals (more than 308,000 men and slightly fewer than 250,000 women) in the UK are living with heart failure (HF). Quality and Outcome Framework (QOF) data supports this: in 2012/13 just over 480,000 patients were recorded as having heart failure. The average age of HF patients in general practice in the UK is 77 years.

Prior to 1990, 60–70% of patients died within 5 years of diagnosis, and admission to hospital with worsening symptoms was a regular and recurrent event. Effective treatment has improved care, with a relative reduction in hospitalization in recent years of 30–50%, and smaller but significant decreases in mortality. 4-6

More than £6.8 billion was spent on treating all cardiovascular disease within the NHS in England in 2012/13 with 63% of these costs coming from within secondary care and 21% within primary care. Within secondary care, non-elective inpatient admittance for cardiovascular disease, i.e. emergency admissions, had the greatest expenditure with £1,925 million.¹

Case management (CM) is the process of planning, coordinating and reviewing the care of an individual. We used the definition cited by the King's Fund in the UK 'A collaborative process of assessment, planning, facilitation, care coordination, evaluation, and advocacy for options and services to meet an individual's and family's comprehensive health needs through communication and available resources to promote quality cost-effective outcomes.'⁷ The NHS has used less-intensive approaches than the traditional US model, for example, through the use of nurses to support older people and those with long-term conditions at home.⁸ In this review we have focused on CM based on nurse co-ordinated multi-component care of patients which is applicable to the primary care based health systems such as that in the UK.

A previous systematic review and meta-analysis shows that CM is not effective in reducing unplanned hospital admissions for the general older/elderly population. ⁹ However limited data suggests that CM for patients with HF is promising. ¹⁰ This current review aimed to a) identify the evidence of the effectiveness and related costs of CM interventions for HF patients predominantly based in the community and b) to better understand the potential success of CM by examining the components of tested interventions.

Version 21st March 2016

METHODS

Search

Databases and registries

A search strategy was developed using keywords for the electronic databases according to their specific subject headings or searching structure. The search strategy was run from 1985 – 2012 in the OVID databases - Medline®, Excerpta Medica Database (Embase), Cumulative Index to Nursing and Allied Health Literature (CINAHL®), PsycINFO® on the 2nd July 2014. (Appendix one) The search strategy was modified to search internet sites such as the Agency for Healthcare Research and Quality (AHRQ) and the King's Fund. A pragmatic update of these searches was conducted on the 20/11/15 using the full search strategy and run in Medline and Medline in process only.

Other sources

Once the included papers were determined, both backwards (reference list of paper) and forwards citation searching (via google scholar) was performed to identify any other potentially relevant studies.

All authors of included studies in the field were contacted with data queries and to identify additional relevant studies.

Eligibility criteria:

Randomised controlled trials (RCTs), and other controlled studies (NRCTs) (controlled trials, controlled before and after studies, analytic cohorts, comparative studies) were included as determined by our eligibility criteria. We were aware from our previous work that not all community-based studies were randomised and felt it was important to be more inclusive in order to understand why CM may work for HF. CM interventions needed to be initiated either whilst as an inpatient or on discharge from acute care hospitals including the emergency department (ED), or in the community, and then continue on in the community. Only studies including adults with HF in Organisation for Economic Co-operation and Development (OECD) countries were included as the outcomes were more likely to be comparable for synthesis, and relevant to the UK situation. Studies were included as long as one of the outcomes of interest was unplanned hospital (re)admissions, ED attendance, length of hospital stay (LOS) as well as related costs of the interventions. Other outcomes of interest were primary health care resources e.g. GP visits, visits to other primary care health professionals or

services and prescriptions. Studies written in any language were considered if there was an English abstract available.

Reference management and study selection

EndNote and Excel were used to manage the references. Duplicates were removed from the Endnote file. References underwent a two stage process of screening using the inclusion and exclusion criteria by two reviewers independently (AH, AK, RJ). Firstly, a screen of titles and abstracts (if abstract available) and secondly screening of the full paper was conducted. Where there was continued disagreement between reviewers about including or excluding a paper, a third reviewer made the final decision (SP or RJ).

In addition to the included quantitative intervention papers, we identified relevant reviews from the search. Any potentially relevant conference proceedings were followed up, firstly by searching in Medline to see if the study had been published. If the study was not published, the authors were contacted where possible to check if the studies were likely to be published within the work frame of this review.

Data extraction and assessment of risk of bias

Data were extracted into a custom-designed table which included description of trial type, participants, intervention, controls, outcome measures and results. Based on the Kings Fund definition of case management we devised taxonomy of intervention components. (table 1) As part of this data extraction process, the intervention and control treatments were also described by their component parts e.g. monitoring signs and symptoms using the framework of the CM definition.

Quantitative data concerning the outcomes of interest were extracted into the Cochrane Revman software. The Cochrane risk of bias tool was used to record trial bias for RCTs and the EPOC Risk of bias tool was used for non-randomised controlled trials (NRCTs). These processes were performed by one author and checked by a second. (AH, AK) Any disagreements were resolved by discussion and if necessary with a third author. (RJ, SP)

Data analysis

Trials were divided as previously described by Huntley *et al* 2013 into hospital-initiated CM and community-initiated CM.⁹ Where there were data from three or more studies, effect sizes were calculated and presented in forest plots as rate ratios ((re)admissions) or mean differences (LOS) using Revman software. If the heterogeneity of the combined data was greater than 50%, a random effects model was used for analysis.

We conducted pre-specified sensitivity analysis in response to the risk of bias assessment of studies, removing high risk of bias studies as appropriate; the results of both analyses are presented.¹³ We conducted pre-specified subgroup-analysis to explore the effects of CM duration (3,6 and 12 months plus) on hospital admission and LOS. There was insufficient detail in trials to perform sub-analysis by severity of HF or intensity of intervention.

Data were assessed narratively in respect of the components of interventions using the CM definition cited above as guidance.⁸ (table 1) In addition, where possible post-hoc subgroup-analysis was conducted in Revman in which interventions with components of interest were compared with those that did not have these components.

RESULTS

The systematic review yielded 22 studies with data published over 32 papers of which 17 were RCTs and five were NRCTs all published in the English language. ¹⁴⁻⁴⁵ (Figure one) No relevant studies were identified in a pragmatic update using the full search strategy run in Medline and Medline in process only in November 2015. Seventeen of these studies described hospital-initiated CM (n=4794)^{14,15,17,18,20-24,26-28,31,32, and five described community-initiated CM of HF (n=3832). ^{38,42-45} The PRISMA checklist was used to ensure the quality of our systematic review manuscript.}

Profile of patients (table 2)

The range of female participants in the trials was 1-58%, but the majority of trials had relatively even gender divide. Co and multi-morbidity were common. Eight of the 22 trials gave no detail on ethnicity of participants; in four studies the trialists used white/non-white and English speaking/non-English speaking categories. In the remaining 10 studies, a fuller profile was described. Twelve of the 22 trials

 were conducted in the USA and the ethnicity profile reflected that including Spanish speaking/Hispanic, American Indian, Black, African American, Asian and White participants.

The majority of trials described the severity of HF using New York Heart Association (NYHA) classification. Twelve of the trials gave a breakdown of numbers or percentages in the I-IV classes with some trials only giving numbers of participants for the III and IV class. In these trials the percentage range of III and IV class patients was 6-98%. Four trials gave mean and median values of NYHA status, one trial used the APR-DRG severity of illness scale, and five trials did not describe disease severity.

Profile of interventions (table 2)

The majority of studies (n=15) described the intervention being delivered by a case manager/ specialist nurse with no specific mention of other health professionals, and the remaining seven studies described a case manager/speciality nurse working as part of a multi-disciplinary team.

All but two studies compared CM with usual care although the control group was not always described. The two remaining studies were comparative: one RCT comparing CM with specialist clinics and one RCT comparing CM with telemedicine plus CM. ^{32,42},

The duration of the CM interventions in the studies was 1-24 months with the majority having a 3 or 6 month duration. The majority of studies were conducted face to face or a combination of in-person and by phone. Four interventions were conducted purely by phone. ^{21,22,26,42} Outcomes were measured to match the total duration of intervention in the majority of studies. For many of the studies the intensity of interventions was not stated explicitly. When intensity was described it was always a tapered approach after an initial intensive period.

Risk of bias (Figure 2a&b)

The degree of risk of bias was starkly different between the RCTs and NRCTs. All five of the NRCTs were rated at high risk or unknown risk for most domains. 35,36,37, 44,45

The majority of the RCTs were rated at low risk for most domains with the exception of the domain of blinding of the participants and personnel which is not applicable to this type of intervention. Three RCTs were assessed as at high risk for at least one domain: both Hancock and Wade gave no description of the randomisation process or allocation concealment, Riegel 2002 was randomised at physician level and patients were chosen by physician preference.^{21, 42,43} Four of the five community-

Version 21st March 2016

initiated trials (2 RCTs and 2 NRCTs) were assessed to be at high risk of bias, and in some studies did not present usable data. 35-37,42

All the intervention studies reported unplanned hospital (re)admissions ¹⁴⁻⁴⁵ and 17 reported length of time in hospital. ^{14, 15, 17, 18, 20-22, 24, 26-28, 35, 38, 42-45} There were few data on A and E attendance and primary care resource use. However, only some of the data could be used in meta-analysis with the main reasons being that data were presented in different formats where neither confidence intervals, standard errors nor raw data were given. Due to heterogeneity of data all analysis was conducted using a random effects model.

Unplanned HF (re)admissions data (figure 3)

Hospital-initiated CM

Thirteen of the hospital-initiated CM trials had data that could be used in a meta-analysis of which 12 were RCTs. The pooled data from the RCTs showed a rate ratio of readmissions was 0.74 [95% CI 0.60, 0.92] p=0.008 I² =69% in favour of hospital- initiated CM. (figure 3a) A sensitivity analysis was conducted, removing Riegel 2002 (RCT with high risk of bias for randomisation domain); this had a minimal effect on the rate ratio and heterogeneity 0.77 [0.61, 0.96] p=0.02 I² =68%.^{21,35} Sub-analysis looking at 3, 6 and 12-18 month data did not produce a clear time-related effect which is most likely due to heterogeneity within and between studies.(figure 3b,c,d). There was one hospital-initiated CM trial which compared CM with specialist clinics which reported no differences in hospital readmissions between the two groups.³²

Community initiated CM

Of the four community initiated trials (2 RCTs and 2 NRCTs) comparing admissions between CM with usual care, two reported no significant differences^{38,43} and two reported statistically significant reductions in favour of CM.^{44,45} One further trial compared CM, with telehealth and CM and reported no differences in admissions but data were not presented.⁴²

Length of hospital stay (LOS) (figure 4)

Hospital-initiated CM

Nine of the hospital-initiated CM trials had data that could be used in a meta-analysis of which 8 were RCTs. The pooled data from the RCTs showed that mean LOS was reduced in the CM group

compared to usual care mean difference (MD) -1.28 days [-2.04,-0.52] p=0.001 I^2 =63%. (figure 4a) A sensitivity analysis was conducted removing Riegel 2002 which had an important effect on the rate ratio and heterogeneity MD -1.76 [-2.29,-1.23] p<0.00001 I^2 = 14%. ^{21,35} Sub-analysis looking at 3, 6 and 12-18 month data suggests this effect is short-term (first 3 months) but a longer time-related effect was difficult to assess due to lack of data.(figure 4b,c,d).

The one study comparing CM with specialist clinic care, reported that CM patients accumulated 592 and clinic patients 547 *all-cause* hospitalizations (p = 0.087) associated with 3067 versus 4410 days of hospital stay (p < 0.01 for rate and duration of hospital stay).³²

Community initiated CM

Of the 4 community initiated trials comparing CM with usual care, two did not report LOS ^{38, 43} one reported median values in favour of CM⁴⁴ and the remaining one reported a mean reduction in LOS. ⁴⁵ (table 2). The one comparative trial between community-initiated CM, and telehealth and CM did not report any useful data. ⁴²

Intervention components (table 1 and appendix 2)

Fourteen intervention components were identified and grouped as per the CM definition in methods and prevalence determined for both hospital and community –initiated CM studies with a usual care control group (table 2).

Hospital-initiated CM (appendix 2)

Data from individual studies which contained components of family involvement showed an overall reduction in hospital readmissions in comparison with usual care and a reduction in hospital readmissions observationally in comparison with interventions which did not contain these components (rate ratio of 0.56 [0.34, 0.92] p=0.003). However, post-hoc analysis comparing these studies in which the component was present with those studies in which the component was absent did not yield any statistically significant differences (p=0.15). (appendix 2a) The same calculations for medication review, referral to other services, and assessment of home environment, CM meetings and patient-directed access did not indicate any specific effect of these components of hospital-initiated CM on rates of admission. (appendix 2 b-g) The majority of the interventions included

Version 21st March 2016

education/self-management and there were insufficient data from studies without this component to allow comparison.

Community initiated CM

There were insufficient data to conduct any subgroup-analysis on any of the remaining components of hospital-initiated CM, community initiated studies or the LOS data.

Outpatient health care resources

Only six of the included studies measured outpatient resource use. In some studies, outpatient resource data were all-cause and not HF-specific. In some studies primary and secondary use was combined. 23,24,35,38,42,45 Two of these studies also reported ED attendance. All but one of these studies reported no difference between intervention and control group for these measures with the exception of Lowery which showed a statistically significant greater use of outpatient resources in the usual care group (optional primary care visits 1yr 16.75(13.62),10.43(9.6) p<0.001, 2yr 14.27(11.98),9.35(9.97) p<0.001).

Costs (table 3)

Nine of the seventeen hospital initiated trials described cost data. Of these, six reported no statistically significant difference between CM (3 or 6 month duration) and usual care. ^{17,18,22,24, 26 35,} and three reported costs in favour of CM although data from Stauffer was brief. ^{15,32,37} One of these was 12-18 months ³² and two were 3 months in duration. It was difficult from the intervention descriptions to determine their intensity. There were no cost data reported from the community-initiated trials.

DISCUSSION

This systematic review confirms that hospital-initiated CM can be successful in reducing unplanned hospital readmissions, and reducing LOS in hospital in the short term for people with HF. There were only five community-initiated CM studies (3RCTs and 2NRCTs) of which four were at high risk of bias. This limited evidence suggests no effect of community-initiated CM on hospital admissions. A

 minority of trials report cost comparisons with usual care and most of those show no difference. There were limited data on the effect of CM on other health care resources.

Many factors are likely to modify the effect of CM on use of emergency care seen in these studies. It is generally accepted that CM is more appropriate for people with severe HF and poorer general health. However it was difficult to compare the health status of the study participants in both hospital-initiated and community-initiated trials as in some studies there was little detail, others gave median and mean figures for NYHA status, and the presentation format and detail of co-morbidities varied. All the included studies have been conducted within the past 12 years so it is important to put these results in the context of overall improved treatment and reduction in hospital admissions since the early 1990's. 4-6

Seventeen studies described hospital-initiated CM and five described community initiated CM of heart failure although often the participants were identified via hospital clinic records. Overall the meta-analysis showed that CM reduced readmissions and hospital LOS. This may be explained by the fact that in most of the trials the participants were identified via hospital contact and therefore were likely to have had a recent exacerbation of their HF and to be at increased risk of re-admission in the post-discharge period. In addition, it is likely that interventions are acting at a time of highest risk as reflected by HF mortality in first year of diagnosis. Therefore once they were assessed and given extra support they were stable for a period of time. Previous work by Roland *et al* 2005 suggests that admission rates in people aged 65 with two or more emergency admissions in 12 months fall in subsequent years without any intervention and account for fewer than 10% of admissions in the following year and thus effectiveness of admission avoidance schemes cannot be judged by tracking admission rates without careful comparison with a control group. The data from trials of community-initiated CM was lacking both in the number of studies, and the fact there were limited useable data that showed no effect on unplanned hospital admissions. It is likely that these patients were likely to be in more stable health.

A meta-review of a wide range of HF disease management programs by Savard 2011 reports that nine previous systematic reviews (2001-2009) identified significant reductions in HF admissions with reductions in risk ranging from 30-56%. ⁴⁷ However the authors caution that these reviews are limited by inadequate reporting in the population, setting, intervention and comparator components. They

 Version 21st March 2016

report that reviewers have not taken into account statistical, clinical and methodological heterogeneity in interventions.⁴⁷ Our review focussed specifically on CM avoiding some of these limitations and indicates a reduction in HF readmissions with hospital-initiated CM in the range of 10-30%.

Wakefield *et al* in 2013 looked at common components of a range of HF care programs focusing mainly on disease management and education investigated in RCTs, and 10/35 of the discussed studies were included in our review. They described patient education, symptom management by health professionals and by patients, and medication adherence strategies as the most commonly occurring elements of care. A literature review by Jaarsma 2012 looked at 70 'home care' controlled studies (mostly RCTs) which encompassed 9 of our included CM studies covering a wide range of approaches such as telemedicine, hospital at home and health buddies for patients with HF. They identified a multi-disciplinary team, continuity of care, care plans, optimising titration of medication, education/counselling of patients and caregivers and increased access as important. Unfortunately we had insufficient data to perform sub-analysis on the component of education/self-management. Previous systematic reviews have investigated the role of the lay caregiver in HF patient management. These suggest that better relationship quality and communication were related to reduced mortality, increased health status and less distress and improved patient self-care outcomes. Our review adds to this evidence base by suggesting that more family involvement in CM may also reduce unscheduled readmissions.

Education about HF and about its pharmacological and non-pharmaceutical treatment has been well-reviewed both as an individual approach and as part of complex interventions, and is considered to be essential for improving many patient outcomes. ^{49, 53,54}A recent mixed method study, suggests asking patients with HF to write down their learning needs before the education increases their chances of receiving education based on their individual needs. ⁵⁵ Qualitative interviews with health professionals caring for HF patients suggest that communication with, and education by specialist nurses facilitated by continuity of care is essential to good care of HF patients. The authors also highlight the role of the specialist nurse in multi-disciplinary team communication and functioning; essentially describing the role of the specialist nurse as a case manager. ⁵⁶

Our review of CM suggests that the evidence for its cost-effectiveness is lacking with most studies that have performed cost comparisons with usual care show no advantage. Previous work by de Bruin

et al 2001 looked at cost effectiveness of disease management for a range of chronic conditions and concluded that the data is most positive for HF with 5 out of the 8 included studies showing cost-effectiveness.⁵⁷

Strengths and limitations

The contribution of our high quality systematic review to the above is that we have focused on CM which is based on nurse co-ordinated multi-component care of patients which is applicable to the primary care based health systems such as that in the UK. We have focused on HF (re)admissions and LOS as opposed to all-cause data which many of the previous reviews have used.

By examining the components of CM we have a profile of the components most likely to lead to the success of CM of patients with heart failure in terms of reducing (re)admissions and hospital LOS.

Our review has high-lighted the potential importance of family involvement albeit in post-hoc analysis.

The limitations of this review are that majority of the community-initiated CM studies were of low quality with the exception of one low risk of bias RCT, and provided limited evidence. Whilst funnel plot analysis was not appropriate with our data we acknowledge that there may be publication bias on this topic. This was counteracted by the fact that the hospital-initiated studies comprised of predominantly community-based case management. There is a lack of cost data and analysis in the included papers. This point needs to be emphasised for future trials. It is possible that cost effectiveness will be more likely with intervention for patients with more severe HF.

CONCLUSIONS

Hospital-initiated CM reduces unplanned hospital admissions, and length of stay for people with HF in the short term. Cost data is limited. There was limited evidence for community-initiated CM which suggested it does not reduce hospital admission. Further research is needed to determine the individual components of CM that contribute to reduced admissions.

Version 21st March 2016

Contributorship statement

Alyson Huntley. Main systematic reviewer, worked across all stages of the review from inception to completed draft.

Rachel Johnson. Cardiology and primary care expertise. Worked on screening, selection of studies, commenting on analysis, and development and checking of final document content.

Anna King. Second reviewer, involved in screening, selection, data checking and commenting on developing and final document content.

Richard Morris. Statistical expertise, advising on data analysis and commenting on the developing and final document content.

Sarah Purdy. Primary care and admission avoidance expertise. Advised throughout project, third reviewer for screening process and commenting on the developing and final document content.

Competing interests

None declared

Funding

This project was funded by the National School of Primary Health Care project no.238.

Data sharing statement

Full data extraction tables and data analysis files are available on request

REFERENCES

- Cardiovascular disease statistics 2014 British Heart Foundation accessed at https://www.bhf.org.uk/~/media/files/publications/research/bhf_cvd-statistics-2014_web_2.pdf on the 7/10/15.
- Quality and Outcomes Framework Achievement, prevalence and exceptions data 2012/13. 29
 October 2013. Accessed at http://www.hscic.gov.uk/catalogue/PUB12262/qual-outc-fram-12-13-rep.pdf on the7/10/15
- de Giuli F, Khaw KT, Cowie MR, et al. Incidence and outcome of persons with a clinical diagnosis of heart failure in a general practice population of 696,884 in the United Kingdom. Eur J Heart Fail 2005; 7(3):295–302.
- 4. Stewart S, MacIntyre K, Hole DJ *et al.* More 'malignant' than cancer? Five-year survival following a first admission for heart failure. *Eur J Heart Fail* 2001; 3:315–322.
- Stewart S, Ekman I, Ekman T et al. Population impact of heart failure and the most common forms of cancer: a study of 1 162 309 hospital cases in Sweden (1988 to 2004). Circ Cardiovasc Qual Outcomes 2010; 3:573–580.523.
- Jhund PS, Macintyre K, Simpson CR et al. Long-term trends in first hospitalization for heart failure and subsequent survival between 1986 and 2003: a population study of 5.1 million people. Circulation 2009; 119:515–523.
- Ross S, Curry N, Goodwin N. Case Management. What Is It and How It Can Best Be Implemented. London, UK: The Kings Fund, 2011. Accessed at http://www.kingsfund.org.uk/publications/case_management.html on the7/10/15.
- Department of Health. Supporting People with Long Term Conditions. Liberating the Talents
 of Nurses Who Care for People with Long Term Conditions. London, UK: Department of
 Health, 2005. Accessed at
 www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH
 _4102469 on 01 /12/11.
- Huntley AL, Thomas R, Mann M et al. Is case management effective in reducing the risk of unplanned hospital admissions for older people? A systematic review and meta-analysis. Fam Pract 2013; 30(3):266-75

10. Purdy S. Interventions to reduce unplanned hospital admission: a series of systematic reviews. Report accessed at http://www.bristol.ac.uk/medialibrary/sites/primaryhealthcare/migrated/documents/unplannedadmissions.pdf on 07/10/15.

- List of OECD Member countries accessed at http://www.oecd.org/about/membersandpartners/list-oecd-member-countries.htm on 07/10/15.
- Chapter 8: Assessing risk of bias in included studies accessed at http://handbook.cochrane.org/chapter_8/8_assessing_risk_of_bias_in_included_studies.htm on 07/10/15.
- Chapter 9- 9.7 Sensitivity analyses accessed at http://handbook.cochrane.org/chapter_9/9_7_sensitivity_analyses.htmon07/10/15.
- 14. Rich, Vinson JM, Sperry JC *et al.* Prevention of readmission in elderly patients with congestive heart failure: results of a prospective, randomized pilot study. *J Gen Intern Med* 1993; 8(11): 585-90. **MAIN PUBLICATION**
- Rich MW, Beckham V, Wittenberg C et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. NEJM 1995; 333(18): 1190-5.

MAIN PUBLICATION

16. Rich MW, Gray DB, Beckham V *et al.* Effect of a multidisciplinary intervention on medication compliance in elderly patients with congestive heart failure. *Am J Med* 1996; 101 (3): 270-6.

EXTRA

 Stewart S, Pearson S, Luke CG et al. Effects of home-based intervention on unplanned readmissions and out-of-hospital deaths. J Am Geriatr Soc 1998; 46(2):174-80. MAIN PUBLICATION

- Stewart S, Marley JE, Horowitz JD. Effects of a multidisciplinary, home-based intervention on unplanned readmissions and survival among patients with chronic congestive heart failure: a randomised controlled study. *Lancet* 1999; 354 (9184): 1077-83. MAIN PUBLICATION
- 19. Stewart S, Horowitz JD. Home-based intervention in congestive heart failure: long-term implications on readmission and survival. *Circulation* 2002; 105 (24): 2861-6. **EXTRA**
- Blue L, Lang E, McMurray et al. Randomised controlled trial of specialist nurse intervention in heart failure. BMJ 2001; 323(7315): 715-718. MAIN PUBLICATION

- 21. Riegel B, Carlson B, Kopp Z *et al.* Effect of a standardized nurse case-management telephone intervention on resource use in patients with chronic heart failure. *Arch Intern Med* 2002; 162(6):705-12. **MAIN PUBLICATION**
- 22. Laramee S, Levinsky SK, Sargent J et al. Case management in a heterogeneous congestive heart failure population: a randomized controlled trial. Arch Intern Med 2003; 163(7): 809-17.
 MAIN PUBLICATION
- 23. DeBusk RF, Miller NH, Parker KM *et al.* Care management for low-risk patients with heart failure: a randomized, controlled trial. *Ann Intern Med* 2004; 141(8):606-13. **MAIN PUBLICATION**
- 24. Naylor MD, Brooten DA, Campbell RL et al. Transitional Care of Older Adults Hospitalized with Heart Failure: A Randomized, Controlled Trial. J Am Geriatr Soc 2004; 52(5):675-684.
 MAIN PUBLICATION
- McCauley KM, Bixby MB, Naylor MD. Advanced practice nurse strategies to improve outcomes and reduce cost in elders with heart failure. *Dis Manag* 2006; 9(5):302-10. EXTRA
- 26. Riegel B, Carlson B, Glaser D, Romero T.
 Randomized controlled trial of telephone case management in Hispanics of Mexican origin wit h heart failure. J Card Fail 2006; 12(3):211-9. MAIN PUBLICATION
- 27. Thompson DR, Roebuck A, Stewart S. Effects of a nurse-led, clinic and home-based intervention on recurrent hospital use in chronic heart failure. Eur J Heart Failure 2005; 7(3): 377-384. MAIN PUBLICATION
- 28. Jaarsma T, van der Wal MH, Lesman-Leegte I et al. A Coordinating Study Evaluating Outcomes of and I. Counseling in Heart Failure Effect of moderate or intensive disease management program on outcome in patients with heart failure: Coordinating Study Evaluating Outcomes of Advising and Counseling in Heart Failure (COACH). Arch Intern Med 2008; 168(3): 316-24. MAIN PUBLICATION
- 29. Postmus D, Pari AA, Jaarsma T *et al.* A trial-based economic evaluation of 2 nurse-led disease management programs in heart failure. *Am Heart J* 2011; 162(6): 1096-104. **EXTRA**
- 30. Inglis SC, Pearson S, Treen S *et al.* Extending the horizon in chronic heart failure: effects of multidisciplinary, home-based intervention relative to usual care. *Circulation* 2006; 114 (23): 2466-73. **EXTRA**

- 31. Brotons C, Falces C, Alegre J *et al.* Randomized clinical trial of the effectiveness of a home based intervention in patients with heart failure: the IC-DOM study. *Rev Esp Cardiol* 2009; 62(4):400-8. **MAIN PUBLICATION**
- 32. Stewart S, Carrington MJ, Marwick TH et al. Impact of home versus clinic-based management of chronic heart failure: the WHICH? (Which Heart Failure Intervention Is Most Cost-Effective & Consumer Friendly in Reducing Hospital Care) multicenter, randomized trial.
 J Am Coll Cardiol 2012; 60(14);1239-48. MAIN PUBLICATION
- 33. Stewart S, Carrington MJ, Horowitz JD et al.
 Prolonged impact of home versus clinicbased management of chronic heart failure:
 extended follow-up of a pragmatic, multicentre randomized trial cohort. Int J Cardiol 2014;
 174(3):600-10. EXTRA
- 34. Stewart S, Carrington MJ, Chan YK. Prolonged benefits of nurse-led, home-based intervention versus a specialist heart failure clinic: Extended follow-up of the WHICH? Trial Cohort. Eur J Heart Failure 2014; 16(106). EXTRA
- 35. Riegel B, Carlson B, Glaser D *et al.* Which patients with heart failure respond best to multidisciplinary disease management? *J Cardiac Fail* 2000; 6 (4): 290-9. **MAIN PUBLICATION**
- 36. Russell D, Rosati RJ, Sobolewski S *et al.* Implementing a transitional care program for highrisk heart failure patients: findings from a community-based partnership between a certified home healthcare agency and regional hospital. *J Healthcare Qual* 2011; 33(6):17-23. **MAIN PUBLICATION**
- 37. Stauffer BD, Fullerton C, Fleming N et al.
 Effectiveness and cost of transitional care program for heart failure:
 a prospective study with concurrent controls. Arch Intern Med 2011; 171(14):1238-43. MAIN
 PUBLICATION
- 38. Peters-Klimm F, Campbell S, Hermann K *et al.* Case management for patients with chronic systolic heart failure in primary care: The HICMan exploratory randomised controlled trial.

 Trials 2010; 11: 56. MAIN PUBLICATION
- 39. Peters-Klimm F, Muller-Tasch T, Schellberg D *et al.* Rationale, design and conduct of a randomised controlled trial evaluating a primary care-based complex intervention to improve

- the quality of life of heart failure patients: HICMan (Heidelberg Integrated Case Management).

 BMC Cardiovascular Disorders 2007: 7, (25). PROTOCOL
- 40. Freund F, Baldauf A, Muth C *et al.* Practice-based home visit and telephone monitoring of chronic heart failure patients: rationale, design and practical application of monitoring lists in the HICMan trial *Zeitschrift fur Evidenz Fortbildung und Qualitat im Gesundheitswesen* 2011; 105 (6): 434-45. **EXTRA**
- 41. Peters-Klimm F, Campbell S, Muller-Tasch T *et al.* Primary care-based multifaceted, interdisciplinary medical educational intervention for patients with systolic heart failure: lessons learned from a cluster randomised controlled trial. *Trials* 2009; 10:68. **EXTRA**
- 42. Wade MJ, Desai AS, Spettell CM *et al.* Telemonitoring with case management for seniors with heart failure. *Am J Manag Care* 2011; 17(3): e71-9. **MAIN PUBLICATION**
- 43. Hancock HC, Close H, Mason JM et al. Feasibility of evidence-based diagnosis and management of heart failure in older people in care: a pilot randomised controlled trial. BMC Geriatr 2012 Nov 14;12:70.
- 44. Bonarek-Hessamfar M, Benchimol D, Lauribe P et al.
 Multidisciplinary network in heart failure management in a community
 based population: results and benefits at 2 years. Int J Cardiol 2009 May 1;134(1):120-2.
- 45. Lowery J, Hopp F, Subramanian U et al. Evaluation of a nurse practitioner disease management model for chronic heart failure: a multi-site implementation study. Congest Heart Fail 2012; 18(1):64-71.
- 46. Roland M, Dusheiko M, Gravelle H *et al.* Follow up of people aged 65 and over with a history of emergency admissions: analysis of routine admission data. *BMJ* 2005; 330(7486):289-92.
- 47. Savard LA, Thompson DR, Clark AM.
 A metareview of evidence on heart failure disease management programs:
 the challenges of describing and synthesizing evidence on complex interventions. *Trials* 2011;
 12:194.
- 48. Wakefield BJ, Boren SA, Groves PS *et al.* Heart failure care management programs: a review of study interventions and meta-analysis of outcomes. *J Cardiovasc Nurs* 2013; 28(1):8-19
- 49. Jaarsma T, Brons M, Kraai I *et al.* Components of heart failure management in home care; a literature review. *Eur J Cardiovasc Nurs* 2013; 12(3):230-41.

Version 21st March 2016

- 50. Hooker SA, Grigsby ME, Riegel B *et al.* The Impact of Relationship Quality on Health-Related Outcomes in Heart Failure Patients and Informal Family Caregivers: An Integrative Review. *J Cardiovasc Nurs* 2015;30(4 Suppl 1):S52-63.
- 51. Buck HG, Harkness K, Wion R *et al.* Caregivers' contributions to heart failure self-care: a systematic review. *Eur J Cardiovasc Nurs* 2015; 14(1):79-89.
- 52. Clark AM, Spaling M, Harkness K *et al.* Determinants of effective heart failure self-care: a systematic review of patients' and caregivers' perceptions. *Heart* 2014;100(9):716-21.
- 53. Spatola CF, Cocchieri A, De Marinis MG *et al.* Educational interventions in patients with heart failure: a review of the literature. Article in Italian *Ig Sanita Pubbl* 2013; 69(5):557-74.
- 54. Boyde M, Turner C, Thompson DR *et al.* Educational interventions for patients with heart failure: a systematic review of randomized controlled trials. *J Cardiovasc Nurs* 2011; 26(4):E27-35.
- 55. Ross A, Ohlsson U, Blomberg K *et al.* Evaluation of an intervention to individualise patient education at a nurse-led heart failure clinic: a mixed-method study. *J Clin Nurs* 2015; 24(11-12):1594-602.
- 56. Glogowska M, Simmonds R, McLachlan S et al. Managing Patients With Heart Failure: A Qualitative Study of Multidisciplinary Teams With Specialist Heart Failure Nurses. Ann Fam Med 2015; 13(5):466-71.
- 57. de Bruin SR, Heijink R, Lemmens LC et al. Impact of disease management programs on healthcare expenditures for patients with diabetes, depression, heart failure or chronic obstructive pulmonary disease: a systematic review of the literature. Health Policy 2011; 101(2):105-21.
- 58. Chapter 10 Recommendations on testing for funnel plot asymmetry accessed at http://handbook.cochrane.org/chapter_10/10_4_3_1_recommendations_on_testing_for_funne l plot asymmetry.htm on 07/10/15.

Table 1: Components of CM interventions

Definition & total prevalence of Components of CM interventions	Number of hospital- initiated CM vs. usual care with component present (total studies=16)	Number of community-initiated CM vs. usual care with component present (total studies=4)
Assessment /Evaluation		
Monitoring signs & symptoms (n=18) Encompasses general care of CHF patients which is likely to include establishing a relationship with patient over visits, physical and cardiac status checking, lifestyle assessment, general medication check and screening tests e.g. depression, dementia	14	4
Medication review	6	2
(n=8) Review and adjustment of medication by experienced case manager (nurse), pharmacist, GP or consultant often using a		
combination of these health professionals.		
Assessment of home environment (n=4) Assessment carried out by case manager to identify any issues or potential issues with home environment e.g. stairs	4	0
Planning		
CM meetings/feedback to other HPs	3	2
(n=5) Group meetings of health professionals involved in CHF patients care with the aim of reporting on and planning for patients care. adients care.		
Appointment organisation (n=2)	2	0
Case manager checking medical appointments, ensuring ability to go etc.		
Advance care planning (n=1)	1	
Facilitation		
Education/self-management (n=18) (n=18) Education (n=18) Educating CHF patients about their condition, treatment and what to expect. The aim of this is to assist self-management (care with assistance of health professionals) and self-care (patient engaging in activities to promote their health and well-being).	15	3
Patient-directed access (n=6) The ability of CHF patients to initiate care from the case manager or case management service.	6	0
Care co-ordination		
Referral to (n=14) When the case manager refers the patient to other health or social care professionals, this can be GP hospital consultant, social care or tests.	11	3
Advocacy for options & services		
Equipment (n=4) Description of thems to assist action! a health age push as all acceptors, unintaing eaches and macaused unter heitles.	3	1
Provision of items to assist patient's health care such as pill counters, weighing scales and measured water bottles. Physical therapy (n=1)	1	0
CHF patient receiving physical therapy/rehabilitation		
Support group (n=1) CHF attending or being offered the opportunity of a support group.	1	0
Other		
Family involvement (n=9) When the patient's family in terms of information, education or involvement e.g. goal setting in	8	1
patients' care or active monitoring Emotional support (n=1)	0	1
Case manager providing emotional support to CHF patient.		

Table 2: study characteristics of intervention studies

Non-pharmacological comprehensive multidisciplinary treatment strategy NPCM (n=63) Components of intervention at home: Monitoring signs& symptoms Medication review (nurse) Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Usual care UC (n=35) Components of intervention: Visits by home nurse Usual care (n=140) As above for Rich 1993 Usual care (n=48) Components of intervention:	Intervention vs. control No. of readmissions (%) 21[CI 21.7, 44.9 (33.3%), 16 [29.2,62.2] (45.7%) Total hospital days 272, 200 Mean no of days 4.3(SD1.1),5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4) P=0.04
comprehensive multidisciplinary treatment strategy NPCM (n=63) Components of intervention at home: Monitoring signs& symptoms Medication review (nurse) Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Components of intervention: Visits by home nurse Usual care (n=140) As above for Rich 1993 Usual care (n=48)	21[Cl 21.7, 44.9](33.3%), 16 [29.2,62.2] (45.7%) Total hospital days 272, 200 Mean no of days 4.3(SD1.1),5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
comprehensive multidisciplinary treatment strategy NPCM (n=63) Components of intervention at home: Monitoring signs& symptoms Medication review (nurse) Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Components of intervention: Visits by home nurse Usual care (n=140) As above for Rich 1993 Usual care (n=48)	21[Cl 21.7, 44.9](33.3%), 16 [29.2,62.2] (45.7%) Total hospital days 272, 200 Mean no of days 4.3(SD1.1),5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
treatment strategy NPCM (n=63) Components of intervention at home: Monitoring signs& symptoms Medication review (nurse) Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Usual care (n=140) As above for Rich 1993 Usual care (n=48)	[29.2,62.2] (45.7%) Total hospital days 272, 200 Mean no of days 4.3(SD1.1).5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
home: Monitoring signs& symptoms Medication review (nurse) Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Usual care (n=140) As above for Rich 1993 Usual care (n=48)	272, 200 Mean no of days 4.3(SD1.1).5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
Monitoring signs& symptoms Medication review (nurse) Gducation /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	As above for Rich 1993 Usual care (n=48)	Mean no of days 4.3(SD1.1),5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
Education /Self-management support Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	As above for Rich 1993 Usual care (n=48)	4.3(SD1.1),5.7(SD2.0) No. of readmissions 24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
Assessment of home environment Patient directed access to study personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	As above for Rich 1993 Usual care (n=48)	24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
Personnel Nurse-directed multidisciplinary intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	As above for Rich 1993 Usual care (n=48)	24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
intervention (n=142) As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	As above for Rich 1993 Usual care (n=48)	24,54 p=0.04 Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
As above for Rich 1993 Home-based intervention (n=49) Components of intervention at home: Monitoring signs& symptoms	Usual care (n=48)	Total hospital days 556,865 Mean no. of days 3.9(10), 6.2(11.4)
Components of intervention at home: Monitoring signs& symptoms	i i	556,865 Mean no. of days 3.9(10), 6.2(11.4)
Components of intervention at home: Monitoring signs& symptoms	i i	3.9(10), 6.2(11.4)
Components of intervention at home: Monitoring signs& symptoms	i i	
Components of intervention at home: Monitoring signs& symptoms	i i	
home: Monitoring signs& symptoms	Components of intervention:	No. of readmissions 36,63 (p=0.03)
	Disease management	No. of patients experiencing a
Education (pharmacist)/self-		readmission 24,31 (p=0.12)
management support Medication review (pharmacist)		LOS in days
Referral to GP Family involvement		261,452 (p=0.05)
Equipment Multidisciplinary home base	Usual care (n=100)	6mths
intervention (n=100)	Components of intervention:	No. of readmissions
Components of intervention at home:	Contact with other health and social professionals	(event rates give p=0.02)
Monitoring signs& symptoms Referral to other health & social care	Appointment with GP or cardiac clinic or both	Rate of readmissions 0.14(0.1,018), 0.34 (0.19,0.49)
Appointment organisation	or port	p=0·031
Assessment of home environment Family involvement		LOS in days
Education/self-management support Medication review		460,1174 0.9 (0.6,1.2) 2.9(1.9,3.9) p=0.004
(nurse/GP/cardiologist)		18mths
		No. of readmissions 64,125 p=0.02
		Mean number of hospital days 10.5 (14.4), 21.1(24.1) days per
Specialist nurse intervention (Usual care (n=75)	patient p=0.004 No.(%) of readmissions
n=82)	Components of intervention:	12(14), 26(32)
Components of intervention at home:	GP care	Hazard ratio 0.38 (0.19,0.76) p=0.0044
Monitoring signs& symptoms Education/Self-management support		LOS in days
Referral to other health & social care Appointment organisation		3.43 (12.2), 7.46(16.6) CI 0.6 (0.41, 0.88) p=0.0051)
Medication review (nurse, cardiologist)		
	Ilsual care (n=228)	Readmission rates
Telephonic case management	Components of intervention:	3mths 14.6, 22.8 p=0.06 [calculation 19 people vs. 52 exp 1
Telephonic case management (n=130)	INOL KIIOWII	[calculation 19 people vs. 52 exp 1 or more ad] 6mths 17.7, 27.6 p=0.06
(n=130) Components of intervention at		[calculation 23 people vs. 63
(n=130) Components of intervention at home: Monitoring signs& symptoms		people exp 1 or more ad]
(n=130) Components of intervention at home: Monitoring signs & symptoms Self-management support Referral to other HPs (includ.GP) &		LOS in days 3mths 0.85(2.3), 1.6(3.9) p=0.56
	(n=130) Components of intervention at	(n=130) Components of intervention: Not known Not known Monitoring signs& symptoms Self-management support

1	
2	
3	
4	
5	
6	
7	
1	
8	
9	
1	0
1	1
1	2
1	3
1	4
1	- 5
1	2
١	O
1	1
1	8
1	9
2	012345678901234567890123456
2	1
2	2
2	3
2	1
2	_
2	S
2	0
2	1
2	8
2	9
3	0
3	1
3	2
3	_ ব
3	1
2	-
ა ი	S
3	ნ 7
_	-
	8
3	9
4	
4	
4	
4	
4	Δ
4	
4	0
4	1
4	8
4	9
5	0
5	1
5	2
5	3
5	4
5	5
J	S
5	6
5	1
5	8
_	a

Laramee ²² 2003	Age: 70.6(11.4), 70.8(12.2) yrs. % female:	Case management (n=131 data available)	Usual care (n=125 data available)	No. of readmissions 3mth period
USA	42,50% Ethnicity:	Components of intervention at	Components of intervention: Not known	18(14) vs.21(17) ns
n=287 randomised	Not reported Disease Status (SD):	home: Monitoring signs& symptoms		LOS in days
Patients admitted to hospital for CHF were screened.	NYHA I 10(7),35(26) II 76(55),47(36)	Education/Self-management support Family involvement Equipment		in hospital for those patients with ≥1 readmission
screened.	III 50(36),46(35) IV 3(2),4(3) Note P<0.001	Patient-directed access to CM		6.9(6.5), 9.5(9.8) ns
De Busk ²³ 2004	Age: <60yrs 15,14%, 6-70yrs 22,24%,70-80yrs	Case management (n=228)	Usual care (n=234)	Total no. of readmissions in one year
USA n=462 randomised	40,37%, >80yrs 21,26% % Female :52,45%	Components of intervention at home:	Components of interventions: not known	76,86 no stats available
Patients who were	Ethnicity: American Indian 0,1%	Monitoring signs& symptoms Education /Self-management support		
admitted with a provisional diagnosis of HF from	Asian 4,8% Black 2,2%	CM meetings/feedback to other health providers		
Kaiser Permanente medical centres in	White 5,6% Hispanic 3,3%			
California	Disease Status: NHYA I-II 50,50% III-IV 50,50%			
Naylor ^{24,25} 2004	Age: 76.4(6.9),75.6(6.5) % Female:60, 56%	Transitional care intervention with advanced practice nurses (APNs)	Usual care (n=121)	No. of readmissions 40 vs. 72 ns
USA n=239 patients	Ethnicity: African American 34,38%, White 66, 62%	(n=118)	Components of intervention: Care from standard home care	\$175,840,\$498,110
randomised	Disease Status: Functional Status (Moinpur C	Components of intervention at home:	services Patient–directed access to home care	Total hospital days (all cause)
Patients aged 65yrs+ admitted to 6 study	1992) Personal 17.1(5.8),16.9(5.8)	Monitoring signs& symptoms Education/Self-management support	services	588, 970 Per patient, mean ±SD 5.0 ± 7.3
hospitals from home with a diagnosis of HF were	Social 8.4(2.6),8.6(2.6) Total 25.5(8),25.4(7.8)	Family involvement CM meetings/feedback to other health		8.0 ±2.3 ns Per hospitalized patient,
screened for participation.	·	providers Patient–directed access to CM		mean± SD 11.1±7.2 14.5±13.4 ns
Riegel ²⁶ 2006	Age: 71.6910.8),72.7(11.2) % Female:58,49.2%	Telephonic case management (n=69)	Usual care (n=65)	Readmission rates (%) (no. of people) all ns
San Diego USA	Ethnicity: Hispanic patients Speak/read only Spanish	Components of intervention at	Components of intervention: Disease management	1mth 8.7,13.8% [Calculation 6.003,8.97]
n=135 randomised	60.9,65.1% Disease Status: NYHA	home: Monitoring signs& symptoms	Information	3mth 21.7,26.2% [calculation 14.49,17.03]
Self-identified Hispanics	II 17.4,20% III 44.9,47.7%	self-management support Referral to other HPs (includ.GP) &		6mth 31.9,33.8% [calculation 22.011,21.97]
were identified at 2 community hospitals close to US-Mexico border	IV 37.7,32.3%	social care Family involvement		LOS in days (mean) all ns
to de Mexico Border				1mth 0.59(2.3),1.41(5.5) 3mth 2.19(5.4),2.4(6.2) 6mth 3.65(7.8),3.4(7.1)
Thompson ²⁷ 2005	Age: 73(14),72(12) % Female:38,27%	Clinic & home based intervention (n=58)	Usual care (n=48)	No. of patients experiencing one or more readmissions
UK	Ethnicity:No details Disease Status:	Components of intervention at	Components of intervention at home:	13,21
Randomisation was at GP practice level	NYHA III &IV 76,73%	home: Monitoring signs& symptoms	unknown	Total no. of readmissions 15,45
Patients recruited from 2 North of England general		Education/self-management support Family involvement In outpatient clinic		Total no of hospital days
hospitals following an admission		Monitoring signs& symptoms Education/self-management support		P<0.01 for all at 6 mths
admission		Family involvement Referral to other health and social care		1 -0.0 Fibrial at 6 mail
Jaarsma ^{28,29} 2008	Age: 71(11),70(12),72(11) % Female:34,39,40%	Basic nurse support (BNS) (n=340)	Usual care group (n=339)	No. of readmissions 121,134,120 ns
The Netherlands	Ethnicity: No detail Disease Status: NYHA	Components of intervention: Outpatients	Components of intervention: Disease management	LOS in days
n=1049 randomised	II 51,48,54% III 47,48,42%	Education /Self-management support Patient directed access to HF nurse		(medians) 8.0(4,14),9.5(5,17),12(5,19.5)
All patients had been admitted to hospital with	IV3,4,4%	Intensive nurse support (INS)		P<0.01 between BNS grp & control but ns between INS grp &
symptoms of HF		(n=344) Components of intervention at	_	control
		home: Patient directed access to HF nurse		
		Referral to other health & social care Education/Self-management support Equipment		
Brotons ³¹ 2009	Age: 76.6(7.5),76(8.9)yrs. % Female: 54.2,56.1%	Home based intervention (n=144)	Usual care (n=139)	No. of readmissions 52, 62 ns
Spain n=283 randomised	Ethnicity: Not reported Disease Status: NHYA	Components of intervention at home:	Components of intervention: not known	Mean no. of readmissions
Patients were recruited by	I 42.4,55.4% II 52.1,37.4%	Monitoring signs& symptoms Education/Self-management support		1.01,1.3 ns
well-trained nurses at 2 university hospitals	III 4.9,5.8% IV 0.7,1.4%	Medication review (nurse, physician, cardiologist)		
		Referral to physician or cardiologist as necessary		

1	
3	
4	
5	
6	
7 8	
8	
1	0 1
1	
1	3
1	4
1	5
1	6
1	ر و
1	34567890123
2	0
2	1
2	2
2	3
2	4
2	5
2	4 5 6 7 8 9
2	8
2	9
-2	n
3	1
3	1 2
3 3 3 3 3 3	1 2 3 4 5 6 7
3 3 3 3 3 3 3	12345678
3 3 3 3 3 3 3 3	123456789
333333334	1234567890
333333344	12345678901
3333333444	123456789012
333333344	1234567890123
3333333444444	123456789012345
3333333344444444	1234567890123456
33333333444444444	12345678901234567
333333334444444444	123456789012345678
33333333444444444444	1234567890123456789
33333333444444444445	12345678901234567890
3333333344444444445555	1234567890123456789012
33333333444444444455555	12345678901234567890123
33333333344444444444555555	123456789012345678901234
33333333444444444455555555	1234567890123456789012345
33333333444444444455555555	123456789012345678901234
3 3 3 3 3 3 3 3 4 4 4 4 4 4 4 4 4 4 5 5 5 5	123456789012345678901234567
3 3 3 3 3 3 3 3 4 4 4 4 4 4 4 4 4 4 5 5 5 5	123456789012345678901234567

Stewart 32	Home vs. Clinic	Home based intervention (n=143)	Clinic-based intervention (n=137)	Rates of readmissions/100
2012	Age: 70 (15), 73(13) yrs	interest (ii 140)	(ii 107)	days/patient
WHICH trial	% Female:27,28%	Components of intervention at	Components of intervention:	
Australia	Ethnicity: No details	home:	In clinic	0.52±0.76,0.53±1.02 ns
	Disease Status: NYHA II or III	Monitoring signs& symptoms	Disease management	
n=280 randomised	83,88%	Family involvement	Assessment of home environment Family involvement?	Mean days of hospitalisation
Patients admitted to	Months since CHF diagnosis 34.6(55.3),44.8(71.0)	CM meetings/feedback to other health providers	Referral to other health or social care	Mean days of nospitalisation
participating hospitals	34.6(55.3),44.6(71.0)	Referral to other health or social care	CM meetings/feedback to other	4.96±8.57,3.62±6.36 ns
were screened for study		Assessment of home environment	health providers	
eligibility		Medication review (nurse, pharmacist,	·	At 12-18mths
11	ļ	physician, cardiologist)		
Hospital Initiated CM - NRCTs				
Riegel ³⁵	Age: 74.44 yrs. (10.65), 70.77	Multidisciplinary disease	Usual Care (n=120)	Readmission rates
2000	(11.77)	management (DM) (n=120)	Osual Care (II-120)	3mths
JSA	% female:55, 55%		Components of intervention at	0.22 (0.52),0.13 (.45) (ns)
n=240 were randomised	Ethnicity: No details	Components of intervention at	home:	6mths
	Disease Status: NYHA	home:	Disease management	0.32(0.58),0.23(0.53) (ns)
Patients were recruited from 5 hospitals following	I 19.2,24.2% II 26.7,18.3%	Monitoring signs& symptoms Support group		LOS in days
a hospitalisation for HF.	III 43.3,44.2%	Referral to specialist RN visits		3mths
a noopitalisation for the	IV 10.8,13.3%	receital to openialist KIV Visits		0.89 (3.34), 0.48(1.64) (ns)
				6mths
				1.31(3.77), 1.08(3.46) (ns)
Russell ³⁶	Age:79.4(10.7),79.9(10.7)	Transitional care service (n=223)	Usual home care services (n=224)	Readmissions
2011	% Female:55.6,57.6 (numbers)	Components of intervention at	Components of intervention of	Upadinated add 20 de
JSA n=447	Ethnicity: White non-Hispanic 56.9,58.4	home:	Components of intervention at home:	Unadjusted odds ratio 30 days 0.58 (0.38,0.88) p<0.01
n -1 1	African-American 17.0,16.5	Self-management support	Nurse visits	σ.σσ (σ.σσ,σ.σσ) μ~σ.στ
Patients were referred	Hispanic 14.8,14.	Referral to other health & social care	Physical therapy (44.6)	Adjusted odds ratio 30 days
from a single large not-for-	Asian/other 11.2,10.7	Assessment of home environment	Home health aide service (27.7)	0.57 (0.38,0.87) p<0.01
profit general medical and	Disease Status: Patients with	CM meetings/feedback to other health		
surgical hospital	a primary or secondary	providers		
	diagnosis of CHF	Advance care planning Physical therapy		
Stauffer ³⁷	Age: 78.9(8.3), 81.4(8.3)	Nurse-led transitional care	Control group (n=84)	Readmission rate at 30 days
2011	% Female:58.1, 54.8%	intervention (n=56)	Control group (ii 64)	12.6 (7.4,17.8) difference -12.6, %
JSA	Ethnicity: Hispanic ethnicity		Components of interventions:	change -48%; 16.4(14,18.7)
	7.1,3.6%	Components of intervention at	unknown	difference -1.6 % change 11%
n=140	Disease Status:	home:		
Patients were screened	APR-DRG severity of illness	Monitoring signs& symptoms		
for eligibility within 48 hours of hospital	1 5.4,1.2% 2 44.6,31%	Education/Self-management support Family involvement		
admission	3 37.5,57.1%	Referral (Assessing availability of		
44111661611	4 12.5,10.7%	social care)		
		Patient directed access to study		
		personnel		
Community-initiated CM-				
RCTs				
Peters-Klimm 39	Baseline characteristics of	Case management (n=97)	Usual care (n=100)	No. of admissions
2010	participants: CM,UC	. ,		(baseline 36 vs. 35)
Germany	Age: 70.4yrs (10.0), 68.9 (9.7)	Components of intervention at	Components of control	
	% Female:29, 26%	home:	intervention:	18 vs. 9 at 12 mths (ns)
n=199 at randomisation	Ethnicity: No details Disease Status: NYHA	Monitoring signs& symptoms Education/Self-management support	Disease management Education	No. of patients experiencing one
n= 199 at randomisation	I 1(1.0), 5 (5)	Medication review (CM/GP)	Education	or more CHF admissions
Recruitment was via	II 63 (64.9),67 (67)	Referral to GP		11 vs. 7 at 12 mths (ns)
general practice by mail.	III 33 (34),27 (27)			
•	IV 0, 1(1.0)			
	Mean yrs. with CHD			
Wade⁴²	6.2(4.6) (n=79),6.8(6.3)(n=74 Age: 75.8, 77.7 yrs.	Case management (n=152)	Tele health with case management	No data available for primer:
wade 2011	**Mage: 75.8, 77.7 yrs.** **Mage: 75.8, 77.7 yr	Case management (n=152)	(THCM) (n=164)	No data available for primary outcome but described as ns
JSA	Ethnicity: Black/African	Components of intervention at	(511) (11-10-1)	oatoonie but described as its
n=2,200 were randomised	American	home:	Components of intervention:	The participant population overall
	24,20.4%	Referral to other health & social care	Disease management	had 42% fewer inpatient days
Aetna Medicare	Disease Status (SD):	Equipment	Education	during the intervention period
Advantage members with medical & pharmacy	No detail		Referral to other health & social care	compared with the previous year. No data
medical & pharmacy benefits were identified	1			ivo uata
through analysis of claims	1			
Hancock ⁴³	Age: 85.1 (6.7), 81.8 (7.1)yrs	Case management (n=16)	Routine GP-led care (n=12)	No. of admissions
2012	% Female:56%, 58%	1		at 6 & 12 months
JK 	Ethnicity:100% white British Disease Status:	Components of interventions at	Components of intervention:	0.0 = 1.0==1h=
n=28 randomised	Disease Status:	home: Monitoring signs and symptoms	Disease management	0, 0 at 6mths 0.0 at 12 mths
Residents from 33 of 35	10:1:4:1, 5:4:1:1	Education		0,0 at 12 mins
ong-term residential &	,	CM meetings/feedback to other health		
nursing homes	1	providers		
-	1	Medication review		
Community initiated	ļ	(CM/GP/cardiologist)		
Community –initiated CM -NRCTs				
СМ -NRCTS Bonarek- Hessamfar ⁴⁴	Age: Median 78, 80 yrs.	Co-ordinated care via	Usual care (n=233)	No. of patients experiencing at
2008	% Female: No details	multidisciplinary network (n=129)		least one admission
rance	Ethnicity: No details		Components of intervention: not	26,58
1=362	Disease Status: NHYA	Components of intervention at	known	Total no. of admissions
1-302		home:	i	35,96
Compared patients	Median of			
Compared patients ncluded prospectively	III, IV	Monitoring signs& symptoms		Median LOS
Compared patients ncluded prospectively from Jan 1st 2004- Dec		Monitoring signs& symptoms Education (diet)		Median LOS 9.2,11.7 days
Compared patients ncluded prospectively from Jan 1st 2004- Dec 31st 2005 from GP list		Monitoring signs& symptoms		Median LOS


Lowery ⁴⁵	Age: 65.4(0.51),67.4(0.45) yrs.	Nurse-practitioner-led disease	Usual care (n=510)	Mean no. of readmissions	
2012	% female:1, 1%	management model (n=457)		1yr	
USA	Ethnicity:		Components of intervention: not	0.7(0.32), 0.23(0.65) p<0.001	
n=1043	White 71.2,79.9%	Components of intervention at	known	(417,428)	
	Black 24,16.1%	home:		2yr	
Intervention implemented	Other 4.8,4.0%	Location was lead tertiary centre,		0.15 (0.58), 0.13(0.42) ns	
in 4 Midwest VA medical	Disease Status: No details	other medical centres (some primary		(384,382)	
centres from the same		care) or one affiliated outpatient clinic.			
region & one affiliated				Mean no. of days in hospital	
outpatient clinic and 2 VA		Monitoring signs& symptoms		1yr	
medical centres served as		Education /Self-management support		0.37(2.25),0.97(3.15) p=0.0014	
control.		Referral to other health & social care		2yr	
		Family involvement		0.86 (3.98),0.66(2.74) ns	
				I .	1

Key: AF atrial fibrillation, CHF Chronic heart failure, COPD chronic obstructive pulmonary disease, CM case management or case manager, HOCM/RCM Hypertrophic Obstructive/Restrictive Cardiomyopathy, GP general practitioner, UC usual care, NYHA, LV left ventricular SNF skilled nursing facility, PAD peripheral arterial disease.

Table 3: Available cost data from studies (n=9)

Study	Cost data Intervention vs. control
	(ns= not statistically significant)
Rich 1995	3 month data
	Study intervention cost US\$216 per person
	Hospital readmissions \$2178 vs. 3,236 p=0.03
Stewart 1998	6 month data
otomati 1000	Cost of study intervention Aus\$190 per person
	Mean cost of hospital based care
	\$3200 (1800-4600), 5400 (3200-6800) ns
	Cost of community based care
	\$620(460,740), 680(550,800) ns
Stewart 1999	6 and 18 month data
Otomari 1000	6 months
	Total hospital based care Aus\$490,300 vs. 922,600 ns
	18 month data
	Total hospital based care \$ 5100 (6800) vs.10,600(13000) ns
Laramee 2003	3 month data
Laramee 2003	Total care costs
	Mean(US\$) 23,054 vs.25,536 ns
Naylor 2004	12 month data
Naylor 2004	CHF readmissions US \$175.840 vs. 498.110?
	Physician's office (outpatients)\$4,549, 5,169 ns
	ER visits \$1780, vs. 5650 ns
	Home visits (all cause)
	Visiting nurse \$11021, 64,531 p<0.001
	APN \$104,019 vs. 0
	Physical therapist \$7,120 vs. 10,918 ns
	Social worker \$178.vs. 534 ns
	Home health aides \$9,167 vs. 11,081 ns
	Total home visits \$138,649 vs. 97.883 p<0.001
	Total costs \$725,903, 1,163,810 ns
Stewart 2012	12-18 month data
Stewart 2012	
	Costs per patient
	Au\$1813(220) vs. 1829(174) ns
	Total costs Au \$3.93 million vs. 5.53 million p=0.03 for median costs per day
D'1 0000	3 & 6 month data
Riegel 2000	
	Total costs
	3mths US\$ 632 (2,378) vs. 317 (1,188) ns
D' I 0000	6mths \$1,024(3,017) vs. 686(2,225) ns
Riegel 2006	1, 3 & 6 month data
	HF inpatient costs all ns
	1mth US\$1012(4022) vs. 2830 (13,896)
	3mth \$3045(7784) vs. 4130 (14,468)
	6mth \$5567(13,137) vs. 6151 (16,650)
Stauffer 2011	1 month data
	'under the current payment system, the intervention reduced the hospital financial
	contribution on average by US\$227for each Medicare patient with HF'

Figure 1: PRISMA Flow Diagram



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.

215x279mm (300 x 300 DPI)

Figure 2b EPOC Risk of bias for Non-randomised controlled trials (NRCTs)

Study name	Allocation sequence adequatel y generated	Allocation adequatel y concealed	Baseline outcome measurement s similar	Baseline characteristic s similar	Incomplet e outcome data	Knowledge of the allocated intervention s	Protected against contamination	Selectiv e outcome reporting	Other risks of bias
Bonarek- Hessamfa r 2008	High risk	High risk	Unclear risk	High risk	Unclear risk	Low risk	Unclear	High risk	High risk
Lowery	High	High	Unclear	Low	High	Low	Unclear	Low	High
2012	risk	risk	risk	risk	risk	risk	risk		risk
Riegel 2000	High risk	High risk	Unclear risk	Unclear Risk/High risk	Unclear risk	Low risk	Unclear risk	Low risk	High risk
Russell	High	High	Unclear	Low	Unclear	Low	Unclear	Low	High
2011	risk	risk	risk	risk	risk	risk	risk		risk
Stauffer	High	High	Unclear	Low	Unclear	Low	Unclear	Low	Low
2011	risk	risk	risk	risk	risk	risk	risk	risk	risk

297x209mm (300 x 300 DPI)

Figure 3: CHF admissions data

a) Overall for hospital -initiated CM

			CM	Usual Care		Rate Ratio	Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Blue 2001	-0.129	0.248	84	81	7.8%	0.88 [0.54, 1.43]	-
Brotons 2009	-0.211	0.188	144	139	9.3%	0.81 [0.56, 1.17]	-
DeBusk 2004	-0.098	0.157	228	234	10.1%	0.91 [0.67, 1.23]	+
Jaarsma 2008	0.076	0.151	344	339	10.3%	1.08 [0.80, 1.45]	+
Laramee 2003	-0.201	0.321	131	125	6.2%	0.82 [0.44, 1.53]	-
Naylor 2004	-0.563	0.197	118	121	9.1%	0.57 [0.39, 0.84]	-
Rich 1993	-0.136	0.332	63	35	6.0%	0.87 [0.46, 1.67]	
Rich 1995	-0.825	0.245	142	140	7.9%	0.44 [0.27, 0.71]	-
Riegel 2000	0.054	0.164	140	140	0.0%	1.06 [0.77, 1.46]	
Riegel 2002	-0.675	0.219	130	228	8.5%	0.51 [0.33, 0.78]	-
Riegel 2006	0.112	0.24	69	65	8.0%	1.12 [0.70, 1.79]	· ·
Stewart 1999	0.054	0.164	140	140	10.0%	1.06 [0.77, 1.46]	+
Thompson 2005	-1.288	0.298	58	48	6.7%	0.28 [0.15, 0.49]	
Total (95% CI)			1651	1695	100.0%	0.74 [0.60, 0.92]	•
Heterogeneity: Tau ² :	0.10: Chi ² = 35.5	9 df = 1	1 (P =	0.00025; (*= 6	9%		Land In the state of the state
Test for overall effect							0.01 0.1 1 10 100
	The second second					1	Favours [experimental] Favours [control]

Sensitivity analysis: (Riegel 2002 removed) I2=68% RR 0.77 [0.61, 0.96] p=0.02

b) 3 month data for hospital -initiated CM

Study or Subgroup	log[Rate Ratio]	SE	Case management Total		Weight	Rate Ratio IV, Random, 95% CI	Year	Rate Ratio IV, Random, 95% CI
Rich 1993	-0.136	0.332	63	35	16.1%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	26.2%	0.44 [0.27, 0.71]	1995	
Riegel 2000	0.486	0.318	120	120		Not estimable	2000	
Riegel 2002	-0.445	0.268	130	228	22.9%	0.64 (0.38, 1.08)	2002	-
_aramee 2003	-0.201	0.321	131	125	17.0%	0.82 [0.44, 1.53]	2003	
Riegel 2006	-0.109	0.313	69	65	17.8%	0.90 [0.49, 1.66]	2006	-
Total (95% CI)			535	593	100.0%	0.67 [0.51, 0.90]		•
Heterogeneity: Tau ² =	0.02; Chi ² = 4.92	df = 4	(P = 0.30); P = 19%					ter de de de
Fest for overall effect	Z = 2.73 (P = 0.00	(36)					F	0.01 0.1 1 10 100 svours [Case management] Favours [Usual care]

Sensitivity analysis: (Riegel 2002 removed) I² = 39% RR 0.70 [0.48, 1.01] p=0.06

c) 6 month data for hospital -initiated CM

			Case management	Usual Care		Rate Ratio	Rate Ratio	
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% C	IV, Random, 95	% CI
Riegel 2000	0.054	0.164	140	140		Not estimable		
Riegel 2002	-0.675	0.219	130	228	25.5%	0.51 [0.33, 0.78	-	
Riegel 2006	0.112	0.24	69	65	24.4%	1.12 [0.70, 1.79		
Stewart 1999	-0.551	0.152	100	100	28.6%	0.58 (0.43, 0.78		
Thompson 2005	-1.200	0.290	50	40	21.5%	0.20 (0.15, 0.49	i —	
Total (95% CI)			357	441	100.0%	0.56 [0.35, 0.89]	•	
Heterogeneity: Tau ^a =	= 0.17; Chi ² = 14.0	6. df = 3	(P = 0.003); P= 79%				to. J.	J
Test for overall effect	Z = 2.43 (P = 0.01)					0.01 0.1 1 Favours (experimental) Favo	10 10 urs [control]

Sensitivity analysis: (Riegel 2002 removed) I² = 85% RR 0.57 [0.30, 1.21] p=0.1

d) 12-18 months for hospital -initiated CM

	og[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Bluc 2001	0.129	0.248	84	81	9.5%	0.88 [0.54, 1.43]	-
Brotons 2009	-0.211	0.188	144	139	14.5%	0.81 [0.56, 1.17]	-
DeBusk 2004	-0.098	0.157	228	234	18.5%	0.91 [0.67, 1.23]	+
Jaarsma 2008	0.076	0.151	344	339	19.4%	1.08 [0.80, 1.45]	-
Naylor 2004	-0.563	0.197	118	121	13.6%	0.57 [0.39, 0.84]	-
Stewart 1999	-0.279	0.122	100	100	24.5%	0.76 [0.60, 0.96]	-
Total (95% CI)			1018	1014	100.0%	0.83 [0.70, 0.98]	•

209x297mm (300 x 300 DPI)

Figure 4: CHF length of stay

a) Overall for hospital -initiated CM

	Case n	nanager	nent	Usi	ial car	e		Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Blue 2001	3.42	12.2	82	7.48	16.6	75	2.5%	-4.04 [-8.63, 0.55]	
Laramee 2003	6.9	8.5	131	9.5	9.8	125	9.1%	-2.60 [-4.65, -0.55]	-
Rich 1993	4.3	1.1	63	5.7	2	35	21.2%	-1.40 [-2.12, -0.68]	•
Rich 1995	3.9	10	142	6.2	11.4	140	6.8%	-2.30 [-4.80, 0.20]	
Riegel 2000	1.31	3.77	120	1.08	3.46	120	0.0%	0.23 [-0.69, 1.15]	
Riegel 2002	1.1	3.1	130	2.1	4.6	228	20.3%	-1.00 [-1.80, -0.20]	
Riegel 2006	3.65	7.8	69	3.4	7.1	65	6.8%	0.25 [-2.27, 2.77]	-
Stewart 1999	0.9	0.9	100	2.9	2.9	100	22.5%	-2.00 [-2.60, -1.40]	
Stewart 2012	4.96	8.57	143	3.62	6.36	137	10.9%	1.34 [-0.42, 3.10]	*
Total (95% CI)			860			905	100.0%	-1.28 [-2.04, -0.52]	•
Heterogeneity: Tau2=	0.58; Chi	P= 18.7	5, df = 7	(P = 0.0	009); P	= 63%			-do do do do
Test for overall effect	Z = 3.29 (P = 0.00	11)					Fa	-20 -10 0 10 20 avours [case management] Favours [usual care]

Sensitivity analysis: (Riegel 2002 removed) I² = 14% MD -1.76[-2.29,-1.23] p<0.00001

b) 3 month data for hospital -initiated CM

	Casen	anagen	nent	Ust	ial car	e		Mean Difference		Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
Rich 1993	4.3	1.1	63	5.7	2	35	37.0%	-1.40 [-2.12, -0.68]	1993	•
Rich 1995	3.9	10	142	6.2	11.4	140	5.5%	-2.30 [-4.80, 0.20]	1995	
Riegel 2000	0.89	3.34	120	0.48	1.64	120		Not estimable	2000	
Riegel 2002	0.85	2.3	130	1.6	3.9	228	41.0%	-0.75 [-1.39, -0.11]	2002	-
Laramee 2003	6.9	6.5	131	9.5	9.8	125	8.0%	-2.60 [-4.65, -0.55]	2003	
Riegel 2006	2.19	5.4	69	2.4	6.2	65	8.5%	-0.21 [-2.18, 1.76]	2006	+
Total (95% CI)			535			593	100.0%	-1.18 [-1.79, -0.57]		•
Heterogeneity: Tau*=	0.13; Chi	= 5.54.	df= 4 (P = 0.24); z = 1	28%			-	J. t 1 t t
Test for overall effect:	Z = 3.77	P = 0.00	02)						F	-10 -5 0 5 10 loace management(Favours (Usual care)

Sensitivity analysis: (Riegel 2002 removed) I² = 6% MD -1.46 [-2.15,-0.77] p<0.0001

c) 6 month data for hospital -initiated CM

	Case m	anager	nent	Usi	ial car	e		Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI
Riegel 2000	1.31	3.77	120	1.08	3.46	120		Not estimable	
Riegel 2002	1.1	3.1	130	2.1	4.6	228	44.9%	-1.00 [-1.80, -0.20]	-
Riegel 2006	3.65	7.8	69	3.4	7.1	65	12.1%	0.25 [-2.27, 2.77]	
Stewart 1999	0.9	0.9	49	2.9	2.9	48	43.0%	-2.00 [-2.86, -1.14]	-
Total (95% CI)			248			341	100.0%	-1.28 [-2.25, -0.30]	•
Heterogeneity: Tau*=	0.38; Chi	= 4.47	df= 2 (P = 0.11); P=	55%		-	
Test for overall effect								Egun	-10 -5 0 5 10 ours Icase managementi Favours IUsual carel

Sensitivity analysis: (Riegel 2002 removed so n=2) I² = 63% MD -1.20 [-3.31,0.91] p=0.26

209x297mm (300 x 300 DPI)

2

4

5 6

7 8 9

10

11 12

13

14 15

16

17 18

19

20 21

22

23 24

25

26 27

28

29 30

31

32 33

34

35

36 37

38 39

40

41 42

43

44

45 46

47

48 49

50

51 52

53

54 55

56

Appendix 1:

Database: Medline In-process - Current week, Medline 1950 to present

Search Strategy:

- 1 randomized controlled trial.pt. (376608)
- 2 random\$.tw. (717987)
- 3 control\$.tw. (2630023)
- 4 intervention \$.tw. (556397)
- 5 evaluat\$.tw. (2214967)
- 6 or/1-5 (5041451)
- 7 Qualitative Research/ (20094)
- 8 semi-structured questionnaire.mp. (1162)
- 9 observation methods.mp. (152)
- 10 Observation/mt [Methods] (635)
- 11 Nvivo.mp. (639)
- 12 interview/ (25018)
- 13 Personal Narratives/ (877)
- 14 Focus Groups/ (16824)
- 15 patient experience*.mp. (8525)
- 16 or/7-15 (70071)
- 17 exp Heart Failure/ (87270)
- 18 exp Heart Failure, Diastolic/ (496)
- 19 exp heart failure, systolic/ (789)
- 20 exp Ventricular Dysfunction/ (26332)
- 21 chronic heart failure.mp. (11341)
- 22 congestive heart failure.mp. (33082)
- 23 cardiac failure.mp. (10151)
- 24 LV dysfunction.mp. (2827)
- 25 left ventricular dysfunction.mp. (9373)
- 26 left ventricular impairment.mp. (188)
- 27 diastolic impairment.mp. (121)
- 28 systolic impairment.mp. (93)
- 29 or/17-28 (135885)
- 30 exp Case Management/ (8326)
- 31 exp Patient Care Planning/ (52319)
- 32 organisation of care.mp. (367)
- 33 community matron.mp. (44)

- 34 "Continuity of Patient Care"/ (14497)
- 35 Community Health Nursing/ (18371)
- 36 transit* care.mp. (580)

2

4 5

6

7 8

9

10 11

12

13 14

15

16 17

18

19 20

21

22 23

24

25 26

27

28 29

30

31 32

33

34 35

36

37

38 39

40 41

42

43

44 45

46

47 48

49 50

51

52


53 54

55

56 57


- 37 Interdisciplinary Communication/ (10602)
- 38 Patient Discharge/ (18977)
- 39 discharge plan.mp. (176)
- 40 exp Patient Care Management/ (535496)
- 41 Comprehensive Health Care/ (6078)
- 42 exp Managed Care Programs/ (38918)
- 43 Primary Health Care/ (54234)
- 44 Community Health Services/ (26923)
- 45 General Practitioners/ (1943)
- 46 Family Practice/ (60223)
- 47 Physicians, Family/ (14745)
- 48 multidisciplinary.mp. (44988)
- 49 or/30-48 (674050)
- 50 6 or 16 (5094326)
- 51 29 and 50 (47009)
- 52 49 and 51 (2590)
- 53 52 not (case report/ or case study/ or letter/ or editorial/ or expert opinion.mp.) (2489)
- 54 53 not (Algeria\$ or Egypt\$ or Liby\$ or Morocc\$ or Tunisia\$ or Western Sahara\$ or Angola\$ or Benin or Botswana\$ or Burkina Faso or Burundi or Cameroon or Cape Verde or Central African Republic or Chad or Comoros or Congo or Diibouti or Eritrea or Ethiopia\$ or Gabon or Gambia\$ or Ghana or Guinea or Keny\$ or Lesotho or Liberia or Madagasca\$ or Malawi or Mali or Mauritania or Mauritius or Mayotte or Mozambiq\$ or Namibia\$ or Niger or Nigeria\$ or Reunion or Rwand\$ or Saint Helena or Senegal or Seychelles or Sierra Leone or Somalia or South Africa\$ or Sudan or Swaziland or Tanzania or Togo or Ugand\$ or Zambia\$ or Zimbabw\$ or China or Chinese or Hong Kong or Macao or Mongolia\$ or Taiwan\$ or Belarus or Moldov\$ or Russia\$ or Ukraine or Afghanistan or Armenia\$ or Azerbaijan or Bahrain or Cyprus or Cypriot or Georgia\$ or Iran\$ or Iraq\$ or Israel\$ or Jordan\$ or Kazakhstan or Kuwait or Kyrgyzstan or Leban\$ or Oman or Pakistan\$ or Palestin\$ or Qatar or Saudi Arabia or Syria\$ or Tajikistan or Turkmenistan or United Arab Emirates or Uzbekistan or Yemen or Bangladesh\$ or Bhutan or British Indian Ocean Territory or Brunei Darussalam or Cambodia\$ or India\$ or Indonesia\$ or Lao or People's Democratic Republic or Malaysia\$ or Maldives or Myanmar or Nepal or Philippin\$ or Singapore or Sri Lanka or Thai\$ or Timor Leste or Vietnam or Albania\$ or Andorra or Bosnia\$ or Herzegovina\$ or Bulgaria\$ or Croatia\$ or Estonia or Faroe Islands or Greenland or Liechtenstein or Lithuani\$ or Macedonia or Malta or maltese or Romania or Serbia\$ or Montenegro or Slovenia or Svalbard or

Argentina\$ or Belize or Bolivia\$ or Brazil\$ or chile or Chilean or Colombia\$ or Costa Rica\$ or



Appendix 2: Subgroup-analysis of unplanned readmission data by intervention component- hospital initiated studies

a) Family involvement



Test for subgroup differences: $Chi^2 = 2.09$, df = 1 (P = 0.15), $I^2 = 52.1\%$


Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.98 df=1 (p=0.32) l²-0%

b) education & self-management

			Case management	Usual care	Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.20.1 Education and	self-managemer	nt					
Rich 1993	-0.136	0.332	63	35	0.87 [0.46, 1.67]	1993	+
Rich 1995	-0.825	0.245	142	140	0.44 [0.27, 0.71]	1995	+
Blue 2001	-0.129	0.248	84	81	0.88 [0.54, 1.43]	2001	+
Riegel 2002	-0.675	0.219	130	228	0.51 [0.33, 0.78]	2002	+
Laramee 2003	-0.201	0.321	131	125	0.82 [0.44, 1.53]	2003	+
DeBusk 2004	-0.098	0.157	228	234	0.91 [0.67, 1.23]	2004	+
Naylor 2004	-0.563	0.197	118	121	0.57 [0.39, 0.84]	2004	+
Thompson 2005	-0.1288	0.298	58	48	0.88 [0.49, 1.58]	2005	+
Riegel 2006	0.112	0.24	69	65	1.12 [0.70, 1.79]	2006	+
Jaarsma 2008	0.076	0.151	344	339	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	139	0.81 [0.56, 1.17]	2009	+
1.20.2 No education a	and self-manager	nent					
Stewart 1999	-0.279	0.122	100	100	0.76 [0.60, 0.96]	1999	+
Riegel 2000	0.054	0.164	140	140	1.06 [0.77, 1.46]	2000	
Stewart 2012	0.018	0.071	143	137	Not estimable	2012	
							0.002 0.1 i 10 50b
						Fa	avours [case management] Favours [Usual care]

There were inadequate data in the no education/self-management group to perform a comparison.

c) medication review




Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.36 df=1 (p=0.55) I²-0%

d) Referral to other services

			Case management	Usual care		Rate Ratio		Rate Ratio
Study or Subgroup	log[Rate Ratio]	SE	Total	Total	Weight	IV, Random, 95% CI	Year	IV, Random, 95% CI
1.16.1 Referral to oth	ers							
Stewart 1999	-0.279	0.122	100	100	18.6%	0.76 [0.60, 0.96]	1999	
Riegel 2000	0.486	0.318	120	120	0.0%	1.63 [0.87, 3.03]	2000	
Blue 2001	-0.129	0.248	84	81	12.8%	0.88 [0.54, 1.43]	2001	
Riegel 2002	-0.445	0.268	120	120	11.9%	0.64 [0.38, 1.08]	2002	
Thompson 2005	-1.288	0.298	58	48	10.8%	0.28 [0.15, 0.49]	2005	
Riegel 2006	0.112	0.24	69	65	13.1%	1.12 [0.70, 1.79]	2006	
Jaarsma 2008	0.076	0.151	344	339	17.3%	1.08 [0.80, 1.45]	2008	+
Brotons 2009	-0.211	0.188	144	0	15.5%	0.81 [0.56, 1.17]	2009	 +
Russell 2011	0	0	0	0		Not estimable	2011	
Stewart 2012	0.018	0.071	143			Not estimable	2012	
Subtotal (95% CI)			919	753	100.0%	0.77 [0.59, 1.00]		•
Heterogeneity: Tau² =	0.09; Chi ² = 19.6	7, $df = 6$	6 (P = 0.003); I ² = 69%					
Test for overall effect:	Z = 1.93 (P = 0.05	i)						
1.16.2 No referral								
Rich 1993	-0.136	0.332	63	35	13.7%	0.87 [0.46, 1.67]	1993	
Rich 1995	-0.825	0.245	142	140	19.6%	0.44 [0.27, 0.71]	1995	
Laramee 2003	-0.201	0.321	131	125	14.3%	0.82 [0.44, 1.53]	2003	
DeBusk 2004	-0.098	0.157	228	234	28.3%	0.91 [0.67, 1.23]	2004	-
Naylor 2004	-0.563	0.197	118	121	24.1%	0.57 [0.39, 0.84]	2004	
Subtotal (95% CI)			682	655	100.0%	0.69 [0.51, 0.93]		•
Heterogeneity: Tau ² =	0.06; Chi ² = 8.13 .	df = 4	(P = 0.09); I ² = 51%					
Test for overall effect:	Z = 2.47 (P = 0.01)						
								0.01 0.1 1 10 100
							Fa	vours [case management] Favours [usual care]
Test for subgroup diffe	erences: $Chi^2 = 0$.	28. df=	= 1 (P = 0.60), P = 0%					


Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.36 df=1 (p=0.55) l²=0%

e) Assessment of home environment



Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.46 df=1 (p=0.50) l²=0%


f) Case management/health professional meetings



rest for subgroup differences; Chin= 0.01, di = 1 (P = 0.91), in= 0%

Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.07 df=1 (p=0.07) l²=0%

g) Patient-directed access



Sensitivity analysis (removing Riegel 2002) Test for subgroup differences Chi² 0.25 df=1 (p=0.61) I²=0%



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4-5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	N/A
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Appendix one
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6-7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	6-7
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ² for each meta-analysis http://bmjopen.bmj.com/site/about/guidelines.xhtml	6-7



PRISMA 2009 Checklist

		Page 1 of 2	
Section/topic	#	Checklist item	Reported on page
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	6-7
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7 & Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	7-8
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	9
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	9-11
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	9-11 plus figures 3 & 4 and appendix 2
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	9-11 plus figures 3 & 4
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9-11 plus figures 3 & 4
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	12
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	12,14-15
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	15



PRISMA 2009 Checklist

FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	21

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. For me 10 doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

