

BMJ Open

The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Study

Journal:	BMJ Open
Manuscript ID	bmjopen-2017-016144
Article Type:	Protocol
Date Submitted by the Author:	27-Jan-2017
Complete List of Authors:	Goldsmith, Petra; Royal Liverpool and Broadgreen University Hospitals NHS Trust, Renal Transplant Surgery; University of Liverpool, Institute of Infection and Global Health Bottomley, Matthew; Oxford University Hospitals NHS Foundation Trust Okechukwu, Okidi; Central Manchester University Hospitals NHS Foundation Trust Ross, Victoria; NHS Greater Glasgow and Clyde Ghita, Ryan; NHS Greater Glasgow and Clyde Wandless, David; NHS Grampian Falconer, Stuart; NHS Lothian Papachristos, Stavros; Central Manchester University Hospitals NHS Foundation Trust Nash, Philip; King's College Hospital NHS Foundation Trust Androshchuk, Vitaliy; Oxford University Hospitals NHS Foundation Trust Clancy, Marc; NHS Greater Glasgow and Clyde; University of Glasgow
Primary Subject Heading:	Renal medicine
Secondary Subject Heading:	Surgery
Keywords:	Renal transplantation < NEPHROLOGY, TRANSPLANT SURGERY, TRANSPLANT MEDICINE

SCHOLARONE™
Manuscripts

STUDY PROTOCOL: The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Study

Authors:

Petra M Goldsmith^{1,2}

Matthew J Bottomley³

Okidi Okechukwu⁴

Victoria C Ross⁵

Ryan Ghita⁵

David Wandless⁶

Stuart J Falconer⁷

Stavros Papachristos⁴

Philip Nash⁸

Vitaliy Androshchuk³

Marc Clancy^{7,9}

Corresponding Author:

Petra M Goldsmith

Royal Liverpool University Hospitals NHS Trust

Prescot Street

LIVERPOOL

L7 8XP

United Kingdom

petra.goldsmith@googlemail.com

07795122575

On behalf of the Transplant Audit Collaborative

1. Renal Transplant Unit, Royal Liverpool University Hospitals NHS Trust, Liverpool, UK
2. University of Liverpool
3. Department of Nephrology, Oxford University Hospitals NHS Foundation Trust, Oxford, UK
4. Department of Transplantation, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK
5. Department of Transplantation, NHS Greater Glasgow and Clyde, Glasgow, UK
6. Department of Nephrology, NHS Grampian, Aberdeen, UK
7. Department of Transplantation, NHS Lothian, Edinburgh, UK
8. Department of Nephrology, King's College Hospital NHS Foundation Trust, London, UK
9. University of Glasgow, Glasgow, UK

WORD COUNT 1,537

ABSTRACT

Introduction: High intra-patient variability (IPV) in tacrolimus trough levels has been shown to be associated with higher rates of renal transplant failure. There is no consensus on what level of IPV constitutes a risk of graft loss. The establishment of such a threshold could help to guide clinicians in identifying at risk patients to receive targeted interventions to improve IPV and thus outcomes.

Methods and Analysis: A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study. The aim is to recruit a minimum of 1,600 patients with follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss. The study also aims to determine any national or regional trends in IPV and any demographic associations.

Ethics and Dissemination: Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The study will be registered locally in each participating centre in line with local Research and Development protocols. It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

ARTICLE SUMMARY

Strengths and Limitations of this Study

1. This is a multicentre collaborative study comprised of units across the UK which are both Transplant Centres and referring Nephrology Units
2. The sample size will be the largest investigating IPV to date
3. The study investigates retrospective data and requires a lengthy follow up period which will lead to some exclusions
4. There will be local variations in the laboratory assay that cannot be corrected for in the IPV calculations

INTRODUCTION

The addition of calcineurin inhibitors (CNI) as maintenance immunosuppressants has improved renal transplant 1 year survival rates since the 1980s.[1] Tacrolimus emerged as a viable alternative to Ciclosporin in the 1990s.[1] In 2005, a meta-analysis was published on randomised trial data comparing Tacrolimus and Ciclosporin as primary immunosuppressants in renal transplant, observing a 44% reduction in death censored graft loss with Tacrolimus over Ciclosporin.[2] In 2007, the Symphony Study reported favourable graft survival and function, and reduced biopsy proven rejection with low dose Tacrolimus over low dose Ciclosporin, Sirolimus or standard dose Tacrolimus.[3]

CNIs have a narrow therapeutic index: too little exposure places a transplant recipient at increased risk of acute rejection and donor specific antibody formation. Too much exposure and a transplant recipient is placed at increased risk of malignancy, infection, nephrotoxicity and unacceptable side effects such as tremor.[4-7]

Trough levels are used as a proxy for oral bioavailability of CNIs and vary both between patients (inter-patient variability) and for an individual over time (intra-patient variability, IPV). Between individuals, age, gender, ethnicity, body mass index, genetic polymorphisms in CYP3A5 and CYP3A4, drug interactions, adherence, liver function and lifestyle choices account for the differences. Similarly, IPV is affected by adherence, gastrointestinal metabolism and motility, diarrhoea, food and drug interactions, synchronicity of dose administration and blood test and variability of the laboratory assay.[8-15]

An emerging body of evidence is being established indicating favourable graft function, survival and fewer rejection episodes up to 1 year post transplant for patients demonstrating low intra-patient variability.[16-18] Similarly, high IPV has been associated with poorer outcomes and graft survival.[20-21] Donor age and previous transplants appear to be risk factors for a high intra-patient variability.[18] However, little data exists on the long term impact of high IPV and studies have not yet been able to draw conclusions about risk thresholds of variability because of limitations in sample size.

Objectives:

- To establish important baseline data about national and regional trends in IPV
- To investigate demographic associations and other characteristics for patients in high and low variability groups
- To formulate a “danger” threshold for IPV, above which a patient is deemed at risk of adverse outcomes so they can then be targeted for intervention prior to organ damage or failure

METHODS AND ANALYSIS

A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. It is the first Collaborative of its kind, facilitating the development of this largest study examining IPV to date. The TAC is comprised of junior doctors with an interest in Nephrology and/or Transplantation. It is supported by Consultant Medics and Surgeons in these fields to undertake research and audit projects related to transplantation. Any UK NHS Trust

involved in the after care of renal transplant recipients is eligible to register for this study providing they have a transplant or nephrology doctor willing to enrol in the TAC.

Patient Selection

Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study (table 1). To be enrolled, patients are required to have follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss.

Table 1: Inclusion / Exclusion Criteria

(T1=6-12 months post-transplant; T2=most recent 12 months)

INCLUSION CRITERIA	EXCLUSION CRITERIA
Age ≥ 18 at time of transplant	Age ≤ 17 at time of transplant
A functioning graft* at 2 years	Failed graft before 2 years
Renal only transplant	Non-renal transplant
Short acting Tacrolimus preparation only during study period	Modified release preparation at any point during study period
Tacrolimus as primary immunosuppressant	Other primary immunosuppressant (e.g. ciclosporin, sirolimus, other non-steroid)
Alive minimum 2 years following transplant	Pregnancy at any point during study period
	Death prior to 2 years following transplant
	Non-standard transplant (ABO or HLA incompatible; requiring desensitisation)
	Patients with fewer than 4 tacrolimus trough levels for T1 and T2

Sample Size

A large sample size is needed to provide meaningful numbers to establish variability risk cut offs. It is estimated that a minimum of 8 UK based centres will participate in the study. If each centre, on average, supplies data for 200 patients, a minimum dataset of 1,600 patients will be achieved.

It is, however, recognised that there will be a significant variation between the numbers of eligible patients available to each centre (tertiary transplant units will naturally have access to larger numbers than referring District General Hospitals). The set number of 200, therefore, is only to be used as a guide and it is recognised that this may not be achievable for smaller, secondary nephrology units. Similarly, we welcome larger numbers from any centre able to do so.

Data Collection and Information Governance

The TAC has established a standard minimum dataset (see Table 2) which each site will use as the basis for data collection. A template database will be provided to a representative of the study team from each NHS Trust participating in the study. This database will be password protected and held on secure local Trust servers. All data should be entered onto this database – the use of hard copy proformas associated with this audit is discouraged to avoid breaching data protection policies.

Table 2: Minimum Dataset

(T1=6-12 months post-transplant; T2=most recent 12 months)

DATA SET

- Date of transplant
- Recipient and Donor Ages
- Gender
- Ethnicity
- Primary Renal Diagnosis
- Previous transplants
- Mismatch Grade (A, B, DR)
- Type of donor
- eGFR at the end of T1 and T2
- Creatinine at 12 months and at the end of T2
- All Tacrolimus trough levels during T1 and T2
- Urinary protein creatinine ratio at the of end T1 and T2
- Graft Failure
- Delayed Graft Function
- Tacrolimus dosing at month 12 and at the end of T2
- Other immunosuppression at T1 and T2 – steroid Y/N; Antiproliferative Y/N
- Induction agent
- Denovo Donor Specific Antibody status post-transplant and level (mean fluorescence intensity)
- Biopsy proven acute rejection
- Death

The data collection database will be anonymised and will contain no patient identifiable data. Each hospital will be issued with patient study identifiers for all patients included in the study. A separate password protected spreadsheet of the study identifiers with the corresponding hospital numbers should be securely stored locally by each participating centre for local reference only should there be any difficulties or queries regarding data collection.

The anonymised master database will be compiled by a dedicated, named member of the collaborative. It will be shared with all members of the Transplant Audit Collaborative using secure NHS email only (either local Trust email or NHS.net) and will be held on secure, password protected Trust servers only.

A full data analysis will be conducted in conjunction with an appropriately qualified medical statistician.

Permissions and Registration

Each participating centre will be expected to complete a site registration form. A named member of the TAC from each participating centre will be responsible for gaining all necessary local Trust permissions and study registrations as required by local Research and Development and Audit offices.

Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The findings of the study are not expected to impact upon individual patient care.

ETHICS AND DISSEMINATION

As this study does not fulfil the NHS Health Research Authority and Medical Research Council’s criteria for research, formal ethical approval is not needed. However, all local NHS Trust approvals and registrations will be sought from each participating centre.

It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

CONTRIBUTORSHIP STATEMENT

All authors fulfil the ICMJE criteria for authorship and agree to be accountable for all aspects of the work. In brief:

Petra M Goldsmith wrote the protocol, produced revisions and was involved in the original concept, study design and implementation.

Matthew J Bottomley made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Okidi Okechukwu made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Victoria C Ross made revisions to the protocol and was involved in the original concept, study design and implementation.

Ryan Ghita made revisions to the protocol and was involved in the original concept, study design and implementation.

David Wandless made revisions to the protocol and was involved in the original concept, study design and implementation.

Stuart J Falconer made revisions to the protocol and was involved in the original concept, study design and implementation.

Stavros Papachristos made revisions to the protocol and was involved in the original concept, study design and implementation.

Philip Nash made revisions to the protocol and was involved in the original concept, study design and implementation.

Vitaliy Androshchuk made revisions to the protocol and was involved in the original concept, study design and implementation.

Marc Clancy is senior author and made revisions to the protocol and was involved in the original concept, study design and implementation.

COMPETING INTERESTS

Declaration of Interest

The Transplant Audit Collaborative acknowledges financial support from Astellas Pharma UK Ltd in the facilitation of meetings, the provision of training opportunities for its members and in funding publication costs. Astellas Pharma UK Ltd has been given no editorial capacity in this paper, neither will they be given access to raw data, nor influence any aspect of the study which will be conducted transparently and without bias. Astellas Pharma UK Ltd will not be involved in the study design, interpretation of data or in the authorship of disseminations arising from the work being undertaken.

FUNDING

Funding for this publication will be provided by Astellas Pharma UK Ltd.

DATA SHARING STATEMENT

At the time of writing this protocol, data has not been collected and thus data sharing is not applicable at this time.

REFERENCES:

1. Hong JC, Kahan BD. Immunosuppressive agents in organ transplantation: past, present and future. *Semin Nephrol* 2000; 20: 108-25
2. Webster AC et al. Tacrolimus versus ciclosporin as primary immunosuppression for kidney transplant recipients: meta-analysis and meta-regression of randomised trial data. *BMJ* 2005; 331: 810
3. Ekberg H et al. Reduced Exposure to Calcineurin Inhibitors in renal Transplantation. *NEJM* 2007; 357; 25: 2562
4. Hesselink DA, et al. The pharmacogenetics of calcineurin inhibitors: one step closer toward individualized immunosuppression? *Pharmacogenomics* 2005;6:323–37.
5. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000712/WC500022234.pdf EMA/618604/2008. 10 December 2012.
6. Schiff J, et al. *Clin J Am Soc Nephrol* 2007;2:374–384.
7. Venkataramanan R, et al. Clinical pharmacokinetics of tacrolimus. *Clin Pharmacokinet* 1995;29:404–430.
8. Antignac M, et al. Population pharmacokinetics and bioavailability of tacrolimus in kidney transplant patients. *Br J Pharmacol* 2007;64:750–757.
9. de Jonge H, et al. New Insights Into the Pharmacokinetics and Pharmacodynamics of the Calcineurin Inhibitors and Mycophenolic Acid: Possible Consequences for Therapeutic Drug Monitoring in Solid Organ Transplantation. *Ther Drug Monit* 2009;31:416–435.
10. Scholten EM, et al. AUC-guided dosing of tacrolimus prevents progressive systemic overexposure in renal transplant recipients. *Kidney Int* 2005;67:2440–2447.
11. Dew MA et al. Rates and Risk Factors for Nonadherence to the Medical Regimen After Adult Solid Organ Transplantation. *Transplantation* 2007;83:858–873.
12. Pashaei N, et al. CYP3A5 Genotype Is Not Related to the Inpatient Variability of Tacrolimus Clearance. *Ther Drug Monit* 2011;33:369–371.
13. Staatz CE, et al. Effect of CYP3A and ABCB1 Single Nucleotide Polymorphisms on the Pharmacokinetics and Pharmacodynamics of Calcineurin Inhibitors: Part II. *Clin Pharmacokinet* 2010;49:141–175.

14. Elens L, et al. A New Functional CYP3A4 Intron 6 Polymorphism Significantly Affects Tacrolimus Pharmacokinetics in Kidney Transplant Recipients. *Clin Chem* 2011;57:1574–1583.

15. Waiser J, et al. Impact of the variability of cyclosporin A trough levels on long-term renal allograft function. *Nephrol Dial Transplant* 2002;17:1310–1317

16. Hsiao M, et al. Monitoring Nonadherence and Acute Rejection With Variation in Blood Immunosuppressant Levels in Pediatric Renal Transplantation. *Transplantation* 2011;92: 918–922.

17. Whalen HR, et al. *Transplantation* 2016; (Epub ahead of print).

18. Shuker N, et al. Intra-patient variability in tacrolimus exposure: Causes, consequences for clinical management. *Transplantation Reviews* 2015;29: 78–84.

19. Sapir-Pichhadze R et al. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney International* 2014; 85: 1404-1411

20. Borra CPL et al. High within-patient variability in the clearance of tacrolimus is a risk factor for poor long-term outcome after kidney transplantation. *Nephrol Dial Transplant* 2010; 25: 2757-2763

BMJ Open

The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Retrospective Cohort Study Protocol



Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-016144.R1
Article Type:	Protocol
Date Submitted by the Author:	18-Apr-2017
Complete List of Authors:	Goldsmith, Petra; Royal Liverpool and Broadgreen University Hospitals NHS Trust, Renal Transplant Surgery; University of Liverpool, Institute of Infection and Global Health Bottomley, Matthew; Oxford University Hospitals NHS Foundation Trust Okechukwu, Okidi; Central Manchester University Hospitals NHS Foundation Trust Ross, Victoria; NHS Greater Glasgow and Clyde Ghita, Ryan; NHS Greater Glasgow and Clyde Wandless, David; NHS Grampian Falconer, Stuart; NHS Lothian Papachristos, Stavros; Central Manchester University Hospitals NHS Foundation Trust Nash, Philip; King's College Hospital NHS Foundation Trust Androshchuk, Vitaliy; Oxford University Hospitals NHS Foundation Trust Clancy, Marc; NHS Greater Glasgow and Clyde; University of Glasgow
Primary Subject Heading:	Renal medicine
Secondary Subject Heading:	Surgery
Keywords:	Renal transplantation < NEPHROLOGY, TRANSPLANT SURGERY, TRANSPLANT MEDICINE

SCHOLARONE™
Manuscripts

The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Retrospective Cohort Study Protocol

Authors:

Petra M Goldsmith^{1,2}

Matthew J Bottomley³

Okidi Okechukwu⁴

Victoria C Ross⁵

Ryan Ghita⁵

David Wandless⁶

Stuart J Falconer⁷

Stavros Papachristos⁴

Philip Nash⁸

Vitaliy Androshchuk³

Marc Clancy^{7,9}

Corresponding Author:

Petra M Goldsmith

Royal Liverpool University Hospitals NHS Trust

Prescot Street

LIVERPOOL

L7 8XP

United Kingdom

petra.goldsmith@googlemail.com

07795122575

On behalf of the Transplant Audit Collaborative

1. Renal Transplant Unit, Royal Liverpool University Hospitals NHS Trust, Liverpool, UK
2. University of Liverpool
3. Department of Nephrology, Oxford University Hospitals NHS Foundation Trust, Oxford, UK
4. Department of Transplantation, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK
5. Department of Transplantation, NHS Greater Glasgow and Clyde, Glasgow, UK
6. Department of Nephrology, NHS Grampian, Aberdeen, UK
7. Department of Transplantation, NHS Lothian, Edinburgh, UK
8. Department of Nephrology, King's College Hospital NHS Foundation Trust, London, UK
9. University of Glasgow, Glasgow, UK

WORD COUNT 1,787

ABSTRACT

Introduction: High intra-patient variability (IPV) in tacrolimus trough levels has been shown to be associated with higher rates of renal transplant failure. There is no consensus on what level of IPV constitutes a risk of graft loss. The establishment of such a threshold could help to guide clinicians in identifying at risk patients to receive targeted interventions to improve IPV and thus outcomes.

Methods and Analysis: A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study. The aim is to recruit a minimum of 1,600 patients with follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss. The study also aims to determine any national or regional trends in IPV and any demographic associations.

Ethics and Dissemination: Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The study will be registered locally in each participating centre in line with local Research and Development protocols. It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ARTICLE SUMMARY

Strengths and Limitations of this Study

1. This is a multicentre collaborative study comprised of units across the UK which are both Transplant Centres and referring Nephrology Units
2. The sample size will be the largest investigating IPV to date
3. The study utilises a unified method across all participating centres, which will enable meaningful comparison between centres
4. The study investigates retrospective data and requires a lengthy follow up period which will lead to some exclusions
5. There will be local variations in the laboratory assay that cannot be corrected for in the IPV calculations

INTRODUCTION

The addition of calcineurin inhibitors (CNI) as maintenance immunosuppressants has improved renal transplant 1 year survival rates since the 1980s.[1] Tacrolimus emerged as a viable alternative to Ciclosporin in the 1990s.[1] In 2005, a meta-analysis was published on randomised trial data comparing Tacrolimus and Ciclosporin as primary immunosuppressants in renal transplant, observing a 44% reduction in death censored graft loss with Tacrolimus over Ciclosporin.[2] In 2007, the Symphony Study reported favourable graft survival and function, and reduced biopsy proven rejection with low dose Tacrolimus over low dose Ciclosporin, Sirolimus or standard dose Tacrolimus.[3]

CNIs have a narrow therapeutic index: too little exposure places a transplant recipient at increased risk of acute rejection and donor specific antibody formation. Too much exposure and a transplant recipient is placed at increased risk of malignancy, infection, nephrotoxicity and unacceptable side effects such as tremor.[4-7]

Trough levels are used as a proxy for oral bioavailability of CNIs and vary both between patients (inter-patient variability) and for an individual over time (intra-patient variability, IPV). Between individuals, age, gender, ethnicity, body mass index, genetic polymorphisms in CYP3A5 and CYP3A4, drug interactions, adherence, liver function and lifestyle choices account for the differences. Similarly, IPV is affected by adherence, gastrointestinal metabolism and motility, diarrhoea, food and drug interactions, synchronicity of dose administration and blood test and variability of the laboratory assay.[8-15]

An emerging body of evidence is being established indicating favourable graft function, survival and fewer rejection episodes up to 1 year post transplant for patients demonstrating low intra-patient variability.[16-18] Similarly, high IPV has been associated with poorer outcomes and graft survival.[19-20] Donor age and previous transplants appear to be risk factors for a high intra-patient variability.[18] However, little data exists on the long term impact of high IPV and studies have not yet been able to draw conclusions about risk thresholds of variability because of limitations in sample size.

Objectives:

- To establish important baseline data about national and regional trends in IPV
- To investigate demographic associations and other characteristics for patients in high and low variability groups
- To establish whether there exists a “danger” threshold for IPV, above which a patient is deemed at risk of graft loss or dysfunction, so they can then be targeted for intervention prior to organ damage or failure

METHODS AND ANALYSIS

A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. It is the first Collaborative of its kind, facilitating the development of this largest study examining IPV to date. The TAC is comprised of junior doctors with an interest in Nephrology and/or Transplantation. It is supported by Consultant Medics and Surgeons in these fields to undertake research and audit projects related to transplantation. Any UK NHS Trust

involved in the after care of renal transplant recipients is eligible to register for this study providing they have a transplant or nephrology doctor willing to enrol in the TAC.

Patient Selection

Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study (table 1). To be enrolled, patients are required to have follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss.

Table 1: Inclusion / Exclusion Criteria

(T1=6-12 months post-transplant; T2=most recent 12 months)

INCLUSION CRITERIA	EXCLUSION CRITERIA
Age ≥ 18 at time of transplant	Age ≤ 17 at time of transplant
A functioning graft* at 2 years	Failed graft before 2 years
Renal only transplant	Non-renal transplant
Short acting Tacrolimus preparation only during study period	Modified release preparation at any point during study period
Tacrolimus as primary immunosuppressant	Other primary immunosuppressant (e.g. ciclosporin, sirolimus, other non-steroid)
Alive minimum 2 years following transplant	Pregnancy at any point during study period
	Death prior to 2 years following transplant
	Non-standard transplant (ABO or HLA incompatible; requiring desensitisation)
	Patients with fewer than 4 tacrolimus trough levels for T1 and T2

Sample Size

A large sample size is needed to provide meaningful numbers to establish variability risk cut offs. It is estimated that a minimum of 8 UK based centres will participate in the study. If each centre, on average, supplies data for 200 patients, a minimum dataset of 1,600 patients will be achieved.

It is, however, recognised that there will be a significant variation between the numbers of eligible patients available to each centre (tertiary transplant units will naturally have access to larger numbers than referring District General Hospitals). The set number of 200, therefore, is only to be used as a guide and it is recognised that this may not be achievable for smaller, secondary nephrology units. Similarly, we welcome larger numbers from any centre able to do so.

Data Collection and Information Governance

The TAC has established a standard minimum dataset (see Table 2) which each site will use as the basis for data collection. A template database will be provided to a representative of the study team from each NHS Trust participating in the study. This database will be password protected and held on secure local Trust servers. All data should be entered onto this database – the use of hard copy proformas associated with this audit is discouraged to avoid breaching data protection policies.

Table 2: Minimum Dataset

(T1=6-12 months post-transplant; T2=most recent 12 months)

DATA SET

- Date of transplant
- Recipient and Donor Ages
- Gender
- Ethnicity
- Primary Renal Diagnosis
- Previous transplants
- Mismatch Grade (A, B, DR)
- Type of donor
- eGFR at the end of T1 and T2
- Creatinine at 12 months and at the end of T2
- All Tacrolimus trough levels during T1 and T2
- Urinary protein creatinine ratio at the of end T1 and T2
- Graft Failure
- Delayed Graft Function
- Tacrolimus dosing at month 12 and at the end of T2
- Other immunosuppression at both T1 and T2:
 - steroid Y/N;
 - MMF Y/N;
 - Azathioprine Y/N
- Induction agent
- Denovo Donor Specific Antibody status post-transplant and level (mean fluorescence intensity)

The data collection database will be anonymised and will contain no patient identifiable data. Each hospital will be issued with patient study identifiers for all patients included in the study. A separate password protected spreadsheet of the study identifiers with the corresponding hospital numbers should be securely stored locally by each participating centre for local reference only should there be any difficulties or queries regarding data collection.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The anonymised master database will be compiled by a dedicated, named member of the collaborative. It will be shared with all members of the Transplant Audit Collaborative using secure NHS email only (either local Trust email or NHS.net) and will be held on secure, password protected Trust servers only.

Data collection has been underway since March 2017 and is scheduled for completion in June 2017. A period of data analysis will then follow until September 2017 at which time the results will be disseminated as detailed in the Ethics and Dissemination section below.

Data Analysis

Studies have shown that high IPV is associated with poorer renal transplant outcomes¹⁷ but no research group has yet established or described an IPV level at which the risk of such outcomes is significantly increased. With our large sample size, we will be able to stratify our group into quartiles or quintiles based on IPV, enabling us to compare outcomes between the groups.

IPV will be calculated using the mean absolute deviation as described by Shuker et al (18). Individual subjects will be stratified into groups based on observed variability during T1. Intergroup comparisons will be made using both univariate and multivariate analyses for the clinically relevant end points including graft loss, graft dysfunction (assessed by eGFR and new onset proteinuria) and biopsy proven rejection episodes. The univariate predictive value of T1 IPV for these outcomes will be evaluated by receiver operator curve (ROC) assessment.

Categorical variables will be compared using Chi Square and Fischer's exact test where appropriate. Continuous variables will be assessed using T test for parametric and Mann Whitney U test for non-parametric data.

Multivariate analysis will use Cox regression survival analysis to compare event free survival, corrected for potential confounders including age, gender and ethnicity.

Permissions and Registration

Each participating centre will be expected to complete a site registration form. A named member of the TAC from each participating centre will be responsible for gaining all necessary local Trust permissions and study registrations as required by local Research and Development and Audit offices.

Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The findings of the study are not expected to impact upon individual patient care.

ETHICS AND DISSEMINATION

As this study does not fulfil the NHS Health Research Authority and Medical Research Council's criteria for research, formal ethical approval is not needed. However, all local NHS Trust approvals and registrations will be sought from each participating centre.

It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

CONTRIBUTORSHIP STATEMENT

All authors fulfil the ICMJE criteria for authorship and agree to be accountable for all aspects of the work. In brief:

Petra M Goldsmith wrote the protocol, produced revisions and was involved in the original concept, study design and implementation.

Matthew J Bottomley made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Okidi Okechukwu made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Victoria C Ross made revisions to the protocol and was involved in the original concept, study design and implementation.

Ryan Ghita made revisions to the protocol and was involved in the original concept, study design and implementation.

David Wandless made revisions to the protocol and was involved in the original concept, study design and implementation.

Stuart J Falconer made revisions to the protocol and was involved in the original concept, study design and implementation.

Stavros Papachristos made revisions to the protocol and was involved in the original concept, study design and implementation.

Philip Nash made revisions to the protocol and was involved in the original concept, study design and implementation.

Vitaliy Androshchuk made revisions to the protocol and was involved in the original concept, study design and implementation.

Marc Clancy is senior author and made revisions to the protocol and was involved in the original concept, study design and implementation.

COMPETING INTERESTS

Declaration of Interest

The Transplant Audit Collaborative acknowledges financial support from Astellas Pharma UK Ltd in the facilitation of meetings, the provision of training opportunities for its members and in funding publication costs. Astellas Pharma UK Ltd has been given no editorial capacity in this paper, neither will they be given access to raw data, nor influence any aspect of the study which will be conducted transparently and without bias. Astellas Pharma UK Ltd will not be involved in the study design, interpretation of data or in the authorship of disseminations arising from the work being undertaken.

FUNDING

Funding for this publication will be provided by Astellas Pharma UK Ltd.

DATA SHARING STATEMENT

At the time of writing this protocol, data has not been collected and thus data sharing is not applicable at this time.

REFERENCES:

1. Hong JC, Kahan BD. Immunosuppressive agents in organ transplantation: past, present and future. *Semin Nephrol* 2000; 20: 108-25

2. Webster AC et al. Tacrolimus versus ciclosporin as primary immunosuppression for kidney transplant recipients: meta-analysis and meta-regression of randomised trial data. *BMJ* 2005; 331: 810

3. Ekberg H et al. Reduced Exposure to Calcineurin Inhibitors in renal Transplantation. *NEJM* 2007; 357; 25: 2562

4. Hesselink DA, et al. The pharmacogenetics of calcineurin inhibitors: one step closer toward individualized immunosuppression? *Pharmacogenomics* 2005;6:323–37.

5. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000712/WC500022234.pdfEMA: EMA/618604/2008. 10 December 2012.

6. Schiff J, et al. *Clin J Am Soc Nephrol* 2007;2:374–384.

7. Venkataramanan R, et al. Clinical pharmacokinetics of tacrolimus. *Clin Pharmacokinet* 1995;29:404–430.

8. Antignac M, et al. Population pharmacokinetics and bioavailability of tacrolimus in kidney transplant patients. *Br J Pharmacol* 2007;64:750–757.

9. de Jonge H, et al. New Insights Into the Pharmacokinetics and Pharmacodynamics of the Calcineurin Inhibitors and Mycophenolic Acid: Possible Consequences for Therapeutic Drug Monitoring in Solid Organ Transplantation. *Ther Drug Monit* 2009;31:416–435.

10. Scholten EM, et al. AUC-guided dosing of tacrolimus prevents progressive systemic overexposure in renal transplant recipients. *Kidney Int* 2005;67:2440–2447.

11. Dew MA et al. Rates and Risk Factors for Nonadherence to the Medical Regimen After Adult Solid Organ Transplantation. *Transplantation* 2007;83:858–873.

12. Pashaei N, et al. CYP3A5 Genotype Is Not Related to the Inpatient Variability of Tacrolimus Clearance. *Ther Drug Monit* 2011;33:369–371.

13. Staatz CE, et al. Effect of CYP3A and ABCB1 Single Nucleotide Polymorphisms on the Pharmacokinetics and Pharmacodynamics of Calcineurin Inhibitors: Part II. *Clin Pharmacokinet* 2010;49:141–175.

14. Elens L, et al. A New Functional CYP3A4 Intron 6 Polymorphism Significantly Affects Tacrolimus Pharmacokinetics in Kidney Transplant Recipients. *Clin Chem* 2011;57:1574–1583.
15. Waiser J, et al. Impact of the variability of cyclosporin A trough levels on long-term renal allograft function. *Nephrol Dial Transplant* 2002;17:1310–1317
16. Hsiao M, et al. Monitoring Nonadherence and Acute Rejection With Variation in Blood Immunosuppressant Levels in Pediatric Renal Transplantation. *Transplantation* 2011;92: 918–922.
17. Whalen HR, et al. High Inpatient Tacrolimus Variability is Associated with Worse Outcomes in Renal Transplantation Using Low-Dose Tacrolimus Immunosuppressive Regime. *Transplantation* 2017; 101 (2): 430-436.
18. Shuker N, et al. Intra-patient variability in tacrolimus exposure: Causes, consequences for clinical management. *Transplantation Reviews* 2015;29: 78–84.
19. Sapir-Pichhadze R et al. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney International* 2014; 85: 1404-1411
20. Borra CPL et al. High within-patient variability in the clearance of tacrolimus is a risk factor for poor long-term outcome after kidney transplantation. *Nephrol Dial Transplant* 2010; 25: 2757-2763

BMJ Open

The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Retrospective Cohort Study Protocol



Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-016144.R2
Article Type:	Protocol
Date Submitted by the Author:	05-Jun-2017
Complete List of Authors:	Goldsmith, Petra; Royal Liverpool and Broadgreen University Hospitals NHS Trust, Renal Transplant Surgery; University of Liverpool, Institute of Infection and Global Health Bottomley, Matthew; Oxford University Hospitals NHS Foundation Trust Okechukwu, Okidi; Central Manchester University Hospitals NHS Foundation Trust Ross, Victoria; NHS Greater Glasgow and Clyde Ghita, Ryan; NHS Greater Glasgow and Clyde Wandless, David; NHS Grampian Falconer, Stuart; NHS Lothian Papachristos, Stavros; Central Manchester University Hospitals NHS Foundation Trust Nash, Philip; King's College Hospital NHS Foundation Trust Androshchuk, Vitaliy; Oxford University Hospitals NHS Foundation Trust Clancy, Marc; NHS Greater Glasgow and Clyde; University of Glasgow
Primary Subject Heading:	Renal medicine
Secondary Subject Heading:	Surgery
Keywords:	Renal transplantation < NEPHROLOGY, TRANSPLANT SURGERY, TRANSPLANT MEDICINE

SCHOLARONE™
Manuscripts

The Impact of Intra-Patient Variability (IPV) in Tacrolimus Trough Levels on Long-Term Renal Transplant Function: Multicentre Collaborative Retrospective Cohort Study Protocol

Authors:

Petra M Goldsmith^{1,2}

Matthew J Bottomley³

Okidi Okechukwu⁴

Victoria C Ross⁵

Ryan Ghita⁵

David Wandless⁶

Stuart J Falconer⁷

Stavros Papachristos⁴

Philip Nash⁸

Vitaliy Androshchuk³

Marc Clancy^{7,9}

Corresponding Author:

Petra M Goldsmith

Royal Liverpool University Hospitals NHS Trust

Prescot Street

LIVERPOOL

L7 8XP

United Kingdom

petra.goldsmith@googlemail.com

07795122575

On behalf of the Transplant Audit Collaborative

1. Renal Transplant Unit, Royal Liverpool University Hospitals NHS Trust, Liverpool, UK
2. University of Liverpool
3. Department of Nephrology, Oxford University Hospitals NHS Foundation Trust, Oxford, UK
4. Department of Transplantation, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK
5. Department of Transplantation, NHS Greater Glasgow and Clyde, Glasgow, UK
6. Department of Nephrology, NHS Grampian, Aberdeen, UK
7. Department of Transplantation, NHS Lothian, Edinburgh, UK
8. Department of Nephrology, King's College Hospital NHS Foundation Trust, London, UK
9. University of Glasgow, Glasgow, UK

WORD COUNT 2,217

ABSTRACT

Introduction: High intra-patient variability (IPV) in tacrolimus trough levels has been shown to be associated with higher rates of renal transplant failure. There is no consensus on what level of IPV constitutes a risk of graft loss. The establishment of such a threshold could help to guide clinicians in identifying at risk patients to receive targeted interventions to improve IPV and thus outcomes.

Methods and Analysis: A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study. The aim is to recruit a minimum of 1,600 patients with follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss. The study also aims to determine any national or regional trends in IPV and any demographic associations.

Ethics and Dissemination: Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The study will be registered locally in each participating centre in line with local Research and Development protocols. It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ARTICLE SUMMARY

Strengths and Limitations of this Study

1. This is a multicentre collaborative study comprised of units across the UK which are both Transplant Centres and referring Nephrology Units
2. The sample size will be the largest investigating IPV to date
3. The study utilises a unified method across all participating centres, which will enable meaningful comparison between centres
4. The study investigates retrospective data and requires a lengthy follow up period which will lead to some exclusions
5. There will be minor local variations in the laboratory assay that cannot be corrected for in the IPV calculations
6. Confounding factors such as geographical bias, repatriation as a cause of lost to follow up, under-representation of poorly compliant patients who do not attend appointments, frequency of sampling and temporary medications are beyond the scope of this study

INTRODUCTION

The addition of calcineurin inhibitors (CNI) as maintenance immunosuppressants has improved renal transplant 1 year survival rates since the 1980s.[1] Tacrolimus emerged as a viable alternative to Ciclosporin in the 1990s.[1] In 2005, a meta-analysis was published on randomised trial data comparing Tacrolimus and Ciclosporin as primary immunosuppressants in renal transplant, observing a 44% reduction in death censored graft loss with Tacrolimus over Ciclosporin.[2] In 2007, the Symphony Study reported favourable graft survival and function, and reduced biopsy proven rejection with low dose Tacrolimus over low dose Ciclosporin, Sirolimus or standard dose Tacrolimus.[3]

CNIs have a narrow therapeutic index: too little exposure places a transplant recipient at increased risk of acute rejection and donor specific antibody formation. Too much exposure and a transplant recipient is placed at increased risk of malignancy, infection, nephrotoxicity and unacceptable side effects such as tremor.[4-7]

Trough levels are used as a proxy for oral bioavailability of CNIs and vary both between patients (inter-patient variability) and for an individual over time (intra-patient variability, IPV). Between individuals, age, gender, ethnicity, body mass index, genetic polymorphisms in CYP3A5 and CYP3A4, drug interactions, adherence, liver function and lifestyle choices account for the differences. Similarly, IPV is affected by adherence, gastrointestinal metabolism and motility, diarrhoea, food and drug interactions, synchronicity of dose administration and blood test and variability of the laboratory assay.[8-15]

An emerging body of evidence is being established indicating favourable graft function, survival and fewer rejection episodes up to 1 year post transplant for patients demonstrating low intra-patient variability.[16-18] Similarly, high IPV has been associated with poorer outcomes and graft survival.[19-20] Donor age and previous transplants appear to be risk factors for a high intra-patient variability.[18] However, little data exists on the long term impact of high IPV and studies have not yet been able to draw conclusions about risk thresholds of variability because of limitations in sample size.

Objectives:

- To establish important baseline data about national and regional trends in IPV
- To investigate demographic associations and other characteristics for patients in high and low variability groups
- To establish whether there exists a “danger” threshold for IPV, above which a patient is deemed at risk of graft loss or dysfunction, so they can then be targeted for intervention prior to organ damage or failure

Outcomes:

Primary Outcomes:	Recent 12 months’ IPV	IPV months 6-12	Change in IPV
Secondary Outcomes:	Ethnicity	Recipient Age	Change in IPV
	Previous transplants	DR mismatch	Graft function
	Graft survival	Gender	

Confounders:

We acknowledge the potential for confounding factors that are outside the scope of this study to address. These include frequency of tacrolimus level sampling (and the reasons why this might be increased) and conversely under-representation of poorly compliant patients who do not attend appointments. These confounders may be affected by hospital admissions, temporary medication use (such as oral antibiotics) and those patients with a modified Tacrolimus target.

METHODS AND ANALYSIS

A multicentre Transplant Audit Collaborative (TAC) has been established to conduct a retrospective study examining tacrolimus IPV and renal transplant outcomes. It is the first Collaborative of its kind, facilitating the development of this largest study examining IPV to date. The TAC is comprised of junior doctors with an interest in Nephrology and/or Transplantation. It is supported by Consultant physicians and surgeons in these fields to undertake research and audit projects related to transplantation. Any UK NHS Trust involved in the after care of renal transplant recipients is eligible to register for this study providing they have a transplant or nephrology doctor willing to enrol in the TAC.

Patient Selection

Patients in receipt of a renal transplant at participating centres between 2009 and 2014 and fulfilling the inclusion criteria will be included in the study (table 1). To be enrolled, patients are required to have follow-up spanning at least 2 years in order to determine a threshold IPV above which a renal transplant recipient would be considered at increased risk of graft loss.

Table 1: Inclusion / Exclusion Criteria

(T1=6-12 months post-transplant; T2=most recent 12 months)

INCLUSION CRITERIA	EXCLUSION CRITERIA
Age ≥18 at time of transplant	Age ≤17 at time of transplant
A functioning graft* at 2 years	Failed graft before 2 years
Renal only transplant	Non-renal transplant
Short acting Tacrolimus preparation only during study period	Modified release preparation at any point during study period
Tacrolimus as primary immunosuppressant	Other primary immunosuppressant (e.g. ciclosporin, sirolimus, other non-steroid)
Alive minimum 2 years following transplant	Pregnancy at any point during study period
	Death prior to 2 years following transplant
	Non-standard transplant (ABO or HLA incompatible; requiring desensitisation)
	Patients with fewer than 4 tacrolimus trough levels for T1 and T2

Sample Size

A large sample size is needed to provide meaningful numbers to establish variability risk cut offs. It is estimated that a minimum of 8 UK based centres will participate in the study. If each centre, on average, supplies data for 200 patients, a minimum dataset of 1,600 patients will be achieved.

It is, however, recognised that there will be a significant variation between the numbers of eligible patients available to each centre (tertiary transplant units will naturally have access to larger numbers than referring District General Hospitals). The set number of 200, therefore, is only to be used as a guide and it is recognised that this may not be achievable for smaller, secondary nephrology units. Similarly, we welcome larger numbers from any centre able to do so.

Data Collection and Information Governance

The TAC has established a standard minimum dataset (see Table 2) which each site will use as the basis for data collection. A template database will be provided to a representative of the study team from each NHS Trust participating in the study. This database will be password protected and held on secure local Trust servers. All data should be entered onto this database – the use of hard copy proformas associated with this audit is discouraged to avoid breaching data protection policies.

Table 2: Minimum Dataset

(T1=6-12 months post-transplant; T2=most recent 12 months)

DATA SET
<ul style="list-style-type: none">• Date of transplant• Recipient and Donor Ages• Gender• Ethnicity• Primary Renal Diagnosis• Previous transplants• Mismatch Grade (A, B, DR)• Type of donor• eGFR at the end of T1 and T2• Creatinine at 12 months and at the end of T2• All Tacrolimus trough levels during T1 and T2• Urinary protein creatinine ratio at the of end T1 and T2• Graft Failure• Delayed Graft Function• Tacrolimus dosing at month 12 and at the end of T2• Other immunosuppression at both T1 and T2:<ul style="list-style-type: none">○ steroid Y/N;○ MMF Y/N;○ Azathioprine Y/N• Induction agent• Denovo Donor Specific Antibody status post-transplant and level (mean fluorescence intensity)

The data collection database will be anonymised and will contain no patient identifiable data. Each hospital will be issued with patient study identifiers for all patients included in the study. A separate password protected spreadsheet of the study identifiers with the corresponding hospital numbers should be securely stored locally by each participating centre for local reference only should there be any difficulties or queries regarding data collection.

The anonymised master database will be compiled by a dedicated, named member of the collaborative. It will be shared with all members of the Transplant Audit Collaborative using secure NHS email only (either local Trust email or NHS.net) and will be held on secure, password protected Trust servers only.

Data collection has been underway since March 2017 and is scheduled for completion in September 2017. A period of data analysis will then follow until December 2017 at which time the results will be disseminated as detailed in the Ethics and Dissemination section below.

Data Analysis

Studies have shown that high IPV is associated with poorer renal transplant outcomes¹⁷ but no research group has yet established or described an IPV level at which the risk of such outcomes is significantly increased. With our large sample size, we will be able to stratify our group into quartiles or quintiles based on IPV, enabling us to compare outcomes between the groups.

IPV will be calculated using the mean absolute deviation as described by Shuker et al (18). Individual subjects will be stratified into groups based on observed variability during T1. Intergroup comparisons will be made using both univariate and multivariate analyses for the clinically relevant end points including graft loss, graft dysfunction (assessed by eGFR and new onset proteinuria) and biopsy proven rejection episodes. The univariate predictive value of T1 IPV for these outcomes will be evaluated by receiver operator curve (ROC) assessment.

Categorical variables will be compared using Chi Square and Fischer’s exact test where appropriate. Continuous variables will be assessed using T test for parametric and Mann Whitney U test for non-parametric data.

Multivariate analysis will use Cox regression survival analysis to compare event free survival, corrected for potential confounders including age, gender and ethnicity. Where there is loss to follow up after the two year period, data will be censored according to last known status at the time of last creatinine or tacrolimus level (whichever is the latter).

Permissions and Registration

Each participating centre will be expected to complete a site registration form. A named member of the TAC from each participating centre will be responsible for gaining all necessary local Trust permissions and study registrations as required by local Research and Development and Audit offices.

Consent will not be sought from patients whose data are utilised in this study as no additional procedures or information will be required from participants beyond that which would normally take place as part of clinical care. The findings of the study are not expected to impact upon individual patient care.

ETHICS AND DISSEMINATION

As this study does not fulfil the NHS Health Research Authority and Medical Research Council’s criteria for research, formal ethical approval is not needed. However, all local NHS Trust approvals and registrations will be sought from each participating centre.

It is anticipated that the results of this audit will be disseminated locally, in participating NHS Trusts, through national and international meetings and publications in peer reviewed journals.

CONTRIBUTORSHIP STATEMENT

All authors fulfil the ICMJE criteria for authorship and agree to be accountable for all aspects of the work. In brief:

Petra M Goldsmith wrote the protocol, produced revisions and was involved in the original concept, study design and implementation.

Matthew J Bottomley made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Okidi Okechukwu made substantial revisions to the protocol and was involved in the original concept, study design and implementation.

Victoria C Ross made revisions to the protocol and was involved in the original concept, study design and implementation.

Ryan Ghita made revisions to the protocol and was involved in the original concept, study design and implementation.

David Wandless made revisions to the protocol and was involved in the original concept, study design and implementation.

Stuart J Falconer made revisions to the protocol and was involved in the original concept, study design and implementation.

Stavros Papachristos made revisions to the protocol and was involved in the original concept, study design and implementation.

Philip Nash made revisions to the protocol and was involved in the original concept, study design and implementation.

Vitaliy Androshchuk made revisions to the protocol and was involved in the original concept, study design and implementation.

Marc Clancy is senior author and made revisions to the protocol and was involved in the original concept, study design and implementation.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

COMPETING INTERESTS

Declaration of Interest

The Transplant Audit Collaborative acknowledges financial support from Astellas Pharma UK Ltd in the facilitation of meetings, the provision of training opportunities for its members and in funding publication costs. Astellas Pharma UK Ltd has been given no editorial capacity in this paper, neither will they be given access to raw data, nor influence any aspect of the study which will be conducted transparently and without bias. Astellas Pharma UK Ltd will not be involved in the study design, interpretation of data or in the authorship of disseminations arising from the work being undertaken.

FUNDING

Funding for this publication will be provided by Astellas Pharma UK Ltd.

DATA SHARING STATEMENT

At the time of writing this protocol, data has not been collected and thus data sharing is not applicable at this time.

REFERENCES:

1. Hong JC, Kahan BD. Immunosuppressive agents in organ transplantation: past, present and future. *Semin Nephrol* 2000; 20: 108-25
2. Webster AC et al. Tacrolimus versus ciclosporin as primary immunosuppression for kidney transplant recipients: meta-analysis and meta-regression of randomised trial data. *BMJ* 2005; 331: 810
3. Ekberg H et al. Reduced Exposure to Calcineurin Inhibitors in renal Transplantation. *NEJM* 2007; 357; 25: 2562
4. Hesselink DA, et al. The pharmacogenetics of calcineurin inhibitors: one step closer toward individualized immunosuppression? *Pharmacogenomics* 2005;6:323–37.
5. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000712/WC500022234.pdf EMA/618604/2008. 10 December 2012.
6. Schiff J, et al. *Clin J Am Soc Nephrol* 2007;2:374–384.
7. Venkataramanan R, et al. Clinical pharmacokinetics of tacrolimus. *Clin Pharmacokinet* 1995;29:404–430.
8. Antignac M, et al. Population pharmacokinetics and bioavailability of tacrolimus in kidney transplant patients. *Br J Pharmacol* 2007;64:750–757.
9. de Jonge H, et al. New Insights Into the Pharmacokinetics and Pharmacodynamics of the Calcineurin Inhibitors and Mycophenolic Acid: Possible Consequences for Therapeutic Drug Monitoring in Solid Organ Transplantation. *Ther Drug Monit* 2009;31:416–435.
10. Scholten EM, et al. AUC-guided dosing of tacrolimus prevents progressive systemic overexposure in renal transplant recipients. *Kidney Int* 2005;67:2440–2447.
11. Dew MA et al. Rates and Risk Factors for Nonadherence to the Medical Regimen After Adult Solid Organ Transplantation. *Transplantation* 2007;83:858–873.
12. Pashaei N, et al. CYP3A5 Genotype Is Not Related to the Inpatient Variability of Tacrolimus Clearance. *Ther Drug Monit* 2011;33:369–371.
13. Staatz CE, et al. Effect of CYP3A and ABCB1 Single Nucleotide Polymorphisms on the Pharmacokinetics and Pharmacodynamics of Calcineurin Inhibitors: Part II. *Clin Pharmacokinet* 2010;49:141–175.

14. Elens L, et al. A New Functional CYP3A4 Intron 6 Polymorphism Significantly Affects Tacrolimus Pharmacokinetics in Kidney Transplant Recipients. *Clin Chem* 2011;57:1574–1583.

15. Waiser J, et al. Impact of the variability of cyclosporin A trough levels on long-term renal allograft function. *Nephrol Dial Transplant* 2002;17:1310–1317

16. Hsiau M, et al. Monitoring Nonadherence and Acute Rejection With Variation in Blood Immunosuppressant Levels in Pediatric Renal Transplantation. *Transplantation* 2011;92: 918–922.

17. Whalen HR, et al. High Inpatient Tacrolimus Variability is Associated with Worse Outcomes in Renal Transplantation Using Low-Dose Tacrolimus Immunosuppressive Regime. *Transplantation* 2017; 101 (2): 430-436.

18. Shuker N, et al. Intra-patient variability in tacrolimus exposure: Causes, consequences for clinical management. *Transplantation Reviews* 2015;29: 78–84.

19. Sapir-Pichhadze R et al. Time-dependent variability in tacrolimus trough blood levels is a risk factor for late kidney transplant failure. *Kidney International* 2014; 85: 1404-1411

20. Borra CPL et al. High within-patient variability in the clearance of tacrolimus is a risk factor for poor long-term outcome after kidney transplantation. *Nephrol Dial Transplant* 2010; 25: 2757-2763