

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Factors Associated with Healthcare Workers Willingness to Participate in Disasters in Sana'a, Yemen

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-030547
Article Type:	Original research
Date Submitted by the Author:	05-Apr-2019
Complete List of Authors:	Al-Hunaishi, Weiam; University of Malaya, Department of Social and Preventive Medicine Hoe, Victor; Universiti Malaya Chinna, Karuthan ; Universiti Malaya
Keywords:	Disaster preparedness, willingness, self-efficacy, interpersonal, healthcare worker

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Factors associated with Healthcare Workers Willingness to Participate in Disasters in**
6
7 **Sana'a, Yemen**
8
9

10 **Authors' names, credentials/degrees, email addresses and institutional affiliations:**
11

- 12
13 1. **MPH. MD. Weiam Al-Hunaishi:** Department of Social & Preventive Medicine, Faculty of
14 Medicine, University of Malaya, Kuala Lumpur, Malaysia. (e-mail: waam59@hotmail.com)
15
16
17 2. **Prof. Dr. Victor CW Hoe:** Centre for Occupational and Environmental Health, Department
18 of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur,
19 Malaysia. (e-mail: victor@ummc.edu.my)
20
21
22 3. **Associate Prof. Dr. Karuthan Chinna:** Julius Centre University of Malaya, Department of
23 Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur,
24 Malaysia. (e-mail: karuthan@um.edu.my)
25
26
27
28
29
30

31 **Correspondence and reprint requests:** to Weiam Al-Hunaishi (e-mail: Waam59@hotmail.com,
32 telephone number: +60173479802, mail address: Department of Social and Preventive Medicine
33 Faculty of Medicine, University of Malaya 50603 Kuala Lumpur, Malaysia).
34
35
36
37

38 **List of Abbreviations:**
39

40		
41	CI	Confidence Interval
42		
43	HCW	Healthcare workers
44		
45	IQR	Interquartile range
46		
47	SD	Standard Deviation
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

Abstract

Objectives: Willingness to participate in disasters has been indicated as a principal issue when overlooked. Hence, it requires addressing during disaster preparedness training courses to insure health service coverage. The study aims to assess healthcare workers (HCW) willingness to participate in biological and natural disasters, and its associated factors.

Design: This is a cross-sectional study, in which a self-administered questionnaire was distributed to 1093 HCW. The data was analysed using multiple logistic regression with significance level $p < 0.05$. Ethical clearance and consent of the participants were duly obtained.

Setting: In three public Hospitals that provide tertiary level healthcare, in Sana'a City, Yemen.

Participants: Nurses and doctors (HCW).

Results: There were 692 (response rate 63.3%) completed the questionnaire, of which 55.1% were nurses and 44.9% were doctors. The study found that self-efficacy was associated with willingness to participate in disaster response for any type of disasters (Odds Ratio [OR]=1.328, 95% Confidence Interval [95%CI]: 1.206 to 1.464), natural disasters (OR=1.138, 95%CI: 1.064 to 1.217), and influenza pandemic (OR=1.112, 95%CI: 1.049 to 1.180). The results further show that willingness is associated with HCW being young, male and having higher educational qualifications.

Conclusion: Self-efficacy has been found to be an important factor associated with willingness. Improving self-efficacy through training in disaster preparedness may increase willingness of HCW to participate in disaster.

Key words: Disaster preparedness; willingness; self-efficacy; interpersonal; healthcare worker.

Strength and limitations of this study

- This study is the first study that attempt to assess HCW willingness to participate in different types of disasters in Sana'a City, Yemen.
- Willingness levels were found to be different in different type of disasters; the lowest willingness rate was in influenza pandemic, which considered concerning in the fragile state of Yemen and the increased number of outbreaks in the country.
- The study was survey based, using a convenience sample, however, it represented HCW in Sana'a City in term of gender and living city.
- The study data was a conversion of Likert data to binary data, so there might be some degree of bias.
- Beside the limitations that result from employing a cross-sectional design in the study, it was conducted in a city with considerable political unrest, therefore, a subsequent study is recommended.

Introduction:

In various parts of the world, disasters destroyed communities and infrastructures, causing huge material and human losses.¹ Similarly, hospitals and health centres are also affected by the disasters, which restrict its work of relieving the disaster-stricken community. Disasters is defined as “a situation or event which overwhelms local capacity, necessitating a request to a national or international level for external assistance; an unforeseen and often sudden event that causes great damage, destruction and human suffering”.² By 2025, more than half of world population will live in cities, particularly in urban cities located along a seismic fault lines, flood and other natural disasters prone areas.³

According to the Global Facility for Disaster Reduction and Recovery (GFDRR), Yemen is one of the priority countries in the Middle East and North Africa region, mainly due to its vulnerability to disasters.⁴⁻⁶ Natural disasters are recurrent in Yemen, which includes storms, landslides, earthquake and floods., Floods due to the monsoonal rainfall, is responsible for the majority of the mortality due to disaster, this is followed by storms and landslides as repeated disasters.⁶ Currently in Yemen disaster management only mainly focuses on responding to post disasters damage and there is a lack of disaster preparedness such training and mock drills. The insufficient training makes it difficult to maintain preparedness.^{4,7,8}

During disasters, HCW are expected to provide health care assistance to people suffering from the disasters alongside with caring for their usual patients. Some are also required to care for their dependents. Most of the previous studies reported an anticipated decrease in health workforce during disaster as not all HCW are willing to participate in a disaster. Systematic reviews have found that willingness was associated with factors like the nature and type of event; competing obligations between personal and professional needs; the

1
2
3 work environment and climate including personal safety; and the relationship between
4
5 knowledge and perceptions of efficacy.^{9,10}
6
7

8
9
10 Disaster preparedness activities and routine disasters trainings has been found to
11 improve knowledge, skills and attitude preparedness of disaster.¹¹ However, the factors that
12 affect willingness to participate in disasters are beyond just having knowledge on disasters
13 management.⁹⁻¹²
14
15
16

17
18 In a recent systematic review, they found that only few researches have been
19 conducted in the Middle East on willingness to participate in disasters. They recommend that
20 further research should be conducted based on behavioural theories to better understand the
21 Middle East context of willingness to participate in a disaster.¹³ In another systematic review
22 of 70 studies on HCW willingness to participate in disasters only 12 studies were form Asia
23 and none of those studies was from the Middle East.¹² Therefore, the objective of this study is
24 to determine the associations between socio-demographic, professional, and intrapersonal
25 factors associated with doctors' and nurses' willingness to participate in disasters in response
26 in Sana'a, Yemen.
27
28
29
30
31
32
33
34
35
36
37
38

39 **Methods:**

40 **Study design, population and instrument**

41
42 This is a cross-sectional study conducted in three public hospitals in the Sana'a
43 Governorate, Yemen; Al-Thora, Al-Jumhuri, and Al-Kuwait hospitals, which have a more
44 than total of 4000 nurses and doctors which were included in this study. Out of this 1093
45 HCW were selected through convenient sampling by approaching the healthcare providers
46 individually. The sample size of 1093 was determined based the "perception of responsibility
47 to participate", which is an element of the self-efficacy construct. The determined sample size
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 was 614, which was inflated by 78 percent based on the response rate of the previous
4
5 studies.¹⁴
6
7

8 The questionnaire was developed by the researcher based on information from
9
10 previous studies and opinion of national and international experts.¹⁴⁻¹⁷ The questionnaire was
11
12 pilot tested on doctors and nurses from a different hospital; additional feedback from local
13
14 experts was sort, to ensure adequate coverage and understanding of key topics items on the
15
16 questionnaire. The questionnaire consists of socio-demographic, professional, and
17
18 intrapersonal factors, and willingness to participate in a disaster. The professional,
19
20 intrapersonal and willingness questions was measured on a five-point Likert scale (1=
21
22 strongly disagree, 2= disagree, 3=somewhat agree, 4= agree, and 5= strongly agree). A binary
23
24 variable was created based on “less or equal two” as low and “more than 2” as high.
25
26
27
28

29 There were four items in the self-efficacy construct. The descriptive statistics and
30
31 inter-item correlation values of the self-efficacy are presented in Table 1.
32
33

34
35 [\[Table 1\]](#)
36
37

38 Based on Table 1, there was moderate level of agreement in all the four items. The
39
40 highest correlation for each item with at least one other item in the construct was between 0.3
41
42 and 0.9. in factor analysis, the Kaiser-Meyer-Olkin (KMO) value was 0.658 ($p < 0.001$). A
43
44 single factor was formed from the four items computed and was saved as incentive to be used
45
46 in further analysis. Accordingly, to calculate the self- efficacy, four questions was asked. The
47
48 sum outcome of the four answers ranged from 4 to 20 (Median 16, Interquartile range [IQR]
49
50 5).
51
52
53

54
55 Self-efficacy theory was chosen to build the construct as it illustrates beliefs that
56
57 drives actions to face and solve problems faced to achieve goals. In case of disaster, the
58
59 theory could be applied to as HCW coping with fear and threat and adapting new behaviours
60

1
2
3 because of their beliefs in their competences. This belief is derived from their successful
4 performances, observing colleagues and managers positive behaviour, convincing by a
5 superior person, and calming the physiological and emotional pressure caused by the threat.¹⁸
6
7
8
9
10
11
12

13 **Data collection**

14
15
16 Ethical approval was obtained from the Ministry of Health and Population of Yemen.
17 Informed consent was obtained from the respondent and confidentiality of personal
18 disclosures was re-assured. The self-administered questionnaires were distributed and
19 retrieved between February and March 2018. A final 767 questionnaires were returned for
20 analyses; 75 questionnaires were omitted due to missing values.
21
22
23
24
25
26
27
28
29
30

31 **Statistical analysis**

32
33 The data was analysed using Statistical Package for Social Science (SPSS) version 22.
34 Descriptive outputs were generated describing the median, interquartile range, frequency
35 counts and percentages. Chi square test and multiple logistic regression was performed to test
36 the hypothesis of the study on $p < 0.05$ for all statistical tests. The variables selected to be
37 included in the multivariate analysis were variables with $p < 0.25$ in the univariate analysis,
38 and gender, age, and type of profession were also included.
39
40
41
42
43
44
45
46
47

48 **Patient and Public Involvement**

49
50
51 Patients and the general public were not involved in the design of this research. HCW
52 were involved during the pilot study in order to test the understanding of the written
53 questionnaire. In addition, the data was access by limited number of the research team.
54
55
56
57
58
59
60

Results:

Socio-demographics Characteristics

Responses are collected from 692 HCW (response rate 63.3%), where 311 (44.9%) are doctors and 381 (55.1%) nurses. Most of the female participants are nurses 64.5%. More than half of the doctors (56.6%) have more than five years of experience at their current place of work, as compare to only 40.9 per cent of the nurses. The average age of participants is 31.96 (Standard Deviation [SD] 7.46) years across the two occupations. Out of the 65.9% who have dependents, i.e., child or elderly persons, 73.9% reported to have support to care for their dependent in a case of disaster. The percentages are almost equally distributed among doctors and nurses.

Professional and Intrapersonal Characteristics

Table 2 describes the socio-demographic, professional, interpersonal, and willingness characteristics of respondents. Only 39.2% of the HCW had have any previous training in disaster. This is the same for previous work experience, where only 35.8% of the participants had previous work experience in a disaster situation.

The doctors and nurses report a high trust in work safety, in family, in colleague's preparedness to react, and in hospital preparedness to react in case of a disaster, with trust in colleague's preparedness as the highest percentage (88.3%). Similarly, HCW have a high median score (16; IQR 2) in self-efficacy construct.

Respondents willingness

Ninety percent of the participants express high willingness to participate in any type of disasters. However, they were less willing to participate in natural disasters (77.3%) and influenza pandemic (66.0%) (Table2).

[\[Table 2\]](#)

Factors associated with willingness to participate in any type disaster

There was an association between participants' gender and willingness to participate in any type of disaster, with male being more willing compared to female (crude OR=2.161, 95%CI: 1.307 to 3.573) (Table 3). Those with high trust in work safety (crude OR=3.284, 95%CI: 1.937 to 5.567), trust in colleague's preparedness (crude OR=2.592, 95%CI: 1.401 to 4.795) and self-efficacy (crude OR=1.358; 95%CI: 1.247 to 1.479) were also found to be more willing to participate in any type of disaster in the univariate analysis (Table 3).

In the final model, having trust in work safety (adjusted OR=2.535, 95%CI: 1.357 to 4.736) and self-efficacy (adjusted OR=1.319, 95%CI: 1.197 to 1.453) were found to be associated with general willingness with any type of disasters (Table 4).

Factors associated with willingness to participate in natural disaster

In the univariate analysis, there was an association between some of the socio-demographic characteristics (age, gender, education level, type of profession and work duration) with willingness to participate in natural disasters (Table 3). Participants with bachelor and postgraduate degrees has a higher odd of willingness to participate in natural disaster compared to those with diploma education. Those in the age group of

1
2
3 between 31 and 45 were found to be more willing when compared to participants that are 30
4 years old and younger (Table 3). The main interpersonal factors associated with willingness
5 in natural disasters were participants' trust in colleagues' preparedness in case of a natural
6 disaster, and self-efficacy (Table 3).
7
8
9
10
11

12
13 In multivariate, being male (adjusted OR=1.639, 95%CI: 1.102 to 2.439) and self-
14 efficacy (adjusted OR=1.143, 95%CI: 1.069 to 1.221) were significantly associated with
15 willingness to participate in natural disasters (Table 4).
16
17
18
19
20
21
22

23 **Factors associated with willingness to participate in influenza pandemic disaster**

24
25 For willingness to participate in influenza pandemic, the univariate results reveal that
26 being male, having a dependent with no support (compared to participants without
27 dependent), having previous experience and self-efficacy were associated with willingness to
28 participate in pandemic (Table 3).
29
30
31
32
33
34

35 In the final model, having previous experience (adjusted OR=1.528, 95%CI: 1.058 to
36 2.207) and self-efficacy (adjusted OR=1.114, 95%CI: 1.050 to 1.182) were found to be
37 associated with willingness to participate in influenza pandemic (Table 4).
38
39
40
41
42

43 [\[Table 3, Table 4\]](#)
44
45
46
47
48

49 **Discussion:**

50
51 Having one hundred percent participation of HCW is difficult to obtain as much as it
52 is vital in case of disasters. Previous studies suggested that between 65 and 97 percent of
53 HCW were willing to participate in a natural disaster, and between 54 and 86 percent in an
54 influenza pandemic.^{11,15,19,20} The reason in this difference in the levels of willingness between
55
56
57
58
59
60

1
2
3 the two types of disasters is due to the great distinction in their nature. The outcome of the
4 interaction between HCW and the socioenvironmental determinants of these disasters leads to
5 having different willingness levels. According to Conner et al, the weighted risk resulting
6 from this interaction plays a major role in HCW willingness. In case of pandemics, the fear of
7 the inability to control biohazards and watching colleagues inquiring a communicable disease
8 after contact with affected persons was a suggested reason for the low willingness levels.¹⁰
9
10 Another qualitative study in Australia highlighted that in different types of disasters
11 emergency nurse's willingness to attend to work is shaped by the weighted risk to self and
12 surrounding people and the pressure formed from the period dealing with the disaster.¹²
13
14 Natural disasters may not directly affect HCW or their families like influenzas pandemic
15 which may be a reason for the higher level of willingness to attend to work compared to
16 pandemics.
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 The findings of this study are consistent with the previous studies regarding their
32 willingness level to different types of disasters. More HCW were willing to participate in
33 natural disaster (77.3%) as compared to influenza pandemic (66.0%). Even though, there
34 were a higher percentage of HCW willingness from this study to participate in both natural
35 disaster and influenza pandemic were higher compared to other studies, Yemen's high
36 vulnerability, due its topography and current economic, political and health status, could
37 affect the country health status and cause huge adverse health impacts in case of disasters.
38
39 According to the Vulnerability Matrix, one third of Yemen's districts are highly vulnerable.⁵
40
41
42
43
44
45
46
47
48
49

50 In addition, due to the armed conflicts in Yemen that started in early 2015, it had
51 become difficult to deal with any disaster in case it occurred. According to the World Health
52 Organization (WHO), the healthcare system in Yemen is in a critical situation with about
53 50% of the health facilities in Yemen are either partially or total damage as a result of the
54 natural disasters and conflicts. Furthermore, of the 3507 healthcare facilities in Yemen,
55
56
57
58
59
60

1
2
3 almost 300 healthcare facilities have been destroyed.^{5,21} Therefore, any possible decrease in
4 the health work force during disasters in Yemen must be put in considerations during
5 response to disasters. The existed drained health sector is based on a vulnerable health system
6 in term of its structure and healthcare staff number and distribution in the country.²² Thus,
7 training and preparing healthcare staff to act during disasters is critical and could help
8 achieving Priority 4 in the United Nations' Sendai Framework for Disaster Risk Reduction
9 2015- 2030 in enhancing disaster preparedness for effective response and to "Build Back
10 Better" in recovery, rehabilitation and reconstruction of disasters, and prevent any possible
11 humanitarian catastrophe.²³

24 The results show that willingness to participate in events of natural disasters may be
25 encouraged by many suggested factors. Factors such as respondents being male are
26 accompanied with higher willingness to participate in natural disasters. which is similar to
27 previous studies conducted in Jordan, China and United nations of America.^{14,15,24,25}
28 However, having less female health workers willing to participate in disasters highlight a
29 need to address this group during health preparedness programs. This lack in female HCW
30 willingness could be due to the gender and cultural barriers in Yemen. These barriers need to
31 be addressed to better motivate females' workers participation in case of disasters. This is
32 important area for research as studies on willingness moving forward.

33 As in previous studies, nurses and doctors with previous experience in influenza
34 pandemic are more willing to participate in influenza pandemic.^{24,26} HCW which has
35 experience with a previous disaster have better knowledge in disaster management, coping
36 strategies, and ways to protect themselves and their families; which may be a reason for
37 having higher odds of willingness. According to Alzahrani et al., Most of the nurses have a
38 higher level in Mass gatherings management such as communicating effectively during
39 emergencies in Saudi Arabia due to Previous experience, he also found a need for further
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 trainings by the nurses.²⁷ Chokahi et al., found that simulation training, attending conferences
4 and previous experience of increase paediatric surgeons feeling of preparedness.²⁶
5
6
7

8 This study found that trust in work safety during any type of disaster play key role in
9 willingness to participate, which is similar to results from previous studies.^{11,24,28} Work safety
10 is important in order for HCP to feel safe and do their work, a study by Stergachis et al, found
11 that majority of the participants reported their fear or concern for self in case of influenza
12 pandemic and during earthquake scenarios as one of the major barriers to willingness.²⁹
13 Considering work safety in disaster preparedness is especially important in Yemen after
14 notifications made by Aldahrai et al. and Naser et al. on the current low work place safety
15 and the needs of increasing safety and security standards in hospitals in Yemen Health
16 establishments, regardless to external or internal catastrophes; unsafe work place could lead
17 to increase optional absenteeism during disaster.^{7,8} Thus, putting information on work safety
18 and protection devices provided during several types of disasters is advisable.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 Many studies have established the association between self-efficacy and willingness
35 to participate in disasters in previous.^{10,14,15} Similarly, this study have identified that self-
36 efficacy plays an important role in willingness to participate in natural disasters, in influenza
37 pandemic and any type of disasters. It indicates that elements tested for self-efficacy like
38 participants increase familiarity with their role, responsibility to react, and confidence of
39 ability to deal with various types of disasters could be a key factor to increase HCP self-
40 efficacy, thus, increase their willingness. Disaster preparedness trainings are encouraged to
41 contain materials that explain response to different type of disasters, doctors' and nurses' role
42 in disasters and how it makes a different in response to disasters. Supportive measures to
43 increase HCW self-efficacy such as immediate communication with needed information and
44 rewarding for efforts in disasters are suggested.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strength and limitation

Regardless of the limitation of this study, it had substantial strengths. It is the first study explored the patterns of the willingness status in the Yemen and the factors associated with it. Also, study was distributed in person to the HCP to ensure obtaining higher response rate. In addition, it was using self-administered questionnaire to minimise information bias such as interviewer bias.

This study had limitations, as the study was performed under a political insecurity/fragile state of the country a list of workers of HCP working at the hospitals was difficult to obtain. As a result, a universal sample was undertaken which could limit the study representativeness. Other than the self-administered questionnaire limitations, the study findings are limited to staff working at the tertiary level public hospitals in urban areas of Yemen.

Conclusion:

Increased the likelihood of willingness to participate in disaster play a key role in guarantee optimum number of work force. This study indicates that one's socio-demographic, professional and intrapersonal factors plays a role in increase his/her willingness in general and across different type of disasters. Significant differences were revealed between participants' willingness in natural disasters by gender and self-efficacy, and participant's willingness in influenza pandemics by previous experience and self-efficacy. This result suggests integrating disaster management into an earlier education levels of doctors' and nurses' educational establishments. Other areas for preparedness may include increase hospitals safety and resilience; Hence that willingness to participate in natural disasters' response was influence by demographic characteristic of the healthcare personnel, others

1
2
3 found outside the bracket should be motivated to participate with a reward package such as
4
5 incentives and hazard allowance. Further studies should be conducted in both urban and rural
6
7 settings with relatively peaceful atmosphere.
8
9

10
11
12
13 **Acknowledgement:** We thank Dr. Susan Connor, Dr. Karen Hammad and
14
15
16 Dr..Mohammed Khalil for their Support and advice during the study development, all
17
18 University Malaya academic staff, the team who collected the data, and all the participants
19
20 who contributed in this study.
21
22

23
24 **Competing interests:** The authors have no financial disclosures to declare and no
25
26 conflicts of interest to report.
27
28

29
30 **Funding:** This research did not receive any specific grant from funding agencies in the
31
32 public, commercial, or not-for-profit sectors.
33
34

35
36 **Data sharing statement:** No additional data are available.
37
38

39 40 **Reference:**

- 41
42
43 1. United Nations Economic and Social Commission for Asia and the Pacific
44 (UNESCAP). Disasters in Asia and the Pacific: 2015 year in review. United Nations
45 report Economic and social commission for Asia and the Pacific 2015.
46
47
- 48
49 2. Below R, Wirtz A, Guha-Sapir D. Disaster category classification and peril
50 terminology for operational purposes. 2009.
51
52
- 53
54 3. United Nations Office for Disaster Risk Reduction (UNISDR). (2012). Making
55 cities resilient report 2012. Available from:
56 http://www.unisdr.org/files/28240_rcreport.pdf.
57
- 58
59 4. Environment MoWa. Yemen: National progress report on the implementation of the
60 Hyogo Framework for Action (2013-2015). 2015.

- 1
2
3 5. World Health Organization. WHO annual report 2017: Yemen. Cairo: WHO
4 Regional Office for the Eastern Mediterranean; 2018. Licence: CC BYNC-SA 3.0
5 IGO.
6
7
- 8
9 6. The International Bank for Reconstruction and Development / The World Bank.
10 Natural Disasters in the Middle East and North Africa: A Regional Overview. 2014.
11
- 12
13 7. Aladhrai SA, Djalali A, Della Corte F, Alsabri M, El-Bakri NK, Ingrassia PL.
14 Impact of the 2011 Revolution on Hospital Disaster Preparedness in Yemen.
15 Disaster Medicine and Public Health Preparedness. 2015;9(4):396-402.
16
- 17
18 8. Naser WN, Ingrassia PL, Aladhrae S, Abdulraheem WA. A Study of Hospital
19 Disaster Preparedness in South Yemen. Prehospital and Disaster Medicine.
20 2018;33(2):133.
21
- 22
23 9. Chaffee M. Willingness of health care personnel to work in a disaster: an integrative
24 review of the literature. Disaster Medicine and Public Health Preparedness.
25 2009;3(1):42-56.
26
- 27
28 10. Connor SB. When and why health care personnel respond to a disaster: the state of
29 the science. Prehospital and Disaster Medicine. 2014;29(3):270-4.
30
- 31
32 11. Gowing JR, Walker KN, Elmer SL, Cummings EA. Disaster Preparedness among
33 Health Professionals and Support Staff: What is Effective? An Integrative Literature
34 Review. Prehospital and Disaster Medicine. 2017;32(3):321-8.
35
- 36
37 12. Arbon P, Cusack L, Ranse J, Shaban RZ, Considine J, Kako M, et al. Exploring
38 staff willingness to attend work during a disaster: a study of nurses employed in
39 four Australian emergency departments. Australasian Emergency Nursing Journal :
40 AENJ. 2013;16(3):103-9.
41
- 42
43 13. Ejeta LT, Ardalan A, Paton D. Application of Behavioral Theories to Disaster and
44 Emergency Health Preparedness: A Systematic Review. PLOS Currents Disasters.
45 2015;7. Doi: 10.1371/currents.dis.31a8995ced321301466db400f1357829.
46
- 47
48 14. Burke RV, Goodhue CJ, Chokshi NK, Upperman JS. Factors associated with
49 willingness to respond to a disaster: a study of healthcare workers in a tertiary
50 setting. Prehospital and Disaster Medicine. 2011;26(4):244-50.
51
52
53
54
55
56
57
58
59
60

15. Charney RL, Rebmann T, Flood RG. Hospital Employee Willingness to Work during Earthquakes Versus Pandemics. *Journal of Emergency Medicine*. 2015;49(5):665-74. Doi: <https://doi.org/10.1016/j.jemermed.2015.07.030>.
16. Balicer RD, Omer SB, Barnett DJ, Everly GS, Jr. Local public health workers' perceptions toward responding to an influenza pandemic. *BMC Public Health*. 2006;6:99.
17. Shapira S, Aharonson-Daniel L, Bar-Dayyan Y, Sykes D, Adini B. Knowledge, perceptions, attitudes and willingness to report to work in an earthquake: A pilot study comparing Canadian versus Israeli hospital nursing staff. *International Emergency Nursing*. 2016;25:7-12.
18. Bandura A. *Self-efficacy: The exercise of control*: Macmillan; 1997.
19. Arbon P, Ranse J, Cusack L, Considine J, Shaban RZ, Woodman RJ, et al. Australasian emergency nurses' willingness to attend work in a disaster: a survey. *Australasian Emergency Nursing Journal : AENJ*. 2013;16(2):52-7.
20. Brice JH, Gregg D, Sawyer D, Cyr JM. Survey of Hospital Employees' Personal Preparedness and Willingness to Work Following a Disaster. *Southern Medical Journal*. 2017;110(8):516-22.
21. United Nations Office for the Coordination of Humanitarian Affairs (UNOCHA). *Humanitarian Bulletin Yemen/ Health system on the verge of collapse*2015. Available from: <http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-30-november-2015-enar>.
22. Qirbi N, Ismail SA. Health system functionality in a low-income country in the midst of conflict: the case of Yemen. *Health Policy and Planning*. 2017;32(6):911-22.
23. United Nations - Headquarters. *Sendai Framework for Disaster Risk Reduction 2015-2030*. 2015. Available from: <https://www.unisdr.org/we/inform/publications/43291>.
24. Loke AY, Fung WM, Liu X. Mainland China nurses' willingness to report to work in a disaster. *American Journal of Disaster Medicine*. 2013;8(4):273-82. Doi: 10.5055/ajdm.2013.0134.

- 1
2
3 25. Al Khalaileh MA, Bond E, Alasad JA. Jordanian nurses' perceptions of their
4 preparedness for disaster management. *International emergency nursing*.
5 2012;20(1):14-23.
6
7
8
9 26. Chokshi NK, Behar S, Nager AL, Dorey F, Upperman JS. Disaster management
10 among pediatric surgeons: preparedness, training and involvement. *American*
11 *Journal of Disaster Medicine*. 2008;3(1):5-14.
12
13
14 27. Alzahrani F, Yiannis K. Emergency nurse disaster preparedness during mass
15 gatherings: a cross-sectional survey of emergency nurses' perceptions in hospitals in
16 Mecca, Saudi Arabia. *BMJ Open*. 2017;7(4):e013563.
17
18
19 28. Ogedegbe C, Nyirenda T, Delmoro G, Yamin E, Feldman J. Health care workers
20 and disaster preparedness: barriers to and facilitators of willingness to respond.
21 *International Journal of Emergency Medicine*. 2012;5(1):29.
22
23
24 29. Stergachis A, Garberson L, Lien O, D'Ambrosio L, Sangare L, Dold C. Health care
25 workers' ability and willingness to report to work during public health emergencies.
26 *Disaster Medicine and Public Health Preparedness*. 2011;5(4):300-8.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Descriptive statistics and inter-term correlation for items in self-efficacy

Items	Descriptive statistics		Inter-item correlation			
	Mean	SD	item 1	item 2	item 3	item 4
Item 1: Ability to perform work	3.36	1.074	1.000	0.592	0.267	0.259
Item 2: Familiarity with their role	3.62	1.085	0.592	1.000	0.325	0.324
Item 3: Responsibility to participate	4.23	0.929	0.267	0.325	1.000	0.390
Item 4: Ability report to work	4.03	1.062	0.259	0.324	0.390	1.000

For peer review only

Table 2: Socio-demographic, professional, interpersonal, and willingness Characteristics of Respondents

Variable		Frequency (n= 692)	Percentage (%)	
Age (years)	≤30	360	52.0	
	31-45	294	42.5	
	≥46	38	5.5	
Gender	Male	419	60.5	
	Female	273	39.5	
Marital-status	Single	258	37.3	
	Married	420	60.7	
	Divorce	8	1.2	
	Widow	6	0.9	
Education-level	Diploma	280	40.5	
	Bachelor	275	39.7	
	Post graduate education	Master	104	15.0
		Professional	33	4.8
		Doctors	311	44.9
Profession type	Specialist medical practitioner	126	8.2	
	General medical practitioner	185	26.7	
	Nurses	381	55.1	
Work duration (years)	≤5	360	52	
	6-10	213	30.8	
	11-15	76	11.0	
	≥16	43	6.2	
Dependent	With dependent	Elder	65	9.4
		Child	276	39.9
		Both	115	16.6
		No dependent	236	34.1
	^a Support	No	119	26.1
Yes		337	73.9	
^b Previous training	With previous training	271	39.2	
	Without previous training	421	60.8	
^b Previous experience	With previous experience	248	35.8	
	Without previous training	444	64.2	
Trust in work safety in case of disaster	High trust in work safety	563	81.4	
	Low trust in work safety	129	18.6	
Trust in family preparedness in case of a disaster	High trust in family preparedness	544	78.6	
	Low trust in family preparedness	148	21.4	
Trust in colleague preparedness to react in disaster	High trust in colleague preparedness	611	88.3	
	Low trust in colleague preparedness	81	11.7	

Trust in hospital preparedness to react in disaster	High trust in hospital preparedness	522	75.4
	Low trust in hospital preparedness	170	24.6
Self-efficacy score (Median; Inter-Quartile)		16	5
Willingness to participate in Any type of disaster	High	623	90.0
	Low	69	10.0
Willingness to participate in Natural disaster	High	535	77.3
	Low	157	22.7
Willingness to participate in Influenza and Pandemic	High	457	66.0
	Low	235	34.0

^aOnly for participants with dependants. The percentages are only within participants with dependents.

^bPrevious experience and training in disasters

Table 3: Univariate association of crude socio-demographic, professional and interpersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters	Willingness to participate in natural disasters	Willingness to participate in influenza pandemic
	Odds ratio (95% C.I.)	Odds ratio (95% C.I.)	Odds ratio (95% C.I.)
Age			
≤30	1	1	1
31-45	1.056 (0.628-1.775)	1.639 (1.126-2.387)	1.138 (0.822-1.576)
≥46	0.733 (0.269-1.997)	1.967 (0.798-4.850)	1.354 (0.650-2.820)
Gender			
Female	1	1	1
Male	2.161 (1.307-3.573)	2.254 (1.571-3.234)	1.505 (1.093-2.072)
Marital status			
Single	1	1	1
Married	0.899 (0.532-1.521)	1.128 (0.781-1.629)	1.090 (0.787-1.510)
Divorce and widow	0.615 (0.130-2.913)	1.160 (0.314-4.291)	1.362 (0.416-4.466)
Education Level			
Up to Diploma	1	1	1
Up to Bachelor	1.380 (0.775-2.458)	1.962 (1.317-2.922)	1.221 (0.858-1.738)
Up to Postgraduate	0.847 (0.450-1.596)	2.279 (1.351-3.842)	0.998 (0.651-1.528)
Profession type			
Nurses	1	1	1
General medical practitioner	1.473 (0.781-2.776)	1.554 (1.013-2.384)	0.756 (0.524-1.091)
Specialist medical practitioner	0.965 (0.507-1.835)	2.407 (1.377-4.208)	0.942 (0.614-1.446)
Work duration			
≤5	1	1	1
6-10	1.006 (0.563-1.798)	1.079 (0.721-1.614)	0.975 (0.683-1.392)
11-15	1.028 (0.438-2.415)	2.337 (1.118-4.882)	1.059 (0.626-1.791)
≥16	0.456 (0.196-1.063)	0.650 (0.329-1.286)	1.198 (0.603-2.379)
Presence of dependent			
No dependent	1	1	1
Dependent with no support	0.673 (0.346-1.308)	1.149 (0.683-1.933)	1.617 (1.000-2.612)
Dependent with support	1.253 (0.707-2.222)	1.305 (0.880-1.935)	1.283 (0.907-1.815)
^aPrevious training			
No	1	1	1
Yes	1.653 (0.959-2.849)	1.090 (0.756-1.573)	1.147 (0.829-1.585)
^aPrevious experience			
No	1	1	1
Yes	1.216 (0.714-2.070)	0.974 (0.673-1.411)	1.857 (1.317-2.619)
Trust in work safety			

Low	1	1	1
High	3.284 (1.937-5.567)	1.280 (0.825-1.987)	1.192 (0.800-1.774)
Trust in family preparedness			
Low	1	1	1
High	1.338 (0.756-2.369)	1.292 (0.850-1.963)	0.990 (.0674-1.454)
Trust in colleague's preparedness			
Low	1	1	1
High	2.592 (1.401-4.795)	1.974 (1.200-3.246)	1.392 (0.866-2.237)
Trust in hospital preparedness			
Low	1	1	1
High	1.392 (0.807-2.400)	1.212 (0.810-1.814)	1.158 (0.807-1.663)
^bSelf-efficacy	1.358 (1.247-1.479)	1.184 (1.115-1.257)	1.135 (1.076-1.197)

^aPrevious experience and training in disasters

^b*Continuous measure, with one unit increase in self-efficacy*

Table 4: Multivariate association of adjusted professional and interpersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters	Willingness to participate in natural disasters	Willingness to participate in influenza pandemic
	Odds ratio (95% C.I.)	Odds ratio (95% C.I.)	Odds ratio (95% C.I.)
Age			
≤30	1	1	1
31-45	1.137 (0.609-2.121)	1.432 (0.928-2.209)	1.008 (0.695-1.461)
≥46	0.440 (0.136-1.418)	1.087 (0.409-2.886)	1.062 (0.484-2.332)
Gender			
Female	1	1	1
Male	1.456 (0.807-2.628)	1.639 (1.102-2.439)	1.131 (0.793-1.612)
Education Level			
Up to Diploma	-	1	-
Up to Bachelor	-	1.706 (0.993-2.932)	-
Up to Postgraduate	-	1.177 (0.522-2.657)	-
Profession type			
Nurses	1	1	1
General medical practitioner	1.285 (0.438-3.767)	1.581 (0.691-3.614)	0.751 (0.468-1.205)
Specialist medical practitioner	1.392 (0.578-3.352)	0.939 (0.515-1.713)	0.697 (0.473-1.026)
Presence of dependent			
No dependent	-	-	1
Dependent with no support	-	-	1.537 (0.910-2.598)
Dependent with support	-	-	1.154 (0.791-1.685)
^aPrevious experience			
No	-	-	1
Yes	-	-	1.528 (1.058-2.207)
Trust in work safety			
Low	1	-	-
High	2.535 (1.357-4.736)	-	-
Trust in colleague's preparedness			
Low	1	1	-
High	1.199 (0.576-2.496)	1.363 (0.791-2.351)	-
^bSelf-efficacy			
	1.319 (1.197-1.453)	1.143 (1.069-1.221)	1.114 (1.050-1.182)

^aPrevious experience and training in disasters

^bContinuous measure, with one unit increase in self-efficacy

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	Main document (1)
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	Main document (1)
Background /	#2	Explain the scientific background and rationale for the	Main document

1	rationale		investigation being reported	(3)
2				
3				
4	Objectives	#3	State specific objectives, including any prespecified	Main document
5			hypotheses	(4)
6				
7				
8				
9	Study design	#4	Present key elements of study design early in the paper	Main document
10				(4)
11				
12				
13				
14	Setting	#5	Describe the setting, locations, and relevant dates,	Main document
15			including periods of recruitment, exposure, follow-up,	(4-6)
16			and data collection	
17				
18				
19				
20				
21				
22	Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods	Main document
23			of selection of participants.	(4)
24				
25				
26				
27		#7	Clearly define all outcomes, exposures, predictors,	Main document
28			potential confounders, and effect modifiers. Give	(5), table1
29			diagnostic criteria, if applicable	(Page 18)
30				
31				
32				
33				
34				
35	Data sources /	#8	For each variable of interest give sources of data and	Main document
36	measurement		details of methods of assessment (measurement).	(5)
37			Describe comparability of assessment methods if there	
38			is more than one group. Give information separately for	
39			exposed and unexposed groups if applicable.	
40				
41				
42				
43				
44				
45				
46				
47	Bias	#9	Describe any efforts to address potential sources of bias	Main document
48				(13)
49				
50				
51				
52	Study size	#10	Explain how the study size was arrived at	Main document
53				(4,5)
54				
55				
56				
57				
58	Quantitative	#11	Explain how quantitative variables were handled in the	Main document
59				
60				

1	variables	analyses. If applicable, describe which groupings were	(6)
2			
3		chosen, and why	
4			
5			
6	Statistical	#12a Describe all statistical methods, including those used to	Main document
7			
8	methods	control for confounding	(6)
9			
10			
11		#12b Describe any methods used to examine subgroups and	Main document
12			
13		interactions	(6)
14			
15			
16		#12c Explain how missing data were addressed	Main document
17			
18			(6)
19			
20			
21			
22		#12d If applicable, describe analytical methods taking account	Main document
23			
24		of sampling strategy	(4, 6)
25			
26			
27		#12e Describe any sensitivity analyses	
28			
29			
30	Participants	#13a Report numbers of individuals at each stage of study—	Main document
31			
32		eg numbers potentially eligible, examined for eligibility,	(4, 6)
33			
34		confirmed eligible, included in the study, completing	
35			
36		follow-up, and analysed. Give information separately for	
37			
38		for exposed and unexposed groups if applicable.	
39			
40			
41			
42		#13b Give reasons for non-participation at each stage	Main document
43			
44			
45			(4, 6, 13)
46			
47			
48		#13c Consider use of a flow diagram	
49			
50			
51	Descriptive data	#14a Give characteristics of study participants (eg	Main document
52			
53		demographic, clinical, social) and information on	(7,8), table 2
54			
55		exposures and potential confounders. Give information	(page 19, 20)
56			
57		separately for exposed and unexposed groups if	
58			
59			
60			

applicable.

1			
2			
3			
4		#14b	Indicate number of participants with missing data for
5			each variable of interest
6			Table 2 (page
7			19, 20)
8			
9	Outcome data	#15	Report numbers of outcome events or summary
10			Measures. Give information separately for exposed and
11			unexposed groups if applicable.
12			Main document
13			(7), table 2
14			(page 19, 20)
15			
16	Main results	#16a	Give unadjusted estimates and, if applicable,
17			confounder-adjusted estimates and their precision (eg,
18			95% confidence interval). Make clear which confounders
19			were adjusted for and why they were included
20			Main document
21			(8, 9), table 3,
22			4 (page 21-23)
23			
24			
25			
26		#16b	Report category boundaries when continuous variables
27			were categorized
28			Main document
29			5
30			
31			
32		#16c	If relevant, consider translating estimates of relative risk
33			into absolute risk for a meaningful time period
34			
35			
36			
37	Other analyses	#17	Report other analyses done—e.g., analyses of
38			subgroups and interactions, and sensitivity analyses
39			
40			
41			
42	Key results	#18	Summarise key results with reference to study objectives
43			Main document
44			(6-9)
45			
46			
47			
48	Limitations	#19	Discuss limitations of the study, taking into account
49			sources of potential bias or imprecision. Discuss both
50			direction and magnitude of any potential bias.
51			Main document
52			(12, 13)
53			
54			
55	Interpretation	#20	Give a cautious overall interpretation considering
56			objectives, limitations, multiplicity of analyses, results
57			Main document
58			(9-12)
59			
60			

1 from similar studies, and other relevant evidence.

2
3
4 Generalisability [#21](#) Discuss the generalisability (external validity) of the Main document
5 study results (13)
6
7

8
9 Funding [#22](#) Give the source of funding and the role of the funders for Main document
10 the present study and, if applicable, for the original study (14)
11
12 on which the present article is based
13
14
15

16
17 The STROBE checklist is distributed under the terms of the Creative Commons Attribution License
18 CC-BY. This checklist can be completed online using <https://www.goodreports.org/>, a tool made by
19 the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

Factors associated with Healthcare Workers Willingness to Participate in Disasters: a cross-sectional study in Sana'a, Yemen

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-030547.R1
Article Type:	Original research
Date Submitted by the Author:	02-Aug-2019
Complete List of Authors:	Al-Hunaishi, Weiam; University of Malaya, Department of Social and Preventive Medicine Hoe, Victor; Universiti Malaya Chinna, Karuthan ; Taylor's University, School of Medicine-SRI
Primary Subject Heading:	Public health
Secondary Subject Heading:	Global health, Health services research, Public health
Keywords:	Disaster preparedness, willingness, self-efficacy, healthcare worker, intrapersonal

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Factors associated with Healthcare Workers Willingness to Participate in Disasters: a**
6
7 **cross-sectional study in Sana'a, Yemen**
8
9

10 **Authors' names, credentials/degrees, email addresses and institutional affiliations:**
11

- 12
13 1. **MPH. MD. Weiam Al-Hunaishi (WA):** Department of Social & Preventive Medicine,
14 Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. (e-mail:
15 waam59@hotmail.com)
16
17
18
19 2. **Prof. Dr. Victor CW Hoe (VCWH):** Centre for Occupational and Environmental Health,
20 Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya,
21 Kuala Lumpur, Malaysia. (e-mail: victor@ummc.edu.my)
22
23
24 3. **Associate Prof. Dr. Karuthan Chinna (KC):** School of Medicine-SRI, Faculty of Health &
25 Medical Sciences, Taylor's University, Malaysia. (e-mail: karuthan.chinna@taylors.edu.my)
26
27
28
29
30

31 **Correspondence and reprint requests:** to Weiam Al-Hunaishi (e-mail: Waam59@hotmail.com,
32 telephone number: +60173479802, mail address: Department of Social and Preventive Medicine
33 Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia).
34
35
36
37

38 **Contributorship statement:**
39

- 40
41 • WA – conceptualized the study, conducted the data collection, analysis, and interpretation of
42 data for the work; draft and approve the manuscript for publication.
43
44
45 • VCWH – conceptualized the study, interpretation of data for the work; revise and approve the
46 manuscript for publication.
47
48
49 • KC – conceptualized the study, supervised the data analysis, interpretation of data for the
50 work.
51
52
53
54
55
56
57
58
59
60

1
2
3 **List of Abbreviations:**
4
5

6	CI	Confidence Interval
7		
8	HCW	Healthcare workers
9		
10	IQR	Interquartile range
11		
12	SD	Standard Deviation
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

For peer review only

Abstract

Objectives: Willingness to participate in disasters are usually overlooked and not addressed in disaster preparedness training courses to ensure health service coverage. This will lead to issue during the disaster. This study aims to assess healthcare workers (HCW) willingness to participate in biological and natural disasters, and to identify its associated factors.

Design: This is a cross-sectional study using a self-administered questionnaire. The questionnaire was distributed to 1093 HCW. The data was analysed using multiple logistic regression with significance level $p < 0.05$. Ethical clearance and consent of the participants were duly obtained.

Setting: In three public Hospitals that provide tertiary level healthcare in Sana'a City, Yemen.

Participants: Nurses and doctors (HCW).

Results: There were 692 (response rate 63.3%) completed the questionnaire, of which 55.1% were nurses and 44.9% were doctors. The study found that self-efficacy was associated with willingness to participate in disaster response for any type of disasters (Odds Ratio [OR]=1.328, 95% Confidence Interval [95%CI]: 1.206 to 1.464), natural disasters (OR=1.138, 95%CI: 1.064 to 1.217), and influenza pandemic (OR=1.112, 95%CI: 1.049 to 1.180). The results further show that willingness is associated with HCW being young, male and having higher educational qualifications.

Conclusion: Self-efficacy has been found to be an important factor associated with willingness. Improving self-efficacy through training in disaster preparedness may increase willingness of HCW to participate in a disaster.

Key words: Disaster preparedness; willingness; self-efficacy; intrapersonal; healthcare worker.

Strength and limitations of this study

- This study is the first study that attempt to assess HCW willingness to participate in different types of disasters in Sana'a City, Yemen.
- Implementing a cross-sectional design and using a convenience sample has made it difficult to establish causal association.
- The study samples represents HCW in Sana'a City in term of gender and locality.
- The face to face approach, which included introduction of the study and its objectives, ensured a higher response rate and minimised the risk of selection bias.

Introduction:

In various parts of the world, disasters destroy communities and infrastructures, causing huge material and human losses.¹ Similarly, hospitals and health centres are also affected by the disasters, which restrict its work of relieving disaster-stricken communities. By 2025, more than half of world population will live in cities, particularly in urban cities located along a seismic fault lines, flood and other natural disasters prone areas.² Therefore, healthcare worker (HCW) should be prepared to manage the influx of patient during possibly life threatening circumstances. Healthcare worker is defined as “the one who delivers care and services to the sick and ailing either directly as doctors and nurses or indirectly as aides, helpers, laboratory technicians, or even medical waste handlers”.³

According to the Global Facility for Disaster Reduction and Recovery (GFDRR), Yemen is one of the priority countries in the Middle East and North Africa region, mainly due to its vulnerability to disasters.⁴⁻⁶ Natural disasters are recurrent in Yemen, which includes storms, landslides, earthquake and floods. Floods due to the monsoonal rainfall, is responsible for the majority of the mortality due to disaster.⁶ Currently in Yemen, disaster management only mainly focuses on responding to post disasters damage, and there is a lack of disaster preparedness, such training and mock drills. The insufficient training makes it difficult to maintain preparedness.^{4,7,8} A competent prepared healthcare worker could better mitigate and respond to the community health needs during crises, which in turn will elevate health outcomes.

During disasters, HCW are expected to provide health care assistance to people suffering from the disasters alongside with caring for their usual patients. Some are also required to care for their dependents. Most of the previous studies reported an anticipated decrease in health workforce during disaster as not all HCW are willing to participate in a

1
2
3 disaster. Therefore, understanding the factors associated with willingness to participate in a
4
5 disaster would allow more effective planning for disaster situation. Systematic reviews have
6
7 found that willingness was associated with factors like the nature and type of event;
8
9 competing obligations between personal and professional needs; the work environment and
10
11 climate including personal safety, and the relationship between knowledge and perceptions of
12
13 efficacy.^{9,10}
14
15

16
17 Disaster preparedness activities and routine disasters trainings has been found to
18
19 improve knowledge, skills and attitude preparedness of disaster.¹¹ However, the factors that
20
21 affect willingness to participate in disasters are beyond just having knowledge on disasters
22
23 management.⁹⁻¹²
24
25

26
27 In a recent systematic review, they found only few studies that have been conducted
28
29 in the Middle East on willingness to participate in disasters. They recommend that further
30
31 research should be conducted based on behavioural theories to better understand the Middle
32
33 East context of willingness to participate in a disaster.¹³ In another systematic review of 70
34
35 studies on HCW willingness to participate in disasters, only 12 studies were form Asia and
36
37 none of those studies was from the Middle East.¹¹ Therefore, the objective of this study is to
38
39 determine the associations between socio-demographic, professional, and intrapersonal
40
41 factors associated with doctors' and nurses' willingness to participate in disasters in response
42
43 in Sana'a, Yemen. The study factors was built based on the self-efficacy behavioural theory.
44
45
46
47

48 **Methods:**

49 **Study design, population and instrument**

50
51 This is a cross-sectional study conducted in three public hospitals in the Sana'a
52
53 Governorate, Yemen; Al-Thora, Al-Jumhuri, and Al-Kuwait hospitals, which have more
54
55 than a total of 4,000 nurses and doctors, which was the focus of this study. Out of this, 1093
56
57
58
59
60

1
2
3 HCW were selected through convenient sampling by approaching the healthcare workers
4
5 individually. The sample size calculation was based on a 95% confidence interval with a
6
7 power of set to 80%. The sample size of 1093 was determined based on the variable
8
9 “perception of responsibility to participate” from a pre-study, which is an element of the self-
10
11 efficacy construct with the highest sample size to achieve an association.¹⁴ The determined
12
13 sample size was 614, which was inflated by 78 percent based on the lowest response rate of
14
15 previous studies to reach 1093.¹⁵
16
17
18
19

20 The questionnaire was developed by the researcher based on information from
21
22 previous studies and opinion of national and international experts (supplementary materials).¹⁵⁻
23
24 ¹⁸ The questionnaire was pre tested for validity and reliability. The questionnaire was
25
26 distributed to national and international expert together with a questions on the validity of the
27
28 questionnaire with a scale of 1 to 4 (1 not relevant to 4 highly relevant). It tested the
29
30 questionnaire consistency, relatedness, representativeness and clarity of wording. As a result,
31
32 six items of the questionnaire, which was related to the knowledge construct, was deleted and
33
34 changes on the questions was reworded. These questions have undergone forward and
35
36 backward-translation form English to Arabic then from Arabic to English. Then the
37
38 researcher used a face to face meeting with three HCW, where the study objective was
39
40 explained, then the participants were asked to give their comments on what they think of the
41
42 questions.
43
44
45
46
47

48 The questionnaire was pilot tested on 20 doctors and nurses from a hospital, other
49
50 than hospitals that have been chosen for the study. The internal consistency was assessed
51
52 using Cronbach’s alpha. Self- efficacy was 0.801, which considered good inter-correlation.
53
54 Therefore, the self-efficacy four items were used in the final survey to build the construct as
55
56 one of the intrapersonal factors. The pilot study have not published yet. The questionnaire
57
58 consists of socio-demographic, professional, and intrapersonal factors, and willingness to
59
60

1
2
3 participate in a disaster. The professional, intrapersonal and willingness questions was
4 measured on a five-point Likert scale (1=strongly disagree, 2=disagree, 3=somewhat agree,
5 4=agree, and 5=strongly agree). A binary variable was created based on “less or equal two”
6 as low and” more than 2” as high.
7
8
9
10
11
12

13 **Self-efficacy theory**

14
15 Self-efficacy theory was chosen to build the self-efficacy construct as it illustrates
16 beliefs that drive actions to face and solve problems faced to achieve intended goals. In case
17 of disaster, the theory could be applied to HCW who are coping with fear and threat and
18 adapting new behaviours because of their beliefs in their competences. This belief is derived
19 from their successful performances, observing colleagues and managers positive behaviour,
20 convincing by a superior person, and calming the physiological and emotional pressure
21 caused by the threat.¹⁹
22
23
24
25
26
27
28
29
30
31
32
33
34

35 **Data collection**

36
37 Ethical approval was obtained from the Ministry of Health and Population of Yemen.
38 Informed consent was obtained from the respondent and confidentiality of personal
39 disclosures was re-assured. The self-administered questionnaires were distributed and
40 retrieved between February and March 2018. A final 767 questionnaires were returned for
41 analyses; 75 questionnaires were omitted due to missing values.
42
43
44
45
46
47
48
49
50
51

52 **Statistical analysis**

53
54 The data was analysed using Statistical Package for Social Science (SPSS) version 22.
55 Descriptive outputs were generated describing the median, interquartile range, frequency
56 counts and percentages. Chi square test and multiple logistic regression was performed to test
57
58
59
60

1
2
3 the hypothesis of the study on $p < 0.05$ for all statistical tests. The variables selected to be
4 included in the multivariate analysis were variables with $p < 0.25$ in the univariate analysis,
5 and gender, age, and type of profession were also included.
6
7
8
9
10
11
12

13 **Patient and Public Involvement**

14
15
16 Patients and the general public were not involved in the conduct of this research.
17
18 HCW were involved during the pilot study in order to test the understanding of the written
19 questionnaire. In addition, the data was assessable by the researchers and is stored in a secure
20 file in the computer and online storage.
21
22
23
24
25
26
27
28

29 **Results:**

30 **Socio-demographics Characteristics**

31
32
33 Responses were collected from 692 HCW (response rate 63.3%), where 311 (44.9%)
34 were doctors and 381 (55.1%) were nurses. Most of the female participants were nurses
35 64.5%. More doctors (56.6%) had >5 years of experience at their current place of work, as
36 compare to nurses (40.9%). The average age of participants was 31.96 (Standard Deviation
37 [SD] 7.46) years across the two occupations. Out of the 65.9% who had dependents, i.e.,
38 having to take-care of child or elderly persons, 73.9% reported to have support to care for
39 their dependent in a case of disaster. The percentages were almost equally distributed
40 between doctors and nurses.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Professional and Intrapersonal Characteristics

Table 1 describes the socio-demographic, professional, intrapersonal, and willingness characteristics of respondents. Only 39.2% of the HCW had have any previous training in disaster. This was the same for previous work experience, where only 35.8% of the participants had previous work experience in a disaster situation.

The doctors and nurses reported a high trust in work safety, in family, in colleague's preparedness to react, and in hospital preparedness to react in case of a disaster, with trust in colleague's preparedness as the highest percentage (88.3%). Similarly, HCW had a high median score (16; IQR 2) in self-efficacy construct.

Respondents' willingness

Ninety percent of the participants expressed high willingness to participate in any type of disasters. However, they were less willing to participate in natural disasters (77.3%) and influenza pandemic (66.0%) (Table 1).

[\[Table 1\]](#)

Analysis of Self-efficacy's factors

There were four items in the self-efficacy construct. After the data collection, the inter-correlation value of self- efficacy was tested before running the main analysis. The descriptive statistics and inter-item correlation values of the self-efficacy are presented in Table 2.

[\[Table 2\]](#)

1
2
3 Based on Table 2, there was moderate level of agreement in all the four items. The
4
5 highest correlation for each item with at least one other item in the construct was between 0.3
6
7 and 0.9. In factor analysis, the Kaiser-Meyer-Olkin (KMO) value was 0.658 ($p < 0.001$), KMO
8
9 value close to 1, so the variables are suitable for factor analysis. Therefore, a single factor
10
11 was formed from the four items computed and was saved as incentive to be used in further
12
13 analysis. Accordingly, to calculate the self- efficacy, four questions was asked. The sum
14
15 outcome of the four answers ranged from 4 to 20 (Median 16, Interquartile range [IQR] 5).
16
17
18
19
20
21

22 **Factors associated with willingness to participate in any type disaster**

23
24
25 There was an association between participants' gender and willingness to participate
26
27 in any type of disaster, with male being more willing compared to female (crude OR=2.161,
28
29 95%CI: 1.307 to 3.573) (Table 3). Those with high trust in work safety (crude OR=3.284,
30
31 95%CI: 1.937 to 5.567), trust in colleague's preparedness (crude OR=2.592, 95%CI: 1.401 to
32
33 4.795) and self-efficacy (crude OR=1.358; 95%CI: 1.247 to 1.479) were also found to be
34
35 more willing to participate in any type of disaster in the univariate analysis (Table 3).
36
37
38

39 In the final model, having trust in work safety (adjusted OR=2.535, 95%CI: 1.357 to
40
41 4.736) and self-efficacy (adjusted OR=1.319, 95%CI: 1.197 to 1.453) were found to be
42
43 associated with general willingness with any type of disasters (Table 4).
44
45
46
47
48

49 **Factors associated with willingness to participate in natural disaster**

50
51
52 In the univariate analysis, there was an association between some of the socio-
53
54 demographic characteristics (age, gender, education level, type of profession and work
55
56 duration) with willingness to participate in natural disasters. Participants with bachelor and
57
58 postgraduate degrees had a higher odd of willingness to participate in natural disaster
59
60

1
2
3 disasters compared to those with diploma education. Those in the age group of between 31
4
5 and 45 were found to be more willing when compared to participants who were ≤ 30 years
6
7 old. The main intrapersonal factors associated with willingness in natural disasters were
8
9 participants' trust in colleagues' preparedness in case of a natural disaster, and self-efficacy
10
11 (Table 3).
12
13
14

15 In multivariate, being male (adjusted OR=1.639, 95%CI: 1.102 to 2.439) and self-
16
17 efficacy (adjusted OR=1.143, 95%CI: 1.069 to 1.221) were significantly associated with
18
19 willingness to participate in natural disasters (Table 4).
20
21
22
23
24

25 **Factors associated with willingness to participate in influenza pandemic disaster**

26
27
28 For willingness to participate in influenza pandemic, the univariate results revealed
29
30 that being male, having a dependent with no support (compared to participants without
31
32 dependent), having previous experience and self-efficacy were associated with willingness to
33
34 participate in pandemic (Table 3).
35
36
37

38 In the final model, having previous experience (adjusted OR=1.528, 95%CI: 1.058 to
39
40 2.207) and self-efficacy (adjusted OR=1.114, 95%CI: 1.050 to 1.182) were found to be
41
42 associated with willingness to participate in influenza pandemic (Table 4).
43
44

45 [[Table 3](#), [Table 4](#)]
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion:

This study contained a main question that is to test the strength and direction of association between the independent variables- socio-demographic, professional, and intrapersonal variables- with the variables of willingness to participate in three different types of disasters. The results showed that trust in work safety and self-efficacy were associated with disaster participation willingness after in the multivariate analysis. Gender and self-efficacy were found to be significantly associated with willingness to participate in natural disaster. While previous experience and self-efficacy were statistically significant with willingness of HCW to participate in influenza pandemic.

Having one hundred percent participation of HCW is difficult to obtain as much as it is vital in case of disasters. Previous studies suggested that between 65 and 97 percent of HCW were willing to participate in a natural disaster, and between 54 and 86 percent in an influenza pandemic.^{11,16,20,21} The reason in this difference in the levels of willingness between the two types of disasters is due to the great distinction in their nature. The outcome of the interaction between HCW and the socio-environmental determinants of these disasters leads to having different willingness levels. According to Connor et al., the weighted risk resulting from this interaction plays a major role in HCW willingness.¹⁰ In case of pandemics, the fear of the inability to control biohazards and watching colleagues inquiring a communicable disease after contact with affected persons was a suggested reason for the low willingness levels.¹⁰ Another qualitative study in Australia highlighted that in different types of disasters emergency nurses' willingness to attend to work is shaped by the weighted risk to self and surrounding people and the pressure formed from the period dealing with the disaster.¹² Natural disasters may not directly affect HCW or their families like influenza pandemic

1
2
3 which may be a reason for the higher level of willingness to attend to work compared to
4
5
6 pandemics.
7

8 The findings of this study are consistent with the previous studies regarding their
9
10 willingness level to different types of disasters. More HCW were willing to participate in
11
12 natural disaster (77.3%) as compared to influenza pandemic (66.0%). Even though, there
13
14 were a higher percentage of HCW willingness from this study to participate in both natural
15
16 disaster and influenza pandemic were higher compared to other studies, Yemen's high
17
18 vulnerability, due its topography and current economic, political and health status, could
19
20 affect the country health status and cause huge adverse health impacts in case of disasters.
21
22 According to the Vulnerability Matrix, one third of Yemen's districts are highly vulnerable.⁵
23
24
25
26

27 In addition, due to the armed conflicts in Yemen that started in early 2015, it had
28
29 become difficult to deal with any disaster in case it occurred. According to the World Health
30
31 Organization (WHO), the healthcare system in Yemen is in a critical situation with about
32
33 50% of the health facilities in Yemen are either partially or total damage as a result of the
34
35 natural disasters and conflicts. Furthermore, of the 3,507 healthcare facilities in Yemen,
36
37 almost 300 healthcare facilities have been destroyed.^{5,22} Therefore, any possible decrease in
38
39 the health work force during disasters in Yemen must be put in considerations during
40
41 response to disasters. The existed drained health sector is based on a vulnerable health system
42
43 in term of its structure and healthcare staff number and distribution in the country.²³ Thus,
44
45 training and preparing healthcare staff to act during disasters is critical and could help
46
47 achieving Priority 4 in the United Nations' Sendai Framework for Disaster Risk Reduction
48
49 2015-2030 in enhancing disaster preparedness for effective response and to "Build Back
50
51 Better" in recovery, rehabilitation and reconstruction of disasters, and prevent any possible
52
53 humanitarian catastrophe.²⁴
54
55
56
57
58
59
60

1
2
3 The results show that willingness to participate in events of natural disasters may be
4 encouraged by many suggested factors. Factors such as respondents being male are
5 accompanied with higher willingness to participate in natural disasters. The findings is
6 similar to previous studies conducted in Jordan, China and United States of America.^{15,16,25,26}
7
8 However, having less female health workers willing to participate in disasters highlight a
9 need to address this group during health preparedness programs. This lack in female HCW
10 willingness could be due to the gender and cultural barriers in Yemen. These barriers need to
11 be addressed to better motivate females workers participation in case of disasters. This is
12 important area for research as studies on willingness moving forward.
13
14

15
16 As in previous studies, nurses and doctors with previous experience in influenza
17 pandemic are more willing to participate in influenza pandemic.^{14, 25} HCW which has
18 experience with a previous disaster have better knowledge in disaster management, coping
19 strategies, and ways to protect themselves and their families; which may be a reason for
20 having higher odds of willingness. According to Alzahrani et al., Most of the nurses have a
21 higher level in Mass gatherings management such as communicating effectively during
22 emergencies in Saudi Arabia due to Previous experience, he also found a need for further
23 trainings by the nurses.²⁷ Chokahi et al., found that simulation training, attending conferences
24 and previous experience of increase paediatric surgeons feeling of preparedness.¹⁴
25
26

27
28 This study found that trust in work safety during any type of disaster play key role in
29 willingness to participate, which is similar to results from previous studies.^{11,25,28} Work safety
30 is important in order for HCW to feel safe and do their work, a study by Stergachis et al.,
31 found that majority of the participants reported their fear or concern for self in case of
32 influenza pandemic and during earthquake scenarios as one of the major barriers to
33 willingness.²⁹ Considering work safety in disaster preparedness is especially important in
34 Yemen after notifications made by Aldahrai et al., and Naser et al., on the current low work
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 place safety and the needs of increasing safety and security standards in hospitals in Yemen
4
5 Health establishments, regardless to external or internal catastrophes; unsafe work place
6
7 could lead to increase optional absenteeism during disaster.^{7,8} Thus, putting information on
8
9 work safety and protection devices provided during several types of disasters is advisable.
10
11
12

13 Many studies have established the association between self-efficacy and willingness
14
15 to participate in disasters in previous.^{10,15,16} Similarly, this study have identified that self-
16
17 efficacy plays an important role in willingness to participate in natural disasters, in influenza
18
19 pandemic and any type of disasters. It indicates that elements tested for self-efficacy like
20
21 participants increase familiarity with their role, responsibility to react, and confidence of
22
23 ability to deal with various types of disasters could be a key factor to increase HCW self-
24
25 efficacy, thus, increase their willingness. Disaster preparedness trainings are encouraged to
26
27 contain materials that explain response to different type of disasters, doctors' and nurses' role
28
29 in disasters and how it makes a different in response to disasters. Supportive measures to
30
31 increase HCW self-efficacy such as immediate communication with needed information and
32
33 rewarding for efforts in disasters are suggested.
34
35
36
37
38

39 Although the study has faced challenges in the sense of collecting the data during
40
41 political unrest in the country, the data was collected from three of the major public hospital.
42
43 The quality of data was insured by explaining the study and its objectives, and answering any
44
45 question that may arise. This prevented difference in understanding the questions and
46
47 increased participants' inclusion in the study. All and all the study suggested the vital value
48
49 of increasing self-efficacy and its element in order to obtain more willingness.
50
51
52
53
54
55
56
57
58
59
60

Strength and limitation

This is the first study explored the patterns and associated factors of the willingness status to participate in disasters among healthcare workers in Yemen. The response rate was also high as the questionnaire was distributed personally by the researcher. The questionnaire was pilot tested; this was to ensure that the validity of questionnaire.

This study had limitations, as the study was performed under a political insecurity/fragile state of the country a list of workers of HCW working at the hospitals was difficult to obtain. As a result, a universal sample was undertaken which could limit the study representativeness. This is a cross-sectional study using a self-administered questionnaire, where the actual willingness of the respondents cannot be ascertained. The respondents may answer positively due to social desirability bias. Other than the self-administrated questionnaire limitations, the study findings are limited to staff working at the tertiary level public hospitals in urban areas of Yemen.

Conclusion:

Increased the likelihood of willingness to participate in disaster play a key role in guarantee optimum number of work force. This study indicates that one's socio-demographic, professional and intrapersonal factors plays a role in increase his/her willingness in general and across different type of disasters. Significant differences were revealed between participants' willingness in natural disasters by gender and self-efficacy, and participant's willingness in influenza pandemics by previous experience and self-efficacy. This result suggests integrating disaster management into an earlier education levels of doctors' and nurses' educational establishments. This could be achieved by adding early exposure of

1
2
3 healthcare workers to relevant disaster experience which would further boost their
4
5 willingness to participate in disaster response.
6
7

8 Other areas for preparedness may include increase hospitals safety and resilience;
9
10 hence that willingness to participate in natural disasters' response was influence by
11
12 demographic characteristic of the healthcare personnel, others found outside the bracket
13
14 should be motivated to participate with a reward package such as incentives and hazard
15
16 allowance. Further studies should be conducted in both urban and rural settings with
17
18 relatively peaceful atmosphere.
19
20
21
22
23
24

25 **Acknowledgement:** We thank Dr. Susan Connor, Dr. Karen Hammad and
26
27 Dr..Mohammed Khalil for their Support and advice during the study development, all
28
29 University Malaya academic staff, the team who collected the data, and all the participants
30
31 who contributed in this study.
32
33
34

35 **Competing interests:** None declared
36
37

38 **Funding:** The authors have no financial disclosures to declare. This research did not
39
40 receive any specific grant from funding agencies in the public, commercial, or not-for-profit
41
42 sectors.
43
44
45

46 **Data sharing statement:** No additional data are available.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reference:

1. United Nations Economic and Social Commission for Asia and the Pacific. Disasters in Asia and the Pacific: 2015 year in review. United Nations report. Economic and social commission for Asia and the Pacific. 2015. [21 p.]. Available from: <https://www.unescap.org/resources/disasters-asia-and-pacific-2015-year-review>
2. Below R, Wirtz A, Guha-Sapir D. Disaster category classification and peril terminology for operational purposes. Centre for Research on the Epidemiology of Disasters. 2009. Available from: <https://www.cred.be/node/564>
3. Joseph, B., & Joseph, M. (2016). The health of the healthcare workers. Indian journal of occupational and environmental medicine, 20(2), 71–72. doi:10.4103/0019-5278.197518.
4. Ministry of Water and Environment. Yemen: National progress report on the implementation of the Hyogo Framework for Action (2013-2015). 2015. Available from: <http://preventionweb.net/go/40141>
5. World Health Organization. WHO annual report 2017: Yemen. Cairo: WHO Regional Office for the Eastern Mediterranean; 2018. Licence: CC BYNC-SA 3.0 IGO.
6. Banerjee, Aditi; Bhavnani, Rakhi; Burtonboy, Catherine H, et al. Natural Disasters in the Middle East and North Africa: A Regional Overview. The International Bank for Reconstruction and Development / The World Bank. [101 p.]. 2014; Report No.: 81658.
7. Aladhrai SA, Djalali A, Della Corte F, Alsabri M, El-Bakri NK, Ingrassia PL. Impact of the 2011 Revolution on Hospital Disaster Preparedness in Yemen. Disaster Medicine and Public Health Preparedness. 2015;9(4):396-402.
8. Naser WN, Ingrassia PL, Aladhrae S, Abdulraheem WA. A Study of Hospital Disaster Preparedness in South Yemen. Prehospital and Disaster Medicine. 2018;33(2):133.
9. Chaffee M. Willingness of health care personnel to work in a disaster: an integrative review of the literature. Disaster Medicine and Public Health Preparedness. 2009;3(1):42-56.

10. Connor SB. When and why health care personnel respond to a disaster: the state of the science. *Prehospital and Disaster Medicine*. 2014;29(3):270-4.
11. Gowing JR, Walker KN, Elmer SL, Cummings EA. Disaster Preparedness among Health Professionals and Support Staff: What is Effective? An Integrative Literature Review. *Prehospital and Disaster Medicine*. 2017;32(3):321-8.
12. Arbon P, Cusack L, Ranse J, Shaban RZ, Considine J, Kako M, et al. Exploring staff willingness to attend work during a disaster: a study of nurses employed in four Australian emergency departments. *Australasian Emergency Nursing Journal : AENJ*. 2013;16(3):103-9.
13. Ejeta LT, Ardalan A, Paton D. Application of Behavioral Theories to Disaster and Emergency Health Preparedness: A Systematic Review. *PLOS Currents Disasters*. 2015;7. Doi: 10.1371/currents.dis.31a8995ced321301466db400f1357829.
14. Chokshi NK, Behar S, Nager AL, Dorey F, Upperman JS. Disaster management among pediatric surgeons: preparedness, training and involvement. *American Journal of Disaster Medicine*. 2008;3(1):5-14.
15. Burke RV, Goodhue CJ, Chokshi NK, Upperman JS. Factors associated with willingness to respond to a disaster: a study of healthcare workers in a tertiary setting. *Prehospital and Disaster Medicine*. 2011;26(4):244-50.
16. Charney RL, Rebmann T, Flood RG. Hospital Employee Willingness to Work during Earthquakes Versus Pandemics. *Journal of Emergency Medicine*. 2015;49(5):665-74. Doi: <https://doi.org/10.1016/j.jemermed.2015.07.030>.
17. Balicer RD, Omer SB, Barnett DJ, Everly GS, Jr. Local public health workers' perceptions toward responding to an influenza pandemic. *BMC Public Health*. 2006;6:99.
18. Shapira S, Aharonson-Daniel L, Bar-Dayyan Y, Sykes D, Adini B. Knowledge, perceptions, attitudes and willingness to report to work in an earthquake: A pilot study comparing Canadian versus Israeli hospital nursing staff. *International Emergency Nursing*. 2016;25:7-12.
19. Bandura A. *Self-efficacy: The exercise of control*: Macmillan; 1997.
20. Arbon P, Ranse J, Cusack L, Considine J, Shaban RZ, Woodman RJ, et al. Australasian emergency nurses' willingness to attend work in a disaster: a survey. *Australasian Emergency Nursing Journal : AENJ*. 2013;16(2):52-7.

- 1
2
3 21. Brice JH, Gregg D, Sawyer D, Cyr JM. Survey of Hospital Employees' Personal
4 Preparedness and Willingness to Work Following a Disaster. *Southern Medical*
5 *Journal*. 2017;110(8):516-22.
6
7
- 8 22. United Nations Office for the Coordination of Humanitarian Affairs. *Humanitarian*
9 *Bulletin Yemen/ Health system on the verge of collapse*. 2015. [4 p.]. Available
10 from: [http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-](http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-30-november-2015-enar)
11 [30-november-2015-enar](http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-30-november-2015-enar)
12
13
- 14 23. Qirbi N, Ismail SA. Health system functionality in a low-income country in the
15 midst of conflict: the case of Yemen. *Health Policy and Planning*. 2017;32(6):911-
16 22.
17
18
- 19 24. United Nations - Headquarters. *Sendai Framework for Disaster Risk Reduction*
20 *2015-2030*. United Nations Office for Disaster Risk Reduction. 2015. [32 p.].
21 Available from: <https://www.unisdr.org/we/inform/publications/43291>
22
23
- 24 25. Loke AY, Fung WM, Liu X. Mainland China nurses' willingness to report to work
25 in a disaster. *American Journal of Disaster Medicine*. 2013;8(4):273-82. Doi:
26 [10.5055/ajdm.2013.0134](https://doi.org/10.5055/ajdm.2013.0134).
27
28
- 29 26. Al Khalaileh MA, Bond E, Alasad JA. Jordanian nurses' perceptions of their
30 preparedness for disaster management. *International emergency nursing*.
31 *2012;20(1):14-23*.
32
33
- 34 27. Alzahrani F, Yiannis K. Emergency nurse disaster preparedness during mass
35 gatherings: a cross-sectional survey of emergency nurses' perceptions in hospitals in
36 Mecca, Saudi Arabia. *BMJ Open*. 2017;7(4):e013563.
37
38
- 39 28. Ogedegbe C, Nyirenda T, Delmoro G, Yamin E, Feldman J. Health care workers
40 and disaster preparedness: barriers to and facilitators of willingness to respond.
41 *International Journal of Emergency Medicine*. 2012;5(1):29.
42
43
- 44 29. Stergachis A, Garberson L, Lien O, D'Ambrosio L, Sangare L, Dold C. Health care
45 workers' ability and willingness to report to work during public health emergencies.
46 *Disaster Medicine and Public Health Preparedness*. 2011;5(4):300-8.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Socio-demographic, professional, intrapersonal, and willingness Characteristics of Respondents

Variable		Frequency (n= 692)	Percentage (%)
Age (years)	≤30	360	52.0
	31-45	294	42.5
	≥46	38	5.5
Gender	Male	419	60.5
	Female	273	39.5
Marital-status	Single	258	37.3
	Married	420	60.7
	Divorce	8	1.2
	Widow	6	0.9
Education-level	Diploma	280	40.5
	Bachelor	275	39.7
	Post graduate education	137	19.8
	Master	104	15.0
Professional	33	4.8	
Profession type	Doctors	311	44.9
	Specialist medical practitioner	126	8.2
	General medical practitioner	185	26.7
	Nurses	381	55.1
Work duration (years)	≤5	360	52
	6-10	213	30.8
	11-15	76	11.0
	≥16	43	6.2
Dependent	With dependent	456	65.9
	Elder	65	9.4
	Child	276	39.9
	Both	115	16.6
	No dependent	236	34.1
^a Support	No	119	26.1
	Yes	337	73.9
^b Previous training	With previous training	271	39.2
	Without previous training	421	60.8
^b Previous experience	With previous experience	248	35.8
	Without previous training	444	64.2
Trust in work safety in case of disaster	High trust in work safety	563	81.4
	Low trust in work safety	129	18.6
Trust in family preparedness in case of a disaster	High trust in family preparedness	544	78.6
	Low trust in family preparedness	148	21.4
Trust in colleague preparedness to react in disaster	High trust in colleague preparedness	611	88.3
	Low trust in colleague preparedness	81	11.7

Trust in hospital preparedness to react in disaster	High trust in hospital preparedness	522	75.4
	Low trust in hospital preparedness	170	24.6
Self-efficacy score (Median; Inter-Quartile)		16	5
Willingness to participate in Any type of disaster	High	623	90.0
	Low	69	10.0
Willingness to participate in Natural disaster	High	535	77.3
	Low	157	22.7
Willingness to participate in Influenza and Pandemic	High	457	66.0
	Low	235	34.0

^aOnly for participants with dependants. The percentages are only within participants with dependents.

^bPrevious experience and training in disasters

Table 2: Descriptive statistics and inter-term correlation for items in self-efficacy

Items	Descriptive statistics		Inter-item correlation			
	Mean	SD	item 1	item 2	item 3	item 4
Item 1: Ability to perform work	3.36	1.074	1.000	0.592	0.267	0.259
Item 2: Familiarity with their role	3.62	1.085	0.592	1.000	0.325	0.324
Item 3: Responsibility to participate	4.23	0.929	0.267	0.325	1.000	0.390
Item 4: Ability report to work	4.03	1.062	0.259	0.324	0.390	1.000

For peer review only

Table 3: Univariate association of crude socio-demographic, professional and intrapersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters		Willingness to participate in natural disasters		Willingness to participate in influenza pandemic	
	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value
Age						
≤30	1	1	1	1	1	1
31-45	1.056 (0.628-1.775)	0.838	1.639 (1.126-2.387)	0.010	1.138 (0.822-1.576)	0.437
≥46	0.733 (0.269-1.997)	0.544	1.967 (0.798-4.850)	0.142	1.354 (0.650-2.820)	0.418
Gender						
Female	1	1	1	1	1	1
Male	2.161 (1.307-3.573)	0.003	2.254 (1.571-3.234)	<0.001	1.505 (1.093-2.072)	0.012
Marital status						
Single	1	1	1	1	1	1
Married	0.899 (0.532-1.521)	0.692	1.128 (0.781-1.629)	0.521	1.090 (0.787-1.510)	0.605
Divorce and widow	0.615 (0.130-2.913)	0.541	1.160 (0.314-4.291)	0.824	1.362 (0.416-4.466)	0.610
Education Level						
Up to Diploma	1	1	1	1	1	1
Up to Bachelor	1.380 (0.775-2.458)	0.274	1.962 (1.317-2.922)	0.001	1.221 (0.858-1.738)	0.268
Up to Postgraduate	0.847 (0.450-1.596)	0.608	2.279 (1.351-3.842)	0.002	0.998 (0.651-1.528)	0.992
Profession type						
Nurses	1	1	1	1	1	1
General medical practitioner	1.473 (0.781-2.776)	0.231	1.554 (1.013-2.384)	0.044	0.756 (0.524-1.091)	0.135
Specialist medical practitioner	0.965 (0.507-1.835)	0.913	2.407 (1.377-4.208)	0.002	0.942 (0.614-1.446)	0.785
Work duration						

≤5	1	1	1	1	1	1	1
6-10	1.006 (0.563-1.798)	0.983	1.079 (0.721-1.614)	0.711	0.975 (0.683-1.392)	0.889	
11-15	1.028 (0.438-2.415)	0.949	2.337 (1.118-4.882)	0.024	1.059 (0.626-1.791)	0.832	
≥16	0.456 (0.196-1.063)	0.069	0.650 (0.329-1.286)	0.216	1.198 (0.603-2.379)	0.606	
Presence of dependent							
No dependent	1	1	1	1	1	1	
Dependent with no support	0.673 (0.346-1.308)	0.242	1.149 (0.683-1.933)	0.601	1.617 (1.000-2.612)	0.050	
Dependent with support	1.253 (0.707-2.222)	0.439	1.305 (0.880-1.935)	0.185	1.283 (0.907-1.815)	0.158	
^aPrevious training							
No	1	1	1	1	1	1	
Yes	1.653 (0.959-2.849)	0.070	1.090 (0.756-1.573)	0.644	1.147 (0.829-1.585)	0.408	
^aPrevious experience							
No	1	1	1	1	1	1	
Yes	1.216 (0.714-2.070)	0.471	0.974 (0.673-1.411)	0.889	1.857 (1.317-2.619)	<0.001	
Trust in work safety							
Low	1	1	1	1	1	1	
High	3.284 (1.937-5.567)	<0.001	1.280 (0.825-1.987)	0.271	1.192 (0.800-1.774)	0.388	
Trust in family preparedness							
Low	1	1	1	1	1	1	
High	1.338 (0.756-2.369)	0.317	1.292 (0.850-1.963)	0.231	0.990 (0.674-1.454)	0.959	
Trust in colleague's preparedness							
Low	1	1	1	1	1	1	
High	2.592 (1.401-4.795)	0.002	1.974 (1.200-3.246)	0.007	1.392 (0.866-2.237)	0.172	
Trust in hospital preparedness							
Low	1	1	1	1	1	1	
High	1.392 (0.807-2.400)	0.234	1.212 (0.810-1.814)	0.351	1.158 (0.807-1.663)	0.426	
^bSelf-efficacy							
	1.358 (1.247-1.479)	<0.001	1.184 (1.115-1.257)	<0.001	1.135 (1.076-1.197)	<0.001	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

^aPrevious experience and training in disasters

^b*Continuous measure, with one unit increase in self-efficacy*

For peer review only

Table 4: Multivariate association of adjusted professional and intrapersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters		Willingness to participate in natural disasters		Willingness to participate in influenza pandemic	
	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value
Age						
≤30	1	1	1	1	1	1
31-45	1.137 (0.609-2.121)	0.687	1.432 (0.928-2.209)	0.105	1.008 (0.695-1.461)	0.967
≥46	0.440 (0.136-1.418)	0.169	1.087 (0.409-2.886)	0.867	1.062 (0.484-2.332)	0.880
Gender						
Female	1	1	1	1	1	1
Male	1.456 (0.807-2.628)	0.212	1.639 (1.102-2.439)	0.015	1.131 (0.793-1.612)	0.498
Education Level						
Up to Diploma	-	-	1	1	-	-
Up to Bachelor	-	-	1.706 (0.993-2.932)	0.053	-	-
Up to Postgraduate	-	-	1.177 (0.522-2.657)	0.695	-	-
Profession type						
Nurses	1	1	1	1	1	1
General medical practitioner	1.285 (0.438-3.767)	0.261	1.581 (0.691-3.614)	0.278	0.751 (0.468-1.205)	0.235
Specialist medical practitioner	1.392 (0.578-3.352)	0.644	0.939 (0.515-1.713)	0.838	0.697 (0.473-1.026)	0.067
Presence of dependent						
No dependent	-	-	-	-	1	1
Dependent with no support	-	-	-	-	1.537 (0.910-2.598)	0.108
Dependent with support	-	-	-	-	1.154 (0.791-1.685)	0.457

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

^aPrevious experience						
No	-	-	-	-	1	1
Yes	-	-	-	-	1.528 (1.058-2.207)	0.024
Trust in work safety						
Low	1	1	-	-	-	-
High	2.535 (1.357-4.736)	0.004	-	-	-	-
Trust in colleague's preparedness						
Low	1	1	1	1	-	-
High	1.199 (0.576-2.496)	0.686	1.363 (0.791-2.351)	0.265	-	-
^bSelf-efficacy	1.319 (1.197-1.453)	<0.001	1.143 (1.069-1.221)	<0.001	1.114 (1.050-1.182)	<0.001
^a Previous experience and training in disasters						
^b Continuous measure, with one unit increase in self-efficacy						

Peer review only

SECTION ONE:

Please tick (✓) in the appropriate box

1. **Age:** _____

2. **Gender :**

Male

Female

3. **Marital status:**

Single

Married

Divorce

Widow/er

4. **Highest Education level completed :**

Diploma

Bachelor

Masters

Professional

5. **What is your primary job category?**

Specialist medical practitioner

General medical practitioner

Nurse

6. **Duration of work at the hospital:** _____ Years.

7. Do you have any of the following responsibilities (you can choose more than one choice):

Children Elders None

8. Do you have any one who can take care of the previously mentioned responsibilities in case of your absence?

Yes No

SECTION TWO

Please tick (√) in the appropriate box:

1. Have you participated in a disaster before?

Yes No

2. Have attended any previous disaster trainings?

Yes No

SECTION THREE:

Part A:

NO	Questions	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am confident of my personal safety at work in case of a disaster.					

2.	I am assured that my family is prepared to function in in my absence during a disaster.					
3.	I am sure that my colleagues are able to perform their duties during a disaster.					
4.	The hospital is prepared to provide effective response in case of a disaster.					

PART B:

NO	Statement	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am able to treat patients of different type of disasters.					
2.	I am confident that I can perform my role in the hospital following any type of disasters.					
3.	I feel that it is my duty to work in the event of a disaster.					
4.	I will be able to report to work at the hospital during an event of a disaster.					

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

SECTION FOUR:

NO	Statement	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am willing to participate in any type of disaster regardless of its severity					
2.	I am willing to participate in natural disasters (earthquake, floods or cyclone)					
3.	I am you willing to participate in influenza pandemic					

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	Main document (1)
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	Main document (1)
Background /	#2	Explain the scientific background and rationale for the	Main document

1	rationale		investigation being reported	(3)
2				
3				
4	Objectives	#3	State specific objectives, including any prespecified	Main document
5			hypotheses	(4)
6				
7				
8				
9	Study design	#4	Present key elements of study design early in the paper	Main document
10				(4)
11				
12				
13				
14	Setting	#5	Describe the setting, locations, and relevant dates,	Main document
15			including periods of recruitment, exposure, follow-up,	(4-6)
16			and data collection	
17				
18				
19				
20				
21				
22	Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods	Main document
23			of selection of participants.	(4)
24				
25				
26				
27		#7	Clearly define all outcomes, exposures, predictors,	Main document
28			potential confounders, and effect modifiers. Give	(5), table1
29			diagnostic criteria, if applicable	(Page 18)
30				
31				
32				
33				
34				
35	Data sources /	#8	For each variable of interest give sources of data and	Main document
36	measurement		details of methods of assessment (measurement).	(5)
37			Describe comparability of assessment methods if there	
38			is more than one group. Give information separately for	
39			exposed and unexposed groups if applicable.	
40				
41				
42				
43				
44				
45				
46				
47	Bias	#9	Describe any efforts to address potential sources of bias	Main document
48				(13)
49				
50				
51				
52	Study size	#10	Explain how the study size was arrived at	Main document
53				(4,5)
54				
55				
56				
57				
58	Quantitative	#11	Explain how quantitative variables were handled in the	Main document
59				
60				

1	variables	analyses. If applicable, describe which groupings were	(6)
2			
3		chosen, and why	
4			
5			
6	Statistical	#12a Describe all statistical methods, including those used to	Main document
7			
8	methods	control for confounding	(6)
9			
10			
11		#12b Describe any methods used to examine subgroups and	Main document
12			
13		interactions	(6)
14			
15			
16		#12c Explain how missing data were addressed	Main document
17			
18			(6)
19			
20			
21			
22		#12d If applicable, describe analytical methods taking account	Main document
23			
24		of sampling strategy	(4, 6)
25			
26			
27		#12e Describe any sensitivity analyses	
28			
29			
30	Participants	#13a Report numbers of individuals at each stage of study—	Main document
31			
32		eg numbers potentially eligible, examined for eligibility,	(4, 6)
33			
34		confirmed eligible, included in the study, completing	
35			
36		follow-up, and analysed. Give information separately for	
37			
38		for exposed and unexposed groups if applicable.	
39			
40			
41			
42		#13b Give reasons for non-participation at each stage	Main document
43			
44			
45			(4, 6, 13)
46			
47			
48		#13c Consider use of a flow diagram	
49			
50			
51	Descriptive data	#14a Give characteristics of study participants (eg	Main document
52			
53		demographic, clinical, social) and information on	(7,8), table 2
54			
55		exposures and potential confounders. Give information	(page 19, 20)
56			
57		separately for exposed and unexposed groups if	
58			
59			
60			

applicable.

1		applicable.	
2			
3			
4		#14b Indicate number of participants with missing data for	Table 2 (page
5			
6		each variable of interest	19, 20)
7			
8			
9	Outcome data	#15 Report numbers of outcome events or summary	Main document
10			
11		measures. Give information separately for exposed and	(7), table 2
12			
13		unexposed groups if applicable.	(page 19, 20)
14			
15			
16	Main results	#16a Give unadjusted estimates and, if applicable,	Main document
17			
18		confounder-adjusted estimates and their precision (eg,	(8, 9), table 3,
19			
20		95% confidence interval). Make clear which confounders	4 (page 21-23)
21			
22		were adjusted for and why they were included	
23			
24			
25			
26		#16b Report category boundaries when continuous variables	Main document
27			
28		were categorized	5
29			
30			
31			
32		#16c If relevant, consider translating estimates of relative risk	
33			
34		into absolute risk for a meaningful time period	
35			
36			
37	Other analyses	#17 Report other analyses done—e.g., analyses of	
38			
39		subgroups and interactions, and sensitivity analyses	
40			
41			
42	Key results	#18 Summarise key results with reference to study objectives	Main document
43			
44			(6-9)
45			
46			
47			
48	Limitations	#19 Discuss limitations of the study, taking into account	Main document
49			
50		sources of potential bias or imprecision. Discuss both	(12, 13)
51			
52		direction and magnitude of any potential bias.	
53			
54			
55	Interpretation	#20 Give a cautious overall interpretation considering	Main document
56			
57		objectives, limitations, multiplicity of analyses, results	(9-12)
58			
59			
60			

1 from similar studies, and other relevant evidence.

2
3
4 Generalisability [#21](#) Discuss the generalisability (external validity) of the Main document
5 study results (13)
6
7

8
9 Funding [#22](#) Give the source of funding and the role of the funders for Main document
10 the present study and, if applicable, for the original study (14)
11 on which the present article is based
12
13
14
15

16
17 The STROBE checklist is distributed under the terms of the Creative Commons Attribution License
18 CC-BY. This checklist can be completed online using <https://www.goodreports.org/>, a tool made by
19 the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

Factors associated with Healthcare Workers Willingness to Participate in Disasters: a cross-sectional study in Sana'a, Yemen

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-030547.R2
Article Type:	Original research
Date Submitted by the Author:	04-Sep-2019
Complete List of Authors:	Al-Hunaishi, Weiam; University of Malaya, Department of Social and Preventive Medicine Hoe, Victor; Universiti Malaya Chinna, Karuthan ; Taylor's University, School of Medicine-SRI
Primary Subject Heading:	Public health
Secondary Subject Heading:	Global health, Health services research, Public health
Keywords:	Disaster preparedness, willingness, self-efficacy, healthcare worker, intrapersonal

SCHOLARONE™
Manuscripts

1
2
3
4
5 **Factors associated with Healthcare Workers Willingness to Participate in Disasters: a**
6
7 **cross-sectional study in Sana'a, Yemen**
8
9

10 **Authors' names, credentials/degrees, email addresses and institutional affiliations:**
11

- 12
13 1. **MPH. MD. Weiam Al-Hunaishi (WA):** Department of Social & Preventive Medicine,
14 Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. (e-mail:
15 waam59@hotmail.com)
16
17
18
19 2. **Prof. Dr. Victor CW Hoe (VCWH):** Centre for Occupational and Environmental Health,
20 Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya,
21 Kuala Lumpur, Malaysia. (e-mail: victor@ummc.edu.my)
22
23
24
25 3. **Associate Prof. Dr. Karuthan Chinna (KC):** School of Medicine-SRI, Faculty of Health &
26 Medical Sciences, Taylor's University, Malaysia. (e-mail: karuthan.chinna@taylors.edu.my)
27
28
29
30

31 **Correspondence and reprint requests:** to Weiam Al-Hunaishi (e-mail: Waam59@hotmail.com,
32 telephone number: +60173479802, mail address: Department of Social and Preventive Medicine
33 Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia).
34
35
36
37

38 **Contributorship statement:**
39

- 40
41 • WA – conceptualized the study, conducted the data collection, analysis, and interpretation of
42 data for the work; draft and approve the manuscript for publication.
43
44
45 • VCWH – conceptualized the study, interpretation of data for the work; revise and approve the
46 manuscript for publication.
47
48
49
50 • KC – conceptualized the study, supervised the data analysis, interpretation of data for the
51 work.
52
53
54
55
56
57
58
59
60

1
2
3 **List of Abbreviations:**
4
5

6	CI	Confidence Interval
7		
8	HCW	Healthcare workers
9		
10	IQR	Interquartile range
11		
12	SD	Standard Deviation
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

For peer review only

Abstract

Objectives: Willingness to participate in disasters are usually overlooked and not addressed in disaster preparedness training courses to ensure health service coverage. This will lead to issues during the disaster's response. This study, therefore, aims to assess healthcare workers willingness to participate in biological and natural disasters, and to identify its associated factors.

Design: This is a cross-sectional study using a self-administered questionnaire. The questionnaire was distributed to 1093 healthcare workers. The data was analysed using multiple logistic regression with significance level $p < 0.05$. Ethical clearance and consent of the participants were duly obtained.

Setting: In three public Hospitals that provide tertiary level healthcare in Sana'a City, Yemen.

Participants: There were 692 nurses and doctors (response rate 63.3%) completed the questionnaires.

Results: Almost half of the participants 55.1% were nurses and 44.9% were doctors. The study found that self-efficacy was associated with willingness to participate in disaster response for any type of disasters (Odds Ratio [OR]=1.328, 95% Confidence Interval [95%CI]: 1.206 to 1.464), natural disasters (OR=1.138, 95%CI: 1.064 to 1.217), and influenza pandemic (OR=1.112, 95%CI: 1.049 to 1.180). The results further show that willingness is associated with healthcare workers being young, male and having higher educational qualifications.

Conclusion: Self-efficacy has been found to be an important factor associated with willingness. Improving self-efficacy through training in disaster preparedness may increase willingness of healthcare workers to participate in a disaster.

1
2
3 **Key words:** Disaster preparedness; willingness; self-efficacy; intrapersonal; healthcare
4
5 worker.
6
7

8 **Strength and limitations of this study** 9

- 10
11 - This study is the first study that attempts to assess Healthcare workers willingness to
12 participate in different types of disasters in Sana'a City, Yemen.
13
14 - Implementing a cross-sectional design and using a convenience sample has made it
15 difficult to establish causal association.
16
17 - The study sample represents healthcare workers in Sana'a City in terms of gender and
18 locality.
19
20 - The face to face approach, which included introduction of the study and its objectives,
21 ensured a higher response rate and minimised the risk of selection bias.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction:

In various parts of the world, disasters destroy communities and infrastructures, causing huge material and human losses.¹ Similarly, hospitals and health centres are also affected by the disasters, which restrict its work of relieving disaster-stricken communities. By 2025, more than half of the world population will live in cities, particularly in urban cities located along seismic fault lines, flood and other natural disaster-prone areas.² Therefore, healthcare workers (HCW) should be prepared to manage the influx of patients during possibly life threatening circumstances. A healthcare worker is defined as “the one who delivers care and services to the sick and ailing either directly as doctors and nurses or indirectly as aides, helpers, laboratory technicians, or even medical waste handlers”.³

According to the Global Facility for Disaster Reduction and Recovery (GFDRR), Yemen is one of the priority countries in the Middle East and North African region, mainly due to its vulnerability to disasters.⁴⁻⁶ Natural disasters, which includes storms, landslides, earthquake and floods are recurrent in Yemen. Floods due to the monsoonal rainfall, is responsible for most of the mortality due to disasters.⁶ Currently in Yemen, disaster management mainly focuses on responding to post disasters damage, and there is a lack of disaster preparedness, such as training and mock drills. The insufficient training makes it difficult to maintain preparedness.^{4,7,8} A competent prepared healthcare worker could better mitigate and respond to the community health needs during crises, which in turn will elevate health outcomes.

During disasters, HCW are expected to provide health care assistance to people suffering from the disasters alongside with caring for their usual patients. Some are also required to care for their dependents. Most of the previous studies reported an anticipated decrease in health workforce during a disaster as not all HCW are willing to participate in a

1
2
3 disaster. Therefore, understanding the factors associated with willingness to participate in a
4
5 disaster would allow more effective planning for a disaster situation. Systematic reviews have
6
7 found that willingness was associated with factors like the nature and type of event;
8
9 competing obligations between personal and professional needs; the work environment and
10
11 climate including personal safety, and the relationship between knowledge and perceptions of
12
13 efficacy.^{9,10}
14
15

16
17 Disaster preparedness activities and routine disasters trainings have been found to
18
19 improve the knowledge, skills and attitude preparedness of disasters.¹¹ However, the factors
20
21 that affect the willingness to participate in disasters are beyond just having knowledge on
22
23 disaster management.⁹⁻¹²
24
25

26
27 In a recent systematic review, they found only few studies that have been conducted
28
29 in the Middle East on the willingness to participate in disasters. They recommend that further
30
31 research should be conducted based on behavioural theories to better understand the Middle
32
33 East context of willingness to participate in a disaster.¹³ In another systematic review of 70
34
35 studies on HCW willingness to participate in disasters, only 12 studies were from Asia and
36
37 none of those studies were from the Middle East.¹¹ Therefore, the objective of this study is to
38
39 determine the associations between socio-demographic, professional, and intrapersonal
40
41 factors associated with the doctors' and nurses' willingness to participate in disasters
42
43 response in Sana'a, Yemen. The study factors are built based on the self-efficacy behavioural
44
45 theory.
46
47
48
49

50 **Methods:**

51 **Study design, population and instrument**

52
53 This is a cross-sectional study conducted in three public hospitals in the Sana'a
54
55 Governorate, Yemen; Al-Thora, Al-Jumhuri, and Al-Kuwait hospitals, which have more
56
57
58
59
60

1
2
3 than a total of 4,000 nurses and doctors, which is the focus of this study. Out of this, 1093
4
5 HCW were selected through convenient sampling by approaching the healthcare workers
6
7 individually. The sample size calculation was determined using the OpenEpi software. It was
8
9 based on a 95% confidence interval with a power of set to 80%. The sample size of 1093 was
10
11 determined based on the variable “perception of responsibility to participate” from a pre-
12
13 study, which is an element of the self-efficacy construct with the highest sample size to
14
15 achieve an association.¹⁴ The determined sample size was 614, which was inflated by 78
16
17 percent based on the lowest response rate of previous studies to reach 1093.¹⁵
18
19
20
21

22 The questionnaire was developed by the researcher based on information from
23
24 previous studies and opinions of national and international experts (supplementary
25
26 martial).¹⁵⁻¹⁸ The questionnaire was pre tested for validity and reliability. The questionnaire
27
28 was distributed to national and international experts together with questions on the validity of
29
30 the questionnaire with a scale of 1 to 4 (1 not relevant to 4 highly relevant). It tested the
31
32 questionnaire’s consistency, relatedness, representativeness and clarity of wording. From the
33
34 results of the pre-test, six items of the questionnaire that was related to the knowledge
35
36 construct, was deleted and changes on the questions were reworded. The revised
37
38 questionnaires were then forward and backward-translated form English to Arabic and from
39
40 Arabic to English. Following that, three other HCW were asked to give their opinion on the
41
42 questions based on the objective of the study.
43
44
45
46
47

48 The final questionnaire was pilot tested on 20 doctors and nurses from a hospital,
49
50 other than hospitals that have been chosen for the study. The internal consistency was
51
52 assessed using Cronbach’s alpha. Self- efficacy was 0.801 and this is considered good inter-
53
54 correlation. Therefore, the self-efficacy four items were used in the final survey to build the
55
56 construct as one of the intrapersonal factors (unpublished).
57
58
59
60

1
2
3 The questionnaire consists of socio-demographic, professional, and intrapersonal
4 factors, and willingness to participate in a disaster. The professional, intrapersonal and
5 willingness questions were measured on a five-point Likert scale (1=strongly disagree,
6 2=disagree, 3=somewhat agree, 4=agree, and 5=strongly agree). A binary variable was
7 created based on “less or equal two” as low and “more than 2” as high.
8
9
10
11
12
13
14

15 **Self-efficacy theory**

16
17 The self-efficacy theory was chosen to build the self-efficacy construct as it illustrates
18 beliefs that drive actions to face and solve problems that are faced to achieve the intended
19 goals. In case of disaster, the theory could be applied to HCW who are coping with fear and
20 threat and adapting new behaviours because of their beliefs in their competences. This belief
21 is derived from their successful performances, observing colleagues and managers positive
22 behaviours, convincing by a superior person, and calming the physiological and emotional
23 pressure caused by the threat.¹⁹
24
25
26
27
28
29
30
31
32
33
34
35
36

37 **Data collection**

38
39 Ethical approval was obtained from the Ministry of Health and Population of Yemen.
40 Informed consent was obtained from the respondents and confidentiality of personal
41 disclosures was re-assured. The self-administered questionnaires were distributed and
42 retrieved between February and March 2018. A final 767 questionnaires were returned for
43 analyses; 75 questionnaires were omitted due to missing values.
44
45
46
47
48
49
50
51
52
53

54 **Statistical analysis**

55
56
57 The data was analysed using Statistical Package for Social Science (SPSS) version 22.
58 Descriptive outputs were generated describing the median, interquartile range, frequency
59
60

counts and percentages. Chi square test and multiple logistic regression was performed to test the hypothesis of the study on $p < 0.05$ for all statistical tests. The variables selected to be included in the multivariate analysis were variables with $p < 0.25$ in the univariate analysis, and gender, age, and type of profession were also included.

Patient and Public Involvement

Patients and the general public were not involved in the conduct of this research. HCW were involved during the pilot study in order to test the understanding of the written questionnaire. In addition, data was assessable by the researchers and is stored in a secure file in the computer and online storage.

Results:

Socio-demographics Characteristics

Responses were collected from 692 HCW (response rate 63.3%), where 311 (44.9%) were doctors and 381 (55.1%) were nurses. Most of the female participants were nurses at 64.5%. More doctors (56.6%) had >5 years of experience at their current place of work, as compared to nurses (40.9%). The average age of participants was 31.96 (Standard Deviation [SD] 7.46) years across the two occupations. Out of the 65.9% who had dependents, i.e., having to take-care of child or elderly persons, 73.9% reported to have support to care for their dependents in a case of a disaster. The percentages were almost equally distributed between doctors and nurses.

Professional and Intrapersonal Characteristics

Table 1 describes the socio-demographic, professional, intrapersonal, and willingness characteristics of respondents. Only 39.2% of the HCW had any previous training in dealing with disasters. This was the same for previous work experience, where only 35.8% of the participants had previous work experience in a disaster situation.

The doctors and nurses reported a high trust in work safety, in family, in colleague's preparedness to react, and in their hospital's preparedness to react in case of a disaster, with trust in colleague's preparedness as the highest percentage (88.3%). Similarly, HCW had a high median score (16; IQR 2) in self-efficacy construct.

Respondents' willingness

Ninety percent of the participants expressed high willingness to participate in any type of disasters. However, they were less willing to participate in natural disasters (77.3%) and influenza pandemic (66.0%) (Table 1).

[Table 1]

Analysis of Self-efficacy's factors

There were four items in the self-efficacy construct. After the data collection, the inter-correlation value of self-efficacy was tested before running the main analysis. The descriptive statistics and inter-item correlation values of the self-efficacy construct are presented in Table 2.

[Table 2]

1
2
3 Based on Table 2, there was a moderate level of agreement in all the four items. The
4
5 highest correlation for each item with at least one other item in the construct was between 0.3
6
7 and 0.9. In factor analysis, the Kaiser-Meyer-Olkin (KMO) value was 0.658 ($p < 0.001$), KMO
8
9 value close to 1, so the variables are suitable for factor analysis. Therefore, a single factor
10
11 was formed from the four items computed, and it was saved as incentive to be used in further
12
13 analysis. Accordingly, to calculate the self- efficacy, four questions was asked. The sum
14
15 outcome of the four answers ranged from 4 to 20 (Median 16, Interquartile range [IQR] 5).
16
17
18
19
20
21

22 **Factors associated with willingness to participate in any type disaster**

23
24
25 There was an association between participants' gender and willingness to participate
26
27 in any type of disaster, with males being more willing compared to females (crude
28
29 OR=2.161, 95%CI: 1.307 to 3.573) (Table 3). Those with high trust in work safety (crude
30
31 OR=3.284, 95%CI: 1.937 to 5.567), trust in colleague's preparedness (crude OR=2.592,
32
33 95%CI: 1.401 to 4.795) and self-efficacy (crude OR=1.358; 95%CI: 1.247 to 1.479) were
34
35 also found to be more willing to participate in any type of disaster in the univariate analysis
36
37 (Table 3).
38
39
40

41 In the final model, having trust in work safety (adjusted OR=2.535, 95%CI: 1.357 to
42
43 4.736) and self-efficacy (adjusted OR=1.319, 95%CI: 1.197 to 1.453) were found to be
44
45 associated with general willingness with any type of disasters (Table 4).
46
47
48
49
50

51 **Factors associated with willingness to participate in natural disaster**

52
53
54 In the univariate analysis, there was an association between some of the socio-
55
56 demographic characteristics (age, gender, education level, type of profession and work
57
58 duration) with willingness to participate in natural disasters. Participants with bachelor and
59
60

1
2
3 postgraduate degrees had a higher odd of willingness to participate in natural disasters
4
5 compared to those with a diploma education. Those in the age group of between 31 and 45
6
7 years were found to be more willing when compared to participants who were ≤ 30 years old.
8
9
10 The main intrapersonal factors associated with willingness in natural disasters were
11
12 participants' trust in colleagues' preparedness in case of a natural disaster, and self-efficacy
13
14 (Table 3).
15

16
17 In multivariate, being male (adjusted OR=1.639, 95%CI: 1.102 to 2.439) and self-
18
19 efficacy (adjusted OR=1.143, 95%CI: 1.069 to 1.221) were significantly associated with
20
21 willingness to participate in natural disasters (Table 4).
22
23
24
25
26
27

28 **Factors associated with willingness to participate in influenza pandemic disaster**

29
30 For willingness to participate in an influenza pandemic, the univariate results revealed
31
32 that being male, having a dependent with no support (compared to participants without a
33
34 dependent), having previous experience and self-efficacy were associated with willingness to
35
36 participate in a pandemic (Table 3).
37
38
39

40 In the final model, having previous experience (adjusted OR=1.528, 95%CI: 1.058 to
41
42 2.207) and self-efficacy (adjusted OR=1.114, 95%CI: 1.050 to 1.182) were found to be
43
44 associated with willingness to participate in influenza pandemic (Table 4).
45
46
47

48 [Table 3, Table 4]
49
50
51
52
53
54
55
56
57
58
59
60

Discussion:

The study's main question is to test the strength and direction of the association between the independent variables- socio-demographic, professional, and intrapersonal variables- with the variables of willingness to participate in three different types of disasters. The results showed that trust in work safety and self-efficacy were associated with disaster participation willingness after in the multivariate analysis. Gender and self-efficacy were found to be significantly associated with willingness to participate in natural disasters. However, previous experience and self-efficacy were statistically significant with willingness of HCW to participate in an influenza pandemic.

It is difficult to obtain a one hundred percent participation of HCW even though it is vital in case of disasters. Previous studies suggested that between 65 and 97 percent of HCW were willing to participate in a natural disaster, and between 54 and 86 percent in an influenza pandemic.^{11,16,20,21} The reason for this difference in the levels of willingness between the two types of disasters is due to the great distinction in their nature. The outcome of the interaction between HCW and the socio-environmental determinants of these disasters leads to having different willingness levels. According to Connor et al., (2014) the weighted risk resulting from this interaction plays a major role in HCW willingness.¹⁰ In the case of pandemics, the fear of the inability to control biohazards and watching colleagues acquiring a communicable disease after contact with affected persons was a suggested reason for the low willingness levels.¹⁰ Another qualitative study in Australia highlighted that in different types of disasters emergency nurses' willingness to attend to work is shaped by the weighted risk to self and surrounding people and the pressure formed from the period dealing with the disaster.¹² Natural disasters may not directly affect HCW or their families like the influenza

1
2
3 pandemic which may be a reason for the higher level of willingness to attend to work
4
5 compared to pandemics.
6
7

8 The findings of this study are consistent with the previous studies regarding their
9
10 willingness level to different types of disasters. More HCW were willing to participate in
11
12 natural disasters (77.3%) as compared to influenza pandemics (66.0%). Nevertheless, even
13
14 though there was a higher percentage of HCW willingness in this study to participate in both
15
16 types of disasters compared to other studies, Yemen's high vulnerability, due its topography
17
18 and current economic, political and health status, could affect the country's health status and
19
20 cause huge adverse health impacts when disasters occur. According to the Vulnerability
21
22 Matrix, one third of Yemen's districts are highly vulnerable.⁵
23
24
25

26
27 Subsequently, due to the armed conflicts in Yemen that started in early 2015, it has
28
29 become difficult to deal with any disaster that may occur. According to the World Health
30
31 Organization (WHO), the healthcare system in Yemen is in a critical situation with about
32
33 50% of the health facilities in Yemen being either partially or totally damaged as a result of
34
35 the natural disasters and conflicts. Furthermore, of the 3,507 healthcare facilities in Yemen,
36
37 almost 300 healthcare facilities have been destroyed.^{5,22} Therefore, any possible decrease in
38
39 the health work force during disasters in Yemen must be considered. The existing drained
40
41 health sector is based on a vulnerable health system in terms of its structure and healthcare
42
43 staff number and distribution in the country.²³ Thus, training and preparing healthcare staff to
44
45 act during disasters is critical and could help achieve Priority 4 in the United Nations' Sendai
46
47 Framework for Disaster Risk Reduction 2015-2030 in enhancing disaster preparedness for
48
49 effective response and to "Build Back Better" in recovery, rehabilitation and reconstruction
50
51 of disasters, and further prevent any possible humanitarian catastrophe.²⁴
52
53
54
55
56

57 The results show that willingness to participate in events of natural disasters may be
58
59 encouraged by many suggested factors. Factors such as respondents being male are
60

1
2
3 accompanied with higher willingness to participate in natural disasters. The findings are
4 similar to previous studies conducted in Jordan, China and the United States of
5 America.^{15,16,25,26} However, having less female health workers who are willing to participate
6 in disasters highlight a need to address this group during health preparedness programs. This
7 lack in female HCW willingness could be due to the gender and cultural barriers in Yemen.
8 These barriers need to be addressed to better motivate female workers participation in
9 disasters. This is an important area for research as studies on willingness are moving forward.

10
11
12
13
14
15
16
17
18
19
20 As noted in previous studies, nurses and doctors with previous experience in influenza
21 pandemics are ever willing to participate in influenza pandemics.^{14, 25} HCW who have
22 experience working with previous disaster have a better knowledge in disaster management,
23 coping strategies, and ways to protect themselves and their families; which may be a reason
24 for having higher odds of willingness to participate. According to Alzahrani et al., (2017)
25 most of the nurses have a higher level in Mass gatherings management such as
26 communicating effectively during emergencies in Saudi Arabia due to Previous experience.²⁷
27 He also found a need for further trainings by the nurses. Similarly, Chokshi et al., (2008)
28 found that simulation training, attending conferences and previous experience increase
29 paediatric surgeons' feeling of preparedness.¹⁴

30
31
32
33
34
35
36
37
38
39
40
41
42
43 This study found that the trust in work safety during any type of disaster plays a key
44 role in the willingness to participate, which is similar to the results from previous
45 studies.^{11,25,28} Work safety is important in order for HCW to feel safe and do their work. A
46 study by Stergachis et al., (2011) found that the majority of the participants reported their fear
47 or concern for self in case of influenza pandemic and during earthquake scenarios as one of
48 the major barriers to willingness to participate.²⁹ In Yemen, work safety in disaster
49 preparedness is concerning after notifications made by Aladhrai et al.,(2015) and Naser et
50 al., (2018) on the current low work place safety and the needs of increasing safety and
51
52
53
54
55
56
57
58
59
60

1
2
3 security standards in Yemen's Health establishments.^{7,8} Whether it is external or internal
4 catastrophe, unsafe work place could lead to increase optional absenteeism during disaster.
5
6 Thus, disseminating information on work safety and protection devices during several types
7
8 of disasters is advisable.
9
10

11
12
13 Many studies have established the association between self-efficacy and willingness
14 to participate in disasters.^{10,15,16} Similarly, this study has identified that self-efficacy plays an
15 important role in the willingness to participate in natural disasters, in influenza pandemic and
16 any type of disasters. It indicates that elements tested for self-efficacy like participants
17 increase in familiarity with their role, responsibility to react, and being confident of their
18 ability to deal with various types of disasters could be a key factor to increase HCW self-
19 efficacy. Thus, their willingness to participate will also increase. Disaster preparedness
20 trainings are encouraged to contain materials that explain the responses to the different types
21 of disasters, doctors' and nurses' role in disasters and how it makes a difference in
22 responding to disasters. Supportive measures to increase HCW self-efficacy such as
23 immediate communication with much needed information and rewards for efforts in disasters
24 are suggested.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Although the study faced challenges in the collection of data during the political
42 unrest in the country, data was collected from three of the major public hospitals. The quality
43 of data was ensured by researcher explaining the study and its objectives and answering any
44 questions that may arise. This prevented differences in understanding the questions and
45 increased participants' inclusion in the study. All in all, the study suggested the vital value of
46 increasing self-efficacy and its elements in order to obtain more willingness participants.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations

This is the first study to explore the patterns and associated factors of the willingness status to participate in disasters among healthcare workers in Yemen. The response rate was also high as the questionnaire was distributed personally by the researcher. The questionnaire was pilot tested to ensure the validity of the questionnaire.

This study had some limitations. As the study was performed under a political insecurity/fragile state of the country and a list of workers of HCW working at the hospitals was difficult to obtain. As a result, a universal sample was undertaken which could limit the study of representativeness. This is a cross-sectional study using a self-administered questionnaire, where the actual willingness of the respondents cannot be ascertained. The respondents may answer positively due to social desirability bias. Other than the self-administrated questionnaire limitations, the findings of the study are limited to staff working at the tertiary level public hospitals in urban areas of Yemen.

Conclusion:

Increasing the likelihood of willingness to participate in disasters play a key role in guarantying the optimum number of work force. This study indicates that one's socio-demographic, professional and intrapersonal factors play a role in increasing his/her willingness in general and across different types of disasters. Significant differences were revealed between participants' willingness in natural disasters by gender and self-efficacy, and participant's willingness in influenza pandemics by previous experience and self-efficacy. This result suggests integrating disaster management to doctors' and nurses' in the early stages of their educational curriculums. This could be achieved by adding early

1
2
3 exposure of healthcare workers to relevant disaster experiences which would further boost
4
5 their willingness to participate in disaster response.
6
7

8 Other areas for preparedness may include increasing hospitals safety and resilience;
9
10 hence that willingness to participate in natural disasters' response was influenced by the
11
12 demographic characteristic of the healthcare personnel. Additionally, others found outside the
13
14 bracket should be motivated to participate with a reward package such as incentives and
15
16 hazard allowance. Further studies should also be conducted in both urban and rural settings
17
18 with a relatively peaceful atmosphere.
19
20
21
22
23
24

25 **Acknowledgement:** We thank Dr. Susan Connor, Dr. Karen Hammad and Dr.
26
27 Mohammed Khalil for their support and advice during the study development, all of the
28
29 University Malaya academic staff, the team who collected the data, and all the participants
30
31 who contributed in this study.
32
33
34

35 **Competing interests:** None declared
36
37

38 **Funding:** The authors have no financial disclosures to declare. This research did not
39
40 receive any specific grant from funding agencies in the public, commercial, or not-for-profit
41
42 sectors.
43
44
45

46 **Data sharing statement:** This is an anonymous gathered survey; the data are the
47
48 responses from the participants. The data have been grouped and presented in the manuscript.
49
50
51
52
53
54
55
56
57
58
59
60

Reference:

1. United Nations Economic and Social Commission for Asia and the Pacific. Disasters in Asia and the Pacific: 2015 year in review. United Nations report. Economic and social commission for Asia and the Pacific. 2015. [21 p.]. Available from: <https://www.unescap.org/resources/disasters-asia-and-pacific-2015-year-review>
2. Below R, Wirtz A, Guha-Sapir D. Disaster category classification and peril terminology for operational purposes. Centre for Research on the Epidemiology of Disasters. 2009. Available from: <https://www.cred.be/node/564>
3. Joseph, B., & Joseph, M. (2016). The health of the healthcare workers. *Indian journal of occupational and environmental medicine*, 20(2), 71–72. doi:10.4103/0019-5278.197518.
4. Ministry of Water and Environment. Yemen: National progress report on the implementation of the Hyogo Framework for Action (2013-2015). 2015. Available from: <http://preventionweb.net/go/40141>
5. World Health Organization. WHO annual report 2017: Yemen. Cairo: WHO Regional Office for the Eastern Mediterranean; 2018. Licence: CC BYNC-SA 3.0 IGO.
6. Banerjee, Aditi; Bhavnani, Rakhi; Burtonboy, Catherine H, et al. Natural Disasters in the Middle East and North Africa: A Regional Overview. The International Bank for Reconstruction and Development / The World Bank. [101 p.]. 2014; Report No.: 81658.
7. Aladhrai SA, Djalali A, Della Corte F, Alsabri M, El-Bakri NK, Ingrassia PL. Impact of the 2011 Revolution on Hospital Disaster Preparedness in Yemen. *Disaster Medicine and Public Health Preparedness*. 2015;9(4):396-402.
8. Naser WN, Ingrassia PL, Aladhrae S, Abdulraheem WA. A Study of Hospital Disaster Preparedness in South Yemen. *Prehospital and Disaster Medicine*. 2018;33(2):133.
9. Chaffee M. Willingness of health care personnel to work in a disaster: an integrative review of the literature. *Disaster Medicine and Public Health Preparedness*. 2009;3(1):42-56.

10. Connor SB. When and why health care personnel respond to a disaster: the state of the science. *Prehospital and Disaster Medicine*. 2014;29(3):270-4.
11. Gowing JR, Walker KN, Elmer SL, Cummings EA. Disaster Preparedness among Health Professionals and Support Staff: What is Effective? An Integrative Literature Review. *Prehospital and Disaster Medicine*. 2017;32(3):321-8.
12. Arbon P, Cusack L, Ranse J, Shaban RZ, Considine J, Kako M, et al. Exploring staff willingness to attend work during a disaster: a study of nurses employed in four Australian emergency departments. *Australasian Emergency Nursing Journal : AENJ*. 2013;16(3):103-9.
13. Ejeta LT, Ardalan A, Paton D. Application of Behavioral Theories to Disaster and Emergency Health Preparedness: A Systematic Review. *PLOS Currents Disasters*. 2015;7. Doi: 10.1371/currents.dis.31a8995ced321301466db400f1357829.
14. Chokshi NK, Behar S, Nager AL, Dorey F, Upperman JS. Disaster management among pediatric surgeons: preparedness, training and involvement. *American Journal of Disaster Medicine*. 2008;3(1):5-14.
15. Burke RV, Goodhue CJ, Chokshi NK, Upperman JS. Factors associated with willingness to respond to a disaster: a study of healthcare workers in a tertiary setting. *Prehospital and Disaster Medicine*. 2011;26(4):244-50.
16. Charney RL, Rebmann T, Flood RG. Hospital Employee Willingness to Work during Earthquakes Versus Pandemics. *Journal of Emergency Medicine*. 2015;49(5):665-74. Doi: <https://doi.org/10.1016/j.jemermed.2015.07.030>.
17. Balicer RD, Omer SB, Barnett DJ, Everly GS, Jr. Local public health workers' perceptions toward responding to an influenza pandemic. *BMC Public Health*. 2006;6:99.
18. Shapira S, Aharonson-Daniel L, Bar-Dayyan Y, Sykes D, Adini B. Knowledge, perceptions, attitudes and willingness to report to work in an earthquake: A pilot study comparing Canadian versus Israeli hospital nursing staff. *International Emergency Nursing*. 2016;25:7-12.
19. Bandura A. *Self-efficacy: The exercise of control*: Macmillan; 1997.
20. Arbon P, Ranse J, Cusack L, Considine J, Shaban RZ, Woodman RJ, et al. Australasian emergency nurses' willingness to attend work in a disaster: a survey. *Australasian Emergency Nursing Journal : AENJ*. 2013;16(2):52-7.

- 1
2
3 21. Brice JH, Gregg D, Sawyer D, Cyr JM. Survey of Hospital Employees' Personal
4 Preparedness and Willingness to Work Following a Disaster. *Southern Medical*
5 *Journal*. 2017;110(8):516-22.
6
7
- 8 22. United Nations Office for the Coordination of Humanitarian Affairs. *Humanitarian*
9 *Bulletin Yemen/ Health system on the verge of collapse*. 2015. [4 p.]. Available
10 from: [http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-](http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-30-november-2015-enar)
11 [30-november-2015-enar](http://reliefweb.int/report/yemen/yemen-humanitarian-bulletin-issue-6-issued-30-november-2015-enar)
12
13
- 14 23. Qirbi N, Ismail SA. Health system functionality in a low-income country in the
15 midst of conflict: the case of Yemen. *Health Policy and Planning*. 2017;32(6):911-
16 22.
17
18
- 19 24. United Nations - Headquarters. *Sendai Framework for Disaster Risk Reduction*
20 *2015-2030*. United Nations Office for Disaster Risk Reduction. 2015. [32 p.].
21 Available from: <https://www.unisdr.org/we/inform/publications/43291>
22
23
- 24 25. Loke AY, Fung WM, Liu X. Mainland China nurses' willingness to report to work
25 in a disaster. *American Journal of Disaster Medicine*. 2013;8(4):273-82. Doi:
26 [10.5055/ajdm.2013.0134](https://doi.org/10.5055/ajdm.2013.0134).
27
28
- 29 26. Al Khalaileh MA, Bond E, Alasad JA. Jordanian nurses' perceptions of their
30 preparedness for disaster management. *International emergency nursing*.
31 2012;20(1):14-23.
32
33
- 34 27. Alzahrani F, Yiannis K. Emergency nurse disaster preparedness during mass
35 gatherings: a cross-sectional survey of emergency nurses' perceptions in hospitals in
36 Mecca, Saudi Arabia. *BMJ Open*. 2017;7(4):e013563.
37
38
- 39 28. Ogedegbe C, Nyirenda T, Delmoro G, Yamin E, Feldman J. Health care workers
40 and disaster preparedness: barriers to and facilitators of willingness to respond.
41 *International Journal of Emergency Medicine*. 2012;5(1):29.
42
43
- 44 29. Stergachis A, Garberson L, Lien O, D'Ambrosio L, Sangare L, Dold C. Health care
45 workers' ability and willingness to report to work during public health emergencies.
46 *Disaster Medicine and Public Health Preparedness*. 2011;5(4):300-8.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Socio-demographic, professional, intrapersonal, and willingness**Characteristics of Respondents**

Variable		Frequency (n= 692)	Percentage (%)
Age (years)	≤30	360	52.0
	31-45	294	42.5
	≥46	38	5.5
Gender	Male	419	60.5
	Female	273	39.5
Marital-status	Single	258	37.3
	Married	420	60.7
	Divorce	8	1.2
	Widow	6	0.9
Education-level	Diploma	280	40.5
	Bachelor	275	39.7
	Post graduate education	137	19.8
	Master	104	15.0
Professional	33	4.8	
Profession type	Doctors	311	44.9
	Specialist medical practitioner	126	8.2
	General medical practitioner	185	26.7
	Nurses	381	55.1
Work duration (years)	≤5	360	52
	6-10	213	30.8
	11-15	76	11.0
	≥16	43	6.2
Dependent	With dependent	456	65.9
	Elder	65	9.4
	Child	276	39.9
	Both	115	16.6
	No dependent	236	34.1
^a Support	No	119	26.1
	Yes	337	73.9
^b Previous training	With previous training	271	39.2
	Without previous training	421	60.8
^b Previous experience	With previous experience	248	35.8
	Without previous training	444	64.2
Trust in work safety in case of disaster	High trust in work safety	563	81.4
	Low trust in work safety	129	18.6
Trust in family preparedness in case of a disaster	High trust in family preparedness	544	78.6
	Low trust in family preparedness	148	21.4
Trust in colleague preparedness to react in disaster	High trust in colleague preparedness	611	88.3
	Low trust in colleague preparedness	81	11.7

Trust in hospital preparedness to react in disaster	High trust in hospital preparedness	522	75.4
	Low trust in hospital preparedness	170	24.6
Self-efficacy score (Median; Inter-Quartile)		16	5
Willingness to participate in Any type of disaster	High	623	90.0
	Low	69	10.0
Willingness to participate in Natural disaster	High	535	77.3
	Low	157	22.7
Willingness to participate in Influenza and Pandemic	High	457	66.0
	Low	235	34.0

^aOnly for participants with dependants. The percentages are only within participants with dependents.

^bPrevious experience and training in disasters

Table 2: Descriptive statistics and inter-term correlation for items in self-efficacy

Items	Descriptive statistics		Inter-item correlation			
	Mean	SD	item 1	item 2	item 3	item 4
Item 1: Ability to perform work	3.36	1.074	1.000	0.592	0.267	0.259
Item 2: Familiarity with their role	3.62	1.085	0.592	1.000	0.325	0.324
Item 3: Responsibility to participate	4.23	0.929	0.267	0.325	1.000	0.390
Item 4: Ability report to work	4.03	1.062	0.259	0.324	0.390	1.000

For peer review only

Table 3: Univariate association of crude socio-demographic, professional and intrapersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters		Willingness to participate in natural disasters		Willingness to participate in influenza pandemic	
	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value
Age						
≤30	1	1	1	1	1	1
31-45	1.056 (0.628-1.775)	0.838	1.639 (1.126-2.387)	0.010	1.138 (0.822-1.576)	0.437
≥46	0.733 (0.269-1.997)	0.544	1.967 (0.798-4.850)	0.142	1.354 (0.650-2.820)	0.418
Gender						
Female	1	1	1	1	1	1
Male	2.161 (1.307-3.573)	0.003	2.254 (1.571-3.234)	<0.001	1.505 (1.093-2.072)	0.012
Marital status						
Single	1	1	1	1	1	1
Married	0.899 (0.532-1.521)	0.692	1.128 (0.781-1.629)	0.521	1.090 (0.787-1.510)	0.605
Divorce and widow	0.615 (0.130-2.913)	0.541	1.160 (0.314-4.291)	0.824	1.362 (0.416-4.466)	0.610
Education Level						
Up to Diploma	1	1	1	1	1	1
Up to Bachelor	1.380 (0.775-2.458)	0.274	1.962 (1.317-2.922)	0.001	1.221 (0.858-1.738)	0.268
Up to Postgraduate	0.847 (0.450-1.596)	0.608	2.279 (1.351-3.842)	0.002	0.998 (0.651-1.528)	0.992
Profession type						
Nurses	1	1	1	1	1	1
General medical practitioner	1.473 (0.781-2.776)	0.231	1.554 (1.013-2.384)	0.044	0.756 (0.524-1.091)	0.135
Specialist medical practitioner	0.965 (0.507-1.835)	0.913	2.407 (1.377-4.208)	0.002	0.942 (0.614-1.446)	0.785
Work duration						

≤5	1	1	1	1	1	1	1
6-10	1.006 (0.563-1.798)	0.983	1.079 (0.721-1.614)	0.711	0.975 (0.683-1.392)	0.889	
11-15	1.028 (0.438-2.415)	0.949	2.337 (1.118-4.882)	0.024	1.059 (0.626-1.791)	0.832	
≥16	0.456 (0.196-1.063)	0.069	0.650 (0.329-1.286)	0.216	1.198 (0.603-2.379)	0.606	
Presence of dependent							
No dependent	1	1	1	1	1	1	
Dependent with no support	0.673 (0.346-1.308)	0.242	1.149 (0.683-1.933)	0.601	1.617 (1.000-2.612)	0.050	
Dependent with support	1.253 (0.707-2.222)	0.439	1.305 (0.880-1.935)	0.185	1.283 (0.907-1.815)	0.158	
^aPrevious training							
No	1	1	1	1	1	1	
Yes	1.653 (0.959-2.849)	0.070	1.090 (0.756-1.573)	0.644	1.147 (0.829-1.585)	0.408	
^aPrevious experience							
No	1	1	1	1	1	1	
Yes	1.216 (0.714-2.070)	0.471	0.974 (0.673-1.411)	0.889	1.857 (1.317-2.619)	<0.001	
Trust in work safety							
Low	1	1	1	1	1	1	
High	3.284 (1.937-5.567)	<0.001	1.280 (0.825-1.987)	0.271	1.192 (0.800-1.774)	0.388	
Trust in family preparedness							
Low	1	1	1	1	1	1	
High	1.338 (0.756-2.369)	0.317	1.292 (0.850-1.963)	0.231	0.990 (0.674-1.454)	0.959	
Trust in colleague's preparedness							
Low	1	1	1	1	1	1	
High	2.592 (1.401-4.795)	0.002	1.974 (1.200-3.246)	0.007	1.392 (0.866-2.237)	0.172	
Trust in hospital preparedness							
Low	1	1	1	1	1	1	
High	1.392 (0.807-2.400)	0.234	1.212 (0.810-1.814)	0.351	1.158 (0.807-1.663)	0.426	
^bSelf-efficacy							
	1.358 (1.247-1.479)	<0.001	1.184 (1.115-1.257)	<0.001	1.135 (1.076-1.197)	<0.001	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

^aPrevious experience and training in disasters

^b*Continuous measure, with one unit increase in self-efficacy*

For peer review only

Table 4: Multivariate association of adjusted professional and intrapersonal characteristics with willingness in any type of disaster, in natural disasters, or in influenza pandemic

Variable	Willingness to participate in any type of disasters		Willingness to participate in natural disasters		Willingness to participate in influenza pandemic	
	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value	Odds ratio (95% C.I.)	p-value
Age						
≤30	1	1	1	1	1	1
31-45	1.137 (0.609-2.121)	0.687	1.432 (0.928-2.209)	0.105	1.008 (0.695-1.461)	0.967
≥46	0.440 (0.136-1.418)	0.169	1.087 (0.409-2.886)	0.867	1.062 (0.484-2.332)	0.880
Gender						
Female	1	1	1	1	1	1
Male	1.456 (0.807-2.628)	0.212	1.639 (1.102-2.439)	0.015	1.131 (0.793-1.612)	0.498
Education Level						
Up to Diploma	-	-	1	1	-	-
Up to Bachelor	-	-	1.706 (0.993-2.932)	0.053	-	-
Up to Postgraduate	-	-	1.177 (0.522-2.657)	0.695	-	-
Profession type						
Nurses	1	1	1	1	1	1
General medical practitioner	1.285 (0.438-3.767)	0.261	1.581 (0.691-3.614)	0.278	0.751 (0.468-1.205)	0.235
Specialist medical practitioner	1.392 (0.578-3.352)	0.644	0.939 (0.515-1.713)	0.838	0.697 (0.473-1.026)	0.067
Presence of dependent						
No dependent	-	-	-	-	1	1
Dependent with no support	-	-	-	-	1.537 (0.910-2.598)	0.108
Dependent with support	-	-	-	-	1.154 (0.791-1.685)	0.457

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

^aPrevious experience						
No	-	-	-	-	1	1
Yes	-	-	-	-	1.528 (1.058-2.207)	0.024
Trust in work safety						
Low	1	1	-	-	-	-
High	2.535 (1.357-4.736)	0.004	-	-	-	-
Trust in colleague's preparedness						
Low	1	1	1	1	-	-
High	1.199 (0.576-2.496)	0.686	1.363 (0.791-2.351)	0.265	-	-
^bSelf-efficacy	1.319 (1.197-1.453)	<0.001	1.143 (1.069-1.221)	<0.001	1.114 (1.050-1.182)	<0.001
^a Previous experience and training in disasters						
^b Continuous measure, with one unit increase in self-efficacy						

Peer review only

SECTION ONE:

Please tick (✓) in the appropriate box

1. **Age:** _____

2. **Gender :**

Male

Female

3. **Marital status:**

Single

Married

Divorce

Widow/er

4. **Highest Education level completed :**

Diploma

Bachelor

Masters

Professional

5. **What is your primary job category?**

Specialist medical practitioner

General medical practitioner

Nurse

6. **Duration of work at the hospital:** _____ Years.

7. Do you have any of the following responsibilities (you can choose more than one choice):

Children Elders None

8. Do you have any one who can take care of the previously mentioned responsibilities in case of your absence?

Yes No

SECTION TWO

Please tick (✓) in the appropriate box:

1. Have you participated in a disaster before?

Yes No

2. Have attended any previous disaster trainings?

Yes No

SECTION THREE:

Part A:

NO	Questions	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am confident of my personal safety at work in case of a disaster.					

2.	I am assured that my family is prepared to function in in my absence during a disaster.					
3.	I am sure that my colleagues are able to perform their duties during a disaster.					
4.	The hospital is prepared to provide effective response in case of a disaster.					

PART B:

NO	Statement	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am able to treat patients of different type of disasters.					
2.	I am confident that I can perform my role in the hospital following any type of disasters.					
3.	I feel that it is my duty to work in the event of a disaster.					
4.	I will be able to report to work at the hospital during an event of a disaster.					

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

SECTION FOUR:

NO	Statement	Strongly disagree (1)	Disagree (2)	Somewhat agree (3)	Agree (4)	Strongly agree (5)
1.	I am willing to participate in any type of disaster regardless of its severity					
2.	I am willing to participate in natural disasters (earthquake, floods or cyclone)					
3.	I am you willing to participate in influenza pandemic					

Reporting checklist for cross sectional study.

Based on the STROBE cross sectional guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cross sectional reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

		Reporting Item	Page Number
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	Main document (1)
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	Main document (1)
Background /	#2	Explain the scientific background and rationale for the	Main document

1	rationale		investigation being reported	(3)
2				
3				
4	Objectives	#3	State specific objectives, including any prespecified	Main document
5			hypotheses	(4)
6				
7				
8				
9	Study design	#4	Present key elements of study design early in the paper	Main document
10				(4)
11				
12				
13				
14	Setting	#5	Describe the setting, locations, and relevant dates,	Main document
15			including periods of recruitment, exposure, follow-up,	(4-6)
16			and data collection	
17				
18				
19				
20				
21				
22	Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods	Main document
23			of selection of participants.	(4)
24				
25				
26				
27		#7	Clearly define all outcomes, exposures, predictors,	Main document
28			potential confounders, and effect modifiers. Give	(5), table1
29			diagnostic criteria, if applicable	(Page 18)
30				
31				
32				
33				
34				
35	Data sources /	#8	For each variable of interest give sources of data and	Main document
36	measurement		details of methods of assessment (measurement).	(5)
37			Describe comparability of assessment methods if there	
38			is more than one group. Give information separately for	
39			exposed and unexposed groups if applicable.	
40				
41				
42				
43				
44				
45				
46				
47	Bias	#9	Describe any efforts to address potential sources of bias	Main document
48				(13)
49				
50				
51				
52	Study size	#10	Explain how the study size was arrived at	Main document
53				(4,5)
54				
55				
56				
57				
58	Quantitative	#11	Explain how quantitative variables were handled in the	Main document
59				
60				

1	variables	analyses. If applicable, describe which groupings were	(6)
2			
3		chosen, and why	
4			
5			
6	Statistical	#12a Describe all statistical methods, including those used to	Main document
7			
8	methods	control for confounding	(6)
9			
10			
11		#12b Describe any methods used to examine subgroups and	Main document
12			
13		interactions	(6)
14			
15			
16		#12c Explain how missing data were addressed	Main document
17			
18			(6)
19			
20			
21			
22		#12d If applicable, describe analytical methods taking account	Main document
23			
24		of sampling strategy	(4, 6)
25			
26			
27		#12e Describe any sensitivity analyses	
28			
29			
30	Participants	#13a Report numbers of individuals at each stage of study—	Main document
31			
32		eg numbers potentially eligible, examined for eligibility,	(4, 6)
33			
34		confirmed eligible, included in the study, completing	
35			
36		follow-up, and analysed. Give information separately for	
37			
38		for exposed and unexposed groups if applicable.	
39			
40			
41			
42		#13b Give reasons for non-participation at each stage	Main document
43			
44			(4, 6, 13)
45			
46			
47			
48		#13c Consider use of a flow diagram	
49			
50			
51	Descriptive data	#14a Give characteristics of study participants (eg	Main document
52			
53		demographic, clinical, social) and information on	(7,8), table 2
54			
55		exposures and potential confounders. Give information	(page 19, 20)
56			
57		separately for exposed and unexposed groups if	
58			
59			
60			

applicable.

1			
2			
3			
4		#14b	Indicate number of participants with missing data for
5			each variable of interest
6			Table 2 (page
7			19, 20)
8			
9	Outcome data	#15	Report numbers of outcome events or summary
10			Main document
11			(7), table 2
12			
13			unexposed groups if applicable.
14			(page 19, 20)
15			
16	Main results	#16a	Give unadjusted estimates and, if applicable,
17			Main document
18			(8, 9), table 3,
19			95% confidence interval). Make clear which confounders
20			4 (page 21-23)
21			
22			were adjusted for and why they were included
23			
24			
25			
26		#16b	Report category boundaries when continuous variables
27			Main document
28			
29			were categorized
30			5
31			
32		#16c	If relevant, consider translating estimates of relative risk
33			
34			into absolute risk for a meaningful time period
35			
36			
37	Other analyses	#17	Report other analyses done—e.g., analyses of
38			subgroups and interactions, and sensitivity analyses
39			
40			
41			
42	Key results	#18	Summarise key results with reference to study objectives
43			Main document
44			(6-9)
45			
46			
47			
48	Limitations	#19	Discuss limitations of the study, taking into account
49			Main document
50			(12, 13)
51			
52			sources of potential bias or imprecision. Discuss both
53			direction and magnitude of any potential bias.
54			
55	Interpretation	#20	Give a cautious overall interpretation considering
56			Main document
57			(9-12)
58			objectives, limitations, multiplicity of analyses, results
59			
60			

1 from similar studies, and other relevant evidence.

2
3
4 Generalisability [#21](#) Discuss the generalisability (external validity) of the Main document
5
6 study results (13)

7
8
9 Funding [#22](#) Give the source of funding and the role of the funders for Main document
10
11 the present study and, if applicable, for the original study (14)
12
13 on which the present article is based

14
15
16 The STROBE checklist is distributed under the terms of the Creative Commons Attribution License
17
18 CC-BY. This checklist can be completed online using <https://www.goodreports.org/>, a tool made by
19
20 the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)
21
22