



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

"It's like a phone call to your brain..." Understanding how children communicate their concept of pain: a qualitative study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-033199
Article Type:	Research
Date Submitted by the Author:	25-Jul-2019
Complete List of Authors:	Pate, Joshua; Macquarie University, Health Professions Noblet, Tim; Macquarie University, Health Professions Hush, Julia ; Macquarie University, Health Professions Hancock, Mark; Macquarie University, Health Professions Sandells, Renee; Children's Hospital at Westmead Pounder, Meg; Children's Hospital at Westmead Pacey, Verity; Macquarie University Faculty of Medicine and Health Sciences
Keywords:	concept of pain, pain education, paediatric pain, qualitative interviews, drawing task

SCHOLARONE™
Manuscripts

Title:

“It’s like a phone call to your brain...” Understanding how children communicate their concept of pain: a qualitative study.

Authors:

Joshua W. Pate^{1,2}, MRes, Senior Physiotherapist and PhD Candidate, joshua.pate@mq.edu.au

Tim Noblet¹, MSc (MSKMed), Senior Physiotherapist and PhD Candidate,
tim.noblet@mq.edu.au

Julia M. Hush¹, PhD, Professor, julia.hush@mq.edu.au

Mark J. Hancock¹, PhD, Professor, mark.hancock@mq.edu.au

Renee Sandells², MPsych (Clin), Clinical Psychologist, renee.sandells@health.nsw.gov.au

Meg Pounder², BSc (Physiotherapy), Senior Physiotherapist, meg.pounder@health.nsw.gov.au

Verity Pacey^{1,2}, PhD, Senior Lecturer, verity.pacey@mq.edu.au

Affiliations:

¹ Macquarie University, New South Wales, Australia

² The Children’s Hospital at Westmead, New South Wales, Australia

Corresponding Author:

Joshua Pate

Ground floor 75 Talavera Road

Macquarie University 2109

Phone: +61 2 9850 6626

Fax: +61 2 9850 6630

Email: joshua.pate@mq.edu.au

Institutional URL: <https://researchers.mq.edu.au/en/persons/joshua-pate>

Keywords:

concept of pain; pain education; paediatric pain; qualitative interviews; drawing task

Previous presentations of the research:

Poster of one subtheme of the manuscript presented for IASP 2018 in Boston, titled: “Explaining pain to children with and without pain: what terminology should we use? Qualitative interviews describing pain-related neuroanatomy.”

Acknowledgments:

We thank all the children and their families for their openness and for giving their valuable time, energy and efforts to participate in this study. We also acknowledge Natasha Haynes, Nurse Practitioner from The Children’s Hospital at Westmead Pain Team, who helped to recruit participants and supported the study.

Funding:

J.W.P. is a Ph.D. Student funded by a Macquarie University Research Training Program (RTP) full-time scholarship.

Conflicts of Interest:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

For peer review only

Abstract:

Objective: A person's concept of pain can be defined as how they understand what pain actually is, what function it serves, and what biological processes are thought to underpin it. This study aimed to explore the concept of pain in children with and without persistent pain.

Design: In-depth, face-to-face interviews with drawing tasks were conducted with 16 children (aged 8-12 years) in New South Wales, Australia. Thematic analysis was used to analyse and synthesize the data.

Setting: Children with persistent pain were identified from a pain clinic waiting list in Australia, and children without pain were identified through advertising flyers and email bulletins at a university and hospital.

Participants: Eight children had persistent pain and eight children were pain-free.

Results: Four themes emerged from the data: 'my pain-related knowledge', 'pain in the world around me', 'pain in me', and 'communicating my concept of pain'. A conceptual framework of the potential interactions between the themes resulting from the analysis is proposed. The concept of pain of Australian children aged 8-12 years varied depending on their knowledge, experiences and literacy levels. For example, when undertaking a drawing task, most children with persistent pain drew emotional elements to describe pain, whereas most children who were pain-free did not.

Conclusions: Gaining an in-depth understanding of a child's previous pain-related experiences and knowledge is important to facilitate clear and meaningful pain education. The use of age-appropriate language, in combination with novel assessment and education tasks such as drawing and discussing vignettes, allowed children to communicate their individual concept of pain.

1

2

3 **Article Summary:**

4

5 Strengths and limitations of this study:

6

- 7
- 8 • One strength of this study was the use of novel data collection methods, such as drawing
 - 9 tasks and vignettes, as a means of data triangulation.
 - 10
 - 11 • Another strength was the large variations in pain experience amongst the group of 8 to
 - 12 12-year old children with persistent pain; some children had lifelong permanent physical
 - 13 disabilities whereas others had recurrent headaches for just over one year, capturing
 - 14 individual variability in pain experiences.
 - 15
 - 16 • A limitation of this study is the potential lack of transferability. This study only included
 - 17 children from New South Wales, Australia who spoke English. It is unknown if the
 - 18 results would be transferable to other cultures and age groups.
 - 19
 - 20 • A further limitation is that other ‘raindrops’ in the proposed conceptual framework could
 - 21 exist that were not identified and explored in these interviews.
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60

Introduction:

Estimates of point-prevalence of persistent pain in children range from 11% to 54%,^{1 2} which is comparable to that for adults.³ Persistent pain in children has detrimental effects on a child's quality of life⁴ and is a significant socioeconomic and health problem. For example, persistent pain in childhood can predict persistent pain in adulthood⁵ and is associated with reduced pain-related disability,¹ school absence,⁶ poorer reading levels,⁷ and emotional distress.⁸ An increasing number of children are being admitted to hospitals for treatment of persistent pain⁹ and health service costs are high.¹⁰

Multidisciplinary pain management is currently the gold standard for persistent pain management.¹¹ An important part of the multidisciplinary approach is pain education, aiming to change a person's concept of pain and to improve pain and function. A person's concept of pain can be defined as how they understand "what pain actually is, what function it serves, and what biological processes are thought to underpin it";¹² these can be summarized as the 'what, why and how of pain'. Clinically, a range of methods for educational assessments and treatments are used for people with persistent pain including drawing tasks,¹³ which are a robust interview strategy with children.¹⁴ However, these methods have not been applied to research to investigate a child's concept of pain.

Limited aspects of a child's experience of pain have been investigated in previous research. For example, children who are pain-free have been asked to describe sensations relating to a grazed knee, an injection, and a headache.¹⁵ In addition, pain-specific behaviours, pain quality, and pain intensity have been investigated in children with pain.¹⁶ However, a child's concept of pain, as

defined in the previous paragraph, has not been investigated in children with or without persistent pain. Therefore, this remains poorly understood, providing clinicians little guidance on how to target age-appropriate pain education to children with persistent pain as part of multidisciplinary management, or to pain-free children as a strategy to prevent future pain. The aims of this study were (1) to explore the concept of pain of children diagnosed with persistent pain, and (2) to compare the concept of pain between children who have pain with those who are pain-free.

Methods:

The study was approved by the Human Research Ethics Committees of Sydney Children’s Hospital Network (Ref: LNR/17/SCHN/268) and Macquarie University (Ref: 5201700820). All participants’ parents provided informed consent prior to participation in the study.

Study Design

To ensure explicit and comprehensive study reporting, the widely used Consolidated Criteria for Reporting Qualitative Health Research (COREQ)¹⁷ was implemented.

The study utilized 30-minute one-to-one qualitative interviews of children aged 8-12 years. Children were asked to answer questions and complete activities relating to their concept of pain. By adapting grounded theory and utilizing thematic analysis,^{18 19} concepts in the transcripts were inductively identified and recorded and then a coding structure was iteratively developed and refined until the aims of the study were achieved. Themes and sub-themes were developed from data analysis and synthesis.

Participants

Two groups of children aged 8-12 years in the state of New South Wales, Australia, were purposefully recruited; children who had persistent pain (greater than 3 months) and children who were pain-free and healthy by parent-report. Children were excluded from the study if they (1) had received pain education from a health care professional, (2) were deemed medically unsuitable by their treating physician/surgeon, or (3) were unable to participate in a 30-minute interview based on parent-report.

A minimum age for participants was set at 8 years, the youngest age a child can be meaningfully interviewed.²⁰ The maximum age of participation was 12 years, an age prior to when abstract thinking capacities typically develop in the teenage years.²¹ The age range was determined to meet a clinical need highlighted by research team members to provide improved assessment and management resources for pre-teenage children. A pragmatic approach incorporating age limits, rather than a formal assessment of cognitive development, allows for direct applicability in time-limited clinical settings.

A purposive selection strategy was used to identify children of varying demographics and pain characteristics who were likely to meet the eligibility criteria. For all children, a mix of gender, age, and ethnic backgrounds was sought to capture diversity of experiences. For the children with persistent pain, purposive sampling attended to capturing variations in the duration and location of pain. Children were identified from The Children's Hospital at Westmead, Australia

Complex Pain Service waiting list, and through advertising flyers and email bulletins at Macquarie University and The Children’s Hospital at Westmead.

Children were interviewed until data saturation was reached.

Interview Script

An interview script was developed (Table 1) to investigate a child’s concept of pain based on seven domains (e.g. ‘How pain works’) identified by pediatric pain experts in a recent survey as relevant and appropriate.²² The interview script aligned with guidelines for interviewing children,²³⁻²⁵ and strategies supported by these guidelines were used in the interviews. For example, children were asked to choose their three favourite coloured pens as an icebreaker activity; these pens were then used for the drawing activities. In addition, vignettes with familiar media and cartoon characters were used to increase engagement.²⁶ For example, an open-ended question of “How does pain work?” was followed by a child-friendly cartoon video and children were asked, “Which body part decides when you will feel pain?”. This question was explained, and clarification of understanding was ensured by asking response-dependent follow-up questions such as “What else does your brain do?”.

The team of investigators (including physiotherapists, clinical psychologists, and experts in methodology) provided feedback regarding script development. Pilot testing on three healthy pain-free children (aged 7, 8 and 10 years), discussions with education and linguistics experts, and further review by the investigators, resulted in re-wording of several questions to ensure that the questions and language were appropriate for children aged 8-12 years. For example,

clarifying questions using synonyms (such as “replace the word ‘helpful’ with ‘useful’ if needed”, and ask “What word could we use instead of that?”) were incorporated. Following an iterative process, appropriate terminology and language resulting from interviews informed the focus of subsequent interviews until data saturation was reached.

Data Collection

Semi-structured interviews were conducted by J.W.P. (Male physiotherapist, a PhD candidate, and no previous relationship with potential participants). To maximize feelings of physical and emotional comfort, interviews took place at a location chosen by the participant (The Children’s Hospital at Westmead, Macquarie University, or in participants’ homes). Parents/caregivers were invited to be present for each interview but no questions were asked of them.²⁴ Parents were asked not to contribute unless they felt that their child was upset or distressed, at which point the interview was to be terminated and an offer to speak to the Clinical Psychologist would be made.

A PowerPoint presentation displayed prompts on a laptop screen to guide the discussion. The interviews were audio recorded to enable transcription. Field notes of contextual details and non-verbal expressions were recorded for data analysis and interpretation. For respondent checking, a summary of the interview transcript was provided to families for comment and feedback to enhance the analytical framework,²⁷ and a full copy of the transcript was available upon request.

A range of demographic characteristics, including age, sex, ethnicity, school year, postcode [for socioeconomic status: using the Australian Socio-Economic Indexes For Areas (SEIFA) ²⁸], parental education level and working status were collected in a demographics questionnaire for

1
2
3 parents. For children with persistent pain, parent-reported location and duration of pain were also
4
5 collected before the commencement of the interview.
6
7
8
9

10 Data triangulation (using multiple methods to enhance understanding) was achieved by including
11
12 a drawing task during the interviews.¹⁴ For this drawing task, children were asked to “draw
13
14 whatever the word ‘pain’ makes you think of”. This follows methodology of a study
15
16 investigating the concept of death in children.²⁹ The other written task involved labelling a
17
18 diagram of their nervous system (brain, spinal cord and nerves).³⁰ If a child had difficulty with
19
20 this, they were asked to verbally label the diagram, or point to where their own brain, spinal cord
21
22 and nerves are on their own body.
23
24
25
26
27

28 *Analysis*

29
30 Audio recordings from interviews were transcribed verbatim. An adapted grounded theory
31
32 approach¹⁹ was used to analyse and synthesize the data. Data was independently coded line-by-
33
34 line by 3 researchers (J.W.P., V.P., and T.N.) using NVivo software.³¹ Drawings were
35
36 thematically analysed by the investigators as has been done in previous research.^{32 33} Analytical
37
38 themes were inductively developed and agreed upon through an iterative process. Emerging
39
40 themes were then scrutinized by the entire investigative team to attain consensus.
41
42
43
44
45
46

47 *Public and Patient Involvement*

48
49 Patients were involved in piloting the interview script to identify the most appropriate language
50
51 to be used in the interviews and the time required to participate. The thematic analysis was
52
53
54
55
56
57
58
59
60

directly informed by participants as it was an iterative process. Patients and the public were not involved in informing other stages of the research process.

Results:

Information sheets were distributed and 16 recruitment emails were sent to parents of children who contacted the investigators stating that they wanted to participate. All families replied to the invitation and consented to participate. Therefore, 16 participants, 8 with persistent pain and 8 children who were pain-free, were enrolled in the study and completed an interview. The median (IQR) interview duration was 33 (27-36) minutes.

Participants

The participants included 9 males and 7 females, with a median (IQR) age of 10.0 (8.8-11.0) years. All children identified themselves as Australian, with 7 children reporting a second ethnicity including Chinese, Lebanese, Indian, Dutch and Eurasian. The median (IQR) Index of Relative Socio-economic Advantage and Disadvantage (IRSAD) SEIFA scores (Likert scale of 1-5 rating from disadvantaged to advantaged) was 4 (2.8-5). For children with persistent pain, back pain was most common, and pain duration ranged from 1-10 years. Table 2 details the demographic characteristics of participants.

Key themes

Four themes emerged from the data: 'my pain-related knowledge', 'pain in the world around me', 'pain in me', and 'communicating my concept of pain'. Illustrative quotations for each theme are provided in Table 3.

Theme 1: My pain-related knowledge

A child’s knowledge of the purpose of pain, pain-related anatomy, and pain mechanisms varied between individuals, while children consistently used concrete terms to describe their pain-related knowledge. Age influenced a child’s ability to express their concept of pain with younger children responding with uncertainty at times. No knowledge differences were identified between children who had persistent pain and children who were pain-free.

Subtheme: Varied knowledge of pain’s purpose, pain-related anatomy, and pain mechanisms.

When asked if pain had ‘a job in the body’, to identify pain’s purpose, children used familiar terms and referred to the idea of pain keeping them safe by protecting, warning, alerting, or knowing. The words ‘injury’ and ‘pain’ were both commonly associated with the word ‘hurt’, resulting in some children expressing confusion about pain’s purpose. When asked to ‘draw whatever the word “pain” makes you think of’, the children’s drawings frequently included an injury (Figure 2). Children used a variety of examples for injuries in their drawings and explanations, such as breaking an arm or leg, having a sprained ankle, or having fallen over.

Regarding pain-related anatomy, all children noted the importance of the brain and correctly labelled the brain inside the head on the diagram (Figure 1). A range of age-related understanding was demonstrated in this task and no differences were identified between children who had persistent pain and children who were pain-free. Children commonly stated that the

1
2
3 brain controls when and how someone feels pain. Exceptions to this were responses that were
4
5 vague such as "the head", "there is no part", or they were unsure. The spinal cord was rarely
6
7 included when explaining key elements of the nervous system involved in pain; children
8
9 frequently drew this in the back disconnected from the brain or were not sure where it was
10
11 located when questioned. Varied understanding of the term 'nerves' was observed; whilst some
12
13 children said that nerves are "everywhere", others confused nerves with feelings of nervousness
14
15 and some children confused nerves with structures such as veins. No child identified
16
17 interconnections within the nervous system, however when directly asked, approximately half of
18
19 each group said the brain, spinal cord and nerves are connected.
20
21
22
23
24
25

26 Knowledge about pain mechanisms also varied. To explain how pain is experienced, children
27
28 frequently reported ideas of "sending messages" or "telling the brain". Metaphors were also used
29
30 to explain the physiological processes underpinning pain, such as "it's like a phone call to your
31
32 brain". Some children referred to ideas of sensitivity or pain tolerance to justify that everyone's
33
34 pain is different. When offered a scenario whereby a cartoon character, the investigator (J.W.P.),
35
36 and the child all accidentally hit their thumbs with a hammer using the same force, most children
37
38 identified that each person or character would not feel the same pain. The main reasons for this
39
40 were explained as being inherently different people, having different levels of 'toughness' or
41
42 different ages.
43
44
45
46
47
48

49 Subtheme: Influences of age

50
51
52
53
54
55
56
57
58
59
60

Younger children tended to use simpler terminology. Some children aged 8 and 9 years more frequently responded “I don’t know” and appeared less confident of their responses, speaking softly when uncertain. In contrast, older children tended to use more complex terminology, such as phrases like “physically harming”. All children in the study appeared to focus more on concrete ideas than on abstract ideas, such as pain having contextual influences. For example, when asked why pain persists for some people and not for others, the most common response was that the person was not caring for an injury correctly.

Children’s understanding of pain-related abstract concepts was influenced by their age. Reciting words from a question was seen as distinctly different from understanding a concept. For example, when given a vignette of someone with persisting pain without an injury and asked about pain management strategies, children responded by suggesting the opposite to the understood pathophysiology (Table 3). It is also important to note that social influences of parents, teachers, and siblings, (explored in ‘Theme 2’) typically vary for different childhood ages.

Theme 2: Pain in the world around me

A child’s concept of pain is strongly influenced by their environmental and social context, including perceptions of how other people respond to pain. Consequently, differences in the concept of pain between children with and without persistent pain were evident.

Subtheme: Contextual influences on a child’s concept of pain

1
2
3
4
5 A child's exposure to family, school, and media, appeared to influence their concept of pain, to
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A child's exposure to family, school, and media, appeared to influence their concept of pain, to varying extents between individuals. Children reported they learnt about pain from family members, particularly parents, as well as at school and via media on television and the internet. To explain various aspects of their concept of pain, children with persistent pain also used examples of their interactions with health professionals discussing their pain experience despite not having previously received specific pain education. In contrast, children who were pain-free only used examples of family members and people at school.

Subtheme: Perceptions of how other people respond to pain

Children with and without persistent pain considered other people to be 'tough' or 'weak' based on their responses to painful experiences. People who felt a lot of pain were perceived by children as being weaker. People who ignored their pain were perceived as being tougher. The social stigma surrounding pain was also highlighted in phrases describing pain sensitivity. For example, some children viewed age as a determinant of pain sensitivity, generally stating that 'toughness' increases as a person ages until they "get old" when it decreases significantly. Other children considered 'toughness' to be an inherent quality, whilst others did not and emphasized "getting stronger" decreases sensitivity to pain.

Theme 3: Pain in me

1
2
3 A child’s concept of pain was based upon their individual experiences with pain and injury. All
4
5 children clearly described physical aspects of pain, whereas emotional aspects were relatively
6
7 hidden and were revealed only by children with persistent pain upon further verbal probing or
8
9 drawing tasks. The impact of persistent pain was highly individualized and related deeply to their
10
11 past experiences.
12
13

14
15
16
17 Subtheme: Experiences of any pain/injury
18

19
20
21 Children with and without persistent pain explained their concept of pain by using examples of
22
23 pain and injury from their own experiences. Regarding the purpose that pain serves, most
24
25 children reported that pain is sometimes a helpful thing to feel. The most common reason for this
26
27 was a warning to avoid further problems, however, two children described the idea of pain being
28
29 helpful to avoid attending something deemed boring or undesirable such as “a sport” or
30
31 “presentation day”.
32
33

34
35
36
37 Subtheme: The impact of having persistent pain
38

39
40
41 Children with persistent pain expressed different aspects of their concept of pain compared to
42
43 children without pain, with these aspects being dependent on their individual pain experiences.
44
45 Children with persistent pain considered broader aspects of the pain experience when describing
46
47 what ‘pain’ is, such as the impact of surgery, effects of feelings, the idea of healing, and the
48
49 diverse influences of health professionals.
50
51
52
53
54
55
56
57
58
59
60

When children were provided with a vignette and asked if a character could have pain without knowing about it, children were generally uncertain with some children responding purely based on their own experience without reference to the vignette. There appeared to be a difference in explanations between children with and without persistent pain; children who were pain-free responded with regards to the vignette only whereas children with persistent pain used the vignette with reference to their own personal experience.

Children with persistent pain focused more on emotional aspects of pain than children who were pain-free. When undertaking a drawing task, most children with persistent pain drew emotional elements to describe pain, whereas most children who were pain-free did not (Figure 2). Some children were confused when questioned about aspects of their drawing that illustrated emotional pain, mental pain, or nightmares, and were unable to verbally communicate how these types of pain related to a person's overall pain experience.

Further investigating what children think pain is, the discussion around "Is pain real?" varied between children with and without persistent pain. While some children in both groups described some confusion around the term 'real' when asked this question, most children with persistent pain said that pain is not always real. In contrast, most children who were pain-free said that pain is always real. Having persistent pain appeared to influence views on the 'reality' of pain.

Subtheme: The physical is obvious, but the emotional is hidden

Children tended to focus their discussions on physical and visible components of pain-related processes. The most obvious example of this was mentioned previously; that children without persistent pain did not draw emotional elements of pain.

When children discussed possible mechanisms for how different factors can influence pain, most children with persistent pain considered their experiences and stated that they could use self-taught management techniques, such as distracting themselves from pain by listening to music. In stark contrast, children who were pain-free did not mention cognitive strategies such as distraction. Other than attentional strategies, children did not mention any other factors that can influence the experience of pain. All children interviewed appeared unsure about the possible influence of other senses on pain, for example visual input.

Theme 4: Communicating my concept of pain

Children used verbal descriptions, drawings and diagrams to describe their concept of pain with drawing tasks and vignettes aiding some children’s communication during the interviews. No differences in the use of these methods of communicating were identified between children who had persistent pain and children who were pain-free.

Subtheme: Verbal communication

Children’s verbal descriptions of their concept of pain were individual and varied. For example, some children described the physiological processes underpinning pain using terms such as

‘chemicals’ whereas others used terms such as ‘virus’ and ‘disease’. The words ‘pain’, ‘injury’ and ‘hurt’ were used interchangeably by multiple children. No differences between groups were identified for this transposable use of terms.

Subtheme: Individual engagement

The value of the drawing tasks and vignettes appeared to relate to levels of engagement with the task, and this value did not appear related to age. The drawing task was very helpful in enabling some children to articulate thoughts they had not verbally communicated. Asking the follow-up question, “why did you draw that?” gave children an opportunity to expand on their thoughts.

For example, some children did not draw anything when first asked, but later drew representations, such as a face with tears. In contrast, some children did not like the drawing task and did not feel comfortable with this activity, preferring to speak about it and leaving the drawing page mostly blank.

Proposed conceptual framework

The potential interactions between the themes resulting from the analysis and a child’s concept of pain form a proposed conceptual framework that is visually presented in Figure 3.

The first three themes are inputs and combine in varying proportions to produce the output which is how a child communicates their concept of pain.

Discussion:

This study is the first known to the authors to investigate a child’s concept of what pain is, why it exists, and how it is experienced. The primary objective of this study was to gain an insight into the way that children aged 8-12 years, who had not received pain education, conceptualize pain. Four themes emerged from the data: ‘my pain-related knowledge’, ‘pain in the world around me’, ‘pain in me’, and ‘communicating my concept of pain’. Children’s communication of their concept of pain was based on their individual knowledge and experience. Children with persistent pain considered broader aspects of pain and focused more on emotional impacts and influences than children who were pain-free.

Children’s experiences of pain have been reported in a number of previous qualitative studies. For example, one study considered children’s descriptions of types of pain¹⁵ and a recent study of children aged 10-18 years considered personal experiences of neuropathic pain and types of language used.³⁴ Similarly in studies of adults, the experiences of pain have been explored.^{35 36} These studies differed from the present study in that the focus was on the experience of pain, rather than a person’s concept of pain. Children’s conceptual frameworks have been studied using drawings with the concept of death,²⁹ but the present study is the first to apply drawing methodology to investigate the concept of pain in children. Metaphors and explanations about pain that clinicians use with children have been discussed in the literature,³⁷ but the metaphors and explanations that children themselves use have not been explored.

Using the conceptual framework in Figure 3, several important issues are raised for future research regarding clinical practice. Firstly, it is currently unknown the extent to which each of the themes impacts upon a child’s concept of pain. Does a child’s experience have more of an

1
2
3 impact on their concept of pain than the impact of their knowledge or emotions? The relative size
4
5 of each 'raindrop' is not yet known because this study was of a small and somewhat specific
6
7 sample of children aged 8-12 years. Nonetheless, the extent to which each of the themes
8
9 influences a child's concept of pain is likely to be highly idiosyncratic, differing from one child
10
11 to another and also changing over time. Secondly, it is unknown if other factors contribute to the
12
13 composition of the puddle. The 'blank' raindrops in Figure 3 highlight that, for an individual
14
15 child, there could be other factors involved that were not identified in these interviews. Thirdly,
16
17 how can changes in a child's concept of pain 'puddle' be assessed? Conceptual change theory³⁸
18
19 suggests that the 'puddle' can theoretically change with education, however no research exists to
20
21 confirm this. A child's concept of pain assessment tool would be required to investigate this
22
23 further. Further research would also be required to explore whether the findings relate to children
24
25 below the age of 8 years, adolescents, adults and non-English speaking populations.
26
27
28
29
30
31
32

33 A key principle of clinical practice requiring future exploration is how patient care is
34
35 appropriately tailored and individualized based on a child's concept of pain. The results of this
36
37 study may be useful for assessing children prior to pain education, considering the language and
38
39 other communication tools used, to ensure care is individualized based on identified needs. We
40
41 found that descriptions of a child's concept of pain, their experiences, their vocabulary and use
42
43 of different metaphors, levels of engagement, and the interview duration, all varied from child to
44
45 child. The helpfulness of using a variety of media (e.g. drawings, videos, or text) appeared to
46
47 relate more to a child's level of comfort with the medium or level of engagement, rather than
48
49 their age. Therefore, as part of a flexible approach to communicating with children, using a range
50
51 of media is recommended. In particular, drawings can provide more information and/or more
52
53
54
55
56
57
58
59
60

accurate information compared with narrative methods, and they can increase comfort, facilitate memory retrieval, and help children organize verbal reports.³⁹ Both the open-ended drawing and the diagram-labelling task provided helpful information and enhanced other verbal responses in this study. Several children commented at the conclusion of the interviews that they enjoyed discussing the vignettes provided, which supports the established benefits of using vignettes clinically and in qualitative research.⁴⁰ Because a child's concept of pain is not always overtly communicated by the child, particularly the 'hidden' emotional components which affect communication,⁴¹ using vignettes and drawings in pain education or assessments may be particularly beneficial in clinical practice and should be investigated. In addition to these future clinical directions, exploring the effects of different pain durations and pain diagnoses on a child's focus on emotional aspects of pain is warranted.

A further potential clinical implication is that a more flexible and individualized assessment of a child's concept of pain could directly inform what pain education is likely to be most beneficial for a child. Pain education for adults is based on conceptual change theory where the assumption is that misconceptions exist.¹² The results of the current study show that complex concepts are still developing in childhood, and so pain education for children should also incorporate educational approaches where concepts are taught with gradually increasing complexity built on previous conceptual development.⁴² Depending on the different variables that have made up an individual child's concept of pain 'puddle' (Figure 3), one child may require different management approaches, explicitly delivering varying depths of pain education in different ways. For example, unexplored variables may be impacting on a child's concept of pain, such as their intellectual and emotional intelligence, their general attitude, fears, social circumstances,

1
2
3 and other aspects of their history. In addition, it is possible that some children may not require
4
5 any formal pain education, but rather benefit most from reassurance from an adult with influence
6
7 such as a parent, teacher or health professional. Education deemed too complex by a child may
8
9 result in misunderstanding and fear,⁴³ which could potentially aid in the development and/or
10
11 maintenance of persistent pain and disability.⁴⁴
12
13
14
15

16
17 From our findings, a child's broader social and environmental influences, as well as domains
18
19 such as fear-avoidance, have logical links to their concept of pain and warrant further
20
21 investigation. For example, some children stated that they had forgotten about a bruise and "just
22
23 moved on", whereas others use pain as a means to avoid activities. The influence of trusted
24
25 people, such as parents⁴⁵ or health professionals, giving children 'permission' to feel and
26
27 communicate their experience of pain is also a significant consideration for clinical practice.
28
29 Some children with persistent pain explained that they thought about pain sensitivity and
30
31 persistence in terms of the overall strength or weakness of others, or their age. A child's
32
33 perspective on 'toughness' is therefore worth investigating when assessing a child's concept of
34
35 pain.
36
37
38
39
40
41

42 The terminology and understanding of neuroanatomy used in descriptions of a child's concept of
43
44 pain is also important. In the current study children typically conceptualized the nervous system
45
46 without inclusion of the spinal cord and considered it to have two parts; the brain and a
47
48 peripheral component, and uncertainty existed regarding how these two parts are connected.
49
50 When providing education regarding pain-related neuroanatomy to children, phrases such as
51
52 'spinal cord', 'nerves', and 'real', are likely to be poorly understood and potentially
53
54
55
56
57
58
59
60

misinterpreted. Children appear to grasp the concept of the brain receiving warning messages and, based on these messages and many other factors, producing pain to keep the body safe. It is critical to carefully consider language when discussing pain with children, as younger children in this age group tended to only use and understand simpler terminology. Age-appropriate language used in conversations by health professionals may facilitate the therapeutic alliance. For example, if a child uses a particular word or phrase to describe their concept of pain, a patient-centred empathetic communication strategy⁴⁶ would be for the clinician to incorporate the child's language in the discussion rather than using technical terms. In addition, the apparent lack of awareness regarding how sensory input and distraction can influence pain provides some support for educating children at an age-appropriate level about biological mechanisms underpinning pain.

One strength of this study was the use of novel data collection methods, such as drawing tasks and vignettes, as a means of data triangulation.⁴⁷ Another strength was the large variations in pain experience amongst the group of 8 to 12-year old children with persistent pain; some children had lifelong permanent physical disabilities whereas others had recurrent headaches for just over one year, capturing individual variability in pain experiences.⁴⁸ A limitation of this study is the potential lack of transferability. This study only included children from New South Wales, Australia who spoke English. It is unknown if the results would be transferable to other cultures and age groups. A further limitation is that other 'raindrops' in the proposed conceptual framework could exist that were not identified and explored in these interviews.

1
2
3 In conclusion, Australian children aged 8-12 years who have not received formal pain education
4
5 had concepts of pain that depended on a child's knowledge and experiences. The importance of
6
7 gaining an in-depth understanding of a child's previous pain-related experiences and knowledge
8
9 is emphasized, to facilitate clear and meaningful pain education. The use of age-appropriate
10
11 language, in combination with novel assessment and education tasks such as drawing and
12
13 discussing vignettes, allowed children with persistent pain to communicate their individual
14
15 concept of pain. Future research is needed regarding the assessment of a child's concept of pain
16
17 to improve health interactions and outcomes for children with pain.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. King S, Chambers CT, Huguet A, et al. The epidemiology of chronic pain in children and adolescents revisited: a systematic review. *Pain* 2011;152(12):2729-38. doi: 10.1016/j.pain.2011.07.016

2. Swain MS, Henschke N, Kamper SJ, et al. An international survey of pain in adolescents. *BMC Public Health* 2014;14(1):447.

3. Access Economics. The high price of pain: the economic impact of persistent pain in Australia. Sydney: MBF Foundation 2007:9-11.

4. Hunfeld JA, Perquin CW, Duivenvoorden HJ, et al. Chronic pain and its impact on quality of life in adolescents and their families. *J Pediatr Psychol* 2001;26(3):145-53.

5. Hassett AL, Hilliard PE, Goesling J, et al. Reports of chronic pain in childhood and adolescence among patients at a tertiary care pain clinic. *J Pain* 2013;14(11):1390-97.

6. Lisman-van Leeuwen Y, Spee LA, Benninga MA, et al. Prognosis of abdominal pain in children in primary care—a prospective cohort study. *Ann Fam Med* 2013;11(3):238-44.

7. Kosola S, Mundy LK, Sawyer SM, et al. Pain and learning in primary school: a population-based study. *Pain* 2017;158(9):1825-30. doi: 10.1097/j.pain.0000000000000984

8. Zernikow B, Wager J, Hechler T, et al. Characteristics of highly impaired children with severe chronic pain: a 5-year retrospective study on 2249 pediatric pain patients. *BMC Pediatr* 2012;12(1):54.

9. Coffelt TA, Bauer BD, Carroll AE. Inpatient characteristics of the child admitted with chronic pain. *Pediatrics* 2013;132(2):e422-e29.

10. Groenewald CB, Essner BS, Wright D, et al. The economic costs of chronic pain among a cohort of treatment-seeking adolescents in the United States. *J Pain* 2014;15(9):925-33.

11. Kamper SJ, Apeldoorn A, Chiarotto A, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain: Cochrane systematic review and meta-analysis. *BMJ* 2015;350:h444.
12. Moseley GL, Butler DS. Fifteen Years of Explaining Pain: The Past, Present, and Future. *J Pain* 2015;16(9):807-13. doi: 10.1016/j.jpain.2015.05.005
13. Louw A, Zimney K, Puentedura EJ, et al. The efficacy of pain neuroscience education on musculoskeletal pain: A systematic review of the literature. *Physiother Theory Pract* 2016;32(5):332-55. doi: 10.1080/09593985.2016.1194646
14. Driessnack M. Children's drawings as facilitators of communication: a meta-analysis. *J Pediatr Nurs* 2005;20(6):415-23.
15. Harbeck C, Peterson L. Elephants dancing in my head: a developmental approach to children's concepts of specific pains. *Child Dev* 1992;63(1):138-49.
16. Jacobson CJ, Jr., Kashikar-Zuck S, Farrell J, et al. Qualitative Evaluation of Pediatric Pain Behavior, Quality, and Intensity Item Candidates and the PROMIS Pain Domain Framework in Children With Chronic Pain. *J Pain* 2015;16(12):1243-55. doi: 10.1016/j.jpain.2015.08.007
17. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19(6):349-57. doi: 10.1093/intqhc/mzm042
18. Kvale S. *InterViews*. Thousands Oaks: Ca: Sage, 1996.
19. Liamputtong P. *Qualitative research methods*. 3rd ed. Victoria, Australia: Oxford University Press 2009.

20. Arbuckle R, Abetz-Webb L. "Not just little adults": qualitative methods to support the development of pediatric patient-reported outcomes. *Patient* 2013;6(3):143-59. doi: 10.1007/s40271-013-0022-3

21. Christie D, Viner R. ABC of adolescence: Adolescent development. *BMJ* 2005;330(7486):301.

22. Pate JW, Hush JM, Hancock MJ, et al. A Child's Concept of Pain: An International Survey of Pediatric Pain Experts. *Children* 2018;5(1):12.

23. Docherty S, Sandelowski M. Focus on qualitative methods: Interviewing children. *Res Nurs Health* 1999;22(2):177-85.

24. Lobe B, Livingstone S, Olafsson K, et al. Best Practice Research Guide: How to research children and online technologies in comparative perspective: EU Kids Online 2008.

25. Kortessluoma RL, Hentinen M, Nikkonen M. Conducting a qualitative child interview: methodological considerations. *J Adv Nurs* 2003;42(5):434-41.

26. Ritchie J, Lewis J, Nicholls CM, et al. Qualitative research practice: A guide for social science students and researchers: Sage 2013.

27. Popay J, Rogers A, Williams G. Rationale and standards for the systematic review of qualitative literature in health services research. *Qual Health Res* 1998;8(3):341-51.

28. Pink B. Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA), Australia, 2011. Technical Paper. *Australian Bureau of Statistics (ABS), ed Canberra: Australian Government* 2011:54-58.

29. Bonoti F, Leondari A, Mastora A. Exploring children's understanding of death: through drawings and the death concept questionnaire. *Death Stud* 2013;37(1):47-60.

30. Buckley CA, Waring MJ. Using diagrams to support the research process: Examples from grounded theory. *Qual Res* 2013;13(2):148-72.
31. QSR International Pty Ltd. NVivo Qualitative Data Analysis Software (Version 9): QSR International Pty Ltd Doncaster, Vic, Australia, 2006.
32. Tamm ME, Granqvist A. The meaning of death for children and adolescents: A phenomenographic study of drawings. *Death Stud* 1995;19(3):203-22.
33. Angell C, Alexander J, Hunt JA. 'Draw, write and tell': A literature review and methodological development on the 'draw and write' research method. *Early Child Res* 2015;13(1):17-28.
34. Yeung K-K, Engle L, Rabel A, et al. It just feels weird!: a qualitative study of how children aged 10–18 years describe neuropathic pain. *Disabil Rehabil* 2017;39(17):1695-702. doi: 10.1080/09638288.2016.1207107
35. Bunzli S, Watkins R, Smith A, et al. Lives on hold: a qualitative synthesis exploring the experience of chronic low-back pain. *Clin J Pain* 2013;29(10):907-16.
36. Sim J, Madden S. Illness experience in fibromyalgia syndrome: a metasynthesis of qualitative studies. *Soc Sci Med* 2008;67(1):57-67.
37. Coakley R, Schechter N. Chronic pain is like... The clinical use of analogy and metaphor in the treatment of chronic pain in children. *Pediatric Pain Letter* 2013;15(1):1-8.
38. Posner GJ, Strike KA, Hewson PW, et al. Accommodation of a scientific conception: Toward a theory of conceptual change. *Sci Educ* 1982;66(2):211-27.
39. Gross J, Hayne H. Drawing facilitates children's verbal reports of emotionally laden events. *J Exp Psychol Appl* 1998;4(2):163.

40. Barter C, Renold E. 'I wanna tell you a story': exploring the application of vignettes in qualitative research with children and young people. *Int J Soc Res Methodol* 2000;3(4):307-23.

41. Hadjistavropoulos T, Craig KD, Duck S, et al. A biopsychosocial formulation of pain communication. *Psychol Bull* 2011;137(6):910.

42. Novak JD, Cañas AJ. The theory underlying concept maps and how to construct and use them. Pensacola: Institute for Human and Machine Cognition, 2008.

43. Simons LE, Kaczynski KJ. The Fear Avoidance model of chronic pain: examination for pediatric application. *The Journal of Pain* 2012;13(9):827-35.

44. Zale EL, Ditre JW. Pain-related fear, disability, and the fear-avoidance model of chronic pain. *Current opinion in psychology* 2015;5:24-30.

45. Evans S, Tsao JC, Lu Q, et al. Parent-child pain relationships from a psychosocial perspective: a review of the literature. *J Pain Manag* 2008;1(3):237.

46. Linton SJ, Flink IK, Nilsson E, et al. Can training in empathetic validation improve medical students' communication with patients suffering pain? A test of concept. *Pain Rep* 2017;2(3):e600.

47. Krefting L. Rigor in qualitative research: The assessment of trustworthiness. *Am J Occup Ther* 1991;45(3):214-22.

48. Denk F, McMahon SB. Neurobiological basis for pain vulnerability: why me? *Pain* 2017;158:S108-S14.

Table 1. Example questions, drawing activities, and vignettes in the interview script.

<i>Questions</i>	<ul style="list-style-type: none"> • What do you think pain is? <ul style="list-style-type: none"> ◦ What can pain feel like? Any picture words you can think of? • What do you think happens inside your body for you to feel hurt/sore? <ul style="list-style-type: none"> ◦ Or “how do you think pain is made inside your body?” • What do you think of when you hear the word “injury”? • Can someone be injured but have no pain? • Can someone have pain but not be injured? <ul style="list-style-type: none"> ◦ Where did you learn about that? • The job of eyes is to see. The job of ears is to hear. The job of the nose is to _____. And if pain were to have a job, what do you think the job of pain would be? • Why does pain last a long time for some people and not for others? • Do you think pain is always real?
<i>Drawing activities</i>	<ul style="list-style-type: none"> • I want you to draw whatever the word “pain” makes you think of. • Draw where you think your brain, spinal cord and nerves are with 3 different coloured pencils on the diagram.
<i>Vignettes</i>	<ul style="list-style-type: none"> • What do you think is happening inside this man’s body (cartoon character in a video) when he puts his hand on the stove? <ul style="list-style-type: none"> ◦ Why doesn’t his hand just stay on the stove top? • Do you think this cartoon character can have pain without knowing about it? <ul style="list-style-type: none"> ◦ Or “...without being aware of it?” • Let’s pretend this character has had pain for a long time. If he feels sad, do you think this would change his pain? <ul style="list-style-type: none"> ◦ If yes: Do you think his pain would be more/less/ bigger/ smaller/ stronger/quieter? ◦ How do you think he could make his pain feel different/better/worse?

Table 2. Demographics of the participants.

Characteristics		n (%)
<i>Gender</i>		
Male		9 (56)
Female		7 (44)
<i>Age (years)</i>		
8		4 (25)
9		3 (19)
10		4 (25)
11		3 (19)
12		2 (13)
<i>Persistent Pain</i>		
Yes		8 (50)
-	Duration:	
	0-1 year	0 (0)
	1-2 years	5 (63)
	3-5 years	2 (25)
	6-10 years	1 (13)
	11+ years	0 (0)
-	Location:	
	Headache only	2 (25)
	Back only	2 (25)
	Neck only	1 (13)
	Multiple sites	3 (38)
No		8 (50)
<i>Interview Setting</i>		
Home		9 (56)
University		6 (38)
Hospital		1 (6)

Note: Characteristics may not sum to exactly 100% due to the effect of rounding.

Table 3. Quotations to illustrate the subthemes of the four main themes.

Themes and Subthemes	Quotations
1. My pain-related knowledge: - Varied knowledge of pain's purpose, pain-related anatomy, and pain mechanisms.	<p>"...To alert your body that there is a virus and you are sick or something." (Boy, 9, persistent pain)</p> <p>"...It's to know you've hurt your knee and you need to stop playing." (Boy, 8, persistent pain)</p> <p>"The nerves feel it, then they tell the brain, [draws line up to brain outside body] and then the brain tells the nerve maybe." (Boy, 9, persistent pain)</p> <p>"Umm because like your nerves send like... It's like a phone call to your brain saying "I just hurt myself" and then your brain will say, "oh like that's going to hurt" and you go aww!" (Girl, 10, pain-free)</p> <p>"Maybe [people with higher sensitivity] have more nerves? And some people might have a little less." (Boy, 11, pain-free)</p>
- Influences of age	[When asked how a vignette character could make his persistent abdominal pain feel different] "...Umm well not eat as much food if he is eating a lot. Basically, go on a diet, and depending on if he is fit or not, get fit." (Boy, 10, persistent pain)
2. Pain in the world around me: - Contextual influences on a child's concept of pain - Perceptions of how other people respond to pain	<p>"My mum... she has... and my teacher... they might have to get knee replacements. My teacher had one. And my mum has to get one. And my neighbour got one." (Girl, 9, persistent pain)</p> <p>[When asked where he learnt about the role of the brain] "Well I learnt most of it off TV, and a little bit from school." (Boy, 12, persistent pain)</p> <p>"... Some people might be more sensitive than others. Like, I guess some people who play like full contact sports they would be more tough because they are used to the punching and everything. And people who don't do any sports or who are really sensitive naturally – you just go like "that!" (touches hand to desk lightly) it really, really hurts." (Girl, 10, pain-free)</p> <p>"I guess it's just how you're born. And how your nerves work." (Girl, 10, pain-free)</p> <p>"Some people have stronger muscles... It's a matter between weakness and strength." (Boy, 11, persistent pain)</p> <p>[When asked about sensitivity] "It means umm... more vulnerable".</p> <p>"...Umm well not really, because some people have a high pain tolerance. Because if Sarah, Mary and I did that, Mary and Sarah would be crying because they have low pain tolerances. Because I'm kind of tougher than them, I'm the oldest." (Girl, 11, pain-free)</p> <p>[When asked if different people feel the same amount of pain with the same hammer-hitting-thumb-injury] "We wouldn't feel the same amount of pain. Because you're older and I'm younger. I can't hold in the pain as much as you. So I'd feel more than you." (Boy, 8, pain-free)</p> <p>"Well, if you're like 96 or something you'd feel pain like kids... Because you're getting old." (Boy, 8, pain-free)</p> <p>"[Adults] may be more used to it, because they may have hit it before – depending how often! Because they may have had longer time to experience pain on their thumbs." (Boy, 10, persistent pain)</p>
3. Pain in me: - Experiences of any pain/injury	<p>"Well pain is like if you've hurt yourself, it's the physical contact and it hurts. Or like after surgeries you'd have like pain and it hurts to move." (Girl, 12, persistent pain)</p> <p>"It helps us not get into any more trouble... it'll help us to not do [an injury] again. We'll remember last time and [do the task] differently." (Boy, 8, pain-free)</p> <p>"Well if you really didn't like something, like a sport, then you don't have to play it. It'd be helpful maybe." (Girl, 10, pain-free)</p> <p>"If you had, like we have a presentation day that goes for 2 hours at school. If you didn't want to go, if you have pain you can stay home." (Girl, 9, persistent pain)</p>
- The impact of having persistent pain	<p>"You could go see a doctor" (Boy, 8, pain-free)</p> <p>"If they have pain then they can go to a physio..." (Boy, 8, pain-free)</p> <p>"If you maybe have a sore tummy, and you're like "oh no I might have appendicitis!" or something like that. And then you go to the doctors and then they're like "oh no you're just sick" or something like that. You're like "Oh thank goodness, I've only got a fever!" (Girl, 11, pain-free)</p> <p>"Umm... I reckon no... I always know about it [pain]." (Boy, 10, persistent pain)</p> <p>"If you defined pain in the dictionary... it would be something that physically hurts you. So it's a fake thing. Just a warning... It depends what you mean by real?" (Boy, 12, persistent pain)</p> <p>[When asked if pain was always 'real'] "I think it's always real. Because sometimes in your body, like right now when I'm talking, pain can still be in your body even if it doesn't hurt." (Boy, 8, pain-free)</p>
- The physical is obvious, but the emotional is hidden	"I reckon if he [the vignette character] looked at like his favourite thing in the world, and got to play with it or just looked at it at least and just went like "I need it" – it would distract him from his pain because he is focussed on his favourite toy or something, in the world." (Boy, 10, persistent pain)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4. Communicating my concept of pain:	<i>[Pain is] "...when something hurts in your body" (Girl, 10, pain-free)</i>
	<i>"The body is fighting the virus or something." (Boy, 9, persistent pain)</i>
- Verbal communication	
- Individual engagement	<i>[Pointing to parts of the diagram whilst talking] "[Nerves] release chemicals that flow up to your brain. And then your brain sends down another chemical I think, that gives it pain." (Boy, 12, persistent pain)</i>

For peer review only

Figure Legends

Figure 1. Eight examples of labelled diagrams of the brain, spinal cord and nerves by children aged 8-12 years. The top row are drawings of children who were pain-free and the bottom row are drawings of children with persistent pain. No differences between children with and without persistent pain were identified in this drawing task.

Figure 2. Examples of drawings by children when asked to draw whatever the word 'pain' makes them think of. Panels (a) and (b) represent how pain-free children drew injuries. Panels (c) and (d) are examples of how children with persistent pain drew more emotional elements such as hearts and tears.

Figure 3. A proposed conceptual framework of the potential interactions between the themes resulting from the analysis and a child's concept of pain.

Notes:

1. The raindrops without labels highlight that all potential contributors to a child's concept of pain may not have been identified; and
2. The size of the raindrops vary from one child to the next.

1
2
3 **Word Count:** 5054 words
4
5

6
7
8 **Author Statement:**
9

10 Conception and design: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.
11

12 Planning and implementation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.
13

14 Data collection: J.W.P.
15

16 Coding data for thematic analysis: J.W.P., T.N., and V.P.
17

18 Analysis and interpretation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.
19

20 Drafting the article: J.W.P.
21

22 Review and editing: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.
23

24 Overall responsibility: J.W.P. under the supervision of V.P., J.M.H., and M.J.H.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Eight examples of labelled diagrams of the brain, spinal cord and nerves by children aged 8-12 years. The top row are drawings of children who were pain-free and the bottom row are drawings of children with persistent pain. No differences between children with and without persistent pain were identified in this drawing task.




Figure 2. Examples of drawings by children when asked to draw whatever the word 'pain' makes them think of. Panels (a) and (b) represent how pain-free children drew injuries. Panels (c) and (d) are examples of how children with persistent pain drew more emotional elements such as hearts and tears.




Figure 3. A proposed conceptual framework of the potential interactions between the themes resulting from the analysis and a child's concept of pain.

Notes:

1. The raindrops without labels highlight that all potential contributors to a child's concept of pain may not have been identified; and
2. The size of the raindrops vary from one child to the next.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

Developed from Table 1 of: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

No. Item	Guide questions/description	Reported on Page #
Domain 1: Research team and reflexivity		
<i>Personal Characteristics</i>		
1. Interviewer/facilitator	Which author/s conducted the interview or focus group?	Page 9
2. Credentials	What were the researcher’s credentials? E.g. PhD, MD	Page 9
3. Occupation	What was their occupation at the time of the study?	Page 9
4. Gender	Was the researcher male or female?	Page 9
5. Experience and training	What experience or training did the researcher have?	Page 9
<i>Relationship with participants</i>		
6. Relationship established	Was a relationship established prior to study commencement?	Page 9
7. Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	Page 9
8. Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. bias, assumptions, reasons and interests in the research topic	Page 9
Domain 2: Study design		
<i>Theoretical framework</i>		
9. Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	Page 11
<i>Participant selection</i>		
10. Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	Page 8
11. Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	Page 8
12. Sample size	How many participants were in the study?	Page 11
13. Non-participation	How many people refused to participate or dropped out? Reasons?	Page 11
<i>Setting</i>		

14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	Table 2 (Page 31)
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	Page 10
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	Table 2 (Page 31)
<i>Data collection</i>		
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	Table 1 (Page 30)
18. Repeat interviews	Were repeat interviews carried out? If yes, how many?	No, page 7
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	Page 10
20. Field notes	Were field notes made during and/or after the interview or focus group?	Page 10
21. Duration	What was the duration of the interviews or focus group?	Page 11
22. Data saturation	Was data saturation discussed?	Page 8 and 9
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	Page 10
Domain 3: analysis and findings		
<i>Data analysis</i>		
24. Number of data coders	How many data coders coded the data?	3 coders, Page 11
25. Description of the coding tree	Did authors provide a description of the coding tree?	Page 11
26. Derivation of themes	Were themes identified in advance or derived from the data?	Page 11
27. Software	What software, if applicable, was used to manage the data?	Page 11
28. Participant checking	Did participants provide feedback on the findings?	Page 10
<i>Reporting</i>		
29. Quotations presented	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	Table 3, page 32
30. Data and findings consistent	Was there consistency between the data presented and the findings?	Yes
31. Clarity of major themes	Were major themes clearly presented in the findings?	Yes
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	Yes

BMJ Open

Exploring the concept of pain of Australian children with and without pain: a qualitative study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-033199.R1
Article Type:	Original research
Date Submitted by the Author:	13-Sep-2019
Complete List of Authors:	Pate, Joshua; Macquarie University, Health Professions Noblet, Tim; Macquarie University, Health Professions Hush, Julia ; Macquarie University, Health Professions Hancock, Mark; Macquarie University, Health Professions Sandells, Renee; Children's Hospital at Westmead Pounder, Meg; Children's Hospital at Westmead Pacey, Verity; Macquarie University Faculty of Medicine and Health Sciences
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Qualitative research, Communication, Patient-centred medicine
Keywords:	pain science education, concept of pain, paediatric pain, QUALITATIVE RESEARCH, drawing task

SCHOLARONE™
Manuscripts

Title:

Exploring the concept of pain of Australian children with and without pain: a qualitative study.

Authors:

Joshua W. Pate^{1,2}, MRes, Senior Physiotherapist and PhD Candidate, joshua.pate@mq.edu.au

Tim Noblet¹, MSc (MSKMed), Senior Physiotherapist and PhD Candidate,
tim.noblet@mq.edu.au

Julia M. Hush¹, PhD, Professor of Physiotherapy, julia.hush@mq.edu.au

Mark J. Hancock¹, PhD, Professor of Physiotherapy, mark.hancock@mq.edu.au

Renee Sandells², MPsych (Clin), Clinical Psychologist, renee.sandells@health.nsw.gov.au

Meg Pounder², BSc (Physiotherapy), Senior Physiotherapist, meg.pounder@health.nsw.gov.au

Verity Pacey^{1,3}, PhD, Associate Professor of Physiotherapy, verity.pacey@mq.edu.au

Affiliations:

¹ Macquarie University, New South Wales, Australia

² Department of Pain Medicine, The Children’s Hospital at Westmead, New South Wales,
Australia

³ The University of Sydney Children’s Hospital Westmead Clinical School, New South Wales,
Australia

Corresponding Author:

Joshua Pate

Ground floor 75 Talavera Road

Macquarie University 2109

Phone: +61 2 9850 6626

Fax: +61 2 9850 6630

Email: joshua.pate@mq.edu.au

Institutional URL: <https://researchers.mq.edu.au/en/persons/joshua-pate>

Keywords:

concept of pain; pain science education; paediatric pain; qualitative interviews; drawing task

Previous presentations of the research:

Poster of one subtheme of the manuscript presented for IASP 2018 in Boston, titled: “Explaining pain to children with and without pain: what terminology should we use? Qualitative interviews describing pain-related neuroanatomy.”

Acknowledgments:

We thank all the children and their families for their openness and for giving their valuable time, energy and efforts to participate in this study. We also acknowledge Natasha Haynes, Nurse Practitioner from The Children’s Hospital at Westmead Pain Team, who helped to recruit participants and supported the study.

Funding:

J.W.P. is a Ph.D. Student funded by a Macquarie University Research Training Program (RTP) full-time scholarship.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflicts of Interest:

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Abstract:

Objective: A person's concept of pain can be defined as how they understand what pain actually is, what function it serves, and what biological processes are thought to underpin it. This study aimed to explore the concept of pain in children with and without persistent pain.

Design: In-depth, face-to-face interviews with drawing tasks were conducted with 16 children (aged 8-12 years) in New South Wales, Australia. Thematic analysis was used to analyse and synthesize the data.

Setting: Children with persistent pain were identified from a pain clinic waiting list in Australia, and children without pain were identified through advertising flyers and email bulletins at a university and hospital.

Participants: Eight children had persistent pain and eight children were pain-free.

Results: Four themes emerged from the data: 'my pain-related knowledge', 'pain in the world around me', 'pain in me', and 'communicating my concept of pain'. A conceptual framework of the potential interactions between the themes resulting from the analysis is proposed. The concept of pain of Australian children aged 8-12 years varied depending on their knowledge, experiences and literacy levels. For example, when undertaking a drawing task, children with persistent pain tended to draw emotional elements to describe pain, whereas children who were pain-free did not.

Conclusions: Gaining an in-depth understanding of a child's previous pain-related experiences and knowledge is important to facilitate clear and meaningful pain science education. The use of age-appropriate language, in combination with novel assessment and education tasks such as drawing and discussing vignettes, allowed children to communicate their individual concept of pain.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary:
Strengths and limitations of this study:

- One strength of this study was the use of drawing tasks and vignettes, as a means of data triangulation.
- Another strength was the large variations in pain experience amongst the group of 8 to 12-year old children with persistent pain; some participants had lifelong permanent physical disabilities whereas others had recurrent headaches for just over one year, capturing individual variability in pain experiences.
- A limitation of this study is the potential lack of transferability. This study only included children from New South Wales, Australia who spoke English. It is unknown if the results would be transferable to other cultures and age groups.
- A further limitation is that within the proposed conceptual framework, other influences on a child’s concept of pain could exist, however they were not identified and explored in these interviews.

Introduction:

Estimates of point-prevalence of persistent pain (pain that extends beyond the expected period of healing¹) throughout childhood range from 11% to 54%,^{2,3} which is comparable to that for adults.⁴ Persistent pain in children has detrimental effects on a child's quality of life⁵ and is a significant socioeconomic and health problem. For example, persistent pain in childhood can predict persistent pain in adulthood⁶ and is associated with increased pain-related disability,² school absence,⁷ poorer reading levels,⁸ and emotional distress.⁹ An increasing number of children are being admitted to hospitals for treatment of persistent pain¹⁰ and health service costs are high.¹¹

Multidisciplinary pain management is currently the gold standard for persistent pain management.¹² An important part of the multidisciplinary approach is pain science education, aiming to change a person's concept of pain and to improve pain and function by teaching the underlying biopsychosocial mechanisms of pain.¹³ A person's concept of pain can be defined as how they understand "what pain actually is, what function it serves, and what biological processes are thought to underpin it";¹⁴ these can be summarized as the 'what, why and how of pain'. Clinically, a range of methods for educational assessments and treatments are used for adults with persistent pain including drawing tasks.¹⁵ Despite drawing tasks being a robust interview strategy with children,¹⁶ these methods have not been applied to paediatric pain research to investigate a child's concept of pain.¹⁷

Limited research has been conducted that focuses on a child's concept of pain. For example, children who are pain-free have been asked to describe sensations relating to a grazed knee, an

injection, and a headache.¹⁸ In addition, pain-specific behaviours, pain quality, and pain intensity have been investigated in children with pain.¹⁹ However, a child’s concept of pain, as defined in the previous paragraph, has not been specifically investigated in children with or without persistent pain. Therefore, this remains poorly understood, providing clinicians little guidance on how to provide pain science education to children with persistent pain as part of multidisciplinary management, or to children who are pain-free as a strategy to prevent future pain, using language which is clear and unambiguous to a child. The aims of this study were (1) to explore the concept of pain of children diagnosed with persistent pain, and (2) to qualitatively compare the concept of pain between children who have pain with those who are pain-free.

Methods:

The study was approved by the Human Research Ethics Committees of Sydney Children’s Hospital Network (Ref: LNR/17/SCHN/268) and Macquarie University (Ref: 5201700820). All participants’ parents provided informed consent prior to participation in the study, and verbal assent was obtained from the participants prior to the commencement of the interview.

Study Design

This qualitative study utilized one-on interviews, lasting approximately 30-minutes, of children aged 8-12 years. To ensure explicit and comprehensive study reporting, the widely used Consolidated Criteria for Reporting Qualitative Health Research (COREQ)²⁰ was implemented. Children were asked to answer questions and complete activities relating to their concept of pain. By adapting grounded theory and utilizing thematic analysis,^{21 22} concepts in the transcripts were inductively identified and recorded and then a coding structure was iteratively developed and

refined until the aims of the study were achieved. Themes and sub-themes were developed from data analysis and synthesis.

Participants

Two groups of children aged 8-12 years in the state of New South Wales, Australia, were purposefully recruited; children who had persistent pain (greater than 3 months) and children who were pain-free and healthy by parent-report. Parent-report was used to minimise burden on the child being interviewed. The absence, presence, and/or persistence of pain was later confirmed throughout the interview with the child, allowing opportunity to clarify any discrepancies between reporting. Children were excluded from the study if they (1) had received pain science education from a health care professional, (2) were deemed not medically fit to be part of this research study by their treating physician/surgeon, or (3) were unable to participate in a 30-minute interview based on parent-report.

A minimum age for participants was set at 8 years, the youngest age a child can typically self-report and provide meaningful responses in interviews due to their linguistic and cognitive development.²³ The maximum age of participation was 12 years, an age prior to when abstract thinking capacities typically develop in the teenage years.²⁴ The age range was determined to meet a clinical need highlighted by research team members to provide improved assessment and management resources for pre-teenage children. A pragmatic approach incorporating age limits, rather than a formal assessment of cognitive development or language abilities, allows for direct applicability in time-limited clinical settings.

A purposive selection strategy was used to identify children of varying demographics and pain characteristics who were likely to meet the eligibility criteria. For all children, a mix of gender, age, and ethnic backgrounds was sought to capture diversity of experiences. For the children with persistent pain, purposive sampling attended to capturing variations in the duration and location of pain. Children were consecutively identified from The Children’s Hospital at Westmead, Australia Complex Pain Service waiting list, and through advertising flyers and email bulletins at Macquarie University and The Children’s Hospital at Westmead.

Children were interviewed until data saturation was reached. Data saturation was defined as when no additional new information was attained with additional interviews regarding the final themes; the three researchers analysing the data (J.W.P., V.P., and T.N.) met fortnightly throughout data collection and the thematic analysis to determine when data saturation was reached. Data collection was stopped when all three researchers agreed that saturation had taken place.

Interview Script

An interview script was developed (Table 1) to investigate a child’s concept of pain based on seven domains (e.g. ‘How pain works’). These domains were proposed in a recent survey of paediatric pain experts by categorizing items from the revised Neurophysiology of Pain Questionnaire (rNPQ)²⁵ and Explain Pain Target Concepts,²⁶ and they were rated as relevant and appropriate.²⁷ The interview script aligned with guidelines for interviewing children,²⁸⁻³⁰ and strategies supported by these guidelines were used in the interviews. For example, children were asked to choose their three favourite coloured pens as an icebreaker activity; these pens were

then used for the drawing activities. In addition, vignettes with familiar media and cartoon characters were used to increase engagement.³¹ For example, an open-ended question of “How does pain work?” was followed by a child-friendly cartoon video accessible on YouTube³² and children were asked, “Which body part decides when you will feel pain?”. This question was explained, and clarification of understanding was ensured by asking response-dependent follow-up questions such as “What else does your brain do?”.

The team of investigators (including physiotherapists, clinical psychologists, and experts in methodology) provided feedback regarding script development. Pilot testing on three healthy pain-free children (aged 7, 8 and 10 years), discussions with education and linguistics experts, and further review by the investigators, resulted in re-wording of several questions to ensure that the questions and language were appropriate for children aged 8-12 years. For example, clarifying questions using synonyms (such as “replace the word ‘helpful’ with ‘useful’ if needed”, and ask “What word could we use instead of that?”) were incorporated. Following an iterative process, appropriate terminology and language resulting from interviews informed the focus of subsequent interviews until data saturation was reached.

Data Collection

Semi-structured interviews were all conducted by J.W.P. (Male physiotherapist, a PhD candidate, and no previous relationship with potential participants). To maximize feelings of physical and emotional comfort, interviews took place at a location chosen by the participant (The Children’s Hospital at Westmead, Macquarie University, or in participants’ homes).

Parents/caregivers (from now on referred to as “parents”) were invited to be present for each

interview but no questions were asked of them.²⁹ Parents were asked not to contribute unless they felt that their child was upset or appeared distressed, at which point the interview was to be terminated and an offer to speak to the Clinical Psychologist would be made.

A PowerPoint presentation displayed prompts on a laptop screen to guide the discussion. The interviews were audio recorded to enable transcription. Field notes of contextual details and non-verbal expressions were recorded for data analysis and interpretation. For respondent checking, a summary of the interview transcript was provided to families for comment and feedback to enhance the analytical framework,³³ and a full copy of the transcript was available upon request.

A range of demographic characteristics, including age, sex, ethnicity, school year, postcode [for socioeconomic status: using the Australian Socio-Economic Indexes For Areas (SEIFA) ³⁴], parental education level and working status were collected in a demographics questionnaire completed by parents. For children with persistent pain, parent-reported location and duration of pain were also collected before the commencement of the interview.

Data triangulation (using multiple methods to enhance understanding) was achieved by including a drawing task during the interviews.¹⁶ For this drawing task, children were asked to “draw whatever the word ‘pain’ makes you think of”. This follows methodology of a study investigating the concept of death in children.³⁵ To explore the language children use to describe components of the nervous system and their familiarity with the anatomical components of this system, a second written task involved labelling a diagram of their nervous system (brain, spinal cord and nerves).³⁶ If a child had difficulty with this, they were asked to verbally label the

1
2
3 diagram, or point to where their own brain, spinal cord and nerves are on their own body.
4
5
6

7 8 *Analysis* 9

10 Audio recordings from interviews were transcribed by J.W.P verbatim. V.P. and T.N. checked
11 transcripts for accuracy. An adapted grounded theory approach²² was used to analyse and
12 synthesize the data. Data were independently coded line-by-line by three researchers (J.W.P.,
13 V.P., and T.N.) using NVivo software.³⁷ Drawings were thematically analysed by the
14 investigators as has been done in previous research.^{38 39} Analytical themes were inductively
15 developed and fully agreed upon by these three researchers through a rigorous iterative process.
16 The themes were scrutinised by a team of subject experts including paediatric clinical
17 psychologists, physiotherapists, academics and methodological experts, in two meetings during
18 the analytical process and one final meeting to attain 100% consensus.
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 *Public and Patient Involvement* 34

35 Patients were involved in piloting the interview script to identify the most appropriate language
36 to be used in the interviews and the time required to participate. The thematic analysis was
37 directly informed by participants as it was an iterative process. Patients and the general public
38 were not involved in informing other stages of the research process.
39
40
41
42
43
44
45
46

47 **Results:** 48

49 Information sheets were distributed and 16 recruitment emails in total were sent to parents of
50 children who contacted the investigators stating that they wanted to participate. All families
51 replied to the invitation and consented to participate. Therefore, 16 participants, 8 with persistent
52
53
54
55
56
57
58
59
60

pain and 8 children who were pain-free, were enrolled in the study and completed an interview between October 2017 and February 2018 before data saturation was reached. The median (IQR) interview duration was 33 (27-36) minutes. The three researchers (J.W.P., V.P., and T.N.) coding the transcripts agreed on the correctness of the transcripts, with no disagreements identified.

Participants

The participants included 9 males and 7 females, with a median age and interquartile range (IQR) of 10.0 (8.8-11.0) years. All participants identified themselves as Australian, with 7 participants reporting a second ethnicity including Chinese, Lebanese, Indian, Dutch and Eurasian. The median (IQR) Index of Relative Socio-economic Advantage and Disadvantage (IRSAD) SEIFA scores (Likert scale of 1-5 rating from disadvantaged to advantaged) was 4 (2.8-5). Of participants with persistent pain, five children in total reported back pain (63%), two children reported headaches (25%), and one child reported neck pain (13%). Pain duration ranged from 1-10 years with two participants reporting lifelong permanent physical disabilities. No discrepancies between parent- and child-reports of the participant’s pain history were identified. Table 2 details the demographic characteristics of participants. No participants reported feeling upset or appeared distressed during the interviews.

Key themes

Four themes emerged from the data: ‘my pain-related knowledge’, ‘pain in the world around me’, ‘pain in me’, and ‘communicating my concept of pain’. The first three themes focus on the content that participants conveyed, whereas the fourth theme focuses on the method they used to convey this content. Illustrative quotations for each theme are provided in Table 3.

Theme 1: My pain-related knowledge

Participant knowledge of the purpose of pain, pain-related anatomy, and pain mechanisms varied between individuals, while participants consistently used concrete terms to describe their pain-related knowledge. Age influenced the participants' ability to express their concept of pain with younger participants responding with uncertainty at times. Based on the thematic analysis, no differences in participant knowledge were identified between participants who had persistent pain and participants who were pain-free, nor based on their gender or ethnic background.

Subtheme: Varied knowledge of pain's purpose, pain-related anatomy, and pain mechanisms.

When asked if pain had 'a job in the body', to identify pain's purpose, participants used familiar terms and referred to the idea of pain keeping them safe by protecting, warning, alerting, or knowing. The words 'injury' and 'pain' were both commonly associated with the word 'hurt', resulting in some participants expressing confusion about pain's purpose. When asked to 'draw whatever the word "pain" makes you think of', the participants' drawings frequently included an injury (Figure 1). Participants used a variety of examples for injuries in their drawings and explanations, such as breaking an arm or leg, having a sprained ankle, or having fallen over.

Regarding pain-related anatomy, all participants noted the importance of the brain and correctly labelled the brain inside the head on the diagram (Figure 2). A range of understanding was demonstrated in this task and no differences were identified between participants who had

1
2
3 persistent pain and participants who were pain-free. Participants commonly stated that the brain
4 controls when and how someone feels pain. Exceptions to this were responses that were vague
5 such as "the head", "there is no part", or they were unsure. The spinal cord was rarely included
6 when explaining key elements of the nervous system involved in pain; participants frequently
7 drew this in the back disconnected from the brain or were not sure where it was located when
8 questioned. Participants of each age varied in their understanding of the nervous system. Varied
9 understanding of the term 'nerves' was observed; whilst some participants said that nerves are
10 "everywhere", others confused nerves with feelings of nervousness and some participants
11 confused nerves with structures such as veins. No child identified interconnections within the
12 nervous system, however when directly asked, approximately half of each group said the brain,
13 spinal cord and nerves are connected.

14
15 Knowledge about pain mechanisms also varied. To explain how pain is experienced, participants
16 frequently reported ideas of "sending messages" or "telling the brain". Metaphors were also used
17 to explain the physiological processes underpinning pain, such as "it's like a phone call to your
18 brain". Some participants referred to ideas of sensitivity or pain tolerance to justify that
19 everyone's pain is different. When participants were asked if they would feel the same or
20 different pain if a cartoon character, the investigator (J.W.P.) and the participant all accidentally hit
21 their thumbs with a hammer using the same force, most participants identified that each person
22 or character would not feel the same pain. Participants provided explanations for this such as
23 being inherently different people, having different levels of 'toughness' or different ages.

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Subtheme: Influences of age
55
56
57
58
59
60

Younger participants tended to use simpler terminology. Some participants aged 8 and 9 years more frequently responded “I don’t know” and appeared less confident of their responses, speaking softly when uncertain. In contrast, older participants tended to use more complex terminology, such as phrases like “physically harming”. All participants in the study appeared to focus more on concrete ideas than on abstract ideas, such as pain having contextual influences. For example, when asked why pain persists for some people and not for others, the most common response was that the person was not caring for an injury correctly.

Participant understanding of pain-related abstract concepts was influenced by their age, reflecting this age group’s stage of cognitive development. Reciting words from a question was seen as distinctly different from understanding a concept. For example, when given a vignette of someone with persisting pain without an injury and asked about pain management strategies, participants responded by suggesting the opposite to the understood pathophysiology (Table 3).

Theme 2: Pain in the world around me

A child’s concept of pain is strongly influenced by their environmental and social context, including perceptions of how other people respond to pain. Consequently, differences in the concept of pain between participants with and without persistent pain were evident.

Subtheme: Contextual influences on a child’s concept of pain

A child’s exposure to family, school, and media, appeared to influence their concept of pain, to varying extents between individuals. Participants reported they learnt about pain from family members, particularly parents, as well as at school and via media on television and the internet. To explain various aspects of their concept of pain, participants with persistent pain also used examples of their interactions with health professionals discussing their pain experience despite not having previously received specific pain science education. In contrast, participants who were pain-free only used examples of family members and people at school.

Subtheme: Perceptions of how other people respond to pain

Participants with and without persistent pain considered other people to be ‘tough’ or ‘weak’ based on their responses to painful experiences. People who felt a lot of pain were perceived by participants as being weaker. People who ignored their pain were perceived as being tougher. The social stigma surrounding pain was also highlighted in phrases describing pain sensitivity. For example, some participants viewed age as a determinant of pain sensitivity, generally stating that ‘toughness’ increases as a person ages until they “get old” when it decreases significantly. Other participants considered ‘toughness’ to be an inherent quality, whilst others did not and emphasized “getting stronger” decreases sensitivity to pain.

Theme 3: Pain in me

A child’s concept of pain was based upon their individual experiences with pain and injury. All participants clearly described physical aspects of pain, whereas emotional aspects were relatively

hidden and were revealed only by participants with persistent pain upon further verbal probing or drawing tasks. The impact of persistent pain was highly individualized and related deeply to their past experiences.

Subtheme: Experiences of any pain/injury

Participants with and without persistent pain explained their concept of pain by using examples of pain and injury from their own experiences. Regarding the purpose that pain serves, most participants reported that pain is sometimes a helpful thing to feel. The most common reason for this was a warning to avoid further problems, however, two participants (one with persistent pain and one without) described the idea of pain being helpful to avoid attending something deemed boring or undesirable such as “a sport” or “presentation day”.

Subtheme: The impact of having persistent pain

Participants with persistent pain expressed different aspects of their concept of pain compared to participants without pain, with these aspects being dependent on their individual pain experiences. Participants with persistent pain considered broader aspects of the pain experience when describing what ‘pain’ is, such as the impact of surgery, effects of feelings, the idea of healing, and the diverse influences of health professionals.

When participants were provided with a vignette and asked if a character could have pain without knowing about it, participants were generally uncertain with some responding purely

based on their own experience without reference to the vignette. There appeared to be a difference in explanations between participants with and without persistent pain; participants who were pain-free responded with regards to the vignette only whereas participants with persistent pain used the vignette with reference to their own personal experience.

Participants with persistent pain focused more on emotional aspects of pain than participants who were pain-free. When undertaking a drawing task, participants with persistent pain tended to draw emotional elements to describe pain, whereas participants who were pain-free did not (Figure 1). Some participants were confused when questioned about aspects of their drawing that illustrated emotional pain, mental pain, or nightmares, and were unable to verbally communicate how these types of pain related to a person’s overall pain experience.

Further investigating what children think pain is, the discussion around “Is pain real?” varied between participants with and without persistent pain. While some participants in both groups described some confusion around the term ‘real’ when asked this question, most participants with persistent pain said that pain is not always real. In contrast, most participants who were pain-free said that pain is always real. Having persistent pain appeared to influence views on the ‘reality’ of pain.

Subtheme: The physical is obvious, but the emotional is hidden

Participants tended to focus their discussions on physical and visible components of pain-related processes. The most obvious example of this was mentioned previously; that participants without persistent pain did not draw emotional elements of pain.

When participants discussed possible mechanisms for how different factors can influence pain, most participants with persistent pain considered their experiences and stated that they could use self-taught management techniques, such as distracting themselves from pain by listening to music. In stark contrast, participants who were pain-free did not mention cognitive strategies such as distraction. Other than attentional strategies, participants did not mention any other factors that can influence the experience of pain. All children interviewed appeared unsure about the possible influence of other senses on pain, for example visual input.

Theme 4: Communicating my concept of pain

Participants used verbal descriptions, drawings and diagrams to describe their concept of pain with drawing tasks and vignettes aiding some participants' communication during the interviews. No differences in the use of these methods of communicating were identified between participants who had persistent pain and participants who were pain-free.

Subtheme: Verbal communication

Participants' verbal descriptions of their concept of pain were individual and varied. For example, some participants described the physiological processes underpinning pain using terms such as 'chemicals' whereas others used terms such as 'virus' and 'disease'. The words 'pain', 'injury' and 'hurt' were used interchangeably by multiple participants. No differences between groups were identified for this transposable use of terms.

Subtheme: Individual engagement

The value of the drawing tasks and vignettes appeared to relate to levels of engagement with the task, and this value did not appear related to age, gender or ethnic background. The drawing task was very helpful in enabling some participants to articulate thoughts they had not verbally communicated. Asking the follow-up question, “why did you draw that?” gave participants an opportunity to expand on their thoughts. For example, some participants did not draw anything when first asked, but later drew representations, such as a face with tears. In contrast, four participants did not like the drawing task and did not feel comfortable with this activity, preferring to speak about it and leaving the drawing page mostly blank (8 years, male, pain-free; 12 years, male, persistent pain; 11 years, female, pain-free; 8 years, male, pain-free).

Proposed conceptual framework

The potential interactions between the themes resulting from the analysis and a child’s concept of pain form a proposed conceptual framework that is visually presented in Figure 3. The first three themes are inputs and combine in varying proportions to produce the output which is how a child communicates their concept of pain.

Discussion:

This study is the first known to the authors to investigate a child’s concept of what pain is, why it exists, and how it is experienced. The primary objective of this study was to gain insight into the way that children aged 8-12 years, who have not received pain science education, conceptualize

1
2
3 pain. Four themes emerged from the data: ‘my pain-related knowledge’, ‘pain in the world
4 around me’, ‘pain in me’, and ‘communicating my concept of pain’. Participants’
5
6 communication of their concept of pain was based on their individual knowledge and experience.
7
8
9
10 Participants with persistent pain considered broader aspects of pain and focused more on
11
12 emotional impacts and influences than participants who were pain-free.
13
14

15
16
17 Children’s experiences of pain have previously been reported in qualitative studies. As examples,
18
19 one study considered children’s descriptions of types of pain¹⁸, another focused on painful
20
21 experiences⁴⁰, another on children’s pain-related vocabulary⁴¹, and a recent study of children
22
23 aged 10-18 years with neuropathic pain considered personal pain experiences and language
24
25 used.⁴² Similarly in studies of adults, pain experiences have been explored.^{43 44} These studies
26
27 differed from the present study in that the focus was on the experience of pain, rather than a
28
29 person’s concept of pain. Children’s conceptual frameworks have been studied using drawings
30
31 with the concept of death,³⁵ but the present study is the first to apply drawing methodology to
32
33 investigate the concept of pain. Metaphors and explanations about pain that clinicians use with
34
35 children have been discussed in the literature,⁴⁵ but the metaphors and explanations that children
36
37 themselves use have not been explored.
38
39
40
41
42
43

44
45 Using the conceptual framework in Figure 3, several important issues are raised for future
46
47 research regarding clinical practice. First, it is currently unknown the extent to which each of the
48
49 themes impacts upon a child’s concept of pain. Does a child’s experience have more of an
50
51 impact on their concept of pain than the impact of their knowledge or emotions? The relative size
52
53 of each ‘raindrop’ is not yet known. Nonetheless, the extent to which each theme influences a
54
55
56
57
58
59
60

child’s concept of pain is likely to be highly idiosyncratic, differing from one child to another and also changing over time. Second, it is unknown if other factors contribute to the composition of the puddle. The ‘blank’ raindrops in Figure 3 highlight that, for an individual child, there could be other factors involved that were not identified in these interviews. Third, how can changes in a child’s concept of pain ‘puddle’ be assessed? Conceptual change theory⁴⁶ suggests that the ‘puddle’ can theoretically change with education, however no research exists to confirm this. A child’s concept of pain assessment tool would be required to investigate this further. Further research would also be required to explore whether the findings relate to children below the age of 8 years, adolescents, adults and non-English speaking populations.

A key principle of clinical practice requiring future exploration is how patient care is appropriately tailored and individualized based on a child’s concept of pain. The language and other communication tools used in this study may be useful for assessing children prior to pain science education, to ensure care is individualized based on identified needs. The helpfulness of using a variety of media (e.g. drawings, videos, and text) appeared to relate more to a child’s level of comfort with the medium or level of engagement, rather than their age. Therefore, as part of a flexible approach to communicating with children, using a range of media is recommended. In particular, drawings can provide more information and/or more accurate information compared with narrative methods, and they can increase comfort, facilitate memory retrieval, and help children organize verbal reports.⁴⁷ Both the open-ended drawing and the diagram-labelling task provided helpful information and enhanced other verbal responses in this study. Several participants commented that they enjoyed discussing the vignettes provided, which supports the established benefits of using vignettes clinically and in qualitative research.⁴⁸

⁴⁹ Because a child's concept of pain is not always overtly communicated by the child, particularly the 'hidden' emotional components which affect communication,⁵⁰ using vignettes and drawings in pain science education or assessments may be particularly beneficial in clinical practice. In addition to these future clinical directions, exploring the effects of different pain durations and pain diagnoses on a child's focus on emotional aspects of pain is warranted.

A further potential clinical implication is that a more flexible and individualized assessment of a child's concept of pain could directly inform what pain science education is likely to be most beneficial for a child. Pain science education for adults is based on conceptual change theory where the assumption is that misconceptions exist.¹⁴ The results of the current study show that complex concepts are still developing in childhood, and so pain science education for children should also incorporate educational approaches where concepts are taught with gradually increasing complexity built on previous conceptual development.⁵¹ Tailored, individualised pain science education, specifically aimed at addressing the different variables that have made up an individual child's concept of pain 'puddle' (Figure 3), appears warranted. Unexplored variables may be impacting on a child's concept of pain, such as their intellectual and emotional intelligence, their general attitude, fears, social circumstances, perspectives on 'toughness', and other aspects of their history. Each individual child may benefit from tailored education provided in different ways. Some children may not require any formal pain science education, but rather benefit most from reassurance from a trusted adult with influence such as a parent,⁵² teacher or health professional. Education deemed too complex by a child may result in misunderstanding and fear,⁵³ which could potentially aid in the development and/or maintenance of persistent pain and disability.⁵⁴

The terminology and understanding of neuroanatomy used in descriptions of a child’s concept of pain is also important given the use of neuroanatomical terms in pain science education.^{13 14} In the current study, children typically conceptualized the nervous system without inclusion of the spinal cord and considered it to have two parts; the brain and a peripheral component, and uncertainty existed regarding how these two parts are connected. Phrases such as ‘spinal cord’, ‘nerves’, and ‘real’, are likely to be poorly understood and potentially misinterpreted by children. Participants appeared to grasp the concept of the brain receiving warning messages and, based on these messages and many other factors, producing pain to keep the body safe. It is critical to carefully consider language when discussing pain with children,⁴¹ as younger participants in this age group tended to only use and understand simpler terminology. Age-appropriate language used in conversations by health professionals may facilitate the therapeutic alliance. For example, if a child uses a particular word or phrase to describe their concept of pain, a patient-centred empathetic communication strategy⁵⁵ would be for the clinician to incorporate the child’s language in the discussion rather than using technical terms. In addition, the apparent lack of awareness regarding how sensory input and distraction can influence pain provides some support for educating children at an age-appropriate level about biological mechanisms underpinning pain.

This study has several strengths and limitations. One strength of this study was the use of drawing tasks and vignettes as a means of data triangulation.⁵⁶ Another strength was the large variations in pain experience amongst the group of 8 to 12-year old participants with persistent pain; some participants had lifelong permanent physical disabilities whereas others had recurrent

headaches for just over one year, capturing individual variability in pain experiences.⁵⁷ A limitation of this study is the potential lack of transferability. This study only included children from New South Wales, Australia who spoke English. It is unknown if the results would be transferable to other cultures and age groups. A further limitation is that other 'raindrops' in the proposed conceptual framework could exist that were not identified and explored in these interviews and would be suitable to investigate in future research.

In conclusion, Australian children aged 8-12 years who have not received formal pain science education had concepts of pain that depended on a child's knowledge and experiences. The importance of gaining an in-depth understanding of a child's previous pain-related experiences and knowledge is emphasized, to facilitate clear and meaningful pain science education. The use of age-appropriate language, in combination with assessment and education tasks such as drawing and discussing vignettes, allowed participants with persistent pain to communicate their individual concept of pain. Future research quantitatively assessing a child's concept of pain is needed to improve and target pain science education for children with pain.

REFERENCES

1. Friedrichsdorf S, Giordano J, Desai Dakoji K, et al. Chronic pain in children and adolescents: diagnosis and treatment of primary pain disorders in head, abdomen, muscles and joints. *Children* 2016;3(4):42.

2. King S, Chambers CT, Huguet A, et al. The epidemiology of chronic pain in children and adolescents revisited: a systematic review. *Pain* 2011;152(12):2729-38. doi: 10.1016/j.pain.2011.07.016

3. Swain MS, Henschke N, Kamper SJ, et al. An international survey of pain in adolescents. *BMC Public Health* 2014;14(1):447.

4. Access Economics. The high price of pain: the economic impact of persistent pain in Australia. *Sydney: MBF Foundation* 2007:9-11.

5. Hunfeld JA, Perquin CW, Duivenvoorden HJ, et al. Chronic pain and its impact on quality of life in adolescents and their families. *J Pediatr Psychol* 2001;26(3):145-53.

6. Hassett AL, Hilliard PE, Goesling J, et al. Reports of chronic pain in childhood and adolescence among patients at a tertiary care pain clinic. *J Pain* 2013;14(11):1390-97.

7. Lisman-van Leeuwen Y, Spee LA, Benninga MA, et al. Prognosis of abdominal pain in children in primary care—a prospective cohort study. *Ann Fam Med* 2013;11(3):238-44.

8. Kosola S, Mundy LK, Sawyer SM, et al. Pain and learning in primary school: a population-based study. *Pain* 2017;158(9):1825-30. doi: 10.1097/j.pain.0000000000000984

9. Zernikow B, Wager J, Hechler T, et al. Characteristics of highly impaired children with severe chronic pain: a 5-year retrospective study on 2249 pediatric pain patients. *BMC Pediatr* 2012;12(1):54.

10. Coffelt TA, Bauer BD, Carroll AE. Inpatient characteristics of the child admitted with chronic pain. *Pediatrics* 2013;132(2):e422-e29.
11. Groenewald CB, Essner BS, Wright D, et al. The economic costs of chronic pain among a cohort of treatment-seeking adolescents in the United States. *J Pain* 2014;15(9):925-33.
12. Kamper SJ, Apeldoorn A, Chiarotto A, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain: Cochrane systematic review and meta-analysis. *BMJ* 2015;350:h444.
13. Harrison LE, Pate JW, Richardson PA, et al. Best-Evidence for the Rehabilitation of Chronic Pain Part 1: Pediatric Pain. *Journal of Clinical Medicine* 2019;8(9):1267.
14. Moseley GL, Butler DS. Fifteen Years of Explaining Pain: The Past, Present, and Future. *J Pain* 2015;16(9):807-13. doi: 10.1016/j.jpain.2015.05.005
15. Louw A, Zimney K, Puentedura EJ, et al. The efficacy of pain neuroscience education on musculoskeletal pain: A systematic review of the literature. *Physiother Theory Pract* 2016;32(5):332-55. doi: 10.1080/09593985.2016.1194646
16. Driessnack M. Children's drawings as facilitators of communication: a meta-analysis. *J Pediatr Nurs* 2005;20(6):415-23.
17. Robins H, Perron V, Heathcote L, et al. Pain neuroscience education: state of the art and application in pediatrics. *Children* 2016;3(4):43.
18. Harbeck C, Peterson L. Elephants dancing in my head: a developmental approach to children's concepts of specific pains. *Child Dev* 1992;63(1):138-49.
19. Jacobson CJ, Jr., Kashikar-Zuck S, Farrell J, et al. Qualitative Evaluation of Pediatric Pain Behavior, Quality, and Intensity Item Candidates and the PROMIS Pain Domain

Framework in Children With Chronic Pain. *J Pain* 2015;16(12):1243-55. doi: 10.1016/j.jpain.2015.08.007

20. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19(6):349-57. doi: 10.1093/intqhc/mzm042

21. Kvale S. *InterViews*. Thousands Oaks: Ca: Sage, 1996.

22. Liamputtong P. *Qualitative research methods*. 3rd ed. Victoria, Australia: Oxford University Press 2009.

23. Arbuckle R, Abetz-Webb L. "Not just little adults": qualitative methods to support the development of pediatric patient-reported outcomes. *Patient* 2013;6(3):143-59. doi: 10.1007/s40271-013-0022-3

24. Christie D, Viner R. ABC of adolescence: Adolescent development. *BMJ* 2005;330(7486):301.

25. Catley MJ, O'Connell NE, Moseley GL. How good is the neurophysiology of pain questionnaire? A Rasch analysis of psychometric properties. *The journal of pain* 2013;14(8):818-27.

26. Butler D, Moseley G. *Explain pain supercharged*. Adelaide City West 2017

27. Pate JW, Hush JM, Hancock MJ, et al. A Child's Concept of Pain: An International Survey of Pediatric Pain Experts. *Children* 2018;5(1):12.

28. Docherty S, Sandelowski M. Focus on qualitative methods: Interviewing children. *Res Nurs Health* 1999;22(2):177-85.

29. Lobe B, Livingstone S, Olafsson K, et al. *Best Practice Research Guide: How to research children and online technologies in comparative perspective*: EU Kids Online 2008.

30. Kortessluoma RL, Hentinen M, Nikkonen M. Conducting a qualitative child interview: methodological considerations. *J Adv Nurs* 2003;42(5):434-41.
31. Ritchie J, Lewis J, Nicholls CM, et al. Qualitative research practice: A guide for social science students and researchers: Sage 2013.
32. TED-Ed. How does your brain respond to pain? - Karen D. Davis. 2014. YouTube (2:24 - 2:26). <https://www.youtube.com/watch?v=I7wfDenj6CQ> (accessed 1 Oct. 2017).
33. Popay J, Rogers A, Williams G. Rationale and standards for the systematic review of qualitative literature in health services research. *Qual Health Res* 1998;8(3):341-51.
34. Pink B. Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA), Australia, 2011. Technical Paper. *Australian Bureau of Statistics (ABS), ed Canberra: Australian Government* 2011:54-58.
35. Bonoti F, Leondari A, Mastora A. Exploring children's understanding of death: through drawings and the death concept questionnaire. *Death Stud* 2013;37(1):47-60.
36. Buckley CA, Waring MJ. Using diagrams to support the research process: Examples from grounded theory. *Qual Res* 2013;13(2):148-72.
37. QSR International Pty Ltd. NVivo Qualitative Data Analysis Software (Version 9): QSR International Pty Ltd Doncaster, Vic, Australia, 2006.
38. Tamm ME, Granqvist A. The meaning of death for children and adolescents: A phenomenographic study of drawings. *Death Stud* 1995;19(3):203-22.
39. Angell C, Alexander J, Hunt JA. 'Draw, write and tell': A literature review and methodological development on the 'draw and write' research method. *Early Child Res* 2015;13(1):17-28.

40. Ely EA. The experience of pain for school-age children: Blood, Band-Aids, and feelings. *Children's Health Care* 1992;21(3):168-76.

41. Johnson E, Bornman J, Tönsing KM. An exploration of pain-related vocabulary: implications for AAC use with children. *Augmentative and Alternative Communication* 2016;32(4):249-60.

42. Yeung K-K, Engle L, Rabel A, et al. It just feels weird!: a qualitative study of how children aged 10–18 years describe neuropathic pain. *Disabil Rehabil* 2017;39(17):1695-702. doi: 10.1080/09638288.2016.1207107

43. Bunzli S, Watkins R, Smith A, et al. Lives on hold: a qualitative synthesis exploring the experience of chronic low-back pain. *Clin J Pain* 2013;29(10):907-16.

44. Sim J, Madden S. Illness experience in fibromyalgia syndrome: a metasynthesis of qualitative studies. *Soc Sci Med* 2008;67(1):57-67.

45. Coakley R, Schechter N. Chronic pain is like... The clinical use of analogy and metaphor in the treatment of chronic pain in children. *Pediatric Pain Letter* 2013;15(1):1-8.

46. Posner GJ, Strike KA, Hewson PW, et al. Accommodation of a scientific conception: Toward a theory of conceptual change. *Sci Educ* 1982;66(2):211-27.

47. Gross J, Hayne H. Drawing facilitates children's verbal reports of emotionally laden events. *J Exp Psychol Appl* 1998;4(2):163.

48. Barter C, Renold E. 'I wanna tell you a story': exploring the application of vignettes in qualitative research with children and young people. *Int J Soc Res Methodol* 2000;3(4):307-23.

49. Jenkins N, Bloor M, Fischer J, et al. Putting it in context: the use of vignettes in qualitative interviewing. *Qualitative Research* 2010;10(2):175-98.

50. Hadjistavropoulos T, Craig KD, Duck S, et al. A biopsychosocial formulation of pain communication. *Psychol Bull* 2011;137(6):910.
51. Novak JD, Cañas AJ. The theory underlying concept maps and how to construct and use them. Pensacola: Institute for Human and Machine Cognition, 2008.
52. Evans S, Tsao JC, Lu Q, et al. Parent-child pain relationships from a psychosocial perspective: a review of the literature. *J Pain Manag* 2008;1(3):237.
53. Simons LE, Kaczynski KJ. The Fear Avoidance model of chronic pain: examination for pediatric application. *The Journal of Pain* 2012;13(9):827-35.
54. Zale EL, Ditre JW. Pain-related fear, disability, and the fear-avoidance model of chronic pain. *Current opinion in psychology* 2015;5:24-30.
55. Linton SJ, Flink IK, Nilsson E, et al. Can training in empathetic validation improve medical students' communication with patients suffering pain? A test of concept. *Pain Rep* 2017;2(3):e600.
56. Krefting L. Rigor in qualitative research: The assessment of trustworthiness. *Am J Occup Ther* 1991;45(3):214-22.
57. Denk F, McMahon SB. Neurobiological basis for pain vulnerability: why me? *Pain* 2017;158:S108-S14.

Table 1. Example questions, drawing activities, and vignettes in the interview script.

<i>Questions</i>	<ul style="list-style-type: none">• What do you think pain is?<ul style="list-style-type: none">◦ What can pain feel like? Any picture words you can think of?• What do you think happens inside your body for you to feel hurt/sore?<ul style="list-style-type: none">◦ Or “how do you think pain is made inside your body?”• What do you think of when you hear the word “injury”?• Can someone be injured but have no pain?• Can someone have pain but not be injured?<ul style="list-style-type: none">◦ Where did you learn about that?• The job of eyes is to see. The job of ears is to hear. The job of the nose is to _____. And if pain were to have a job, what do you think the job of pain would be?• Why does pain last a long time for some people and not for others?• Do you think pain is always real?
<i>Drawing activities</i>	<ul style="list-style-type: none">• I want you to draw whatever the word “pain” makes you think of.• Draw where you think your brain, spinal cord and nerves are with 3 different coloured pencils on the diagram.
<i>Vignettes</i>	<ul style="list-style-type: none">• What do you think is happening inside this man’s body (cartoon character in a video) when he puts his hand on the stove?<ul style="list-style-type: none">◦ Why doesn’t his hand just stay on the stove top?• Do you think this cartoon character can have pain without knowing about it?<ul style="list-style-type: none">◦ Or “...without being aware of it?”• Let’s pretend this character has had pain for a long time. If he feels sad, do you think this would change his pain?<ul style="list-style-type: none">◦ If yes: Do you think his pain would be more/less/ bigger/ smaller/ stronger/quieter?◦ How do you think he could make his pain feel different/better/worse?

Table 2. Demographics of the participants.

Characteristics		n (%)
<i>Gender</i>		
Male		9 (56)
Female		7 (44)
<i>Age (years)</i>		
8		4 (25)
9		3 (19)
10		4 (25)
11		3 (19)
12		2 (13)
<i>Persistent Pain</i>		
Yes		8 (50)
- Duration:		
0-1 year		0 (0)
1-2 years		5 (63)
3-5 years		2 (25)
6-10 years		1 (13)
11+ years		0 (0)
- Location:		
Headache only		2 (25)
Back only		2 (25)
Neck only		1 (13)
Multiple sites		3 (38)
No		8 (50)
<i>Interview Setting</i>		
Home		9 (56)
University		6 (38)
Hospital		1 (6)

Note: Characteristics may not sum to exactly 100% due to the effect of rounding.

Table 3. Quotations to illustrate the subthemes of the four main themes.

Themes and Subthemes	Quotations
1. My pain-related knowledge: - Varied knowledge of pain's purpose, pain-related anatomy, and pain mechanisms.	<p>"...To alert your body that there is a virus and you are sick or something." (Boy, 9, persistent pain)</p> <p>"...It's to know you've hurt your knee and you need to stop playing." (Boy, 8, persistent pain)</p> <p>"The nerves feel it, then they tell the brain, [draws line up to brain outside body] and then the brain tells the nerve maybe." (Boy, 9, persistent pain)</p> <p>"Umm because like your nerves send like... It's like a phone call to your brain saying "I just hurt myself" and then your brain will say, "oh like that's going to hurt" and you go aww!" (Girl, 10, pain-free)</p> <p>"Maybe [people with higher sensitivity] have more nerves? And some people might have a little less." (Boy, 11, pain-free)</p>
- Influences of age	<p>[When asked how a vignette character could make his persistent abdominal pain feel different] "...Umm well not eat as much food if he is eating a lot. Basically, go on a diet, and depending on if he is fit or not, get fit." (Boy, 10, persistent pain)</p>
2. Pain in the world around me: - Contextual influences on a child's concept of pain	<p>"My mum... she has... and my teacher... they might have to get knee replacements. My teacher had one. And my mum has to get one. And my neighbour got one." (Girl, 9, persistent pain)</p> <p>[When asked where he learnt about the role of the brain] "Well I learnt most of it off TV, and a little bit from school." (Boy, 12, persistent pain)</p>
- Perceptions of how other people respond to pain	<p>"... Some people might be more sensitive than others. Like, I guess some people who play like full contact sports they would be more tough because they are used to the punching and everything. And people who don't do any sports or who are really sensitive naturally – you just go like "that!" (touches hand to desk lightly) it really, really hurts." (Girl, 10, pain-free)</p> <p>"I guess it's just how you're born. And how your nerves work." (Girl, 10, pain-free)</p> <p>"Some people have stronger muscles... It's a matter between weakness and strength." (Boy, 11, persistent pain)</p> <p>[When asked about sensitivity] "It means umm.. more vulnerable".</p> <p>"...Umm well not really, because some people have a high pain tolerance. Because if Sarah, Mary and I did that, Mary and Sarah would be crying because they have low pain tolerances. Because I'm kind of tougher than them, I'm the oldest." (Girl, 11, pain-free)</p> <p>[When asked if different people feel the same amount of pain with the same hammer-hitting-thumb-injury] "We wouldn't feel the same amount of pain. Because you're older and I'm younger. I can't hold in the pain as much as you. So I'd feel more than you." (Boy, 8, pain-free)</p> <p>"Well, if you're like 96 or something you'd feel pain like kids... Because you're getting old." (Boy, 8, pain-free)</p> <p>"[Adults] may be more used to it, because they may have hit it before – depending how often! Because they may have had longer time to experience pain on their thumbs." (Boy, 10, persistent pain)</p>
3. Pain in me: - Experiences of any pain/injury	<p>"Well pain is like if you've hurt yourself, it's the physical contact and it hurts. Or like after surgeries you'd have like pain and it hurts to move." (Girl, 12, persistent pain)</p> <p>"It helps us not get into any more trouble... it'll help us to not do [an injury] again. We'll remember last time and [do the task] differently." (Boy, 8, pain-free)</p> <p>"Well if you really didn't like something, like a sport, then you don't have to play it. It'd be helpful maybe." (Girl, 10, pain-free)</p> <p>"If you had, like we have a presentation day that goes for 2 hours at school. If you didn't want to go, if you have pain you can stay home." (Girl, 9, persistent pain)</p>
- The impact of having persistent pain	<p>"You could go see a doctor" (Boy, 8, pain-free)</p> <p>"If they have pain then they can go to a physio..." (Boy, 8, pain-free)</p> <p>"If you maybe have a sore tummy, and you're like "oh no I might have appendicitis!" or something like that. And then you go to the doctors and then they're like "oh no you're just sick" or something like that. You're like "Oh thank goodness, I've only got a fever!" (Girl, 11, pain-free)</p> <p>"Umm... I reckon no... I always know about it [pain]." (Boy, 10, persistent pain)</p> <p>"If you defined pain in the dictionary... it would be something that physically hurts you. So it's a fake thing. Just a warning... It depends what you mean by real?" (Boy, 12, persistent pain)</p> <p>[When asked if pain was always 'real'] "I think it's always real. Because sometimes in your body, like right now when I'm talking, pain can still be in your body even if it doesn't hurt." (Boy, 8, pain-free)</p>
- The physical is obvious, but the emotional is hidden	<p>"I reckon if he [the vignette character] looked at like his favourite thing in the world, and got to play with it or just looked at it at least and just went like "I need it" – it would distract him from his pain because he is focussed on his favourite toy or something, in the world." (Boy, 10, persistent pain)</p>

4. Communicating my concept of pain:	[Pain is] "...when something hurts in your body" (Girl, 10, pain-free)
	"The body is fighting the virus or something." (Boy, 9, persistent pain)
- Verbal communication	
- Individual engagement	[Pointing to parts of the diagram whilst talking] "[Nerves] release chemicals that flow up to your brain. And then your brain sends down another chemical I think, that gives it pain." (Boy, 12, persistent pain)

For peer review only

1

2

3 **Figure Legends**

4

5

6

7

8 Figure 1. Examples of drawings by children when asked to draw whatever the word ‘pain’ makes

9 them think of. Panels (a) and (b) represent how children who were pain-free drew injuries (9

10 years, female; 10 years female). Panels (c) and (d) are examples of how children with persistent

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

pain drew more emotional elements such as hearts and tears (11 years, male; 10 years, female).

Figure 2. Eight examples of labelled diagrams of the brain, spinal cord and nerves by children

aged 8-12 years. The top row are drawings of children who were pain-free (8 years, male; 8

years, male; 10 years, female; 11 years, male) and the bottom row are drawings of children with

persistent pain (8 years, male; 12 years, male; 9 years, female; 11 years, male). No differences

between children with and without persistent pain were identified in this drawing task.

Figure 3. A proposed conceptual framework of the potential interactions between the themes

resulting from the analysis and a child’s concept of pain.

- Notes:*
- 1. The raindrops (influences on a child’s concept of pain) without labels highlight that all potential contributors to a child’s concept of pain may not have been identified; and*
 - 2. The size of the raindrops vary from one child to the next.*

Word Count: 5054 words

Data availability statement:

Deidentified transcripts and drawings, and the unpublished protocol, will be made available upon reasonable request from Joshua Pate (<https://orcid.org/0000-0002-1049-3916>) until 2024 as per ethical approval. Reuse without ethical approval is not permitted. A data-sharing agreement will require a commitment to using the data only for specified research purposes, to securing the data appropriately and to destroying the data after a nominated period.

Contributorship statement:

Conception and design: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Planning and implementation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Data collection: J.W.P.

Coding data for thematic analysis: J.W.P., T.N., and V.P.

Analysis and interpretation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Drafting the article: J.W.P.

Review and editing: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Overall responsibility: J.W.P. under the supervision of V.P., J.M.H., and M.J.H.




Figure 1




Figure 2



Figure 3. A proposed conceptual framework of the potential interactions between the themes resulting from the analysis and a child's concept of pain.

Notes:

1. The raindrops without labels highlight that all potential contributors to a child's concept of pain may not have been identified; and
2. The size of the raindrops vary from one child to the next.

Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

Developed from Table 1 of: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

No. Item	Guide questions/description	Reported on Page #
Domain 1: Research team and reflexivity		
<i>Personal Characteristics</i>		
1. Interviewer/facilitator	Which author/s conducted the interview or focus group?	Page 9
2. Credentials	What were the researcher's credentials? E.g. PhD, MD	Page 9
3. Occupation	What was their occupation at the time of the study?	Page 9
4. Gender	Was the researcher male or female?	Page 9
5. Experience and training	What experience or training did the researcher have?	Page 9
<i>Relationship with participants</i>		
6. Relationship established	Was a relationship established prior to study commencement?	Page 9
7. Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	Page 9
8. Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. bias, assumptions, reasons and interests in the research topic	Page 9
Domain 2: Study design		
<i>Theoretical framework</i>		
9. Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	Page 11
<i>Participant selection</i>		
10. Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	Page 8
11. Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	Page 8
12. Sample size	How many participants were in the study?	Page 11
13. Non-participation	How many people refused to participate or dropped out? Reasons?	Page 11
<i>Setting</i>		

14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	Table 2 (Page 31)
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	Page 10
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	Table 2 (Page 31)
<i>Data collection</i>		
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	Table 1 (Page 30)
18. Repeat interviews	Were repeat inter iews carried out? If yes, how many?	No, page 7
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	Page 10
20. Field notes	Were field notes made during and/or after the interview or focus group?	Page 10
21. Duration	What was the duration of the interviews or focus group?	Page 11
22. Data saturation	Was data saturation discussed?	Page 8 and 9
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	Page 10
Domain 3: analysis and findings		
<i>Data analysis</i>		
24. Number of data coders	How many data coders coded the data?	3 coders, Page 11
25. Description of the coding tree	Did authors provide a description of the coding tree?	Page 11
26. Derivation of themes	Were themes identified in advance or derived from the data?	Page 11
27. Software	What software, if applicable, was used to manage the data?	Page 11
28. Participant checking	Did participants provide feedback on the findings?	Page 10
<i>Reporting</i>		
29. Quotations presented	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	Table 3, page 32
30. Data and findings consistent	Was there consistency between the data presented and the findings?	Yes
31. Clarity of major themes	Were major themes clearly presented in the findings?	Yes
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	Yes

BMJ Open

Exploring the concept of pain of Australian children with and without pain: a qualitative study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-033199.R2
Article Type:	Original research
Date Submitted by the Author:	01-Oct-2019
Complete List of Authors:	Pate, Joshua; Macquarie University, Health Professions Noblet, Tim; Macquarie University, Health Professions Hush, Julia ; Macquarie University, Health Professions Hancock, Mark; Macquarie University, Health Professions Sandells, Renee; Children's Hospital at Westmead Pounder, Meg; Children's Hospital at Westmead Pacey, Verity; Macquarie University Faculty of Medicine and Health Sciences
Primary Subject Heading:	Paediatrics
Secondary Subject Heading:	Qualitative research, Communication, Patient-centred medicine
Keywords:	pain science education, concept of pain, paediatric pain, QUALITATIVE RESEARCH, drawing task

SCHOLARONE™
Manuscripts

Title:

Exploring the concept of pain of Australian children with and without pain: a qualitative study.

Authors:

Joshua W. Pate^{1,2}, MRes, Senior Physiotherapist and PhD Candidate, joshua.pate@mq.edu.au

Tim Noblet¹, MSc (MSKMed), Senior Physiotherapist and PhD Candidate,
tim.noblet@mq.edu.au

Julia M. Hush¹, PhD, Professor of Physiotherapy, julia.hush@mq.edu.au

Mark J. Hancock¹, PhD, Professor of Physiotherapy, mark.hancock@mq.edu.au

Renee Sandells², MPsych (Clin), Clinical Psychologist, renee.sandells@health.nsw.gov.au

Meg Pounder², BSc (Physiotherapy), Senior Physiotherapist, meg.pounder@health.nsw.gov.au

Verity Pacey^{1,3}, PhD, Associate Professor of Physiotherapy, verity.pacey@mq.edu.au

Affiliations:

¹ Macquarie University, New South Wales, Australia

² Department of Pain Medicine, The Children’s Hospital at Westmead, New South Wales,
Australia

³ The University of Sydney Children’s Hospital Westmead Clinical School, New South Wales,
Australia

Corresponding Author:

Joshua Pate

Ground floor 75 Talavera Road

Macquarie University 2109

Phone: +61 2 9850 6626

Fax: +61 2 9850 6630

Email: joshua.pate@mq.edu.au

Institutional URL: <https://researchers.mq.edu.au/en/persons/joshua-pate>

Keywords:

concept of pain; pain science education; paediatric pain; qualitative interviews; drawing task

Previous presentations of the research:

Poster of one subtheme of the manuscript presented for IASP 2018 in Boston, titled: “Explaining pain to children with and without pain: what terminology should we use? Qualitative interviews describing pain-related neuroanatomy.”

Acknowledgments:

We thank all the children and their families for their openness and for giving their valuable time, energy and efforts to participate in this study. We also acknowledge Natasha Haynes, Nurse Practitioner from The Children’s Hospital at Westmead Pain Team, who helped to recruit participants and supported the study.

Funding:

J.W.P. is a Ph.D. Student funded by a Macquarie University Research Training Program (RTP) full-time scholarship.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflicts of Interest:

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/doi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Abstract:

Objective: A person's concept of pain can be defined as how they understand what pain actually is, what function it serves, and what biological processes are thought to underpin it. This study aimed to explore the concept of pain in children with and without persistent pain.

Design: In-depth, face-to-face interviews with drawing tasks were conducted with 16 children (aged 8-12 years) in New South Wales, Australia. Thematic analysis was used to analyse and synthesize the data.

Setting: Children with persistent pain were identified from a pain clinic waiting list in Australia, and children without pain were identified through advertising flyers and email bulletins at a university and hospital.

Participants: Eight children had persistent pain and eight children were pain-free.

Results: Four themes emerged from the data: 'my pain-related knowledge', 'pain in the world around me', 'pain in me', and 'communicating my concept of pain'. A conceptual framework of the potential interactions between the themes resulting from the analysis is proposed. The concept of pain of Australian children aged 8-12 years varied depending on their knowledge, experiences and literacy levels. For example, when undertaking a drawing task, children with persistent pain tended to draw emotional elements to describe pain, whereas children who were pain-free did not.

Conclusions: Gaining an in-depth understanding of a child's previous pain-related experiences and knowledge is important to facilitate clear and meaningful pain science education. The use of age-appropriate language, in combination with appropriate assessment and education tasks such as drawing and discussing vignettes, allowed children to communicate their individual concept of pain.

Article Summary:
Strengths and limitations of this study:

- One strength of this study was the use of drawing tasks and vignettes in the face-to-face interviews as a means of data triangulation.
- Another strength was the large variations in pain experience amongst the group of 8 to 12-year old children with persistent pain; some participants had lifelong permanent physical disabilities whereas others had recurrent headaches for just over one year, capturing individual variability in pain experiences.
- A limitation of this study is the potential lack of transferability. This study only included children from New South Wales, Australia who spoke English. It is unknown if the results would be transferable to other cultures and age groups.
- A further limitation is that within the proposed conceptual framework, other influences on a child’s concept of pain could exist, however they were not identified and explored in these interviews.

Introduction:

Estimates of point-prevalence of persistent pain (pain that extends beyond the expected period of healing¹) throughout childhood range from 11% to 54%,^{2,3} which is comparable to that for adults.⁴ Persistent pain in children has detrimental effects on a child's quality of life⁵ and is a significant socioeconomic and health problem. For example, persistent pain in childhood could predict persistent pain in adulthood⁶ and is associated with increased pain-related disability,² school absence,⁷ poorer reading levels,⁸ and emotional distress.⁹ An increasing number of children are being admitted to hospitals for treatment of persistent pain¹⁰ and health service costs are high.^{11,12}

Multidisciplinary pain management is currently the gold standard for persistent pain management.¹³ An important part of the multidisciplinary approach is pain science education, aiming to change a person's concept of pain and to improve pain and function by teaching the underlying biopsychosocial mechanisms of pain.¹⁴ A person's concept of pain can be defined as how they understand "what pain actually is, what function it serves, and what biological processes are thought to underpin it";¹⁵ these can be summarized as the 'what, why and how of pain'. Clinically, a range of methods for educational assessments and treatments are used for adults with persistent pain including drawing tasks.¹⁶ Despite drawing tasks being a robust interview strategy with children,¹⁷ these methods have not been applied to paediatric pain research to investigate a child's concept of pain.¹⁸

Limited research has been conducted that focuses on a child's concept of pain. For example, children who are pain-free have been asked to describe sensations relating to a grazed knee, an

injection, and a headache.¹⁹ In addition, pain-specific behaviours, pain quality, and pain intensity have been investigated in children with pain.²⁰ However, a child’s concept of pain, as defined in the previous paragraph, has not been specifically investigated in children with or without persistent pain. Therefore, this remains poorly understood, providing clinicians little guidance on how to provide pain science education to children with persistent pain as part of multidisciplinary management, or to children who are pain-free as a strategy to prevent future pain, using language which is clear and unambiguous to a child. The aims of this study were (1) to explore the concept of pain of children diagnosed with persistent pain, and (2) to qualitatively compare the concept of pain between children who have pain with those who are pain-free.

Methods:

The study was approved by the Human Research Ethics Committees of Sydney Children’s Hospital Network (Ref: LNR/17/SCHN/268) and Macquarie University (Ref: 5201700820). All participants’ parents provided informed consent prior to participation in the study, and verbal assent was obtained from the participants prior to the commencement of the interview.

Study Design

This qualitative study utilized one-on-one interviews, lasting approximately 30-minutes, with children aged 8-12 years. To ensure explicit and comprehensive study reporting, the widely used Consolidated Criteria for Reporting Qualitative Health Research (COREQ)²¹ was implemented. Children were asked to answer questions and complete activities relating to their concept of pain. By adapting grounded theory and utilizing thematic analysis,^{22 23} concepts in the transcripts were inductively identified and recorded and then a coding structure was iteratively developed and

refined until the aims of the study were achieved. Themes and sub-themes were developed from data analysis and synthesis.

Participants

Two groups of children aged 8-12 years in the state of New South Wales, Australia, were purposefully recruited; children who had persistent pain (greater than 3 months) and children who were pain-free and healthy by parent-report. Parent-report was used to minimise burden on the children being interviewed. The absence, presence, and/or persistence of pain was later confirmed throughout the interviews, allowing opportunity to clarify any discrepancies between reporting. Children were excluded from the study if they (1) had received pain science education from a health care professional, (2) were deemed not medically fit to be part of this research study by their treating physician/surgeon, or (3) were unable to participate in a 30-minute interview based on parent-report.

A minimum age for participants was set at 8 years, the youngest age a child can typically self-report and provide meaningful responses in interviews due to their linguistic and cognitive development.²⁴ The maximum age of participation was 12 years, an age prior to when abstract thinking capacities typically develop in the teenage years.²⁵ The age range was determined to meet a clinical need highlighted by research team members to provide improved assessment and management resources for pre-teenage children. A pragmatic approach incorporating age limits, rather than a formal assessment of cognitive development or language abilities, allows for direct applicability in time-limited clinical settings.

A purposive selection strategy was used to identify children of varying demographics and pain characteristics who were likely to meet the eligibility criteria. For all children, a mix of gender, age, and ethnic backgrounds was sought to capture diversity of experiences. For the children with persistent pain, purposive sampling attended to capturing variations in the duration and location of pain. Children were consecutively identified from The Children’s Hospital at Westmead, Australia Complex Pain Service waiting list, and through advertising flyers and email bulletins at Macquarie University and The Children’s Hospital at Westmead.

Children were interviewed until data saturation was reached. Data saturation was defined as when no additional new information was attained with additional interviews regarding the final themes. The three researchers analysing the data (J.W.P., V.P., and T.N.) met fortnightly throughout data collection and the thematic analysis to determine when data saturation was reached. Data collection was stopped when all three researchers agreed that saturation had taken place.

Interview Script

An interview script was developed (Table 1) to investigate a child’s concept of pain based on seven domains (e.g. ‘How pain works’). These domains were proposed in a recent survey of paediatric pain experts by categorizing items from the revised Neurophysiology of Pain Questionnaire (rNPQ)²⁶ and Explain Pain Target Concepts,²⁷ and they were rated as relevant and appropriate.²⁸ The interview script aligned with guidelines for interviewing children,²⁹⁻³¹ and strategies supported by these guidelines were used in the interviews. For example, children were asked to choose their three favourite coloured pens as an icebreaker activity; these pens were

then used for the drawing activities. In addition, vignettes with familiar media and cartoon characters were used to increase engagement.³² For example, an open-ended question of “How does pain work?” was followed by a child-friendly cartoon video accessible on YouTube³³ and children were asked, “Which body part decides when you will feel pain?”. This question was explained, and clarification of understanding was ensured by asking response-dependent follow-up questions such as “What else does your brain do?”.

The team of investigators (including physiotherapists, clinical psychologists, and experts in methodology) provided feedback regarding script development. Pilot testing on three healthy pain-free children (aged 7, 8 and 10 years), discussions with education and linguistics experts, and further review by the investigators, resulted in re-wording of several questions to ensure that the questions and language were appropriate for children aged 8-12 years. For example, clarifying questions using synonyms (such as “replace the word ‘helpful’ with ‘useful’ if needed”, and ask “What word could we use instead of that?”) were incorporated. Following an iterative process, appropriate terminology and language resulting from interviews informed the focus of subsequent interviews until data saturation was reached.

Data Collection

Semi-structured interviews were all conducted by J.W.P. (male physiotherapist, a PhD candidate, and no previous relationship with potential participants). To maximize feelings of physical and emotional comfort, interviews took place at a location chosen by the participant (The Children’s Hospital at Westmead, Macquarie University, or in participants’ homes). Parents/caregivers (from now on referred to as “parents”) were invited to be present for each interview but no

questions were asked of them.³⁰ Parents were asked not to contribute unless they felt that their child was upset or appeared distressed, at which point the interview was to be terminated and an offer to speak to the clinical psychologist would be made.

A PowerPoint presentation displayed prompts on a laptop screen to guide the discussion. The interviews were audio recorded to enable transcription. Field notes of contextual details and non-verbal expressions were recorded for data analysis and interpretation. For respondent checking, a summary of the interview transcript was provided to families for comment and feedback to enhance the analytical framework,³⁴ and a full copy of the transcript was available upon request.

A range of demographic characteristics, including age, sex, ethnicity, school year, postcode [for socioeconomic status: using the Australian Socio-Economic Indexes For Areas (SEIFA) ³⁵], parental education level and working status were collected in a demographics questionnaire completed by parents. For children with persistent pain, parent-reported location and duration of pain were also collected before the commencement of the interview.

Data triangulation (using multiple methods to enhance understanding) was achieved by including a drawing task during the interviews.¹⁷ For this drawing task, children were asked to “draw whatever the word ‘pain’ makes you think of”. This follows methodology of a study investigating the concept of death in children.³⁶ To explore the language children use to describe components of the nervous system and their familiarity with the anatomical components of this system, a second written task involved labelling a diagram of their nervous system (brain, spinal cord and nerves).³⁷ If a child had difficulty with this, they were asked to verbally label the

1
2
3 diagram, or point to where their own brain, spinal cord and nerves are on their own body.
4
5
6

7 8 *Analysis*

9
10 Audio recordings from interviews were transcribed by J.W.P verbatim. V.P. and T.N. checked
11 transcripts for accuracy. An adapted grounded theory approach²³ was used to analyse and
12 synthesize the data. Data were independently coded line-by-line by three researchers (J.W.P.,
13 V.P., and T.N.) using NVivo software.³⁸ Drawings were thematically analysed by the
14 investigators as has been done in previous research.^{39 40} Analytical themes were inductively
15 developed and fully agreed upon by these three researchers through a rigorous iterative process.
16 The themes were scrutinised by a team of subject experts including paediatric clinical
17 psychologists, physiotherapists, academics and methodological experts, in two meetings during
18 the analytical process and one final meeting to attain 100% consensus.
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 *Public and Patient Involvement*

34
35 Patients were involved in piloting the interview script to identify the most appropriate language
36 to be used in the interviews and the time required to participate. The thematic analysis was
37 directly informed by participants as it was an iterative process. Patients and the general public
38 were not involved in informing other stages of the research process.
39
40
41
42
43
44
45
46

47 **Results:**

48
49 Information sheets were distributed and 16 recruitment emails in total were sent to parents of
50 children who contacted the investigators stating that they wanted to participate. All families
51 replied to the invitation and consented to participate. Therefore, 16 participants, 8 with persistent
52
53
54
55
56
57
58
59
60

pain and 8 children who were pain-free, were enrolled in the study and completed an interview between October 2017 and February 2018 before data saturation was reached. The median and interquartile range (IQR) interview duration was 33 (27-36) minutes. The three researchers (J.W.P., V.P., and T.N.) coding the transcripts agreed on the correctness of the transcripts, with no disagreements identified.

Participants

The participants included 9 males and 7 females, with a median (IQR) age of 10.0 (8.8-11.0) years. All participants identified themselves as Australian, with 7 participants reporting a second ethnicity including Chinese, Lebanese, Indian, Dutch and Eurasian. The median (IQR) Index of Relative Socio-economic Advantage and Disadvantage (IRSAD) SEIFA scores (Likert scale of 1-5 rating from disadvantaged to advantaged) was 4 (2.8-5). Of participants with persistent pain, five children in total reported back pain (63%), two children reported headaches (25%), and one child reported neck pain (13%). Pain duration ranged from 1-10 years with two participants reporting lifelong permanent physical disabilities. No discrepancies between parent- and child-reports of the participant’s pain history were identified. Table 2 details the demographic characteristics of participants. No participants reported feeling upset or appeared distressed during the interviews.

Key themes

Four themes emerged from the data: ‘my pain-related knowledge’, ‘pain in the world around me’, ‘pain in me’, and ‘communicating my concept of pain’. The first three themes focus on the

content that participants conveyed, whereas the fourth theme focuses on the method they used to convey this content. Illustrative quotations for each theme are provided in Table 3.

Theme 1: My pain-related knowledge

Participant knowledge of the purpose of pain, pain-related anatomy, and pain mechanisms varied between individuals, while participants consistently used concrete terms to describe their pain-related knowledge. Age influenced the participants' ability to express their concept of pain with younger participants responding with uncertainty at times. Based on the thematic analysis, no differences in participant knowledge were identified between participants who had persistent pain and participants who were pain-free, nor based on their gender or ethnic background.

Subtheme: Varied knowledge of pain's purpose, pain-related anatomy, and pain mechanisms.

When asked if pain had 'a job in the body', to identify pain's purpose, participants used familiar terms and referred to the idea of pain keeping them safe by protecting, warning, alerting, or knowing. The words 'injury' and 'pain' were both commonly associated with the word 'hurt', resulting in some participants expressing confusion about pain's purpose. When asked to 'draw whatever the word "pain" makes you think of', the participants' drawings frequently included an injury (Figure 1). Participants used a variety of examples for injuries in their drawings and explanations, such as breaking an arm or leg, having a sprained ankle, or having fallen over.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Regarding pain-related anatomy, all participants noted the importance of the brain and correctly labelled the brain inside the head on the diagram (Figure 2). A range of understanding was demonstrated in this task and no differences were identified between participants who had persistent pain and participants who were pain-free. Participants commonly stated that the brain controls when and how someone feels pain. Exceptions to this were responses that were vague such as "the head", "there is no part", or they were unsure. The spinal cord was rarely included when explaining key elements of the nervous system involved in pain; participants frequently drew this in the back disconnected from the brain or were not sure where it was located when questioned. Participants of each age varied in their understanding of the nervous system. Varied understanding of the term ‘nerves’ was observed; whilst some participants said that nerves are "everywhere", others confused nerves with feelings of nervousness and some participants confused nerves with structures such as veins. No child identified interconnections within the nervous system, however when directly asked, approximately half of each group said the brain, spinal cord and nerves are connected.

Knowledge about pain mechanisms also varied. To explain how pain is experienced, participants frequently reported ideas of “sending messages” or “telling the brain”. Metaphors were also used to explain the physiological processes underpinning pain, such as “it’s like a phone call to your brain”. Some participants referred to ideas of sensitivity or pain tolerance to justify that everyone's pain is different. When participants were asked if they would feel the same or different pain if a cartoon character, the investigator (J.W.P.), and the participant all accidentally hit their thumbs with a hammer using the same force, most participants identified that each

would feel a different pain. Participants provided explanations for this such as being inherently different people, having different levels of ‘toughness’ or different ages.

Subtheme: Influences of age

Younger participants tended to use simpler terminology. Some participants aged 8 and 9 years more frequently responded “I don’t know” and appeared less confident of their responses, speaking softly when uncertain. In contrast, older participants tended to use more complex terminology, such as phrases like “physically harming”. All participants in the study appeared to focus more on concrete ideas than on abstract ideas, such as pain having contextual influences. For example, when asked why pain persists for some people and not for others, the most common response was that the person was not caring for an injury correctly.

Participants’ understanding of pain-related abstract concepts was influenced by their age, reflecting this age group’s stage of cognitive development. Reciting words from a question was seen as distinctly different from understanding a concept. For example, when given a vignette of someone with persisting pain without an injury and asked about pain management strategies, participants responded by suggesting management strategies directly aligned to ‘fixing’ potential causes of the pain (“not eat as much food if he is eating a lot” and “depending on if he is fit or not, get fit” – Table 3).

Theme 2: Pain in the world around me

A child’s concept of pain is strongly influenced by their environmental and social context, including perceptions of how other people respond to pain. Consequently, differences in the concept of pain between participants with and without persistent pain were evident.

Subtheme: Contextual influences on a child’s concept of pain

A child’s exposure to family, school, and media, appeared to influence their concept of pain, to varying extents between individuals. Participants reported they learnt about pain from family members, particularly parents, as well as at school and via media on television and the internet. To explain various aspects of their concept of pain, participants with persistent pain also used examples of their interactions with health professionals discussing their pain experience despite not having previously received specific pain science education. In contrast, participants who were pain-free only used examples of family members and people at school.

Subtheme: Perceptions of how other people respond to pain

Participants with and without persistent pain considered other people to be ‘tough’ or ‘weak’ based on their responses to painful experiences. People who felt a lot of pain were perceived by participants as being weaker. People who ignored their pain were perceived as being tougher. The social stigma surrounding pain was also highlighted in phrases describing pain sensitivity. For example, some participants viewed age as a determinant of pain sensitivity, generally stating that ‘toughness’ increases as a person ages until they “get old” when it decreases significantly.

Other participants considered ‘toughness’ to be an inherent quality, whilst others did not and emphasized “getting stronger” decreases sensitivity to pain.

Theme 3: Pain in me

A child’s concept of pain was based upon their individual experiences with pain and injury. All participants clearly described physical aspects of pain, whereas emotional aspects were relatively hidden and were revealed only by participants with persistent pain upon further verbal probing or drawing tasks. The impact of persistent pain was highly individualized and related deeply to their past experiences.

Subtheme: Experiences of any pain/injury

Participants with and without persistent pain explained their concept of pain by using examples of pain and injury from their own experiences. Regarding the purpose that pain serves, most participants reported that pain is sometimes a helpful thing to feel. The most common reason for this was a warning to avoid further problems, however, two participants (one with persistent pain and one without) described the idea of pain being helpful to avoid attending something deemed boring or undesirable such as “a sport” or “presentation day”.

Subtheme: The impact of having persistent pain

Participants with persistent pain expressed different aspects of their concept of pain compared to participants without pain, with these aspects being dependent on their individual pain experiences. Participants with persistent pain considered broader aspects of the pain experience when describing what ‘pain’ is, such as the impact of surgery, effects of feelings, the idea of healing, and the diverse influences of health professionals.

When participants were provided with a vignette and asked if a character could have pain without knowing about it, participants were generally uncertain with some responding purely based on their own experience without reference to the vignette. There appeared to be a difference in explanations between participants with and without persistent pain; participants who were pain-free responded with regards to the vignette only whereas participants with persistent pain used the vignette with reference to their own personal experience.

Participants with persistent pain focused more on emotional aspects of pain than participants who were pain-free. When undertaking a drawing task, participants with persistent pain tended to draw emotional elements to describe pain, whereas participants who were pain-free did not (Figure 1). Some participants were confused when questioned about aspects of their drawing that illustrated emotional pain, mental pain, or nightmares, and were unable to verbally communicate how these types of pain related to a person’s overall pain experience.

Further investigating what children think pain is, the discussion around “Is pain real?” varied between participants with and without persistent pain. While some participants in both groups described some confusion around the term ‘real’ when asked this question, most participants

with persistent pain said that pain is not always real. In contrast, most participants who were pain-free said that pain is always real. Having persistent pain appeared to influence views on the 'reality' of pain.

Subtheme: The physical is obvious, but the emotional is hidden

Participants tended to focus their discussions on physical and visible components of pain-related processes. The most obvious example of this was mentioned previously; that participants without persistent pain did not draw emotional elements of pain.

When participants discussed possible mechanisms for how different factors can influence pain, most participants with persistent pain considered their experiences and stated that they could use self-taught management techniques, such as distracting themselves from pain by listening to music. In stark contrast, participants who were pain-free did not mention cognitive strategies such as distraction. Other than attentional strategies, participants did not mention any other factors that can influence the experience of pain. All children interviewed appeared unsure about the possible influence of other senses on pain, for example visual input.

Theme 4: Communicating my concept of pain

Participants used verbal descriptions, drawings and diagrams to describe their concept of pain with drawing tasks and vignettes aiding some participants' communication during the interviews. No differences in the use of these methods of communicating were identified between participants who had persistent pain and participants who were pain-free.

Subtheme: Verbal communication

Participants’ verbal descriptions of their concept of pain were individual and varied. For example, some participants described the physiological processes underpinning pain using terms such as ‘chemicals’ whereas others used terms such as ‘virus’ and ‘disease’. The words ‘pain’, ‘injury’ and ‘hurt’ were used interchangeably by multiple participants. No differences between groups were identified for this transposable use of terms.

Subtheme: Individual engagement

The value of the drawing tasks and vignettes appeared to relate to levels of engagement with the task, and this value did not appear related to age, gender or ethnic background. The drawing task was very helpful in enabling some participants to articulate thoughts they had not verbally communicated. Asking the follow-up question, “why did you draw that?” gave participants an opportunity to expand on their thoughts. For example, some participants did not draw anything when first asked, but later drew representations, such as a face with tears. In contrast, four participants did not like the drawing task and did not feel comfortable with this activity, preferring to speak about it and leaving the drawing page mostly blank (8 years, male, pain-free; 12 years, male, persistent pain; 11 years, female, pain-free; 8 years, male, pain-free).

Proposed conceptual framework

The potential interactions between the themes resulting from the analysis and a child's concept of pain form a proposed conceptual framework that is visually presented in Figure 3.

The first three themes are inputs and combine in varying proportions to produce the output which is how a child communicates their concept of pain.

Discussion:

This study is the first known to the authors to investigate a child's concept of what pain is, why it exists, and how it is experienced. The primary objective of this study was to gain insight into the way that children aged 8-12 years, who have not received pain science education, conceptualize pain. Four themes emerged from the data: 'my pain-related knowledge', 'pain in the world around me', 'pain in me', and 'communicating my concept of pain'. Participants' communication of their concept of pain was based on their individual knowledge and experience. Participants with persistent pain considered broader aspects of pain and focused more on emotional impacts and influences than participants who were pain-free.

Children's experiences of pain have previously been reported in qualitative studies. As examples, one study considered children's descriptions of types of pain¹⁹, another focused on painful experiences⁴¹, another on children's pain-related vocabulary⁴², and a recent study of children aged 10-18 years with neuropathic pain considered personal pain experiences and language used.⁴³ Similarly in studies of adults, pain experiences have been explored.^{44 45} These studies differed from the present study in that the focus was on the experience of pain, rather than a person's concept of pain. Children's conceptual frameworks have been studied using drawings with the concept of death,³⁶ but the present study is the first to apply drawing methodology to

investigate the concept of pain. Metaphors and explanations about pain that clinicians use with children have been discussed in the literature,⁴⁶ but the metaphors and explanations that children themselves use have not been explored.

Using the conceptual framework in Figure 3, several important issues are raised for future research regarding clinical practice. First, it is currently unknown the extent to which each of the themes impacts upon a child’s concept of pain. Does a child’s experience have more of an impact on their concept of pain than the impact of their knowledge or emotions? The relative size of each ‘raindrop’ is not yet known. Nonetheless, the extent to which each theme influences a child’s concept of pain is likely to be highly idiosyncratic, differing from one child to another and also changing over time. Second, it is unknown if other factors contribute to the composition of the puddle. The ‘blank’ raindrops in Figure 3 highlight that, for an individual child, there could be other factors involved that were not identified in these interviews. Third, how can changes in a child’s concept of pain ‘puddle’ be assessed? Conceptual change theory⁴⁷ suggests that the ‘puddle’ can theoretically change with education, however no research exists to confirm this. A child’s concept of pain assessment tool would be required to investigate this further. Further research would also be required to explore whether the findings relate to children below the age of 8 years, adolescents, adults and non-English speaking populations.

A key principle of clinical practice requiring future exploration is how patient care is appropriately tailored and individualized based on a child’s concept of pain. The language and other communication tools used in this study may be useful for assessing a child’s concept of pain prior to pain science education, to ensure care is individualized based on identified needs.⁴⁸

⁴⁹ The helpfulness of using a variety of media (e.g. drawings, videos, and text) appeared to relate more to a child's level of comfort with the medium or level of engagement, rather than their age. Therefore, as part of a flexible approach to communicating with children, using a range of media is recommended. In particular, drawings can provide more information and/or more accurate information compared with narrative methods, and they can increase comfort, facilitate memory retrieval, and help children organize verbal reports.^{50 51} Both the open-ended drawing and the diagram-labelling task provided helpful information and enhanced other verbal responses in this study. Several participants commented that they enjoyed discussing the vignettes provided, which supports the established benefits of using vignettes clinically and in qualitative research.⁵²

⁵³ Because a child's concept of pain is not always overtly communicated by the child, particularly the 'hidden' emotional components which affect communication,⁵⁴ using vignettes and drawings in pain science education or assessments may be particularly beneficial in clinical practice. In addition to these future clinical directions, exploring the effects of different pain durations and pain diagnoses on a child's focus on emotional aspects of pain is warranted.

A further potential clinical implication is that a more flexible and individualized assessment of a child's concept of pain could directly inform what pain science education is likely to be most beneficial for a child. Pain science education for adults is based on conceptual change theory where the assumption is that misconceptions exist.¹⁵ The results of the current study show that complex concepts are still developing in childhood, and so pain science education for children should also incorporate educational approaches where concepts are taught with gradually increasing complexity built on previous conceptual development.⁵⁵ Tailored, individualised pain science education, specifically aimed at addressing the different variables that have made up an

individual child’s concept of pain ‘puddle’ (Figure 3), appears warranted. Unexplored variables may be impacting on a child’s concept of pain, such as their intellectual and emotional intelligence, their general attitude, fears, social circumstances, perspectives on ‘toughness’, and other aspects of their history. Each individual child may benefit from tailored education provided in different ways. Some children may not require any formal pain science education, but rather benefit most from reassurance from a trusted adult with influence such as a parent,⁵⁶ teacher or health professional. Education deemed too complex by a child may result in misunderstanding and fear,⁵⁷ which could potentially aid in the development and/or maintenance of persistent pain and disability.^{58 59}

The terminology and understanding of neuroanatomy used in descriptions of a child’s concept of pain is also important given the use of neuroanatomical terms in pain science education.^{14 15} In the current study, children typically conceptualized the nervous system without inclusion of the spinal cord and considered it to have two parts; the brain and a peripheral component, and uncertainty existed regarding how these two parts are connected. Phrases such as ‘spinal cord’, ‘nerves’, and ‘real’, are likely to be poorly understood and potentially misinterpreted by children. Participants appeared to grasp the concept of the brain receiving warning messages and, based on these messages and many other factors, producing pain to keep the body safe. It is critical to carefully consider language when discussing pain with children,⁴² as younger participants in this age group tended to only use and understand simpler terminology. Age-appropriate language used in conversations by health professionals may facilitate the therapeutic alliance. For example, if a child uses a particular word or phrase to describe their concept of pain, a patient-centred empathetic communication strategy⁶⁰ would be for the clinician to incorporate the child’s

language in the discussion rather than using technical terms. In addition, the apparent lack of awareness regarding how sensory input and distraction can influence pain provides some support for educating children at an age-appropriate level about biological mechanisms underpinning pain.

This study has several strengths and limitations. One strength of this study was the use of drawing tasks and vignettes in the face-to-face interviews as a means of data triangulation.⁶¹ Another strength was the large variations in pain experience amongst the group of 8 to 12-year old participants with persistent pain; some participants had lifelong permanent physical disabilities whereas others had recurrent headaches for just over one year, capturing individual variability in pain experiences.⁶² A limitation of this study is the potential lack of transferability. This study only included children from New South Wales, Australia who spoke English. It is unknown if the results would be transferable to other cultures and age groups. A further limitation is that other 'raindrops' in the proposed conceptual framework could exist that were not identified and explored in these interviews and would be suitable to investigate in future research.

In conclusion, Australian children aged 8-12 years who have not received formal pain science education had concepts of pain that depended on a child's knowledge and experiences. The importance of gaining an in-depth understanding of a child's previous pain-related experiences and knowledge is emphasized, to facilitate clear and meaningful pain science education. The use of age-appropriate language, in combination with assessment and education tasks such as drawing and discussing vignettes, allowed participants with persistent pain to communicate their

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

individual concept of pain. Future research quantitatively assessing a child’s concept of pain is needed to improve and target pain science education for children with pain.

For peer review only

REFERENCES

1. Friedrichsdorf S, Giordano J, Desai Dakoji K, et al. Chronic pain in children and adolescents: diagnosis and treatment of primary pain disorders in head, abdomen, muscles and joints. *Children* 2016;3(4):42.
2. King S, Chambers CT, Huguet A, et al. The epidemiology of chronic pain in children and adolescents revisited: a systematic review. *Pain* 2011;152(12):2729-38. doi: 10.1016/j.pain.2011.07.016
3. Swain MS, Henschke N, Kamper SJ, et al. An international survey of pain in adolescents. *BMC Public Health* 2014;14(1):447.
4. Access Economics. The high price of pain: the economic impact of persistent pain in Australia. Sydney: MBF Foundation 2007:9-11.
5. Hunfeld JA, Perquin CW, Duivenvoorden HJ, et al. Chronic pain and its impact on quality of life in adolescents and their families. *J Pediatr Psychol* 2001;26(3):145-53.
6. Hassett AL, Hilliard PE, Goesling J, et al. Reports of chronic pain in childhood and adolescence among patients at a tertiary care pain clinic. *J Pain* 2013;14(11):1390-97.
7. Lisman-van Leeuwen Y, Spee LA, Benninga MA, et al. Prognosis of abdominal pain in children in primary care—a prospective cohort study. *Ann Fam Med* 2013;11(3):238-44.
8. Kosola S, Mundy LK, Sawyer SM, et al. Pain and learning in primary school: a population-based study. *Pain* 2017;158(9):1825-30. doi: 10.1097/j.pain.0000000000000984
9. Zernikow B, Wager J, Hechler T, et al. Characteristics of highly impaired children with severe chronic pain: a 5-year retrospective study on 2249 pediatric pain patients. *BMC Pediatr* 2012;12(1):54.

10. Coffelt TA, Bauer BD, Carroll AE. Inpatient characteristics of the child admitted with chronic pain. *Pediatrics* 2013;132(2):e422-e29.

11. Groenewald CB, Essner BS, Wright D, et al. The economic costs of chronic pain among a cohort of treatment-seeking adolescents in the United States. *J Pain* 2014;15(9):925-33.

12. Mahrer NE, Gold JI, Luu M, et al. A cost-analysis of an interdisciplinary pediatric chronic pain clinic. *The Journal of Pain* 2018;19(2):158-65.

13. Kamper SJ, Apeldoorn A, Chiarotto A, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain: Cochrane systematic review and meta-analysis. *BMJ* 2015;350:h444.

14. Harrison LE, Pate JW, Richardson PA, et al. Best-Evidence for the Rehabilitation of Chronic Pain Part 1: Pediatric Pain. *Journal of Clinical Medicine* 2019;8(9):1267.

15. Moseley GL, Butler DS. Fifteen Years of Explaining Pain: The Past, Present, and Future. *J Pain* 2015;16(9):807-13. doi: 10.1016/j.jpain.2015.05.005

16. Louw A, Zimney K, Puentedura EJ, et al. The efficacy of pain neuroscience education on musculoskeletal pain: A systematic review of the literature. *Physiother Theory Pract* 2016;32(5):332-55. doi: 10.1080/09593985.2016.1194646

17. Driessnack M. Children's drawings as facilitators of communication: a meta-analysis. *J Pediatr Nurs* 2005;20(6):415-23.

18. Robins H, Perron V, Heathcote L, et al. Pain neuroscience education: state of the art and application in pediatrics. *Children* 2016;3(4):43.

19. Harbeck C, Peterson L. Elephants dancing in my head: a developmental approach to children's concepts of specific pains. *Child Dev* 1992;63(1):138-49.

20. Jacobson CJ, Jr., Kashikar-Zuck S, Farrell J, et al. Qualitative Evaluation of Pediatric Pain Behavior, Quality, and Intensity Item Candidates and the PROMIS Pain Domain Framework in Children With Chronic Pain. *J Pain* 2015;16(12):1243-55. doi: 10.1016/j.jpain.2015.08.007
21. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19(6):349-57. doi: 10.1093/intqhc/mzm042
22. Kvale S. *InterViews*. Thousands Oaks: Ca: Sage, 1996.
23. Liamputtong P. *Qualitative research methods*. 3rd ed. Victoria, Australia: Oxford University Press 2009.
24. Arbuckle R, Abetz-Webb L. "Not just little adults": qualitative methods to support the development of pediatric patient-reported outcomes. *Patient* 2013;6(3):143-59. doi: 10.1007/s40271-013-0022-3
25. Christie D, Viner R. ABC of adolescence: Adolescent development. *BMJ* 2005;330(7486):301.
26. Catley MJ, O'Connell NE, Moseley GL. How good is the neurophysiology of pain questionnaire? A Rasch analysis of psychometric properties. *The journal of pain* 2013;14(8):818-27.
27. Butler D, Moseley G. *Explain pain supercharged*. Adelaide City West 2017
28. Pate JW, Hush JM, Hancock MJ, et al. A Child's Concept of Pain: An International Survey of Pediatric Pain Experts. *Children* 2018;5(1):12.
29. Docherty S, Sandelowski M. Focus on qualitative methods: Interviewing children. *Res Nurs Health* 1999;22(2):177-85.

30. Lobe B, Livingstone S, Olafsson K, et al. Best Practice Research Guide: How to research children and online technologies in comparative perspective: EU Kids Online 2008.

31. Kortessluoma RL, Hentinen M, Nikkonen M. Conducting a qualitative child interview: methodological considerations. *J Adv Nurs* 2003;42(5):434-41.

32. Ritchie J, Lewis J, Nicholls CM, et al. Qualitative research practice: A guide for social science students and researchers: Sage 2013.

33. TED-Ed. How does your brain respond to pain? - Karen D. Davis. 2014. YouTube (2:24 - 2:26). <https://www.youtube.com/watch?v=I7wfDenj6CQ> (accessed 1 Oct. 2017).

34. Popay J, Rogers A, Williams G. Rationale and standards for the systematic review of qualitative literature in health services research. *Qual Health Res* 1998;8(3):341-51.

35. Pink B. Census of Population and Housing: Socio-Economic Indexes for Areas (SEIFA), Australia, 2011. Technical Paper. *Australian Bureau of Statistics (ABS), ed Canberra: Australian Government* 2011:54-58.

36. Bonoti F, Leondari A, Mastora A. Exploring children's understanding of death: through drawings and the death concept questionnaire. *Death Stud* 2013;37(1):47-60.

37. Buckley CA, Waring MJ. Using diagrams to support the research process: Examples from grounded theory. *Qual Res* 2013;13(2):148-72.

38. QSR International Pty Ltd. NVivo Qualitative Data Analysis Software (Version 9): QSR International Pty Ltd Doncaster, Vic, Australia, 2006.

39. Tamm ME, Granqvist A. The meaning of death for children and adolescents: A phenomenographic study of drawings. *Death Stud* 1995;19(3):203-22.

40. Angell C, Alexander J, Hunt JA. 'Draw, write and tell': A literature review and methodological development on the 'draw and write' research method. *Early Child Res* 2015;13(1):17-28.
41. Ely EA. The experience of pain for school-age children: Blood, Band-Aids, and feelings. *Children's Health Care* 1992;21(3):168-76.
42. Johnson E, Bornman J, Tönsing KM. An exploration of pain-related vocabulary: implications for AAC use with children. *Augmentative and Alternative Communication* 2016;32(4):249-60.
43. Yeung K-K, Engle L, Rabel A, et al. It just feels weird!: a qualitative study of how children aged 10–18 years describe neuropathic pain. *Disabil Rehabil* 2017;39(17):1695-702. doi: 10.1080/09638288.2016.1207107
44. Bunzli S, Watkins R, Smith A, et al. Lives on hold: a qualitative synthesis exploring the experience of chronic low-back pain. *Clin J Pain* 2013;29(10):907-16.
45. Sim J, Madden S. Illness experience in fibromyalgia syndrome: a metasynthesis of qualitative studies. *Soc Sci Med* 2008;67(1):57-67.
46. Coakley R, Schechter N. Chronic pain is like... The clinical use of analogy and metaphor in the treatment of chronic pain in children. *Pediatric Pain Letter* 2013;15(1):1-8.
47. Posner GJ, Strike KA, Hewson PW, et al. Accommodation of a scientific conception: Toward a theory of conceptual change. *Sci Educ* 1982;66(2):211-27.
48. Celedon X, Amari A, Ward CM, et al. Children and adolescents with chronic pain and functional disability: use of a behavioral rehabilitation approach. *Current Physical Medicine and Rehabilitation Reports* 2014;2(2):86-92.

49. Williams G, Howard RF, Liossi C. Persistent postsurgical pain in children and young people: prediction, prevention, and management. *Pain Reports* 2017;2(5)

50. Gross J, Hayne H. Drawing facilitates children's verbal reports of emotionally laden events. *J Exp Psychol Appl* 1998;4(2):163.

51. Liamputtong P, Fernandes S. What makes people sick?: The drawing method and children's conceptualisation of health and illness. *Australasian Journal of Early Childhood* 2015;40(1):23-32.

52. Barter C, Renold E. 'I wanna tell you a story': exploring the application of vignettes in qualitative research with children and young people. *Int J Soc Res Methodol* 2000;3(4):307-23.

53. Jenkins N, Bloor M, Fischer J, et al. Putting it in context: the use of vignettes in qualitative interviewing. *Qualitative Research* 2010;10(2):175-98.

54. Hadjistavropoulos T, Craig KD, Duck S, et al. A biopsychosocial formulation of pain communication. *Psychol Bull* 2011;137(6):910.

55. Novak JD, Cañas AJ. The theory underlying concept maps and how to construct and use them. Pensacola: Institute for Human and Machine Cognition, 2008.

56. Evans S, Tsao JC, Lu Q, et al. Parent-child pain relationships from a psychosocial perspective: a review of the literature. *J Pain Manag* 2008;1(3):237.

57. Simons LE, Kaczynski KJ. The Fear Avoidance model of chronic pain: examination for pediatric application. *The Journal of Pain* 2012;13(9):827-35.

58. Zale EL, Ditre JW. Pain-related fear, disability, and the fear-avoidance model of chronic pain. *Current opinion in psychology* 2015;5:24-30.

- 1
2
3 59. Vlaeyen JW, Crombez G, Linton SJ. The fear-avoidance model of pain. *Pain*
4
5 2016;157(8):1588-89.
6
7
8 60. Linton SJ, Flink IK, Nilsson E, et al. Can training in empathetic validation improve medical
9
10 students' communication with patients suffering pain? A test of concept. *Pain Rep*
11
12 2017;2(3):e600.
13
14
15 61. Krefting L. Rigor in qualitative research: The assessment of trustworthiness. *Am J Occup*
16
17 *Ther* 1991;45(3):214-22.
18
19 62. Denk F, McMahon SB. Neurobiological basis for pain vulnerability: why me? *Pain*
20
21 2017;158:S108-S14.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Example questions, drawing activities, and vignettes in the interview script.

<i>Questions</i>	<ul style="list-style-type: none">• What do you think pain is?<ul style="list-style-type: none">◦ What can pain feel like? Any picture words you can think of?• What do you think happens inside your body for you to feel hurt/sore?<ul style="list-style-type: none">◦ Or “how do you think pain is made inside your body?”• What do you think of when you hear the word “injury”?• Can someone be injured but have no pain?• Can someone have pain but not be injured?<ul style="list-style-type: none">◦ Where did you learn about that?• The job of eyes is to see. The job of ears is to hear. The job of the nose is to _____. And if pain were to have a job, what do you think the job of pain would be?• Why does pain last a long time for some people and not for others?• Do you think pain is always real?
<i>Drawing activities</i>	<ul style="list-style-type: none">• I want you to draw whatever the word “pain” makes you think of.• Draw where you think your brain, spinal cord and nerves are with 3 different coloured pencils on the diagram.
<i>Vignettes</i>	<ul style="list-style-type: none">• What do you think is happening inside this man’s body (cartoon character in a video) when he puts his hand on the stove?<ul style="list-style-type: none">◦ Why doesn’t his hand just stay on the stove top?• Do you think this cartoon character can have pain without knowing about it?<ul style="list-style-type: none">◦ Or “...without being aware of it?”• Let’s pretend this character has had pain for a long time. If he feels sad, do you think this would change his pain?<ul style="list-style-type: none">◦ If yes: Do you think his pain would be more/less/ bigger/ smaller/ stronger/quieter?◦ How do you think he could make his pain feel different/better/worse?

Table 2. Demographics of the participants.

Characteristics	n (%)
<i>Gender</i>	
Male	9 (56)
Female	7 (44)
<i>Age (years)</i>	
8	4 (25)
9	3 (19)
10	4 (25)
11	3 (19)
12	2 (13)
<i>Persistent Pain</i>	
Yes	8 (50)
- Duration:	
0-1 year	0 (0)
1-2 years	5 (63)
3-5 years	2 (25)
6-10 years	1 (13)
11+ years	0 (0)
- Location:	
Headache only	2 (25)
Back only	2 (25)
Neck only	1 (13)
Multiple sites	3 (38)
No	8 (50)
<i>Interview Setting</i>	
Home	9 (56)
University	6 (38)
Hospital	1 (6)

Note: Characteristics may not sum to exactly 100% due to the effect of rounding.

Table 3. Quotations to illustrate the subthemes of the four main themes.

Themes and Subthemes	Quotations
1. My pain-related knowledge: - Varied knowledge of pain's purpose, pain-related anatomy, and pain mechanisms.	<p>"...To alert your body that there is a virus and you are sick or something." (Boy, 9, persistent pain)</p> <p>"...It's to know you've hurt your knee and you need to stop playing." (Boy, 8, persistent pain)</p> <p>"The nerves feel it, then they tell the brain, [draws line up to brain outside body] and then the brain tells the nerve maybe." (Boy, 9, persistent pain)</p> <p>"Umm because like your nerves send like... It's like a phone call to your brain saying "I just hurt myself" and then your brain will say, "oh like that's going to hurt" and you go aww!" (Girl, 10, pain-free)</p> <p>"Maybe [people with higher sensitivity] have more nerves? And some people might have a little less." (Boy, 11, pain-free)</p>
- Influences of age	<p>[When asked how a vignette character could make his persistent abdominal pain feel different] "...Umm well not eat as much food if he is eating a lot. Basically, go on a diet, and depending on if he is fit or not, get fit." (Boy, 10, persistent pain)</p>
2. Pain in the world around me: - Contextual influences on a child's concept of pain	<p>"My mum... she has... and my teacher... they might have to get knee replacements. My teacher had one. And my mum has to get one. And my neighbour got one." (Girl, 9, persistent pain)</p> <p>[When asked where he learnt about the role of the brain] "Well I learnt most of it off TV, and a little bit from school." (Boy, 12, persistent pain)</p>
- Perceptions of how other people respond to pain	<p>"... Some people might be more sensitive than others. Like, I guess some people who play like full contact sports they would be more tough because they are used to the punching and everything. And people who don't do any sports or who are really sensitive naturally – you just go like "that!" (touches hand to desk lightly) it really, really hurts." (Girl, 10, pain-free)</p> <p>"I guess it's just how you're born. And how your nerves work." (Girl, 10, pain-free)</p> <p>"Some people have stronger muscles... It's a matter between weakness and strength." (Boy, 11, persistent pain)</p> <p>[When asked about sensitivity] "It means umm.. more vulnerable".</p> <p>"...Umm well not really, because some people have a high pain tolerance. Because if Sarah, Mary and I did that, Mary and Sarah would be crying because they have low pain tolerances. Because I'm kind of tougher than them, I'm the oldest." (Girl, 11, pain-free)</p> <p>[When asked if different people feel the same amount of pain with the same hammer-hitting-thumb-injury] "We wouldn't feel the same amount of pain. Because you're older and I'm younger. I can't hold in the pain as much as you. So I'd feel more than you." (Boy, 8, pain-free)</p> <p>"Well, if you're like 96 or something you'd feel pain like kids... Because you're getting old." (Boy, 8, pain-free)</p> <p>"[Adults] may be more used to it, because they may have hit it before – depending how often! Because they may have had longer time to experience pain on their thumbs." (Boy, 10, persistent pain)</p>
3. Pain in me: - Experiences of any pain/injury	<p>"Well pain is like if you've hurt yourself, it's the physical contact and it hurts. Or like after surgeries you'd have like pain and it hurts to move." (Girl, 12, persistent pain)</p> <p>"It helps us not get into any more trouble... it'll help us to not do [an injury] again. We'll remember last time and [do the task] differently." (Boy, 8, pain-free)</p> <p>"Well if you really didn't like something, like a sport, then you don't have to play it. It'd be helpful maybe." (Girl, 10, pain-free)</p> <p>"If you had, like we have a presentation day that goes for 2 hours at school. If you didn't want to go, if you have pain you can stay home." (Girl, 9, persistent pain)</p>
- The impact of having persistent pain	<p>"You could go see a doctor" (Boy, 8, pain-free)</p> <p>"If they have pain then they can go to a physio..." (Boy, 8, pain-free)</p> <p>"If you maybe have a sore tummy, and you're like "oh no I might have appendicitis!" or something like that. And then you go to the doctors and then they're like "oh no you're just sick" or something like that. You're like "Oh thank goodness, I've only got a fever!" (Girl, 11, pain-free)</p> <p>"Umm... I reckon no... I always know about it [pain]." (Boy, 10, persistent pain)</p> <p>"If you defined pain in the dictionary... it would be something that physically hurts you. So it's a fake thing. Just a warning... It depends what you mean by real?" (Boy, 12, persistent pain)</p> <p>[When asked if pain was always 'real'] "I think it's always real. Because sometimes in your body, like right now when I'm talking, pain can still be in your body even if it doesn't hurt." (Boy, 8, pain-free)</p>
- The physical is obvious, but the emotional is hidden	<p>"I reckon if he [the vignette character] looked at like his favourite thing in the world, and got to play with it or just looked at it at least and just went like "I need it" – it would distract him from his pain because he is focussed on his favourite toy or something, in the world." (Boy, 10, persistent pain)</p>

4. Communicating my concept of pain:	<i>[Pain is] "...when something hurts in your body" (Girl, 10, pain-free)</i>
- Verbal communication	<i>"The body is fighting the virus or something." (Boy, 9, persistent pain)</i>
- Individual engagement	<i>[Pointing to parts of the diagram whilst talking] "[Nerves] release chemicals that flow up to your brain. And then your brain sends down another chemical I think, that gives it pain." (Boy, 12, persistent pain)</i>

For peer review only

1

2

3 **Figure Legends**

4

5

6

7

8 Figure 1. Examples of drawings by children when asked to draw whatever the word ‘pain’ makes

9 them think of. Panels (a) and (b) represent how children who were pain-free drew injuries (9

10 years, female; 10 years female). Panels (c) and (d) are examples of how children with persistent

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

pain drew more emotional elements such as hearts and tears (11 years, male; 10 years, female).

Figure 2. Eight examples of labelled diagrams of the brain, spinal cord and nerves by children

aged 8-12 years. The top row are drawings of children who were pain-free (8 years, male; 8

years, male; 10 years, female; 11 years, male) and the bottom row are drawings of children with

persistent pain (8 years, male; 12 years, male; 9 years, female; 11 years, male). No differences

between children with and without persistent pain were identified in this drawing task.

Figure 3. A proposed conceptual framework of the potential interactions between the themes

resulting from the analysis and a child’s concept of pain.

- Notes:*
- 1. The raindrops (influences on a child’s concept of pain) without labels highlight that all potential contributors to a child’s concept of pain may not have been identified; and*
 - 2. The size of the raindrops vary from one child to the next.*

Word Count: 5054 words

Data availability statement:

Deidentified transcripts and drawings, and the unpublished protocol, will be made available upon reasonable request from Joshua Pate (<https://orcid.org/0000-0002-1049-3916>) until 2024 as per ethical approval. Reuse without ethical approval is not permitted. A data-sharing agreement will require a commitment to using the data only for specified research purposes, to securing the data appropriately and to destroying the data after a nominated period.

Contributorship statement:

Conception and design: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Planning and implementation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Data collection: J.W.P.

Coding data for thematic analysis: J.W.P., T.N., and V.P.

Analysis and interpretation: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Drafting the article: J.W.P.

Review and editing: J.W.P., T.N., J.M.H., M.J.H., R.S., M.P., and V.P.

Overall responsibility: J.W.P. under the supervision of V.P., J.M.H., and M.J.H.




Figure 1




Figure 2




Figure 3. A proposed conceptual framework of the potential interactions between the themes resulting from the analysis and a child's concept of pain.

Notes:

1. The raindrops without labels highlight that all potential contributors to a child's concept of pain may not have been identified; and
2. The size of the raindrops vary from one child to the next.

Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

Developed from Table 1 of: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

No.	Item	Guide questions/description	Reported on Page #
Domain 1: Research team and reflexivity			
<i>Personal Characteristics</i>			
1.	Interviewer/facilitator	Which author/s conducted the interview or focus group?	Page 9
2.	Credentials	What were the researcher's credentials? E.g. PhD, MD	Page 9
3.	Occupation	What was their occupation at the time of the study?	Page 9
4.	Gender	Was the researcher male or female?	Page 9
5.	Experience and training	What experience or training did the researcher have?	Page 9
<i>Relationship with participants</i>			
6.	Relationship established	Was a relationship established prior to study commencement?	Page 9
7.	Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	Page 9
8.	Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. bias, assumptions, reasons and interests in the research topic	Page 9
Domain 2: Study design			
<i>Theoretical framework</i>			
9.	Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	Page 11
<i>Participant selection</i>			
10.	Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	Page 8
11.	Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	Page 8
12.	Sample size	How many participants were in the study?	Page 11
13.	Non-participation	How many people refused to participate or dropped out? Reasons?	Page 11
<i>Setting</i>			

14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	Table 2 (Page 31)
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	Page 10
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	Table 2 (Page 31)
<i>Data collection</i>		
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	Table 1 (Page 30)
18. Repeat interviews	Were repeat inter iews carried out? If yes, how many?	No, page 7
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	Page 10
20. Field notes	Were field notes made during and/or after the interview or focus group?	Page 10
21. Duration	What was the duration of the interviews or focus group?	Page 11
22. Data saturation	Was data saturation discussed?	Page 8 and 9
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	Page 10
Domain 3: analysis and findings		
<i>Data analysis</i>		
24. Number of data coders	How many data coders coded the data?	3 coders, Page 11
25. Description of the coding tree	Did authors provide a description of the coding tree?	Page 11
26. Derivation of themes	Were themes identified in advance or derived from the data?	Page 11
27. Software	What software, if applicable, was used to manage the data?	Page 11
28. Participant checking	Did participants provide feedback on the findings?	Page 10
<i>Reporting</i>		
29. Quotations presented	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	Table 3, page 32
30. Data and findings consistent	Was there consistency between the data presented and the findings?	Yes
31. Clarity of major themes	Were major themes clearly presented in the findings?	Yes
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	Yes