

BMJ Open One-minute and five-minute Apgar scores and child developmental health at 5 years of age: a population-based cohort study in British Columbia, Canada

Neda Razaz,¹ Sven Cnattingius,² Martina Persson,¹ Kristina Tedroff,³ Sarka Lisonkova,⁴ K S Joseph⁴

To cite: Razaz N, Cnattingius S, Persson M, *et al.* One-minute and five-minute Apgar scores and child developmental health at 5 years of age: a population-based cohort study in British Columbia, Canada. *BMJ Open* 2019;**9**:e027655. doi:10.1136/bmjopen-2018-027655

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-027655>).

Received 2 November 2018

Revised 27 February 2019

Accepted 6 March 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Medicine, Karolinska Institutet, Stockholm, Sweden

²Department of Medicine, Solna, Karolinska Institute, Stockholm, Sweden

³Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden

⁴Department of Obstetrics and Gynaecology, University of British Columbia, Vancouver, British Columbia, Canada

Correspondence to

Dr Neda Razaz;
neda.razaz@gmail.com

ABSTRACT

Objectives We investigated the associations between Apgar scores at 1 and 5 min, across the entire range of score values, and child developmental health at 5 years of age.

Setting British Columbia, Canada

Participants All singleton term infants without major congenital anomalies born between 1993 and 2009, who had a developmental assessment in kindergarten between 1999 and 2014.

Main outcomes and measures Developmental vulnerability on one or more domains of the Early Development Instrument and special needs requirements. Adjusted rate ratios (aRRs) and 95% CIs were estimated using log-linear regression.

Results Of the 150 081 children in the study, 45 334 (30.2%) were developmentally vulnerable and 3644 (2.5%) had special needs. There was an increasing trend in developmental vulnerability and special needs with decreasing 1 min and 5 min Apgar scores. Compared with children with an Apgar score of 10 at 5 min, the aRR for developmental vulnerability increased steadily with decreasing Apgar score from 1.02 (95% CI 1.00 to 1.04) for an Apgar score of 9 to 1.57 (95% CI 1.03 to 2.39) for an Apgar score of 2. Among children with 1 min Apgar scores in the 7–10 range, changes in Apgar scores between 1 and 5 min were associated with significant differences in developmental vulnerability. Compared with children who had an Apgar score of 9 at 1 min and 10 at 5 min, children with an Apgar score of 9 at both 1 and 5 min had higher rates of developmental vulnerability (aRR 1.03, 95% CI 1.01 to 1.05). Compared with infants with an Apgar of 10 at both 1 and 5 min, infants with a 1 min score of 10 and a 5 min score of <10 had higher rates of developmental vulnerability (aRR 1.53, 95% CI 1.08 to 2.17).

Conclusion Risks of adverse developmental health and having special needs at 5 years of age are inversely associated with 1 min and 5 min Apgar scores across their entire range.

INTRODUCTION

In 1953, Virginia Apgar proposed a scoring system that enabled a rapid assessment of the

Strengths and limitations of this study

- Ability to access comprehensive health-related and education-related databases at the population level.
- Using a teacher-reported instrument, no reliance was placed on parental report of developmental health.
- There may be some individual differences in teachers' ability to evaluate developmental health on the Early Development Instrument.
- Study was restricted to the comparatively healthy subset of all term live births, as children with severe disabilities may not have enrolled in kindergarten.

clinical status of the newborn infant and identified infants requiring resuscitation on the basis of heart rate, respiration, colour, muscle tone and reflex irritability.¹ Initially, the Apgar score at 1 min was used to assess the need for immediate resuscitation. Subsequently, the Apgar score at 5 min was shown to be a better predictor of neonatal survival than the Apgar score at 1 min. Although the value of a low Apgar score for accurately predicting adverse neurological outcomes at the individual level has been questioned,^{2 3} low Apgar scores are well correlated with both short-term⁴ and long-term outcomes, in both preterm and term infants.^{5–11}

Only the lowest and more compromised Apgar scores have been conventionally regarded as predictive of maladaptive development and morbidity. Nevertheless, a few population-based studies have shown that risks of cerebral palsy, epilepsy, early developmental health status and need for special education are inversely associated with 5 min Apgar scores in a dose-dependent manner across the entire range of scores.^{12–14} Even children with an Apgar score of 9 at 5 or

10 min have an increased risk of adverse neurological outcomes compared with children with 5 min or 10 min Apgar scores of 10.^{12 13} Although approximately 65%–85% of newborns receive a 1 min or a 5 min Apgar score in the 7–9 range,¹³ there is a dearth of information on how this impacts a child's developmental health.

Changes in Apgar score values between 1 and 5 min, and between 5 and 10 min are known to influence risks of cerebral palsy and epilepsy.^{12 15 16} Our recent population-based study demonstrated elevated risks of cerebral palsy and epilepsy among children with a 5 min Apgar score of 7 or 8, even if their 10 min Apgar score was 9 or 10.¹² Although it is recognised that changes in Apgar scores between 1 and 5 min are a useful measure of the response to resuscitation, the long-term significance of changes in such Apgar scores within the 'normal' range (ie, 7–10) is not clear.

In this population-based study, we investigated the associations between Apgar scores at 1 and 5 min across the entire range of score values, and developmental health at 5 years of age. We also analysed the effect of a change in Apgar scores from 1 to 5 min, including changes within the normal range of Apgar scores. Specifically, we were interested in developmental health among children with 1 min Apgar scores in the 7–9 range who received a score less than 10 at 5 min.

METHODS

The study was based on all singleton term infants without major congenital anomalies born between 1993 and 2009, who had a developmental assessment in kindergarten between 1999 and 2014. Information on the study population was obtained from several population-based linked health and demographic databases in British Columbia. The anonymised linked data used in this study included information from the Discharge Abstract Database,¹⁷ which comprised hospital admission and discharge records; the Vital Statistics Birth and Clinical Births¹⁸ databases, which contained information on all births in the province, along with delivery and neonatal health status, including diagnoses based on International Classification of Diseases (ICD-9 or ICD-10-CA) codes; the Census GeoData, which provided socioeconomic status (SES) data expressed as average neighbourhood income quintiles (based on census information from Statistics Canada and quantified using postal codes)¹⁹; the Consolidation File,²⁰ which provided demographic information on study subjects and confirmed residency in the province; and the Early Development Instrument (EDI)²¹ data, which provided information on early childhood developmental health, and were accessed through linkage with the Human Early Learning Partnership.²² The EDI has been routinely administered province-wide in British Columbia every 1–3 years since the 1999/2000 school year, achieving at least 85% participation of kindergarten children from each school district. Teachers completed the EDI for each child in their kindergarten class (age

range 5–7 years) in February. The EDI is designed to tap five core areas of early childhood development^{21–23}: physical health and well-being; social competence; emotional maturity; language and cognitive development; and communication skills and general knowledge (online supplementary table 1).²¹ It consists of 104 binary and Likert-scale items, from which scores between 0 and 10 are calculated for each domain. The EDI also records demographic information on each child and whether the child has identified special needs.

The study population included all singleton term (≥ 37 weeks' gestation) infants born between 01 April 1993 and 31 December 2009, who had documented 1 min and 5 min Apgar scores as well as a completed EDI assessment in kindergarten. Inclusion of infants with these birth dates meant that children were 5–7 years of age between 1999 and 2014 and part of the EDI assessment. The study population was restricted to infants without major congenital anomalies, identified using diagnosis codes from linked hospital records in the year after birth.

Apgar scores at 1 and 5 min were considered as the main exposures and examined both as discrete values from 0 to 10 and also as grouped categories (Apgar values of 0–3, 4–6, 7, 8, 9 and 10). Children with an Apgar score of 0 at 1 or 5 min who did not have a diagnostic code for birth asphyxia (ICD-9: 768.5, 768.6 and 768.9; ICD-10: P21), or an intervention code for either resuscitation or ventilation (Canadian Classification of Health Interventions: 1.GZ.30, 1.GZ31, 1.HZ.30, 1361, 1362, 1363, 1373, 1379 and 1004) were excluded from the study ($n=470$), as information on these cases likely resulted from transcription errors.

Developmental health assessment included whether a child had special needs or was developmentally vulnerable as measured by the EDI. Children were categorised as being developmentally vulnerable if their scores on the EDI fell below the 10th percentile value²⁴ in any of the five domains, based on the national EDI cut-off scores.²⁵ The 10th percentile cut-off has been recommended because it is higher and hence, more sensitive than clinical cut-off points of 3% or 5% for diagnosing developmental delay.²¹ Developmentally vulnerable children may not manifest developmental delays but may be at risk of experiencing challenges in school and society without additional support and care.²⁶ Children with special needs were defined as requiring special assistance because of chronic medical, physical or intellectually disabling conditions.

Other independent variables examined included infant sex (male vs female), birth weight-for-gestational age, age of the child in years at the time of EDI assessment, gestational age at birth in completed weeks (37, 38, 39, 40, 41 and ≥ 42), birth order (1, 2, 3 and +4), marital status (married vs not married) and SES (quintiles). Birth weight-for-gestational age was categorised as: small (<10 th percentile), appropriate (10th–90th percentile) and large (>90 th percentile) for gestational age.²⁷ Each child's family income was derived from the median household income in the child's residential area (based

Table 1 Maternal and birth characteristics according to Apgar score at 5 min among singleton term live births, British Columbia, 1993–2009

Maternal and birth characteristics	Total No. (%)	Apgar 0–3 (n=147) %	Apgar 4–6 (n=1328) %	Apgar 7 (n=2375) %	Apgar 8 (n=7666) %	Apgar 9 (n=101 191) %	Apgar 10 (n=37 374) %
Total	150 081 (100)						
Maternal age (years)							
≤19	6170 (4.11)	0.15	1.41	1.93	5.80	64.17	26.55
20–24	24 637 (16.42)	0.09	1.11	1.77	5.88	64.83	26.32
25–29	43 832 (29.21)	0.10	0.88	1.64	5.19	66.66	25.54
30–34	47 332 (31.54)	0.10	0.80	1.50	4.76	68.45	24.39
≥35	28 081 (18.71)	0.09	0.72	1.39	4.73	69.89	23.17
Missing	29 (0.02)	0	<17.24	<17.24	<17.24	58.62	31.03
Socioeconomic status							
Fifth quintile (highest)	27 519 (18.34)	0.10	0.90	1.64	5.00	66.88	25.47
Fourth quintile	31 282 (20.84)	0.11	0.83	1.69	5.38	66.79	25.21
Third quintile	30 939 (20.61)	0.10	0.86	1.65	5.18	67.47	24.74
Second quintile	31 266 (20.83)	0.06	0.84	1.48	5.08	67.73	24.80
First quintile (lowest)	28 889 (19.25)	0.12	1.00	1.45	4.88	68.25	24.30
Missing	186 (0.12)	0	<2.69	<2.69	3.23	66.13	28.49
Married							
Yes	103 099 (68.70)	0.09	0.78	1.47	4.73	68.43	24.49
No	43 374 (28.90)	0.12	1.13	1.85	6.01	64.93	25.95
Missing	3608 (2.40)	0.17	0.89	1.47	4.93	68.63	23.92
Infant's sex							
Female	73 809 (49.18)	0.08	0.78	1.46	4.91	67.17	25.61
Male	76 272 (50.82)	0.12	0.99	1.70	5.30	67.67	24.22
Birth order							
1	67 516 (44.99)	0.12	1.25	2.09	6.13	67.92	22.49
2	56 025 (37.33)	0.09	0.63	1.24	4.32	67.51	26.22
3	19 239 (12.82)	0.07	0.46	1.05	4.13	66.66	27.63
≥4	7301 (4.86)	<0.07	0.56	0.99	4.34	64.17	29.91
Gestational age (weeks)							
37	8966 (5.97)	0.10	1.08	2.02	6.88	68.02	21.89
38	25 821 (17.20)	0.05	0.74	1.37	4.67	68.21	24.96
39	37 408 (34.03)	0.09	0.76	1.32	4.36	68.57	24.89
40	51 079 (34.03)	0.10	0.82	1.65	5.05	66.31	26.08
41	25 040 (16.68)	0.15	1.22	1.87	6.08	67.38	23.29
42–44	1767 (1.18)	<0.28	1.58	2.09	6.51	61.35	28.3
Birth weight-for-gestational age							
Appropriate	121 035 (80.65)	0.09	0.84	1.51	4.96	67.42	25.18
Small	11 581 (7.72)	0.19	1.35	2.20	6.16	67.04	23.06
Large	17 445 (11.62)	0.08	0.85	1.65	5.47	67.76	24.18
Missing	20 (0.01)	<25.00	<25.00	0	0	40.00	50.00
Child's age at EDI data collection (years)							
Mean (SD)	5.70 (0.32)	5.67 (0.30)	5.65 (0.30)	5.66 (0.30)	5.66 (0.30)	5.65 (0.30)	5.65 (0.30)

EDI, Early Development Instrument.




Figure 1 Rates of vulnerability within the five Early Development Instrument domains by Apgar score at 5 min, British Columbia, Canada.

on postal code) obtained from the 2006 Canadian Census data.^{28–30}

The frequency of each 5 min Apgar score value was calculated within categories of maternal and infant characteristics. Multivariable log-linear regression models with robust variance estimates³¹ were used to examine the association between Apgar scores at 1 and 5 min and developmental vulnerability and special needs. Results were expressed as crude and adjusted rate ratios (aRRs) with 95% CIs. Other variables included in the final models were based on the literature^{24 32} or statistical significance (p value <0.10). The full model included child's sex, child's age at EDI completion, SES, child's first language, birth weight-for-gestational age, birth order and gestational age. Interactions between Apgar scores and other determinants were examined and stratified analyses were carried out when a significant interaction was present.

Patient and public involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in developing plans for or implementation of the study. No patients were asked to advise on interpretation of the findings.

RESULTS

There were 150 081 children (mean age=5.7 years) with a gestational age at birth of ≥37 weeks, without major malformations and complete Apgar and EDI data included in the study. Information on special needs was

available in 148 699 (99.1%) children. Five-minute Apgar scores showed a U-shaped association with gestational age at birth, with low scores more frequent at 37 weeks and ≥42 weeks (table 1). Low 5 min Apgar scores were comparable for most characteristics but more frequent among males, small-for-gestational-age live births, children of mothers who were nulliparous, not married and those with a low SES.

Overall, the prevalence of vulnerability in one or more domains of the EDI was 30.2%, with physical and social domains having the highest rates of vulnerability at 15.2% and 12.7%, respectively (figure 1). There was an increasing trend in the rate of developmental vulnerability with decreasing 1 min and 5 min Apgar scores (p for trend<0.001; table 2). However, this association was much more pronounced for the 5 min Apgar score. Compared with children with an Apgar score of 10 at 5 min, children with a 5 min Apgar score of 2 had a 57% higher rate of developmental vulnerability (aRR 1.57, 95% CI 1.03 to 2.39). Similarly, children with a 5 min Apgar score of 7, 8 or 9 had significantly higher rates of developmental vulnerability compared with children with a 5 min Apgar score of 10 (aRR 1.08, 1.06 and 1.02 for Apgar 7, 8 and 9, respectively; table 2). The association between 5 min Apgar scores and developmental vulnerability was mainly due to the higher rates of vulnerability in the language and emotional domains of the EDI (online supplementary table 2).

In total, 3644 (2.5%) children had special needs (table 3). The proportion of children with special needs

Table 2 Apgar scores at 1 and 5 min and rate ratios for developmental vulnerability among singleton term live births, British Columbia, Canada

Apgar score	Total number of children	Developmental vulnerability		
		No. with outcome (%)	Rate ratio (95% CI)	
			Crude	Adjusted*
1 min Apgar	150081	45334 (30.2)		
0	24	9 (37.5)	1.25 (0.74 to 2.10)	1.08 (0.64 to 1.83)
1	469	161 (34.3)	1.15 (1.00 to 1.31)	1.16 (1.02 to 1.32)
2	1060	329 (31.0)	1.04 (0.93 to 1.15)	1.03 (0.93 to 1.14)
3	1760	546 (31.0)	1.04 (0.95 to 1.13)	1.03 (0.95 to 1.13)
4	2582	814 (31.5)	1.05 (0.97 to 1.14)	1.07 (0.99 to 1.15)
5	4069	1261 (31.0)	1.03 (0.96 to 1.11)	1.05 (0.98 to 1.12)
6	6975	2124 (30.5)	1.02 (0.95 to 1.08)	1.04 (0.98 to 1.11)
7	12019	3648 (30.4)	1.01 (0.95 to 1.08)	1.03 (0.97 to 1.09)
8	38671	11666 (30.2)	1.01 (0.95 to 1.06)	1.02 (0.96 to 1.08)
9	79369	23852 (30.1)	1.00 (0.95 to 1.06)	1.00 (0.95 to 1.06)
10	3083	924 (30.0)	1.00 (reference)	1.00 (reference)
P for trend				<0.001
Per one unit of Apgar				0.99 (0.98 to 0.99)
5 min Apgar				
0	20	7 (35.0)	1.18 (0.65 to 2.15)	1.16 (0.62 to 2.17)
1	16	9 (56.3)	1.90 (1.24 to 2.93)	1.88 (1.27 to 2.77)
2	28	13 (46.4)	1.57 (1.05 to 2.34)	1.57 (1.03 to 2.39)
3	83	30 (36.2)	1.22 (0.92 to 1.63)	1.25 (0.93 to 1.67)
4	106	43 (40.6)	1.37 (1.09 to 1.73)	1.33 (1.06 to 1.67)
5	290	85 (29.3)	0.99 (0.83 to 1.19)	0.98 (0.82 to 1.17)
6	932	306 (32.8)	1.11 (1.01 to 1.22)	1.08 (0.99 to 1.18)
7	2375	740 (31.2)	1.05 (0.99 to 1.12)	1.08 (1.01 to 1.14)
8	7666	2387 (31.1)	1.05 (1.02 to 1.09)	1.06 (1.02 to 1.10)
9	101191	30668 (30.3)	1.03 (1.01 to 1.04)	1.02 (1.00 to 1.04)
10	37374	11046 (29.6)	1.00 (reference)	1.00 (reference)
P for trend				<0.001
Per one unit of Apgar				0.98 (0.97 to 0.99)

*Adjusted for child's sex (male vs female), child's age at EDI completion (years), socioeconomic status (first quintile, second quintile, third quintile, fourth quintile vs fifth quintile), child's first language (other vs English), birth order (2, 3, +4 vs 1), birth weight-for-gestational age (large, small vs appropriate) and gestational age (weeks).
EDI, Early Development Instrument.

increased linearly with decreasing 1 min and 5 min Apgar scores (p for trend<0.001). Compared with children who had a 1 min Apgar score of 10, those with an Apgar score of 2 at 1 min had significantly higher adjusted rates of having special needs (aRR 1.72, 95% CI 1.19 to 2.48), while those with an Apgar score of 5 at 1 min had 1.39 times higher rate of having special needs (95% CI 1.05 to 1.85). Children with 5 min Apgar scores in the 1–8 range had higher adjusted rates for having special needs, which consistently increased with decreasing 5 min Apgar score values: from 1.20 in children with an Apgar score of 8 at 5 min to 5.13 among those with an Apgar score of 1 at

5 min. The aRRs for having special needs among children with 1 min and 5 min Apgar scores in the 0–3 range had wide 95% CIs because of small numbers of children in these categories.

Table 4 presents rates of developmental vulnerability in relation to changes in Apgar score from 1 to 5 min, among children whose 1 min Apgar score was in the normal range (7–10). Among children with a 1 min Apgar score of 7, the rate of developmental vulnerability decreased in a dose–response manner with greater improvement in the Apgar score from 1 to 5 min (p value for dose response=0.02). Larger reductions in developmental

Table 3 Apgar score at 1 and 5 min and rate ratios for special needs status among singleton term live births in British Columbia, Canada

Apgar score	Total number of children	Special needs		
		No. with outcome (%)	Rate ratio (95% CI)	
			Crude	Adjusted*
1 min Apgar	148 699	3644 (2.5)		
0	22	<5 (4.6)	1.94 (0.28 to 13.4)	1.44 (0.23 to 8.97)
1	463	26 (5.6)	2.40 (1.55 to 3.72)	2.23 (1.44 to 3.46)
2	1054	45 (4.3)	1.82 (1.26 to 2.63)	1.72 (1.19 to 2.48)
3	1743	53 (3.0)	1.30 (0.91 to 1.84)	1.23 (0.86 to 1.74)
4	2554	69 (2.7)	1.15 (0.83 to 1.60)	1.09 (0.79 to 1.52)
5	4032	136 (3.4)	1.44 (1.09 to 1.91)	1.39 (1.05 to 1.85)
6	6894	191 (2.8)	1.18 (0.90 to 1.55)	1.16 (0.89 to 1.52)
7	11 903	298 (2.5)	1.07 (0.83 to 1.38)	1.06 (0.82 to 1.37)
8	38 300	946 (2.5)	1.06 (0.83 to 1.34)	1.07 (0.84 to 1.35)
9	78 701	1808 (2.3)	0.98 (0.78 to 1.24)	1.00 (0.79 to 1.26)
10	3033	71 (2.3)	1.00 (reference)	1.00 (reference)
P for trend				<0.001
Per one unit of Apgar				0.99 (0.98 to 0.99)
5 min Apgar				
0	17	<5 (<29.4)	2.51 (0.37 to 16.8)	2.59 (0.41 to 16.3)
1	15	<5 (<33.3)	5.69 (1.56 to 20.7)	5.13 (1.45 to 18.1)
2	28	<5 (<17.9)	6.10 (2.46 to 15.2)	5.17 (2.01 to 13.3)
3	83	9 (10.8)	4.63 (2.49 to 8.61)	3.78 (2.03 to 7.02)
4	103	7 (6.8)	2.90 (1.41 to 5.95)	2.59 (1.25 to 5.35)
5	289	8 (2.8)	1.18 (0.59 to 2.35)	1.10 (0.56 to 2.16)
6	928	36 (3.9)	1.66 (1.19 to 2.30)	1.49 (1.07 to 2.06)
7	2342	74 (3.2)	1.35 (1.07 to 1.70)	1.28 (1.01 to 1.61)
8	7597	225 (3.0)	1.26 (1.09 to 1.46)	1.20 (1.03 to 1.38)
9	100 281	2411 (2.4)	1.03 (0.95 to 1.11)	1.01 (0.94 to 1.09)
10	37 016	867 (2.3)	1.00 (reference)	1.00 (reference)
P for trend				<0.001
Per one unit of Apgar				0.98 (0.97 to 0.99)

*Adjusted for child's sex (male vs female), child's age at EDI completion (years), socioeconomic status (first quintile, second quintile, third quintile, fourth quintile vs fifth quintile), child's first language (other vs English), birth order (2, 3, +4 vs 1), birth weight-for-gestational age (large, small vs appropriate) and gestational age (weeks).
EDI, Early Development Instrument.

vulnerability with greater improvements in 1 min to 5 min Apgar scores were also evident among children with a 1 min Apgar score of 9 (p value for trend=0.009) but not among children with a 1 min Apgar score of 8 (p value for trend=0.36). Children with an Apgar score of 9 at 1 min and 9 at 5 min had higher rates of developmental vulnerability compared with those who had Apgar scores of 9 at 1 min and 10 at 5 min (aRR 1.03, 95% CI 1.01 to 1.05). Furthermore, compared with children who had Apgar scores of 10 at both 1 and 5 min, children whose 1 min Apgar score decreased from 10 to a 5 min Apgar score

of <10, had 1.53 times the rate of developmental vulnerability (aRR 1.53, 95% CI 1.08 to 2.17).

DISCUSSION

In this population-based study, we found graded, continuously increasing risks of developmental vulnerability and special needs at 5 years of age with decreasing 1 min and 5 min Apgar scores. A low Apgar score at 5 min was more strongly associated with developmental vulnerability and special needs than a low Apgar score at 1 min. In particular, children with 'normal' 5 min Apgar scores of 7, 8

Table 4 Rate ratios for developmental vulnerability according to the combination of Apgar scores at 1 and 5 min, singleton term live births, British Columbia, Canada

1 min Apgar	5 min Apgar	Total number of children	Developmental vulnerability			
			No. with outcome (%)	Rate ratio (95% CI)		P for trend
				Crude	Adjusted*	
7	<7	20	9 (45.0)	1.62 (0.99 to 2.65)	1.34 (0.80 to 2.25)	
7	7	172	56 (32.6)	1.18 (0.93 to 1.48)	1.18 (0.94 to 1.47)	
7	8	1987	629 (31.7)	1.14 (1.02 to 1.28)	1.12 (1.01 to 1.23)	
7	9	8700	2637 (30.3)	1.09 (0.99 to 1.20)	1.08 (0.99 to 1.19)	
7	10	1140	317 (27.8)	1.00 (reference)	1.00 (reference)	0.024
8	<8	66	17 (25.8)	0.85 (0.56 to 1.28)	0.71 (0.47 to 1.07)	
8	8	1337	420 (31.4)	1.03 (0.94 to 1.13)	1.01 (0.92 to 1.10)	
8	9	33255	10007 (30.1)	0.99 (0.94 to 1.04)	0.97 (0.93 to 1.02)	
8	10	4013	1222 (30.5)	1.00 (reference)	1.00 (reference)	0.36
9	<9	140	48 (34.3)	1.17 (0.93 to 1.47)	1.10 (0.88 to 1.38)	
9	9	50976	15501 (30.4)	1.03 (1.01 to 1.06)	1.03 (1.01 to 1.05)	
9	10	28253	8303 (29.4)	1.00 (reference)	1.00 (reference)	0.009
10	<10	26	13 (50.0)	1.68 (1.14 to 2.47)	1.53 (1.08 to 2.17)	
10	10	3057	911 (29.8)	1.00 (reference)	1.00 (reference)	0.016†

*Adjusted for child's sex (male vs female), child's age at EDI completion (years), socioeconomic status (first quintile, second quintile, third quintile, fourth quintile vs fifth quintile), child's first language (others vs English), birth order (2, 3, +4 vs 1), birth weight-for-gestational age (large, small vs appropriate) and gestational age (weeks).

†P value for difference in rates.

EDI, Early Development Instrument.

and 9 were more likely to have developmental vulnerability compared with children with 5 min Apgar scores of 10. Similarly, children who had Apgar scores of 7 or 8 at 5 min had higher risks of having special needs compared with those with a 5 min Apgar score of 10. Furthermore, children with a 1 min Apgar score in the normal range (7–10) had an increased risk of developmental vulnerability, if their Apgar score at 5 min was <10. Particularly, noteworthy was a reduction in the Apgar score from 10 at 1 min to 7–9 at 5 min, as this substantially increased the risk of developmental vulnerability.

Our results confirm previous findings from a smaller cohort, which showed that developmental adversity extended in a linear fashion across the full range of Apgar scores.¹³ Both research and clinical practice generally emphasise the increased risks of adverse outcomes associated with very low and less common Apgar scores (ie, <7 or <4). Our results suggest that the negative association between Apgar score and developmental adversity or special needs extends across the full range of scores. Consistent with our findings, previous studies have shown a significant linear relationship between each one-point decrease in 5 min and 10 min Apgar scores and increasing risk of epilepsy, cerebral palsy and needing education in a special school.^{12 14} While profound perinatal events can cause death or obvious neurological deficits, milder insults may sometimes cause subtle cognitive impairment

only detectable as the child grows older and apparent only at a population level.

Our study also showed that changes in Apgar scores from 1 to 5 min were associated with developmental vulnerability. This is in agreement with previous studies showing that changes in Apgar scores immediately after birth influence risks of cerebral palsy and epilepsy.^{12 15 16} To the best of our knowledge, this is the first study that examined risks of developmental adversity in relation to changes in Apgar scores from 1 to 5 min. Current guidelines define 'normal' Apgar scores as 7 or more at 1 min and 8 or more at 5 min, indicating that the baby does not require assistance if scores are within these ranges.³³ However, our results reveal that lower scores within the normal range (7–9) and even a slight reduction in score from 10 at 1 min to 9 at 5 min are both associated with a significant increase in the risk of developmental vulnerability. Similarly, infants who have low Apgar scores for prolonged, or even brief periods are reported to have a higher risk of poor IQ scores at age 18, even if the infants recover subsequently.⁶ The higher developmental vulnerability observed among infants whose optimal Apgar score (of 10) at 1 min falls with time after birth may be important clinically; such a progression may indicate problems with physiological circulatory, respiratory or central nervous system changes that follow delivery. Deterioration in the Apgar score immediately after birth, therefore, warrants

re-evaluation of the infant and close clinical scrutiny in order to exclude congenital abnormalities and drug-induced depression of the central nervous system.

The strengths of our study included the ability to access comprehensive health-related and education-related databases at the population level. By using a teacher-reported instrument, no reliance was placed on parent or self-report of developmental health. Nonetheless, there may be some individual differences in teachers' ability to evaluate developmental health on the EDI.²⁵ Further, our study was restricted to the comparatively healthy subset of all term live births, as children with severe disabilities may not have enrolled in kindergarten or may have enrolled in special needs schools. Furthermore, although the EDI has broad coverage across British Columbia, it is collected less frequently in independent schools (30% coverage). We recognise that the Apgar score as recorded in medical charts represents routine clinical practice,³⁴ and is prone to interobserver variability,³⁴ specifically in intubated newborn babies.³⁵ However, the quality of Apgar score values should not differ between children with and without subsequent diagnosed developmental vulnerability. Nevertheless, measurement errors inherent in routinely recorded Apgar scores (and possibly the EDI) may potentially explain the lack of an evident dose-response relationship between Apgar scores and developmental vulnerability. Finally, we acknowledge that the incidence of adverse outcomes in the setting of normal Apgar scores is rare and a low Apgar in the normal range is a poor predictor of developmental vulnerability for the individual infant.

In summary, our study showed that the risk of developmental vulnerability and special needs at 5 years of age was inversely associated with 1 min and 5 min Apgar scores across their entire range. Furthermore, improvements in Apgar scores between 1 and 5 min among children with a 1 min Apgar score of 7–9 were associated with a lower risk of developmental vulnerability. These results provide clinicians with valuable prognostic information and the justification to carefully monitor infants who are even mildly compromised at 1 and 5 min. Future studies should examine the underlying mechanism by which Apgar scores in the normal range could influence long-term neurodevelopmental outcomes.

Contributors NR conceptualised and designed the study, analysed the data, drafted the initial manuscript and finalised the manuscript based on co-authors' feedback. She had full access to all the data used in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. SC, MP, KT and SL reviewed and commented on the initial and final analyses, provided feedback on the initial draft of the manuscript and approved the final version of the manuscript. KJ assisted with conceptualisation and design of the study, reviewed and commented on the initial and final analyses, provided feedback on the initial draft of the manuscript and approved the final version of the manuscript.

Funding NR is supported by a postdoctoral fellowship award from the Canadian Institutes of Health Research (CIHR). KJ is supported by the BC Children's Hospital Research Institute.

Disclaimer All inferences, opinions and conclusions drawn in this journal article are those of the authors and do not reflect the opinions or policies of the data steward(s).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The University of British Columbia's Clinical Research Ethics Board approved the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Appgar V. A proposal for a new method of evaluation of the newborn infant. *Anesthesia & Analgesia* 1953;32:260-267.
2. Bharti B, Bharti S. A review of the Apgar score indicated that contextualization was required within the contemporary perinatal and neonatal care framework in different settings. *J Clin Epidemiol* 2005;58:121-9.
3. American Academy of Pediatrics Committee on Fetus and Newborn, American College of Obstetricians and Gynecologists Committee on Obstetric Practice. The Apgar Score. *Pediatrics* 2015;136:819-.
4. Li J, Cnattingius S, Gissler M, et al. The 5-minute Apgar score as a predictor of childhood cancer: a population-based cohort study in five million children. *BMJ Open* 2012;2:e001095.
5. Moore EA, Harris F, Laurens KR, et al. Birth outcomes and academic achievement in childhood: A population record linkage study. *Journal of Early Childhood Research* 2014;12:234-50.
6. Odd DE, Rasmussen F, Gunnell D, et al. A cohort study of low Apgar scores and cognitive outcomes. *Arch Dis Child Fetal Neonatal Ed* 2008;93:F115-F120.
7. Ehrenstein V, Pedersen L, Grijsa M, et al. Association of Apgar score at five minutes with long-term neurologic disability and cognitive function in a prevalence study of Danish conscripts. *BMC Pregnancy Childbirth* 2009;9:14.
8. Marschik PB, Einspieler C, Garzaroli B, et al. Events at early development: are they associated with early word production and neurodevelopmental abilities at the preschool age? *Early Hum Dev* 2007;83:107-14.
9. Krebs L, Langhoff-Ross J, Thorngren-Jerneck K. Long-term outcome in term breech infants with low Apgar score—a population-based follow-up. *Eur J Obstet Gynecol Reprod Biol* 2001;100:5-8.
10. Cnattingius S, Norman M, Granath F, et al. Apgar score components at 5 minutes: risks and prediction of neonatal mortality. *Paediatr Perinat Epidemiol* 2017;31:328-37.
11. Tweed EJ, Mackay DF, Nelson SM, et al. Five-minute Apgar score and educational outcomes: retrospective cohort study of 751,369 children. *Arch Dis Child Fetal Neonatal Ed* 2016;101:F121-F126.
12. Persson M, Razaz N, Tedroff K, et al. Five and 10 minute Apgar scores and risks of cerebral palsy and epilepsy: population based cohort study in Sweden. *BMJ* 2018;360:k207.
13. Razaz N, Boyce WT, Brownell M, et al. Five-minute Apgar score as a marker for developmental vulnerability at 5 years of age. *Arch Dis Child Fetal Neonatal Ed* 2016;101:F114-F120.
14. Stuart A, Otterblad Olausson P, Källen K. Apgar scores at 5 minutes after birth in relation to school performance at 16 years of age. *Obstet Gynecol* 2011;118(2 Pt 1):201-8.
15. Sun Y, Vestergaard M, Pedersen CB, et al. Apgar scores and long-term risk of epilepsy. *Epidemiology* 2006;17:296-301.
16. Moster D, Lie RT, Irgens LM, et al. The association of Apgar score with subsequent death and cerebral palsy: A population-based study in term infants. *J Pediatr* 2001;138:798-803.
17. British Columbia Ministry of Health [creator] [2012]. *Discharge Abstract Database (Hospital Separations): Population Data BC* [publisher]; Data Extract, 2012. <http://www.popdata.bc.ca/data>
18. British Columbia Vital Statistics Agency [creator] [2012]. *Vital Statistics Births: Population Data BC* [publisher]; Data Extract BC Vital Statistics Agency, 2012. <http://www.popdata.bc.ca/data>
19. Statistics Canada [creator] (2009). *Statistics Canada Income Band Data. Catalogue Number: 13C0016: Population Data BC* [publisher]; Population Data BC, 2017. <http://www.popdata.bc.ca/data>
20. British Columbia Ministry of Health [creator] [2012]. *Consolidation File (MSP Registration & Premium Billing): Population Data BC* [publisher]; Data Extract, 2012. <http://www.popdata.bc.ca/data>

21. Janus M, Offord DR. Development and psychometric properties of the Early Development Instrument (EDI): a measure of children's school readiness. *Can J Behav Sci* 2007;39:1–22.
22. Human Early Learning Partnership [creator] (2012) Early Development Instrument. *Human Early Learning Partnership*. Vancouver, BC: Population Data BC [publisher]; Data Extract University of British Columbia, School of Population and Public Health, 2012. <http://www.popdata.bc.ca/data>.
23. Duncan GJ, Dowsett CJ, Claessens A, *et al*. School readiness and later achievement. *Dev Psychol* 2007;43:1428–46.
24. Janus M, Duku E. The school entry gap: Socioeconomic, Family, and Health Factors Associated With Children's School Readiness to Learn. *Early Education & Development* 2007;18:375–403.
25. The Offord Centre for Child Studies. Measuring in support of early childhood development: The Normative II report [Report]. 2012 http://www.offordcentre.com/readiness/files/updated_normative_II.pdf (Accessed 01 Apr 2013).
26. Brinkman S, Sayers M, Goldfeld S, *et al*. Population monitoring of language and cognitive development in Australia: The Australian Early Development Index. *Int J Speech Lang Pathol* 2009;11:419–30.
27. Kramer MS, Platt RW, Wen SW, *et al*. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics* 2001;108:e35.
28. Martens P. *Health inequities in Manitoba: is the socioeconomic gap in health widening or narrowing over time?* Winnipeg: Manitoba Centre for Health Policy, University of Manitoba, 2010.
29. Mustard CA, Derksen S, Berthelot J-M, *et al*. Assessing ecologic proxies for household income: a comparison of household and neighbourhood level income measures in the study of population health status. *Health Place* 1999;5:157–71.
30. Krieger N. Overcoming the absence of socioeconomic data in medical records: validation and application of a census-based methodology. *Am J Public Health* 1992;82:703–10.
31. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
32. Santos R, Brownell MD, Ekuma O, *et al*. *The Early Development Instrument (EDI) in Manitoba: linking socioeconomic adversity and biological vulnerability at birth to children's outcomes at age 5*. Winnipeg, MB: Manitoba Centre for Health Policy, University of Manitoba, 2012.
33. AAo P. The APGAR score. *Adv Neonatal Care* 2006;6:220–3.
34. O'Donnell CPF, Kamlin COF, Davis PG, *et al*. Interobserver variability of the 5-minute Apgar score. *J Pediatr* 2006;149:486–9.
35. Lopriore E, van Burk GF, Walther FJ, *et al*. Correct use of the Apgar score for resuscitated and intubated newborn babies: questionnaire study. *BMJ* 2004;329:143–4.