



BMJ Open CCEDRRN COVID-19 Infection Score (CCIS): development and validation in a Canadian cohort of a clinical risk score to predict SARS-CoV-2 infection in patients presenting to the emergency department with suspected COVID-19

Andrew D McRae ¹, Corinne M Hohl,² Rhonda Rosychuk,³ Shabnam Vatanpour,¹ Gelareh Ghaderi,² Patrick M Archambault,⁴ Steven C Brooks,⁵ Ivy Cheng,⁶ Philip Davis,⁷ Jake Hayward,⁸ Eddy Lang ¹, Robert Ohle,⁹ Brian Rowe,⁸ Michelle Welsford,¹⁰ Krishan Yadav,¹¹ Laurie J Morrison,¹² Jeffrey Perry,¹¹ Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN) investigators for the Network of Canadian Emergency Researchers and the Canadian Critical Care Trials Group

To cite: McRae AD, Hohl CM, Rosychuk R, *et al.* CCEDRRN COVID-19 Infection Score (CCIS): development and validation in a Canadian cohort of a clinical risk score to predict SARS-CoV-2 infection in patients presenting to the emergency department with suspected COVID-19. *BMJ Open* 2021;**11**:e055832. doi:10.1136/bmjopen-2021-055832

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055832>).

ADM and CMH are joint first authors.
LJM and JP are joint senior authors.

Received 27 July 2021
Accepted 08 November 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Andrew D McRae;
amcrae@ucalgary.ca

ABSTRACT

Objectives To develop and validate a clinical risk score that can accurately quantify the probability of SARS-CoV-2 infection in patients presenting to an emergency department without the need for laboratory testing.

Design Cohort study of participants in the Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN) registry. Regression models were fitted to predict a positive SARS-CoV-2 test result using clinical and demographic predictors, as well as an indicator of local SARS-CoV-2 incidence.

Setting 32 emergency departments in eight Canadian provinces.

Participants 27 665 consecutively enrolled patients who were tested for SARS-CoV-2 in participating emergency departments between 1 March and 30 October 2020.

Main outcome measures Positive SARS-CoV-2 nucleic acid test result within 14 days of an index emergency department encounter for suspected COVID-19 disease.

Results We derived a 10-item CCEDRRN COVID-19 Infection Score using data from 21 743 patients. This score included variables from history and physical examination and an indicator of local disease incidence. The score had a c-statistic of 0.838 with excellent calibration. We externally validated the rule in 5295 patients. The score maintained excellent discrimination and calibration and had superior performance compared with another previously published risk score. Score cut-offs were identified that can rule-in or rule-out SARS-CoV-2 infection without the need for nucleic acid testing with 97.4% sensitivity (95% CI 96.4 to 98.3) and 95.9% specificity (95% CI 95.5 to 96.0).

Conclusions The CCEDRRN COVID-19 Infection Score uses clinical characteristics and publicly available indicators of disease incidence to quantify a patient's probability of SARS-CoV-2 infection. The score can identify patients at sufficiently

Strengths and limitations of this study

- Patients were enrolled in a large, geographically distributed network of Canadian urban, regional and rural emergency departments, with strict data quality and cleaning protocols to ensure reliability of collected data.
- In addition to clinical variables, we also included the average daily incidence of SARS-CoV-2 infections in a patient's health region, which is an essential predictor of the probability of a patient's risk of COVID-19 infection.
- Some missing data required either multiple imputation or classification of missing categorical variables as being absent, but the overall missingness of data in this registry is very low.
- Although the data collection for the Canadian COVID-19 Emergency Department Rapid Response Network registry relies on abstraction from health records, this approach has been shown to be reliable in our study sites when compared with prospective data collection.
- This risk score was developed using data from patients enrolled in the first 9 months of the pandemic when rates of influenza were low, so the score may need to be revalidated and refined in the future to reflect the influence of influenza, the emergence of variant strains of SARS-CoV-2 and widespread population immunisation on patients' risk of infection.

high risk of SARS-CoV-2 infection to warrant isolation and empirical therapy prior to test confirmation while also identifying patients at sufficiently low risk of infection that they may not need testing.

Trial registration number NCT04702945.



INTRODUCTION

To date, the WHO has reported 190 million diagnosed cases of COVID-19 with 4.2 million fatalities.¹ Despite the availability of vaccines to prevent COVID-19, incomplete population-level immunisation and the emergence of variants of concern means that hospitals around the world need to continue to identify and isolate patients with suspected COVID-19 from the time they arrive in the emergency department until their SARS-CoV-2 test results are available. In acutely ill patients, clinicians may need to initiate empirical therapy immediately. A quantitative risk score that can accurately predict the probability of a positive SARS-CoV-2 test result would guide initial isolation and empirical therapy prior to nucleic acid amplification test (NAAT) test result availability while identifying patients with sufficiently low probability of COVID-19 who may not require testing or isolation.

Many risk prediction tools have been developed to predict the probability of SARS-CoV-2 infection.²⁻¹⁴ A living systematic review of these models concluded that most were generated using poor methodological approaches and none were ready for widespread use.² Most published risk prediction tools, including one identified as promising by the living systematic review, included early laboratory or imaging findings, thus precluding their utility to guide immediate isolation and clinical decisions at the time of first clinical contact. Other risk prediction tools using machine learning included laboratory and imaging results and can only be implemented in hospitals using electronic health records with integrated decision support. None of these models accounted for the prevalence of COVID-19 disease in the local population, which is an important risk predictor, and most only included patients from the early stages of the pandemic.²

The objective of this study is to develop a clinical risk score to predict the probability of a positive SARS-CoV-2 nucleic acid test in a large, generalisable population of patients presenting to emergency departments using only clinical characteristics and indicators of local SARS-CoV-2 incidence. This risk score is intended to guide SARS-CoV-2 testing, isolation and empirical therapy decisions without relying on other laboratory testing or diagnostic imaging. This score could be invaluable in settings that may not have access to adequate resources for timely SARS-CoV-2 testing.

METHODS

This analysis uses data from the Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN; pronounced 'SED-rin'). CCEDRRN is an ongoing multi-centre, pan-Canadian registry that has been enrolling consecutive patients presenting to emergency departments with suspected COVID-19 disease in hospitals in 8 of 10 Canadian provinces since 1 March 2020.¹⁵ Information on the network, including detailed methods and participating sites, is available elsewhere.¹⁵ Sites and enrolment periods are shown in the, online supplemental appendix,

table 1. Additional information on network sites is available in the online supplemental network appendix. This study follows the methodological and reporting recommendations outlined in the transparency in reporting of a multivariable prediction model for individual diagnosis and prognosis criteria.¹⁶ The CCEDRRN data collection form includes prespecified demographic and social variables, vital signs, symptoms and comorbid conditions (derived from the International Severe Acute Respiratory and Emerging Infection Consortium reporting form),^{17 18} exposure risk variables, hospital laboratory and diagnostic imaging test results, SARS-CoV-2 NAAT results and patient outcomes. Data were abstracted at each site using electronic medical record extraction where available as well as manual review of either electronic or paper charts (depending on site-specific documentation practices) by trained research assistants who were blinded to the potential predictor variables at the time of data collection. Reliability of health record data abstraction was evaluated by comparison with prospective data collection in a sample of patients and found to be reliable.¹⁵

Each consecutive, eligible patient enrolled in the registry was assigned a CCEDRRN unique identifier. Trained research assistants entered anonymised participant data into a REDCap database (V.10.9.4; Vanderbilt University, Nashville, Tennessee, USA). Regular data quality checks including verification of extreme or outlying values were performed by each participating site, coordinated by the CCEDRRN coordinating centre.

Participants

We included data from consecutive patients tested for SARS-CoV-2 at 32 CCEDRRN sites. From each site's start date forward, we included consecutive eligible patients aged 18 years and older who had a biological sample (swab, endotracheal aspirate and bronchoalveolar lavage) specimen collected for NAAT on their index emergency department visit or, if admitted, within 24 hours of emergency department arrival. For patients with multiple emergency department encounters involving COVID-19 testing, we only used the first encounter in this analysis.

We excluded patients who had a positive SARS-CoV-2 NAAT within 14 days prior to their emergency department visit, patients with cardiac arrest prior to emergency department arrival and those with missing outcome data.

Predictors

Candidate predictors were chosen based on clinical consensus and availability within the CCEDRRN registry. Predictors included known risk factors for SARS-CoV-2 infection, including work as a healthcare provider, institutional living (ie, long-term care and prison), close personal or household contacts with SARS-CoV-2 infection and symptoms including cough, anosmia or dysgeusia, fever, myalgias and vital signs on emergency department arrival. The full list of candidate variables and their definitions are available in the (online supplemental appendix table 2).

Table 1 Characteristics and selected outcomes of enrolled patients

	Derivation (n=21 743)	Validation (n=5922)
Age in years, median (IQR)	57 (38–73)	56 (37–73)
Female (%)	10992 (50.5)	3085 (52.1)
Arrival from, n (%)		
Home	19879 (91.4)	5429 (91.7)
Long-term care/rehabilitation facility/corrections facility	1000 (4.6)	262 (4.4)
No fixed address/shelter/single room occupancy	574 (2.6)	201 (3.4)
Interhospital transfer	290 (1.3)	30 (0.5)
Risk for infection, n (%)		
Healthcare worker	505 (2.3)	567 (9.6)
Household/caregiver contact	566 (2.6)	161 (2.7)
Institutional exposure (eg, LTC, prison)	1354 (6.2)	213 (3.6)
Microbiology lab	4 (0.0)	8 (0.1)
Travel	924 (4.2)	344 (5.8)
Other	1320 (6.1)	449 (7.6)
Unknown	5415 (24.9)	1856 (31.3)
No documented risk for infection	10028 (46.1)	1075 (18.1)
Arrival vital signs, median (IQR)		
Body temperature	36.7 (36.3–37.1)	36.8 (36.5–37.1)
Heart rate	91 (79–107)	90 (78–105)
Oxygen saturation	97 (95–98)	97 (95–99)
Respiratory rate	18 (18–20)	18 (16–20)
Systolic blood pressure	133 (118–150)	136 (120–149)
Common comorbid conditions, n (%)		
Active malignant neoplasm (cancer)	1678 (7.7)	333 (5.6)
Asthma	1699 (7.8)	468 (7.9)
Atrial fibrillation	1598 (7.3)	402 (6.8)
Chronic kidney disease	1214 (5.6)	321 (5.4)
Chronic lung disease (not asthma/pulmonary fibrosis)	1729 (8)	583 (9.8)
Chronic neurological disorder (not dementia; eg, stroke/TIA, seizure disorder)	1310 (6)	400 (6.8)
Congestive heart failure	1450 (6.7)	368 (6.2)
Coronary artery disease	1591 (7.3)	449 (7.6)
Dementia	734 (3.4)	188 (3.2)
Diabetes	2583 (11.9)	916 (15.5)

Continued

Table 1 Continued

Dialysis	198 (0.9)	28 (0.5)
Dyslipidaemia	2375 (10.9)	543 (9.2)
Hypertension	6320 (29.1)	1697 (28.6)
Hypothyroidism	1344 (6.2)	397 (6.7)
Mild liver disease	280 (1.3)	90 (1.5)
Moderate/severe liver disease	245 (1.1)	88 (1.5)
Obesity (clinical impression)	284 (1.3)	108 (1.8)
Organ transplant	128 (0.6)	19 (0.3)
Rheumatological disorder	1122 (5.2)	258 (4.4)
Other	10075 (46.3)	2174 (36.7)
Past malignant neoplasm (cancer)	936 (4.3)	256 (4.3)
Psychiatric condition/mental health diagnosis	2967 (13.6)	831 (14)
Pulmonary fibrosis	80 (0.4)	26 (0.4)
Symptoms reported, n (%)		
Abdominal pain	2725 (12.5)	540 (9.1)
Altered consciousness/confusion	1456 (6.7)	322 (5.4)
Bleeding (haemorrhage)	330 (1.5)	22 (0.4)
Chest pain (includes discomfort or tightness)	4242 (19.5)	974 (16.4)
Chills	2045 (9.4)	594 (10)
Conjunctivitis	49 (0.2)	26 (0.4)
Cough	7724 (35.5)	2663 (44.9)
Diarrhoea	2140 (9.8)	526 (8.9)
Dizziness/vertigo	1521 (7)	300 (5.1)
Dysgeusia/anosmia	140 (0.6)	33 (0.6)
Ear pain	144 (0.7)	30 (0.5)
Fatigue/malaise	3361 (15.5)	924 (15.6)
Fever	5055 (23.2)	1580 (26.7)
Headache	2144 (9.9)	624 (10.5)
Hemoptysis (bloody sputum)	298 (1.4)	66 (1.1)
Joint pain (arthralgia)	296 (1.4)	82 (1.4)
Lower chest wall indrawing	10 (0)	7 (0.1)
Lymphadenopathy	67 (0.3)	21 (0.4)
Muscle aches (myalgia)	1575 (7.2)	517 (8.7)
Nausea/vomiting	4219 (19.4)	935 (15.8)
No recorded symptoms	2113 (9.7)	431 (7.3)
Runny nose (rhinorrhoea)	1061 (4.9)	501 (8.5)
Seizures	205 (0.9)	42 (0.7)
Shortness of breath (dyspnoea)	8537 (39.3)	2383 (40.2)

Continued

**Table 1** Continued

Skin rash	241 (1.1)	38 (0.6)
Skin ulcers	27 (0.1)	<5
Sore throat	3024 (13.9)	985 (16.6)
Sputum production	1507 (6.9)	401 (6.8)
Wheezing	582 (2.7)	130 (2.2)
Tobacco use, n (%)	1852 (8.5)	616 (10.4)
Illicit substance use, n (%)	1219 (5.6)	353 (6.0)
Oxygen required in ED, n (%)	1919 (8.8)	627 (10.6)
Hospital admission, n (%)	9913 (45.6)	2446 (41.3)
In-hospital death, n (%)	753 (3.5)	213 (3.6)
7-day average incident COVID-19 cases, median (IQR)	1.3 (0.73-2)	0.96 (0.5-1.3)
SARS-CoV-2 positive, n (%)	940 (4.3)	227 (3.8)

ED, emergency department; LTC, long-term care; TIA, transient ischaemic attack.

In addition to these clinical variables, the 7-day average incident COVID-19 case count was calculated for the health region of each participating site using publicly available epidemiological data.¹⁹ For each calendar day within each health region represented in the study, we calculated the average daily incident rate of new infections per 100 000 population over the preceding 7 days. This 7-day average incidence was assigned to each patient based on the date of their index emergency department encounter and the health region of the forward sortation area of their postal code of residence. For patients with no fixed address, we allocated them to the health region of the hospital in which they were tested. As publicly available incident COVID-19 case data were not available for the early pandemic, we imputed values for the first 5 weeks of the pandemic by modelling the reported COVID-19 cases that had accumulated in every health region over time using linear interpolation (0.1% missing).

Outcome

The primary outcome of this analysis was the diagnosis of SARS-CoV-2 infection using a criterion standard of a positive NAAT at the time of index emergency department visit or within 14 days after the index encounter.

Sample size and precision

The 46 candidate predictors had 52 degrees of freedom and with an expected SARS-CoV-2 infection rate of 5%, a sample size of 1040 was sufficient for the derivation cohort based on an anticipated event rate of less than 20% and a requirement for 20 outcomes per degree of freedom.²⁰ Over 21 000 patients were available for the derivation cohort at the time of analysis, providing more than sufficient data for reliable prediction modelling.

Model development and validation

We randomly assigned study sites to the derivation and validation cohorts with the goal of assigning 75% of eligible patients and outcome events to the derivation cohort and 25% to the validation cohort. Thus, the derivation and validation cohorts are geographically distinct. Within the derivation cohort, candidate predictors were examined for collinearity and missing or extreme values. In the presence of collinearity, one predictor was dropped from the set of candidate predictors. Five multiple imputations were used for continuous variables with missing data. Patients with values of 'not recorded' for categorical variables (eg, smoking and need for supplemental oxygen) were assumed to have the reference value (ie, 'no') for that categorical variable. The initial logistic regression model considered all candidate predictors, with continuous predictors fit with restricted cubic splines with three knots. The strengths of associations between predictors and outcome were assessed using an analysis of variance plot to inform the df to allocate to each predictor. The model was fit again with these changes. A fast step-down procedure reduced the model to key predictors based on an Akaike's information criterion stopping rule with a threshold of 120 to enable a model with a relatively small number of predictors that would be clinically easy to use. Internal bootstrap validation with 1000 bootstrap samples was conducted to provide an optimism-corrected C-statistic. Continuous predictors were categorised based on the relationship between the spline function and outcome.

We then developed the points-based CCEDRRN COVID-19 Infection Score (CCIS) using a nomogram to assign integer point values for each variable included in the derived model. Discrimination of the score was evaluated using the c-statistic. Calibration was evaluated using calibration curves and comparison of observed and expected outcomes. Diagnostic performance was evaluated using sensitivity and specificity, predictive values and likelihood ratios at different point thresholds.

We then evaluated the discrimination, calibration and performance characteristics of the CCIS in an external validation cohort of patients from geographically distinct study sites who were not part of the derivation cohort.

Validation of previously published models

We used our combined (derivation and validation) study cohort to externally validate the COvid Rule out Criteria (CORC) score developed by Kline *et al*³ (although we were not able to include race and ethnicity variables as these are not reliably recorded or reported in most Canadian hospitals). We compared measures of discrimination and calibration, along with sensitivity and specificity of risk score values for the CCIS and CORC (with race and ethnicity variables removed). We split each score into categories of low, moderate and

high risk for SARS-CoV-2 infection. Low risk was defined as a score having a sensitivity for ruling out infection of 95% or higher. High risk was defined as a score having a specificity for ruling in infection of 95% or higher. We compared the performance of the two scores by calculating net reclassification improvement across low-risk, moderate-risk and high-risk categories.^{21 22}

All analyses were performed in R²³ using the rms package.²⁴

Role of the funding sources

The funding organisations had no role in the study conduct, data analysis, manuscript preparation or submission.

Patient and public involvement

The CCEDRRN governance structure includes patient representatives on the Executive Committee, Scientific Steering Committee, Protocol Review and Publications Committee, Data Access and Monitoring Committee and Knowledge Translation Committee. The network also has a Patient Engagement Committee composed of patient partners from across Canada. Patient partners provided input into study design and selection of outcomes for all CCEDRRN analyses and provide advice on knowledge sharing and translation strategies.

RESULTS

This analysis is based on 27665 patients consecutively enrolled from 32 participating emergency departments between March and October 2020 (figure 1, online supplemental appendix table 1). Sites and enrolment periods contributing patient data are shown in the online supplemental appendix table 1. Of the included patients, 1167 (4.2%) had a positive SARS-CoV-2 NAAT result, including 1133 who had a positive initial test and 34 who tested positive after a negative (27) or indeterminate (7) initial NAAT.

The study cohort was subdivided into a derivation cohort (21 743 patients from 16 sites, 940 (4.3%) SARS-CoV-2 positive) and a separate external validation cohort (5922 patients from 16 different sites, 227 (3.8%) SARS-CoV-2 positive). Demographic and clinical characteristics of the derivation and validation cohorts are shown in table 1. No continuous variable requiring multiple imputation had more than 3.4% missingness (online supplemental appendix table 2).

In the derivation cohort, we derived a 10-variable model to predict the probability of a patient having a positive SARS-CoV-2 NAAT. The regression coefficients and ORs for each variable in the model are shown in table 2. The c-statistic for the derived model was 0.851 with excellent calibration.

We created a points-based CCIS using rounded regression coefficients with a range of -2 to 9 points (table 2). The c-statistic of the CCIS in the derivation cohort was 0.838 (0.824–0.852) with excellent calibration (figure 2).




Figure 1 Flow diagram of patients through the study . ED, emergency department.

A score of zero or less ruled out a positive SARS-CoV-2 test result in 5996/21 743 patients (27.6%) with a sensitivity of 96.6% (95% CI 95.2 to 97.7). A score of 4 or more was observed in 1338/21 743 patients (6.2%) and had a specificity of 95.6 (95% CI 95.3 to 95.8) indicating a low frequency of false positives (online supplemental appendix table 3).

We then quantified the performance of the CCIS in our external validation cohort. In this cohort, the c-statistic for the points-based risk score was 0.792 (figure 2). A score of zero or less ruled out a positive SARS-CoV-2 test result in 1863/5925 patients (31.4%) with a sensitivity of 94.3% (95% CI 90.4 to 96.9). A score of 4 or more was observed in 174/5925 patients (2.9%) and had a specificity of 97.8 (95% CI 97.4 to 98.1) indicating a low frequency of false positives (table 3).

In a combined cohort of patients (derivation and validation combined), we compared the discrimination and diagnostic performance of the CCIS to the CORC score. The CCIS had a c-statistic of 0.837 compared with 0.750 for the CORC score (with race/ethnicity variables removed) (online supplemental appendix figure 1). A CCIS of zero or less ruled out SARS-CoV-2 infection in 28.4% of patients with a sensitivity of 96.1% (online supplemental appendix table 4), whereas a CORC score of negative one or less ruled out SARS-CoV 2 infection in 9.9% of patients with 97.4% (online supplemental appendix table 5) sensitivity. Compared with the CORC

**Table 2** Adjusted associations between model predictor variables and SARS-CoV-2 nucleic acid test results

Variable/score component	Regression coefficient (SE)	Adjusted OR (95% CI)	Score value
7-day average incident COVID-19 cases			
0–2 daily cases per 100 000 population	–	–	0
2–7.99 daily cases per 100 000 population	1.22 (0.09)	3.38 (2.85 to 4.00)	1
≥8 daily cases per 100 000 population	2.21 (0.10)	9.09 (7.53 to 10.97)	2
Institutional exposure (eg, LTC, prison) or travel from country with known cases within 14 days	0.88 (0.09)	2.40 (2.01 to 2.87)	1
Healthcare worker/microbiology lab	1.10 (0.16)	3.02 (2.22 to 4.10)	1
Household/caregiver contact	1.83 (0.12)	6.25 (4.92 to 7.93)	2
Temperature			
<36 and no self-reported fever	–0.75 (0.3)	0.47 (0.28 to 0.80)	–1
36–37.4 and no self-reported fever	–	–	0
≥37.5 or self-reported fever	1.21 (0.08)	3.36 (2.88 to 3.91)	1
Supplemental oxygen delivered in the ED	0.98 (0.1)	2.66 (2.18 to 3.24)	1
Cough	0.85 (0.08)	2.33 (2.01 to 2.71)	1
Dysgeusia/anosmia	2.03 (0.24)	7.60 (4.76 to 12.15)	2
Muscle aches (myalgia)	0.7 (0.11)	2.02 (1.64 to 2.48)	1
Current tobacco user	–1.13 (0.21)	0.32 (0.21 to 0.49)	–1

ED, emergency department; LTC, long-term care.

score (with race/ethnicity variables removed), the CCIS showed substantial net reclassification improvement (NRI=0.310,) (online supplemental appendix table 6).

DISCUSSION

We have derived and validated a simple clinical risk score, the CCIS, to predict the probability of a positive SARS-CoV-2 NAAT in patients presenting to emergency departments. It uses only clinical variables available at the patient's bedside, along with a common publicly available measure of community COVID-19 incidence. In this study population, the score ruled out SARS-CoV-2 infection with 96.1% sensitivity in almost one-third of patients. It also identified patients at high risk of infection with over 95% specificity.

In addition to clinical variables, we also included the average daily incidence of SARS-CoV-2 infections in a patient's health region, which is an essential predictor of the probability of a patient's risk of COVID-19 infection. Although access to timely incidence data may be challenging in under-resourced health systems, this information is publicly reported in many health jurisdictions. In practice, the local incidence would likely need to be shared within an emergency department on a daily basis. We developed data-driven cut-offs for categorisation of low, moderate and high incidence for calculation the CCIS. Thus, the clinician would only need to know whether local incidence is high, moderate or low to use this score, and the incidence category changes slowly over time. Patients who live and work in separate

health regions could be assigned the higher incidence value at hospital presentation for a conservative risk estimate. Patients in areas with high disease burden will automatically score 2 points, meaning that few patients in these settings will be classified as low risk. Therefore, symptomatic patients would all warrant testing. This underscores the need for liberal isolation and testing practices in settings with high rates of community SARS-CoV-2 transmission.

The CCIS has several important clinical applications. The ability to differentiate patients with high or low probability of COVID-19 disease could guide safe and effective patient isolation or cohorting from the time of hospital arrival, prior to the availability of SARS-CoV-2 test results. Identification of patients with extremely low risk of SARS-CoV-2 infection may even allow safe omission of testing, which will minimise testing resource utilisation in settings with limited testing capacity. Identifying patients with a high probability of SARS-CoV-2 infection can help prioritise use of rapid antigen testing and initiation of effective empiric therapy in critically ill patients prior to availability of NAAT results. By presenting risk estimates and sensitivity for all risk score values, we allow end-users to choose cut-offs for ruling-in and ruling-out SARS-CoV-2 infection that make sense for their setting and application.

Several other risk prediction instruments have been developed to predict positive COVID-19 test results in undifferentiated patients. These tools were developed in studies with substantial methodological limitations and incorporate variables not immediately available at




Figure 2 Distribution and performance of the CCEDRRN COVID-19 infection score in the derivation cohort (left panel) and validation cohorts (right panel): (A) distribution of the score, (B) observed in-hospital mortality across the range of the score, (C) predicted versus observed probability of in-hospital mortality and (D) receiver operating characteristic curve with area under the curve and associated 95% CI. CCEDRRN, Canadian COVID-19 Emergency Department Rapid Response Network.

the time of a patient's hospital arrival, so are not useful to guide early isolation, testing and treatment decisions.² None of these risk prediction tools considered the prevalence of disease in the population. Prevalence can substantially change the approach to testing and cohorting, and this will become increasingly important as prevalence rates drop and selective rather than liberal testing may be more appropriate.

US-based investigators recently reported the development³ and validation²⁵ of the CORC score using only clinical variables. The CORC score contains several similar variables to the CCIS. The CORC score included race and ethnicity as predictor variables, which may limit the generalisability of the CORC score beyond the urban American population in which it was developed, as it does not reflect the international diversity of ethnic

**Table 3** Performance metrics for the CCEDRRN COVID-19 Infection Score for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the validation cohort

Score cut-off	n (%)	Sensitivity (%, 95% CI)	Specificity (%, 95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	17 (0.3)	100 (98.4 to 100)	0.3 (0.2 to 0.5)	1	NA	0 (0)
≤-1	310 (5.2)	99.6 (97.6 to 100)	5.4 (4.9 to 6.1)	1.1	0.1	1 (0.3)
≤0	1863 (31.5)	94.3 (90.4 to 96.9)	32.5 (31.3 to 33.7)	1.4	0.2	13 (0.7)
≤1	3806 (64.3)	78.9 (73.0 to 84.0)	66.0 (64.7 to 67.2)	2.3	0.3	48 (1.3)
≤2	5152 (87.0)	52.9 (46.2 to 60.0)	88.6 (87.7 to 89.4)	4.6	0.5	107 (2.1)
≤3	5748 (97.1)	20.7 (15.6 to 26.6)	97.8 (97.4 to 98.1)	9.3	0.8	180 (3.1)
Rule in:						
≥3	770 (13.0)	52.9 (46.2 to 59.5)	88.6 (87.7 to 89.4)	4.6	0.5	120 (15.6)
≥4	174 (2.9)	20.7 (15.6 to 26.6)	97.8 (97.4 to 98.1)	9.3	0.8	47 (27.0)
≥5	44 (0.7)	7.9 (4.8 to 12.2)	99.5 (99.3 to 99.7)	17.4	0.9	18 (40.9)
≥6	6 (0.1)	0.9 (0.1 to 3.2)	99.9 (99.8 to 100)	12.5	1	2 (33.3)
≥7	1 (<0.1)	0 (0 to 1.6)	100.0 (99.9 to 100)	NA	1	0 (0)

CCEDRRN, Canadian COVID-19 Emergency Department Rapid Response Network; LR+, positive likelihood ratio; LR-, negative likelihood ratio.

backgrounds. Moreover, it is unlikely race or ethnicity represents a biological risk. The association between race and ethnicity and SARS-CoV-2 infection in the CORC score likely reflects other sociodemographic and geographic predictors of the risk of COVID-19 infection in the American population.²⁵ The CCIS uses the 7-day average local incidence as an estimate of population risk. We believe this approach is more generalisable across populations and better reflects individual patients' pretest probability of SARS-CoV-2 infection.²⁶

Strengths and limitations

The cohorts used to derive and validate the rule included comprehensive data on consecutive eligible patients from a large, geographically distributed network of Canadian urban, regional and rural emergency departments. Strict data quality protocols and data cleaning protocols ensured the reliability of collected data. This score may be employed at the time of a patient's arrival to hospital, does not require the use of additional laboratory testing or imaging nor the use of electronic calculators or electronic medical records for implementation.

Some missing data required either multiple imputation or classification of missing categorical variables as being absent. The overall missingness of data in this registry is very low.¹⁵ Although the data collection for the CCEDRRN registry relies on abstraction from health records, this approach has been shown to be reliable in our study sites when compared with prospective data collection.¹⁵

The clinical variables in the model are not likely to be sensitive to changes in geographical changes in SARS-CoV-2 epidemiology. The variable of travel from a country with high incidence may become less

informative as the pandemic has spread globally and 'hot spots' change. However, high-prevalence areas may change over time, meaning that the risk factor of travel from a region with a high prevalence is likely to still be informative.

This risk score was developed using data from patients enrolled in the first nine months of the pandemic when rates of influenza were low. As such, the score may need to be revalidated and refined in the future to reflect the influence of influenza, the emergence of variant strains of SARS-CoV-2 and widespread population immunisation on patients' risk of infection.

CONCLUSION

We derived and successfully validated the CCIS to accurately predict the probability of SARS-CoV-2 nucleic acid test results in patients presenting to emergency departments. The CCIS uses clinical variables, accounts for the incidence of SARS-CoV-2 in the community and is ready for immediate clinical use. This score has potential utility to guide early decisions around SARS-CoV-2 test utilisation, patient isolation and empirical therapy for patients solely based on clinical assessment.

Author affiliations

¹Department of Emergency Medicine, University of Calgary Cumming School of Medicine, Calgary, Alberta, Canada

²Department of Emergency Medicine, The University of British Columbia Faculty of Medicine, Vancouver, British Columbia, Canada

³Department of Paediatrics, University of Alberta Faculty of Medicine & Dentistry, Edmonton, Alberta, Canada

⁴Department of Emergency Medicine, Université Laval Faculté de médecine, Québec, Québec, Canada

⁵Department of Emergency Medicine, Queen's University School of Medicine, Kingston, Ontario, Canada

⁶Department of Emergency Medicine, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada

⁷Department of Emergency Medicine, University of Saskatchewan College of Medicine, Saskatoon, Saskatchewan, Canada

⁸Department of Emergency Medicine, University of Alberta Faculty of Medicine & Dentistry, Edmonton, Alberta, Canada

⁹Department of Emergency Medicine, Northern Ontario School of Medicine, Thunder Bay, Ontario, Canada

¹⁰Department of Emergency Medicine, McMaster University Faculty of Health Sciences, Hamilton, Ontario, Canada

¹¹Department of Emergency Medicine, University of Ottawa Faculty of Medicine, Ottawa, Ontario, Canada

¹²Department of Emergency Medicine, St Michael's Hospital, Toronto, Ontario, Canada

Correction notice This article has been corrected since it was first published. ADM and CMH are joint first authors.

Twitter Krishan Yadav @GameYadav

Acknowledgements We gratefully acknowledge the assistance of Ms Amber Cragg in the preparation of this manuscript. We would like to thank the University of British Columbia clinical coordinating centre staff, legal, ethics, privacy and contract staff and the research staff at each of the participating institutions in the network outlined in the attached supplement. The network would not exist today without the dedication of these professionals. Thank you to all our patient partners who shared their lived experiences and perspectives to ensure that the knowledge we cocreate addresses the concerns of patients and the public. Creating the largest network of collaboration across Canadian Emergency Departments would not have been feasible without the tireless efforts of emergency department chiefs, and research coordinators and research assistants at participating sites. Finally, our most humble and sincere gratitude to all our colleagues in medicine, nursing and the allied health professions who have been on the frontlines of this pandemic from day 1 staffing our ambulances, emergency departments, intensive care units and hospitals bravely facing the risks of COVID-19 to look after our fellow citizens and after one another. We dedicate this network to you (online supplemental appendix, table 1).

Collaborators Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN) investigators for the Network of Canadian Emergency Researchers and the Canadian Critical Care Trials Group: Patrick Fok, Hana Wiemer, Samuel Campbell, Kory Arsenault, Tara Dahn, Kavish Chandra, Patrick Archambault, Joel Turner, Éric Mercier, Greg Clark, Sébastien Robert, Raoul Daoust, Laurie Morrison, Steven Brooks, Ivy Cheng, Krishan Yadav, Michelle Welsford, Rob Ohle, Justin Yan, Rohit Mohindra, Megan Landes, Tomislav Jelic, Ankit Kapur, Phil Davis, Andrew McRae, Brian Rowe, Katie Lin, Stephanie VandenBerg, Jake Hayward, Jaspreet Khangura, Corinne Hohl, Daniel Ting, Maja Stachura, Frank Scheuermeyer, Balijeet Braar, Craig Murray, John Taylor, Ian Martin, Sean Wormsbecker, Matt Bouchard, Lee Graham.

Contributors CMH, ADM, LJM, RR and JP conceived the study, with input on the design and selection of variables from the other contributors. CMH, LJM, PMA, SCB, PD and EL obtained funding on behalf of the Canadian COVID-19 Emergency Department Rapid Response Network (CCEDRRN) investigators. CMH, ADM, PMA, SCB, IC, PD, JH, BR, RO, MW and KY facilitated data collection along with other members of the CCEDRRN and can verify the underlying data. RR and JP developed the analytic plan. SV performed the analysis, with assistance from GG and RR, including accessing and verification of underlying data. All contributors provided input on interpretation of findings. ADM, CMH and RR drafted the manuscript with additional input from all contributors. ADM and CH are the guarantors of this work.

Funding The Canadian Institutes of Health Research (447679), Ontario Ministry of Colleges and Universities (C-655–2129), Saskatchewan Health Research Foundation (5357), Genome BC (COV024 and VAC007) Fondation du CHU de Québec (Octroi No. 4007). Sero-Surveillance and Research (COVID-19 Immunity Task Force Initiative) provided peer-reviewed funding. The BC Academic Health Science Network and BioTalent Canada provided non-peer reviewed funding.

Disclaimer These organisations are not-for-profit and had no role in study conduct, analysis or manuscript preparation.

Competing interests ADM, CMH, RR, PMA, SCB, IC, PD, JH, EL, RO, BR, MW, KY, LJM and JP are coinvestigators on the funding sources listed in the funding

statement and have no additional competing interests. GG and SV have no competing interests.

Patient consent for publication Not applicable.

Ethics Approval The CCEDRRN network protocol was approved with a waiver of informed consent by the University of British Columbia Research Ethics Board (UBC REB H20-01015) and subsequently by the research ethics boards of all participating institutions.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. For investigators who wish to access Canadian COVID-19 Emergency Rapid Response Network data, proposals may be submitted to the network for review and approval by the network's peer-review publication committee, the data access and management committee and the executive committee, as per the network's governance. Information regarding submitting proposals and accessing data may be found at <https://www.ccedrrn.com>

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Andrew D McRae <http://orcid.org/0000-0003-2743-4290>

Eddy Lang <http://orcid.org/0000-0003-0850-4337>

REFERENCES

- World Health Organization. WHO Coronavirus Dashboard, 2021. Available: <https://covid19.who.int/> [Accessed 19 Nov 2021].
- Update to living systematic review on prediction models for diagnosis and prognosis of covid-19. *BMJ* 2021;372:n236.
- Kline JA, Camargo CA, Courtney DM, *et al*. Clinical prediction rule for SARS-CoV-2 infection from 116 U.S. emergency departments 2-22-2021. *PLoS One* 2021;16:e0248438.
- McDonald SA, Medford RJ, Basit MA, *et al*. Derivation with internal validation of a multivariable predictive model to predict COVID-19 test results in emergency department patients. *Acad Emerg Med* 2021;28:206–14.
- Kurstjens S, van der Horst A, Herpers R, *et al*. Rapid identification of SARS-CoV-2-infected patients at the emergency department using routine testing. *Clin Chem Lab Med* 2020;58:1587–93.
- Lippi G, Henry BM, Hoehn J, *et al*. Validation of the Corona-Score for rapid identification of SARS-CoV-2 infections in patients seeking emergency department care in the United States. *Clin Chem Lab Med* 2020;58:e311–3.
- Sung J, Choudry N, Bachour R. Development and validation of a simple risk score for diagnosing COVID-19 in the emergency room. *Epidemiol Infect* 2020;148:e273.
- Schwab P, DuMont Schütte A, Dietz B, *et al*. Clinical predictive models for COVID-19: systematic study. *J Med Internet Res* 2020;22:e21439.
- Plante TB, Blau AM, Berg AN, *et al*. Development and external validation of a machine learning tool to rule out COVID-19 among adults in the emergency department using routine blood tests: a large, multicenter, real-world study. *J Med Internet Res* 2020;22:e24048.
- Soltan AAS, Kouchaki S, Zhu T, *et al*. Rapid triage for COVID-19 using routine clinical data for patients attending Hospital: development and prospective validation of an artificial intelligence screening test. *Lancet Digit Health* 2021;3:e78–87.
- Cabitza F, Campagner A, Ferrari D, *et al*. Development, evaluation, and validation of machine learning models for COVID-19 detection based on routine blood tests. *Clin Chem Lab Med* 2020;59:421–31.



- 12 Jehi L, Ji X, Milinovich A, *et al*. Individualizing risk prediction for positive coronavirus disease 2019 testing: results from 11,672 patients. *Chest* 2020;158:1364–75.
- 13 Joshi RP, Pejaver V, Hammarlund NE, *et al*. A predictive tool for identification of SARS-CoV-2 PCR-negative emergency department patients using routine test results. *J Clin Virol* 2020;129:104502.
- 14 Formica V, Minieri M, Bernardini S, *et al*. Complete blood count might help to identify subjects with high probability of testing positive to SARS-CoV-2. *Clin Med* 2020;20:e114–9.
- 15 Hohl CM, Rosychuk RJ, McRae AD, *et al*. Development of the Canadian COVID-19 emergency department rapid response network population-based registry: a methodology study. *CMAJ Open* 2021;9:E261–70.
- 16 Moons KGM, Altman DG, Reitsma JB, *et al*. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): explanation and elaboration. *Ann Intern Med* 2015;162:W1–73.
- 17 WHO. *Clinical characterisation protocol (CCP), version 3.1/3.2*. Oxford University, 2020.
- 18 ISARIC/WHO Clinical Characterisation Protocol for Severe Emerging Infections [COVID-19] [UPH]. Health Res. Available: /planning-and-improving-research/application-summaries/research-summaries/isaricwho-clinical-characterisation-protocol-for-severe-emerging-infections/ [Accessed 4 Jun 2021].
- 19 Health regional Archive (public view), 2019. Available: <https://resources-covid19canada.hub.arcgis.com/datasets/3aa9f7b1428642998fa399c57dad8045/data?layer=1> [Accessed 4 Jun 2021].
- 20 Visit 6 manual 17: ARIC neurocognitive exam. Available: <https://www2.csc.unc.edu/aric/>
- 21 Leening MJG, Vedder MM, Witteman JCM, *et al*. Net reclassification improvement: computation, interpretation, and controversies: a literature review and clinician's guide. *Ann Intern Med* 2014;160:122–31.
- 22 Kerr KF, Wang Z, Janes H, *et al*. Net reclassification indices for evaluating risk prediction instruments: a critical review. *Epidemiology* 2014;25:114–21.
- 23 R Core Team. *R: a language and environment for statistical computing*. Vienna, Austria: R foundation for statistical computing, 2020.
- 24 Harrell. rms: regression modeling Strategies., version 6.2-0, 2021. Available: <https://cran.r-project.org/web/packages/rms/index.html>
- 25 Nevel AE, Kline JA. Inter-Rater reliability and prospective validation of a clinical prediction rule for SARS-CoV-2 infection. *Acad Emerg Med* 2021;28:Epub ahead of print:761–7.
- 26 Vyas DA, Eisenstein LG, Jones DS. Hidden in Plain Sight - Reconsidering the Use of Race Correction in Clinical Algorithms. *N Engl J Med* 2020;383:874–82.

Appendices

Appendix Table 1. Patients enrolled by CCEDRRN site and time periods for data collection

Site (N patients contributed)	Mar 2020	Apr 2020	May 2020	Jun 2020	Jul 2020	Aug 2020	Sept 2020	Oct 2020	Total
Derivation Cohort	3217	4797	5493	3096	2232	1276	1266	366	21743
Abbotsford Regional Hospital		228	474	385	198				1285
Eagle Ridge Hospital	196	163							359
Foothills, Calgary	437	131							568
Halifax Infirmary/Dalhousie, Nova Scotia	17								17
Hants Community Hospital, Nova Scotia	1								1
Hôpital du Sacré-Coeur de Montreal	27	96	401						524
Hôtel-Dieu de Lévis	1	19	246						266
Jewish General Hospital	754	959	93						1806
Peter Lougheed Centre	321	1119	1169	605	638	552	616	215	5235
Royal Columbian Hospital	236	408	366						1010
Royal University, Saskatoon	132	275	357	296	340	265	193		1858
Saint John Regional Hospital, New Brunswick	98	102							200
South Campus, Calgary	367	598	612	526	586	459	457	151	3756
Sunnybrook Health Sciences Centre			473	593	470				1536
The Ottawa Hospital - Civic Campus	58	24	537						619
Vancouver General Hospital	572	675	765	691					2703
Validation Cohort	2082	2012	695	381	330		422		5922
Cobequid Community Health Centre	6								6
Dartmouth General College, Dartmouth Nova Scotia	7								7
Health Science North, Sudbury Ontario			295	381	330				1006

Hôpital de l'Enfant-Jésus, CHU de Québec-Université Laval	40	99	26						165
IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec	4	5	95						104
Lions Gate Hospital	294	220							514
Mount St Joseph's	236								236
Rockyview, Calgary	368	104							472
Saint Paul's Hospital	541								541
Saskatoon City Hospital, Saskatoon	33	53							86
Secondary Assessment Centers of Dartmouth General and Halifax Infirmary	3	70	66						139
St Paul's Hospital, Saskatoon	84	198							282
Surrey Memorial Hospital	404	927							1331
The Ottawa Hospital - General Campus	62	33	135						230
Toronto Western Hospital							422		422
University of Alberta Hospital		303	78						381

Appendix Table 2. Candidate variables for entry into regression model

Variable	Definition	N (%) Missing
Demographics		
Age	Age in years	0 (0)
Sex	Male, Female, Other	0 (0)
Arrival from	Home + other (not clearly documented)	0 (0)
	Single room + no fixed address + shelter	0 (0)
	Institutional living: long-term care/rehab + correctional	0 (0)
	Inter-hospital transfer	0 (0)
Infection risk		
Travel risk	Travel from country with known cases within 14 days	0 (0)
Institutional exposure	Possible exposure in institutional setting (e. g., Long-term care, prison)	0 (0)
Healthcare worker	Healthcare worker/Microbiology lab employee	0 (0)
Household/caregiver contact	Household contact /caregiver of known positive case	0 (0)
No documented risk	Documented absence of risk factors	0 (0)
Emergency department variables		
ED arrival mode		2 (0)
Ambulance:	arrived by ambulance	
Self/police	self-transported or transported to ED by police	
Arrival heart rate	beats/minute	452 (2.1)
Arrival respiratory rate	breaths/minute	732 (3.4)
Arrival oxygen saturation	%	517 (2.4)
Lowest recorded oxygen saturation in ED	%	445 (2.0)
Fever		847 (3.9)
Temperature <36.0	Temperature <36.0C AND no self-reported fever	
Temperature 36.0-37.4	Temperature 36.0-37.4C AND no self-reported fever	
Temperature ≥37.5 or fever	Temperature ≥37.5 OR self-reported fever	
Respiratory distress	Increased work of breathing documented by treating clinician	1(0)
Supplemental oxygen delivered in the ED	Yes/No	0 (0)
COVID symptoms		
Abdominal pain	Patient-reported symptom as documented by treating clinician	0 (0)
Altered consciousness/confusion	Patient-reported symptom as documented by treating clinician	0 (0)
Bleeding (hemorrhage)	Patient-reported symptom as documented by treating clinician	0 (0)
Chest pain (includes discomfort or tightness)	Patient-reported symptom as documented by treating clinician	0 (0)
Chills	Patient-reported symptom as documented by treating clinician	0 (0)
Conjunctivitis	Patient-reported symptom as documented by treating clinician	0 (0)
Cough	Patient-reported symptom as documented by treating clinician	0 (0)

Diarrhea	Patient-reported symptom as documented by treating clinician	0 (0)
Dizziness/Vertigo	Patient-reported symptom as documented by treating clinician	0 (0)
Dysgeusia/anosmia	Patient-reported symptom as documented by treating clinician	0 (0)
Ear pain	Patient-reported symptom as documented by treating clinician	0 (0)
Fatigue/malaise	Patient-reported symptom as documented by treating clinician	0 (0)
Headache	Patient-reported symptom as documented by treating clinician	0 (0)
Hemoptysis (bloody sputum)	Patient-reported symptom as documented by treating clinician	0 (0)
Joint pain (arthralgia)	Patient-reported symptom as documented by treating clinician	0 (0)
Lymphadenopathy	Patient-reported symptom as documented by treating clinician	0 (0)
Muscle aches (myalgia)	Patient-reported symptom as documented by treating clinician	0 (0)
Nausea/vomiting	Patient-reported symptom as documented by treating clinician	0 (0)
No reported symptoms	Patient-reported symptom as documented by treating clinician	0 (0)
Runny nose (rhinorrhea)	Patient-reported symptom as documented by treating clinician	0 (0)
Seizures	Patient-reported symptom as documented by treating clinician	0 (0)
Shortness of breath (dyspnea)	Patient-reported symptom as documented by treating clinician	0 (0)
Skin rash	Patient-reported symptom as documented by treating clinician	0 (0)
Sore throat	Patient-reported symptom as documented by treating clinician	0 (0)
Sputum production	Patient-reported symptom as documented by treating clinician	0 (0)
Wheezing	Patient-reported symptom as documented by treating clinician	0 (0)
Current tobacco user	Documented current tobacco use	6 (0)
Current illicit user	Documented methamphetamine, opioid or other illicit drug use	6 (0)
7-day average incident COVID-19 cases	Daily reported incidence of new cases in health region, averaged over the seven days preceding hospital arrival. Reported in units of new cases/100,000 population	32 (0.1)

Appendix Table 3. Performance metrics for the CCEDRRN COVID-19 Infection Score for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the derivation cohort

Score	n (%)	Sensitivity (%, 95% CI)	Specificity (%, 95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	51 (0.2)	100 (99.6–100)	0.25 (0.2–0.3)	1.0	NA	0 (0)
≤-1	937 (4.3)	99.89 (99.4–100)	4.5 (4.2–4.8)	1.1	<0.1	1 (0.1)
≤0	5996 (27.6)	96.6 (95.2–97.7)	28.67 (28.1–29.3)	1.4	0.1	32 (0.5)
≤1	13114 (60.3)	86.6 (84.3–88.7)	62.43 (61.8–63.1)	2.3	0.2	126 (1.0)
≤2	18041 (83.0)	67.34 (64.2–70.3)	85.25 (84.8–85.7)	4.6	0.4	307 (1.7)
≤3	20405 (93.9)	45.11 (41.9–48.4)	95.61 (95.3–95.9)	10.3	0.6	516 (2.5)
Rule in:						
≥3	3702 (17.0)	67.34 (64.2–70.3)	85.25 (84.8–85.7)	4.6	0.4	633 (17.1)
≥4	1338 (6.2)	45.11 (41.9–48.4)	95.61 (95.3–95.9)	10.3	0.6	424 (31.7)
≥5	440 (2.0)	23.51 (20.8–26.4)	98.95 (98.8–99.1)	22.3	0.8	221 (50.2)
≥6	122 (0.6)	9.68 (7.9–11.8)	99.85 (99.8–99.9)	65.0	0.9	91 (74.6)
≥7	31 (0.1)	2.77 (1.8–4.0)	99.98 (99.9–100.0)	115.1	1.0	26 (83.9)
≥8	12 (0.1)	1.17 (0.6–2.1)	100 (100.0–100.0)	243.4	1.0	11 (91.7)
≥9	2 (<0.1)	0.21 (<0.1–0.8)	100 (100.0–100.0)	NA	1.0	2 (100)

LR+: Positive Likelihood Ratio; LR-: Negative Likelihood Ratio

Appendix Table 4. Performance metrics for CCEDRRN COVID Infection Score for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the combined cohort

Score cutoff	n (%)	Sensitivity (95% CI)	Specificity (95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	70 (0.3)	100 (99.7,100)	0.3 (0.2,0.3)	1	0	0 (0)
≤-1	1257 (4.5)	99.8 (99.4,100)	4.7 (4.5,5)	1.1	<0.1	2 (0.2)
≤0	7872 (28.5)	96.1 (94.8,97.1)	29.5 (29.0,30.1)	1.4	0.1	46 (0.6)
≤1	16962 (61.3)	85 (82.8,87.0)	63.4 (62.8,63.9)	2.3	0.2	175 (1.0)
≤2	23243 (84.0)	64.8 (62.0,67.5)	86.2 (85.7,86.6)	4.7	0.4	411 (1.8)
≤3	26169 (94.6)	40.3 (37.4,43.2)	96.1 (95.9,96.4)	10.4	0.6	697 (2.7)
Rule in:						
≥3	4422 (16.0)	64.8 (62.0,67.5)	86.2 (85.7,86.6)	4.7	0.4	756 (17.1)
≥4	1496 (5.4)	40.3 (37.4,43.2)	96.1 (95.9,96.4)	10.4	0.6	470 (31.4)
≥5	476 (1.7)	20.2 (18.0,22.6)	99.1 (99.0,99.2)	22.3	0.8	236 (49.6)
≥6	128 (0.5)	8 (6.5,9.7)	99.9 (99.8,99.9)	60.3	0.9	93 (72.7)
≥7	32 (0.1)	2.2 (1.5,3.2)	100 (100,100)	98.4	1.0	26 (81.3)
≥8	12 (<0.1)	0.9 (0.5,1.7)	100 (100,100)	249.8	1.0	11 (91.7)

LR+: Positive likelihood ratio; LR-: Negative likelihood ratio

Appendix Table 5. Performance metrics for the CORC score (race and ethnicity variables removed) for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the combined cohort

Score	n (%)	Sensitivity (95% CI)	Specificity (95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	202 (0.7)	99.9 (99.5–100)	0.8 (0.7–0.9)	1.01	0.11	1 (0.5)
≤-1	2715 (9.8)	97.4 (96.4–98.3)	10.1 (9.8–10.5)	1.08	0.25	30 (1.1)
≤0	9089 (32.9)	90.1 (88.2–91.7)	33.9 (33.3–34.4)	1.36	0.29	116 (1.3)
≤1	17582 (63.9)	72.8 (70.2–75.4)	65.2 (64.6–65.7)	2.09	0.42	317 (1.8)
≤2	23421 (84.7)	51.2 (48.3–54.1)	86.2 (85.8–86.7)	3.72	0.57	569 (2.4)
≤3	26224 (94.8)	27.7 (25.1–30.3)	95.8 (95.5–96)	6.56	0.76	844 (3.2)
Rule in:						
≥3	4244 (15.3)	51.2 (48.3–54.1)	86.2 (85.8–86.7)	3.72	0.57	598 (14.1)
≥4	1441 (5.2)	27.7 (25.1–30.3)	95.8 (95.5–96)	6.56	0.76	323 (22.4)
≥5	358 (1.3)	11.1 (9.3–13)	99.1 (99–99.2)	12.79	0.9	129 (36.0)
≥6	54 (0.2)	3.1 (2.2–4.2)	99.9 (99.9–100)	45.41	0.97	36 (66.7)


LR+: Positive likelihood ratio; LR-: Negative likelihood ratio

Appendix Table 6. Net Reclassification Improvement of the CCEDRRN COVID-19 Infection Score compared to the CORC Score (race and ethnicity variables removed)

Primary Outcome : Covid Positive					
CORC risk category		CCIS risk category			Total
		Low	Medium	High	
	Low	12	18	0	30
	Medium	34	539	241	814
	High	0	94	229	323
	Total	46	651	470	1167
Primary Outcome: Covid Negative					
CORC risk category		CCIS risk category			Total
		Low	Medium	High	
	Low	1593	1092	0	2685
	Medium	6233	15756	706	22695
	High	0	798	320	1118
	Total	7826	17646	1026	26498

<u>COVID Positive</u>		<u>COVID Negative</u>	
Number of outcomes	1167	Number of outcomes	26498
Correct reclassification	259	Correct reclassification	7031
Incorrect reclassification	128	Incorrect reclassification	1798
Net reclassification	131	Net reclassification	5233
Net reclassification improvement (Event)	0.112	Net reclassification improvement (Non-event)	0.197
Total net reclassification improvement			0.310

Appendix Figure 1. Receiver operating characteristic curves for the CCEDRRN COVID Infection Score (CCIS) and the CORC score (race and ethnicity variables removed) in the combined study cohort





Supplementary Table: Contributors to the Canadian COVID-19 Emergency Department Rapid Response Network

1. Purpose

This supplementary table provides details of the support staff at each of the participating institutions in the Canadian COVID-19 Emergency Department Rapid Response Network. This supplementary document should be attached to each peer-reviewed manuscript after the methods manuscript (M1). The purpose is to ensure research staffs and lead coordinators are appropriately recognized for their contributions to the network.

2. List of Support Staff

Table 1. Network coordinating centre staff at the University of British Columbia

Name	Roles	Contributions
Gelareh Ghaderi	Data analyst	Data processing and analysis for manuscripts.
Jeffrey Hau	Data manager	REDCap, data processing and analysis for manuscripts.
Vi Ho	National coordinator	Coordinate with provincial coordinators and training/onboarding of research assistants.
Joe Larkin	Project manager	Project management.
Fiona O'Sullivan	Data analyst	Data processing and analysis for manuscripts.
Serena Small	Research coordinator	Ethics & privacy reviews, data management plan, privacy impact assessment, and qualitative analyses
Amber Cragg	Research manager	Data and manuscript management
Wei Zhao	Data analyst	Data processing and analysis for manuscripts.
Vicky Wu	Data analyst	Data processing and analysis for manuscripts.
Elnaz Bodaghkhani	Research associate	Data and manuscript management

Table 2. Provincial Coordinators

Name	Province	Institutional affiliation	Contributions to CCEDRRN
Corinne DeMone	NS	Dalhousie University, Halifax, Nova Scotia	Research ethics board submission, manages research assistants, data cleaning and quality.
Jacqueline Fraser	NB	Dalhousie University, St. John New Brunswick	Site coordinator as well as research assistant.
Veronique Gélinas	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis	Provincial research coordinator, translation of research material to French, ethics management.

Canadian
COVID-19 ED
Network

Connie Taylor	ON	Queen's University, Kingston	Coordination of research assistants in Ontario, maintenance of REB applications for the province
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg	Lead RA for the province
Aimee Goss	SK	University of Saskatchewan, Saskatoon	Screens records in Saskatoon, data/extraction and entry, coordinates research assistants.
Hina Walia	AB	University of Calgary, Calgary	Provincial coordinator lead for Alberta, oversight of all Alberta sites.
Rajan Bola	BC	University of British Columbia, Vancouver	Provincial coordinator lead for BC, oversight of all BC sites.

Table 3. Institutional research assistant (RA) leads

Institutional RA leads are responsible for data extraction and integrity, communication with provincial leads.

Name	Province	Institutional affiliation(s)
Corinne DeMone	NS	Dartmouth General Hospital, Cobequid Community Health Centre, Hants Community Hospital Secondary Assessment Centers of the Dartmouth General Hospital, and Halifax Infirmary, Halifax
Jacqueline Fraser	NB	Saint John Regional Hospital, Saint John
Alexandra Nadeau	QC	CHU de Québec Université Laval, Quebec City
Audrey Nolet	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis
Xiaoqing Xue	QC	Jewish General Hospital, Montréal
David Iannuzzi	QC	McGill University Health Center, Montréal
Chantal Lanthier	QC	Hôpital du Sacré-Cœur de Montréal, Montréal
Konika Nirmalanathan	ON	University Health Network, Toronto
Vlad Latiu	ON	Kingston General Hospital, Hotel Dieu Hospital, Kingston
Joanna Yeung	ON	Sunnybrook Health Sciences Center, Toronto
Natasha Clayton	ON	Hamilton General Hospital, Juravinski Hospital, Hamilton
Tom Chen	ON	London Health Sciences Centre, London
Jenna Nichols	ON	Health Sciences North, Sudbury
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg
Aimee Goss	SK	St. Paul's Hospital, Royal University Hospital, Saskatoon City Hospital, Saskatoon
Stacy Ruddell	AB	Foothills Medical Centre, Peter Lougheed Centre, Rockyview General Hospital, South Health Campus, Calgary
Natalie Runham	AB	University of Alberta Hospital, Edmonton

Canadian
COVID-19 ED
Network

Name	Province	Institutional affiliation(s)
Karlin Su	AB	Royal Alexandra Hospital/Northeast Community Health Center, Edmonton
Josie Kanu	BC	St. Paul's Hospital, Mount Saint Joseph, Vancouver
Bernice Huynh	BC	Abbotsford Regional Hospital and Cancer Center, Abbotsford
Amanda Swirhun	BC	Royal Columbian Hospital, New Westminster
Tracy Taylor	BC	Eagle Ridge Hospital and Health Care Centre, Port Moody
Mai Hayashi	BC	Royal Inland Hospital, Kamloops
Mackenzie Cheyne	BC	Kelowna General Hospital, Kelowna
Sarim Asim	BC	Surrey Memorial Hospital, Surrey
Katherine Lam	BC	Vancouver General Hospital, Vancouver
Kelsey Compagna	BC	Lions Gate Hospital, Vancouver

Table 4. Contributing Study Sites and Investigators

Lead Investigator	Contributing Site / Code	Member Investigators
Maritime		
Patrick Fok		
Nova Scotia		
Hana Wiemer	Halifax Infirmary/ 902	Patrick Fok
	Dartmouth General Hospital/ 903	Hana Wiemer
	Hants Community Hospital/ 904	Samuel Campbell
	Cobequid Community Health Centre/ 905	Kory Arsenaault
	Secondary Assessment Centers of Dartmouth General and Halifax Infirmary/ 908	Tara Dahn
New Brunswick		
Kavish Chandra	Saint John Regional Hospital/ 901	Kavish Chandra
Quebec		
Patrick Archambault	Hotel-Dieu de Lévis/ 701	Patrick Archambault
	Jewish General Hospital/ 702	Joel Turner
	Centre Hospitalier de l'Université Laval (CHU de Québec)/ 703	Éric Mercier
	L'hôpital Royal Victoria - Royal Victoria Hospital/ 705	Greg Clark
	Hôpital de l'Enfant-Jésus, CHU de Québec/ 706	Éric Mercier
	Hôpital du Saint-Sacrement, CHU de Québec/ 707	Éric Mercier
	Hôpital Saint-François d'Assise, CHU de Québec/ 708	Éric Mercier
	Hôtel-Dieu de Québec, CHU de Québec/ 709	Éric Mercier

Canadian
COVID-19 ED
Network

	IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec/ 710	Sébastien Robert
	Hôpital du Sacré-Coeur de Montreal/ 711	Raoul Daoust
Ontario		
Laurie Morrison & Steven Brooks	Sunnybrook/ 401	Ivy Cheng
	The Ottawa Hospital - Civic Campus/ 403	Krishan Yadav
	The Ottawa Hospital - General Campus/ 404	Krishan Yadav
	Kingston/Queens/ 406	Steven Brooks
	Hamilton General Hospital/ 407	Michelle Welsford
	Health Science North, Sudbury Ontario/ 408	Rob Ohle
	University Hospital – LHSC/ 409	Justin Yan
	North York General Hospital, Toronto/ 410	Rohit Mohindra
	Victoria Hospital – LHSC/ 412	Justin Yan
	Toronto Western Hospital/ 414	Megan Landes
Manitoba		
Tomislav Jelic	Health Sciences Centre/ 307	Tomislav Jelic
Saskatchewan		
Phil Davis	Pasqua Hospital, Regina/ 301	Ankit Kapur
	Regina General Hospital, Regina/ 302	Ankit Kapur
	St Paul's Hospital, Saskatoon/ 303	Phil Davis
	Royal University, Saskatoon/ 304	Phil Davis
	Saskatoon City Hospital, Saskatoon/ 305	Phil Davis
Alberta		
Andrew McRae	University of Alberta Hospital, Edmonton/ 201	Brian Rowe
	Foothills, Calgary/ 202	Katie Lin
	Rockyview, Calgary/ 203	Andrew McRae
	Peter Lougheed Centre/ 204	Andrew McRae
	South Campus, Calgary/ 205	Stephanie VandenBerg
	Northeast Community Health Centre, Edmonton/ 206	Jake Hayward, Jaspreet Khangura
		Royal Alexandra Hospital, Edmonton/ 306
British Columbia		
Corinne Hohl	Vancouver General Hospital/ 101	Daniel Ting
	Lions Gate Hospital/ 102	Maja Stachura
	Saint Paul's Hospital/ 103	Frank Scheuermeyer

Canadian
COVID-19 ED
Network

Mount St Joseph's/ 104	Frank Scheuermeyer
Surrey Memorial Hospital/ 105	Balijeet Braar/ Craig Murray
Royal Columbian Hospital/ 106	John Taylor
Abbotsford Regional Hospital/ 107	Ian Martin
Eagle Ridge Hospital/ 108	Sean Wormsbecker
Victoria General Hospital/ 109	Matt Bouchard
Royal Jubilee Hospital/ 110	Matt Bouchard
Nanaimo General Hospital/ 111	Matt Bouchard
Royal Inland Hospital/ 112	Ian Martin
Kelowna General / Hospital/ 115	Lee Graham

It was not possible for us to recruit Members from Newfoundland and Labrador, Northwest Territories, Nunavut, Prince Edward Island and Yukon at the time of the inception of the registry.

Appendices

Appendix Table 1. Patients enrolled by CCEDRRN site and time periods for data collection

Site (N patients contributed)	Mar 2020	Apr 2020	May 2020	Jun 2020	Jul 2020	Aug 2020	Sept 2020	Oct 2020	Total
Derivation Cohort	3217	4797	5493	3096	2232	1276	1266	366	21743
Abbotsford Regional Hospital		228	474	385	198				1285
Eagle Ridge Hospital	196	163							359
Foothills, Calgary	437	131							568
Halifax Infirmary/Dalhousie, Nova Scotia	17								17
Hants Community Hospital, Nova Scotia	1								1
Hôpital du Sacré-Coeur de Montreal	27	96	401						524
Hôtel-Dieu de Lévis	1	19	246						266
Jewish General Hospital	754	959	93						1806
Peter Lougheed Centre	321	1119	1169	605	638	552	616	215	5235
Royal Columbian Hospital	236	408	366						1010
Royal University, Saskatoon	132	275	357	296	340	265	193		1858
Saint John Regional Hospital, New Brunswick	98	102							200
South Campus, Calgary	367	598	612	526	586	459	457	151	3756
Sunnybrook Health Sciences Centre			473	593	470				1536
The Ottawa Hospital - Civic Campus	58	24	537						619
Vancouver General Hospital	572	675	765	691					2703
Validation Cohort	2082	2012	695	381	330		422		5922
Cobequid Community Health Centre	6								6
Dartmouth General College, Dartmouth Nova Scotia	7								7
Health Science North, Sudbury Ontario			295	381	330				1006

Hôpital de l'Enfant-Jésus, CHU de Québec-Université Laval	40	99	26						165
IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec	4	5	95						104
Lions Gate Hospital	294	220							514
Mount St Joseph's	236								236
Rockyview, Calgary	368	104							472
Saint Paul's Hospital	541								541
Saskatoon City Hospital, Saskatoon	33	53							86
Secondary Assessment Centers of Dartmouth General and Halifax Infirmary	3	70	66						139
St Paul's Hospital, Saskatoon	84	198							282
Surrey Memorial Hospital	404	927							1331
The Ottawa Hospital - General Campus	62	33	135						230
Toronto Western Hospital							422		422
University of Alberta Hospital		303	78						381

Appendix Table 2. Candidate variables for entry into regression model

Variable	Definition	N (%) Missing
Demographics		
Age	Age in years	0 (0)
Sex	Male, Female, Other	0 (0)
Arrival from	Home + other (not clearly documented)	0 (0)
	Single room + no fixed address + shelter	0 (0)
	Institutional living: long-term care/rehab + correctional	0 (0)
	Inter-hospital transfer	0 (0)
Infection risk		
Travel risk	Travel from country with known cases within 14 days	0 (0)
Institutional exposure	Possible exposure in institutional setting (e. g., Long-term care, prison)	0 (0)
Healthcare worker	Healthcare worker/Microbiology lab employee	0 (0)
Household/caregiver contact	Household contact /caregiver of known positive case	0 (0)
No documented risk	Documented absence of risk factors	0 (0)
Emergency department variables		
ED arrival mode		2 (0)
Ambulance:	arrived by ambulance	
Self/police	self-transported or transported to ED by police	
Arrival heart rate	beats/minute	452 (2.1)
Arrival respiratory rate	breaths/minute	732 (3.4)
Arrival oxygen saturation	%	517 (2.4)
Lowest recorded oxygen saturation in ED	%	445 (2.0)
Fever		847 (3.9)
Temperature <36.0	Temperature <36.0C AND no self-reported fever	
Temperature 36.0-37.4	Temperature 36.0-37.4C AND no self-reported fever	
Temperature ≥37.5 or fever	Temperature ≥37.5 OR self-reported fever	
Respiratory distress	Increased work of breathing documented by treating clinician	1(0)
Supplemental oxygen delivered in the ED	Yes/No	0 (0)
COVID symptoms		
Abdominal pain	Patient-reported symptom as documented by treating clinician	0 (0)
Altered consciousness/confusion	Patient-reported symptom as documented by treating clinician	0 (0)
Bleeding (hemorrhage)	Patient-reported symptom as documented by treating clinician	0 (0)
Chest pain (includes discomfort or tightness)	Patient-reported symptom as documented by treating clinician	0 (0)
Chills	Patient-reported symptom as documented by treating clinician	0 (0)
Conjunctivitis	Patient-reported symptom as documented by treating clinician	0 (0)
Cough	Patient-reported symptom as documented by treating clinician	0 (0)

Diarrhea	Patient-reported symptom as documented by treating clinician	0 (0)
Dizziness/Vertigo	Patient-reported symptom as documented by treating clinician	0 (0)
Dysgeusia/anosmia	Patient-reported symptom as documented by treating clinician	0 (0)
Ear pain	Patient-reported symptom as documented by treating clinician	0 (0)
Fatigue/malaise	Patient-reported symptom as documented by treating clinician	0 (0)
Headache	Patient-reported symptom as documented by treating clinician	0 (0)
Hemoptysis (bloody sputum)	Patient-reported symptom as documented by treating clinician	0 (0)
Joint pain (arthralgia)	Patient-reported symptom as documented by treating clinician	0 (0)
Lymphadenopathy	Patient-reported symptom as documented by treating clinician	0 (0)
Muscle aches (myalgia)	Patient-reported symptom as documented by treating clinician	0 (0)
Nausea/vomiting	Patient-reported symptom as documented by treating clinician	0 (0)
No reported symptoms	Patient-reported symptom as documented by treating clinician	0 (0)
Runny nose (rhinorrhea)	Patient-reported symptom as documented by treating clinician	0 (0)
Seizures	Patient-reported symptom as documented by treating clinician	0 (0)
Shortness of breath (dyspnea)	Patient-reported symptom as documented by treating clinician	0 (0)
Skin rash	Patient-reported symptom as documented by treating clinician	0 (0)
Sore throat	Patient-reported symptom as documented by treating clinician	0 (0)
Sputum production	Patient-reported symptom as documented by treating clinician	0 (0)
Wheezing	Patient-reported symptom as documented by treating clinician	0 (0)
Current tobacco user	Documented current tobacco use	6 (0)
Current illicit user	Documented methamphetamine, opioid or other illicit drug use	6 (0)
7-day average incident COVID-19 cases	Daily reported incidence of new cases in health region, averaged over the seven days preceding hospital arrival. Reported in units of new cases/100,000 population	32 (0.1)

Appendix Table 3. Performance metrics for the CCEDRRN COVID-19 Infection Score for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the derivation cohort

Score	n (%)	Sensitivity (%, 95% CI)	Specificity (%, 95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	51 (0.2)	100 (99.6–100)	0.25 (0.2–0.3)	1.0	NA	0 (0)
≤-1	937 (4.3)	99.89 (99.4–100)	4.5 (4.2–4.8)	1.1	<0.1	1 (0.1)
≤0	5996 (27.6)	96.6 (95.2–97.7)	28.67 (28.1–29.3)	1.4	0.1	32 (0.5)
≤1	13114 (60.3)	86.6 (84.3–88.7)	62.43 (61.8–63.1)	2.3	0.2	126 (1.0)
≤2	18041 (83.0)	67.34 (64.2–70.3)	85.25 (84.8–85.7)	4.6	0.4	307 (1.7)
≤3	20405 (93.9)	45.11 (41.9–48.4)	95.61 (95.3–95.9)	10.3	0.6	516 (2.5)
Rule in:						
≥3	3702 (17.0)	67.34 (64.2–70.3)	85.25 (84.8–85.7)	4.6	0.4	633 (17.1)
≥4	1338 (6.2)	45.11 (41.9–48.4)	95.61 (95.3–95.9)	10.3	0.6	424 (31.7)
≥5	440 (2.0)	23.51 (20.8–26.4)	98.95 (98.8–99.1)	22.3	0.8	221 (50.2)
≥6	122 (0.6)	9.68 (7.9–11.8)	99.85 (99.8–99.9)	65.0	0.9	91 (74.6)
≥7	31 (0.1)	2.77 (1.8–4.0)	99.98 (99.9–100.0)	115.1	1.0	26 (83.9)
≥8	12 (0.1)	1.17 (0.6–2.1)	100 (100.0–100.0)	243.4	1.0	11 (91.7)
≥9	2 (<0.1)	0.21 (<0.1–0.8)	100 (100.0–100.0)	NA	1.0	2 (100)

LR+: Positive Likelihood Ratio; LR-: Negative Likelihood Ratio

Appendix Table 4. Performance metrics for CCEDRRN COVID Infection Score for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the combined cohort

Score cutoff	n (%)	Sensitivity (95% CI)	Specificity (95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	70 (0.3)	100 (99.7,100)	0.3 (0.2,0.3)	1	0	0 (0)
≤-1	1257 (4.5)	99.8 (99.4,100)	4.7 (4.5,5)	1.1	<0.1	2 (0.2)
≤0	7872 (28.5)	96.1 (94.8,97.1)	29.5 (29.0,30.1)	1.4	0.1	46 (0.6)
≤1	16962 (61.3)	85 (82.8,87.0)	63.4 (62.8,63.9)	2.3	0.2	175 (1.0)
≤2	23243 (84.0)	64.8 (62.0,67.5)	86.2 (85.7,86.6)	4.7	0.4	411 (1.8)
≤3	26169 (94.6)	40.3 (37.4,43.2)	96.1 (95.9,96.4)	10.4	0.6	697 (2.7)
Rule in:						
≥3	4422 (16.0)	64.8 (62.0,67.5)	86.2 (85.7,86.6)	4.7	0.4	756 (17.1)
≥4	1496 (5.4)	40.3 (37.4,43.2)	96.1 (95.9,96.4)	10.4	0.6	470 (31.4)
≥5	476 (1.7)	20.2 (18.0,22.6)	99.1 (99.0,99.2)	22.3	0.8	236 (49.6)
≥6	128 (0.5)	8 (6.5,9.7)	99.9 (99.8,99.9)	60.3	0.9	93 (72.7)
≥7	32 (0.1)	2.2 (1.5,3.2)	100 (100,100)	98.4	1.0	26 (81.3)
≥8	12 (<0.1)	0.9 (0.5,1.7)	100 (100,100)	249.8	1.0	11 (91.7)

LR+: Positive likelihood ratio; LR-: Negative likelihood ratio

Appendix Table 5. Performance metrics for the CORC score (race and ethnicity variables removed) for ruling in or ruling out SARS-CoV-2 infection at different score cut-off values in the combined cohort

Score	n (%)	Sensitivity (95% CI)	Specificity (95% CI)	LR+	LR-	COVID+ n (%)
Rule out:						
≤-2	202 (0.7)	99.9 (99.5–100)	0.8 (0.7–0.9)	1.01	0.11	1 (0.5)
≤-1	2715 (9.8)	97.4 (96.4–98.3)	10.1 (9.8–10.5)	1.08	0.25	30 (1.1)
≤0	9089 (32.9)	90.1 (88.2–91.7)	33.9 (33.3–34.4)	1.36	0.29	116 (1.3)
≤1	17582 (63.9)	72.8 (70.2–75.4)	65.2 (64.6–65.7)	2.09	0.42	317 (1.8)
≤2	23421 (84.7)	51.2 (48.3–54.1)	86.2 (85.8–86.7)	3.72	0.57	569 (2.4)
≤3	26224 (94.8)	27.7 (25.1–30.3)	95.8 (95.5–96)	6.56	0.76	844 (3.2)
Rule in:						
≥3	4244 (15.3)	51.2 (48.3–54.1)	86.2 (85.8–86.7)	3.72	0.57	598 (14.1)
≥4	1441 (5.2)	27.7 (25.1–30.3)	95.8 (95.5–96)	6.56	0.76	323 (22.4)
≥5	358 (1.3)	11.1 (9.3–13)	99.1 (99–99.2)	12.79	0.9	129 (36.0)
≥6	54 (0.2)	3.1 (2.2–4.2)	99.9 (99.9–100)	45.41	0.97	36 (66.7)


LR+: Positive likelihood ratio; LR-: Negative likelihood ratio

Appendix Table 6. Net Reclassification Improvement of the CCEDRRN COVID-19 Infection Score compared to the CORC Score (race and ethnicity variables removed)

Primary Outcome : Covid Positive					
CORC risk category		CCIS risk category			Total
		Low	Medium	High	
	Low	12	18	0	30
	Medium	34	539	241	814
	High	0	94	229	323
	Total	46	651	470	1167
Primary Outcome: Covid Negative					
CORC risk category		CCIS risk category			Total
		Low	Medium	High	
	Low	1593	1092	0	2685
	Medium	6233	15756	706	22695
	High	0	798	320	1118
	Total	7826	17646	1026	26498

<u>COVID Positive</u>		<u>COVID Negative</u>	
Number of outcomes	1167	Number of outcomes	26498
Correct reclassification	259	Correct reclassification	7031
Incorrect reclassification	128	Incorrect reclassification	1798
Net reclassification	131	Net reclassification	5233
Net reclassification improvement (Event)	0.112	Net reclassification improvement (Non-event)	0.197
Total net reclassification improvement			0.310

Appendix Figure 1. Receiver operating characteristic curves for the CCEDRRN COVID Infection Score (CCIS) and the CORC score (race and ethnicity variables removed) in the combined study cohort





Supplementary Table: Contributors to the Canadian COVID-19 Emergency Department Rapid Response Network

1. Purpose

This supplementary table provides details of the support staff at each of the participating institutions in the Canadian COVID-19 Emergency Department Rapid Response Network. This supplementary document should be attached to each peer-reviewed manuscript after the methods manuscript (M1). The purpose is to ensure research staffs and lead coordinators are appropriately recognized for their contributions to the network.

2. List of Support Staff

Table 1. Network coordinating centre staff at the University of British Columbia

Name	Roles	Contributions
Gelareh Ghaderi	Data analyst	Data processing and analysis for manuscripts.
Jeffrey Hau	Data manager	REDCap, data processing and analysis for manuscripts.
Vi Ho	National coordinator	Coordinate with provincial coordinators and training/onboarding of research assistants.
Joe Larkin	Project manager	Project management.
Fiona O'Sullivan	Data analyst	Data processing and analysis for manuscripts.
Serena Small	Research coordinator	Ethics & privacy reviews, data management plan, privacy impact assessment, and qualitative analyses
Amber Cragg	Research manager	Data and manuscript management
Wei Zhao	Data analyst	Data processing and analysis for manuscripts.
Vicky Wu	Data analyst	Data processing and analysis for manuscripts.
Elnaz Bodaghkhani	Research associate	Data and manuscript management

Table 2. Provincial Coordinators

Name	Province	Institutional affiliation	Contributions to CCEDRRN
Corinne DeMone	NS	Dalhousie University, Halifax, Nova Scotia	Research ethics board submission, manages research assistants, data cleaning and quality.
Jacqueline Fraser	NB	Dalhousie University, St. John New Brunswick	Site coordinator as well as research assistant.
Veronique Gélinas	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis	Provincial research coordinator, translation of research material to French, ethics management.

Canadian
COVID-19 ED
Network

Connie Taylor	ON	Queen's University, Kingston	Coordination of research assistants in Ontario, maintenance of REB applications for the province
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg	Lead RA for the province
Aimee Goss	SK	University of Saskatchewan, Saskatoon	Screens records in Saskatoon, data/extraction and entry, coordinates research assistants.
Hina Walia	AB	University of Calgary, Calgary	Provincial coordinator lead for Alberta, oversight of all Alberta sites.
Rajan Bola	BC	University of British Columbia, Vancouver	Provincial coordinator lead for BC, oversight of all BC sites.

Table 3. Institutional research assistant (RA) leads

Institutional RA leads are responsible for data extraction and integrity, communication with provincial leads.

Name	Province	Institutional affiliation(s)
Corinne DeMone	NS	Dartmouth General Hospital, Cobequid Community Health Centre, Hants Community Hospital Secondary Assessment Centers of the Dartmouth General Hospital, and Halifax Infirmary, Halifax
Jacqueline Fraser	NB	Saint John Regional Hospital, Saint John
Alexandra Nadeau	QC	CHU de Québec Université Laval, Quebec City
Audrey Nolet	QC	Centre intégré de santé et de services sociaux de Chaudière-Appalaches (Hôtel-Dieu de Lévis site), Lévis
Xiaoqing Xue	QC	Jewish General Hospital, Montréal
David Iannuzzi	QC	McGill University Health Center, Montréal
Chantal Lanthier	QC	Hôpital du Sacré-Cœur de Montréal, Montréal
Konika Nirmalanathan	ON	University Health Network, Toronto
Vlad Latiu	ON	Kingston General Hospital, Hotel Dieu Hospital, Kingston
Joanna Yeung	ON	Sunnybrook Health Sciences Center, Toronto
Natasha Clayton	ON	Hamilton General Hospital, Juravinski Hospital, Hamilton
Tom Chen	ON	London Health Sciences Centre, London
Jenna Nichols	ON	Health Sciences North, Sudbury
Kate Mackenzie	MB	Health Sciences Centre, Winnipeg
Aimee Goss	SK	St. Paul's Hospital, Royal University Hospital, Saskatoon City Hospital, Saskatoon
Stacy Ruddell	AB	Foothills Medical Centre, Peter Lougheed Centre, Rockyview General Hospital, South Health Campus, Calgary
Natalie Runham	AB	University of Alberta Hospital, Edmonton

Canadian
COVID-19 ED
Network

Name	Province	Institutional affiliation(s)
Karlin Su	AB	Royal Alexandra Hospital/Northeast Community Health Center, Edmonton
Josie Kanu	BC	St. Paul's Hospital, Mount Saint Joseph, Vancouver
Bernice Huynh	BC	Abbotsford Regional Hospital and Cancer Center, Abbotsford
Amanda Swirhun	BC	Royal Columbian Hospital, New Westminster
Tracy Taylor	BC	Eagle Ridge Hospital and Health Care Centre, Port Moody
Mai Hayashi	BC	Royal Inland Hospital, Kamloops
Mackenzie Cheyne	BC	Kelowna General Hospital, Kelowna
Sarim Asim	BC	Surrey Memorial Hospital, Surrey
Katherine Lam	BC	Vancouver General Hospital, Vancouver
Kelsey Compagna	BC	Lions Gate Hospital, Vancouver

Table 4. Contributing Study Sites and Investigators

Lead Investigator	Contributing Site / Code	Member Investigators
Maritime		
Patrick Fok		
Nova Scotia		
Hana Wiemer	Halifax Infirmary/ 902	Patrick Fok
	Dartmouth General Hospital/ 903	Hana Wiemer
	Hants Community Hospital/ 904	Samuel Campbell
	Cobequid Community Health Centre/ 905	Kory Arsenaault
	Secondary Assessment Centers of Dartmouth General and Halifax Infirmary/ 908	Tara Dahn
New Brunswick		
Kavish Chandra	Saint John Regional Hospital/ 901	Kavish Chandra
Quebec		
Patrick Archambault	Hotel-Dieu de Lévis/ 701	Patrick Archambault
	Jewish General Hospital/ 702	Joel Turner
	Centre Hospitalier de l'Université Laval (CHU de Québec)/ 703	Éric Mercier
	L'hôpital Royal Victoria - Royal Victoria Hospital/ 705	Greg Clark
	Hôpital de l'Enfant-Jésus, CHU de Québec/ 706	Éric Mercier
	Hôpital du Saint-Sacrement, CHU de Québec/ 707	Éric Mercier
	Hôpital Saint-François d'Assise, CHU de Québec/ 708	Éric Mercier
	Hôtel-Dieu de Québec, CHU de Québec/ 709	Éric Mercier

Canadian
COVID-19 ED
Network

	IUCPQ: Institut universitaire de cardiologie et de pneumologie de Québec/ 710	Sébastien Robert
	Hôpital du Sacré-Coeur de Montreal/ 711	Raoul Daoust
Ontario		
Laurie Morrison & Steven Brooks	Sunnybrook/ 401	Ivy Cheng
	The Ottawa Hospital - Civic Campus/ 403	Krishan Yadav
	The Ottawa Hospital - General Campus/ 404	Krishan Yadav
	Kingston/Queens/ 406	Steven Brooks
	Hamilton General Hospital/ 407	Michelle Welsford
	Health Science North, Sudbury Ontario/ 408	Rob Ohle
	University Hospital – LHSC/ 409	Justin Yan
	North York General Hospital, Toronto/ 410	Rohit Mohindra
	Victoria Hospital – LHSC/ 412	Justin Yan
	Toronto Western Hospital/ 414	Megan Landes
Manitoba		
Tomislav Jelic	Health Sciences Centre/ 307	Tomislav Jelic
Saskatchewan		
Phil Davis	Pasqua Hospital, Regina/ 301	Ankit Kapur
	Regina General Hospital, Regina/ 302	Ankit Kapur
	St Paul's Hospital, Saskatoon/ 303	Phil Davis
	Royal University, Saskatoon/ 304	Phil Davis
	Saskatoon City Hospital, Saskatoon/ 305	Phil Davis
Alberta		
Andrew McRae	University of Alberta Hospital, Edmonton/ 201	Brian Rowe
	Foothills, Calgary/ 202	Katie Lin
	Rockyview, Calgary/ 203	Andrew McRae
	Peter Lougheed Centre/ 204	Andrew McRae
	South Campus, Calgary/ 205	Stephanie VandenBerg
	Northeast Community Health Centre, Edmonton/ 206	Jake Hayward, Jaspreet Khangura
		Royal Alexandra Hospital, Edmonton/ 306
British Columbia		
Corinne Hohl	Vancouver General Hospital/ 101	Daniel Ting
	Lions Gate Hospital/ 102	Maja Stachura
	Saint Paul's Hospital/ 103	Frank Scheuermeyer

Canadian
COVID-19 ED
Network

Mount St Joseph's/ 104	Frank Scheuermeyer
Surrey Memorial Hospital/ 105	Balijeet Braar/ Craig Murray
Royal Columbian Hospital/ 106	John Taylor
Abbotsford Regional Hospital/ 107	Ian Martin
Eagle Ridge Hospital/ 108	Sean Wormsbecker
Victoria General Hospital/ 109	Matt Bouchard
Royal Jubilee Hospital/ 110	Matt Bouchard
Nanaimo General Hospital/ 111	Matt Bouchard
Royal Inland Hospital/ 112	Ian Martin
Kelowna General / Hospital/ 115	Lee Graham

It was not possible for us to recruit Members from Newfoundland and Labrador, Northwest Territories, Nunavut, Prince Edward Island and Yukon at the time of the inception of the registry.