

Multigene interactions and the prediction of depression in the Wisconsin Longitudinal Study

Nicholas S Roetker,¹ James A Yonker,¹ Chee Lee,¹ Vicky Chang,¹ Jacob J Basson,³ Carol L Roan,¹ Taissa S Hauser,¹ Robert M Hauser,¹ Craig S Atwood^{2,3,4}

To cite: Roetker NS, Yonker JA, Lee C, *et al*. Multigene interactions and the prediction of depression in the Wisconsin Longitudinal Study. *BMJ Open* 2012;**2**: e000944. doi:10.1136/bmjopen-2012-000944

► Prepublication history and additional materials for this paper are available online. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-000944>).

Received 26 January 2012
Accepted 29 May 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

¹Department of Sociology, University of Wisconsin, Madison, Wisconsin, USA

²Geriatric Research, Education and Clinical Center, William S. Middleton Memorial Veterans Hospital, Madison, WI

³Department of Medicine, University of Wisconsin-Madison School of Medicine and Public Health, Madison, Wisconsin, USA

⁴School of Exercise, Biomedical and Health Sciences, Edith Cowan University, Joondalup, Australia

Correspondence to
Dr Craig S Atwood;
csa@medicine.wisc.edu

ABSTRACT

Objectives: Single genetic loci offer little predictive power for the identification of depression. This study examined whether an analysis of gene–gene ($G \times G$) interactions of 78 single nucleotide polymorphisms (SNPs) in genes associated with depression and age-related diseases would identify significant interactions with increased predictive power for depression.

Design: A retrospective cohort study.

Setting: A survey of participants in the Wisconsin Longitudinal Study.

Participants: A total of 4811 persons (2464 women and 2347 men) who provided saliva for genotyping; the group comes from a randomly selected sample of Wisconsin high school graduates from the class of 1957 as well as a randomly selected sibling, almost all of whom are non-Hispanic white.

Primary outcome measure: Depression as determined by the Composite International Diagnostic Interview—Short-Form.

Results: Using a classification tree approach (recursive partitioning (RP)), the authors identified a number of candidate $G \times G$ interactions associated with depression. The primary SNP splits revealed by RP (*ANKK1* rs1800497 (also known as *DRD2* Taq1A) in men and *DRD2* rs224592 in women) were found to be significant as single factors by logistic regression (LR) after controlling for multiple testing ($p=0.001$ for both). Without considering interaction effects, only one of the five subsequent RP splits reached nominal significance in LR (*FTO* rs1421085 in women, $p=0.008$). However, after controlling for $G \times G$ interactions by running LR on RP-specific subsets, every split became significant and grew larger in magnitude (OR (before) \rightarrow (after): men: *GNRH1* novel SNP: (1.43 \rightarrow 1.57); women: *APOC3* rs2854116: (1.28 \rightarrow 1.55), *ACVR2B* rs3749386: (1.11 \rightarrow 2.17), *FTO* rs1421085: (1.32 \rightarrow 1.65), *IL6* rs1800795: (1.12 \rightarrow 1.85)).

Conclusions: The results suggest that examining $G \times G$ interactions improves the identification of genetic associations predictive of depression. 4 of the SNPs identified in these interactions were located in two pathways well known to impact depression: neurotransmitter (*ANKK1* and *DRD2*) and neuroendocrine (*GNRH1* and *ACVR2B*) signalling. This study demonstrates the utility of RP analysis as an

ARTICLE SUMMARY

Article focus

■ Single genetic loci offer little predictive power for the identification of depression. This study examined whether an analysis of $G \times G$ interactions of SNPs in genes associated with depression and age-related diseases would identify significant interactions with increased predictive power for depression.

Key messages

■ Using a classification tree approach (RP), we identified a number of candidate $G \times G$ interactions associated with depression. After controlling for $G \times G$ interactions by running LR on RP-specific subsets, every split became significant and grew larger in magnitude. Four of the SNPs identified in these interactions were located in two pathways well known to impact depression: neurotransmitter (*ANKK1* and *DRD2*) and neuroendocrine (*GNRH1* and *ACVR2B*) signalling.

Strengths and limitations of this study

■ Our results suggest that examining $G \times G$ interactions improves the identification of genetic associations predictive of depression. This study demonstrates the utility of RP analysis as an efficient and powerful exploratory analysis technique for uncovering genetic and molecular pathway interactions associated with disease aetiology.

efficient and powerful exploratory analysis technique for uncovering genetic and molecular pathway interactions associated with disease aetiology.

INTRODUCTION

Depression is a widespread mental disorder associated with a host of undesirable health, social and economic outcomes. One in six Americans is diagnosed with depression in his or her lifetime.¹ While many environmental

factors—such as socioeconomic status, childhood abuse and major life events—have important ties with depression, so too does gender and many genetic and epigenetic factors, making the disorder heterogeneous in nature.² Another major risk factor for depression is age, with depression reaching its highest levels in adults aged 80 years and older.³

It has been demonstrated from twin studies that genetic factors typically account for 40%–70% of the risk for developing major depressive disorder, and adoption studies have confirmed the role of genetic risk factors in the development of major depressive disorder (see Zubenko *et al*⁴ and references therein). Genetic studies, including recent genome-wide association studies (GWAS), have identified genetic alterations in over 50 genes known to be associated with depression.⁵ However, individually, the genetic alterations found within these genes (primarily single nucleotide polymorphisms (SNPs)) have little predictive value. There is a similar lack of predictive value from GWAS of other major age-related diseases.⁶

Given this lack of predictive power among individual genetic alterations for depression together with the complex nature of ageing-related diseases, it would seem prudent to examine epistatic effects on this age-related condition. In this respect, we have previously demonstrated that gene-gene ($G \times G$) interactions greatly modulate risk for complex age-related diseases.^{7,8} Recent studies of depression also have identified epistatic effects. In particular, associations have been identified between *BDNF* Val66Met (brain-derived neurotrophic factor; rs6265) and *5-HTTLPR* (serotonin transporter-linked promoter region⁹); *GSK3B* rs6782799 (glycogen synthase kinase 3 β), *BDNF* rs7124442 and *BDNF* Val66Met¹⁰; *BDNF* Val66Met and SNPs in *NTRK2* (neurotrophic tyrosine kinase receptor 2¹¹); and *5-HTTLPR* short allele and a chromosome 4 gene.¹² The machine learning tool recursive partitioning (RP) has recently been used by Wong *et al*¹³ to assess complex $G \times G$ interactions in depression. Wong *et al* note that RP is useful in that it quickly explores high-dimensional data for non-linear effects that are non-biased and easily interpretable.

The goals of this study were therefore to (1) explore $G \times G$ interactions that might better predict the genetic factors involved in the aetiology of depression and (2) to further demonstrate the utility of machine learning algorithms (RP) to identify genetic interactions. Using genotypic data from the Wisconsin Longitudinal Study (WLS), we identified associations between dopaminergic genes and depression in men and women, as well as $G \times G$ interactions involving neuroendocrine signalling pathways, with increased significance compared with single genetic associations.

METHODS

Study participants and surveys

Data were collected from the WLS, a random sample originally comprised 10,317 men and women who

graduated from Wisconsin high schools in 1957. Later in 1977, the WLS began interviewing one randomly selected sibling of each graduate, when possible. The cohort reflects the ancestral makeup of the late-1950s Wisconsin population in that participants are almost entirely non-Hispanic white men and women. In general, the sample is broadly representative of older white Americans with at least a high school education.¹⁴ Further characteristics of the WLS cohort may be found in detail elsewhere.¹⁵ Health and psychological well-being phenotypic data were taken from mail and phone surveys given in 2004–2005. Inclusion criteria for depression included any member of the WLS cohort who was depressed according to the Composite International Diagnostic Interview–Short-Form. Individuals who answered YES to the question ‘Have you ever had a time in life lasting two weeks or more when nearly every day you felt sad, blue, depressed, or when you lost interest in most things like work, hobbies, or things you usually liked to do for fun?’ and whose depression was not caused by alcohol, drugs, medications or physical illness were asked further depression symptom questions. Symptom questions asked whether the 2-week period was accompanied with (1) any weight loss, (2) trouble sleeping, (3) feeling tired, (4) feeling bad upon waking, (5) losing interest, (6) trouble concentrating or (7) thoughts about death. Those answering YES to three or more of these symptom questions were classified as having depression.¹⁶ Those answering YES to two or fewer symptom questions and all those answering NO to the initial stem question were classified as controls.

Genotyping

Seven thousand one hundred and one participants (4569 graduates and 2532 siblings) provided saliva samples in Oragene DNA sample collection kits (DNA Genotek, Kanata, Canada) from which DNA was extracted and genotyped for 78 SNPs that were selected based on their association with depression and age-related conditions and diseases (see supplementary information 1). Genotyping was performed by KBioscience (Hoddesdon, UK) with use of a homogeneous Fluorescent Resonance Energy Transfer technology coupled to competitive allele-specific PCR. All SNP genotypes described in our results were in Hardy–Weinberg equilibrium and their frequencies matched those reported in the literature for European samples.

Statistical analysis

Analyses were limited to the 4811 pooled graduates and siblings for whom we had depression and genotype information (note: individuals with more than 10% missing genotype data were not included). The average age among this sample was just younger than 65 years in 2004. Eighty per cent were married, and the average amount of post-high school educational attainment was 2 years. Median household income in 1993 was \$56 700.

Recursive partitioning

RP is a data mining tool for revealing trends that relate a dependent variable (depressed vs non-depressed) to various predictor variables (SNPs). Zhang and Bonney¹⁷ have shown how RP can be used in genetic association studies to identify disease genes. RP helps control for heterogeneity in the population and confounding factors by allowing for the segregation of the sample population according to any condition. Thus, RP is a useful way to handle complex data sets that might confound regression analysis due to the complexity of the relationship between the independent and dependent variables and due to missing information.

RP classification trees (using R package rpart) were used to identify potential interactions among the 78 SNPs in relation to depression. The trees split the data along branches according to the criteria determined by the rpart package algorithm, which is originally based off the work of Breiman's classification and regression trees algorithm.¹⁸ Basically, the classification and regression trees algorithm first considers all depressed and non-depressed subjects pooled together in a heterogeneous root node. Based on considering every possible 'yes–no' binary partition that can be made by each independent variable, the single split, which maximises homogeneity between the two resulting subnodes as compared with the root node, is made. Each subnode can then be treated independently as a new root node for all subsequent splits, and the pattern continues until every subject constitutes a terminal node, resulting in a very large and complex tree. A 10-part cross-validation procedure seeking to minimise misclassification and complexity determines optimal pruning. See Therneau and Atkinson¹⁹ for specific details of the rpart package. Priors were set to 0.5, 0.5. The use surrogate parameter was set to 0 so that subjects missing the primary split variable do not progress further down the tree, and maxsurrogate was set to 0 to cut computation time in half. The threshold complexity parameter was set to 0.01. Tree nodes were re-created in Microsoft Visio to display percentage depressed and the default number of controls/cases as presented by the rpart.

Logistic regression

Variables found in association with depression based on RP analysis were considered in single-factor logistic regression (LR) models, separate by gender, using the specific dichotomous splitting of genotypes as designated by RP trees. Regression models for all seven SNP splits were first run on the full data set to represent single main factor effects. Then, each split was run on the respective subset of data as represented by the preceding RP split criteria. Thus, we attempt to mirror RP splits within a more formal LR framework in order to measure the significance of interactions presented by the trees. Multiple testing of 78 SNPs in RP for both men and women followed by 14 LR models resulted in a modified false discovery rate (FDR) significance level of 0.009.

RESULTS

Of the 4811 participants (2464 women and 2347 men) under examination in this study, we identified 713 participants (481 women and 232 men) with depression (14.8%). Given that the independent variable gender (when included as a factor in the full data set) was the primary split on RP trees; that women are over two times as likely to be diagnosed with depression than men and since the female aetiology of depression has been reported to be associated with unique social, psychological, and biological factors,²⁰ all subsequent analyses were performed by gender.

Recursive partitioning analysis

To examine multigene interactions for association with depression, we screened our data set using RP. The two-factor RP tree (*ANKK1/GNRH1*) was the optimised pruning for men (figure 1), while the five-factor tree (*DRD2/APOC3/ACVR2B/FTO/IL6*) was the optimised pruning for women (figure 2). For more detailed information on the seven SNPs found by RP, see supplementary information 2.

The best overall split for men was *ANKK1* rs1800497 (historically known as the *DRD2* Taq1A allele), where the incidence of depression increased 2.2-fold in those with




Figure 1 Recursive partitioning tree of Composite International Diagnostic Interview–Short-Form depression in men of the Wisconsin Longitudinal Study. Upper and lower numbers in nodes represent the percentage of participants with depression and the number of controls/cases in that node, respectively. Blue and purple boxes/circles indicate lower and higher rates of depression relative to the primary node, respectively. Split information indicates gene, single nucleotide polymorphism (SNP), and genotype criteria, respectively. M1 is subset of data referenced in table 1. Sensitivity: 0.526, specificity: 0.598, accuracy: 0.591. Due to missing genotype information, we lose approximately 1.5% of participants per split. *rs1800497 is historically known as the *DRD2* Taq1A allele.




Figure 2 Recursive partitioning tree of Composite International Diagnostic Interview—Short-Form depression in women of the Wisconsin Longitudinal Study. Upper and lower numbers in nodes represent the percentage of participants with depression and the number of controls/cases in that node, respectively. Blue and purple boxes/circles indicate lower and higher rates of depression relative to the primary node, respectively. Split information indicates gene, single nucleotide polymorphism (SNP), and genotype criteria, respectively. F1–F4 are subsets referenced in table 1. Sensitivity: 0.607, specificity: 0.563, accuracy: 0.572. Due to missing genotype information, we lose approximately 1.4% of participants per split.

no C-alleles compared with those with one or two C-alleles. Considering interaction between *ANKK1* and *GNRH1* widened the disparity in incidence, where those with at least one C-allele in both *ANKK1* rs1800497 and the novel SNP in *GNRH1* had a 2.7-fold lower incidence than those without a C-allele in *ANKK1* rs1800497.

For women, the best overall split was *DRD2* rs2242592, where those with one or two C-alleles had 1.3-fold higher incidence of depression compared with those without any C-alleles. $G \times G$ interactions associated with the highest incidence of depression included *DRD2* rs2242592 T/T + *APOC3* rs45537037 T/T + *ACVR2B* rs3749386 C/C or T/T, accounting for a 1.4-fold increase in depression compared with baseline incidence.

Single main factor effects

Specific SNP interactions identified by RP were next analysed by LR (see table 1 for full data). The primary SNP splits in men and women were significant at the modified FDR level. Men with no C-alleles for *ANKK1* rs1800497 had 2.55 times higher odds ($p=0.001$ (95% CI 1.44 to 4.51)) of depression compared with men with at least one C-allele. Women with at least one C-allele for *DRD2* rs2242592 had 1.32 times higher odds ($p=0.006$ (95% CI 1.08 to 1.62)) of depression compared with women with no C-alleles. One other split reached nominal significance; women homozygous (C/C or T/T) for *FTO* rs1421085 had 1.32 times higher odds

($p=0.008$ (95% CI 1.08 to 1.62)) for depression than women with a heterozygous genotype. SNP splits of *GNRH1*, *APOC3*, *ACVR2B*, and *IL6* did not significantly associate with depression.

Gene–gene interactions enhance predictability for depression

Specific SNP interactions identified by RP were next analysed by LR as RP-specific subsets (see table 1, RP-subsetted data). All five of the secondary and tertiary RP splits were found to be significant at the modified FDR level when considered as subsets. Among only men with at least one C-allele in *ANKK1* rs1800497, those with no C-allele in the novel SNP of *GNRH1* had 1.57 times higher odds ($p=0.002$ (95% CI 1.18 to 2.08)) for depression than men with one or two C-alleles. For the subset of women in the first right-hand split of figure 2, those homozygous for *FTO* rs1421085 had 1.65 times higher odds ($p=0.0005$ (95% CI 1.24 to 2.18)) for depression than women with a heterozygous genotype. For the remaining subset of women in the second right-hand split of figure 2, those homozygous for *IL6* rs1800795 had 1.85 times higher odds ($p=0.006$ (95% CI 1.19 to 2.89)) for depression than women with a heterozygous genotype. For the subset of women in the first left-hand split of figure 2, those with no C-alleles for *APOC3* rs45537037 had 1.55 times higher odds ($p=0.004$ (95% CI 1.15 to 2.09)) for depression than women with one or two C-alleles. For the subset of women in the second left-hand split of figure 2, those homozygous for *ACVR2B* rs3749386 had 2.17 times higher odds ($p=0.001$ (95% CI 1.37 to 3.44)) for depression than women with a heterozygous genotype.

DISCUSSION

Using RP as a screening tool to find potential multigene interactions, followed by verification by LR, our data demonstrate that multigene interactions predict depression with a greater certainty than single main factor associations. RP provided us with primary dichotomous genotype splits in men and women (*ANKK1* rs1800497 and *DRD2* rs2242592, respectively) that were both significant in LR models at the modified FDR level (table 1). When considering the five subsequent RP splits over the entire data set with LR, only one reached a nominal level of significance (barely), which was *FTO* rs1421085 in women. However, after running LR on specific subsets of data according to the pattern of RP branches, every split was found to be significant and every OR grew larger (table 1; OR (before) → (after): male left: 1.43 → 1.57, female left 1: 1.28 → 1.55, female left 2: 1.11 → 2.17, female right 1: 1.32 → 1.65, female right 2: 1.12 → 1.85). Thus, RP provides two unique and important criteria: dichotomous genotype splitting instructions and $G \times G$ interaction patterns. These criteria go beyond the traditional single-factor SNP approach to genetic association studies and allow identification of important multigene pathways that more suitably characterise the aetiology of complex diseases.

Table 1 Single-factor LR models based directly off male and female RP tree split criteria (see figures 1 and 2)

Gender	RP split	Gene	SNP	Genotypes	Full data		RP-subsetted data		
					OR (95% CI)	p Value	Subset	OR (95% CI)	p Value
Male	Primary	ANKK1#	rs1800497	T/T versus C/C + C/T	2.55 (1.44 to 4.51)	0.001*	—	—	—
	Left	GNRH1	novel SNP	T/T versus C/C + T/C	1.43 (1.09 to 1.88)	0.011	M1	1.57 (1.18 to 2.08)	0.002*
Female	Primary	DRD2	rs2242592	C/C + T/C versus T/T	1.32 (1.08 to 1.62)	0.006*	—	—	—
	Left 1	APOC3	rs2854116	T/T versus C/C + T/C	1.28 (1.04 to 1.57)	0.018	F1	1.55 (1.15 to 2.09)	0.004*
	Left 2	ACVR2B	rs3749386	C/C + T/T versus T/C	1.11 (0.91 to 1.36)	0.302	F2	2.17 (1.37 to 3.44)	0.001*
	Right 1	FTO	rs1421085	C/C + T/T versus T/C	1.32 (1.08 to 1.62)	0.007*	F3	1.65 (1.24 to 2.18)	0.0005*
	Right 2	IL6	rs1800795	C/C + G/G versus C/G	1.12 (0.92 to 1.37)	0.269	F4	1.85 (1.19 to 2.89)	0.006*

Each SNP split was first run on the full data set to represent single main factor effects ('full data') for both men and women. Then, the same SNP splits were run on specific subsets of data per RP tree splits (M1, F1–F4; 'RP-subsetted data').

M1: LR analysis was run for only those with genotype DRD2 rs1800497 C/C or C/T.

F1: LR analysis was run for only those with genotype DRD2 rs2242592 T/T.

F2: LR analysis was run for only those with genotypes DRD2 rs2242592 T/T and APOC3 rs2854116 T/T.

F3: LR analysis was run for only those with genotype DRD2 rs2242592 C/C or T/C.

F4: LR analysis was run for only those with genotypes DRD2 rs2242592 C/C or T/C and FTO rs1421085 T/C.

#rs1800497 is historically known as the DRD2 Taq1A allele.

*p<0.009

LR, logistic regression; RP, recursive partitioning; SNP, single nucleotide polymorphism.

The utility of RP and LR for identification of gene–gene interactions

With recent advances in genotyping allowing for high-dimensional SNP identification, it is now possible to examine genetic data sets for single main factor effects and also for $G \times G$ interactions. The requirement for $G \times G$ analyses as a better predictor of age-related diseases is obvious from the standpoint that humans are complex biological systems composed of numerous molecular interactions and from recent studies indicating disease risk is modulated by $G \times G$ interactions.⁷ Notwithstanding this, the development of analytical tools for the identification of $G \times G$ interactions has not kept pace with the technological advances in identifying genetic alterations among individuals. In this respect, we have previously used multifactor dimensionality reduction (MDR), LR and linkage disequilibrium (LD) to identify $G \times G$ interactions among a small set of SNPs.⁷ However, large data sets require a screening tool to identify potential multigene interactions. In this study, we have used RP to screen for multigene interactions, a data mining technique that is currently underused in genetic studies. RP serves as an efficient and powerful exploratory analysis technique, especially when looking for interactions in data sets with a large number of independent variables. This screening allows for the identification of $G \times G$ interactions (with greater explanatory power) that might otherwise not have been identified and that can then be confirmed using more traditional statistical techniques. As illustrated in this paper, this data mining methodology has the advantage of identification of genetic interactions *between* pathways involved in the aetiology of depression, in keeping with the etiological heterogeneity of this disorder (see later).

Our study provides proof of principle for the use of RP in higher dimensional analyses such as GWAS, where a comprehensive list of SNPs may fully explore genetic predisposition to depression and other age-related disease. The WLS is an ideal candidate for future GWAS studies, given its large sample size, rich covariate composition and longitudinal nature.

In this genetic study, we aimed to identify underlying genetic predispositions to depression and thus have not yet tested environmental, health, socio-behavioral or other non-genetic factors. Future analyses using RP to examine the impact of these factors on the development of depression would be anticipated to identify gene–non-genetic factor interactions. Indeed, the predictive gains of $G \times G$ analyses were stronger for men than for women, despite the fact that depression occurs disproportionately in women (~2:1 female-to-male ratio^{21–25}). This suggests that environmental factors may be needed in addition to genetic factors in understanding the aetiological pathways for women. Indeed, biological factors such as hormonal changes related to reproductive status^{26 27} may impact environmental factors such as psychosocial experiences (trauma, stress, interpersonal relationships, etc) and general health issues in the development of depression.

Genetic and biological correlates of depression

Numerous studies have identified SNPs that associate with depression. Many of the SNPs associated with depression from other studies were not significantly associated in our study. This is perhaps not surprising since a single factor is unlikely to provide consistent association especially in a complex condition such as depression, where multiple pathways intersect in regulating the risk of the disease. For example, if a SNP within the serotonin pathway also requires a SNP in the glutamatergic pathway in order for the patient to present with depression, the presence of either SNP in the absence of the other will not be predictive of depression. Moreover, as indicated by Shi and Weinberg,²⁸ since the human genome contains genetic redundancy, disruption of a single gene may be selectively neutral, but the malfunction of several genes in a pathway might result in expression of a particular phenotype.

Both the primary splits in men and women were SNPs linked with *DRD2* (dopamine receptor D2), a gene that has previously been linked with depression and social phobia.^{29–31} The primary male genotype split rs1800497, technically found in gene *ANKKI*, is historically known as the *DRD2* Taq1A allele because of its known association with decreased dopamine receptor D2 density (in those with T-alleles).^{32–35} The Taq1A allele has also been previously associated with depressive symptoms in children, where those with the A1 allele (T) were more likely to have depressive symptoms.³⁶ We saw a similar association between A1 and depression in WLS men, where those with two A1 alleles had 2.6 times higher odds for depression compared with those with one or no A1 alleles. The primary split in women (*DRD2* rs2242592) has previously been found to be associated with schizophrenia, where the C-allele was associated with higher susceptibility for the disease.³⁷ Interestingly, this same study also found the Taq1A allele to also associate with schizophrenia.

The secondary and tertiary right-hand splits in the female RP tree—*FTO* (fat mass and obesity associated) rs1421085 and *IL6* (interleukin 6) rs1800795—have also been found to relate with mental illness and depression in previous studies.^{38–39} There is evidence that activin receptor signalling also is involved in affective disorders, especially when considering interaction with GABAergic pathways.⁴⁰ Although we did not see an interaction between SNPs in GABA/activin receptor genes and depression, *ACVR2B* was associated with depression in women. No previous associations between depression and *APOC3*, *ACVR2B*, or *GNRH1* have been reported.

That these genetic variants are associated with *neuroendocrine* pathways (*GNRH1*, *ACVR2B*) that are known to regulate *neurotransmitter* release and cognitive behaviour^{39–40} supports these associations as relevant to the aetiology of depression and underlines the benefits of using RP to identify meaningful G × G interactions associated with disease.

Limitations

Given the numerous genetic and non-genetic influences that are linked to depression and the small number of SNPs analysed, it is not surprising that predictability from our models was low (although our predictability was superior to previous studies examining only single main factors). Also, the predictive value of our statistical models was further limited due to user bias in selection of SNPs (from nearly 2 million SNPs in the human genome) used in this study. As a result of this, interactions we have found could potentially be moderated by another gene that we have not considered in this study. Nonetheless, we identified significant G × G interactions between known, and newly identified, loci associated with depression. Importantly, four of the seven SNPs identified in these interactions were primarily located in two pathways well known to impact depression: neurotransmitter and neuroendocrine signalling.

The results from the RP analyses conducted in this study were confirmed by LR, demonstrating the utility of RP as a screening tool for identifying meaningful G × G interactions. Future development of algorithms for RP analysis should maximise the distance between branches of the next best split (ie, rpart) and consider subsequent future split combinations that could potentially result in trees with ‘better’ overall predictability.

Summary

Our data indicate that G × G interaction analyses allow for enhanced predictability of conditions and diseases of ageing. RP is an efficient and powerful exploratory analysis technique for elucidating G × G interactions in large data sets and combined with LR provides an important statistical analysis for the identification of well-supported G × G interactions. We predict that such analytical methods will play an increasingly important role in the identification of epistatic effects in future GWAS. Finally, our studies illustrate how RP analyses can be used to find interacting pathways involved in the aetiology of a disease or condition such as depression.

Acknowledgements This research uses data from the Wisconsin Longitudinal Study (WLS) of the University of Wisconsin-Madison. Since 1991, the WLS has been supported principally by the National Institute on Aging (AG09775, AG21079 and AG33285), with additional support from the Vilas Estate Trust, the National Science Foundation, the Spencer Foundation and the Graduate School of the University of Wisconsin-Madison. A public use file of data from the WLS is available from the Wisconsin Longitudinal Study, University of Wisconsin-Madison, 1180 Observatory Drive, Madison, Wisconsin 53706 and at <http://www.ssc.wisc.edu/wlsresearch/data/>. This material is the result of work supported with resources at the William S. Middleton Memorial Veterans Hospital, Madison, Wisconsin. The opinions expressed herein are those of the authors. The contents do not represent the views of the Department of Veterans Affairs or the US Government. This is Geriatrics Research, Education and Clinical Center VA paper # 2012–03.

Contributors CSA, RMH and TSH conceptualised the study. RMH, TSH, CLR, NSR, CL and CSA collected saliva samples and performed genotyping analyses. NSR, JAY, CL, VC and JJB performed statistical analyses on the Wisconsin Longitudinal Study data set. CSA and RMH directed the statistical analyses. NSR and CSA drafted the manuscript. All authors critically reviewed the manuscript and approved the final version.

Funding The study was supported by the National Institute on Aging (AG09775, AG21079 and AG33285).

Competing interests None.

Patient consent Obtained.

Ethics approval Ethics approval was provided by the Social Sciences Institutional Review Board, UW-Madison.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement WLS public release data is available for download at <http://www.ssc.wisc.edu/wlsresearch>. Information on obtaining WLS genotypic data is available at this site. All WLS data is available free of charge.

REFERENCES

1. CDC. *Anxiety and Depression*. Atlanta, GA: Centers for Disease Control and Prevention, 2009. <http://www.cdc.gov/Features/dsBRFSSDepressionAnxiety/> (accessed 18 Oct 2010).
2. Eley TC, Sugden K, Corsico A, *et al*. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Mol Psychiatry* 2004;9:908–15.
3. Mirowsky J, Ross CE. Age and depression. *J Health Soc Behav* 1992;33:187–205.
4. Zubenko GS, Zubenko WN, Spiker DG, *et al*. Malignancy of recurrent, early-onset major depression: a family study. *Am J Med Genet* 2001;105:690–9.
5. Raymer KA, Waters RF, Price CR. Proposed multigenic Composite Inheritance in major depression. *Med Hypotheses* 2005;65:158–72.
6. Harold D, Abraham R, Hollingworth P, *et al*. Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nat Genet* 2009;41:1088–93.
7. Haasl R, Ahmadi MR, Meethal S, *et al*. A luteinizing hormone receptor intronic variant is significantly associated with decreased risk of Alzheimer's disease in males carrying an apolipoprotein E epsilon4 allele. *BMC Med Genet* 2008;9:37.
8. Basson J, Wilson A, Haasl R, *et al*. Multi locus interactions in steroidogenic pathway genes predict Alzheimer's disease. *Alzheimer's Dementia* 2008;4(Suppl 2):T579, P3–199.
9. Pezawas L, Meyer-Lindenberg A, Goldman AL, *et al*. Evidence of biologic epistasis between BDNF and SLC6A4 and implications for depression. *Mol Psychiatry* 2008;13:709–16.
10. Zhang K, Yang C, Xu Y, *et al*. Genetic association of the interaction between the BDNF and GSK3B genes and major depressive disorder in a Chinese population. *J Neural Transm* 2010;117:393–401.
11. Lin E, Hong CJ, Hwang JP, *et al*. Gene-gene interactions of the brain-derived neurotrophic-factor and neurotrophic tyrosine kinase receptor 2 genes in geriatric depression. *Rejuvenation Res* 2009;12:387–93.
12. Neff CD, Abkevich V, Potter J, *et al*. Evidence for epistasis between SLC6A4 and a chromosome 4 gene as risk factors in major depression. *Am J Med Genet B Neuropsychiatr Genet* 2010;153B:321–2.
13. Wong ML, Dong C, Andreev V, *et al*. Prediction of susceptibility to major depression by a model of interactions of multiple functional genetic variants and environmental factors. *Mol Psychiatry* 2012;17:624–33.
14. Carr D, Khodyakov D. End-of-life health care planning among young-old adults: an assessment of psychosocial influences. *J Gerontol B Psychol Sci Soc Sci* 2007;62:S135–S41.
15. Sewell WH. *As we age: the Wisconsin longitudinal study, 1957-2001*. Madison, WI: Center for Demography and Ecology University of Wisconsin-Madison, 2001.
16. Nelson CB, Kessler RC, Mroczek D. *Scoring the World Health Organization's Composite International Diagnostic Interview short form (CIDI-SF; v1.0 NCV98)*. 2001. http://www.hcp.med.harvard.edu/nfs/ftpdir/cidif_readme.pdf (accessed 27 Jan 2012).
17. Zhang H, Bonney G. Use of classification trees for association studies. *Genet Epidemiol* 2000;19:323–32.
18. Breiman L, Friedman J, Olshen R, *et al*. *Classification and Regression Trees*. Boca Raton: Chapman and Hall/CRC, 1984.
19. Therneau T, Atkinson E, eds. *Technical Report Series No. 61, An Introduction to Recursive Partitioning Using the RPART Routines*. Rochester, MN: Department of Health Science Research, Mayo Clinic, 1997.
20. Accortt EE, Freeman MP, Allen JJB. Women and major depressive disorder: clinical perspectives on causal pathways. *J Womens Health (Larchmt)* 2008;17:1583–90.
21. Kessler RC, McGonagle KA, Swartz M, *et al*. Sex and depression in the National Comorbidity Survey. I: lifetime prevalence, chronicity and recurrence. *J Affect Disord* 1993;29:85–96.
22. Kessler RC, McGonagle KA, Nelson CB, *et al*. Sex and depression in the National Comorbidity Survey. II: cohort effects. *J Affect Disord* 1994;30:15–26.
23. Kessler RC, McGonagle KA, Zhao S, *et al*. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994;51:8–19.
24. Weissman MM, Bland R, Joyce PR, *et al*. Sex differences in rates of depression: cross-national perspectives. *J Affect Disord* 1993;29:77–84.
25. Weissman MM, Olfson M. Depression in women: implications for health care research. *Science* 1995;269:799–801.
26. Pajer K. New strategies in the treatment of depression in women. *J Clin Psychiatry* 1995;56(Suppl 2):30–7.
27. Paykel ES. Depression in women. *Br J Psychiatry Suppl* 1991:22–9. Review.
28. Shi M, Weinberg CR. How much are we missing in SNP-by-SNP analyses of genome-wide association studies? *Epidemiology* 2011;22:845–7.
29. Lawford BR, Young R, Noble EP, *et al*. The D2 dopamine receptor (DRD2) gene is associated with co-morbid depression, anxiety and social dysfunction in untreated veterans with post-traumatic stress disorder. *Eur Psychiatry* 2006;21:180–5.
30. Klimek V, Schenck JE, Han H, *et al*. Dopaminergic abnormalities in amygdaloid nuclei in major depression: a postmortem study. *Biol Psychiatry* 2002;52:740–8.
31. Schaefer FR, Liebowitz MR, Abi-Dargham A, *et al*. Low dopamine D(2) receptor binding potential in social phobia. *Am J Psychiatry* 2000;157:457–9.
32. Noble EP, Blum K, Ritchie T, *et al*. Allelic association of the D2 dopamine receptor gene with receptor-binding characteristics in alcoholism or gene ism. *Arch Gen Psychiatry* 1991;48:648–54.
33. Thompson J, Thomas N, Singleton A, *et al*. D2 dopamine receptor gene (DRD2) Taq1 A polymorphism: reduced dopamine D2 receptor binding in the human striatum associated with the A1 allele. *Pharmacogenetics* 1997;7:479–84.
34. Pohjalainen T, Rinne JO, Nagren K, *et al*. The A1 allele of the human D2 dopamine receptor gene predicts low D2 receptor availability in healthy volunteers. *Mol Psychiatry* 1998;3:256–60.
35. Jonsson EG, Nothen MM, Grunhage F, *et al*. Polymorphisms in the dopamine D2 receptor gene and their relationships to striatal dopamine receptor density of healthy volunteers. *Mol Psychiatry* 1999;4:290–6.
36. Hayden EP, Klein DN, Dougherty LR, *et al*. The dopamine D2 receptor gene and depressive and anxious symptoms in childhood: associations and evidence for gene-environment correlation and gene-environment interaction. *Psychiatr Genet* 2010;20:304–10.
37. Dubertret C, Bardel C, Ramoz N, *et al*. A genetic schizophrenia-susceptibility region located between the ANKK1 and DRD2 genes. *Prog Neuropsychopharmacol Biol Psychiatry* 2010;34:492–9.
38. Kivimaki M, Jokela M, Hamer M, *et al*. Examining overweight and obesity as risk factors for common mental disorders using fat mass and obesity-associated (FTO) Genotype-Instrumented Analysis. *Am J Epidemiol* 2011;173:421–9.
39. Bull SJ, Huezo-Diaz P, Binder EB, *et al*. Functional polymorphisms in the interleukin-6 and serotonin transporter genes, and depression and fatigue induced by interferon-alpha and ribavirin treatment. *Mol Psychiatry* 2009;14:1095–104.
40. Zheng F, Adelsberger H, Muller MR, *et al*. Alzheimer C. Activin tunes GABAergic neurotransmission and modulates anxiety-like behavior. *Mol Psychiatry* 2008;14:332–46.

Supplementary Table 1. Single Nucleotide Polymorphisms Assessed in the WLS

Gene	Encodes	SNP	Associated disease/behavior
<i>A2M</i>	alpha-2-macroglobulin	rs669	Alzheimer's disease (1)
<i>ACVR2A</i>	activin receptor IIA	rs1424954	pre-eclampsia (2)
<i>ACVR2B</i>	activin receptor IIB	rs3749386	--
<i>ADIPOQ</i>	adiponectin, C1Q and collagen domain containing	rs1501299	diabetes II (3, 4), obesity (5, 6), breast cancer (7)
<i>ADIPOQ</i>	adiponectin, C1Q and collagen domain containing	rs2241766	diabetes II (3, 4), obesity (8), breast cancer (7)
<i>ACVRL1</i>	activin receptor-like kinase 1	rs2071219	brain arteriovenous malformations (9)
<i>APOC-3</i>	apolipoprotein C-III	rs2854116	nonalcoholic fatty liver disease (10)
<i>ApoE</i>	apolipoprotein E	rs429358	Alzheimer's disease (11, 12)
<i>ApoE</i>	apolipoprotein E	rs7412	Alzheimer's disease (11, 12)
<i>AR</i>	androgen receptor	rs6152	male pattern baldness (13)
<i>BCKDHB</i>	branched chain keto acid dehydrogenase E1, beta polypeptide	rs4502885	premature ovarian failure (14)
<i>BDNF</i>	brain-derived neurotrophic factor	rs6265	depression (15-17), alcohol dependence-related depression (18), bipolar disorder (19), schizophrenia (20), cognition (21), BMI (22)
<i>BDNF</i>	brain-derived neurotrophic factor	rs908867	antidepressant response (23)
<i>BRCA1</i>	breast cancer 1, early onset	rs1799966	breast cancer (24)
<i>BRCA2</i>	breast cancer 2, early onset homolog	rs144848	breast cancer (24)
<i>CH25H</i>	cholesterol 25-hydroxylase	rs3802657	--
<i>CHRM2</i>	cholinergic receptor, muscarinic 2	rs2061174	alcohol dependence, depression (25)
<i>CHRM2</i>	cholinergic receptor, muscarinic 2	rs8191992	cognition (26)
<i>COMT</i>	catechol-O-methyltransferase	rs4680	ADHD (27), substance abuse (28-31), depression (32), antidepressant response (33), bipolar disorder (34), cognition (35)
<i>CTSD</i>	cathepsin D	rs17571	Alzheimer's disease (36)
<i>CYP11A1</i>	cytochrome P450, family 11, subfamily A, polypeptide 1	rs8039957	breast cancer (37)
<i>CYP11B2</i>	cytochrome P450, family 11, subfamily B, polypeptide 2	rs1799998	stroke (38), cardiovascular disease (39)
<i>DAT1</i>	human dopamine transporter	rs11564774	ADHD (40)
<i>DAT1</i>	human dopamine transporter	rs2963238	alcohol-withdrawal seizures (41)
<i>DISC1</i>	disrupted in schizophrenia 1	rs821616	schizophrenia (42), cognitive aging (43)
<i>DRD2</i>	dopamine receptor D2	rs17529477	--
<i>DRD2/ANKK1</i>	dopamine receptor D2/ ankyrin repeat and kinase domain containing 1	rs1800497	obesity, drug addiction (44)
<i>DRD2</i>	dopamine receptor D2	rs2242592	schizophrenia (45)
<i>DRD2</i>	dopamine receptor D2	rs4245147	--
<i>DRD2</i>	dopamine receptor D2	rs6277	schizophrenia (46), PTSD (47)
<i>DRD4</i>	dopamine receptor D4	rs1800955	ADHD (48), heroine addiction (49)
<i>DTNBP1</i>	dystrobrevin-binding protein 1	rs1018381	schizophrenia (50), cognitive ability (51)
<i>DTNBP1</i>	dystrobrevin-binding protein 1	rs760761	schizophrenia (52)
<i>ESR1</i>	estrogen receptor 1	rs7761133	--
<i>ESR1</i>	estrogen receptor 1	rs3853248	--
<i>FADS2</i>	fatty acid desaturase 2	rs1535	breastfeeding & IQ (53)

<i>FADS2</i>	fatty acid desaturase 2	rs174575	breastfeeding & IQ (53)
<i>FMR1</i>	fragile X mental retardation 1	rs1805420	--
<i>FSH</i>	follicle stimulating hormone	rs6169	--
<i>FSHR</i>	follicle stimulating hormone receptor	rs6166	sterility (54), osteoporosis (55)
<i>FST</i>	follistatin	rs12152850	--
<i>FST</i>	follistatin	rs3797297	--
<i>FTO</i>	fat mass and obesity associated	rs1421085	obesity (56-58), mental disorders (59)
<i>GABBR2</i>	γ -aminobutyric acid B receptor 2	rs1435252	nicotine addiction (60)
<i>GABBR2</i>	γ -aminobutyric acid B receptor 2	rs2779562	nicotine addiction (60)
<i>GNRH1</i>	gonadotropin-releasing hormone	novel SNP	Alzheimer's disease (61)
<i>HERC</i>	hect domain and RLD 2	rs12913832	eye color (62, 63)
<i>HFE</i>	hemochromatosis	rs1799945	hemochromatosis(64)
<i>HSD17B1</i>	estradiol 17 β -dehydrogenase 1	rs12602084	steroid metabolism (65)
<i>HSD17B1</i>	estradiol 17 β -dehydrogenase 1	rs592389	vasomotor symptoms (66), cognition (67)
<i>5-HTR1A</i>	5-hydroxytryptamine (serotonin) receptor 1A	rs878567	mood disorders (68)
<i>5-HTR2A</i>	5-hydroxytryptamine (serotonin) receptor 2A	rs6312	--
<i>5-HTR2A</i>	5-hydroxytryptamine (serotonin) receptor 2A	rs6314	antidepressant response (69), bipolar disorder (70)
<i>5-HTR2A</i>	5-hydroxytryptamine (serotonin) receptor 2A	rs7997012	antidepressant response (71)
<i>5-HTR2C</i>	5-hydroxytryptamine (serotonin) receptor 2C	rs6318	bipolar disorder (72), depression (73)
<i>5-HTT</i>	5-hydroxytryptamine transporter	rs25533	antidepressant response (74)
<i>5-HTT</i>	5-hydroxytryptamine transporter	rs8076005	depressive symptoms (75)
<i>IGF1</i>	insulin-like growth factor 1	rs12313279	--
<i>IL1A</i>	interleukin 1, alpha	rs17561	chronic rhinosinusitis (76), BMI (77)
<i>IL6</i>	interleukin 6	rs1800795	arthritis (78), breast cancer (79), diabetes (80), depression (81)
<i>INHA</i>	inhibin alpha	rs2059693	testicular cancer (82)
<i>INHA</i>	inhibin alpha	rs35118453	--
<i>INHBA</i>	inhibin beta A	rs2237436	--
<i>INHBB</i>	inhibin beta B	rs11902591	--
<i>KIBRA</i>	kidney and brain protein (WWC1)	rs17070145	Alzheimer's disease (83), episodic memory (84)
<i>LEPR</i>	leptin receptor	rs1137100	diabetes II (85), atherosclerosis (86)
<i>LHR</i>	luteinizing hormone receptor	rs4073366	Alzheimer's disease (87)
<i>MAOA</i>	monoamine oxidase A	rs3788862	pain (88)
<i>OXTR</i>	oxytocin receptor	rs2254298	autism (89, 90), social loneliness (91), depressive symptoms & anxiety (92)
<i>PCK1</i>	phosphoenolpyruvate carboxykinase 1	rs707555	diabetes II (93)
<i>PGR</i>	progesterone receptor	rs1042838	ovarian cancer (94), migraine (95), menstruation (96), pregnancy loss (97)
<i>SNAP25</i>	synaptosomal-associated protein 25	rs363050	intelligence (98, 99)
<i>SSADH</i>	succinic semialdehyde dehydrogenase	rs2760118	--
<i>StAR</i>	steroidogenic acute regulatory protein	rs3990403	--
<i>TFAM</i>	transcription factor A, mitochondrial	rs1937	Alzheimer's disease (100)
<i>TFAM</i>	transcription factor A, mitochondrial	rs2306604	Parkinson's disease (101)
<i>TPH1</i>	first tryptophan hydroxylase isoform	rs1799913	heroin addiction (102)

REFERENCES

1. Zappia M, Manna I, Serra P, Cittadella R, Andreoli V, La Russa A, et al. Increased risk for Alzheimer disease with the interaction of MPO and A2M polymorphisms. *Arch Neurol*. 2004 Mar;61(3):341-4.
2. Moses EK, Fitzpatrick E, Freed KA, Dyer TD, Forrest S, Elliott K, et al. Objective prioritization of positional candidate genes at a quantitative trait locus for pre-eclampsia on 2q22. *Molecular Human Reproduction*. 2006 August 2006;12(8):505-12.
3. Hara K, Boutin P, Mori Y, Tobe K, Dina C, Yasuda K, et al. Genetic Variation in the Gene Encoding Adiponectin Is Associated With an Increased Risk of Type 2 Diabetes in the Japanese Population. *Diabetes*. 2002 February 1, 2002;51(2):536-40.
4. Gu HF, Abulaiti A, Ostenson CG, Humphreys K, Wahlestedt C, Brookes AJ, et al. Single nucleotide polymorphisms in the proximal promoter region of the adiponectin (APM1) gene are associated with type 2 diabetes in Swedish caucasians. *Diabetes*. 2004 Feb;53 Suppl 1:S31-5.
5. Bouatia-Naji N, Meyre D, Lobbens S, Seron K, Fumeron F, Balkau B, et al. ACDC/adiponectin polymorphisms are associated with severe childhood and adult obesity. *Diabetes*. 2006 Feb;55(2):545-50.
6. Yang W-S, Yang Y-C, Chen C-L, Wu I-L, Lu J-Y, Lu F-H, et al. Adiponectin SNP276 is associated with obesity, the metabolic syndrome, and diabetes in the elderly. *The American Journal of Clinical Nutrition*. 2007 August 1, 2007;86(2):509-13.
7. Kaklamani VG, Sadim M, Hsi A, Offit K, Oddoux C, Ostrer H, et al. Variants of the adiponectin and adiponectin receptor 1 genes and breast cancer risk. *Cancer Res*. 2008 May 1;68(9):3178-84.
8. Beckers S, Peeters AV, De Freitas F, Mertens IL, Verhulst SL, Haentjens D, et al. Association Study and Mutation Analysis of Adiponectin Shows Association of Variants in APM1 with Complex Obesity in Women. *Annals of Human Genetics*. 2009;73(5):492-501.
9. Young WL, Kwok P-Y, Pawlikowska L, Lawton MT, Kim H, Hysi PG, et al. Arteriovenous Malformation. *Journal of Neurosurgery*. 2007;106(4):731-2.
10. Petersen KF, Dufour S, Hariri A, Nelson-Williams C, Foo JN, Zhang X-M, et al. Apolipoprotein C3 Gene Variants in Nonalcoholic Fatty Liver Disease. *New England Journal of Medicine*. 2010;362(12):1082-9.
11. Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261(5123):921-3.
12. Saunders AM, Strittmatter WJ, Schmechel D, George-Hyslop PH, Pericak-Vance MA, Joo SH, et al. Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology*. 1993;43(8):1467-72.
13. Ellis JA, Stebbing M, Harrap SB. Polymorphism of the Androgen Receptor Gene is Associated with Male Pattern Baldness. 2001;116(3):452-5.
14. Kang H, Lee SK, Cho S-W, Lee S-H, Kwack K. Branched chain [alpha]-keto acid dehydrogenase, E1-[beta] subunit gene is associated with premature ovarian failure. *Fertility and Sterility*. 2008;89(3):728-31.

15. Hwang J-P, Tsai S-J, Hong C-J, Yang C-H, Lirng J-F, Yang Y-M. The Val66Met polymorphism of the brain-derived neurotrophic-factor gene is associated with geriatric depression. *Neurobiology of Aging*. 2006;27(12):1834-7.
16. Czira ME, Wersching H, Baune BT, Berger K. Brain-derived neurotrophic factor gene polymorphisms, neurotransmitter levels, and depressive symptoms in an elderly population. *Age (Dordr)*. 2011 Sep 7.
17. Suchanek R, Owczarek A, Kowalczyk M, Kucia K, Kowalski J. Association between C-281A and val66met functional polymorphisms of BDNF gene and risk of recurrent major depressive disorder in Polish population. *J Mol Neurosci*. 2011 Mar;43(3):524-30.
18. Su N, Zhang L, Fei F, Hu H, Wang K, Hui H, et al. The brain-derived neurotrophic factor is associated with alcohol dependence-related depression and antidepressant response. *Brain Research*. 2011;1415(0):119-26.
19. Sears C, Markie D, Olds R, Fitches A. Evidence of associations between bipolar disorder and the brain-derived neurotrophic factor (BDNF) gene. *Bipolar Disorders*. 2011;13(7-8):630-7.
20. Zakharyan R, Boyajyan A, Arakelyan A, Gevorgyan A, Mrazek F, Petrek M. Functional variants of the genes involved in neurodevelopment and susceptibility to schizophrenia in an Armenian population. *Human Immunology*. 2011;72(9):746-8.
21. McAllister TW, Tyler AL, Flashman LA, Rhodes CH, McDonald BC, Saykin AJ, et al. Polymorphisms in the brain-derived neurotrophic factor gene influence memory and processing speed one month after brain injury. *J Neurotrauma*. 2012 Apr 10;29(6):1111-8.
22. Hong K-W, Lim J-E, Go MJ, Shin Cho Y, Ahn Y, Han B-G, et al. Recapitulation of the Association of the Val66Met Polymorphism of BDNF Gene With BMI in Koreans. *Obesity*. 2011.
23. Gratacos M, Soria V, Urretavizcaya M, Gonzalez JR, Crespo JM, Bayes M, et al. A brain-derived neurotrophic factor (BDNF) haplotype is associated with antidepressant treatment outcome in mood disorders. *Pharmacogenomics J*. 2007;8(2):101-12.
24. Johnson N, Fletcher O, Palles C, Rudd M, Webb E, Sellick G, et al. Counting potentially functional variants in BRCA1, BRCA2 and ATM predicts breast cancer susceptibility. *Human Molecular Genetics*. 2007 May 1, 2007;16(9):1051-7.
25. Wang JC, Hinrichs AL, Stock H, Budde J, Allen R, Bertelsen S, et al. Evidence of common and specific genetic effects: association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Human Molecular Genetics*. 2004 September 1, 2004;13(17):1903-11.
26. Jones KA, Porjesz B, Almasy L, Bierut L, Goate A, Wang JC, et al. Linkage and linkage disequilibrium of evoked EEG oscillations with CHRM2 receptor gene polymorphisms: implications for human brain dynamics and cognition. *International Journal of Psychophysiology*. 2004;53(2):75-90.
27. Eisenberg J, Mei-Tal G, Steinberg A, Tartakovsky E, Zohar A, Gritsenko I, et al. Haplotype relative risk study of catechol-O-methyltransferase (COMT) and attention deficit hyperactivity disorder (ADHD): Association of the high-enzyme activity val allele with adhd impulsive-hyperactive phenotype. *American Journal of Medical Genetics*. 1999;88(5):497-502.
28. Wang T, Franke P, Neidt H, Cichon S, Knapp M, Lichtermann D, et al. Association study of the low-activity allele of catechol-O-methyltransferase and alcoholism using a family-based approach. *Mol Psychiatry*. 2001 Jan;6(1):109-11.
29. Horowitz R, Kotler M, Shufman E, Aharoni S, Kremer I, Cohen H, et al. Confirmation of an excess of the high enzyme activity COMT val allele in heroin addicts in a family-based haplotype relative risk study. *American Journal of Medical Genetics*. 2000;96(5):599-603.

30. Vandenberg DJ, Rodriguez LA, Miller IT, Uhl GR, Lachman HM. High-activity catechol-O-methyltransferase allele is more prevalent in polysubstance abusers. *American Journal of Medical Genetics*. 1997;74(4):439-42.
31. Li T, Chen C-k, Hu X, Ball D, Lin S-K, Chen W, et al. Association analysis of the DRD4 and COMT genes in methamphetamine abuse. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2004;129B(1):120-4.
32. Åberg E, Fandiño-Losada A, Sjöholm LK, Forsell Y, Lavebratt C. The functional Val158Met polymorphism in catechol-O-methyltransferase (COMT) is associated with depression and motivation in men from a Swedish population-based study. *Journal of Affective Disorders*. 2011;129(1–3):158-66.
33. Baune BT, Hohoff C, Berger K, Neumann A, Mortensen S, Roehrs T, et al. Association of the COMT val158met Variant with Antidepressant Treatment Response in Major Depression. *Neuropsychopharmacology*. 2007;33(4):924-32.
34. Lee S-Y, Chen S-L, Chen S-H, Huang S-Y, Tzeng N-S, Chang Y-H, et al. The COMT and DRD3 genes interacted in bipolar I but not bipolar II disorder. *World Journal of Biological Psychiatry*. 2011;12(5):385-91.
35. Dickinson D, Elvevåg B. Genes, cognition and brain through a COMT lens. *Neuroscience*. 2009;164(1):72-87.
36. Albayrak Ö, Tirniceriu A, Riemenschneider M, Kurz A, Scherag A, Egensperger R. The Cathepsin D (224C/T) Polymorphism Confers an Increased Risk to Develop Alzheimer's Disease in Men. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2010 March 1, 2010;65A(3):219-24.
37. Yaspan BL, Breyer JP, Cai Q, Dai Q, Elmore JB, Amundson I, et al. Haplotype Analysis of CYP11A1 Identifies Promoter Variants Associated with Breast Cancer Risk. *Cancer Research*. 2007 June 15, 2007;67(12):5673-82.
38. Tu Y, Cui G, Xu Y, Bao X, Wang X, Wang D. Genetic polymorphism of CYP11B2 gene and stroke in the Han Chinese population and a meta-analysis. *Pharmacogenet Genomics*. 2011;21(3):115-20.
39. White PC, Hautanen A, Kupari M. Aldosterone synthase (CYP11B2) polymorphisms and cardiovascular function. *J Steroid Biochem Mol Biol*. 1999;69(1-6):409-12.
40. Brookes K, Xu X, Chen W, Zhou K, Neale B, Lowe N, et al. The analysis of 51 genes in DSM-IV combined type attention deficit hyperactivity disorder: association signals in DRD4, DAT1 and 16 other genes. *Mol Psychiatry*. 2006;11(10):934-53.
41. Le Strat Y, Ramoz N, Pickering P, Burger V, Boni C, Aubin H-J, et al. The 3' Part of the Dopamine Transporter Gene DAT1/SLC6A3 Is Associated With Withdrawal Seizures in Patients With Alcohol Dependence. *Alcoholism: Clinical and Experimental Research*. 2008;32(1):27-35.
42. Callicott JH, Straub RE, Pezawas L, Egan MF, Mattay VS, Hariri AR, et al. Variation in DISC1 affects hippocampal structure and function and increases risk for schizophrenia. *Proc Natl Acad Sci U S A*. 2005;102(24):8627-32.
43. Thomson PA, Harris SE, Starr JM, Whalley LJ, Porteous DJ, Deary IJ. Association between genotype at an exonic SNP in DISC1 and normal cognitive aging. *Neuroscience Letters*. 2005;389(1):41-5.
44. Blum K, Braverman ER, Wood RC, Gill J, Li C, Chen TJ, et al. Increased prevalence of the Taq I A1 allele of the dopamine receptor gene (DRD2) in obesity with comorbid substance use disorder: a preliminary report. *Pharmacogenetics*. 1996 Aug;6(4):297-305.
45. Dubertret C, Bardel C, Ramoz N, Martin P-M, Deybach J-C, Adès J, et al. A genetic schizophrenia-susceptibility region located between the ANKK1 and DRD2 genes. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2010;34(3):492-9.
46. Monakhov M, Golimbet V, Abramova L, Kaleda V, Karpov V. Association study of three polymorphisms in the dopamine D2 receptor gene and schizophrenia in the Russian population. *Schizophrenia Research*. 2008;100(1–3):302-7.

47. Voisey J, Swagell CD, Hughes IP, Morris CP, van Daal A, Noble EP, et al. The DRD2 gene 957C>T polymorphism is associated with Posttraumatic Stress Disorder in war veterans. *Depression and Anxiety*. 2009;26(1):28-33.
48. Yang J-W, Jang W-S, Hong SD, Ji YI, Kim DH, Park J, et al. A case-control association study of the polymorphism at the promoter region of the DRD4 gene in Korean boys with attention deficit-hyperactivity disorder: Evidence of association with the - 521 C/T SNP. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2008;32(1):243-8.
49. Lai JH, Zhu YS, Huo ZH, Sun RF, Yu B, Wang YP, et al. Association study of polymorphisms in the promoter region of DRD4 with schizophrenia, depression, and heroin addiction. *Brain Research*. 2010;1359:227-32.
50. Funke B, Finn CT, Plocik AM, Lake S, DeRosse P, Kane JM, et al. Association of the DTNBP1 Locus with Schizophrenia in a U.S. Population. *The American Journal of Human Genetics*. 2004;75(5):891-8.
51. Burdick KE, Lencz T, Funke B, Finn CT, Szeszko PR, Kane JM, et al. Genetic variation in DTNBP1 influences general cognitive ability. *Human Molecular Genetics*. 2006 15 May 2006;15(10):1563-8.
52. Riley B, Kuo P-H, Maher BS, Fanous AH, Sun J, Wormley B, et al. The dystrobrevin binding protein 1 (DTNBP1) gene is associated with schizophrenia in the Irish Case Control Study of Schizophrenia (ICCS) sample. *Schizophrenia Research*. 2009;115(2-3):245-53.
53. Caspi A, Williams B, Kim-Cohen J, Craig IW, Milne BJ, Poulton R, et al. Moderation of breastfeeding effects on the IQ by genetic variation in fatty acid metabolism. *Proc Natl Acad Sci U S A*. 2007 Nov 20;104(47):18860-5.
54. Binder H, Dittrich R, Hager I, Muller A, Oeser S, Beckmann MW, et al. Association of FSH receptor and CYP19A1 gene variations with sterility and ovarian hyperstimulation syndrome. *Reproduction*. 2008;135(1):107-16.
55. Rendina D, Gianfrancesco F, De Filippo G, Merlotti D, Esposito T, Mingione A, et al. FSHR gene polymorphisms influence bone mineral density and bone turnover in postmenopausal women. *Eur J Endocrinol*. 2010;163(1):165-72.
56. Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet*. 2007;39(6):724-6.
57. Tung Y-CL, Yeo GSH. From GWAS to biology: lessons from FTO. *Annals of the New York Academy of Sciences*. 2011;1220(1):162-71.
58. Chabris C, Hebert B, Benjamin D, Beauchamp J, Cesarini D, van der Loos M, et al. Most reported genetic associations with general intelligence are probably false positives. *Psychological Science*. 2011;In press.
59. Kivimaki M, Jokela M, Hamer M, Geddes J, Ebmeier K, Kumari M, et al. Examining Overweight and Obesity as Risk Factors for Common Mental Disorders Using Fat Mass and Obesity-Associated (FTO) Genotype-Instrumented Analysis. *Am J Epidemiol*. 2011 Feb 15;173(4):421-9.
60. Beuten J, Ma JZ, Payne TJ, Dupont RT, Crews KM, Somes G, et al. Single- and Multilocus Allelic Variants within the GABAB Receptor Subunit 2 (GABAB2) Gene Are Significantly Associated with Nicotine Dependence. *The American Journal of Human Genetics*. 2005;76(5):859-64.
61. Wilson AC. Investigations into gonadotropin-releasing hormone receptor signaling in extrahypothalamic neurons of the brain: Implications for normal cognition and the pathogenesis of Alzheimer's disease: Thesis. University of Wisconsin, Madison, USA; 2009.
62. Eiberg H, Troelsen J, Nielsen M, Mikkelsen A, Mengel-From J, Kjaer K, et al. Blue eye color in humans may be caused by a perfectly associated founder mutation in a regulatory element located within the HERC2 gene inhibiting OCA2 expression. *Human Genetics*. 2008;123(2):177-87.

63. Sturm RA, Duffy DL, Zhao ZZ, Leite FPN, Stark MS, Hayward Nicholas K, et al. A Single SNP in an Evolutionary Conserved Region within Intron 86 of the HERC2 Gene Determines Human Blue-Brown Eye Color. *The American Journal of Human Genetics*. 2008;82(2):424-31.
64. Beutler E. The significance of the 187G (H63D) mutation in hemochromatosis. *Am J Hum Genet*. 1997 Sep;61(3):762-4.
65. Sun T, Oh WK, Jacobus S, Regan M, Pomerantz M, Freedman ML, et al. The Impact of Common Genetic Variations in Genes of the Sex Hormone Metabolic Pathways on Steroid Hormone Levels and Prostate Cancer Aggressiveness. *Cancer Prevention Research*. 2011 December 1, 2011;4(12):2044-50.
66. Crandall CJ, Crawford SL, Gold EB. Vasomotor Symptom Prevalence Is Associated with Polymorphisms in Sex Steroid-Metabolizing Enzymes and Receptors. *The American Journal of Medicine*. 2006;119(9, Supplement 1):S52-S60.
67. Kravitz HM, Meyer PM, Seeman TE, Greendale GA, Sowers MR. Cognitive Functioning and Sex Steroid Hormone Gene Polymorphisms in Women at Midlife. *The American Journal of Medicine*. 2006;119(9, Supplement 1):S94-S102.
68. Brezo J, Bureau A, Merette C, Jomphe V, Barker ED, Vitaro F, et al. Differences and similarities in the serotonergic diathesis for suicide attempts and mood disorders: a 22-year longitudinal gene-environment study. *Mol Psychiatry*. 2010;15(8):831-43.
69. Wilkie MJV, Smith G, Day RK, Matthews K, Smith D, Blackwood D, et al. Polymorphisms in the SLC6A4 and HTR2A genes influence treatment outcome following antidepressant therapy. *Pharmacogenomics J*. 2008;9(1):61-70.
70. McAuley EZ, Fullerton JM, Blair IP, Donald JA, Mitchell PB, Schofield PR. Association between the serotonin 2A receptor gene and bipolar affective disorder in an Australian cohort. *Psychiatr Genet*. 2009 Oct;19(5):244-52.
71. McMahon FJ, Buervenich S, Charney D, Lipsky R, Rush AJ, Wilson AF, et al. Variation in the Gene Encoding the Serotonin 2A Receptor Is Associated with Outcome of Antidepressant Treatment. *The American Journal of Human Genetics*. 2006;78(5):804-14.
72. Mazza M, Mandelli L, Martinotti G, Di Nicola M, Tavian D, Negri G, et al. Further evidence supporting the association between 5HT2C gene and bipolar disorder. *Psychiatry Res*. 2010 Dec 30;180(2-3):151-2.
73. Lerer B, Macciardi F, Segman RH, Adolfsson R, Blackwood D, Blairy S, et al. Variability of 5-HT2C receptor cys23ser polymorphism among European populations and vulnerability to affective disorder. *Mol Psychiatry*. 2001 Sep;6(5):579-85.
74. Lin E, Chen PS, Chang HH, Gean P-W, Tsai HC, Yang YK, et al. Interaction of serotonin-related genes affects short-term antidepressant response in major depressive disorder. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2009;33(7):1167-72.
75. Su S, Zhao J, Bremner JD, Miller AH, Tang W, Bouzyk M, et al. Serotonin Transporter Gene, Depressive Symptoms, and Interleukin-6 / CLINICAL PERSPECTIVE. *Circulation: Cardiovascular Genetics*. 2009 December 1, 2009;2(6):614-20.
76. Mfuna Endam L, Cormier C, Bosse Y, Filali-Mouhim A, Desrosiers M. Association of IL1A, IL1B, and TNF Gene Polymorphisms With Chronic Rhinosinusitis With and Without Nasal Polyposis: A Replication Study. *Arch Otolaryngol Head Neck Surg*. 2010 February 1, 2010;136(2):187-92.
77. Um J-Y, Rim H-K, Kim S-J, Kim H-L, Hong S-H. Functional Polymorphism of IL-1 Alpha and Its Potential Role in Obesity in Humans and Mice. *PLoS ONE*. 2011;6(12):e29524.
78. Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, et al. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *The Journal of Clinical Investigation*. 1998;102(7):1369-76.

79. Slattery ML, Curtin K, Sweeney C, Wolff RK, Baumgartner RN, Baumgartner KB, et al. Modifying Effects of IL-6 Polymorphisms on Body Size-Associated Breast Cancer Risk. *Obesity*. 2008;16(2):339-47.
80. Huth C, Heid IM, Vollmert C, Gieger C, Grallert H, Wolford JK, et al. IL6 Gene Promoter Polymorphisms and Type 2 Diabetes. *Diabetes*. 2006 October 2006;55(10):2915-21.
81. Bull SJ, Huezo-Diaz P, Binder EB, Cubells JF, Ranjith G, Maddock C, et al. Functional polymorphisms in the interleukin-6 and serotonin transporter genes, and depression and fatigue induced by interferon-alpha and ribavirin treatment. *Mol Psychiatry*. 2009 Dec;14(12):1095-104.
82. Purdue MP, Graubard BI, Chanock SJ, Rubertone MV, Erickson RL, McGlynn KA. Genetic Variation in the Inhibin Pathway and Risk of Testicular Germ Cell Tumors. *Cancer Research*. 2008 April 15, 2008;68(8):3043-8.
83. Rodríguez-Rodríguez E, Infante J, Llorca J, Mateo I, Sánchez-Quintana C, García-Gorostiaga I, et al. Age-dependent association of KIBRA genetic variation and Alzheimer's disease risk. *Neurobiology of Aging*. 2009;30(2):322-4.
84. Almeida OP, Schwab SG, Lautenschlager NT, Morar B, Greenop KR, Flicker L, et al. KIBRA genetic polymorphism influences episodic memory in later life, but does not increase the risk of mild cognitive impairment. *Journal of Cellular and Molecular Medicine*. 2008;12(5a):1672-6.
85. Salopuro T, Pulkkinen L, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, et al. Genetic variation in leptin receptor gene is associated with type 2 diabetes and body weight: The Finnish Diabetes Prevention Study. *Int J Obes Relat Metab Disord*. 2005;29(10):1245-51.
86. Saukko M, Kesaniemi YA, Ukkola O. Leptin receptor Lys109Arg and Gln223Arg polymorphisms are associated with early atherosclerosis. *Metab Syndr Relat Disord*. 2010;8(5):425-30.
87. Haasl R, Ahmadi MR, Meethal S, Gleason C, Johnson S, Asthana S, et al. A luteinizing hormone receptor intronic variant is significantly associated with decreased risk of Alzheimer's disease in males carrying an apolipoprotein E epsilon4 allele. *BMC Medical Genetics*. 2008;9(1):37.
88. Kim H, Lee H, Rowan J, Brahim J, Dionne R. Genetic polymorphisms in monoamine neurotransmitter systems show only weak association with acute post-surgical pain in humans. *Molecular Pain*. 2006;2(1):24.
89. Lerer E, Levi S, Salomon S, Darvasi A, Yirmiya N, Ebstein RP. Association between the oxytocin receptor (OXTR) gene and autism: relationship to Vineland Adaptive Behavior Scales and cognition. *Mol Psychiatry*. 2007;13(10):980-8.
90. Liu X, Kawamura Y, Shimada T, Otowa T, Koishi S, Sugiyama T, et al. Association of the oxytocin receptor (OXTR) gene polymorphisms with autism spectrum disorder (ASD) in the Japanese population. *J Hum Genet*. 2010;55(3):137-41.
91. Lucht MJ, Barnow S, Sonnenfeld C, Rosenberger A, Grabe HJ, Schroeder W, et al. Associations between the oxytocin receptor gene (OXTR) and affect, loneliness and intelligence in normal subjects. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2009;33(5):860-6.
92. Thompson RJ, Parker KJ, Hallmayer JF, Waugh CE, Gotlib IH. Oxytocin receptor gene polymorphism (rs2254298) interacts with familial risk for psychopathology to predict symptoms of depression and anxiety in adolescent girls. *Psychoneuroendocrinology*. 2011;36(1):144-7.
93. Dong Y, Zhang H, Wang X, Feng Q, Chen X, Su Q. A Leu184Val polymorphism in PCK1 gene is associated with type 2 diabetes in Eastern Chinese population with BMI < 23 kg/m². *Diabetes Research and Clinical Practice*. 2009;83(2):227-32.
94. Pearce CL, Hirschhorn JN, Wu AH, Burt NP, Stram DO, Young S, et al. Clarifying the PROGINS Allele Association in Ovarian and Breast Cancer Risk: A Haplotype-Based Analysis. *Journal of the National Cancer Institute*. 2005 January 5, 2005;97(1):51-9.

95. Joshi G, Pradhan S, Mittal B. Role of the oestrogen receptor (ESR1 PvuII and ESR1 325 C→G) and progesterone receptor (PROGINS) polymorphisms in genetic susceptibility to migraine in a North Indian population. *Cephalalgia*. 2010 March 1, 2010;30(3):311-20.
96. Taylor KC, Small CM, Epstein MP, Sherman SL, Tang W, Wilson MM, et al. Associations of Progesterone Receptor Polymorphisms with Age at Menarche and Menstrual Cycle Length. *Hormone Research in Paediatrics*. 2010;74(6):421-7.
97. Su MT, Lee IW, Chen YC, Kuo PL. Association of progesterone receptor polymorphism with idiopathic recurrent pregnancy loss in Taiwanese Han population. *J Assist Reprod Genet*. 2011 Mar;28(3):239-43.
98. Gosso MF, de Geus EJC, van Belzen MJ, Polderman TJC, Heutink P, Boomsma DI, et al. The SNAP-25 gene is associated with cognitive ability: evidence from a family-based study in two independent Dutch cohorts. *Mol Psychiatry*. 2006;11(9):878-86.
99. Gosso MF, De Geus EJC, Polderman TJC, Boomsma DI, Heutink P, Posthuma D. Common variants underlying cognitive ability: further evidence for association between the SNAP-25 gene and cognition using a family-based study in two independent Dutch cohorts. *Genes, Brain and Behavior*. 2008;7(3):355-64.
100. Zhang Q, Yu J-T, Wang P, Chen W, Wu Z-C, Jiang H, et al. Mitochondrial transcription factor A (TFAM) polymorphisms and risk of late-onset Alzheimer's disease in Han Chinese. *Brain Research*. 2011;1368(0):355-60.
101. Gaweda-Walerych K, Safranow K, Maruszak A, Bialecka M, Klodowska-Duda G, Czyzewski K, et al. Mitochondrial transcription factor A variants and the risk of Parkinson's disease. *Neuroscience Letters*. 2010;469(1):24-9.
102. Nielsen DA, Barral S, Proudnikov D, Kellogg S, Ho A, Ott J, et al. TPH2 and TPH1: association of variants and interactions with heroin addiction. *Behav Genet*. 2008 Mar;38(2):133-50.
103. Cichon S, Winge I, Mattheisen M, Georgi A, Karpushova A, Freudenberg J, et al. Brain-specific tryptophan hydroxylase 2 (TPH2): a functional Pro206Ser substitution and variation in the 5'-region are associated with bipolar affective disorder. *Human Molecular Genetics*. 2008 January 1, 2008;17(1):87-97.
104. Goenjian AK, Bailey JN, Walling DP, Steinberg AM, Schmidt D, Dandekar U, et al. Association of TPH1, TPH2, and 5HTTLPR with PTSD and depressive symptoms. *Journal of Affective Disorders*. 2012(in Press).

Supplementary Table 2. Depression-Associated SNP Identified in the WLS

Gene	Encodes	SNP	Alleles	Chr#/Location	Residue	Associated disease/behavior
<i>ACVR2B</i>	activin receptor IIB	rs3749386	T/C	3/intron 1	--	left-right axis malformations*(1)
<i>APOC3</i>	apolipoprotein C-III	rs2854116	T/C	11/promoter (-455)	--	nonalcoholic fatty liver disease(2)
<i>DRD2/ANKK1</i>	dopamine receptor D2/ankyrin repeat and kinase domain containing 1	rs1800497	C/T	11/exon (ANKK1)	Glu713Lys	obesity, drug addiction (3)
<i>DRD2</i>	dopamine receptor D2	rs2242592	T/C	11/3'	--	schizophrenia (4)
<i>FTO</i>	fat mass and obesity associated	rs1421085	T/C	16/intron 1	--	obesity (5-7), mental disorders (8)
<i>GNRH1</i>	gonadotropin-releasing hormone	novel SNP	T/C	8/promoter	--	Alzheimer's disease (9)
<i>IL6</i>	interleukin 6	rs1800795	C/G	7/promoter (-174)	--	arthritis (10), breast cancer (11), diabetes (12), depression (13)

Gene association only

References

1. Kosaki R, Gebbia M, Kosaki K, Lewin M, Bowers P, Towbin JA, et al. Left–right axis malformations associated with mutations in ACVR2B, the gene for human activin receptor type IIB. *American Journal of Medical Genetics*. 1999;82(1):70-6.
2. Petersen KF, Dufour S, Hariri A, Nelson-Williams C, Foo JN, Zhang X-M, et al. Apolipoprotein C3 Gene Variants in Nonalcoholic Fatty Liver Disease. *New England Journal of Medicine*. 2010;362(12):1082-9.
3. Blum K, Braverman ER, Wood RC, Gill J, Li C, Chen TJ, et al. Increased prevalence of the Taq I A1 allele of the dopamine receptor gene (DRD2) in obesity with comorbid substance use disorder: a preliminary report. *Pharmacogenetics*. 1996 Aug;6(4):297-305.
4. Dubertret C, Bardel C, Ramoz N, Martin P-M, Deybach J-C, Adès J, et al. A genetic schizophrenia-susceptibility region located between the ANKK1 and DRD2 genes. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2010;34(3):492-9.
5. Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet*. 2007;39(6):724-6.
6. Tung Y-CL, Yeo GSH. From GWAS to biology: lessons from FTO. *Annals of the New York Academy of Sciences*. 2011;1220(1):162-71.
7. Chabris C, Hebert B, Benjamin D, Beauchamp J, Cesarini D, van der Loos M, et al. Most reported genetic associations with general intelligence are probably false positives. *Psychological Science*. 2011;In press.
8. Kivimaki M, Jokela M, Hamer M, Geddes J, Ebmeier K, Kumari M, et al. Examining Overweight and Obesity as Risk Factors for Common Mental Disorders Using Fat Mass and Obesity-Associated (FTO) Genotype-Instrumented Analysis. *Am J Epidemiol*. 2011 Feb 15;173(4):421-9.
9. Wilson AC. Investigations into gonadotropin-releasing hormone receptor signaling in extrahypothalamic neurons of the brain: Implications for normal cognition and the pathogenesis of Alzheimer's disease: Thesis. University of Wisconsin, Madison, USA; 2009.
10. Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, et al. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *The Journal of Clinical Investigation*. 1998;102(7):1369-76.
11. Slattery ML, Curtin K, Sweeney C, Wolff RK, Baumgartner RN, Baumgartner KB, et al. Modifying Effects of IL-6 Polymorphisms on Body Size-Associated Breast Cancer Risk. *Obesity*. 2008;16(2):339-47.
12. Huth C, Heid IM, Vollmert C, Gieger C, Grallert H, Wolford JK, et al. IL6 Gene Promoter Polymorphisms and Type 2 Diabetes. *Diabetes*. 2006 October 2006;55(10):2915-21.
13. Bull SJ, Huezio-Diaz P, Binder EB, Cubells JF, Ranjith G, Maddock C, et al. Functional polymorphisms in the interleukin-6 and serotonin transporter genes, and depression and fatigue induced by interferon-alpha and ribavirin treatment. *Mol Psychiatry*. 2009 Dec;14(12):1095-104.

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported
Objectives	3	State specific objectives, including any prespecified hypotheses
Methods		
Study design	4	Present key elements of study design early in the paper
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up (b) For matched studies, give matching criteria and number of exposed and unexposed
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	Describe any efforts to address potential sources of bias
Study size	10	Explain how the study size was arrived at
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, explain how loss to follow-up was addressed (e) Describe any sensitivity analyses
Results		
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Summarise follow-up time (eg, average and total amount)
Outcome data	15*	Report numbers of outcome events or summary measures over time
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
Discussion		
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.