

A population-based study of nearly 15 000 observations among Swedish women and men during 1973–2003

Christina Halford,¹ Catharina Welin,² Johan Bogefeldt,¹ Thorne Wallman,^{1,3} Annika Rosengren,² Annika Bardel,^{1,3} Saga Johansson,^{2,4} Henry Eriksson,² Kurt Svärdsudd¹

To cite: Halford C, Welin C, Bogefeldt J, *et al.* A population-based study of nearly 15 000 observations among Swedish women and men during 1973–2003. *BMJ Open* 2012;**2**:e001353. doi:10.1136/bmjopen-2012-001353

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-001353>).

Received 15 May 2012
Accepted 26 September 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article

Correspondence to: Dr Christina Halford; Christina.halford@pubcare.uu.se

ABSTRACT

Objectives: Global self-rated health (SRH) has become extensively used as an outcome measure in population health surveillance. The aim of this study was to analyse the effects of age and secular trend (year of investigation) on SRH.

Design: Prospective cohort study, using population-based data from eight ongoing cohort studies, with sampling performed between 1973 and 2003.

Setting: Sweden.

Participants: 11 880 women and men, aged 25–99 years, providing 14 470 observations.

Primary outcome measure: Global SRH.

Results: In multiple ordinal logistic regression analyses, adjusted for the effects of covariates, there were independent effects of age ($p < 0.0001$) and of year of investigation ($p < 0.0001$) on SRH. In women the association was linear, showing lower levels of SRH with increased age, and more recent year of investigation. In men the association was curvilinear, and thus more complex. The final model explained 76.2% of the SRH variance in women and 74.5% of the variance in men.

Conclusions: SRH was strongly and inversely associated with age in both sexes, after adjustment for other outcome-affecting variables. There was a strongly significant effect of year of investigation indicating a change in SRH, in women towards lower levels over calendar time, in men with fluctuations across time.

INTRODUCTION

Global self-rated health (SRH), a one-item simple question in which respondents are asked to rate their overall health, has predictive validity in relation to functional ability,¹ healthcare utilisation,² morbidity³ and mortality.^{4 5} Owing to the simplicity of use, and predictive validity held in relation to future health SRH has increasingly become used as an outcome measure in medical health research, and in population health surveillance.

ARTICLE SUMMARY

Article focus

- To investigate the effects of age and secular trends on self-rated health (SRH) adjusted for the effects of a wide range of potential confounders in a large population-based study of adult women and men.

Key messages

- SRH is affected by age but differently in men and women.
- There is a secular trend regarding SRH, such that SRH was poorer in more recent years of investigation, after controlling for age and a large number of potential confounders.
- The secular trends differed among women and men.

Strengths and limitations of this study

- Strengths included a large sample size and observation time which spanned three decades in men and one decade in women.
- Limitations included that all age groups were not represented on each measurement occasion, and data were generally cross-sectional.

A number of factors are known to influence SRH. The relation to age is somewhat controversial. Non-comparative SRH has been found to decrease with age,^{6–11} although better SRH among the oldest old^{12 13} and no association between age and SRH,¹⁴ has also been reported.

Age reflects effects of biological processes of ageing internal to the individual, and effects of contextually determined factors of importance to health and well-being.^{15 16} Although increased knowledge concerning effects of contextually determined factors on SRH may have important public health implications, studies investigating effects of cohort or year of investigation on health are few, and results inconclusive. Poorer SRH, as well as better SRH, in more recent cohorts, and

little change between cohorts in age-specific SRH, has been reported.^{6 11 16–19}

The aim of this study was to investigate the effects of age (individual) and secular trends, measured as year of investigation (contextual), on SRH, adjusted for the effects of a wide range of potential confounders, in a large population-based study of adult women and men.

METHODS

Study population

Analyses were based on data from eight ongoing Swedish cohort studies, with baseline investigations performed between 1973 and 2003. The study population has previously been described in detail.^{20 21} Briefly, random samples based on predefined specifications concerning age, sex and area of residence, were drawn from the national population register. The men born in 1913 subsample consisted of a random third of the male

population aged 60 in the city of Gothenburg, Sweden, in 1973, and the men born in 1923 subsample consisted of a random 10th of the male population aged 50 in Gothenburg (table 1). Survivors in the samples were invited to re-examinations in 1980, 1988 and 1993.

The men born in 1943 subsample consisted of a random third of 50-year-old men living in Gothenburg in 1993, re-examined in 2003, and the women and men born in 1953 subsamples consisted of a random third of women and men living in Gothenburg in 2003. The Eskil subsample consisted of a random sample of 625 men aged 30–54 and living in the city of Eskilstuna, Sweden in 1986. The Uppsala Public Health Cohort was based on random samples of 1000 men and women 25 years or older from each of the six municipalities of Uppsala County in 1993. The BEDA II subsample was based on a re-examination in 1997 of a random sample drawn in 1979 of 1746 women born during 1915–1941 and living in Gothenburg. The Uppsala-Örebro Women

Table 1 Characteristics of the cohorts included in the study population

Subpopulation	Investigation		Sex	Age range	Sample size	Responders	Response rate (%)	Investigation procedure
	Year	Place						
Men born in 1913	1973	Gothenburg	Men	60	1009	830	82.3	Questionnaires +medical examination
Men born in 1913	1980	Gothenburg	Men	67	923	707	76.6	Questionnaires +medical examination
Men born in 1913	1988	Gothenburg	Men	75	702	463	66.1	Questionnaires +medical examination
Men born in 1913	1993	Gothenburg	Men	80	447	272	60.9	Questionnaires +medical examination
Men born in 1923	1973	Gothenburg	Men	50	292	226	77.4	Questionnaires +medical examination
Men born in 1923	1980	Gothenburg	Men	57	278	188	67.6	Questionnaires +medical examination
Men born in 1923	1988	Gothenburg	Men	65	265	162	61.1	Questionnaires +medical examination
Men born in 1923	1993	Gothenburg	Men	70	226	143	63.3	Questionnaires +medical examination
ESKIL	1986	Eskilstuna	Men	30–54	625	459	73.8	Postal questionnaire
Public Health Cohort Uppsala	1993	Uppsala	Women	25	2999	2249	75.0	Postal questionnaire
Public Health Cohort Uppsala	1993	Uppsala	Men	25	3001	2156	71.8	Postal questionnaire
BEDA II	1997	Gothenburg	Women	56–82	994	908	91.3	Questionnaires +medical examination
Uppsala-Örebro Women Study	1995	Uppsala	Women	35–64	4200	2991	71.2	Postal questionnaire
Men born in 1943	1993	Gothenburg	Men	50	1463	798	54.5	Questionnaires +medical examination
Men born in 1943	2003	Gothenburg	Men	60	749	655	87.4	Questionnaires +medical examination
Women born in 1953	2003	Gothenburg	Women	50	994	668	67.4	Questionnaires +medical examination
Men born in 1953	2003	Gothenburg	Men	50	993	595	59.9	Questionnaires +medical examination
Total					20 160	14 470	71.8	

Study sample was based on random samples of 600 women aged 35–64 from each of the seven counties in the Uppsala-Örebro Health Care Region, Sweden.

All samples were by definition representative of their underlying general populations. No exclusions were made. The combined samples consisted of 20 160 subjects of whom 3590 were part of more than one sub-sample. Overall, 14 470 (71.8%) observations were obtained, based on 12 000 unique individuals. Of these 10 451 (6808 women and 3644 men) participated once, 964 men twice, 254 men three times and 330 men on four occasions. Additional information on the study population is given in [table 1](#).

Measurements

The data used in this report were obtained by questionnaire. For the variables used here the same questionnaires were used in all studies. Educational level was classified on a four-point scale ranging from ‘compulsory education only’ (=1), to ‘college or university education’ (=4). Occupational status was measured on a four-point nominal scale as ‘gainfully employed’ (including students), ‘unemployed’, ‘on sick leave or disability pension’ or ‘old age retirement’. Marital status was classified as married/cohabiting or not (the latter including response alternatives never married, divorced and widowed).

SRH was measured using the well-being subscale of The Gothenburg Quality of Life instrument (GQL).²² Respondents were asked to rate their health on a seven-point Likert scale with response alternatives ranging from ‘very poor’ (=1) to ‘excellent, could not be better’ (=7), with no verbal labels for the intervening steps. Symptom reporting was assessed based on the Complaint score subscale of GQL, in which subjects are asked ‘Have you been troubled by any of the following symptoms during the past 3 months?’, followed by a list of 30 general symptoms with response alternatives ‘yes’ (=1) or ‘no’ (=0) for each symptom. The Complaint score was obtained as the sum across the 30 symptoms. Complaint score is not intended to measure specific diseases, but rather the tendency to report symptoms, an aspect of quality of life.

Leisure time physical activity was reported on a four-point ordinal scale with response alternatives ‘sedentary’, ‘moderately active’, ‘active’ or ‘vigorously active’.²³ Smoking habits were classified as ‘current smoker’ or ‘non-smoker’ (including never smoked and ex-smoker). In addition, in some of the cohorts a five-point smoking variable was available, where smoking habits were classified as ‘never smoked’ (=1), ‘ex-smoker’ (=2), ‘currently smoking 1–14 g of tobacco per day’ (=3), ‘smoking 15–24 g/day’ (=4), or ‘smoking 25 g or more per day’, with one cigarette equalling 1 g, one cheroot 2 g, one cigar 5 g and pipe tobacco 50 g, divided by the number of days the pack lasted.²³

Ethical considerations

Informed consent was obtained from all participants, verbal in the earlier studies, and written later on, as required first by the Research Ethics Committees at Gothenburg and Uppsala Universities and later by the National Research Ethics Board, that approved the study.

Statistical analyses

Data were analysed using the SAS software, V.9.2²⁴ Data concerning age, sex and examination year were complete, except for one individual where age was missing. Not all variables were measured in all subpopulations. The number of available observations for each variable is shown in [table 2](#). The overall proportion of missing data in subpopulations where the variables were measured was less than 2%. Missing data were not replaced. Simple differences between groups were assessed with Student’s t test or the χ^2 test.

To make full use of SRH as a seven-level ordinal variable, multiple ordinal logistic regression was used, providing cumulative OR across the seven SRH levels for each independent variable, CI and Wald’s χ^2 estimates. The latter is the test parameter and may thus be used to rank the impact or importance of the independent variables on SRH.

The analyses were performed in two steps. In the first step, screening bivariate ordinal logistic regression analyses were performed to identify candidate variables for the final multiple regression model. There were no significant differences in results depending on whether the five-point ordinal smoking variable or the dichotomous smoking variable was used. Since the dichotomous variable was more complete, it was used throughout the analyses.

In the second step, two models were used, one based on age and year of investigation only and the other on all significant variables from step one entered as independent variables. The final full-scale model was based on multiple ordinal logistic regression with backward elimination of non-significant covariates. Possible non-linearity of the effects of age and year of investigation on SRH in the two models was tested by inclusion of the variables age and year of investigation, respectively, raised to the power of 2 and 3 (second-degree and third-degree polynomial functions). Moreover, potential effects of interaction between age and year of investigation were tested, but found to be non-significant.

Two measures of degree of explanation were provided by the SAS logistic procedure, one based on concordance between observed and model computed results, the other based on the receiver operator characteristics (ROC) method. The model fit between crude age and the year of investigation on SRH results and those computed in the final analytical model was assessed by scrutiny and found to be excellent.

The full-scale regression model was also tested with linear regression analysis. The results were generally the

Table 2 Characteristics of the study population

	N	Women		Men		p Value
		n	mean (SD) or %	n	mean (SD) or %	
Number of observations	14470	6816	47.1	7654	52.9	
Age	14469	6816	52.3 (12.6)	7653	56.5 (13.0)	<0.0001
Educational level	14120	6722		7398		<0.0001
Compulsory school		2223	33.1	2773	37.5	
Vocational school		1608	23.9	1905	25.8	
Upper secondary school		1286	19.1	1166	15.8	
University/college		1605	23.9	1554	21.0	
Occupational status	14187	6575		7612		<0.0001
Employed		4444	67.6	4511	59.3	
Unemployed		271	4.1	286	3.8	
Sick-leave/disability pension		713	10.8	624	8.2	
Old-age retirement		1147	17.4	2191	28.8	
Married/cohabiting	14338	4990	74.0	5927	78.0	<0.0001
Smoking habits	14330	6735		7595		0.001
Never smoked or ex-smoker		5002	74.3	5455	71.8	
Current smoker		1733	25.7	2140	28.2	
Leisure time physical activity	13787	6678		7109		<0.0001
Sedentary		1076	16.1	1181	16.6	
Moderately active		4663	69.8	4511	63.4	
Active		890	13.3	1304	18.3	
Vigorously active		49	0.7	113	1.6	
Complaint score (range 0–30)	11365	3777	7.8(5.4)	7588	5.3(4.7)	<0.0001
Self-rated health	14020	6568		7452		<0.0001
1 ('very bad')		132	2.0	136	1.8	
2		253	3.9	204	2.7	
3		387	5.9	423	5.7	
4		1084	16.5	947	12.7	
5		1486	22.6	1373	18.4	
6		1805	27.5	2199	29.5	
7 ('Could not be better')		1421	21.6	2170	29.1	

same as with ordinal logistic regression, but with somewhat lower statistical power. However, the result from the multiple linear regression model was used to compute the regression surfaces in figure 1. All tests were two-tailed. Significance levels were set at $p < 0.10$ in the first, screening analysis step, and $p < 0.05$ in the second, final analysis step.

RESULTS

Characteristics of the study population

Study population characteristics are presented in table 2. The mean age was 54.5 years. Approximately one-quarter of the study population reported college or university education, the majority were employed and married or cohabiting. One-quarter of the participants reported current smoking, and approximately 80% were sedentary or moderately physically active during leisure time. Mean level of SRH was 5.4 (range 1.0–7.0), and mean complaint score 6.1 (range 1.0–30.0). Women were significantly younger than men, had higher level of education, were less often married/cohabiting, less often current smokers, less physically active, reported more symptoms and had lower SRH levels than men.


Effects of age and year of investigation on SRH

In table 3, the effects on SRH of age and year of investigation are shown in women and men according to two models. In model 1, age was significantly inversely associated with SRH in both sexes ($p < 0.0001$), as was year of investigation in women according to a second-degree and in men according to a third-degree polynomial function (non-linear relationships).

In model 2, the linear association between age and SRH among women remained significant ($p < 0.0001$), as did year of investigation, when the covariates were entered into the final model. In men, age and year of investigation, both according to a third-degree polynomial function remained significant, when the covariates were entered in the model.

Table 3 also shows Wald χ^2 estimates, indicating the rank order of variable impact on SRH. In both sexes complaint score had the highest impact on SRH of all variables, followed by sick leave or disability pension, leisure time physical activity, marital status and, finally, education among women and unemployment among men. The final regression model explained 76.2% of the SRH variance in women, and 74.5% of the SRH variance in men.

Figure 1 Association between age, year of investigation (proxy for secular trend) and self-rated health (SRH) in women and men, adjusted for the influence on SRH of education, physical exercise during leisure time, marital status, complaint score, on sick leave/disability pension and being unemployed. Dark bars=data available, white bars=no data available.



Effect of participating in more than one investigation

All women participated only once, but 1548 men participated in more than one examination. An analysis was therefore made with all 1548 male observations from more than one examination, and one with the 3644

men who participated only once. The results were almost identical, the only difference being that in analyses of those participating only once age was significant but year of investigation was not (because of no intraindividual variation?) and among those participating more

Table 3 Effects of age and year of investigation on self-rated health (SRH), based on ordinal logistic regression models

	OR	95% CI	Wald	p Value
Women				
Model 1				
Age	0.98	0.98 to 0.99	75.3	<0.0001
Year of investigation	0.73	0.58 to 0.92	7.4	<0.007
Year of investigation ²	1.00	1.00 to 1.01	4.5	0.03
Model 2*				
Age	0.98	0.98 to 0.98	67.3	<0.0001
Year of investigation	0.95	0.93 to 0.96	42.0	<0.0001
Complaint score	0.82	0.81 to 0.83	889.5	<0.0001
Sick-leave/disability pension	0.35	0.28 to 0.42	106.0	<0.0001
Leisure time physical activity	1.51	1.35 to 1.69	52.4	<0.0001
Married	1.19	1.04 to 1.37	6.5	0.01
Education	1.07	1.01 to 1.13	4.7	0.03
Men				
Model 1				
Age	0.70	0.63 to 0.78	39.7	<0.0001
Age ²	1.01	1.00 to 1.01	31.2	<0.0001
Age ³	1.00	1.00 to 1.00	25.5	<0.0001
Year of investigation	0.94	0.90 to 0.98	7.7	<0.006
Year of investigation ²	1.01	1.00 to 1.01	12.5	<0.0005
Year of investigation ³	1.00	1.00 to 1.00	17.7	<0.0001
Model 2*				
Age	0.70	0.61 to 0.79	30.1	<0.0001
Age ²	1.01	1.00 to 1.01	20.6	<0.0001
Age ³	1.00	1.00 to 1.00	14.8	0.0001
Year of investigation	0.87	0.83 to 0.92	23.9	<0.0001
Year of investigation ²	1.01	1.01 to 1.01	23.9	<0.0001
Year of investigation ³	1.00	1.00 to 1.00	25.5	<0.0001
Complaint score	0.80	0.79 to 0.80	1606.3	<0.0001
Sick-leave/disability pension	0.36	0.31 to 0.43	144.6	<0.0001
Leisure time physical activity	1.48	1.38 to 1.59	114.2	<0.0001
Married	1.33	1.20 to 1.48	27.4	<0.0001
Unemployed	0.79	0.63 to 1.00	3.9	<0.05

*The following variables were removed in the backwards elimination procedure. *Women*: current smoking (p=0.49), unemployed (p=0.07). *Men*: current smoking (p=0.54), education (p=0.20)

than once age was non-significant, while year of investigation was.

Effects of time differences

The observations on women covered a shorter time period (1993–2003) than those on men (1973–2003). To explore if this time difference could explain the difference in time trends between men and women an analysis was performed among men with data covering the time period 1993–2003. The results showed the same determinants for men as those shown in table 3 and thus a curvilinear time trend, as opposed to the linear trend in women.

Visualisation of the analysis model

In figure 1 the analysis model is visualised. In women mean SRH levels fell gradually and linearly from 5.5 among the youngest to 4.6 among the oldest women. Furthermore, mean SRH levels fell across year of investigation, from 5.2 in 1993 to 4.9 in 2003. In men, the association between age and year of investigation on the one hand, and SRH on the other followed a third-degree polynomial function with peaks at 25 years of age and at the age interval 50–70 along the age axis, as well as in 1973 and in the late 1990s along the year of investigation axis.

DISCUSSION

In line with a majority of previous studies investigating effects of age on SRH in adult population-based studies,^{6 11 16} in this population-based study of 12 000 adult women and men, there was a significant, independent relationship between SRH and age in both sexes, resulting in lower SRH with increasing age. The association was independent of year of investigation and remained when adjusted for socio-demographic variables, lifestyle variables and complaint score. In women the association between SRH and age followed a linear function. In men the association was more complex. In a recent study, based on employees from the French national gas and electricity company an overall pattern of age-related gradual decline in SRH was observed, with distinctly better SRH observed in women and men during the year preceding and the year following retirement from the work force.⁸ In the present population-based study, no distinct changes in SRH around retirement age were observed, which suggests that the suggested associations between retirement and SRH may apply in specific populations and contexts, but not necessarily in the population as a whole.

Moreover, there was a significant association between year of investigation and SRH, which was independent of age and covariates. In women, the association between SRH and year of investigation followed a linear function, with gradually decreasing SRH levels over time, and in men the association followed a third-degree polynomial function with fluctuations but no persistent

decrease over time. Results from previous studies concerning associations between year of investigation and SRH are controversial.^{6 11 16 17} However, these studies differ in terms of whether or not medical health and life-style-related variables are controlled in analyses, and also in terms of study-population characteristics. Contextually determined perceptions of health may, for example, be expected to differ between countries. Present results are in line with results from a recent study,⁶ in which poorer age-specific SRH in more recent years of investigation was observed in a Danish study population, but differ with regard to results from the USA, British and German study populations where no age-specific differences were observed during the same time period.

The main strength of the study concerns sample size. Data were based on eight ongoing population studies, covering more than 14 000 observations scattered over a 30-year period. The sample size was thus larger than in previous studies investigating effect of year of investigation on SRH, and observation time spanned up to three decades in men and one decade in women. The determinant variables were measured with the same instruments in all cohorts.

A further strength concerns the use of ordinal regression modelling.^{25 26} SRH is an ordered categorical variable. The majority of previous studies investigating predictors of multilevel SRH either used binary logistic regression technique, despite the reduced efficiency and the risk of bias related to choice of cut point this entails, or analytical methods requiring normally distributed SRH data, a requirement rarely satisfied.

The main limitation of the study is that all age groups were not represented on each measurement occasion, and that some model-based results were thus generated for age groups where no actual measurements were obtained, as shown in figure 1. However, this is a problem common in multiple analysis modelling, and not unique to this study.

Moreover, the data were generally cross-sectional. Even so, some of the cohorts were investigated repeatedly and thus form a truly longitudinal cohort. Results were similar for subjects who participated only once and subjects who participated on several occasions, indicating that the results concerning age and secular trends were stable, and that the design used here is equivalent to a longitudinal one.

The attrition rate was a moderate 28%. To test the potential effect of attrition on the secular trend slope, results were re-analysed based on the assumption that non-participants in the early and the late parts of the study period reported considerably lower or higher levels of SRH than participants. However, even the most unfavourable alternative from the hypothesis point of view (non-participants in the early study period reporting 5 SE units lower SRH than participants, and 5 SE units higher SRH at the end of the study period) was associated with a highly significant decrease in mean SRH over

calendar time. We therefore have no reason to believe that the results are afflicted by selection or other bias to such an extent that the conclusions would be affected.

The results from this study regarding the age–SRH relationship are in line with the majority of previous studies,^{6–11} reporting decreasing SRH levels with age. Results concerning associations between SRH and year of investigation suggested an inverse trend in self-ratings of health over time in women, but less so in men. Previous results are controversial, some studies showing increasingly poorer SRH in more recent cohorts, others showing better SRH.^{12–14 16–18} Changes in SRH over time, on the one hand, may reflect changes in patterns of morbidity or, on the other hand, contextually determined differences in norms and expectations in relation to health,^{13 27} and furthermore, may or may not suggest contextually determined changes in healthcare utilisation. Concepts of health and knowledge and expectations concerning health and healthcare may change over time. One important question from a public health perspective concerns to what extent observed trends are associated with changes in morbidity, mortality and healthcare utilisation. Given that life expectancy has increased over the period it is unlikely that the decrease in SRH reflects a major deterioration in health in the population.

The relationship between age and SRH differed between women and men, as did the observed association between year of investigation and SRH. A linear decline in SRH was observed among women, and a more complex age–SRH relation observed among men. Few studies have, to date, investigated potential non-linearity of associations between age or year of investigation, and SRH.^{11 14 17} A non-linear association between age and SRH in women and men,^{11 14} and a linear cohort effect (measured as 10-year birth cohorts) on SRH,¹⁴ has been reported, but whether women and men differ in functional form of the observed association between age, or cohort, and SRH, was not investigated. In a study of 618 randomly sampled middle-aged women, independent, inverse, linear and quadratic age effects on SRH was reported, with poorer SRH and a more rapid decline in SRH per year in the later cohort.¹⁷

Although similar patterns of determinants of SRH have been observed in women and men,²⁸ gender-related differences in the association between SRH and subsequent mortality have repeatedly been observed,^{29 30} and, accordingly, the relative importance of various contextually related factors of importance for health may be different for women and men.¹⁵ The present results thus underline the importance of investigating effects of contextually related factors, such as age and year of investigation on SRH separately for women and men.

The differing trends among women and men may, furthermore, be explained by the analysis modelling, since the women were less scattered over the years of investigation than men, as shown in figure 1. However, a great deal of effort was devoted to the analysis modelling, with a number of possible models tested. The model with the

best fit between the model data and crude data was finally accepted. It is therefore unlikely that the differing functional form for women and men is an analytical artefact.

The results on SRH predictors were similar for women and men, and in line with previous research on determinants of SRH, in which illness, disease and disability were the strongest predictors of SRH.⁵ Variables included in the final ordinal regression model in the present study explained 76.2% of the variance in SRH in women, and 74.5% of the variance in SRH in men, which implies that the model captures core determinants of global SRH. In our analyses we were restricted to covariates that were available in the datasets common to all subpopulations in this study. The remaining unexplained 25% may be attributable to predictors not included in our model, perhaps different ones in women and men.

Author affiliations

¹Department of Public Health and Caring Sciences, Family Medicine and Preventive Medicine Section, Uppsala University, Uppsala, Sweden

²Department of Heart and Lung Diseases, Sahlgren Academy, Gothenburg, Sweden

³R&D Centre/Centre for Clinical Research, Section of Primary Care, Sörmland County Council, Eskilstuna, Sweden

⁴Department of Epidemiology, AstraZeneca R&D, Mölndal, Sweden

Acknowledgements The study was supported by grants from the Swedish Research Council, the Swedish Research Council for Working Life and Social Research, the Vårdal Foundation, the Sörmland County Council and Gothenburg and Uppsala Universities, Sweden.

Contributors CW, TW, AR, AB, SJ, HE and KS performed the data collection in their respective datasets. CH and KS designed the analyses. JB, CH and KS contributed with SAS programming. CH and KS drafted the manuscript. All authors participated in the critical discussions of the results, identification of relevant reference literature and in the revisions of the manuscript.

Funding The study was supported by grants from the Swedish Research Council, the Swedish Research Council for Working Life and Social Research, the Vårdal Foundation, the Sörmland County Council and Gothenburg and Uppsala Universities, Sweden.

Competing interests None.

Ethics approval Research Ethics Committees at Gothenburg and Uppsala Universities and by the National Research Ethics Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The dataset is based on a research consort of groups in Gothenburg, Uppsala and Eskilstuna. Requests for data access may be made to Kurt Svärdsudd.

REFERENCES

1. Idler EL, Russel LB, Davis D. Survival, functional limitations, and self-rated health in the NHANES I epidemiologic follow-up study, 1992. *Am J Epidemiol* 2000;52:874–83.
2. Mulunpalo S, Vuori I, Oja P, *et al.* Self-rated health status as a health measure: the predictive value of self-reported health status on the use of physician services and on mortality in the working-age population. *J Clin Epidemiol* 1997;50:517–28.
3. Kaplan GA, Goldberg DE, Everson SA, *et al.* Perceived health status and morbidity and mortality: evidence from the Kuopio ischaemic heart disease risk factor study. *Int J Epidemiol* 1996;25:259–65.
4. Idler EL, Benyamini Y. Self-rated health and mortality: a review of twenty-seven community studies. *J Health Soc Behav* 1997;38:21–37.
5. deSalvo KB, Bloser N, Reynolds K, *et al.* Mortality prediction with a single general self-rated health question. A meta-analysis. *J Gen Intern Med* 2006;21:267–75.

6. Sacker A, Worts D, McDonough P. Social influences on trajectories of self-rated health: evidence from Britain, Germany, Denmark and the USA. *J Epidemiol Community Health* 2011;65:130–6.
7. Sargent-Cox KA, Anstey KJ. Patterns of longitudinal change in older adults' self-rated health: The effect of point of reference. *Health Psychol* 2010;29:143–52.
8. Westerlund H, Kivimäki M, Singh-Manouf A, *et al.* Self-rated health before and after retirement in France (GAZEL): a cohort study. *Lancet* 2009;374:1889–96.
9. Krarup Andersen F, Christensen K, Frederiksen H. Self-rated health and age: a cross-sectional and longitudinal study of 11,000 Danes aged 45–102. *Scand J Public Health* 2007;35:164–71.
10. McFadden E, Luben R, Bingham S, *et al.* Social inequalities in self-rated health by age: cross-sectional study of 22 457 middle-aged men and women. *BMC Public Health* 2008;8:230.
11. Chen H, Cohen P, Kasen S. Cohort differences in self-rated health: evidence from a three-decade, community-based, longitudinal study of women. *Am J Epidemiol* 2007;166:439–46.
12. Liang J, Shaw BA, Krause N, *et al.* How does self-assessed health change with age? A study of older adults in Japan. *J Gerontol B-Psychol* 2005;60:S224–32.
13. Idler EL. Age differences in self-assessments of health: age changes, cohort differences, or survivorship? *J Gerontol B-Psychol* 1993;48:S289–300.
14. Hoeymans N, Feskens EJM, van den Bos GAM, *et al.* Age, time, and cohort effects on functional status and self-rated health in elderly men. *Am J Public Health* 1997;87:1620–5.
15. Liu H. Marital dissolution and self-rated health: age trajectories and birth cohort variations. *Soc Sci Med* 2012;74:1107–16.
16. Yang Y, Lee LC. Sex and race disparities in health: cohort variations in life course patterns. *Soc Forces* 2009;87:2093–124.
17. Salomon JA, Nordhagen S, Oza S, *et al.* Are Americans feeling less healthy? The puzzle of trends in self-rated health. *Am J Epidemiol* 2009;170:343–51.
18. Svedberg P, Gatz M, Lichtenstein P, *et al.* Self-rated health in a longitudinal perspective: a 9-year follow-up twin study. *J Gerontol B-Psychol* 2005;60B:S331–40.
19. Zack MM, Moriarty DG, Stroup DF, *et al.* Worsening trends in adult health-related quality of life and self-rated health—United States, 1993–2001. *Public Health Rep* 2004;119:493–505.
20. Wallman T, Wedel H, Palmer E, *et al.* Sick-leave track record and other potential predictors of a disability pension. A population based study of 8,218 men and women followed for 16 years. *BMC Public Health* 2009;9:104.
21. Wallman T, Wedel H, Johansson S, *et al.* The prognosis for individuals on disability retirement. An 18-year mortality follow-up study of 6887 men and women sampled from the general population. *BMC Public Health* 2006;6:103.
22. Tibblin G, Tibblin B, Peciva S, *et al.* 'The Göteborg Quality of Life Instrument'—an assessment of well-being and symptoms among men born 1913 and 1923. *Scand J Prim Health* 1990 (Suppl 1):33–8.
23. Rose GA, Blackburn H, Gillum RF, *et al.* *Cardiovascular survey methods*. Geneva: The World Health Organization, 1982.
24. SAS (Statistical Analysis System). <http://www.sas.com/technologies/analytics/statistics/statfactsheet.pdf> (accessed 25 Jan 2010).
25. Scott SC, Goldberg MS, Mayo NE. Statistical assessment of ordinal outcomes in comparative studies. *J Clin Epidemiol* 1997;50:45–55.
26. Armstrong B, Sloan M. Ordinal regression models for epidemiologic data. *Am J Epidemiol* 1989;129:191–204.
27. Salomon JA, Tandon A, Murray CJL. Comparability of self-rated health: cross sectional multi-country survey using anchoring vignettes. *BMJ* 2004;328:258–61.
28. Undén A-L, Elofsson S. Do different factors explain self-rated health in men and women? *Gender Med* 2006;3:295–308.
29. Spiers N, Jagger C, Clarke M, *et al.* Are gender differences in the relationship between self-rated health and mortality enduring? Results from three birth cohorts in Melton Mowbray, United Kingdom. *Gerontologist* 2003;43:406–11.
30. Deeg DJH, Kriegsman DMW. Concepts of self-rated health: specifying the gender difference in mortality risk. *Gerontologist* 2003;43:376–86.

Correction

Halford C, Welin C, Bogefeldt J, *et al.* A population-based study of nearly 15 000 observations among Swedish women and men during 1973–2003. *BMJ Open* 2012;**2**:e001353. Part of the article title was erroneously omitted. The full title should read 'Effects of age and secular trends on self-rated health. A population-based study of nearly 15 000 observations among Swedish women and men during 1973–2003'.

BMJ Open 2012;**2**:e001353corr1. doi:10.1136/bmjopen-2012-001353corr1

Correction

Kausto J, Solovieva S, Virta LJ, *et al.* Partial sick leave associated with disability pension: propensity score approach in register-based cohort study. *BMJ Open* 2012;2:e001752. Incorrect figures were published in table 5. The corrected table is published below.

BMJ Open 2012;2:e001752corr1. doi:10.1136/bmjopen-2012-001752corr1

Table 5 Absolute and relative risk reduction (a risk difference between partial and full sick leave groups, ARR* and RRR*) of the use of partial and full disability pension with 95% CIs in the matched subsample

	Full disability pension				Partial disability pension					
	ARR (%)	95% CI	RRR (%)	95% CI	NNT	ARR (%)	95% CI	RRR (%)	95% CI	NNT
Total	6	3 to 9	41	24 to 55	16	-8	-10 to (-5)	-159	-264 to (-84)	-14
Men	10	4 to 16	47	20 to 65	10	-5	-10 to (-0)	-102	-302 to (-2)	-21
Women	4	0.5 to 7	32	5 to 52	26	-9	-12 to (-6)	-180	-315 to (-89)	-12
Mental disorders	8	3 to 13	42	17 to 59	12	-10	-14 to (-7)	-361	-795 to (-138)	-10
Musculoskeletal disorders	6	2 to 11	51	19 to 70	15	-7	-12 to (-2)	-94	-202 to (-24)	-15
Traumas and tumours	2	-4 to 8	17	-54 to 56	53	0	-8 to 2	-95	-409 to 25	-31

*Negative values indicate an increase in risk.

*NNT, number needed to treat to prevent full disability pension.