

Social media use among patients DEN and caregivers: a scoping review

Michele P Hamm, Annabritt Chisholm, Jocelyn Shulhan, Andrea Milne, Shannon D Scott, Lisa M Given, Lisa Hartling

To cite: Hamm MP, Chisholm A. Shulhan J. et al. Social media use among patients and caregivers: a scoping review. BMJ Open 2013;3:e002819. doi:10.1136/bmjopen-2013-002819

Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2013-002819).

Received 1 March 2013 Revised 11 April 2013 Accepted 15 April 2013

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see http://bmjopen.bmj.com

For numbered affiliations see end of article.

Correspondence to

Dr Michele P Hamm; michele.hamm@ualberta.ca

ABSTRACT

Objective: To map the state of the existing literature evaluating the use of social media in patient and caregiver populations.

Design: Scoping review.

Data sources: Medline. CENTRAL. ERIC. PubMed. CINAHL Plus Full Text, Academic Search Complete, Alt Health Watch, Health Source, Communication and Mass Media Complete, Web of Knowledge and ProQuest (2000-2012).

Study selection: Studies reporting primary research on the use of social media (collaborative projects, blogs/microblogs, content communities, social networking sites, virtual worlds) by patients or

Data extraction: Two reviewers screened studies for eligibility; one reviewer extracted data from relevant studies and a second performed verification for accuracy and completeness on a 10% sample. Data were analysed to describe which social media tools are being used, by whom, for what purpose and how they are being evaluated.

Results: Two hundred eighty-four studies were included. Discussion forums were highly prevalent and constitute 66.6% of the sample. Social networking sites (14.8%) and blogs/microblogs (14.1%) were the next most commonly used tools. The intended purpose of the tool was to facilitate self-care in 77.1% of studies. While there were clusters of studies that focused on similar conditions (eq. lifestyle/weight loss (12.7%). cancer (11.3%)), there were no patterns in the objectives or tools used. A large proportion of the studies were descriptive (42.3%); however, there were also 48 (16.9%) randomised controlled trials (RCTs). Among the RCTs, 35.4% reported statistically significant results favouring the social media intervention being evaluated; however, 72.9% presented positive conclusions regarding the use of social media. **Conclusions:** There is an extensive body of literature examining the use of social media in patient and caregiver populations. Much of this work is descriptive; however, with such widespread use, evaluations of effectiveness are required. In studies that have examined effectiveness, positive conclusions are often reported, despite non-significant findings.

INTRODUCTION

The use of social media in healthcare has been widely advocated¹⁻⁸; however, there is little

ARTICLE SUMMARY

Article focus

- The use of social media in healthcare has been widely advocated, but there is little evidence describing the current state of the science and whether or not these tools can be used to benefit patient populations.
- We mapped the state of the existing literature evaluating the use of social media in patient and caregiver populations.

Key messages

- There is an extensive and rapidly growing body of literature available investigating the use of social media in patient and caregiver populations.
- Most studies have been descriptive; however, with such widespread use, evaluations of effectiveness are needed.
- In studies that have examined effectiveness. positive conclusions are often reported, despite the non-significant findings.

Strengths and limitations of this study

- Our search was comprehensive and we included an extensive body of literature, across conditions, populations and study designs.
- Social media is constantly evolving, leading to challenges in keeping the search updated.
- A more in-depth analysis is needed on specific topics, conditions and populations to guide the use and implementation of social media interventions.

evidence describing the current state of the science and whether or not these tools can be used to benefit patient populations. It is clear, though, that in addition to seeking out traditional sources of healthcare information, patients are increasingly active online.⁹ In 2011, looking for healthcare information was the third most common online activity, ¹⁰ and in September 2012, 72% of adult internet users sought support and medical information online. 11 In 2012, 67% of internet users were using social media for any purpose¹² and 26% were using it for health issues. 11 As social media continues to evolve, its momentum shows no sign of diminishing, instead finding new niches with unique applications.

Social media can be defined as a group of online applications that allow for the creation and exchange of user-generated content, and can be categorised into five groups: (1) collaborative projects (eg, Wikipedia); (2) blogs or microblogs (eg, Blogger, Twitter); (3) content communities (eg, YouTube); (4) social networking sites (eg, Facebook) and (5) virtual gaming or social worlds (eg, HumanSim).¹³ The collaborative environment to which social media belongs represents a shift in technology and functionality from 'Web 1.0', in which static online content and applications were created and published by individuals, to 'Web 2.0', in which there is continuous modification and participation by all users.¹³ Table 1 provides an overview of the categories of social media tools.

Advocates of the use of social media in healthcare suggest that these tools allow for personalisation, presentation and participation—three key elements that make them highly effective. 14 The content can be tailored to the priorities of the users; the versatility of the different platforms creates numerous options for the presentation of information, and the collaborative nature of social media allows for a meaningful contribution from all user groups. The idea of a synergistic relationship between social media users is one of the main perceived advantages of using these platforms. 15 However, criticisms of the use of social media in healthcare have also arisen. The availability of misinformation is a risk, as healthcare providers are unable to control the content that is posted or discussed. I 16 17 Inappropriate substitution of online information or advice for in-person visits to a healthcare provider can also potentially lead to harmful results, and this has been cited as a limitation of the use of social media and of the internet generally. 1 18 Negative uses of social media have also been highlighted in the context of professionalism and confidentiality, 19 use by children and youth due to a limited capacity for self-regulation and vulnerability to peer

influence, 20 and promotion of high-risk behaviours, such as suicide-related behaviours, drug use and eating disordered behaviours. $^{21-24}$

The objectives of this study were to map the existing literature examining the use of social media in patient and caregiver populations, to determine the extent and type of evidence available to inform more focused knowledge syntheses and to identify gaps for future research. The specific questions guiding this scoping review were: (1) What social media tools are being used to improve health outcomes in patient populations? (2) For what purposes are social media tools being used in patient populations (eg, to improve health literacy, to improve self-care)? (3) For what patient populations and disease conditions are social media tools being used? (4) What types of evidence and research designs (ie, qualitative, quantitative) have been used to examine social media tools?

METHODS

This scoping review on the use of social media in patient and caregiver populations was conducted in parallel with a review on the use of social media in health-care professional and trainee populations²⁵; therefore, the literature search and screening for study eligibility were conducted concurrently. The review followed a protocol that we developed a priori.

Search strategy

A research librarian searched 11 databases in January 2012: Medline, CENTRAL, ERIC, PubMed, CINAHL Plus Full Text, Academic Search Complete, Alt Health Watch, Health Source, Communication and Mass Media Complete, Web of Knowledge and ProQuest. Dates were restricted to 2000 or later, corresponding to the advent of Web 2.0. No language or study design restrictions

Tool	Description	Examples
Collaborative projects	Enable the joint and simultaneous creation of content by many end-users	Wikis (eg, Wikipedia) Social bookmarking applications (eg, Mendeley)
Blogs or	Websites that display date-stamped entries. They are usually managed	Wordpress
microblogs	by one person but provide the opportunity to interact with others through the addition of comments	Twitter (microblog)
Content communities	Allow for the sharing of media content between users, including text, photos, videos and presentations	BookCrossing Flickr YouTube Slideshare
Social networking sites	Enable users to connect by creating personal information profiles that can be accessed by friends and colleagues, and by sending emails and instant messages between each other	Facebook MySpace LinkedIn
Virtual worlds	Platforms that replicate a 3D environment in which users can appear in the form of personalised avatars and interact with each other as they would in real life	Second Life

were applied. The search strategy for Medline is provided in the online supplementary appendix.

Study selection

Two reviewers independently screened titles and abstracts of studies for eligibility. The full text of studies assessed as 'relevant' or 'unclear' was then independently evaluated by two reviewers using a standard form. Discrepancies were resolved by consensus or adjudication by a third party.

Studies were included if they reported primary research (quantitative or qualitative), focused on healthcare issues related to patients or caregivers, and examined the use of a social media tool. Social media was defined according to Kaplan and Haenlein's classification scheme, 13 including: collaborative projects, blogs or microblogs, content communities, social networking sites and virtual worlds. We excluded studies that examined mobile health (eg, tracking or medical reference apps), one-way transmission of content (eg, podcasts) and real-time exchanges mediated by technology (eg, Skype, chat rooms). Electronic discussion forums and bulletin boards were included as they incorporate user-generated content and were judged to fall within the spectrum of social media. Outcomes were not defined a priori as they were to be incorporated into our description of the field. The most likely categories for objectives and outcomes were adapted from those outlined in Coulter and Ellins^{,26} ²⁷ proposed framework for strategies to inform, educate and involve patients.

Data extraction

Data were extracted using standardised forms and entered into Microsoft Excel (Microsoft, Redmond, Washington, USA) by one reviewer and a 10% sample was checked for accuracy and completeness by another. Reviewers resolved discrepancies through consensus. Extracted data included study and population characteristics, description of the social media tools used, objective of the tools, outcomes measured and authors' conclusions. Studies that examined social media as one component of a complex intervention were noted as such. Additional data were collected for randomised controlled trials (RCTs), including the primary outcome and its statistical significance.

Data synthesis

Data were synthesised descriptively to map different aspects of the literature as outlined in our key questions. Studies were grouped according to tool, audience and study design, with data from RCTs being examined in more detail. As discussion forums were not included in our original classification scheme, findings are presented both for all included studies and for studies that investigated tools other than discussion forums. Descriptive statistics were calculated using StataIC 11 (StataCorp, College Station, Texas, USA).

RESULTS

Two hundred eighty-four studies were included in the review. Figure 1 outlines the flow of studies through the inclusion process and table 2 provides a description of the included studies. Most studies (179/284; 63.0%) were conducted in North America, with more than half of the total sample (154/284; 54.2%) carried out in the USA and 8.8% (25/284) conducted in Canada. The median start date was in 2006 (range 1997-2011); when studies evaluating discussion forums were excluded, the start date was more recent (median 2008, range 2000-2011). Studies tended to be fairly short, with a median duration of 5 months (range 1-117 months). Nearly all included studies were published as journal articles (255/284; 89.8%); however, when studies of discussion forums were excluded, the proportion of dissertations written on the use of social media increased (14/284 to 12/95; 4.9–12.6%).

Social media tools used

The social media tools studied are outlined in table 3. The use of discussion boards and online support groups (combined as discussion forums due to their common structure and intent) dominated the literature, encompassing 189 (66.6%) included studies. Social networking sites (42/284; 14.8%) and blogs or microblogs (40/284; 14.1%) were also commonly evaluated, followed by content communities (16/284; 5.6%), collaborative projects (6/284; 2.1%) and virtual worlds (6/284; 2.1%). In




Figure 1 Flow diagram of included studies.

Variable	Total—n (%)	Excluding discussion forums—n (%)	
Total—N	284	95	
Continent of corresponding author			
Asia	12 (4.2)	5 (5.3)	
Australia	14 (4.9)	3 (3.2)	
Europe	78 (27.5)	19 (20.0)	
North America	179 (63.0)	67 (70.5)	
Not reported	1 (0.4)	1 (1.1)	
Study start date—median (range)	2006 (1997–2011)	2008 (2000–2011)	
Study duration—median (range)	5 months (1–117)	3 months (1–117)	
Sample size—median (range)	124 (1–16703)*	130 (2–16703)*	
Publication type	,	,	
Journal article	255 (89.8)	75 (79.0)	
Abstract	15 (S.3)	8 (8.4)	
Dissertation	14 (4.9)	12 (12.6)	
Study design		` '	
Quantitative			
Randomised controlled trial	48 (16.9)	6 (6.3)	
Non-randomised controlled trial	6 (2.1)	1 (1.1)	
Controlled before-after	1 (0.4)	_ ` ′	
Observational	11 (3.9)	3 (3.2)	
Cross-sectional	63 (22.2)	33 (34.7)	
Qualitative	,	,	
Case study	1 (0.4)	_	
Case series	3 (1.1)	2 (2.1)	
Ethnography	3 (1.1)	2 (2.1)	
Grounded theory	6 (2.1)	2 (2.1)	
Phenomenology	6 (2.1)	1 (1.1)	
Qualitative (other/not specified)	46 (16.2)	16 (16.8)	
Mixed methods	33 (11.6)	9 (9.5)	
Other	,	,	
Content analysis	57 (20.1)	20 (21.1)	
Authors' conclusions	,	,	
Positive	186 (65.5)	56 (59.0)	
Neutral	65 (22.9)	23 (24.2)	
Negative	15 (5.3)	10 (10.5)	
Indeterminate	18 (6.3)	6 (6.3)	

116 (40.9%) included studies, the social media tool was included as part of a complex intervention. Where existing and publicly available social media applications were studied, Facebook (16/284; 5.6%), YouTube (12/284; 4.2%) and Twitter (10/284; 3.5%) were evaluated most frequently (figure 2).

Purposes of social media use

The most common intended use of social media was for self-care, which was described as an objective of the tool in 219 (77.1%) studies (table 3). This was particularly relevant to discussion forums, in which 166/189 (87.8%) studies were related to self-care. Other tools were often established with similar functions to discussion forums: they provided a platform on which users could post and share their experiences with peers. Collaborative projects were often used to address health

literacy, and social networking sites were commonly used for patient safety purposes, largely for documentation of adverse events. While there were few studies that addressed clinical decision-making, these were almost exclusively conducted using discussion forums.

We categorised the outcomes measured in each of the studies under patients' knowledge, patients' experience, use of services and costs, health behaviour and status and others (table 4). Measures of patients' experience, specifically peer-to-peer communication (135/284; 47.5%), were most common and were often outcomes related to social support among members of an online community. Measures of psychological well-being (eg, reports of anxiety levels) and changes in self-care activities (eg, increases in physical activity) in relation to use of the tool were also commonly evaluated (78/284 and 63/284; or 27.5% and 22.2%, respectively).

Table 3 Description and objectives of social media tools used (N=284)								
		Objective—n (%)						
Tool	Total—n (%)	Health literacy	Clinical decision making	Self-care	Patient safety	Others		
Total—n (%)		47 (16.6)	7 (2.5)	219 (77.1)	19 (6.7)	39 (13.7)		
Collaborative project	6 (2.1)	5 (83.3)	-	_	_	1 (16.7)		
Blog or microblog	40 (14.1)	11 (27.5)	_	24 (60.0)	4 (10.0)	9 (22.5)		
Content community	16 (5.6)	8 (50.0)	-	5 (31.3)	2 (12.5)	4 (25.0)		
Social networking site	42 (14.8)	10 (23.8)	1 (2.4)	24 (57.1)	8 (19.1)	9 (21.4)		
Virtual world	6 (2.1)	3 (50.0)	_	3 (50.0)	1 (16.7)	1 (16.7)		
Discussion forum	189 (66.6)	23 (12.2)	6 (3.2)	166 (87.8)	3 (1.6)	17 (9.0)		
Component of a complex intervention	116 (40.9)	16 (13.8)	3 (2.6)	108 (93.1)	4 (3.5)	3 (2.6)		
Percentages do not add up to 100 due to the possibility of multiple tools and multiple objectives per study.								

Social media user groups


A wide range of conditions were covered in the included studies (figure 3). The largest proportion fell under the lifestyle and weight loss category (36/284; 12.7%), followed by cancer (32/284; 11.3%) and studies in the general population (22/284; 7.8%). The general population studies tended to be surveys focused on usage, demographics and user preferences relevant to social media use for health-related purposes. No strong trends emerged showing differences between user groups in the objective of the type of social media tool or the specific application used (data not shown). In nearly all conditions investigated, the social media tool studied was intended to facilitate self-care. One exception was seen in the case of infectious disease, where 7/12 (58.3%) relevant studies were focused on health literacy. This was mainly driven by large-scale strategies to provide updates on influenza or H1N1. For specific applications used, there were clusters of studies that examined conditionspecific modalities. Social networking sites were common in studies of diabetes and metabolic syndrome due to the use of TuDiabetes, an online community targeted to those affected by diabetes. Similarly, Twitter was

commonly used in the context of H1N1/influenza, and PatientsLikeMe was used for a group of chronic conditions including amyotrophic lateral sclerosis, fibromyalgia, HIV, mood disorders, multiple sclerosis and Parkinson's disease. Aside from these small clusters, most studies across all conditions were conducted using discussion forums.

Evaluation of social media use

The majority of the included studies were descriptive: 63 (22.2%) were cross-sectional and 57 (20.1%) used content analysis to outline how social media is being applied (table 2). Qualitative studies comprised 22.9% (65/284) of the total sample; mixed methods studies 11.6% (33/284); observational studies 3.9% (11/284) and experimental studies 19.4% (55/284). Of the 33 mixed methods studies, 11 included a cross-sectional component and 20 included content analyses. Forty-eight RCTs were conducted, 45 of which were evaluating discussion forums as at least one component of the intervention. Of the remaining RCTs, one evaluated a blog, one evaluated Second Life and one made use of Facebook and Twitter.

Figure 2 Specific social media tools described in included studies.



Outcomes	Total—n (%)	Excluding discussion forums—n (%)	
Total—N	284	95	
Patients' knowledge			
Conditions and complications	54 (19.0)	22 (23.2)	
Self-care	60 (21.1)	17 (17.9)	
Treatment options	22 (7.8)	10 (10.5)	
Comprehension	2 (0.7)	1 (1.1)	
Patients' experience			
Satisfaction	69 (24.3)	21 (22.1)	
Clinician–patient communication	39 (13.7)	16 (16.8)	
Peer-to-peer communication	135 (47.5)	44 (46.3)	
Quality of life	20 (7.0)	2 (2.1)	
Psychological well-being	78 (27.5)	21 (22.1)	
Self-efficacy	32 (11.3)	4 (4.2)	
Involvement and empowerment	22 (7.8)	6 (6.3)	
Use of services and costs			
Hospital admission rates	4 (1.4)	2 (2.1)	
Emergency admission rates	2 (0.7)	_	
Number of visits to general practitioners	7 (2.5)	2 (2.1)	
Cost effectiveness	4 (1.4)	3 (3.2)	
Health behaviour and status			
Self-care activities	63 (22.2)	15 (15.8)	
Treatment adherence	13 (4.6)	1 (1.1)	
Severity of disease or symptoms	17 (6.0)	4 (4.2)	
Physical functioning	21 (7.4)	6 (6.3)	
Mental functioning	25 (8.8)	8 (8.4)	
Clinical indicators	23 (8.1)	3 (3.2)	
Others			
Attitudes and preferences	14 (4.9)	7 (7.4)	
Content and accuracy	33 (11.6)	21 (22.1)	
Usability	9 (3.2)	2 (2.1)	
Usage and demographics	106 (37.3)	34 (35.8)	

Overall, 186/284 (65.5%) studies concluded that there was evidence for the utility of social media, while only 15/284 (5.3%) concluded that there was not (table 5). The subset of RCTs was examined in more detail; while 35/48 (72.9%) studies presented positive conclusions, only 16/35 (45.7%) reported a statistically significant effect in relation to the primary outcome (figure 4). All but one study with significant findings evaluated the use of a discussion forum; the other study evaluated a blog. Clusters of conditions appeared in the RCTs: six studies were related to lifestyle and weight loss, three were related to tobacco and substance use, two were in mental health and six were in other conditions (diabetes, irritable bowel syndrome, multiple sclerosis, hearing loss and breast cancer). The primary outcome in each of these studies was related to health behaviour and status, except two that evaluated patients' experience and one that measured website use. The social media tool was one component of a complex intervention in all studies, making it difficult to tease out any effect specific to its use. However, improvements were found in outcomes such as changes in body weight and activity levels, tobacco or substance use and quality of life.

DISCUSSION

There is an extensive and rapidly growing body of literature available for investigating the use of social media in patient and caregiver populations. While diversity exists in terms of the tools used, their intended purposes and the conditions studied, the majority of studies evaluate discussion forums. This could point to the popularity of discussion forums among patients and caregivers in addressing their healthcare concerns; however, it may also be indicative of the behaviours or preferences of the site designers.

While general tools with broad applications (ie, discussion forums) are commonly used, the promise of social media lies in its adaptability. Unique applications such as PatientsLikeMe and TuDiabetes have evolved out of the need to address the specific concerns of particular online communities, demonstrating the success that can be realised through tailoring a tool to the requirements of a chosen target audience. Conversely, a general tool such as Twitter has shown that it can not only be applied to a variety of different purposes but also has found a specific niche in disseminating public health alerts. The ability of these platforms to be customised for different

Figure 3 Word cloud representing the conditions included in the study populations. The size of each term is proportional to its representation in the review.



purposes is highly consistent with the principles underlying successful knowledge translation interventions. ³⁰

Most studies were descriptive, but our sample also included 48 RCTs. Nearly all of the trials evaluated the effectiveness of the discussion forums, leaving a research gap in the evaluation of the performance of other social media tools. Given the rapid proliferation of social media, a plethora of platforms are being used and an investigation of their benefits and harms is a logical progression of the research agenda. Similarly, the next steps in research could focus on isolating the effect of the social media tool, particularly as it relates to improved patient outcomes. All of the included RCTs evaluated a complex intervention, of which the social media tool was just one component. More focused efforts to determine whether social media has an impact on its own, or whether any observed effects are attributable to the intervention overall or to the non-social media components, would be a research priority. Similarly, a more in-depth examination of how the social media interventions are implemented, and specifically how and to what extent health or other professionals are involved, would contribute to a better understanding of their use. Further, additional research is needed to clarify whether the use of social media truly confers an advantage, or if the novelty of the medium is solely responsible for its use.³¹ The contrast between the statistical significance of the primary outcome in the RCTs and the positive conclusions reported suggests that issues such as selective outcome reporting (eg, choice of groups to compare), misrepresentation of conclusions (eg, focus on change over time within a group, rather than differences between groups) and a spin in reporting (eg, emphasis on a positive trend) may play a more substantial role in the promotion of social media use than actual effectiveness. The fact that most interventions were evaluated by their developers may have also influenced the positive conclusions reported.

Much of the research to this point has focused on measures of communication between peers or on social support, but our sample also included trials measuring the impact of social media on health behaviour and

Figure 4 Authors' conclusions by statistical significance and sample size among randomised controlled trials. Each bubble represents one study and its size is proportional to the number of individuals evaluated.




Table 5 Social media objectives by authors' conclusions (N=284)							
		Objective—n (%)					
Conclusions	Total—n (%)	Health literacy	Clinical decision making	Self-care	Patient safety	Others	
Total—n (%)		47 (16.6)	7 (2.5)	219 (77.1)	19 (6.7)	39 (13.7)	
Positive	186 (65.5)	28 (59.6)	6 (85.7)	149 (68.0)	14 (73.7)	21 (53.8)	
Neutral	65 (22.9)	12 (25.5)	1 (14.3)	47 (21.5)	1 (5.3)	13 (33.3)	
Negative	15 (5.3)	5 (10.6)	_	7 (3.2)	3 (15.8)	3 (7.7)	
Indeterminate	18 (6.3)	2 (4.3)	_	16 (7.3)	1 (5.3)	2 (5.1)	

status. With applications that directly target health outcomes, social media could present a cost-effective and wide-reaching modality for administering certain types of interventions. This could be particularly advantageous when logistics make arranging in-person appointments difficult, for example, in hard to reach populations, or when geography is an issue. These studies also suggest that social media has the potential to move beyond providing supportive online communities and could have widespread utility within the healthcare setting. However, these applications are dependent on further evidence of effectiveness.

LIMITATIONS

Social media is a relatively new concept that is continually undergoing transformation. As such, there is no universal definition, adding complexity to the process of determining study eligibility. The constantly changing nature of social media also proved challenging in defining the literature search, and the novelty of the topic made it difficult to keep the search updated due to a steady influx of new reports. However, as the focus of this scoping review was to identify broad categories of social media uses, the addition of studies published after the literature search would be unlikely to change the results.

While this scoping review focused on the peerreviewed literature to identify how social media is being used by patient and caregiver populations, it may not encompass all of the work that has been done in the area, or cover the extent of the impact that social media has had on healthcare. Much of the driving force behind the use of social media has come from outside the academic community; therefore, certain constructs such as the role that Facebook plays in advocacy and community, and patient empowerment resulting from the use of Twitter have not been captured. Additionally, certain movements that have shaped social media use in healthcare, such as the ePatient movement³² and Citizen Science,³³ were not included within the scope of our review. While we endeavoured to be as comprehensive as possible in covering the published literature, our included patient population may not be representative of people who use social media for health generally.

As our inclusion criteria were intentionally broad, we included a number of different study designs, encompassing both quantitative and qualitative research. While

this introduced challenges in addressing the nuances of each type of study, the end result is a comprehensive overview of the state of the literature. Further syntheses of the evidence in specific topics, clinical areas and populations will be able to provide more focus on some of these details.

CONCLUSIONS

This scoping review provides a map of the existing literature evaluating the use of social media in patient and caregiver populations. The available evidence is extensive, and most studies to date have been descriptive in nature. Given such widespread use of social media, evaluations of effectiveness are also needed. While positive conclusions are commonly reported, these may not be reflective of the actual findings.

Author affiliations

¹Alberta Research Centre for Health Evidence, Department of Pediatrics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

²Department of Pediatrics, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

³Faculty of Nursing, University of Alberta, Edmonton, Alberta, Canada ⁴School of Information Studies, Research Institute for Professional Practice, Learning and Education, Faculty of Education, Charles Sturt University, Wagga Wagga, NSW, Australia

Acknowledgements The authors would like to thank Walie Aktary for his assistance in data extraction, and Terry Klassen, Sharon Straus, Donna Angus and Liz Whamond for their contributions to the design of this review.

Contributors MPH, SDS, LMG and LH designed the study. MPH coordinated the project and is the guarantor. MPH, AC and JS screened articles and performed data extraction. AM contributed to the conception of the study and conducted the literature search. MPH, AC, JS and LH interpreted the data. MPH drafted and all authors critically reviewed the manuscript. All authors read and approved the manuscript.

Funding This study was supported by a Knowledge Synthesis Grant from the Canadian Institutes of Health Research, grant number 262961. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

REFERENCES

 Hawn C. Take two aspirin and tweet me in the morning: how Twitter, Facebook, and other social media are reshaping health care. Health Affair 2009;28:361–8.

- Allison M. Can web 2.0 reboot clinical trials? Nat Biotechnol 2009:27:895-902
- 3. Bonilla-Warford N. Many social media options exist for optometrists. Optometry 2010;81:613-14.
- Brownstein CA. Brownstein JS. Williams DS. et al. The power of social networking in medicine. Nat Biotechnol 2009;27:888-90.
- Eytan T, Benabio J, Golla V, et al. Social media and the health system. Perm J 2011;15:71-4.
- Spallek H, O'Donnell J, Clayton M, et al. Paradigm shift of annoying distraction: emerging implications of web 2.0 for clinical practice. Appl Clin Inf 2010:1:96-115.
- Morris K. Tweet, post, share—a new school of health communication. Lancet Infect Dis 2011;11:500-1.
- Villagran M. Methodological diversity to reach patients along the margins, in the shadows, and on the cutting edge. Patient Educ Counsel 2011;82:292-7.
- Timimi FK. Medicine, morality and health care social media. BMC Medicine 2012;10:83.
- Fox S. Health topics. American Life Project, 2011, http://pewinternet. org/Reports/2011/HealthTopics/Summary-of-Findings/Looking-forhealth-information.aspx (accessed 29 Jan 2013).
- Fox S, Duggan M. Health online 2013. Pew Research Center, 2013. http://www.pewinternet.org/~/media//Files/Reports/PIP HealthOnline. pdf (accessed 4 Apr 2013).
- Duggan M, Brenner J. The demographics of social media users— 2012. Pew Research Center, 2013. http://pewinternet.org/Reports/ 2013/Social-media-users.aspx (accessed 8 Mar 2013).
- Kaplan AM, Haenlein M. Users of the world, unite! The challenges and opportunities of social media. Bus Horiz 2010:53:59-68.
- Centers for Disease Control and Prevention. The health communicator's social media toolkit. http://www.cdc.gov/ healthcommunication/ToolsTemplates/SocialMediaToolkit BM.pdf (accessed 17 Sep 2012).
- Sarasohn-Kahn J. The wisdom of patients: health care meets online social media. California Health care Foundation, 2008. http://www. chcf.org/publications/2008/04/the-wisdom-of-patients-health-caremeets-online-social-media (accessed 17 Sep 2012).
- Crocco AG, Villasis-Keever M, Jadad AR. Two wrongs don't make a right: harm aggravated by inaccurate information on the Internet. Pediatrics 2002;109:522-3.
- Chou WS, Prestin A, Lyons C, et al. Web 2.0 for health promotion: reviewing the current evidence. Am J Public Health 2013;103:e9-e18.

- Crocco AG, Villasis-Keever M, Jadad AR, Analysis of cases of harm associated with use of health information on the internet. JAMA 2002:287:2869-71.
- Greysen SR, Kind T, Chretien KC. Online professionalism and the mirror of social media. J Gen Intern Med 2010:25:1227-9.
- 20 McBride DL. Risks and benefits of social media for children and adolescents. J Pediatr Nurs 2011;26:498-9.
- Collings SC, Fortune S, Currey N, et al. Media influences on suicidal behaviour: an interview study of young people in New Zealand. Auckland: The National Centre of Mental Health Research. Information and Workforce Development, 2011.
- Vickers H. Social networks and media coverage are blamed for series of teenage suicides in Russia. BMJ 2012;344:e3110.
- Strasburger VC, Donnerstein E. Children, adolescents, and the media in the 21st century. Adolesc Med 2000;11:51-68.
- Strasburger VC. Children, adolescents and the media: what we know, what we don't know and what we need to find out (quickly!). Arch Dis Child 2009;94:657.
- Hamm MP. Chisholm A. Shulhan J. et al. Use of social media by health care professionals and trainees: a scoping review. Acad Med 2013. In press.
- Coulter A, Ellins J. Patient-focused interventions: a review of the evidence. London: Health Foundation, 2006. http://www. pickereurope.org/Filestore/Publications/QEI Review AB.pdf (accessed 17 Sep 2012).
 Coulter A, Ellins J. Effectiveness of strategies for informing,
- educating, and involving patients. BMJ 2007;335:24-7.
- Buscemi N, Hartling L, Vandermeer B, et al. Single data extraction generated more errors than double data extraction in systematic reviews. J Clin Epidemiol 2006;59:697-703.
- Tricco A, Tetzlaff J, Pham B, et al. Non-Cochrane vs. Cochrane reviews were twice as likely to have positive conclusion statements: cross-sectional study. J Clin Epidemiol 2009;62:380-6.
- Graham ID, Logan J, Harrison MB, et al. Lost in translation: time for a map? J Contin Health Educ 2006;26:13-24.
- Rogers EM. Diffusion of Innovations. 5th edn. New York: The Free Press, 2003.
- 32. Debronkart D. How the e-patient community helped save my life: an essay by Dave deBronkart. BMJ 2013;346:f1990.
- Swan M. Crowdsourced health research studies: an important emerging complement to clinical trials in the public health research ecosystem. J Med Internet Res 2012;14:e46.

Appendix. Search strategy for Medline

<u>Database:</u> Medline via Ovid <1946 to Present>

Search Title: Social Media Scoping Review 1.4 all SD filters | Medline – 15Dec2011 – AM

Date Searched: 13 January 2012

<u>Limits:</u> Year of publication ≥2000; RCT/CCT, SR, observational, qualitative study filters applied

Results: 5,468 (Ovid duplicate removal function applied)

Internet and social media related MeSH [Medical Subject Headings]

- 1. exp Internet/
- 2. Electronic Mail/
- 3. Mass Media/td, ut
- 4. Hypermedia/
- 5. Online Systems/td, ut
- 6. Medical Informatics/
- 7. User-Computer Interface/
- 8. Computer-Assisted Instruction/
- 9. Computers/td, ut
- 10. Search Engine/
- 11. Computer Communication Networks/
- 12. Information Dissemination/
- 13. Therapy, Computer-Assisted/
- 14. "Marketing of Health Services"/
- 15. Social Marketing/
- 16. exp Social Environment/
- 17. **Internet.mp. and (or/12-16)** [Internet combined with broader social network/computer terms]

Internet and social medial related keywords

- 18. (digital adj5 platform*).mp.
- 19. (website* or web site* or webpage* or web page*).mp.
- 20. Googl*.mp.
- 21. Facebook*.mp.
- 22. YouTube.mp.
- 23. Second Life.mp.
- 24. PatientsLikeMe.mp.
- 25. WebMD.mp.
- 26. elluminate.mp.
- 27. flickr.mp.
- 28. moodle.mp.
- 29. picsearch.mp.
- 30. skype.mp.
- 31. ustream.mp.
- 32. zotero.mp.
- 33. ((e or electronic) adj3 newsletter*).mp.
- 34. (viral adj5 market*).mp.
- 35. (banner adj5 ad*).mp.
- 36. ("Web 2.0" or "Web 2").mp.
- 37. "Health 2.0".mp.

- 38. "Medicine 2.0".mp.
- 39. (Social adj3 network*).mp.
- 40. linkedin.mp.
- 41. blog*.mp.
- 42. wiki*.mp.
- 43. podcast*.mp.
- 44. really simple syndicat*.mp.
- 45. (rss adj3 (reader* or feed*)).mp.
- 46. (forum* adj3 (internet or web* or chat*)).mp.
- 47. content communit*.mp.
- 48. user generated content.mp.
- 49. microblog*.mp.
- 50. (twitter or tweet*).mp.
- 51. (("peer to peer" adj5 network*) or P2P).mp.
- 52. (social adj3 media*).mp.
- 53. i-phone*.mp.
- 54. myspace.mp.
- 55. smartphone*.mp.

56. or/1-11,17-55 [Internet/social media MeSH and keywords] (92,578)

Health care education/promotion terms

- 57. exp Health/
- 58. "Delivery of Health Care"/
- 59. health behavior/
- 60. exp Health Education/
- 61. exp Health Promotion/
- 62. Patient Care/
- 63. Patient Participation/
- 64. medical education/
- 65. ((patient* or physician* or nurse* or pharm* or "health care profession*") adj2 (teach* or train* or instruction* or intervention* or program* or inform* or educat* or outcome*)).mp.
- 66. **or/57-65** [Health promotion/health outcome terms] (**624,172**)
- 67. and/56,66 [social media + health promotion/outcome terms] (15,219)

Search filters to stream out non-research papers

RCT Filter

- 68. randomized controlled trial.pt.
- 69. controlled clinical trial.pt.
- 70. randomized.ab.
- 71. placebo.ab.
- 72. exp Clinical Trials as Topic/
- 73. randomly.ab.
- 74. trial.ti.
- 75. or/68-74
- 76. exp animals/ not humans.sh.
- 77. **75 not 76** [Cochrane RCT filter to max sensitivity and precision] (730,963)

SR Filter

- 78. meta analysis.mp,pt.
- 79. review.pt.
- 80. search*.tw.
- 81. or/78-80 [HIRU SR filter to balance sensitivity and specificity] (1,779,109)

Observational Study Filter

- 82. epidemiologic studies/
- 83. exp Case-Control Studies/
- 84. exp Cohort Studies/
- 85. case control.tw.
- 86. (cohort adj (study or studies)).tw.
- 87. cohort analy*.tw.
- 88. (follow up adj (study or studies)).tw.
- 89. (observational adj (study or studies)).tw.
- 90. longitudinal.tw.
- 91. retrospective.tw.
- 92. cross sectional.tw.
- 93. Cross-Sectional Studies/
- 94. **or/82-93** [SIGN observational study filter] (1,508,983)

Qualitative Research Filter

- 95. interview*.tw.
- 96. experience*.mp.
- 97. qualitative.tw.
- 98. **or/95-97** [HIRU qualitative study filter] (**756,921**)
- **99.** or/77,81,94,98 [combination of all search filters] (**4,143,826**)
- 100. and/67,99 [combination of social media terms + health ed terms + SD filters] (6,589)
- 101. limit 100 to humans (**6,234**)
- 102. limit 101 to yr="2000 -Current" (5,524)
- 103. remove duplicates from 102 (**5,468**)

Included Studies

- 1. Ache KA, Wallace LS. Human papillomavirus vaccination coverage on YouTube. American Journal of Preventive Medicine. 2008;35(4):389-92.
- 2. Adams SA. Sourcing the crowd for health services improvement: The reflexive patient and "share-your-experience" websites. Social Science & Medicine. 2011;72(7):1069-76.
- 3. Anttila M, Koivunen M, Valimaki M. Information technology-based standardized patient education in psychiatric inpatient care. Journal of Advanced Nursing. 2008;64(2):147-56.
- 4. Armstrong N, Powell J. Patient perspectives on health advice posted on Internet discussion boards: a qualitative study. Health Expectations. 2009;12(3):313-20.
- 5. Atkinson NL, Saperstein SL, Pleis J. Using the internet for health-related activities: findings from a national probability sample. Journal of Medical Internet Research. 2009;11(1):e4.
- 6. Baptist AP, Thompson M, Grossman KS, Mohammed L, Sy A, Sanders GM. Social media, text messaging, and email-preferences of asthma patients between 12 and 40 years old. Journal of Asthma. 2011;48(8):824-30.
- 7. Barker KK. Electronic support groups, patient-consumers, and medicalization: the case of contested illness. Journal of Health & Social Behavior. 2008;49(1):20-36.
- 8. Barr J, McLeod S. They never see how hard it is to be me: siblings' observations of strangers, peers and family. International Journal of Speechlanguage Pathology. 2010;12(2):162-71.
- 9. Barrera M, Jr.;, Glasgow RE, McKay HG, Boles SM, Feil EG. Do Internet-based support interventions change perceptions of social support?: An experimental trial of approaches for supporting diabetes self-management. American Journal of Community Psychology. 2002;30(5):637-54.
- 10. Bedgood R, Sadurski R, Schade RR. The use of the internet in data assimilation in rare diseases. Digestive Diseases & Sciences. 2007;52(2):307-12.
- 11. Bender JL, Jimenez-Marroquin MC, Jadad AR. Seeking support on facebook: a content analysis of breast cancer groups. Journal of Medical Internet Research. 2011;13(1):e16.
- 12. Bers MU, Gonzalez-Heydrich J, Raches D, DeMaso DR. Zora: a pilot virtual community in the pediatric dialysis unit. Studies in Health Technology & Informatics. 2001;84(Pt:1):1-4.
- 13. Bialous SA, Sarna L, Wells M, Elashoff D, Wewers ME, Froelicher ES. Characteristics of nurses who used the Internet-based nurses QuitNet for smoking cessation. Public Health Nursing. 2009;26(4):329-38.
- 14. Binks M, T. vM. Utilization patterns and user characteristics of an ad libitum Internet weight loss program. Journal of Medical Internet Research. 2010;12(1):e9.
- 15. Black AR, Peacock N. Pleasing the masses: messages for daily life management in African American women's popular media sources. American Journal of Public Health. 2011;101(1):144-50.
- 16. Black EW, Thompson LA, Saliba H, Dawson K, Black NM. An analysis of healthcare providers' online ratings. Informatics in Primary Care. 2009;17(4):249-53.
- 17. Bondy SJ, Bercovitz KL. Non-smoking worksites in the residential construction sector: using an online forum to study perspectives and practices. Tobacco Control. 2011;20(3):189-95.
- 18. Bonniface L, Green L. Finding a new kind of knowledge on the HeartNET website. Health Information & Libraries Journal. 2007;24:Suppl-76.
- 19. Booth AO, Nowson CA, Matters H. Evaluation of an interactive, Internet-based weight loss program: a pilot study. Health Education Research. 2008;23(3):371-81.

- 20. Bosak KA, Yates B, Pozehl B. Effects of an Internet physical activity intervention in adults with metabolic syndrome. Western Journal of Nursing Research. 2010;32(1):5-22.
- 21. Botella C, Etchemendy E, Castilla D, Banos RM, Garcia-Palacios A, Quero S, et al. An ehealth system for the elderly (Butler Project): a pilot study on acceptance and satisfaction. Cyberpsychology & Behavior. 2009;12(3):255-62.
- 22. Boyington AR, Dougherty MC, Liao YM. Analysis of interactive continence health information on the Web. Journal of Wound, Ostomy, & Continence Nursing. 2003;30(5):280-6.
- 23. Braner DA, Lai S, Hodo R, Ibsen LA, Bratton SL, Hollemon D, et al. Interactive Web sites for families and physicians of pediatric intensive care unit patients: a preliminary report. Pediatric Critical Care Medicine. 2004;5(5):434-9.
- 24. Broom A. Virtually he@lthy: the impact of internet use on disease experience and the doctor-patient relationship. Qualitative Health Research. 2005;15(3):325-45.
- 25. Bryce CL, Zickmund S, Hess R, McTigue KM, Olshansky E, Fitzgerald K, et al. Value versus user fees: perspectives of patients before and after using a web-based portal for management of diabetes. Telemedicine Journal & E-Health. 2008;14(10):1035-43.
- 26. Buchanan H, Coulson NS. Accessing dental anxiety online support groups: an exploratory qualitative study of motives and experiences. Patient Education & Counseling. 2007;66(3):263-9.
- 27. Bunde M, Suls J, Martin R, Barnett K. Hystersisters online: social support and social comparison among hysterectomy patients on the internet. Annals of Behavioral Medicine. 2006;31(3):271-8.
- 28. Chou MH, Lin MF, Hsu MC, Wang YH, Hu HF. Exploring the self-learning experiences of patients with depression participating in a multimedia education program. Journal of Nursing Research. 2004;12(4):297-306.
- 29. Civan A, Pratt W. Threading together patient expertise. AMIA. 2007; Annual: Symposium-4.
- 30. Coulson NS, Buchanan H, Aubeeluck A. Social support in cyberspace: a content analysis of communication within a Huntington's disease online support group. Patient Education & Counseling. 2007;68(2):173-8.
- 31. Cox MF, Scharer K, Clark AJ. Development of a Web-based program to improve communication about sex. CIN: Computers, Informatics, Nursing. 2009;27(1):18-25.
- 32. Cunningham JA, T.; vM, Fournier R. An online support group for problem drinkers: AlcoholHelpCenter.net. Patient Education & Counseling. 2008;70(2):193-8.
- 33. Curioso WH, Kurth AE. Access, use and perceptions regarding Internet, cell phones and PDAs as a means for health promotion for people living with HIV in Peru. BMC Medical Informatics & Decision Making. 2007;7:24.
- 34. Danaher BG, Smolkowski K, Seeley JR, Severson HH. Mediators of a successful web-based smokeless tobacco cessation program. Addiction. 2008;103(10):1706-12.
- 35. Davison KP, Pennebaker JW, Dickerson SS. Who talks? The social psychology of illness support groups. American Psychologist. 2000;55(2):205-17.
- 36. DeBar LL, Dickerson J, Clarke G, Stevens VJ, Ritenbaugh C, Aickin M. Using a website to build community and enhance outcomes in a group, multi-component intervention promoting healthy diet and exercise in adolescents. Journal of Pediatric Psychology. 2009;34(5):539-50.
- 37. Dickerson SS. Technology--patient interactions: Internet use for gaining a healthy context for living with an implantable cardioverter defibrillator. Heart & Lung. 2005;34(3):157-68.

- 38. Dickerson SS, Flaig DM, Kennedy MC. Therapeutic connection: help seeking on the Internet for persons with implantable cardioverter defibrillators. Heart & Lung. 2000;29(4):248-55.
- 39. Donelle L, Hoffman-Goetz L. Health literacy and online health discussions of North American Black women. Women & Health. 2008;47(4):71-90.
- 40. Ewing LJ, Long K, Rotondi A, Howe C, Bill L, Marsland AL. Brief report: A pilot study of a web-based resource for families of children with cancer. Journal of Pediatric Psychology. 2009;34(5):523-9.
- 41. Fehring RJ, Schneider M, Raviele K. Pilot evaluation of an internet-based natural family planning education and service program. JOGNN Journal of Obstetric, Gynecologic, & Neonatal Nursing. 2011;40(3):281-91.
- 42. Feil EG, Glasgow RE, Boles S, McKay HG. Who participates in Internet-based self-management programs? A study among novice computer users in a primary care setting. Diabetes Educator. 2000;26(5):806-11.
- 43. Ferney SL, Marshall AL, Eakin EG, Owen N. Randomized trial of a neighborhood environment-focused physical activity website intervention. Preventive Medicine. 2009;48(2):144-50.
- 44. Flicker S, Goldberg E, Read S, Veinot T, McClelland A, Saulnier P, et al. HIV-positive youth's perspectives on the Internet and e-health. Journal of Medical Internet Research. 2004;6(3):e32.
- 45. Fogel J, Albert SM, Schnabel F, Ditkoff BA, Neugut AI. Internet use and social support in women with breast cancer. Health Psychology. 2002;21(4):398-404.
- 46. Fox NJ, Ward KJ, O'Rourke AJ. The 'expert patient': empowerment or medical dominance? The case of weight loss, pharmaceutical drugs and the Internet. Social Science & Medicine. 2005;60(6):1299-309.
- 47. Freeman E, Barker C, Pistrang N. Outcome of an online mutual support group for college students with psychological problems. Cyberpsychology & Behavior. 2008;11(5):591-3.
- 48. Frisby G, Bessell TL, Borland R, Anderson JN. Smoking cessation and the Internet: a qualitative method examining online consumer behavior. Journal of Medical Internet Research. 2002;4(2):E8-Nov.
- 49. Frost J, Okun S, Vaughan T, Heywood J, Wicks P. Patient-reported outcomes as a source of evidence in off-label prescribing: analysis of data from PatientsLikeMe. Journal of Medical Internet Research. 2011;13(1):e6.
- 50. Frost JH, Massagli MP, Wicks P, Heywood J. How the Social Web Supports patient experimentation with a new therapy: The demand for patient-controlled and patient-centered informatics. AMIA. 2008;Annual:Symposium-21.
- 51. Frost JH, Massagli MP. Social uses of personal health information within PatientsLikeMe, an online patient community: what can happen when patients have access to one another's data. Journal of Medical Internet Research. 2008;10(3):e15.
- 52. Gavin J, Rodham K, Poyer H. The presentation of "pro-anorexia" in online group interactions. Qualitative Health Research. 2008;18(3):325-33.
- 53. Ghahari S, Packer TL, Passmore AE. Development, standardisation and pilot testing of an online fatigue self-management program. Disability & Rehabilitation. 2009;31(21):1762-72.
- 54. Giles DC, Newbold J. Self- and other-diagnosis in user-led mental health online communities. Qualitative Health Research. 2011;21(3):419-28.

- 55. Glasgow RE, Boles SM, McKay HG, Feil EG, Barrera M, Jr. The D-Net diabetes self-management program: long-term implementation, outcomes, and generalization results. Preventive Medicine. 2003;36(4):410-9.
- 56. Gold BC, Burke S, Pintauro S, Buzzell P, Harvey-Berino J. Weight loss on the web: A pilot study comparing a structured behavioral intervention to a commercial program. Obesity. 2007;15(1):155-64.
- 57. Gow RW, Trace SE, Mazzeo SE. Preventing weight gain in first year college students: an online intervention to prevent the "freshman fifteen". Eating Behaviors. 2010;11(1):33-9.
- 58. Graham AL, Cobb NK, Raymond L, Sill S, Young J. Effectiveness of an internet-based worksite smoking cessation intervention at 12 months. Journal of Occupational & Environmental Medicine. 2007;49(8):821-8.
- 59. Greene JA, Choudhry NK, Kilabuk E, Shrank WH. Online social networking by patients with diabetes: a qualitative evaluation of communication with Facebook. Journal of General Internal Medicine. 2011;26(3):287-92.
- 60. Hagan JC, III;, Kutryb MJ. Cataract and intraocular implant surgery concerns and comments posted at two internet eye care forums. Missouri Medicine. 2009;106(1):78-82.
- 61. Hall W, Irvine V. E-communication among mothers of infants and toddlers in a community-based cohort: a content analysis. Journal of Advanced Nursing. 2009;65(1):175-83.
- 62. Halliday L, Boughton M. Premature menopause: exploring the experience through online communication. Nursing & Health Sciences. 2009;11(1):17-22.
- 63. Hawkins RP, Pingree S, Shaw B, Serlin RC, Swoboda C, Han JY, et al. Mediating processes of two communication interventions for breast cancer patients. Patient Education & Counseling. 2010;81:Suppl-53.
- 64. Hess RF, Weinland JA, Beebe K. "I am not alone": a survey of women with peripartum cardiomyopathy and their participation in an online support group. CIN: Computers, Informatics, Nursing. 2010;28(4):215-21.
- 65. Himmel W, Meyer J, Kochen MM, Michelmann HW. Information needs and visitors' experience of an Internet expert forum on infertility. Journal of Medical Internet Research. 2005;7(2):e20.
- 66. Huber J, Ihrig A, Peters T, Huber CG, Kessler A, Hadaschik B, et al. Decision-making in localized prostate cancer: lessons learned from an online support group. BJU International. 2011;107(10):1570-5.
- 67. Hudson DB, Campbell-Grossman C, Keating-Lefler R, Carraher S, Gehle J, Heusinkvelt S. Online support for single, low-income, African American mothers. MCN, American Journal of Maternal Child Nursing. 2009;34(6):350-5.
- 68. Hurling R, Catt M, Boni MD, Fairley BW, Hurst T, Murray P, et al. Using internet and mobile phone technology to deliver an automated physical activity program: randomized controlled trial. Journal of Medical Internet Research. 2007;9(2):e7.
- 69. Hwang KO, Ottenbacher AJ, Green AP, Cannon-Diehl MR, Richardson O, Bernstam EV, et al. Social support in an Internet weight loss community. International Journal of Medical Informatics. 2010;79(1):5-13.
- 70. Im EO, Lee SH, Chee W. "Being conditioned, yet becoming strong": Asian American women in menopausal transition. Journal of Transcultural Nursing. 2011;22(3):290-9.
- 71. Im EO, Lee SH, Chee W. Black women in menopausal transition. JOGNN Journal of Obstetric, Gynecologic, & Neonatal Nursing. 2010;39(4):435-43.

- 72. Im EO, Lee B, Chee W. Shielded from the real world: perspectives on Internet cancer support groups by Asian Americans. Cancer Nursing. 2010;33(3):E10-E20.
- 73. Im EO, Lee B, Chee W, Dormire S, Brown A. A national multiethnic online forum study on menopausal symptom experience. Nursing Research. 2010;59(1):26-33.
- 74. Im EO, Lim HJ, Lee SH, Dormire S, Chee W, Kresta K. Menopausal symptom experience of Hispanic midlife women in the United States. Health Care for Women International. 2009;30(10):919-34.
- 75. Im EO, Chee W, Lim HJ, Liu WM. An online forum exploring needs for help of patients with cancer: gender and ethnic differences. Oncology Nursing Forum. 2008;35(4):653-60.
- 76. Im EO, Chee W, Lim HJ, Liu Y, Kim HK. Midlife women's attitudes toward physical activity. JOGNN Journal of Obstetric, Gynecologic, & Neonatal Nursing. 2008;37(2):203-13.
- 77. Im EO, Guevara E, Chee W. The pain experience of Hispanic patients with cancer in the United States. Oncology Nursing Forum. 2007;34(4):861-8.
- 78. Im EO, Chee W, Tsai HM, Lin LC, Cheng CY. Internet cancer support groups: a feminist analysis. Cancer Nursing. 2005;28(1):1-7.
- 79. Johnson F, Wardle J. The association between weight loss and engagement with a web-based food and exercise diary in a commercial weight loss programme: a retrospective analysis. International Journal of Behavioral Nutrition & Physical Activity. 2011;8:83.
- 80. Kamel Boulos MN, Toth-Cohen S. The University of Plymouth Sexual Health SIM experience in Second Life: evaluation and reflections after 1 year. Health Information & Libraries Journal. 2009;26(4):279-88.
- 81. Kelders SM, Van Gemert-Pijnen JE, Werkman A, Seydel ER. Evaluation of a web-based lifestyle coach designed to maintain a healthy bodyweight. Journal of Telemedicine & Telecare. 2010;16(1):3-7.
- 82. Kennedy A, Rogers A, Sanders C, Gately C, Lee V. Creating 'good' self-managers?: facilitating and governing an online self care skills training course. BMC Health Services Research. 2009:9:93.
- 83. Kerr C, Murray E, Noble L, Morris R, Bottomley C, Stevenson F, et al. The potential of Web-based interventions for heart disease self-management: a mixed methods investigation. Journal of Medical Internet Research. 2010;12(4):e56.
- 84. Kishimoto K, Fukushima N. Use of anonymous Web communities and websites by medical consumers in Japan to research drug information. Yakugaku Zasshi Journal of the Pharmaceutical Society of Japan. 2011;131(5):685-95.
- 85. Kosma M, Cardinal BJ, McCubbin JA. A pilot study of a web-based physical activity motivational program for adults with physical disabilities. Disability & Rehabilitation. 2005;27(23):1435-42.
- 86. Kouri P, Turunen H, Tossavainen K, Saarikoski S. Pregnant families' discussions on the Net-from virtual connections toward real-life community. Journal of Midwifery & Women's Health. 2006;51(4):279-83.
- 87. Lagu T, Hannon NS, Rothberg MB, Lindenauer PK. Patients' evaluations of health care providers in the era of social networking: an analysis of physician-rating websites. Journal of General Internal Medicine. 2010;25(9):942-6.
- 88. Laurent MR, Vickers TJ. Seeking health information online: does Wikipedia matter? Journal of the American Medical Informatics Association. 2009;16(4):471-9.

- 89. Lee SY, Hawkins R. Why do patients seek an alternative channel? The effects of unmet needs on patients' health-related Internet use. Journal of Health Communication. 2010;15(2):152-66.
- 90. Lindsay S, Smith S, Bellaby P, Baker R. The health impact of an online heart disease support group: a comparison of moderated versus unmoderated support. Health Education Research. 2009;24(4):646-54.
- 91. Linkletter M, Gordon K, Dooley J. The choking game and YouTube: a dangerous combination. Clinical Pediatrics. 2010;49(3):274-9.
- 92. Linssen C, Schook RM, The AM, Lammers E, Festen J, Postmus PE. A web site on lung cancer: who are the users and what are they looking for? Journal of Thoracic Oncology: Official Publication of the International Association for the Study of Lung Cancer. 2007;2(9):813-8.
- 93. Ljotsson B, Falk L, Vesterlund AW, Hedman E, Lindfors P, Ruck C, et al. Internet-delivered exposure and mindfulness based therapy for irritable bowel syndrome--a randomized controlled trial. Behaviour Research & Therapy. 2010;48(6):531-9.
- 94. Lorig KR, Ritter PL, Dost A, Plant K, Laurent DD, McNeil I. The Expert Patients Programme online, a 1-year study of an Internet-based self-management programme for people with long-term conditions. Chronic Illness. 2008;4(4):247-56.
- 95. Lorig KR, Ritter PL, Laurent DD, Plant K. Internet-based chronic disease self-management: a randomized trial. [Erratum appears in Med Care. 2007 Mar;45(3):276]. Medical Care. 2006;44(11):964-71.
- 96. Lou CH, Zhao Q, Gao ES, Shah IH. Can the Internet be used effectively to provide sex education to young people in China? Journal of Adolescent Health. 2006;39(5):720-8.
- 97. Lu HY, Shaw BR, Gustafson DH. Online health consultation: examining uses of an interactive cancer communication tool by low-income women with breast cancer. International Journal of Medical Informatics. 2011;80(7):518-28.
- 98. Lynch M. Healthy habits or damaging diets: an exploratory study of a food blogging community. Ecology of Food & Nutrition. 2010;49(4):316-35.
- 99. McKay HG, King D, Eakin EG, Seeley JR, Glasgow RE. The diabetes network internet-based physical activity intervention: a randomized pilot study. Diabetes Care. 2001;24(8):1328-34.
- 100. Meglic M, Furlan M, Kuzmanic M, Kozel D, Baraga D, Kuhar I, et al. Feasibility of an eHealth service to support collaborative depression care: results of a pilot study. Journal of Medical Internet Research. 2010;12(5):e63.
- 101. Miller EA, Pole A, Bateman C. Variation in health blog features and elements by gender, occupation, and perspective. Journal of Health Communication. 2011;16(7):726-49.
- 102. Moncrieff J, Cohen D, Mason JP. The subjective experience of taking antipsychotic medication: a content analysis of Internet data. Acta Psychiatrica Scandinavica. 2009;120(2):102-11.
- 103. Moreno MA, Parks M, Richardson LP. What are adolescents showing the world about their health risk behaviors on MySpace? Medgenmed [Computer File]: Medscape General Medicine. 2007;9(4):9.
- 104. Morgan PJ, Lubans DR, Collins CE, Warren JM, Callister R. 12-month outcomes and process evaluation of the SHED-IT RCT: an internet-based weight loss program targeting men. Obesity. 2011;19(1):142-51.

- 105. Morita T, Narimatsu H, Matsumura T, Kodama Y, Hori A, Kishi Y, et al. A study of cancer information for cancer patients on the internet. International Journal of Clinical Oncology. 2007;12(6):440-7.
- 106. Motl RW, Dlugonski D, Wojcicki TR, McAuley E, Mohr DC. Internet intervention for increasing physical activity in persons with multiple sclerosis. Multiple Sclerosis. 2011;17(1):116-28.
- 107. Munoz RF, Barrera AZ, Delucchi K, Penilla C, Torres LD, Perez-Stable EJ. International Spanish/English Internet smoking cessation trial yields 20% abstinence rates at 1 year. Nicotine & Tobacco Research. 2009;11(9):1025-34.
- 108. Muramoto ML, Wassum K, Connolly T, Matthews E, Floden L. Helpers program: A pilot test of brief tobacco intervention training in three corporations. American Journal of Preventive Medicine. 2010;38(3:Suppl):Suppl-26.
- 109. Nahm ES, Barker B, Resnick B, Covington B, Magaziner J, Brennan PF. Effects of a social cognitive theory-based hip fracture prevention web site for older adults. CIN: Computers, Informatics, Nursing. 2010;28(6):371-9.
- 110. Nakayama K, Nishio A, Yokoyama Y, Setoyama Y, Togari T, Yonekura Y. When and why do people post questions about health and illness on Web 2.0-based Q&A sites in Japan. Studies in Health Technology & Informatics. 2009;146:731.
- 111. Neal DM, McKenzie PJ. Putting the pieces together: endometriosis blogs, cognitive authority, and collaborative information behavior. Journal of the Medical Library Association. 2011;99(2):127-34.
- 112. Neve MJ, Collins CE, Morgan PJ. Dropout, nonusage attrition, and pretreatment predictors of nonusage attrition in a commercial Web-based weight loss program. Journal of Medical Internet Research. 2010;12(4):e69.
- 113. Nguyen HQ, Donesky-Cuenco D, Wolpin S, Reinke LF, Benditt JO, Paul SM, et al. Randomized controlled trial of an internet-based versus face-to-face dyspnea self-management program for patients with chronic obstructive pulmonary disease: pilot study. Journal of Medical Internet Research. 2008;10(2):e9.
- 114. Nicholas DB, Darch J, McNeill T, Brister L, O'Leary K, Berlin D, et al. Perceptions of online support for hospitalized children and adolescents. Social Work in Health Care. 2007;44(3):205-23.
- 115. Noar SM, Clark A, Cole C, Lustria ML. Review of interactive safer sex Web sites: practice and potential. [Review] [44 refs]. Health Communication. 2006;20(3):233-41.
- 116. Nordqvist C, Hanberger L, Timpka T, Nordfeldt S. Health professionals' attitudes towards using a Web 2.0 portal for child and adolescent diabetes care: qualitative study. Journal of Medical Internet Research. 2009;11(2):e12.
- 117. O'Dea B, Campbell A. Healthy connections: online social networks and their potential for peer support. Studies in Health Technology & Informatics. 2011;168:133-40.
- 118. O'Grady LA. Consumer e-health education in HIV/AIDS: a pilot study of a web-based video workshop. BMC Medical Informatics & Decision Making. 2006;6:10.
- 119. Orizio G, Schulz P, Gasparotti C, Caimi L, Gelatti U. The world of e-patients: A content analysis of online social networks focusing on diseases. Telemedicine Journal & E-Health. 2010;16(10):1060-6.
- 120. Owen JE, Klapow JC, Roth DL, Shuster JL, Jr.;, Bellis J, Meredith R, et al. Randomized pilot of a self-guided internet coping group for women with early-stage breast cancer. Annals of Behavioral Medicine. 2005;30(1):54-64.

- 121. Papadaki A, Scott JA. The Mediterranean eating in Scotland experience project: evaluation of an Internet-based intervention promoting the Mediterranean diet. British Journal of Nutrition. 2005;94(2):290-8.
- 122. Patten CA, Croghan IT, Meis TM, Decker PA, Pingree S, Colligan RC, et al. Randomized clinical trial of an Internet-based versus brief office intervention for adolescent smoking cessation. Patient Education & Counseling. 2006;64(1-3):249-58.
- 123. Pletneva N, Cruchet S, Simonet MA, Kajiwara M, Boyer C. Results of the 10 HON survey on health and medical internet use. Studies in Health Technology & Informatics. 2011;169:73-7.
- 124. Pullen CH, Hageman PA, Boeckner L, Walker SN, Oberdorfer MK. Feasibility of Internet-delivered weight loss interventions among rural women ages 50-69. Journal of Geriatric Physical Therapy. 2008;31(3):105-12.
- 125. Rada R. Entry requirements and membership homogeneity in online patient groups. Medical Informatics & the Internet in Medicine. 2007;32(3):215-23.
- 126. Rezailashkajani M, Roshandel D, Ansari S, Zali MR. A web-based patient education system and self-help group in Persian language for inflammatory bowel disease patients. International Journal of Medical Informatics. 2008;77(2):122-8.
- 127. Richardson CR, Buis LR, Janney AW, Goodrich DE, Sen A, Hess ML, et al. An online community improves adherence in an internet-mediated walking program. Part 1: results of a randomized controlled trial. Journal of Medical Internet Research. 2010;12(4):e71.
- 128. Rotondi AJ, Anderson CM, Haas GL, Eack SM, Spring MB, Ganguli R, et al. Web-based psychoeducational intervention for persons with schizophrenia and their supporters: one-year outcomes. Psychiatric Services. 2010;61(11):1099-105.
- 129. Rydell SA, French SA, Fulkerson JA, Neumark-Sztainer D, Gerlach AF, Story M, et al. Use of a Web-based component of a nutrition and physical activity behavioral intervention with Girl Scouts. Journal of the American Dietetic Association. 2005;105(9):1447-50.
- 130. Sajadi KP, Goldman HB. Social networks lack useful content for incontinence. Urology. 2011;78(4):764-7.
- 131. Saver BG, Gustafson D, Taylor TR, Hawkins RP, Woods NF, Dinauer S, et al. A tale of two studies: the importance of setting, subjects and context in two randomized, controlled trials of a web-based decision support for perimenopausal and postmenopausal health decisions. Patient Education & Counseling. 2007;66(2):211-22.
- 132. Savige GS. E-learning: a nutritionally ripe environment. [Review] [15 refs]. Food & Nutrition Bulletin. 2005;26(2:Suppl 2):Suppl-4.
- 133. Schembri G, Schober P. The Internet as a diagnostic aid: the patients' perspective. International Journal of STD & AIDS. 2009;20(4):231-3.
- 134. Schroder S, Zollner YF, Schaefer M. Drug related problems with Antiparkinsonian agents: consumer Internet reports versus published data. Pharmacoepidemiology & Drug Safety. 2007;16(10):1161-6.
- 135. Schultz PN, Stava C, Beck ML, Vassilopoulou-Sellin R. Internet message board use by patients with cancer and their families. Clinical Journal of Oncology Nursing. 2003;7(6):663-7.
- 136. Schulz PJ, Rubinelli S, Mariotti G, Keller N. Meeting the ranging of informational needs of chronic low back pain sufferers: conceptual design and rationale of the interactive website ONESELF. Disability & Rehabilitation. 2009;31(25):2118-24.

- 137. Schulz PJ, Rubinell S, Hartung U. An internet-based approach to enhance self-management of chronic low back pain in the italian-speaking population of Switzerland: results from a pilot study. International Journal of Public Health. 2007;52(5):286-94.
- 138. Schwinn TM, Schinke SP, Di Noia J. Preventing drug abuse among adolescent girls: outcome data from an internet-based intervention. Prevention Science. 2010;11(1):24-32.
- 139. Severson HH, Gordon JS, Danaher BG, Akers L. ChewFree.com: evaluation of a Webbased cessation program for smokeless tobacco users. Nicotine & Tobacco Research. 2008;10(2):381-91.
- 140. Shigaki CL, Smarr KL, Yang G, Donovan-Hanson K, Siva C, Johnson RA, et al. Social interactions in an online self-management program for rheumatoid arthritis. Chronic Illness. 2008;4(4):239-46.
- 141. Simon C, Schramm S. Cancer and the computerized family: towards a clinical ethics of "indirect" Internet use. Medicine, Health Care & Philosophy. 2008;11(3):337-41.
- 142. Simon GE, Ludman EJ, Goodale LC, Dykstra DM, Stone E, Cutsogeorge D, et al. An online recovery plan program: can peer coaching increase participation? Psychiatric Services. 2011;62(6):666-9.
- 143. Simovska V. The changing meanings of participation in school-based health education and health promotion: the participants' voices. Health Education Research. 2007;22(6):864-78.
- 144. Smarr KL, Musser DR, Shigaki CL, Johnson R, Hanson KD, Siva C. Online self-management in rheumatoid arthritis: a patient-centered model application. Telemedicine Journal & E-Health. 2011;17(2):104-10.
- 145. Smerecnik C, Schaalma H, Gerjo K, Meijer S, Poelman J. An exploratory study of Muslim adolescents' views on sexuality: Implications for sex education and prevention. BMC Public Health. 2010;10:533.
- 146. Smith L, Weinert C. Telecommunication support for rural women with diabetes. Diabetes Educator. 2000;26(4):645-55.
- 147. Sood A, Sarangi S, Pandey A, Murugiah K. YouTube as a source of information on kidney stone disease. Urology. 2011;77(3):558-62.
- 148. Sparud-Lundin C, Ranerup A, Berg M. Internet use, needs and expectations of web-based information and communication in childbearing women with type 1 diabetes. BMC Medical Informatics & Decision Making. 2011;11:49.
- 149. Spittaels H, I.; DB, Vandelanotte C. Evaluation of a website-delivered computer-tailored intervention for increasing physical activity in the general population. Preventive Medicine. 2007;44(3):209-17.
- 150. Springer A, Reck CA, Huber C, Horcher E. Online hypospadias support group data analysis. Journal of Pediatric Surgery. 2011;46(3):520-4.
- 151. Steinberg PL, Wason S, Stern JM, Deters L, Kowal B, Seigne J. YouTube as source of prostate cancer information. Urology. 2010;75(3):619-22.
- 152. Stevens VJ, Funk KL, Brantley PJ, Erlinger TP, Myers VH, Champagne CM, et al. Design and implementation of an interactive website to support long-term maintenance of weight loss. Journal of Medical Internet Research. 2008;10(1):e1.
- 153. Stinson J, McGrath P, Hodnett E, Feldman B, Duffy C, Huber A, et al. Usability testing of an online self-management program for adolescents with juvenile idiopathic arthritis. Journal of Medical Internet Research. 2010;12(3):e30.

- 154. Sublet V, Spring C, Howard J, National Institute for Occupational Safety Health. Does social media improve communication? Evaluating the NIOSH science blog. American Journal of Industrial Medicine. 2011;54(5):384-94.
- 155. Tate DF, Wing RR, Winett RA. Using Internet technology to deliver a behavioral weight loss program. JAMA. 2001;285(9):1172-7.
- 156. Thoren E, Svensson M, Tornqvist A, Andersson G, Carlbring P, Lunner T. Rehabilitative online education versus internet discussion group for hearing aid users: a randomized controlled trial. Journal of the American Academy of Audiology. 2011;22(5):274-85.
- 157. van den Berg MH, Ronday HK, Peeters AJ, Voogt-van der Harst EM, Munneke M, Breedveld FC, et al. Engagement and satisfaction with an Internet-based physical activity intervention in patients with rheumatoid arthritis. Rheumatology. 2007;46(3):545-52.
- 158. van den Berg MH, Ronday HK, Peeters AJ, S.; lC, van der Giesen FJ, Breedveld FC, et al. Using internet technology to deliver a home-based physical activity intervention for patients with rheumatoid arthritis: A randomized controlled trial. Arthritis Care & Research. 2006;55(6):935-45.
- 159. van dV, Drossaert CH, Taal E, van de Laar MA. Patient preferences for a hospital-based rheumatology Interactive Health Communication Application and factors associated with these preferences. Rheumatology. 2011;50(9):1618-26.
- 160. van Uden-Kraan CF, Drossaert CH, Taal E, Seydel ER, van de Laar MA. Patient-initiated online support groups: motives for initiation, extent of success and success factors. Journal of Telemedicine & Telecare. 2010;16(1):30-4.
- 161. van Uden-Kraan CF, Drossaert CH, Taal E, Smit WM, Moens HJ, Siesling S, et al. Health-related Internet use by patients with somatic diseases: frequency of use and characteristics of users. Informatics for health & social care. 2009;34(1):18-29.
- 162. van Uden-Kraan CF, Drossaert CH, Taal E, Seydel ER, van de Laar MA. Participation in online patient support groups endorses patients' empowerment. Patient Education & Counseling. 2009;74(1):61-9.
- 163. van Uden-Kraan CF, Drossaert CH, Taal E, Seydel ER, van de Laar MA. Self-reported differences in empowerment between lurkers and posters in online patient support groups. Journal of Medical Internet Research. 2008;10(2):e18.
- 164. van Uden-Kraan CF, Drossaert CH, Taal E, Shaw BR, Seydel ER, van de Laar MA. Empowering processes and outcomes of participation in online support groups for patients with breast cancer, arthritis, or fibromyalgia. Qualitative Health Research. 2008;18(3):405-17.
- 165. Verheijden M, Bakx JC, Akkermans R, van den HH, Godwin NM, Rosser W, et al. Webbased targeted nutrition counselling and social support for patients at increased cardiovascular risk in general practice: randomized controlled trial. Journal of Medical Internet Research. 2004;6(4):e44.
- 166. Ware LJ, Hurling R, Bataveljic O, Fairley BW, Hurst TL, Murray P, et al. Rates and determinants of uptake and use of an internet physical activity and weight management program in office and manufacturing work sites in England: cohort study. Journal of Medical Internet Research. 2008;10(4):e56.
- 167. Webber KH, Tate DF, Quintiliani LM. Motivational interviewing in internet groups: a pilot study for weight loss. Journal of the American Dietetic Association. 2008;108(6):1029-32.

- 168. Weitzman ER, Adida B, Kelemen S, Mandl KD. Sharing data for public health research by members of an international online diabetes social network. PLoS ONE [Electronic Resource]. 2011;6(4):e19256.
- 169. Whittemore R, Grey M, Lindemann E, Ambrosino J, Jaser S. Development of an Internet coping skills training program for teenagers with type 1 diabetes. CIN: Computers, Informatics, Nursing. 2010;28(2):103-11.
- 170. Wicks P, Massagli M, Frost J, Brownstein C, Okun S, Vaughan T, et al. Sharing health data for better outcomes on PatientsLikeMe. Journal of Medical Internet Research. 2010;12(2):e19.
- 171. Wjst M. When air is rare: behind the scenes of an asthma web site. Journal of Asthma. 2001;38(5):399-404.
- 172. Yu J, Taverner N, Madden K. Young people's views on sharing health-related stories on the Internet. Health & Social Care in the Community. 2011;19(3):326-34.
- 173. Zabinski MF, Wilfley DE, Pung MA, Winzelberg AJ, Eldredge K, Taylor CB. An interactive internet-based intervention for women at risk of eating disorders: a pilot study. International Journal of Eating Disorders. 2001;30(2):129-37.
- 174. Zabinski MF, Pung MA, Wilfley DE, Eppstein DL, Winzelberg AJ, Celio A, et al. Reducing risk factors for eating disorders: targeting at-risk women with a computerized psychoeducational program. International Journal of Eating Disorders. 2001;29(4):401-8.
- 175. Jin S-AA. Leveraging Avatars in 3D Virtual Environments ("Second Life") for Interactive Learning: The Moderating Role of the Behavioral Activation System "vs." Behavioral Inhibition System and the Mediating Role of Enjoyment. Interactive Learning Environments. 2011;19(5):467-86.
- 176. Selkie EM, Benson M, Moreno M. Adolescents' Views regarding Uses of Social Networking Websites and Text Messaging for Adolescent Sexual Health Education. American Journal of Health Education. 2011;42(4):205-12.
- 177. Lynch M. Social Media in Health Research: An Example from Childcare Provider Message Boards. Contemporary Issues in Early Childhood. 2011;12(2):187-90.
- 178. Craig Rushing SN. Use of Media Technologies by Native American Teens and Young Adults: Evaluating Their Utility for Designing Culturally-Appropriate Sexual Health Interventions Targeting Native Youth in the Pacific Northwest. ProQuest LLC PhD Dissertation, Portland State University. 2010.
- 179. Osvaldsson K. Bullying in Context: Stories of Bullying on an Internet Discussion Board. Children and Society. 2011;25(4):317-27.
- 180. Ferney SL, Marshall AL. Website Physical Activity Interventions: Preferences of Potential Users. Health Education Research. 2006;21(4):560-6.
- 181. Caines MJ. Challenges in Fibromyalgia Management: A Study of Anxiety, Depression, and Motivation Using Distance Learning and Social Networking. Online Submission PhD Dissertation, A T Still University. 2010.
- 182. Greidanus E, Everall RD. Helper Therapy in an Online Suicide Prevention Community. British Journal of Guidance and Counselling. 2010;38(2):191-204.
- 183. Webber KH, Tate DF, Ward DS, Bowling JM. Motivation and Its Relationship to Adherence to Self-Monitoring and Weight Loss in a 16-Week Internet Behavioral Weight Loss Intervention. Journal of Nutrition Education and Behavior. 2010;42(3):161-7.
- 184. Richards D. Features and Benefits of Online Counselling: Trinity College Online Mental Health Community. British Journal of Guidance and Counselling. 2009;37(3):231-42.

- 185. Brouwer W, Oenema A, Crutzen R, de Nooijer J, de Vries NK, Brug J. What Makes People Decide to Visit and Use an Internet-Delivered Behavior-Change Intervention?: A Qualitative Study among Adults. Health Education. 2009;109(6):460-73.
- 186. Santor DA, Poulin C, LeBlanc JC, Kusumakar V. Online Health Promotion, Early Identification of Difficulties, and Help Seeking in Young People. Journal of the American Academy of Child and Adolescent Psychiatry. 2007;46(1):50.
- 187. Stamelou M, Edwards MJ, Espay AJ, Fung VS, Hallett M, Lang AE, et al. Movement disorders on YouTube--caveat spectator. N Engl J Med. 2011;365(12):1160-1. Epub 2011/10/14.
- 188. Yanagisawa M, Rhodes M, Zimmern P. Mesh social networking: a patient-driven process. BJU Int. 2011;108(10):1539-41. Epub 2011/10/11.
- 189. Gamage DG, Fuller CA, Cummings R, Tomnay JE, Chung M, Chen M, et al. Advertising sexual health services that provide sexually transmissible infection screening for rural young people what works and what doesn't. Sex Health. 2011;8(3):407-11. Epub 2011/08/20.
- 190. Cowie GA, Hill S, Robinson P. Using an online service for breastfeeding support: what mothers want to discuss. Health Promot J Austr. 2011;22(2):113-8. Epub 2011/08/09.
- 191. Adam BD, Murray J, Ross S, Oliver J, Lincoln SG, Rynard V. hivstigma.com, an innovative web-supported stigma reduction intervention for gay and bisexual men. Health Educ Res. 2011;26(5):795-807. Epub 2011/01/19.
- 192. Holthe T, Walderhaug S. Older people with and without dementia participating in the development of an individual plan with digital calendar and message board. Journal of Assistive Technologies. 2010;4(2):15-25.
- 193. Viiala M, Pietila A, Kankkunen P. Peer support reinforcing health: discussion board messages for people suffering from eating disorders in the Internet [Finnish]. Nursing Evidence / Tutkiva Hoitotyo. 2010;8(3):12-9.
- 194. Weinert C. Social support in cyberspace for women with chronic illness. Rehabilitation Nursing. 2000;25(4):129.
- 195. Ryan S. Trust and participation in online usenet self-help communities. International Journal of Self Help & Self Care. 2006;5(1):43-72.
- 196. Anderson K, Weinert C. Women to women 2: a computer-based support and educational intervention... proceedings of the Communicating Nursing Research Conference and WIN Assembly, "Responding to Societal Imperatives Through Discovery and Innovation", held April 10-12, 2003, Scottsdale, Arizona. Communicating Nursing Research. 2003;36:373-.
- 197. Fredriksen EH, Moland KM, Sundby J. "Listen to your body": A qualitative text analysis of internet discussions related to pregnancy health and pelvic girdle pain in pregnancy. Patient Education & Counseling. 2008;73(2):294-9.
- 198. Atkinson MJ, Lohs J, Kuhagen I, Kaufman J, Bhaidani S. A promising method for identifying cross-cultural differences in patient perspective: the use of Internet-based focus groups for content validation of new Patient Reported Outcome assessments. Health & Quality of Life Outcomes. 2006;4:64-14.
- 199. Kontos EZ, Emmons KM, Puleo E, Viswanath K. Communication Inequalities and Public Health Implications of Adult Social Networking Site Use in the United States. Journal of Health Communication. 2010;15:216-35.
- 200. Stallard P, Velleman S, Richardson T. Computer Use and Attitudes Towards Computerised Therapy Amongst Young People and Parents Attending Child and Adolescent Mental Health Services. Child & Adolescent Mental Health. 2010;15(2):80-4.

- 201. Nicholas DB, Lach L, King G, Scott M, Boydell K, Sawatzky BJ, et al. Contrasting Internet and Face-to-Face Focus Groups for Children with Chronic Health Conditions: Outcomes and Participant Experiences. International Journal of Qualitative Methods. 2010;9(1):105-21.
- 202. Wikgren M. Everyday health information exchange and citation behaviour in Internet discussion groups. New Review of Information Behaviour Research. 2003;4(1):225-39.
- 203. Sneijder P, Te Molder HFM. 'Health Should Not Have to Be a Problem': Talking Health and Accountability in an Internet Forum on Veganism. Journal of Health Psychology. 2004;9(4):599-616.
- 204. Baum LS. Internet Parent Support Groups for Primary Caregivers of a Child with Special Health Care Needs. Pediatric Nursing. 2004;30(5):381-401.
- 205. Seçkin G. Internet Technology in Service of Personal Health Care Management: Patient Perspective. Journal of Technology in Human Services. 2009;27(2):79-92.
- 206. Gao F, Zhang M, Sadri S. Newspapers Use More Sources Compared To Health Blogs in H1N1/Swine Flu Coverage. Newspaper Research Journal. 2011;32(2):89-96.
- 207. Lupianez-Villanueva F, Mayer MA, Torrent J. Opportunities and challenges of Web 2.0 within the health care systems: An empirical exploration. Informatics for health & social care. 2009;34(3):117-26.
- 208. Townsend L, Gearing R. P03-163 The influence of health beliefs and social stigma on choice of internet support groups over formal mental health services. European Psychiatry. 2011;26:1332-.
- 209. Zufferey MC, Schulz PJ. Potentialities and limits of Internet health communication in chronic care: results from a qualitative study. Social Semiotics. 2010;20(1):61-75.
- 210. Stoddard JL, Augustson EM. Smokers who use Internet and smokers who don't: Data from the Health Information and National Trends Survey (HINTS). Nicotine & Tobacco Research. 2006;8:77-85.
- 211. Noel JG, Epstein J. Social Support and Health Among Senior Internet Users: Results of an Online Survey. Journal of Technology in Human Services. 2003;21(3):35-54.
- 212. Colineau N, Paris C. Talking about your health to strangers: understanding the use of online social networks by patients. New Review of Hypermedia & Multimedia. 2010;16(1/2):141-60
- 213. Rains SA, Keating DM. The Social Dimension of Blogging about Health: Health Blogging, Social Support, and Well-being. Communication Monographs. 2011;78(4):511-34.
- 214. Smedberg Å. To Design Holistic Health Service Systems on the Internet. International Journal of Biological & Medical Sciences. 2009;4(2):93-9.
- 215. Lim Fat MJ, Doja A, Barrowman N, Sell E. YouTube Videos as a Teaching Tool and Patient Resource for Infantile Spasms. Journal of Child Neurology. 2011;26(7):804-9.
- 216. Stoller EP, Webster NJ, Blixen CE, McCormick RA, Perzynski AT, Kanuch SW, et al. Lay Management of Chronic Disease: A Qualitative Study of Living with Hepatitis C Infection. American Journal of Health Behavior. 2009;33(4):376-90.
- 217. Bränström R, Penilla C, J.; P-SE, F. MR. Positive Affect and Mood Management in Successful Smoking Cessation. American Journal of Health Behavior. 2010;34(5):553-62.
- 218. Ikemba CM, Kozinetz CA, Feltes TF, Fraser JCD, McKenzie ED, Shah N, et al. Internet Use in Families With Children Requiring Cardiac Surgery for Congenital Heart Disease. Pediatrics. 2002;109(3):419.
- 219. Jayanti RK. A Netnographic Exploration. Journal of Advertising Research. 2010;50(2):181-96.

- 220. Beard L, Morra D, Wilson K, Keelan J. A Survey of Health-Related Activities on Second Life 2009. p. 12-.
- 221. Donelle L, Hoffman-Goetz L. An Exploratory Study of Canadian Aboriginal Online Health Care Forums. Health Communication. 2008;23(3):270-81.
- 222. Whitten P, Smith S, Munday S, LaPlante C. Communication Assessment of the Most Frequented Breast Cancer Websites: Evaluation of Design and Theoretical Criteria. Journal of Computer-Mediated Communication. 2008;13(4):880-911.
- 223. Ye Y. Correlates of Consumer Trust in Online Health Information: Findings From the Health Information National Trends Survey. Journal of Health Communication. 2011;16(1):34-49.
- 224. Stoddard JL, Augustson EM, Moser RP. Effect of adding a virtual community (bulletin board) to smokefree.gov: randomized controlled trial. 2008. p. 14-.
- 225. Bessière K, Pressman S, Kiesler S, Kraut R. Effects of internet use on health and depression: a longitudinal study. 2010. p. 4-.
- 226. Hu Y, Sundar SS. Effects of Online Health Sources on Credibility and Behavioral Intentions. Communication Research. 2010;37(1):105-32.
- 227. Kummervold PE, Chronaki CE, Lausen B, Prokosch H-U, Rasmussen J, Santana S, et al. eHealth Trends in Europe 2005-2007: A Population-Based Survey 2008. p. 4-.
- 228. Shaw BR, McTavish F, Hawkins R, Gustafson DH, Pingree S. Experiences of Women with Breast Cancer: Exchanging Social Support over the CHESS Computer Network. Journal of Health Communication. 2000;5(2):135-59.
- 229. Wen K-Y, McTavish F, Kreps G, Wise M, Gustafson D. From Diagnosis to Death: A Case Study of Coping With Breast Cancer as Seen Through Online Discussion Group Messages. Journal of Computer-Mediated Communication. 2011;16(2):331-61.
- 230. Dutta MJ, Feng H. Health Orientation and Disease State as Predictors of Online Health Support Group Use. Health Communication. 2007;22(2):181-9.
- 231. Weaver JB, Thompson NJ, Weaver SS, Hopkins GL. Healthcare non-adherence decisions and internet health information. 2009. p. 1373-80.
- 232. Gustafson DH, Hawkins R, McTavish F, Pingree S, Wei Chih C, Volrathongchai K, et al. Internet-Based Interactive Support for Cancer Patients: Are Integrated Systems Better? Journal of Communication. 2008;58(2):238-57.
- 233. Weber KM, Solomon DH. Locating Relationship and Communication Issues Among Stressors Associated With Breast Cancer. Health Communication. 2008;23(6):548-59.
- 234. Bylund CL. Mothers'Involvement in Decision Making During the Birthing Process: A Quantitative Analysis of Women's Online Birth Stories. Health Communication. 2005;18(1):23-39.
- 235. Baek TH, Yu H. Online health promotion strategies and appeals in the USA and South Korea: a content analysis of weight-loss websites. Asian Journal of Communication. 2009;19(1):18-38.
- 236. Nordfeldt S, Hanberger L, Berterö C. Patient and parent views on a Web 2.0 Diabetes Portal--the management tool, the generator, and the gatekeeper: qualitative study. 2010. p. 9-.
- 237. Wright K. Perceptions of On-Line Support Providers: An Examination of Perceived Homophily, Source Credibility, Communication and Social Support Within On-line Support Groups. Communication Quarterly. 2000;48(1):44-59.

- 238. McDowell H, Eunkyung K, Shaw BR, Jeong Yeob H, Gumieny L. Predictors and Effects of Training on an Online Health Education and Support System for Women with Breast Cancer. Journal of Computer-Mediated Communication. 2010;15(3):412-26.
- 239. Caiata Zufferey M, Schulz PJ. Self-management of chronic low back pain: An exploration of the impact of a patient-centered website. 2009. p. 27-32.
- 240. Chou W-yS, Hunt YM, Hesse BW, Beckjord EB, Moser RP. Social Media Use in the United States: Implications for Health Communication 2009. p. 9-.
- 241. Yoo JH, Su Ahn J, Taewoong C. Sociocultural Determinants of Negative Emotions Among Dementia Caregivers in the United States and in Korea: A Content Analysis of Online Support Groups. Howard Journal of Communications. 2010;21(1):1-19.
- 242. Weisgerber C. Turning to the Internet for Help on Sensitive Medical Problems. Information, Communication & Society. 2004;7(4):554-74.
- 243. Owen JE, Boxley L, Goldstein MS, Lee JH, Breen N, Rowland JH. Use of Health-Related Online Support Groups: Population Data from the California Health Interview Survey Complementary and Alternative Medicine Study. Journal of Computer-Mediated Communication. 2010;15(3):427-46.
- 244. Arduser L. Warp and Weft: Weaving the Discussion Threads of an Online Community. Journal of Technical Writing & Communication. 2011;41(1):5-31.
- 245. Cousineau TM, Rancourt D, Green TC. Web Chatter Before and After the Women's Health Initiative Results: A Content Analysis of On-line Menopause Message Boards. Journal of Health Communication. 2006;11(2):133-47.
- 246. Stroever SJ, Mackert MS, McAlister AL, Hoelscher DM. Using social media to communicate child health information to low-income parents. Preventing Chronic Disease. 2011;8(6):A148.
- 247. Heaivilin N, Gerbert B, Page JE, Gibbs JL. Public Health Surveillance of Dental Pain via Twitter. Journal of Dental Research. 2011;90(9):1047-51.
- 248. Shah SGS, Robinson I. Patients' perspectives on self-testing of oral anticoagulation therapy: Content analysis of patients' internet blogs. BMC Health Services Research. 2011;11.
- 249. Sanematsu H. FUN WITH FACEBOOK: THE IMPACT OF FOCUS GROUPS ON THE DEVELOPMENT OF AWARENESS CAMPAIGNS FOR ADOLESCENT HEALTH. Journal of Adolescent Health. 2011;48(2):S44-S5.
- 250. Kolpa E, Moreno M. RELATIONSHIP OF WEIGHT REFERENCES TO MENTAL HEALTH AND STRESS REFERENCES ON COLLEGE FRESHMAN FACEBOOK PROFILES. Journal of Adolescent Health. 2011;48(2):S92-S3.
- 251. Prier KW, Smith MS, Giraud-Carrier C, Hanson CL. Identifying Health-Related Topics on Twitter An Exploration of Tobacco-Related Tweets as a Test Topic. In: Salerno JYSJNDCSK, editor. Social Computing, Behavioral-Cultural Modeling and Prediction2011. p. 18-25.
- 252. Benker T, Arikan Y, Benker T. Turkish Patients' Use of Internet and Social Media for Healthcare and Drug Side Effect Information. Drug Safety. 2011;34(10):1003-.
- 253. Househ M. Sharing sensitive personal health information through Facebook: the unintended consequences. Studies in health technology and informatics. 2011;169:616-20.
- 254. Kang H, Veach PM, LeRoy BS. Concerns of South Korean Patients and Family Members Affected with Genetic Conditions: A Content Analysis of Internet Website Messages. Journal of Genetic Counseling. 2010;19(3):280-95.

- 255. Scanfeld D, Scanfeld V, Larson EL. Dissemination of health information through social networks: Twitter and antibiotics. American Journal of Infection Control. 2010;38(3):182-8.
- 256. Fisher RE, Donziger M, Lahr L. Analysis of the MyLifeLine.org Cancer Foundation user database: Does a cancer patient website for internet communication between cancer patients and their wellwisher community assist in cancer patient support? Journal of Clinical Oncology. 2009;27(15).
- 257. Acaster SL, Wild D. A NOVEL COMPARISON OF QUALITATIVE DATA SOURCES: CONTENT ANALYSIS OF SEMI STRUCTURED PATIENT INTERVIEWS VERSUS WEBLOGS (BLOGS). Value in Health. 2009;12(3):A31-A.
- 258. Fernandez-Luque L, Elahi N, Grajales FJ, 3rd;. An analysis of personal medical information disclosed in YouTube videos created by patients with multiple sclerosis. Studies in health technology and informatics. 2009;150:292-6.
- 259. Seeman N. Inside the health blogosphere: quality, governance and the new innovation leaders. Healthcare quarterly (Toronto, Ont). 2009;12(1):99-4.
- 260. Bacque M-F, Gsell-Herold G. The blogs of cancer patients: Self-reorganization through writing activity. Psycho-Oncology. 2008;17:S102-S.
- 261. Czarnecka-Kujawa K, Abdalian R, Grover SC. The quality of open access and open source internet material in gastroenterology: Is wikipedia appropriate for knowledge transfer to patients? Gastroenterology. 2008;134(4):A325-A6.
- 262. Hartoonian N, Ormseth S, Bantum EOC, Owen J. Process and outcome evaluation of a social-networking website for health promotion. Annals of Behavioral Medicine. 2008;35:S63-S.
- 263. Chung DS, Kim S. Blogging activity among cancer patients and their companions: Uses, gratifications, and predictors of outcomes. Journal of the American Society for Information Science and Technology. 2008;59(2):297-306.
- 264. Harrison S, Barlow J, Williams G. The content and interactivity of health support group websites. Health Education Journal. 2007;66(4):371-81.
- 265. Mehta SA. What can physicians learn from the blogs of patients with uveitis? Ocular Immunology and Inflammation. 2007;15(6):421-3.
- 266. Tsai CC, Tsai SH, Zeng-Treitler Q, Liang BA, Mackey T. Patient-centered consumer health social network websites: a pilot study of quality of user-generated health information. AMIA Annual Symposium proceedings / AMIA Symposium AMIA Symposium. 2007:1137.
- 267. Overberg R, Toussaint P, Zwetsloot-Schonk B. Illness stories on the Internet: Features of websites disclosing breast cancer patients' illness stories in the Dutch language. Patient Education and Counseling. 2006;61(3):435-42.
- 268. Coleman J, Olsen SJ, Sauter PK, Baker D, Hodgin MB, Stanfield C, et al. The effect of a Frequently Asked Questions module on a pancreatic cancer Web site patient/family chat room. Cancer Nursing. 2005;28(6):460-8.
- 269. Ezpeleta D. Virtual medical advice to unknown patients by e-mail: an analysis of 219 cases of headache. Cephalalgia. 2005;25(10):962-.
- 270. Rutherford P, Forte P, Bowen T, Rainbow R, Morrison A. Detailed analysis of a patient information website bulletin board What do patients want to know? American Journal of Kidney Diseases. 2004;43(4):A40-A.
- 271. Zhang J, Shen J, Wen QX, Li AP, Medimond. Research report on Red-Ribbon live webcast program of China reproductive health website Analysis on the dialogue between the specialists and Internet users 2004. 101-5 p.

- 272. Stout PA, Villegas J, Kim H. Enhancing learning through use of interactive tools on health-related websites. Health Education Research. 2001;16(6):721-33.
- 273. Yoon S. Application of Social Network Analysis and Text Mining to Characterize Network Structures and Contents of Microblogging Messages: An Observational Study of Physical Activity-Related Tweets [3451533]. United States -- New York: Columbia University; 2011.
- 274. Admane L. E-patients and Social Media: Impact of Online Experience on Perceived Quality of Care [1502921]. United States -- Arizona: Arizona State University; 2011.
- 275. Lao L. Evaluation of a social networking based SNAP-Ed nutrition curriculum on behavior change [1501411]. United States -- Rhode Island: University of Rhode Island; 2011.
- 276. Phua J. The social groups approach to quitting smoking: An examination of smoking cessation online and offline through the influence of social norms, social identification, social capital and social support [3466083]. United States -- California: University of Southern California; 2011.
- 277. Stroever S. The use of social media to communicate child health information to low-income parents: A formative study [1494830]. United States -- Texas: The University of Texas School of Public Health; 2011.
- 278. Vicdan H. Constitution of the market through social media: Dialogical co-production of medicine in a virtual health community organization [3423915]. United States -- Texas: The University of Texas Pan American; 2010.
- 279. Yamauchi E. LoveLife, MYMsta, and Mizz B: Evaluating a mobile phone-based social network to prevent HIV among youth in South Africa [1476700]. United States -- North Carolina: The University of North Carolina at Chapel Hill; 2010.
- 280. Chew C. Pandemics in the Age of Twitter: A Content Analysis of the 2009 H1N1 Outbreak [MR72542]. Canada: University of Toronto (Canada); 2010.
- 281. Bresolin J. Social Networks for Social Change: Looking into Morality and Accountability during times of National Crises in the United States [MR73799]. Canada: University of Ottawa (Canada); 2010.
- 282. Moturu S. Quantifying the trustworthiness of user-generated social media content [3371224]. United States -- Arizona: Arizona State University; 2009.
- 283. Hether H. Social media and health: Social support and social capital on pregnancy-related social networking sites [3389472]. United States -- California: University of Southern California; 2009.
- 284. Corley C. Social network simulation and mining social media to advance epidemiology [3385778]. United States -- Texas: University of North Texas; 2009.