

BMJ Open Quantifying the prevalence of frailty in English hospitals

J Soong,^{1,2} AJ Poots,¹ S Scott,³ K Donald,³ T Woodcock,¹ D Lovett,¹ D Bell¹

To cite: Soong J, Poots AJ, Scott S, *et al*. Quantifying the prevalence of frailty in English hospitals. *BMJ Open* 2015;**5**:e008456. doi:10.1136/bmjopen-2015-008456

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-008456>).

Received 15 April 2015

Revised 31 July 2015

Accepted 21 August 2015



► <http://dx.doi.org/10.1136/bmjopen-2015-008457>



CrossMark

¹NIHR CLAHRC Northwest London, Imperial College London, London, UK

²Royal College of Physicians, London, UK

³Oliver Wyman, London, UK

Correspondence to

Dr J Soong;
johnsoong@imperial.ac.uk

ABSTRACT

Objectives: Population ageing has been associated with an increase in comorbid chronic disease, functional dependence, disability and associated higher health care costs. Frailty Syndromes have been proposed as a way to define this group within older persons. We explore whether frailty syndromes are a reliable methodology to quantify clinically significant frailty within hospital settings, and measure trends and geospatial variation using English secondary care data set Hospital Episode Statistics (HES).

Setting: National English Secondary Care Administrative Data HES.

Participants: All 50 540 141 patient spells for patients over 65 years admitted to acute provider hospitals in England (January 2005—March 2013) within HES.

Primary and secondary outcome measures: We explore the prevalence of Frailty Syndromes as coded by International Statistical Classification of Diseases, Injuries and Causes of Death (ICD-10) over time, and their geographic distribution across England. We examine national trends for admission spells, inpatient mortality and 30-day readmission.

Results: A rising trend of admission spells was noted from January 2005 to March 2013 (daily average admissions for month rising from over 2000 to over 4000). The overall prevalence of coded frailty is increasing (64 559 spells in January 2005 to 150 085 spells by Jan 2013). The majority of patients had a single frailty syndrome coded (10.2% vs total burden of 13.9%). Cognitive impairment and falls (including significant fracture) are the most common frailty syndromes coded within HES. Geographic variation in frailty burden was in keeping with known distribution of prevalence of the English elderly population and location of National Health Service (NHS) acute provider sites. Overtime, in-hospital mortality has decreased (>65 years) whereas readmission rates have increased (esp. >85 years).

Conclusions: This study provides a novel methodology to reliably quantify clinically significant frailty. Applications include evaluation of health service improvement over time, risk stratification and optimisation of services.

INTRODUCTION

People are living longer. At present, it is estimated that 16.1% of the European population is over the age of 65 years (>65 years),

Strengths and limitations of this study

- This study is the first to attempt to use frailty syndromes as an operational definition within an English secondary care data set.
- The methodology uses whole population routinely collected data, with robust trend analysis examining coding reliability.
- This study is a retrospective analysis reliant on the accuracy, reliability and retrospective nature of coding within Hospital Episode Statistics.

and this number is expected to rise to 22% by 2031.¹ In the developed world, the increase is greatest in those over 80 years, and this equates to approximately 3 million people in the UK.² In health terms patients >65 years now constitute two-thirds of the general hospital population, account for 40% of all hospital bed days and 65% of National Health Service (NHS) spend.³ Recent analysis suggests population ageing contributes directly to the increase in emergency admissions to hospitals.⁴

Associated with this demographic shift there has been an increase in comorbid chronic disease, functional dependence, disability, poorer quality of life and higher health care costs.^{5 6} Patients in this category are often considered frail. Currently, there is no universally agreed operational definition for frailty.⁷ Frailty has been described as a clinical phenotype or a biophysical syndrome of accumulated deficit (frailty index). Phenotypic models describe frailty as specific clinical syndrome encompassing a cluster of characteristics, namely unintentional weight loss, exhaustion, weakness, slowness and low physical activity.⁸ The frailty index is characterised by decreased resistance to stressors resulting from the accumulation of deficit across multiple physiological systems, culminating in an increased risk of adverse outcomes.^{9 10} Methodologies to reliably identify the 'frail' at-risk cohort within secondary care, both at patient and population level, are a current research priority.^{11–13}

In clinical practice the terms Geriatric Giants,¹⁴ Geriatric Syndromes^{15 16} or Frailty Syndromes¹⁷ are often used to describe clinically vulnerable group within the elderly. They likely represent high order clinical manifestations of multifactorial processes resultant from the accumulation and interaction of deficits and environmental factors. They include cognitive impairment, falls, mobility problems, pressure ulcers and incontinence. These syndromes, more prevalent in the elderly, confer a higher risk of death,⁸ institutionalisation,¹⁸ disability and poor quality of life.¹⁵ They are arguably the consequences of frailty, or the manifestation of clinically significant frailty.¹⁹ Current National guidelines for the care of the older person in acute care recommend using frailty syndromes as a possible methodology to assess for frailty.^{17 20}

In this study, we measure the trends for all hospital admissions, in-hospital death and readmissions for those over 65 years. We describe Frailty Syndromes^{17 20} as an operational definition within the English secondary care data set Hospital Episode Statistics (HES) in order to examine the frailty burden between 2005 and 2012. In addition we describe the geospatial variation of frailty in English secondary healthcare settings. We compare our results with the existing literature on frailty prevalence and discuss possible applications of this methodology.

METHODS

Data sources

HES is a national administrative database containing patient-level records of all admissions to NHS hospitals in England.²¹ It has high levels of data completeness and rigorous data cleaning processes to ensure data quality. Each record in HES corresponds to a finished consultant episode, during which a patient is under the care of an individual consultant. These episodes were aggregated into hospital spells covering a patient's total length of stay in a hospital (ie, a hospital admission) using established methodology.²²

HES contains 20 fields per record for diagnoses codes that are defined using the tenth revision of the International Statistical Classification of Diseases, Injuries and Causes of Death (ICD-10). The first of these is the primary diagnosis, with the rest available for coding of comorbidities or complications. HES does not contain present-on-admission flags. We reviewed HES for ICD-10 diagnostic codes that could be grouped for frailty syndromes (see online supplementary appendix 1) in all 20 fields. We included only inpatients at acute non specialist hospital trusts, with elective and non-elective admissions for those 65 years and over >65 years. We excluded hyper-specialist hospitals and mental health units as they have a very different case-mix and data quality.²³ Thus, we defined frailty as the presence of at least one frailty syndrome and within the cohort of patients greater than 65 years old.

Annual trend profiles were created for the grouped ICD-10 diagnostic codes from January 2005 to March

2013 to determine coding reliability and shifts (see online supplementary appendix 2). The spells were aggregated both by age-band (65–74; 75–84; >85 years) and monthly. Monthly data are visualised as simple line plots in the first instance. Office of National Statistics (ONS) databases were queried for population size estimates or census data where available.

Study population

All hospital admissions for >65 years to English acute trusts between January 2005 and March 2013 (N=50 540 141 patient spells) were available for analysis.

Temporal analysis

To analyse the variation present in these time-series data, statistical process control is used to separate special cause variation (signal) from common cause variation, an inherent property of all systems. The XmR chart is used as it is a method that is not dependent on data distributions or underlying assumptions.²⁴ When analysing count data, daily averages for months were calculated to correct for unequal 'areas of opportunity'; for example, a count of February admissions will be lower by virtue of fewer days in February, and daily averages account for the difference in available days. For percentage data, such a correction is attained through division by the denominator—all spells and all spells with frailty. Adjustments for seasonal variation are made, and seasonalised reference lines are plotted, for more natural interpretation of the charts. In this work, a standard rule set for detection of signal is adopted, using Microsoft Excel to construct the charts.²⁴

Geospatial analysis

Geo-location is the identification of real-world geographic location of an object. Postcodes of provider sites were used to geo-locate sites, and map elements were derived from open source data provided by Office for National Statistics. Geo-locations aggregated to Primary Care Trust (PCT) level were attached to counts of frailty syndromes for patients >65 years admitted to NHS acute providers in 2012 as this is the applicable unit for these data. Choropleths are thematic maps that shade or colour areas to represent classified values of specific phenomena. ESRI ArcMap V.10.2 software was used to create a choropleth map. Annual trend profiles for inpatient mortality and non-elective readmission within 30 days were plotted. This temporal range of April 2006 to December 2012 was selected due to changes in structure of health geographies within England in 2006,²⁶ and to allow a sufficient follow up period to more accurately reflect the clinical outcomes listed above.

RESULTS

Between January 2005 and March 2013, there was a rising trend with daily average admissions for month increasing from over 2000 to over 4000 (figure 1A).




Figure 1 (A) Daily average admission spells for month and percentage total frailty burden for England NHS acute trusts. (B) The number and percentage of spells for patients >65 years by age-band admitted to English acute providers. LNPL, Lower Natural Process Limits; NHS, National Health Service; UNPL, Upper Natural Process Limits.

There has been an increase in all age bands over this period, 65–74 increasing from 161 641 to 235 756, 75–84 increasing from 162 817 to 233 870 and >85 increasing from 71 396 to 137 991 (figure 1B). The relative proportion of total admissions has remained constant each age band at 40%, 40% and 20%, respectively. Examination of ONS data, (see online supplementary appendix 4) finds that in the general UK population the number of >64 years old in the population increased from 8 031 000 in 2005, to 9 051 179 in 2013. In 2005, the 65–74s represented 52% of those >65-year; in 2013 it was 54%; 75–84s were 36% and 33% 2005–2013; and >85s were 12% and 13%.

Analysis of trends shows that the coded overall frailty burden, based on the coding of at least one frailty syndrome, has increased from 12% to 14% between January 2005 and March 2013. There is evidence of seasonal peaks during winter, partly explained by similar patterns in admission spells (figure 2).

The coding of the frailty syndromes has increased between 2005 and 2013. Most patients had one frailty syndrome coded (figure 3) and the most common frailty syndromes described between 2005 and 2013 were cognitive impairment and falls (including significant fracture) with cognitive impairment increasing to the same levels as falls representing approximately 10% of all spells in the those >65 years. Anxiety and/or depression has increased particularly from 2010 (2.4%) to

2013 (>4%) (figure 4). There is a persistent and steady rise in coding for mobility problems.

Evaluating the frailty syndromes individually, the very elderly (>85 years) represent between 40% and 50% of the spells coded for that syndrome, with rising trend. The exception to this was anxiety and/or depression syndrome, which exhibited a rising trend in the 65–74 s, and the 75–84 s accounted for the largest group (see online supplementary appendix 3). Age-band stratification shows that cognitive impairment and falls in age-bands >85 years and 75–84 years account for a large majority of coded frailty syndromes within this cohort. These four groups accounted for 60.2% of frailty syndromes coded over this time period (N=7 399 671)

Geographic variation in the frailty burden across admission spells in England was seen based on the 2012 HES data (figure 5). For patients >65-year admitted to England Acute providers, the highest levels of frailty are seen in the Northeast, Central and South Coast. The top five PCTs for highest admissions numbers are Nottingham City, Halton & St Helens, Warrington, Waltham Forrest and Wolverhampton city.

Between April 2006 to December 2012, 1 160 299 (3.4%) spells were associated with inpatient mortality, though a decreasing trend is observed for example, April 2006 (N=15 042) to April 2012 (N=14 437) (figure 6A). Non-elective re-admission rates within 30 days of discharge have increased for all admissions > 65 years




Figure 2 The percentage of admissions to English acute providers coded with at least one frailty syndrome. LNPL, Lower Natural Process Limits; UNPL, Upper Natural Process Limits.

from approximately 11–12% (figure 6B). The rates of readmission increased across the age bands >65 years (10%), 75–84 (12%) and >85 (14%). Though the

overall number of very elderly (>85 years) with non-elective 30-day readmission is lower than the other two age-bands, they have more readmissions (figure 7).




Figure 3 Trends for the prevalence of count of frailty syndromes and total frailty burden for patients >65 years admitted to NHS acute provider hospitals between April 2006 and December 2012. NHS, National Health Service.




Figure 4 Trends for the prevalence of frailty syndromes for patients >65 years admitted to NHS acute provider hospitals between January 2005 and March 2013. NHS, National Health Service.

DISCUSSION

Frailty is often defined as a clinical state in which there is an increase in an individual's vulnerability for adverse events and harm when exposed to a stressor.²⁵ It is distinct but related to disability and comorbidity.²⁶⁻²⁷ Some approaches to the measurement of frailty have been characteristically biophysical with emphasis on detection of the consequences of sarcopaenia and chronic inflammation-malnutrition.⁸ Another approach is to measure frailty in relation to the clinical consequences of accumulated loss and insufficiency in ageing individuals (ie, the relationship to mortality and adverse outcomes).²⁸ Both approaches appear complementary²⁹ and overlap, though not completely.³⁰ Frailty measurement is problematic in the acute care setting. High levels of disease acuity on top of chronic multimorbidity, multidimensional complexity and diagnostic uncertainty are challenging for healthcare systems, with increasing evidence and concern for compromised patient safety, quality of care and experience.³¹⁻³⁴

We have examined the prevalence of frailty syndromes within English HES data from both a temporal and geo-spatial point of view. Temporal analysis, it allows us to observe shifts in diagnostic coding, and observe trend in signal changes over time. Spatial analysis allows us to explore geographic heterogeneity of frailty syndrome

prevalence, with consequent implications for service provision and equity of care.

Comparison with ONS data, the corresponding admissions to English acute providers for patients with frailty syndromes is larger than might be expected by demographic shift associated with ageing. Additionally, 75–84s make up approximately one-third of the population of those over 65 years, but have 40% of the admissions, and >85s are approximately 13% of the population of those over 65 years but have 20% of the admissions.

This study has focused on patients admitted to hospital >65 years in England to better understand the impact of frailty syndromes. To the authors knowledge, this is the first study to examine the prevalence of frailty syndromes for patients >65 years across England. This study confirms increasing number of >65 years admitted to hospital (elective and non-elective). The relative burden of coded frailty syndromes has increased over this period with cognitive impairment increasing to similar levels to falls. Anxiety and/or depression is also increasing in this group.

When complex systems fail (biological or otherwise), high-order functions can be first disrupted.³⁵ Frailty syndromes represent the clinical manifestation of high-order disruption, providing a useful clinical marker of multidimensional deficit accumulation. The overall




Figure 5 Percentage of spells for patients >65 years with admission to NHS acute trusts with at least one frailty syndrome by PCT by quintiles (numerator=admission spells with at least one frailty syndrome; denominator=total admission spells to NHS acute trusts within English PCT). NHS, National Health Service; PCT, Primary Care Trust.

prevalence rate of frailty syndromes found in this study is 13.9%. Between 2005 and 2013, though there has been an increase in the numbers of patients admitted >65 years, the percentage by age band has remained stable, thus not suggesting major drift towards older age groups within the older population. However, within the >65 years group, frailty syndromes are more prevalent with the older age bands.

Prevalence rates of frailty vary depending on population and operational definition used in reported studies. Reported prevalence in community dwelling adults varies tremendously (from 4.0% to 59.1%).³⁶ A recent systematic review reported pooled frailty prevalence across 21 community dwelling study cohorts as 10.7%

(N=61 500).³⁶ The recent Survey of Health, Ageing and Retirement in Europe (SHARE) study reported frailty prevalence as 4.1% in community dwelling adults >50 years (N=16 584) in 10 European countries (prevalence of 17% in those over 65 years).³⁷ In the UK, the Hertfordshire Cohort Study³⁸ reported an overall prevalence of 6.3% for 638 community dwelling 64–74-year-olds, while the English Longitudinal study of ageing³⁹ reported a prevalence of 8% and 13% for 3055 community dwelling over 65-year-olds (using the Phenotype⁸ and Frailty Index¹⁰ definitions, respectively).

The prevalence of inpatient frailty in our study was lower than expected from smaller reported clinical studies within secondary care (range 24.7%–80%):




Figure 6 (A) Percentage of spells with inpatient mortality admitted to English providers and (B) non-elective 30-day readmission in patients >65 years admitted to English acute providers.

($n=220$ >70 years admitted to acute geriatric ward from Emergency department,⁴⁰ ($n=6701$) 40% (Phenotype) and 32.5% (SOF;⁴¹ ($n=1388$ >70 years admitted to cardiology service,⁴² ($n=900$ 827 % (Phenotype) and 63% (Frailty Scale⁴³); ($n=298$ >75 years admitted to five different specialist wards, 50–80% (Groningen Frailty Index^{44–45}); ($n=307$ >75 years with diagnosed non-ST elevation myocardial infarction,⁴⁶ 48.5% ($n=2305$ >65 years Clinical Frailty Scale⁴⁷); ($n=752$ medical inpatients >75 years.⁴⁸ In the UK, two recent studies^{12–13} reported frailty prevalence for $n=667$ patients >70 years admitted to Acute Medical Units (AMU) at 69% (ISAR,⁴⁹ 17.9% (Phenotype), 66.4% (SOF), 24.9% (Avila-Funes), 24.1% (Rothman) and 30.9% (Frailty index). Importantly, these studies mainly consisted of non-elective admissions, while our study cohort comprised of elective and non-elective admissions to hospital. However, it may be that this methodology truly underestimates the prevalence of frailty within HES.

Not all frailty syndromes are observed, within HES, to be equally prevalent, nor do they appear to be increasing at the same rate. The observed differences and increase in frailty syndromes in this study (figure 4) may reflect improvements in coding practice within HES due to the introduction of Healthcare Resource Group (HRG V.4 introduced in April 2007) and Payment by Results (since April 2009). The national dementia strategy was also published in 2009. However, this observed rising trend may also reflect a genuine increase in number of diagnosis. Correlation with clinical data sets

for comparison is consequently a necessary research priority.

The frailty syndromes are more prevalent in the very elderly (>85), with a rising trend. The exception to this is anxiety and/or depression, where the most prevalent age-band is 75–84 years, which exhibits a declining trend, while the increase in this anxiety and/or depression from 2010 appears to mainly be in the 65–74 age-band, a pattern noted independently by the HSCIC.⁵⁰ Correlation with clinical data sets is warranted to ensure accuracy.

This analysis suggests that coexistence of multiple frailty syndromes is uncommonly coded within HES; even though we used coded frailty syndromes within all 20 of HES diagnostic domains, incomplete coding may still be a cause, as not all morbidities will be acknowledged and coded for each admission, only those deemed relevant to care at that time. However, it has been noted that accumulation of deficit beyond a certain level is incompatible with survival,⁵¹ and thus multimorbidity would have a ceiling effect. Further investigation on multiple frailty syndromes could be profitable.

Inpatient mortality trends in this population exhibit seasonality with peaks during winter, which persist after adjustment for number of admissions (spells). These peaks, coupled with rising 30-day readmissions (particularly in the very elderly) suggest differences in service provision over the year. A question arises here: is this




Figure 7 Number and percentage non-elective readmissions in patient >65 years admitted to NHS acute providers by age-band. NHS, National Health Service.

seasonality appropriate for the UK population and the provision of care?

Geographic variation in frailty burden appears to be in keeping with known distribution of prevalence of the English elderly population and location of NHS acute provider sites, particularly within urban areas. Healthcare providers and commissioners should consider their local populations when planning services, where frailty may be a larger consideration than other locations. Further study into environmental factors in relation to frailty is a necessary next step.

Limitations

This study is a retrospective analysis reliant on data coded from hospital data warehouses, and subsequently cleaned into HES. As such, its validity is dependent on accuracy of data coding. Including all 20 diagnostic coding fields may help to mitigate this, but correlation with clinical data sets may be warranted for local investigations. Resultant prevalence rates described may underestimate frailty syndromes in this population.

Anxiety and/or depression was only recently recognised as a geriatric syndrome by the Education Committee Writing Group of the American Geriatrics Society.¹⁶ It appears to fulfil several criteria that makes it an attractive putative candidate for a frailty syndrome:⁵¹ poor mental health is often associated with chronic physical deficits,⁵² it appears to increase with age (figure 4),

it is associated with adverse outcome,⁵³ it is neither too rare or too common (figure 4) Recent study has linked it to frailty^{52 54} in older persons, though comprehensive study of its relationship to adverse outcomes with relation to frailty is still lacking. Further study, including correlation with clinical data sets, is warranted.

Conclusion

To our knowledge this study is the first to attempt to use frailty syndromes as an operational definition within an English secondary care data set. While the study is dependent on the accuracy, reliability and retrospective nature of coding within HES, its strengths include being a whole population analysis, with robust trend analysis examining coding reliability. It utilises routinely collected data and is comprehensive in its coding of frailty within all of the diagnostic coding positions in the HES data set. Future studies to correlate with clinical data sets are needed to further investigate the phenomena discovered in this study.

This study provides a methodology to reliably quantify frailty. Applications include the ability to evaluate the effect of interventions over time allowing for health service quality improvement. Geographic analysis allows providers and payers to highlight areas of need, unmet or otherwise for more intelligent targeting of resources, from a public health or clinical perspective. A reliable and quantifiable metric for frailty enables the

development of risk-prediction models and clinical scoring systems that will aid targeted interventions to vulnerable populations that will benefit most.

Contributors JS conceived study, designed analysis, interpreted results and wrote first draft. AJP designed analysis, interpreted results, contributed to ongoing writing. SS and KD designed analysis. TW designed SPC analysis. DL designed GIS analysis. DB conceived study, designed analysis, interpreted results and contributed to ongoing writing.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer This article presents independent research commissioned by the National Institute for Health Research (NIHR) under the Collaborations for Leadership in Applied Health Research and Care (CLAHRC) programme for North West London. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. JS received a Research Fellowship award from Chelsea and Westminster Health Charity. SS and KD seconded from Oliver Wyman to support this project. The lead author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Competing interests None declared.

Ethics approval As per Governance Arrangements for Research Ethics Committees (GAFREC), Research limited to secondary use of information previously collected in the course of normal care (without an intention to use it for research at the time of collection), provided that the patients or service users are not identifiable to the research team in carrying out the research.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- United Nations Department of Economics and Social Affairs. Demographic Yearbook 64th issue. 2013;64:1–912.
- Cornwell J. *The Care of Frail Older People with Complex Needs: Time for a revolution*. The Sir Roger Bannister Health Summit, Leeds Castle: The King's Fund, 2012.
- Department of Health. Improving care and saving money: learning the lessons on prevention and early intervention for older people. 2010.
- Blunt I, Bardsley M, Dixon J. Trends in emergency admissions in England 2004–2009: is greater efficiency breeding inefficiency. London: The Nuffield Trust, 2010.
- Marengoni A, Angleman S, Melis R, *et al*. Aging with multimorbidity: a systematic review of the literature. *Ageing Res Rev* 2011;10:430–9.
- Barnett K, Mercer SW, Norbury M, *et al*. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.
- Rodriguez-Manas L, Feart C, Mann G, *et al*. Searching for an operational definition of frailty: a Delphi method based consensus statement: the frailty operative definition-consensus conference project. *J Gerontol A Biol Sci Med Sci* 2013;68:62–7.
- Fried LP, Tangen CM, Walston J, *et al*. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56: M146–56.
- Ferrucci L, Guralnik JM, Studenski S, *et al*. Designing randomized, controlled trials aimed at preventing or delaying functional decline and disability in frail, older persons: a consensus report. *J Am Geriatr Soc* 2004;52:625–34.
- Rockwood K, Howlett SE, MacKnight C, *et al*. Prevalence, attributes, and outcomes of fitness and frailty in community-dwelling older adults: report from the Canadian study of health and aging. *J Gerontol A Biol Sci Med Sci* 2004;59:1310–17.
- Ellis G, Whitehead MA, O'Neill D, *et al*. Comprehensive geriatric assessment for older adults admitted to hospital. *Cochrane Database of Systematic Reviews* 2011;(7):CD006211.
- Edmans J, Bradshaw L, Gladman JRF, *et al*. The Identification of Seniors at Risk (ISAR) score to predict clinical outcomes and health service costs in older people discharged from UK acute medical units. *Age Ageing* 2013;42:747–53.
- Wou F, Gladman JR, Bradshaw L, *et al*. The predictive properties of frailty-rating scales in the acute medical unit. *Age Ageing* 2013;42:776–81.
- Isaacs B. *The challenge of geriatric medicine*. USA: Oxford University Press, 1992.
- Inouye SK, Studenski S, Tinetti ME, *et al*. Geriatric syndromes: clinical, research, and policy implications of a core geriatric concept. *J Am Geriatr Soc* 2007;55:780–91.
- [No authors listed]. Core competencies for the care of older patients: recommendations of the American Geriatrics Society. The Education Committee Writing Group of the American Geriatrics Society. *Acad Med* 2000;75:252–5.
- Acute Care Toolkit 3. *Acute medical care for frail older people*. London: Royal College of Physicians, 2012.
- Anpalahan M, Gibson SJ. Geriatric syndromes as predictors of adverse outcomes of hospitalization. *Intern Med J* 2008;38:16–23.
- Clegg A, Young J, Iliffe S, *et al*. Frailty in elderly people. *Lancet* 2013;381:752–62.
- Banerjee J, Conroy S, Cooke MW. Quality care for older people with urgent and emergency care needs in UK emergency departments. *Emerg Med J* 2013;30:699–700.
- HES Online: Hospital Episode Statistics. 2013. <http://www.hscic.gov.uk/hes> (accessed Nov 2014).
- Health and social care information centre. Methodology to create provider and CIP spells from HES APC data: 1–9, 2014. http://www.hscic.gov.uk/media/11859/Provider-Spells-Methodology/pdf/Spells_Methodology.pdf (accessed Nov 2014).
- Bottle A, Aylin P. Comorbidity scores for administrative data benefited from adaptation to local coding and diagnostic practices. *J Clin Epidemiol* 2011;64:1426–33.
- Lloyd P, Provost SM. The Health Care Data Guide: Learning from Data for Improvement. John Wiley & Sons October 2011:1–480.
- Morley JE, Vellas B, van Kan GA, *et al*. Frailty consensus: a call to action. *J Am Med Dir Assoc* 2013;14:392–7.
- Fried LP, Ferrucci L, Darer J, *et al*. Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci* 2004;59:M255–63.
- Theou O, Rockwood MR, Mitnitski A, *et al*. Disability and co-morbidity in relation to frailty: how much do they overlap? *Arch Gerontol Geriatr* 2012;55:e1–8.
- Rockwood K. What would make a definition of frailty successful? *Age Ageing* 2005;34:432–4.
- Cesari M, Gambassi G, van Kan GA, *et al*. The frailty phenotype and the frailty index: different instruments for different purposes. *Age Ageing* 2014;43:10–12.
- Blodgett J, Theou O, Kirkland S, *et al*. Frailty in NHANES: Comparing the frailty index and phenotype. *Arch Gerontol Geriatr* 2015;60:464–70.
- Young J, Hood C, Woolley R, *et al*. *Report of the National Audit of Dementia Care in General Hospitals 2011*. Royal College of Psychiatrists, 2011.
- Arora VM, McGory ML, Fung CH. Quality indicators for hospitalization and surgery in vulnerable elders. *J Am Geriatr Soc* 2007;55(Suppl 2):S347–58.
- Trembl J, Husk J, Lowe D, *et al*. Falling Standards, Broken Promises. Report of the national audit of falls and bone health in older people 2010. London: Royal College of Physicians, 2011.
- Wagg A, Harari D, Husk J, *et al*. *National Audit of Continence Care—Combined Organisational and Clinical Report*. London: Royal College of Physicians, 2010.
- Rockwood K, Mitnitski A, Song X, *et al*. Long-term risks of death and institutionalization of elderly people in relation to deficit accumulation at age 70. *J Am Geriatr Soc* 2006;54:975–9.
- Collard RM, Boter H, Schoevers RA, *et al*. Prevalence of frailty in community-dwelling older persons: a systematic review. *J Am Geriatr Soc* 2012;60:1487–92.
- Santos-Eggimann B, Cuenoud P, Spagnoli J, *et al*. Prevalence of frailty in middle-aged and older community-dwelling Europeans living in 10 countries. *J Gerontol A Biol Sci Med Sci* 2009;64: 675–81.
- Syddall H, Roberts HC, Evandrou M, *et al*. Prevalence and correlates of frailty among community-dwelling older men and women: findings from the Hertfordshire Cohort Study. *Age Ageing* 2010;39:197–203.

39. Hubbard RE, Lang IA, Llewellyn DJ, *et al.* Frailty, body mass index, and abdominal obesity in older people. *J Gerontol A Biol Sci Med Sci* 2010;65:377–81.
40. Joosten E, Demuyneck M, Detroyer E, *et al.* Prevalence of frailty and its ability to predict in hospital delirium, falls, and 6-month mortality in hospitalized older patients. *BMC Geriatr* 2014;14:1.
41. Ensrud KE, Ewing SK, Taylor BC, *et al.* Comparison of 2 frailty indexes for prediction of falls, disability, fractures, and death in older women. *Arch Intern Med* 2008;168:382–9.
42. Purser JL, Kuchibhatla MN, Fillenbaum GG, *et al.* Identifying frailty in hospitalized older adults with significant coronary artery disease. *J Am Geriatr Soc* 2006;54:1674–81.
43. Rockwood K, Stadnyk K, MacKnight C, *et al.* A brief clinical instrument to classify frailty in elderly people. *Lancet* 1999;353:205–6.
44. Andela RM, Dijkstra A, Slaets JP, *et al.* Prevalence of frailty on clinical wards: description and implications. *Int J Nurs Pract* 2010;16:14–19.
45. Steverink N, Slaets J, Schuurmans H, *et al.* Measuring frailty: developing and testing of the Groningen Frailty Indicator (GFI). *Gerontologist* 2001;41:236–7.
46. Ekerstad N, Swahn E, Janzon M, *et al.* Frailty is independently associated with short-term outcomes for elderly patients with non-ST-segment elevation myocardial infarction. *Circulation* 2011;124:2397–404.
47. Rockwood K, Song X, MacKnight C, *et al.* A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489.
48. Evans SJ, Sayers M, Mitnitski A, *et al.* The risk of adverse outcomes in hospitalized older patients in relation to a frailty index based on a comprehensive geriatric assessment. *Age Ageing* 2014;43:127–32.
49. McCusker J, Bellavance F, Cardin S, *et al.* Detection of older people at increased risk of adverse health outcomes after an emergency visit: the ISAR screening tool. *J Am Geriatr Soc* 1999;47:1229–37.
50. Health and Social Care Information Centre. Anxiety: hospital admissions highest in women in their late 60s. URL: <http://www.hscic.gov.uk/article/3916/Anxiety-hospital-admissions-highest-in-women-in-their-late-60s> (accessed Nov 2014).
51. Searle SD, Mitnitski A, Gahbauer EA, *et al.* A standard procedure for creating a frailty index. *BMC Geriatr* 2008;8:24.
52. Ni Mhaolain AM, Fan CW, Romero-Ortuno R, *et al.* Frailty, depression, and anxiety in later life. *Int Psychogeriatr* 2012;24:1265–74.
53. Dent E, Hoogendijk EO. Psychosocial factors modify the association of frailty with adverse outcomes: a prospective study of hospitalised older people. *BMC Geriatr* 2014;14:108.
54. Mezuk B, Edwards L, Lohman M, *et al.* Depression and frailty in later life: a synthetic review. *Int J Geriatr Psychiatry* 2012;27:879–92.

Correction

Soong J, Poots AJ, Scott S, *et al.* Quantifying the prevalence of frailty in English hospitals. *BMJ Open* 2015;5:e008456. The corresponding author's email address is incorrect in this paper. The correct address is j.soong@imperial.ac.uk

BMJ Open 2015;5:e008456corr1. doi:10.1136/bmjopen-2015-008456corr1



CrossMark

Appendix 1

Frailty Syndrome	ICD-10 Diagnostic Code
Anxiety and Depression	F320 F320- F320-- F320-D F3200 F3200- F3200A F3200D F3201 F3201A F3201D F3207 F320X F321 F321 1 F321- F321-- F321-D F3210 F3210- F3210A F3210D F3211 F3211- F32110 F32111 F3211A F3211D F3219 F322 F322 D F322- F322-D F32211 F3229 F322X F323 F323 D F323- F323-- F323-D F3230 F3231 F3239 F324 F325 F326 F327 F328 F328 A F328- F3289 F328A F329 F329 A F329 D F329- F329-- F329-A F329-D F329. F329/ F3290 F3292 F3293 F3295 F3296 F3298 F3299 F329A F329D F329J2 F329M F329Q F32X F32X- F33#- F330 F330- F330-D F3300 F3300A F3301 F3301A F3301D F331 F331 1 F331- F331-D F3310 F3310- F3310A F3310D F3311 F3311- F3311A F3311D F332 F332- F332-- F332-D F3320 F3329 F333 F333- F333-D F3330 F3331 F3333 F334 F334- F335 F336 F337 F338 F338- F338-D F3380 F339 F339 A F339- F339-- F339-D F3396 F33X F380 F380- F3800 F3800A F3800D F381 F381- F3810 F3810A F3810D F388 F388- F38X F410 F410- F410-- F4100 F4101 F4103 F410D F411 F411- F411-D F412 F412- F412-- F4122 F412D F413 F413- F418 F418- F419 F419- F419-- F4193 F4199 F419X F41X F430 F430- F430-D F4300 F4301 F4302 F431 F431- F431-- F432 F432 0 F432 2 F432 3 F432 5 F432- F432-- F432-D F4320 F4320A F4320D F4320X F4321 F4321- F4321A F4321D F4322 F4322- F4322A F4322D F4323 F4323A F4323D F4324 F4325 F4325- F4325A F4325D F4328 F4328A F4328D F4329 F432X F438 F438- F439 F439- F43X F440 F440- F441 F441- F442 F442- F4422 F443 F443- F444 F444- F445 F445- F446 F446- F447 F447- F448 F448- F4480 F4481 F4481A F4481D F4482 F4488 F449 F449-
Delirium	F050 F050 A F050- F051 F051 A F051 D F051- F051-A F051-D F0513 F051D F058 F058- F058-- F059 F059 D F059- F059--
Dementia	F000 F000 A F000 D F000* F000+ F000- F000-A F000-D F0000 F00001 F00002 F0000A F0001 F00010 F0001A F0002 F0002A F0003 F00031 F00032 F0004 F00040 F00041 F00042 F0004A F0009 F0009A F000a F001 F001 0 F001 1 F001 A F001 D F001* F001+ F001- F001-A F001-D F0010 F00101 F00102 F0010A F0011 F00111 F00112 F0011A F0012 F00122 F0012A F0013 F00130 F00131 F00132 F0014 F00140 F00141 F00142 F0014A F001A F001AG F001D F002 F002 A F002 D F002* F002*A F002+ F002- F002-A F002-D F0020 F0020A F0021 F00211 F0022 F0023 F0023A F0024 F0024A F002A F008 F009 F009 * F009 A F009 D F009* F009+ F009- F009-A F009-D F009.A F0090 F00901 F0090A F0091 F00912 F0091A

		F0092 F0092A F0093 F0093A F0094 F0094A F009A F009A\ F009AGF009D F009DGF009X F009XA F00A-A F00X F00X- F010 F010* F010- F010-D F0100 F01001 F01002 F0100A F0100D F0101 F01012 F0101A F0101D F0102 F0102A F0102D F0103 F0104 F01042 F0104A F0104D F011 F011 A F011 D F011- F011-- F011-A F011-D F0110 F01100 F01101 F01102 F0110A F0111 F01111 F01112 F0111A F0112 F01120 F01121 F01122 F0113 F01131 F01132 F0114 F01141 F01142 F0114A F0114D F0117 F0119 F011A F011D F012 F012 A F012 D F012- F012-D F0120 F0120A F0121 F01211 F01232 F0124 F012A F013 F013 A F013 D F013* F013- F013-D F0130 F01301 F01302 F0130A F0131 F01310 F01312 F0133 F01330 F0134 F01340 F01341 F01342 F018 F018 A F018- F018-A F0180 F0181 F0182 F0183 F0184 F018D F019 F019 * F019 A F019 D F019* F019- F019-- F019-A F019-D F0190 F0191 F01910 F0192 F01921 F0192A F0193 F0194 F01941 F01942 F0197 F0199 F019A F019D F019N F019Z8 F01X F01X- F02. F020 F020 A F020 D F020* F020- F020-A F020-D F0200 F02001 F0200A F0201 F02012 F0202 F0203 F0203A F0204 F0204A F020A F020D F021 F021 A F021* F021- F021-A F0210 F0211 F0214 F021A F022 F022 A F022 D F022* F022- F022-A F0220 F0220A F0222 F0223 F0224 F022A F023 F023 A F023 D F023* F023+ F023- F023-A F023-D F0230 F02301 F0230A F0231 F0231A F0232 F02320 F02321 F0232A F0233 F02331 F0233A F0234 F02341 F02342 F0234A F023A F023AGF023D F023X F023XA F024 F024 A F024* F024-A F0240 F0241 F02412 F0242A F0243 F0244 F024A F028 F028 ! F028 * F028 A F028 D F028* F028+ F028- F028-A F028-D F0280 F02801 F0280A F0281 F02811 F0281A F0282 F02821 F0282A F0283 F0284 F0284A F028A F028D F028XA F029 F02X F03- F030 F0300 F03011 F0304 F03X F03X * F03X A F03X D F03X* F03X+ F03X- F03X-- F03X-A F03X-D F03X0 F03X0* F03X00 F03X01 F03X02 F03X0D F03X1 F03X11 F03X12 F03X2 F03X20 F03X2A F03X2D F03X3 F03X4 F03X41 F03X42 F03X6 F03X9 F03XD F03XG F03XI F03XS F03XZ F04X F04X- R410 R410 D R410- R410-- R4100 R4104 R4109 R410D R410L R410X R411 R411- R411X R412 R412- R413 R413- R413-- R418 R418 D R418- R418-- R4185
Functional Dependence	Z741	Z741- Z742 Z742- Z7421 Z743 Z743- Z748 Z748- Z749 Z749- Z74X Z750 Z750- Z7500 Z751 Z751- Z751-- Z751-D Z7511 Z7513 Z752 Z752- Z7520 Z753 Z753- Z754 Z754- Z7548 Z755 Z755- Z755-D Z7555 Z758 Z758- Z759 Z759- Z75X
Falls and Fractures	R55X	R55X D R55X* R55X+ R55X- R55X-- R55X-D R55X7 R55XA R55XD R55XX S320 S320 0 S320- S320-D S3200 S3200D S3201 S3202 S3205 S3206 S3209 S320D S321 S321 0 S321 D S321- S3210 S3210D S3211 S32130 S322 S322-

	S3220	S3221	S323	S323 0	S323-	S3230	S3230D	S3231
	S3236	S324	S324 0	S324-	S3240	S3240A	S3240D	S3241
	S324D	S325	S325 0	S325 D	S325-	S325-D	S3250	S3250-
	S3250A	S3250D	S3251	S3252	S3254	S3255	S3256	S3258
	S3259	S327	S327 0	S327-	S3270	S3270D	S3271	S328
	S328 0	S328-	S328-D	S3280	S3280D	S3281	S3288	S32X
	S330	S330-	S331	S331-	S331-D	S3310	S331D	S332
	S332-	S3320	S333	S333-	S3330	S3331	S333D	S334
	S334-	S3340	S335	S335-	S3350	S336	S336-	S337
	S337-	S3370	S33X	S420	S420 0	S420-	S420-A	S4200
	S4200D	S4201	S4201D	S4206	S421	S421 0	S421-	S4210
	S4210-	S4210D	S4211	S4212	S4213	S422	S422 0	S422-
	S4220	S4220-	S4220D	S4221	S42210	S4222	S4220	S423
	S423 0	S423 D	S423-	S4230	S4230D	S4231	S4231D	S4232
	S42340	S4236	S4239	S423D	S424	S424 0	S424-	S4240
	S4240D	S4241	S4241D	S4244	S4248	S4249	S427	S427-
	S4270	S4270D	S4271	S428	S428-	S4280	S4281	S429
	S429 0	S429-	S4290	S4290D	S4291	S4299	S430	S430 0
	S430-	S430--	S4300	S4302	S4309	S430D	S431	S431-
	S4310	S4316	S431D	S432	S432-	S4320	S433	S433-
	S4330	S434	S434-	S4340	S4341	S434D	S435	S435-
	S436	S436-	S436D	S437	S437-	s620	S620 0	S620-
	S6200	S6200D	S6201	S6204	S6208	S621	S621 0	S621-
	S6210	S6211	S6211D	S6218	S622	S622 0	S622-	S6220
	S6220D	S6221	S6221D	S6228	S623	S623 0	S623-	S623--
	S6230	S6230D	S6231	S6231D	S6234	S6236	S6239	S624
	S624 0	S624-	S6240	S6240D	S6241	S6241D	S6244	S625
	S626-	S627	S627 0	S6271	S6274	S628	S628 0	S628-
	S6280	S6280-	S6280D	S6281	S6288	S6289	S6280	S629
	S720	S720 0	S720-	S720-D	S720.0	S7200	S7200-	S72000
	S72009	S7200A	S7200D	S7201	S7201D	S7203	S7204	S7205
	S7208	S7209	S720A	S720D	S721	S721 0	S721-	S7210
	S72100	S7210D	S7211	S7215	S7219	S721D	S7210	S722
	S722 0	S722-	S7220	S7220D	S7221	S72210	S7221D	S7222
	S723	S723 0	S723 1	S723-	S7230	S7230D	S7231	S7236
	S723D	S724	S724 0	S724-	S7240	S7240A	S7240D	S7241
	S7246	S727	S727-	S7270	S7271	S728	S728 0	S728-
	S7280	S7280D	S7281	S728D	S729	S729 0	S729-	S7290
	S7290D	S7291	S7295	S7299	S729D	S72X	S730	S730-
	S730-D	S7300	S730D	S731	S731-	S7310	S7315	S731D
	S73X	S73X-	W000	W000-	W0009	W000A	W001	W001-
	W0010	W0012	W0019	W002	W002-	W002A	W003	W003-
	W0033	W003A	W004	W004-	W0040	W0049	W004A	W004D
	W005	W005-	W006	W006-	W007	W007-	W008	W008-
	W0080	W008A	W009	W009-	W0090	W0099	W009A	W010
	W010	AW010	DW010-	W010-A		W0100	W0101	W0103
	W0104	W0108	W0109	W010A	W011	W011-	W0111	W0118
	W0119	W011A	W012	W012-	W012--	W0120	W0122	W0123

	W0128 W0129 W012A W012X W013 W013- W0130 W0131
	W0139 W013A W014 W014- W0140 W0141 W0148 W0149
	W014A W015 W015- W0150 W0152 W0158 W0159 W015A
	W016 W016- W0160 W016A W017 W017- W018 W018-
	W0180 W0181 W0182 W0185 W0188 W0189 W018A W019
	W019- W0190 W0191 W0192 W0195 W0198 W0199 W019A
	W020 W020- W020A W021 W021- W022 W022- W023
	W023- W0230 W0239 W023A W024 W024- W024A W025
	W025- W026 W026- W027 W028 W028- W0280 W0281
	W0282 W028A W029 W029- W0290 W0291 W0293 W0299
	W029A W030 W030- W0300 W0301 W0309 W030A W031
	W031- W0319 W031A W032 W032- W0320 W0329 W032A
	W033 W033- W0330 W0331 W0333 W0339 W033A W034
	W034- W0349 W035 W035- W036 W036- W037 W037-
	W038 W038- W0380 W0383 W038A W039 W039- W0390
	W0398 W0399 W039A W040 W040- W0409 W040A W041
	W041- W0410 W0419 W042 W042- W0429 W043 W043-
	W044 W044- W045 W045- W046 W0460 W0469 W047
	W048 W048- W049 W049- W0491 W0499 W049A W050
	W050- W0504 W0509 W050A W051 W051- W0519 W051A
	W052 W052- W0528 W0529 W052A W053 W053- W054
	W054- W0549 W054A W055 W055- W056 W056- W057
	W057- W058 W058- W0581 W0589 W058A W059 W059-
	W0598 W0599 W059A W060 W060- W0600 W0601 W0604
	W0608 W0609 W060A W061 W061- W061-A W0611
	W0619 W061A W062 W062- W062-- W0624 W0628 W0629
	W062A W063 W063- W064 W064- W065 W065- W065A
	W066 W066- W067 W068 W068- W0689 W069 W069-
	W0690 W0691 W0692 W0699 W069A W070 W070- W0700
	W0701 W0706 W0708 W0709 W070A W071 W071- W0711
	W0718 W0719 W071A W072 W072- W0720 W0728 W0729
	W072A W073 W073- W074 W074- W075 W075- W0752
	W0759 W076 W076- W077 W077- W078 W078- W0782
	W079 W079- W0790 W0798 W0799 W079A W080 W080-
	W0808 W0809 W080A W081 W081- W0810 W0819 W082
	W082 AW082- W0829 W082A W083 W083- W0830 W084
	W084- W085 W085- W0850 W085A W086 W086- W0860
	W087 W087- W088 W088- W0889 W089 W089- W0899
	W089A W090 W090 AW090- W0900 W0901 W0909 W090A
	W091 W091- W092 W092- W0920 W0921 W092A W093
	W093- W0939 W093A W094 W094- W095 W095- W0959
	W095A W096 W096- W097 W097- W098 W098- W0981
	W0988 W0989 W098A W099 W099- W0990 W0991 W0999
	W099A W100 W100- W100-A W1000 W1008 W1009
	W100A W101 W101- W1011 W1012 W1019 W101A W102
	W102- W1029 W102A W103 W103 DW103- W1030 W1039
	W103A W104 W104- W1049 W105 W105- W1052 W1058
	W1059 W105A W106 W106 DW106- W1062 W107 W107-


	W108 W108- W1082 W1085 W1089 W108A W109 W109- W1090 W1098 W1099 W109A W109D W110 W110- W1100 W1103 W1109 W110A W111 W111- W1110 W112 W112 D W112- W113 W113 DW113- W113-D W1130 W1139 W114 W114- W115 W115- W116 W116- W116A W117 W117- W118 W118- W1182 W1183 W1188 W119 W119- W1191 W1192 W1193 W1198 W1199 W119A W120 W120- W120A W121 W121- W122 W122- W123 W123- W124 W124- W125 W125- W126 W126- W126A W127 W127- W128 W128- W129 W129- W1292 W1299 W129A W130 W130- W1300 W1304 W1308 W1309 W130A W131 W131- W131A W132 W132- W1329 W133 W133- W1339 W134 W134- W1349 W135 W135- W136 W136- W1360 W137 W137- W138 W138- W1389 W138A W139 W139- W1390 W1392 W1393 W1399 W139A W140 W140- W140A W141 W141- W142 W142- W143 W143- W144 W144- W1449 W145 W145- W146 W146- W147 W147- W148 W148- W1482 W148A W149 W149- W1490 W1499 W149A W150 W150- W151 W151- W152 W152- W153 W153- W1530 W154 W154- W155 W156 W156- W157 W158 W158- W159 W159- W1590 W160 W160- W161 W161- W162 W162- W163 W163- W164 W164- W165 W165- W166 W166- W167 W167- W168 W168- W169 W169- W170 W170- W1700 W1701 W1708 W1709 W170A W171 W171- W172 W172- W1720 W1729 W172A W173 W173- W1730 W1739 W173A W174 W174- W1740 W1749 W174A W175 W175- W1752 W175A W176 W176- W1762 W1769 W176A W177 W177- W178 W178- W1780 W1781 W1782 W1789 W178A W179 W179- W1790 W1791 W1792 W1798 W1799 W179A W180 W180- W180-A W1800 W1801 W1802 W1803 W1804 W1808 W1809 W180A W180E W181 W181- W1810 W1811 W1819 W181A W181D W182 W182- W182-- W1820 W1821 W1822 W1828 W1829 W182A W183 W183- W1830 W1831 W1839 W183A W184 W184- W1840 W1848 W1849 W184A W185 W185- W1851 W1858 W1859 W185A W186 W186- W1869 W187 W187- W1879 W188 W188- W1880 W1881 W1882 W1883 W1888 W1889 W188A W189 W189- W1890 W1891 W1892 W1893 W1894 W1895 W1898 W1899 W189A W190 W190 AW190 DW190- W190-- W190-A W190-D W1900 W1901 W1903 W1905 W1908 W1909 W190A W191 W191- W191-A W1910 W1911 W1918 W1919 W191A W192 W192 DW192+ W192- W192-- W192-A W1921 W1922 W1928 W1929 W192A W193 W193- W1930 W1939 W194 W194* W194- W1940 W1941 W1943 W1948 W1949 W194A W195 W195- W1959 W195A W196 W196- W196A W197 W197- W197A W198 W198- W198-A W1980 W1981 W1982 W1988 W1989 W198A W199 W199 0 W199 DW199- W199-A W199-D W1990 W1991
--	--

	W1992 W1993 W1994 W1995 W1996 W1998 W1999 W199A W199D W19X
Incontinence	R15X R15X A R15X D R15X- R15X-- R15X9 R32X R32X- R32X-- R32X-A R32X-D R32X0 R32X1 R32X3 R32X9 R32XD
Mobility problems	R260 R260- R260D R261 R261- R261D R262 R262 A R262- R2621 R2623 R263 R263- R263D R268 R268- R268-- R2683 R2686 R2689 R268D R269 Z740 Z740 Z Z740- Z740-- Z740-D Z740. Z7400 Z7401 Z7404 Z740C Z740D
Pressure Ulcers	L890 L890- L890-- L890D L891 L891- L891-- L892 L892- L892-- L893 L893- L893-A L899 L899 A L899- L899-- L89X L89X - L89X A L89X D L89X E L89X I L89X J L89X Z L89X- L89X-- L89X-D L89X1 L89X5 L89X9 L89XD
Senility	R54X R54X A R54X D R54X- R54X-D R54X. R54X0 R54X6 R54X7 R54X9 R54XA R54XD R54XI R54XW R54XX

APPENDIX 2:

Anxiety & Depression Coding Prevalence Over Time


All episodes at acute providers, Jan '05 to Mar '13



■ F329 ■ F329- ■ F419 ■ F410 ■ F412 ■ F339 ■ F419- ■ F410-
 ■ F412- ■ F339- ■ F411 ■ F322 ■ F432 ■ F323 ■ F448 ■ F431
 ■ F411- ■ F445 ■ F328 ■ F322- ■ F323- ■ F333 ■ F432- ■ F320-
 ■ F320 ■ F321 ■ F431- ■ F430 ■ F439 ■ F449 ■ F418 ■ F332
 ■ F321- ■ F444 ■ F332- ■ F333- ■ F449- ■ F448- ■ F430- ■ Other


Delirium Coding Prevalence Over Time

All episodes at acute providers, Jan '05 to Mar '13




■ F059 ■ F051 ■ F059- ■ F050 ■ F058 ■ F051- ■ F050-
 ■ F058- ■ F051 A ■ F0513 ■ F051-D ■ F051 D ■ F051-A ■ F059--
 ■ F051D ■ F050 A ■ F058-- ■ F059 D ■ Other

Dementia Coding Prevalence Over Time
All episodes at acute providers, Jan '05 to Mar '13



F03X R410 F03X- R410- F019 R418 F011
 F009 A F019- F009 F009A R413 F011- R418-
 R413- F03X0 F009-A R412 F023 F023 A F028
 F028 A F023A F028A R412- F001 A F0190 F001
 F03X4 F023-A F001A F018 R411 F028-A F0194
 F010 F0110 R411- F002 A Other


Functional Dependence Coding Prevalence Over Time
All episodes at acute providers, Jan '05 to Mar '13



Z751 Z755 Z751- Z755- Z742 Z748 Z749
 Z752 Z753 Z741 Z758 Z754 Z742- Z748-
 Z743 Z749- Z754- Z741- Z750 Z752- Z743-
 Z759 Z758- Z753- Z750- Z759- Z7513 Z7511
 Z7500 Z7520 Z755-D Z751-- Z7421 Other

Falls (& significant fracture) Coding Prevalence Over Time


All episodes at acute providers, Jan '05 to Mar '13



R55X	W199	S7200	R55X-	W190	W010	S7210
S720	W100	W199-	W180	W019	S3250	S4220
W190-	W014	W010-	W191	W060	W192	W109
W189	S4240	S4200	S6230	W180-	S4230	S3200
W100-	W011	S430	W012	W019-	W070	S7230
W018	S721	S7240	W194	Other		


Incontinence Coding Prevalence Over Time

All episodes at acute providers, Jan '05 to Mar '13




R32X	R15X	R32X-	R15X-	R15X D
R32XD	R15X A	R15X9	R32X9	R32X--
R15X--	R32X-A	R32X-D	Other	

Mobility Problems Coding Prevalence Over Time
All episodes at acute providers, Jan '05 to Mar '13




Z740	R268	R262	Z740-	R268-	R262-
R260	R260-	R261	R261-	R260D	Z7400
Z740C	R2686	R261D	R2621	Z740D	R2689
R2683	R2623	Z7401	Z740 Z	R263	R263-
Z740--	Z740.	R263D	R268--	R262 A	Other

Senility Problems Coding Prevalence Over Time
All episodes at acute providers, Jan '05 to Mar '13




R54X	R54X-	R54XD	R54X D	R54XW
R54X0	R54XA	R54X.	R54XX	R54XI
R54X6	R54X A	R54X9	R54X-D	Other



Pressure Ulcers Coding Prevalence Over Time All episodes at acute providers, Jan '05 to Mar '13







- L89X
- L899
- L893-
- L890
- L891
- L89X A
- L89XD
- L893
- L89X D
- L899-
- L890-
- L891-
- L893-
- L892
- L89X--
- L899-
- L892-
- L890-
- L899--
- L891--
- L890--
- L893-A
- L899 A
- L890D
- L892--
- Other


Appendix 3











Appendix 4

<http://www.ons.gov.uk/ons/taxonomy/index.html?nscl=Population+Estimates>

ENGLAND	2005	2013	% change
All ages	50606000	53865817	6.44%
65-74	4189100	5023573	19.92%
75-84	2855100	3043739	6.61%
85+	986800	1237867	25.44%

% of population

ENGLAND	2005	2013	% change
Denominator, all ages	50606000	53865817	6.44%
65-74	8.28%	9.33%	12.66%
75-84	5.64%	5.65%	0.16%
85+	1.95%	2.30%	17.85%

% of >65yo population

ENGLAND	2005	2013	% change
Denominator, o65s	8031000	9305179	15.87%
65-74	52%	54%	3.50%
75-84	36%	33%	-7.99%
85+	12%	13%	8.27%