

# BMJ Open

## The influence of time spent at home on the association between residential food environment and fruit and vegetable intake

Journal:	BMJ Open
Manuscript ID:	bmjopen-2014-006200
Article Type:	Research
Date Submitted by the Author:	23-Jul-2014
Complete List of Authors:	Chum, Antony; St Michael's Hospital, Centre for Research on Inner City Health Farrell, Eddie; St Michael's Hospital, Centre for Research on Inner City Health Labetski, Anna; St Michael's Hospital, Centre for Research on Inner City Health Vaivada, Tyler; McMaster University, Faculty of Health Sciences Bohnert, Arianne; St Michael's Hospital, Centre for Research on Inner City Health Selvaratnam, Inthuja; St Michael's Hospital, Centre for Research on Inner City Health Pinter, Theresa; University of Toronto, Department of Geography Larsen, Kristian; Toronto Public Health, O'Campo, Patricia; St Michael's Hospital, Centre for Research on Inner City Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Epidemiology, Research methods
Keywords:	PUBLIC HEALTH, PREVENTIVE MEDICINE, EPIDEMIOLOGY

SCHOLARONE™  
Manuscripts

# The influence of time spent at home on the association between residential food environment and fruit and vegetable intake

Corresponding author: Antony Chum,  
St Michael's Hospital  
Centre for Research on Inner City Health  
Li Ka Shing Knowledge Institute  
209 Victoria Street, third floor, Toronto, Ontario M5C 1N8

email: [antony.chum@utoronto.ca](mailto:antony.chum@utoronto.ca)

Authors:

- Antony Chum, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada
- Eddie Farrell, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada
- Anna Labetski, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada
- Tyler Vaivada, McMaster University, Toronto Canada
- Arianne Bohnert, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada
- Inthuja Selvaratnam, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada
- Theresa Pinter, University of Toronto, Toronto Canada
- Kristian Larsen, Centre for Research on Inner City Health, University of Toronto, Toronto Canada
- Patricia O'Campo, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada

Word count: 3,105

## ABSTRACT

**Objective:** There is a growing body of research that investigates the residential neighbourhood context as it relates to individual diet. However, previous studies ignore participants' time spent in the residential environment and this may be a problem because time-use studies show that adults' time-use can significantly vary. To better understand the role of exposure duration, we designed a study to examine 'time spent at home' as a moderator to the residential food environment-diet association.

**Design:** Cross-sectional observational study

**Settings:** City of Toronto, Ontario, Canada

**Participants:** 2411 adults aged 25-65

**Primary Outcome Measure:** Frequency of vegetable and fruit intake (VFI) per day

**Results:** To examine how time spent at home may moderate the relationship between residential food environment and VFI, the full sample was split into 3 equal subgroups - short, medium, and long duration spent at home. We detected significant associations between density of food stores in the residential food environment and VFI for subgroups that spend medium and long durations at home (i.e. spending a mean of 8.0 and 12.3 hours at home respectively – no including sleep time), but no associations exist for people that spend the lowest amount of time at home (mean=4.7 hours). Also, no associations were detected in analyses using the full sample.

**Conclusions:** Our study is the first to demonstrate that residential exposure duration may be an important variable to identify hidden population patterns regarding VFI. Time spent at home can impact the association between the residential food environment and individual VFI.

### Strengths:

- our study extends the body of work on the effects of residential food environments on diet by looking at the amount of time that participants are spending in their residential environment
- previous studies ignore participants' time spent in the residential environment and this may be a problem because time-use studies show that adults' time-use can significantly vary

### Limitations:

- cross-sectional observational data limits the study's ability to discern the direction of causation
- our outcome measure, vegetable and fruit intake, is based on the Canadian Community Health Survey 2010, and participants self-reported the frequency per day of fruits and vegetable eaten rather than the number of servings consumed. The self-reported frequency measure may contribute to both under and over-reporting of food intake behaviour.

INTRODUCTION

Low vegetable and fruit intake (VFI) has been linked to a number of chronic diseases including type II diabetes,[1 2] cardiovascular disease,[3 4] and breast,[5 6] esophageal and colon cancers,[7 8]. Individual level determinants of VFI have been well established in the literature, where income and education are positively associated with VFI;[9-11] however, study results of the association between VFI and its potential environmental determinants are decidedly mixed. Research on the food environment has explored the impacts of food retailers on VFI (e.g. supermarkets, fast food outlets, and convenience stores). Studies have shown that living in proximity to supermarkets is associated with improved diet outcomes,[12-19] and poor diet outcomes.[20 21] Studies also show no association between residential proximity to food vendors and VFI.[22 23]

Along with the inconsistent findings described above, the research is also characterized by a lack of consideration to individuals' duration of exposure to their neighbourhood context. By ignoring the temporal dimension of exposure, previous studies may have unintentionally introduced measurement bias because exposure duration may significantly differ between participants. There is a dearth of studies that have explored this problem using multilevel analyses of neighbourhood effects on individual health outcomes. Chum and O'Campo (2013) found that the use of time-weighted multilevel regressions to account for duration of exposure resulted in a) improved strength of association and 2) improved model fit in models of the association between neighbourhood-level factors (including road traffic, access to supermarkets, and fast food restaurants) and cardiovascular disease risk compared to typical multilevel models that do not account for time.[24] There is also evidence to suggest that time spent in the residential neighbourhood varies. According to the Canadian General Social Survey (2010) public use microdata,[25] time spent at home differs significantly by age and income: analysis of variance shows that total minutes spent at home on a typical weekday differs significantly by age groups and income groups ( $p<0.01$ ) - see Figure 1.

Figure 1: Mean duration (in minutes) at home on an average weekday by age and income groups (Canadian General Social Survey 2010) weighted N=28,075,610

Given the variability of time spent at home between individuals, accounting for individuals' time spent in their residential environment may help us avoid model misclassification by better

quantifying exposure to the residential food environment. We hypothesize that people who spend more time in their home environments would rely more heavily on their local food vendors. Therefore, a stronger association between residential food environment and VFI may exist for those who spend more time at home compared to those who spend less time at home. Our study answers the following research questions to explore this potential dose-response relationship:

- 1) Does an association exist between the residential food environment and VFI?
- 2) Is the strength of the association between the residential food environment and VFI stronger for people who spend more time at home?

## MATERIALS AND METHODS

### *Data Sources*

Project NEHW (Neighbourhood effects on Health and Wellbeing) is a cross-sectional investigation of neighbourhood-level determinants of population health that used a three-staged sampling method. In the first stage, 50 city-delineated neighbourhood planning areas (NPA) were randomly selected out of a total of 140 NPAs. In the second stage, 1-2 census tracts (CT) were randomly selected from each of the 50 NPAs sampled, resulting in 87 randomly selected CTs. Lastly, within the 87 CTs, approximately 30 individuals were randomly selected based on residential address. Eligibility criteria are as follows: 1) only 1 resident per household, 2) participants are aged 25 to 65, 3) able to communicate in English, and 4) lived in the neighbourhood for at least 6 months. The response rate was 72%.

Data collection took place between March 2009 and June 2011. 2411 individuals, representing 87 CTs, participated in the study. Data were obtained from in-person interviews, and participants provided written informed consent at the time of their interview. The Research Ethics Board at St Michael's Hospital in Toronto, Canada provided ethics approval for this study. To ensure the generalizability of our findings, and that it is representative of our target population, post-stratification weights were created based on demographic characteristics from the 2006 Canadian Census data for Toronto including sex, total household income, household size, immigrant status, and age. More information about the study methods can be found in a previously published paper.[24]

**Dependent Variable**

The primary outcome measure is frequency of VFI, and was assessed using questions from the U.S. Behavioural Risk Factor Surveillance System.[26] The same questions are also found in the Canadian Community Health Survey (2010).[27] Six questions, similar to questions in a food frequency questionnaire, were asked to determine total frequency of intake for vegetables and fruits.[28 29] Intake was calculated by adding the frequency of intake of fruit, fruit juice, green salad, carrots, potatoes (not including French fries, fried potatoes or potato chips), and other vegetables. VFI is based on self-reported data. Since VFI is non-normally distributed with a positive skew, for the purposes of regression analysis, we categorized VFI into either “less than five times” or “greater than or equal to five times” per day. These categories of VFI have been used in previous studies.[28 29] Frequency of intake was structured around the recommendations of Canada’s Food Guide whereby one instance of VFI was considered to be one serving, and less than 5 servings a day is below recommendation.[30] While frequency of intake may not necessarily equal the number of servings, for the purpose of this study it is treated roughly in the same manner, following the convention of previous studies.[28 29]

**Independent Variables**

Toronto Public Health’s Toronto Healthy Environments Inspection System (THEIS) (2012),[31] provided the location of all food outlets. Within the THEIS database, the type of food retailer was used to examine food access in residential environments. Fast food, healthy food retailers, and less healthy food stores were used in this study. *Fast food restaurants* were classified as a restaurant, food court vendor, cafeteria, food take out, ice cream/yogurt vendor, or hot dog cart. These restaurants were also required to have takeout options available, limited or no wait staff, and customers having to pay prior to receiving food. If the restaurant had restricted access to the public they were removed (e.g. food outlets in arenas where patrons must pay admission to access the food outlet).

*Healthier food retail* was classified as a food store (convenience/variety or supermarket) that sells a significant quantity and diversity of vegetables and/or fruit. If these food outlets did not meet the definition for healthy food retail, they were classified as a *less healthy food store*.

These are typically convenience stores that primarily sell packaged snacks with low nutritional value. To account for retail locations outside of the City of Toronto in bordering municipalities, additional field work was completed to ensure all food outlets within 1km of the residential addresses were included in the analysis.

Geographic Information Systems (GIS) was used to characterize the food environment of our study area. Our measure of residential food outlet availability was created based on a review of GIS methods to measure food environments,[32] where density of food stores within a buffer around an individual's residence (constructed based on 10-15 minutes walking distances) is a commonly accepted measure. In a field study of walking speeds of the general adult population using mobile accelerometers,[33] the median speed of adult walking speed is 1.25 metres/second (interquartile range=0.12), and speed declines with age at the rate of -0.0037 metres/second per year between the ages of 20-60. However, speed was not significantly associated with other individual characteristics including gender, BMI, and blood pressure. Based on these results, we estimated buffers for distances that would be reachable in a 10 and 15 minute walk from the residence of each participant. The buffers created for this study were based on network service areas around each respondents' homes. Network service area is the travelable distance created using street network data (Figure 2).[34-36] Densities for 10 and 15 minute walking distance buffers were calculated in GIS for the number of fast food, less healthy, and healthy food outlets in each of the participants' walking buffers.

Figure 2: Example of a network buffer for 10 and 15 minute walk

### ***Covariates***

A number of individual level covariates are included in the models to adjust for potential confounding. The following categorical variables were examined: age (25-34, 35-44, 45-54, or 55-65), gender, marital status (married/common-law, or single), education (high school or less, some college, completed college), self-rated health (poor, fair, good, and very good/excellent), and family income (quintiles). While ethnicity/race is examined in bivariate analysis (i.e. white, black, south Asian, south-east Asian, West Asian, or other), only visible minority status (i.e. white vs. others) was used instead of more detailed ethnicity categories in multivariable models due to cell size limitations.



*Time Spent at Home*

Individual level data collected from Project NEWH participants provided self-reported time spent at work and time spent sleeping, but not time spent at home. The average of time spent at home per day was estimated based on individual demographic characteristics and time spent at work using the 2010 Canadian General Social Survey (CGSS) through a multivariate regression model. Data were extracted from the CGSS for adults age 25–65 in urban areas. Time spent at home (dependent variable) was modeled using the following independent variables: age, education level, income, gender, marital status, having children under 5, and minutes spent at work. All the above predictors were significantly associated with the time spent at home ( $p < 0.01$ ). The final model had an adjusted r-squared of 0.40. Beta coefficients from the regression using the CGSS data were used to estimate individual time spent at home for the Project NEWH participants. For example, starting with the intercept of 1325.15 minutes, 1) for every minute spent at work 0.68 fewer minutes were spent at home, 2) females spent 37.34 more minutes at home compared to males, 3) persons with children under 6 spent 30.93 more minutes compared to those without, et cetera. Finally, we subtracted out individuals' sleeping duration from the total time at home because individuals have no chance of contact with their residential food environment while they are sleeping.

*Statistical analysis*

This analysis started by evaluating the bivariable relationships between predictors and the VFI outcome using 1) Kruskal-Wallis non-parametric 1-way ANOVA to compare the mean VFI per day between the categorical predictors, and 2)  $\chi^2$  test to compare the categorical predictors to the proportion of individuals that ate fruits and vegetables at least 5 times per day (see table 1 of appendix). For multivariate multilevel modeling, we used the binary outcome (VFI at least 5 times per day vs. not), because 1) the positive skew in the continuous outcome can impact model stability and 2) this binary outcome has been used in previous models of the impact of individuals factors on VFI.[28 29]

Since the data has a 2-level structure, with individuals nested in census tracts, multilevel logistic regression is used to account for the lack of spatial independence.[37] To examine how



time spent at home may moderate the relationship between residential food environment and VFI, the full sample was split into 3 equal subgroups - short, medium, and long duration spent at home, for separate analysis. Eight models are described in this paper. Models 1a and 1b examine the full sample (n=2411). Models 2a and 2b examine the subgroup (n=804) that spends the least amount of time at home (0-6.5 hours/day, mean=4.7 hours), 3a and 3b examine the subgroup (n=804) that spends a medium amount of time at home (6.51-9.7 hours/day, mean=8.0 hours), and 4a and 4b examine the subgroup (n=803) that spends the highest amount of time at home (>9.7 hours/day, mean=12.3 hours). Models with the “a” suffix calculates food outlet density using the 10 minute walking distance network buffer, and models with the “b” suffix uses the 15 minutes walking distance network buffer. All models adjusted for the effects of gender, age, education, marital status, family income, self-rated health, and visible minority status. The binomial outcome of “eating at least 5 fruits or vegetables” is modeled using multilevel logistic regression with random intercept using PROC GLIMMIX in SAS 9.3.

## RESULTS

Participants in Project NEHW were 53% female and 44% self-reported as visible minority, with a mean age of 44 years (table 1 of appendix) and a mean after-tax family income of \$91,330 (median=\$71,000). The proportion of those eating at least 5 VFI per day differed significantly by gender, age, education, employment status, marital status and income in bivariable analysis ( $p < 0.05$ ).

**Table 1. Sample Characteristics and associations between covariates and VFI (n=2411)**

n=2,411	n (proportion percentage)	Mean of Fruit and Vegetable Intake per day	Kruskal-Wallis 1-way ANOVA p-value	Proportion eating 5 or more Fruits and Vegetables per day	chi square p-value
<b>Gender</b>			<b>&lt;0.0001</b>		<b>&lt;0.0001</b>
Male	1,118 (46.4%)	4.20		<b>25.6%</b>	
Female	1,293 (53.6%)	4.59		<b>34.6%</b>	
<b>Age</b>			<b>0.424</b>		<b>0.0001</b>
25-34	529 (21.9%)	4.24		<b>29.4%</b>	

35-44	742 (30.8%)	4.42		27.9%	
45-54	714 (29.6%)	4.42		29.6%	
55-65	426 (17.7%)	4.52		37.4%	
<b>Education</b>			<b>&lt;0.0001</b>		<b>&lt;0.0001</b>
Less than High School	95 (3.9%)	4.52		31.5%	
High School	389 (16.2%)	3.9		23.9%	
Post-Secondary Degree/Diploma	1,409 (58.5%)	4.33		30.0%	
Graduate Degree	514 (21.4%)	4.66		35.9%	
<b>Employment Status</b>			<b>0.0003</b>		<b>0.0104</b>
Employed	1,618 (67.1%)	4.25		28.8%	
On Temporary Leave	352 (14.6%)	4.54		34.0%	
Unemployed	228 (9.5%)	4.84		39.1%	
Not looking for work	212 (8.8%)	4.25		27.5%	
<b>Marital Status</b>			<b>0.0223</b>		<b>&lt;0.0001</b>
Married/Common law	1,542 (63.9%)	4.42		32.6%	
Separated/Divorced	328 (13.6%)	4.27		26.7%	
Widowed	42 (1.7%)	5.17		52.0%	
Never Married	500 (20.7%)	4.16		27.7%	
<b>Income</b>			<b>0.0058</b>		<b>0.0002</b>
\$39,000 or less	697 (28.9%)	4.13		25.1%	
\$40,000 to \$70,999	584 (24.2%)	4.34		30.7%	
\$71,000 to \$109,999	523 (21.7%)	4.42		32.8%	
\$110,000 or more	606 (25.1%)	4.53		34.3%	
<b>Ethnicity</b>			<b>0.0015</b>		<b>0.0021</b>
White	1349 (56.0%)	4.49		32.8%	
Black	339 (14.1%)	4.24		30.2%	

East Asian	221 (9.2%)	4.29		<b>27.3%</b>	
South Asian	224 (9.3%)	4.03		<b>23.4%</b>	
Aboriginal	18 (0.7%)	4.54		<b>42.3%</b>	
West Asian	52 (2.2%)	4.26		<b>26.1%</b>	
Latin	130 (5.4%)	3.68		<b>21.9%</b>	
Other	77 (3.2%)	4.50		<b>34.9%</b>	

All models to follow have been adjusted for the effects of gender, age, education, self-rated health, marital status, visible minority status, and family income. For models where we disregarded time spent at home, no significant associations were found between the food environment and VFI. This was true for food store density measured at scales of both 10 minutes and 15 minutes walking distances (see Table 2 - model 1a and 1b). Next, we conducted subgroup analyses where we divided the full sample of participants into three equal groups with different amounts time spent at home (low, medium and high). For individuals that spent the least amount of time at home (models 2a & 2b), no significant associations were found between the food environment variables and VFI.

For individuals who spent a medium amount of time at home, model results differed by buffer size. At the 10 minute buffer, residents with zero *less healthy* (see chart) food stores compared to those with three or more, had 38% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 3a). Within the 15 minute buffer, individuals with three or more *healthier* food stores compared to those with zero, had 36% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 3b).

For individuals that spent the highest amount of time at home, model results also differed by buffer size. At the 10 minute buffer, residents with access to three or more *healthier* food stores compared to those with none, had 29% increased odds ( $p<0.05$ ) of having five or more VFIs per day. In addition, residents with zero *less healthy* food stores compared with three or more, had 38% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 4a).

Within the 15 minute buffer, residents with 'two *healthy* food stores' and 'three or more

healthy food stores’ compared with zero, both had 61% increased odds ( $p<0.05$ ) of having five or more VFIs per day. In addition, residents with access to zero or one *fast food* stores compared with three or more, had 28% and 27% increased odds ( $p<0.05$ ) of having five or more VFIs per day respectively (model 4b).

**Table 2. Multilevel logistic regression to examine the association between food environment (10 and 15 minute walking distance network buffers) and the odds of eating 5 vegetables and fruits per day adjusting for gender, age, education, self-rated health, marital status, visible minority status, and family income.**

	Full sample (n=2411)		Subgroup A: spending a mean of 4.7 hours at home (n=804)		Subgroup B: spending a mean of 8 hours at home (n=804)		Subgroup C: spending a mean of 12.3 hours at home (n=803)	
	Model 1a <sup>+</sup>	Model 1b <sup>+</sup>	Model 2a <sup>+</sup>	Model 2b <sup>+</sup>	Model 3a <sup>+</sup>	Model 3b <sup>+</sup>	Model 4a <sup>+</sup>	Model 4b <sup>+</sup>
	Odds ratios (95% confidence interval)							
<b>Healthier (ref=no stores)</b>								
1 store	0.843 (0.494-1.438)	1.430 (0.955-2.141)	0.964 (0.553-1.681)	1.544 (0.948-2.515)	0.788 (0.535-1.161)	0.888 (0.654-1.206)	0.825 (0.596-1.141)	0.972 (0.563-1.679)
2 stores	1.042 (0.819-1.326)	1.313 (0.867-1.988)	0.729 (0.477-1.113)	1.285 (0.970-1.701)	1.083 (0.732-1.601)	1.172 (0.943-1.457)	1.159 (0.765-1.757)	<b>1.606 (1.082-2.385)*</b>
3 or more stores	1.249 (0.796-1.958)	1.184 (0.942-1.489)	1.109 (0.773-1.591)	0.774 (0.354-1.691)	1.139 (0.908-1.917)	<b>1.362 (1.030-1.801)*</b>	<b>1.291 (1.048-1.590)*</b>	<b>1.614 (1.108-2.352)*</b>
<b>Less Healthy (ref=3+ stores)</b>								
0 stores	1.045 (0.710-1.537)	1.045 (0.710-1.537)	1.019 (0.711-1.459)	1.012 (0.690-1.483)	<b>1.377 (1.041-1.823)*</b>	1.014 (0.709-1.451)	<b>1.381 (1.043-1.829)*</b>	1.025 (0.716-1.467)
1 store	1.043 (0.621-1.750)	0.985 (0.576-1.685)	1.003 (0.610-1.649)	0.964 (0.566-1.644)	1.192 (0.958-1.483)	0.985 (0.600-1.616)	1.191 (0.956-1.484)	0.990 (0.604-1.625)
2 stores	0.985 (0.576-1.685)	1.043 (0.621-1.750)	1.000 (0.608-1.646)	1.024 (0.612-1.714)	0.895 (0.659-1.217)	1.003 (0.611-1.648)	0.895 (0.659-1.218)	1.015 (0.617-1.668)
<b>Fast food (ref=3+ stores)</b>								
0 stores	1.082 (0.731-1.601)	1.133 (0.665-1.931)	1.175 (0.810-1.703)	1.658 (0.930-2.955)	1.365 (0.940-1.982)	1.175 (0.810-1.703)	1.227 (0.862-1.745)	<b>1.281 (1.037-1.583)*</b>
1 store	1.070 (0.796-1.439)	1.124 (0.811-1.558)	1.126 (0.648-1.957)	1.299 (0.979-1.723)	1.380 (0.862-2.209)	1.126 (0.648-1.957)	1.235 (0.760-2.005)	<b>1.267 (1.027-1.563)*</b>
2 stores	0.865 (0.670-1.116)	0.970 (0.679-1.386)	0.874 (0.506-1.509)	1.003 (0.538-1.870)	0.864 (0.514-1.453)	0.874 (0.506-1.509)	0.763 (0.466-1.251)	0.992 (0.745-1.321)

\* $p<0.05$

<sup>+</sup>The suffix ‘a’ label denotes food outlet densities calculated within 10 minutes walking distance, and the suffix ‘b’ denotes food outlet densities calculated within 15 minutes walking distance.

## DISCUSSION

In summary, the association between density of food stores in the residential food environment and VFI exists for subgroups that spend medium and long durations at home (i.e. at least 6.5 hours in addition to time spent sleeping), but no associations exist for people that spend the lowest amount of time at home. Also, no associations were detected in analyses using the full sample. A plausible explanation for these observations is that people who spend more time at home tends to make use of their local residential food outlets, while those who spend little time at home may purchase food elsewhere as they spend their day in other locations. This is an important finding since no previous studies have differentiated participants by duration of time spent at home, and our study is the first to demonstrate that residential exposure duration may be an important missing variable to identify hidden population patterns. In this study, we show that adult time use can significantly vary across individuals and is a factor that can modify the food environment-VFI association. Given that there are no other similar studies of VFI that have accounted for time use, we cannot meaningfully compare our results to the associations found in other studies at this point. Our study highlights the importance of understanding the duration of residential exposure, and this has implications for future data collection in the context of multilevel research of environmental effects on health.

Furthermore, our study illustrates that extending the network buffer from 10 minute to 15 minute walking distance has the effect of changing which predictors are significantly associated with VFI. Further research needs to be done on optimum buffer sizes to investigate access to the food environment. It should be noted that we did not have data on car ownership, and thus cannot ascertain the mode of transportation taken to their local food store.

This study has a number of limitations. First, quality and pricing data for supermarkets were not collected, both of which can affect customer shopping habits. For example, individuals may be within close proximity of a supermarket but may be unable to afford the groceries, or the quality of the fruits and vegetables may be poor. Second, the VFI outcome variable may be subject to self-reporting and social desirability bias.[38] Our questions regarding VFI is based on the Canadian Community Health Survey 2010, and participants of our study reported the frequency per day of fruits and vegetable eaten rather than the number of servings consumed.

The self-reported frequency measure may contribute to both under and over-reporting of food intake behaviour. As such, it is difficult to determine the actual VFI and make comparisons to Canada’s Food Guide.[28] Third, health-selected migration can occur when healthy individuals are attracted to healthier areas. Similarly, businesses may be more inclined to target neighbourhoods where people are perceived as living healthier lifestyles.[39] Thus, there is a possibility for reverse-causation through the above processes. Fourth, this study is based on a cross-sectional observational design and direct causation for the observed associations cannot be verified except through future longitudinal studies. Fifth, there may be residential confounding that we did not consider in our study, beyond what could be captured by self-rated health, such as the presence of specific medical conditions that may impact VFI, individual mobility issues, and dietary preferences.

Future research on the effects of the neighbourhood food environment on diet should pay greater attention to adult time use. As noted in figure 1, older adults and those with lower household income tend to spend more time at home, and thus they may be more vulnerable to deficits in their residential environments. On the other hand, for adults that spend little time in the home environment (i.e. under 6.5 hours per day not including sleep time), their residential neighbourhood may have little relevance on their diet outcomes. Finally, future research should also consider the effects of the non-residential food environment in addition to the residential environment.

**CONTRIBUTORSHIP STATEMENT**

AC drafted the manuscript and conducted all the analysis, EF conducted geospatial and statistical analysis and compiled the data, AL performed the GIS processing and created the tables for geospatial analysis, TV collected and sorted the time use data and drafted the methods section, AB, IS, and TP drafted the literature and discussion components of the manuscript, KL compiled the grocery stores data, PO designed the original study and obtained the grant through CIHR for funding. AC, EF, AL, TV, AB, IS, TP, KL, and PN interpreted the results and approved the manuscript.

**COMPETING INTERESTS**

There are no competing interests.

**FUNDING**

This study is supported by funding from the Government of Canada: Social Sciences and Humanities Research Council of Canada (SSHRC) and Canadian Institutes of Health Research (CIHR)

**DATA SHARING AGREEMENT**

No additional data are available.

## REFERENCES

1. Cooper AJ, Forouhi NG, Ye Z, et al. Fruit and vegetable intake and type 2 diabetes: EPIC-InterAct prospective study and meta-analysis. *European journal of clinical nutrition* 2012;**66**(10):1082-92
2. Carter P, Gray LJ, Troughton J, Khunti K, Davies MJ. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ: British Medical Journal* 2010;**341**
3. Bazzano LA, He J, Ogden LG, et al. Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *The American journal of clinical nutrition* 2002;**76**(1):93-99
4. Joshipura KJ, Hu FB, Manson JE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Annals of internal medicine* 2001;**134**(12):1106-14
5. Zhang CX, Ho SC, Chen YM, Fu JH, Cheng SZ, Lin FY. Greater vegetable and fruit intake is associated with a lower risk of breast cancer among Chinese women. *International journal of cancer* 2009;**125**(1):181-88
6. Jung S, Spiegelman D, Baglietto L, et al. Fruit and vegetable intake and risk of breast cancer by hormone receptor status. *Journal of the National Cancer Institute* 2013;**105**(3):219-36
7. Yamaji T, Inoue M, Sasazuki S, et al. Fruit and vegetable consumption and squamous cell carcinoma of the esophagus in Japan: the JPHC study. *International Journal of Cancer* 2008;**123**(8):1935-40
8. Aune D, Lau R, Chan DS, et al. Nonlinear reduction in risk for colorectal cancer by fruit and vegetable intake based on meta-analysis of prospective studies. *Gastroenterology* 2011;**141**(1):106-18
9. Dehghan M, Akhtar-Danesh N, Merchant A. Factors associated with fruit and vegetable consumption among adults. *Journal of Human Nutrition and Dietetics* 2011;**24**(2):128-34
10. Ding D, Sallis JF, Norman GJ, et al. Community food environment, home food environment, and fruit and vegetable intake of children and adolescents. *Journal of nutrition education and behavior* 2012;**44**(6):634-38
11. Subar AF, Heimendinger J, Patterson BH, Krebs-Smith SM, Pivonka E, Kessler R. Fruit and vegetable intake in the United States: the baseline survey of the Five A Day for Better Health Program. *American Journal of Health Promotion* 1995;**9**(5):352-60
12. Giskes K, van Lenthe F, Kamphuis C, Huisman M, Brug J, Mackenbach JP. Household and food shopping environments: do they play a role in socioeconomic inequalities in fruit and vegetable consumption? A multilevel study among Dutch adults. *Journal of epidemiology and community health* 2009;**63**(2):113-20
13. Laraia BA, Siega-Riz AM, Kaufman JS, Jones SJ. Proximity of supermarkets is positively associated with diet quality index for pregnancy. *Preventive medicine* 2004;**39**(5):869-75
14. Moore LV, Roux AVD, Nettleton JA, Jacobs DR. Associations of the Local Food Environment with Diet Quality—A Comparison of Assessments based on Surveys and Geographic Information Systems The Multi-Ethnic Study of Atherosclerosis. *American Journal of Epidemiology* 2008;**167**(8):917-24
15. Morland K, Wing S, Roux AD. The contextual effect of the local food environment on residents' diets: the atherosclerosis risk in communities study. *Journal Information* 2002;**92**(11)
16. Rasmussen M, Krølner R, Klepp K-I, et al. Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: quantitative studies. *International Journal of Behavioral Nutrition and Physical Activity* 2006;**3**(1):22



17. Rose D, Richards R. Food store access and household fruit and vegetable use among participants in the US Food Stamp Program. *Public health nutrition* 2004;**7**(08):1081-88

18. Wrigley N, Warm D, Margetts B, Whelan A. Assessing the impact of improved retail access on diet in a 'food desert': a preliminary report. *Urban Studies* 2002;**39**(11):2061-82

19. Zenk SN, Schulz AJ, Hollis-Neely T, et al. Fruit and vegetable intake in African Americans: income and store characteristics. *American journal of preventive medicine* 2005;**29**(1):1-9

20. Gustafson AA, Sharkey J, Samuel-Hodge CD, et al. Perceived and objective measures of the food store environment and the association with weight and diet among low-income women in North Carolina. *Public Health Nutrition* 2011;**14**(6):1032

21. Timperio A, Ball K, Roberts R, Campbell K, Andrianopoulos N, Crawford D. Children's fruit and vegetable intake: associations with the neighbourhood food environment. *Preventive medicine* 2008;**46**(4):331-35

22. Pearson T, Russell J, Campbell MJ, Barker ME. Do 'food deserts' influence fruit and vegetable consumption?—A cross-sectional study. *Appetite* 2005;**45**(2):195-97

23. Pearce J, Hiscock R, Blakely T, Witten K. The contextual effects of neighbourhood access to supermarkets and convenience stores on individual fruit and vegetable consumption. *Journal of Epidemiology and Community Health* 2008;**62**(3):198-201

24. Chum A, O'Campo P. Contextual determinants of cardiovascular diseases: Overcoming the residential trap by accounting for non-residential context and duration of exposure. *Health & Place* 2013. <http://dx.doi.org/10.1016/j.healthplace.2013.07.007> (accessed Aug 18 2013).

25. Social and Aboriginal Statistics Division, Statistics Canada. General Social Survey, Cycle 24: Time Stress and Well-Being, 2010: Public Use Microdata Files. December 2011 ed. Ottawa, Ontario: Statistics Canada, 2011.

26. Pérez CE. Fruit and vegetable consumption. *Health Reports* 2002;**13**(3):23-31

27. Health Statistics Division, Statistics Canada. Canadian Community Health Survey - Annual Component, 2010. June 2011 ed. Ottawa, Ontario: Statistics Canada 2011.

28. Riediger ND, Moghadasian MH. Patterns of fruit and vegetable consumption and the influence of sex, age and socio-demographic factors among Canadian elderly. *Journal of the American College of Nutrition* 2008;**27**(2):306-13

29. Riediger ND, Shooshtari S, Moghadasian MH. The influence of sociodemographic factors on patterns of fruit and vegetable consumption in Canadian adolescents. *Journal of the American Dietetic Association* 2007;**107**(9):1511-18

30. Health Canada. Canada's Food Guide: Government of Canada, 2011.

31. Toronto Public Health. Toronto Healthy Environments Inspection System, Toronto, Ontario: City of Toronto 2012.

32. Charreire H, Casey R, Salze P, et al. Measuring the food environment using geographical information systems: a methodological review. *Public Health Nutrition* 2010;**13**(11):1773

33. Schimpl M, Moore C, Lederer C, et al. Association between Walking Speed and Age in Healthy, Free-Living Individuals Using Mobile Accelerometry—A Cross-Sectional Study. *PLoS One* 2011;**6**(8):e23299

34. Gilliland JA, Rangel CY, Healy MA, et al. Linking Childhood Obesity to the Built Environment: A Multi-level Analysis of Home and School Neighbourhood Factors Associated With Body Mass Index. *Canadian Journal of Public Health* 2012;**103**

35. Smoyer-Tomic KE, Spence JC, Raine KD, et al. The association between neighborhood socioeconomic status and exposure to supermarkets and fast food outlets. *Health & place* 2008;**14**(4):740-54

36. Apparicio P, Cloutier M-S, Shearmur R. The case of Montreal's missing food deserts: evaluation of accessibility to food supermarkets. *International journal of health geographics* 2007;**6**(1):4

- 1  
2  
3 37. Raudenbush SW, Bryk AS. *Hierarchical linear models: Applications and data analysis methods*. 2nd  
4 ed: Sage, 2002.  
5 38. Miller TM, Abdel-Maksoud MF, Crane LA, Marcus AC, Byers TE. Effects of social approval bias on self-  
6 reported fruit and vegetable consumption: a randomized controlled trial. *Nutr J* 2008;**7**(1):18  
7 39. Stark JH, Neckerman K, Lovasi GS, et al. Neighbourhood food environments and body mass index  
8 among New York City adults. *Journal of epidemiology and community health* 2013;**67**(9):736-42  
9  
10  
11  
12  
13

14 Figure 1: Mean duration (in minutes) at home on an average weekday by age and income groups  
15 (Canadian General Social Survey 2010) weighted N=28,075,610  
16  
17

18 Figure 2: Example of a network buffer for a 10 and 15 minute walking radius  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1: Mean duration (in minutes) at home on an average weekday by age and income groups (Canadian General Social Survey 2010) weighted N=28,075,610




Figure 2: Example of a network buffer for a 10 and 15 minute walking radius  
2328x1799mm (72 x 72 DPI)

CHECKLIST

We've ensured that all checklist items below have been included in our study.

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported
Objectives	3	State specific objectives, including any prespecified hypotheses
Methods		
Study design	4	Present key elements of study design early in the paper
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants (b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	Describe any efforts to address potential sources of bias
Study size	10	Explain how the study size was arrived at
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) Cohort study—If applicable, explain how loss to follow-up was addressed Case-control study—If applicable, explain how matching of cases and controls was addressed Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses

Continued on next page

For peer review only

**Results**

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses

**Discussion**

Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results

**Other information**

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based
---------	----	---

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).



# BMJ Open

## The effect of food environments on fruit and vegetable intake as modified by time spent at home: a cross-sectional study

Journal:	BMJ Open
Manuscript ID:	bmjopen-2014-006200.R1
Article Type:	Research
Date Submitted by the Author:	02-Dec-2014
Complete List of Authors:	Chum, Antony; St Michael's Hospital, Centre for Research on Inner City Health Farrell, Eddie; St Michael's Hospital, Centre for Research on Inner City Health Vaivada, Tyler; McMaster University, Faculty of Health Sciences Labetski, Anna; St Michael's Hospital, Centre for Research on Inner City Health Bohnert, Arianne; St Michael's Hospital, Centre for Research on Inner City Health Selvaratnam, Inthuja; St Michael's Hospital, Centre for Research on Inner City Health Larsen, Kristian; Toronto Public Health, Pinter, Theresa; University of Toronto, Department of Geography O'Campo, Patricia; St Michael's Hospital, Centre for Research on Inner City Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Epidemiology, Research methods
Keywords:	PUBLIC HEALTH, PREVENTIVE MEDICINE, EPIDEMIOLOGY

SCHOLARONE™  
Manuscripts

# The effect of food environments on fruit and vegetable intake as modified by time spent at home: a cross-sectional study

Corresponding author: Antony Chum,  
St Michael's Hospital  
Centre for Research on Inner City Health  
Li Ka Shing Knowledge Institute  
209 Victoria Street, third floor, Toronto, Ontario M5C 1N8

email: [antony.chum@utoronto.ca](mailto:antony.chum@utoronto.ca)

## Authors:

Antony Chum, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada  
Eddie Farrell, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada  
Tyler Vaivada, McMaster University, Hamilton Canada  
Anna Labetski, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada  
Arianne Bohnert, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada  
Inthuja Selvaratnam, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada  
Kristian Larsen, Centre for Research on Inner City Health, University of Toronto, Toronto Canada  
Theresa Pinter, University of Toronto, Toronto Canada  
Patricia O'Campo, Centre for Research on Inner City Health, St. Michael's Hospital, Toronto Canada

Keywords: vegetable, fruit, time use, residential, food environment

Word count: 3,684

## ABSTRACT

**Objective:** There is a growing body of research that investigates the residential neighbourhood context as it relates to individual diet. However, previous studies ignore participants' time spent in the residential environment and this may be a problem because time-use studies show that adults' time-use can significantly vary. To better understand the role of exposure duration, we designed a study to examine 'time spent at home' as a moderator to the residential food environment-diet association.

**Design:** Cross-sectional observational study

**Settings:** City of Toronto, Ontario, Canada

**Participants:** 2411 adults aged 25-65

**Primary Outcome Measure:** Frequency of vegetable and fruit intake (VFI) per day

**Results:** To examine how time spent at home may moderate the relationship between residential food environment and VFI, the full sample was split into three equal subgroups - short, medium, and long duration spent at home. We detected significant associations between density of food stores in the residential food environment and VFI for subgroups that spend medium and long durations at home (i.e. spending a mean of 8.0 and 12.3 hours at home respectively – no including sleep time), but no associations exist for people that spend the lowest amount of time at home (mean=4.7 hours). Also, no associations were detected in analyses using the full sample.

**Conclusions:** Our study is the first to demonstrate that time spent at home may be an important variable to identify hidden population patterns regarding VFI. Time spent at home can impact the association between the residential food environment and individual VFI.

### Strengths:

- our study extends the body of work on the effects of residential food environments on diet by looking at the amount of time that participants spend at home
- previous studies ignore participants' time spent at home, and this may be a problem because adults' time-use can significantly vary

### Limitations:

- cross-sectional observational data limits the study's ability to discern the direction of causation
- our outcome measure, vegetable and fruit intake, is based on the Canadian Community Health Survey 2010, and participants self-reported the frequency per day of fruits and vegetable eaten rather than the number of servings consumed. The self-reported frequency measure may contribute to both under and over-reporting of food intake behaviour.

INTRODUCTION

Low vegetable and fruit intake (VFI) has been linked to a number of chronic diseases including type II diabetes,[1 2] cardiovascular disease,[3 4] and breast,[5 6] esophageal and colon cancers.[7 8] Individual level determinants of VFI have been well established in the literature, where income and education are positively associated with VFI;[9-11] however, study results of the association between VFI and its potential environmental determinants are decidedly mixed. Research on the food environment has explored the impacts of food retailers on VFI (e.g. supermarkets, fast food outlets, and convenience stores). Studies have shown that living in proximity to supermarkets is associated with improved diet outcomes,[12-19] and poor diet outcomes.[20 21] Some studies also show no association between residential proximity to food vendors and VFI.[22 23]

Along with the inconsistent findings described above, the research is also characterized by a lack of consideration to individuals' duration of exposure to their neighbourhood context. By ignoring the temporal dimension of exposure, previous studies may have unintentionally introduced measurement bias because exposure duration may significantly differ between participants. There is a dearth of studies that have explored this problem using multilevel analyses of neighbourhood effects on individual health outcomes. Chum and O'Campo[24] found that the use of time-weighted multilevel regressions to account for duration of exposure resulted in i) improved strength of association and ii) improved model fit in models of the association between neighbourhood-level factors (including road traffic, access to supermarkets, and fast food restaurants) and cardiovascular disease risk compared to typical multilevel models that do not account for time. There is also evidence to suggest that time spent in the residential neighbourhood varies. According to the 2010 Canadian General Social Survey (CGSS) public use microdata,[25] time spent at home differs significantly by age and income: analysis of variance shows that total minutes spent at home on a typical weekday differs significantly by age groups and income groups ( $p<0.01$ ). Figure 1 illustrates an increase in time spent at home for older age groups. Although those with lower incomes spend more time at home compared to those in higher income brackets, the income effect is diminished with increased age.

Figure 1: Mean duration (in minutes) at home on an average weekday by age and income groups (Canadian General Social Survey 2010) weighted N=28,075,610

There are a number of studies that examine individuals' exposure to the non-residential food environment and its effect on diet.[26-28] However, the non-residential environment may be less important for those that spend a significant amount of time at home (e.g. individuals in low income and higher age brackets). In fact, Thornton et al[28] examined potential effect modification by employment status on the association between food environment and diet, since time spent at home may differ by employment status.

Given the variability of time spent at home between individuals, accounting for individuals' time use may help us avoid model misclassification by better quantifying exposure to the residential food environment. We hypothesize that people who spend more time in their home environments would rely more heavily on their local food vendors. For those that spend less time outside the home, the significance of exposure to the residential food environment may be more pronounced. Therefore, a stronger association between residential food environment and VFI may exist for those who spend more time at home compared to those who spend less time at home. Our study answers the following research questions to explore this potential dose-response relationship:

- 1) Does an association exist between the residential food environment and VFI?
- 2) Is the strength of the association between the residential food environment and VFI stronger for people who spend more time at home?

## MATERIALS AND METHODS

### *Data Sources*

Project NEHW (Neighbourhood Effects on Health and Wellbeing) is a cross-sectional investigation of neighbourhood-level determinants of population health that used a three-staged sampling method. In the first stage, 50 city-delineated neighbourhood planning areas (NPA) were randomly selected out of a total of 140 NPAs. In the second stage, 1-2 census tracts (CT) were randomly selected from each of the 50 NPAs sampled, resulting in 87 randomly selected CTs. Lastly, within the 87 CTs, approximately 30 individuals were randomly selected based on residential address. Eligibility criteria are as follows: i) only 1 resident per household, ii) participants are aged 25 to 65, iii) able to communicate in English, and iv) lived in the neighbourhood for at least 6 months. The response rate was 72%.

Data collection took place between March 2009 and June 2011. 2411 individuals, representing 87 CTs, participated in the study. Data were obtained from in-person interviews, and participants provided written informed consent at the time of their interview. The Research Ethics Board at St Michael’s Hospital in Toronto, Canada provided ethics approval for this study. To ensure the generalizability of our findings, and that it is representative of our target population, post-stratification weights were created based on demographic characteristics from the 2006 Canadian Census data for Toronto including sex, total household income, household size, immigrant status, and age. More information about the study methods can be found in a previously published paper.[24]

***Dependent Variable***

The primary outcome measure is frequency of VFI, and was assessed using questions from the U.S. Behavioural Risk Factor Surveillance System.[29] The same questions are also found in the Canadian Community Health Survey (2010).[30] Six questions, similar to questions in a food frequency questionnaire, were asked to determine total frequency of intake for vegetables and fruits.[31 32] Intake was calculated by adding the frequency of intake of fruit, fruit juice, green salad, carrots, potatoes (not including French fries, fried potatoes or potato chips), and other vegetables. VFI is based on self-reported data. Since VFI is non-normally distributed with a positive skew, for the purposes of regression analysis, we categorized VFI into either “less than five times” or “greater than or equal to five times” per day. These categories of VFI have been used in previous studies.[31 32] Frequency of intake was structured around the recommendations of Canada’s Food Guide whereby one instance of VFI was considered to be one serving, and less than 5 servings a day is below recommendation.[33] While frequency of intake may not necessarily equal the number of servings, for the purpose of this study it is treated roughly in the same manner, following the convention of previous studies.[31 32]

***Independent Variables***

Toronto Public Health’s Toronto Healthy Environments Inspection System (THEIS) (2012),[34] provided the location of all food outlets. Within the THEIS database, the type of

1  
2  
3 food retailer was used to examine food access in residential environments. Fast food, healthy  
4 food retailers, and less healthy food stores were used in this study. *Fast food restaurants* were  
5 classified as a restaurant, food court vendor, cafeteria, food take out, ice cream/yogurt vendor, or  
6 hot dog cart. These restaurants were also required to have takeout options available, limited or no  
7 wait staff, and customers having to pay prior to receiving food. If the restaurant had restricted  
8 access to the public they were removed (e.g. food outlets in arenas where patrons must pay  
9 admission to access the food outlet).

10  
11  
12 *Healthier food retail* was classified as a food store (convenience/variety or supermarket)  
13 that sells a significant quantity and diversity of vegetables and/or fruit. If these food outlets did  
14 not meet the definition for healthy food retail, they were classified as a *less healthy food store*.  
15 These are typically convenience stores that primarily sell packaged snacks with low nutritional  
16 value. To account for retail locations outside of the City of Toronto in bordering municipalities,  
17 additional fieldwork was completed to ensure all food outlets within 1km of the residential  
18 addresses were included in the analysis.

19  
20  
21 The rationale for examining the relationship between VFI and i) unhealthy food stores  
22 and ii) fast food restaurants was informed by several studies that have found an association  
23 between increased density of these stores and a reduction in vegetable and fruit intake/purchases.  
24 [35 36] Bowman et al [35] found that the mean grams of fruit and non-starchy vegetable intake,  
25 was 148g (standard error = 5g) when no fast food was consumed versus 103g (standard error =  
26 6g) when fast food was consumed during the intake period (significantly different at the level  $p <$   
27 .0001). Mason et al [36] also found reduced odds of purchasing fruits and vegetables above the  
28 median amount for participants in areas with higher density of fast food restaurants and  
29 convenience stores (OR=0.74;  $p < 0.05$ ).

30  
31  
32 Geographic Information Systems (GIS) were used to characterize the food environment  
33 of our study area. Our measure of residential food outlet availability was created based on a  
34 review of GIS methods to measure food environments,[37] where density of food stores within a  
35 buffer around an individual's residence (constructed based on 10-15 minutes walking distances)  
36 is a commonly accepted measure. In a field study of walking speeds of the general adult  
37 population using mobile accelerometers,[38] the median adult walking speed is 1.25  
38 metres/second (interquartile range=0.12), and speed declines with age at the rate of -0.0037  
39



metres/second per year between the ages of 20-60. However, speed was not significantly associated with other individual characteristics including gender, BMI, and blood pressure. Based on these results, we estimated buffers for distances that would be reachable in a 10 and 15 minute walk from the residence of each participant. The buffers created for this study were based on network service areas around each respondent's homes. Network service area is the travelable distance created using street network data (Figure 2).[39-41] Densities for 10 and 15 minute walking distance buffers were calculated in GIS for the number of fast food, less healthy, and healthy food outlets in each of the participant's walking buffers.

Figure 2: Example of a network buffer for 10 and 15 minute walk

**Covariates**

A number of individual level covariates are included in the models to adjust for potential confounding. The following categorical variables were examined: age (25-34, 35-44, 45-54, or 55-65), gender, marital status (married/common-law, or single), education (high school or less, some college, completed college), self-rated health (poor, fair, good, and very good/excellent), and family income (quintiles). While ethnicity/race is examined in bivariate analysis (i.e. white, black, south Asian, south-east Asian, West Asian, or other), only visible minority status (i.e. white vs. others) was used instead of more detailed ethnicity categories in multivariable models due to cell size limitations.

**Time Spent at Home**

Individual level data collected from Project NEWH participants provided self-reported time spent at work and time spent sleeping, but not time spent at home. The average of time spent at home per day was estimated based on individual demographic characteristics and time spent at work using the 2010 Canadian General Social Survey (CGSS) through a multivariate regression model. This method was used to derive time spent at home, and was previously used in a peer-reviewed study.[24] Data were extracted from the CGSS for adults age 25–65 in urban areas. Time spent at home (dependent variable) was modeled using the following independent variables: age, education level, income, gender, marital status, having children under 5, and minutes spent at work. All the above predictors were significantly associated with the time spent

at home ( $p < 0.01$ ). The final model had an adjusted  $r^2$  of 0.40. Beta coefficients from the regression using the CGSS data were used to estimate individual time spent at home for the Project NEWH participants. For example, starting with the intercept of 1325.15 minutes, i) for every minute spent at work 0.68 fewer minutes were spent at home, ii) females spent 37.34 more minutes at home compared to males, iii) persons with children under 6 spent 30.93 more minutes compared to those without, et cetera. Finally, we subtracted out individuals' sleeping duration from the total time at home because individuals have no chance of contact with their residential food environment while they are sleeping.

### ***Statistical analysis***

This analysis started by evaluating the bivariate relationships between predictors and the VFI outcome using i) Kruskal-Wallis non-parametric 1-way ANOVA to compare the mean VFI per day between the categorical predictors, and ii)  $\chi^2$  test to compare the categorical predictors to the proportion of individuals that ate fruits and vegetables at least 5 times per day (see Table 1). For multivariate multilevel modeling, we used the binary outcome (VFI at least 5 times per day vs. not), because i) the positive skew in the continuous outcome can impact model stability and ii) this binary outcome has been used in previous models of the impact of individuals factors on VFI.[31 32]

Since the data has a two-level structure, with individuals nested in CTs, multilevel logistic regression is used to account for the lack of spatial independence.[42] To examine how time spent at home may moderate the relationship between residential food environment and VFI, the full sample was split into three equal subgroups - short, medium, and long duration spent at home, for separate analysis. Eight models are described in this paper. Models 1a and 1b examine the full sample ( $n=2411$ ). Models 2a and 2b examine the subgroup ( $n=804$ ) that spends the least amount of time at home (0-6.5 hours/day, mean=4.7 hours), 3a and 3b examine the subgroup ( $n=804$ ) that spends a medium amount of time at home (6.51-9.7 hours/day, mean=8.0 hours), and 4a and 4b examine the subgroup ( $n=803$ ) that spends the highest amount of time at home ( $>9.7$  hours/day, mean=12.3 hours). Models with the "a" suffix calculates food outlet density using the 10 minute walking distance network buffer, and models with the "b" suffix uses the 15 minutes walking distance network buffer. All models adjusted for the effects of gender, age, education, marital status, family income, self-rated health, and visible minority

status. The binomial outcome of “eating at least 5 fruits or vegetables” is modeled using multilevel logistic regression with random intercept using PROC GLIMMIX in SAS 9.3.

RESULTS

Participants in Project NEHW were 53% female, and 44% visible minority, with a mean age of 44 years (Table 1) and a mean after-tax family income of \$91,330 (median=\$71,000). The proportion of those eating at least five VFI per day differed significantly by gender, age, education, employment status, marital status and income in bivariate analysis ( $p < 0.05$ ). The estimated time spent at home was significantly higher for females, the elderly, those with lower levels of education, those who are unemployed or not in the workforce, those who are divorced or separated, and those in the lowest income brackets.

Table 1. Sample Characteristics and associations between covariates, VFI, and time spent at home in minutes (n=2411)

	n (proportion percentage)	Mean of Fruit and Vegetable Intake per day	Kruskal- Wallis 1- way ANOVA p-value	Proportion eating 5 or more Fruits and Vegetables per day	chi square p-value	Estimated mean time spent at home (in minutes)	1-way ANOVA p- value
Gender			<0.0001		<0.0001		<.0001
Male	1,118 (46.4%)	4.20		25.6%		453.98	
Female	1,293 (53.6%)	4.59		34.6%		530.43	
Age			0.424		0.0001		<.0001
25-34	529 (21.9%)	4.24		29.4%		411.18	
35-44	742 (30.8%)	4.42		27.9%		492.11	
45-54	714 (29.6%)	4.42		29.6%		505.00	
55-65	426 (17.7%)	4.52		37.4%		550.08	

<b>Education</b>			<b>&lt;0.0001</b>		<b>&lt;0.0001</b>		<b>&lt;.0001</b>
Less than High School	95 (3.9%)	4.52		<b>31.5%</b>		631.88	
High School	389 (16.2%)	3.9		<b>23.9%</b>		529.42	
Post-Secondary Degree/Diploma	1,409 (58.5%)	4.33		<b>30.0%</b>		484.37	
Graduate Degree	514 (21.4%)	4.66		<b>35.9%</b>		479.56	
<b>Employment Status</b>			<b>0.0003</b>		<b>0.0104</b>		<b>&lt;.0001</b>
Employed	1,618 (67.1%)	4.25		<b>28.8%</b>		407.42	
On Temporary Leave	352 (14.6%)	4.54		<b>34.0%</b>		453.90	
Unemployed	228 (9.5%)	4.84		<b>39.1%</b>		663.98	
Not looking for work	212 (8.8%)	4.25		<b>27.5%</b>		717.86	
<b>Marital Status</b>			<b>0.0223</b>		<b>&lt;0.0001</b>		<b>0.0043</b>
Married/Common law	1,542 (63.9%)	4.42		<b>32.6%</b>		494.24	
Separated/Divorced	328 (13.6%)	4.27		<b>26.7%</b>		519.65	
Widowed	42 (1.7%)	5.17		<b>52.0%</b>		474.40	
Never Married	500 (20.7%)	4.16		<b>27.7%</b>		485.76	
<b>Income</b>			<b>0.0058</b>		<b>0.0002</b>		<b>&lt;.0001</b>
\$39,000 or less	697 (28.9%)	4.13		<b>25.1%</b>		578.64	
\$40,000 to \$70,999	584 (24.2%)	4.34		<b>30.7%</b>		496.42	
\$71,000 to	523 (21.7%)	4.42		<b>32.8%</b>		447.10	

\$109,999							
\$110,000 or more	606 (25.1%)	4.53		34.3%		438.65	
Ethnicity			0.0015		0.0021		0.5090
White	1349 (56.0%)	4.49		32.8%		493.17	
Black	339 (14.1%)	4.24		30.2%		524.03	
East Asian	221 (9.2%)	4.29		27.3%		497.24	
South Asian	224 (9.3%)	4.03		23.4%		488.84	
Aboriginal	18 (0.7%)	4.54		42.3%		526.12	
West Asian	52 (2.2%)	4.26		26.1%		427.47	
Latin	130 (5.4%)	3.68		21.9%		487.21	
Other	77 (3.2%)	4.50		34.9%		462.42	

All models to follow have been adjusted for the effects of gender, age, education, self-rated health, marital status, visible minority status, and family income. For models where we disregarded time spent at home, no significant associations were found between the food environment and VFI. This was true for food store density measured at scales of both 10 minute and 15 minute walking distances (see Table 2 - model 1a and 1b). Next, we conducted subgroup analyses where we divided the full sample of participants into three equal groups with different amounts of time spent at home (low, medium and high). For individuals that spent the least amount of time at home (models 2a & 2b), no significant associations were found between the food environment variables and VFI.

For individuals who spent a medium amount of time at home, model results differed by buffer size. At the 10 minute buffer, residents with zero *less healthy* food stores compared to those with three or more, had 38% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 3a). Within the 15 minute buffer, individuals with three or more *healthier* food stores

compared to those with zero, had 36% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 3b).

For individuals that spent the highest amount of time at home, model results also differed by buffer size. At the 10 minute buffer, residents with access to three or more *healthier* food stores compared to those with none, had 29% increased odds ( $p<0.05$ ) of having five or more VFIs per day. In addition, residents with zero *less healthy* food stores compared with three or more, had 38% increased odds ( $p<0.05$ ) of having five or more VFIs per day (model 4a).

Within the 15 minute buffer, residents with 'two *healthy* food stores' and 'three or more *healthy* food stores' compared with zero, both had 61% increased odds ( $p<0.05$ ) of having five or more VFIs per day. In addition, residents with access to zero or one *fast food* stores compared with three or more had 28% and 27% increased odds ( $p<0.05$ ) of having five or more VFIs per day, respectively (model 4b).

**Table 2. Multilevel logistic regression to examine the association between food environment (10 and 15 minute walking distance network buffers) and the odds of eating five vegetables and fruits per day adjusting for gender, age, education, self-rated health, marital status, visible minority status, and family income.**

	Full sample (n=2411)		Subgroup A: spending a mean of 4.7 hours at home (n=804)		Subgroup B: spending a mean of 8 hours at home (n=804)		Subgroup C: spending a mean of 12.3 hours at home (n=803)	
	Model 1a <sup>+</sup>	Model 1b <sup>+</sup>	Model 2a <sup>+</sup>	Model 2b <sup>+</sup>	Model 3a <sup>+</sup>	Model 3b <sup>+</sup>	Model 4a <sup>+</sup>	Model 4b <sup>+</sup>
	Odds ratios (95% confidence interval)							
<b>Healthier (ref=no stores)</b>								
1 store	0.843 (0.494-1.438)	1.430 (0.955-2.141)	0.964 (0.553-1.681)	1.544 (0.948-2.515)	0.788 (0.535-1.161)	0.888 (0.654-1.206)	0.825 (0.596-1.141)	0.972 (0.563-1.679)
2 stores	1.042 (0.819-1.326)	1.313 (0.867-1.988)	0.729 (0.477-1.113)	1.285 (0.970-1.701)	1.083 (0.732-1.601)	1.172 (0.943-1.457)	1.159 (0.765-1.757)	<b>1.606</b> <b>(1.082-2.385)*</b>
3 or more stores	1.249 (0.796-1.958)	1.184 (0.942-1.489)	1.109 (0.773-1.591)	0.774 (0.354-1.691)	1.139 (0.908-1.917)	<b>1.362</b> <b>(1.030-1.801)*</b>	<b>1.291</b> <b>(1.048-1.590)*</b>	<b>1.614</b> <b>(1.108-2.352)*</b>
<b>Less Healthy (ref=3+ stores)</b>								
0 stores	1.045 (0.710-1.537)	1.045 (0.710-1.537)	1.019 (0.711-1.459)	1.012 (0.690-1.483)	<b>1.377</b> <b>(1.041-1.823)*</b>	1.014 (0.709-1.451)	<b>1.381</b> <b>(1.043-1.829)*</b>	1.025 (0.716-1.467)
1 store	1.043 (0.621-1.750)	0.985 (0.576-1.685)	1.003 (0.610-1.649)	0.964 (0.566-1.644)	1.192 (0.958-1.483)	0.985 (0.600-1.616)	1.191 (0.956-1.484)	0.990 (0.604-1.625)
2 stores	0.985 (0.576-1.685)	1.043 (0.621-1.750)	1.000 (0.608-1.646)	1.024 (0.612-1.714)	0.895 (0.659-1.217)	1.003 (0.611-1.648)	0.895 (0.659-1.218)	1.015 (0.617-1.668)

Fast food (ref=3+ stores)								
0 stores	1.082 (0.731- 1.601)	1.133 (0.665- 1.931)	1.175 (0.810- 1.703)	1.658 (0.930- 2.955)	1.365 (0.940- 1.982)	1.175 (0.810- 1.703)	1.227 (0.862- 1.745)	<b>1.281 (1.037- 1.583)*</b>
1 store	1.070 (0.796- 1.439)	1.124 (0.811- 1.558)	1.126 (0.648- 1.957)	1.299 (0.979- 1.723)	1.380 (0.862- 2.209)	1.126 (0.648- 1.957)	1.235 (0.760- 2.005)	<b>1.267 (1.027- 1.563)*</b>
2 stores	0.865 (0.670- 1.116)	0.970 (0.679- 1.386)	0.874 (0.506- 1.509)	1.003 (0.538- 1.870)	0.864 (0.514- 1.453)	0.874 (0.506- 1.509)	0.763 (0.466- 1.251)	0.992 (0.745- 1.321)

\* p<0.05

+ The suffix ‘a’ label denotes food outlet densities calculated within 10 minutes walking distance, and the suffix ‘b’ denotes food outlet densities calculated within 15 minutes walking distance.

DISCUSSION

In summary, the association between density of food stores in the residential food environment and VFI exists for subgroups that spend medium and long durations at home (i.e. at least 6.5 hours in addition to time spent sleeping), but no associations exist for people that spend the lowest amount of time at home. Also, no associations were detected in analyses using the full sample. A plausible explanation for these observations is that people who spend more time at home tend to make use of their local residential food outlets, while those who spend little time at home may purchase food elsewhere as they spend their day in other locations. This is an important finding since no previous studies have differentiated participants by duration of time spent at home, and our study is the first to demonstrate that residential exposure duration may be an important missing variable to identify hidden population patterns. In this study, we show that adult time use can significantly vary across individuals and is a factor that can modify the food environment-VFI association. Given that there are no other similar studies of VFI that have accounted for time use, we cannot meaningfully compare our results to the associations found in other studies at this point. Our study highlights the importance of understanding the duration of residential exposure, and this has implications for future data collection in the context of multilevel research of environmental effects on health.

In Thornton et al,[28] employment status was examined for potential effect modification on the association between residential food environment and diet, since people not in the workforce spend more time at home compared to employed people.[27] In other words,



individuals not working may spend less time in the non-residential food environment. Thornton et al. found that supermarkets within 2km of the home were positively associated with vegetable intake, but employment status did not modify this association. In contrast, our study finds that the associations between food environment and VFI were significantly modified by time spent at home. While employment status is significantly associated with time spent at home according to the CGSS (2010) - where full-time workers, part-time workers, and those without regular employment spend a mean of 887, 1021, and 1178 minutes at home respectively (1-way ANOVA,  $p < 0.01$ ) - only 11.7% of the variance ( $\eta^2$ ) in time spent at home can be explained by employment status. Our regression-based estimation of time spent at home used multiple predictors, including i) gender, ii) marital status, iii) age, iv) education, v) parenthood status, vi) income, and vii) time spent at work, which explained 40% of the variance in time spent at home (in the CGSS 2010 data). While Thornton et al. did not find a significant effect modification by employment status, our study found an effect modification by time spent at home, which may be due to the technique used to estimate residential exposure.

Furthermore, our study illustrates that extending the network buffer from 10 minute to 15 minute walking distance has the effect of changing which predictors are significantly associated with VFI. Further research needs to be done on optimum buffer sizes to investigate access to the food environment. It should be noted that we did not have data on car ownership, and thus cannot ascertain the mode of transportation taken to their local food store.

This study has a number of limitations. First, quality and pricing data for supermarkets were not collected, both of which can affect customer shopping habits. For example, individuals may be within close proximity of a supermarket but may be unable to afford the groceries, or the quality of the fruits and vegetables may be poor. Second, the VFI outcome variable may be subject to self-reporting and social desirability bias.[43] Our questions regarding VFI is based on the Canadian Community Health Survey 2010, and participants of our study reported the frequency per day of fruits and vegetables eaten rather than the number of servings consumed. The self-reported frequency measure may contribute to both under and over-reporting of food intake behaviour. As such, it is difficult to determine the actual VFI and make comparisons to Canada's Food Guide.[31] Third, health-selected migration can occur when healthy individuals are attracted to healthier areas. Similarly, businesses may be more inclined to target

neighbourhoods where people are perceived as living healthier lifestyles.[44] Thus, there is a possibility for reverse-causation through the above processes. Fourth, this study is based on a cross-sectional observational design, and direct causation for the observed associations cannot be verified except through future longitudinal studies. Fifth, there may be residential confounding that we did not consider in our study, beyond what could be captured by self-rated health, such as the presence of specific medical conditions that may impact VFI, individual mobility issues, or dietary preferences.

Future research on the effects of the neighbourhood food environment on diet should pay greater attention to adult time use and duration of exposure to various environments. Our study highlights the importance of ensuring adequate access to healthy food stores, especially in areas with vulnerable populations that spend significantly higher amounts of time within their residential neighbourhoods (e.g. individuals who are unemployed or not in the workforce, elderly individuals, and low-income individuals). Neighbourhood-based strategies that change the food environment may be especially beneficial to vulnerable populations that spend more time at home. Public health practitioners may find value in considering the amount of time residents spend at home, and tailoring interventions to individual time use patterns. Future research should consider time spent in both non-residential and residential environments.

**CONTRIBUTORSHIP STATEMENT**

AC drafted the manuscript and conducted all the analysis, EF conducted geospatial and statistical analysis and compiled the data, AL performed the GIS processing and created the tables for geospatial analysis, TV collected and sorted the time use data and drafted the methods section, AB, IS, and TP drafted the literature and discussion components of the manuscript, KL compiled the grocery stores data, PO designed the original study and obtained the grant through CIHR for funding. AC, EF, AL, TV, AB, IS, TP, KL, and PN interpreted the results and approved the manuscript.

**COMPETING INTERESTS**

There are no competing interests.

**FUNDING**

This study is supported by funding from the Government of Canada: Social Sciences and Humanities Research Council of Canada (SSHRC) and Canadian Institutes of Health Research (CIHR).

**DATA SHARING AGREEMENT**

No additional data are available.

## REFERENCES

1. Cooper AJ, Forouhi NG, Ye Z, et al. Fruit and vegetable intake and type 2 diabetes: EPIC-InterAct prospective study and meta-analysis. *European journal of clinical nutrition* 2012;**66**(10):1082-92
2. Carter P, Gray LJ, Troughton J, et al. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ: British Medical Journal* 2010;**341**
3. Bazzano LA, He J, Ogden LG, et al. Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *The American journal of clinical nutrition* 2002;**76**(1):93-99
4. Joshipura KJ, Hu FB, Manson JE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Annals of internal medicine* 2001;**134**(12):1106-14
5. Zhang CX, Ho SC, Chen YM, et al. Greater vegetable and fruit intake is associated with a lower risk of breast cancer among Chinese women. *International journal of cancer* 2009;**125**(1):181-88
6. Jung S, Spiegelman D, Baglietto L, et al. Fruit and vegetable intake and risk of breast cancer by hormone receptor status. *Journal of the National Cancer Institute* 2013;**105**(3):219-36
7. Yamaji T, Inoue M, Sasazuki S, et al. Fruit and vegetable consumption and squamous cell carcinoma of the esophagus in Japan: the JPHC study. *International Journal of Cancer* 2008;**123**(8):1935-40
8. Aune D, Lau R, Chan DS, et al. Nonlinear reduction in risk for colorectal cancer by fruit and vegetable intake based on meta-analysis of prospective studies. *Gastroenterology* 2011;**141**(1):106-18
9. Dehghan M, Akhtar-Danesh N, Merchant A. Factors associated with fruit and vegetable consumption among adults. *Journal of Human Nutrition and Dietetics* 2011;**24**(2):128-34
10. Ding D, Sallis JF, Norman GJ, et al. Community food environment, home food environment, and fruit and vegetable intake of children and adolescents. *Journal of nutrition education and behavior* 2012;**44**(6):634-38
11. Subar AF, Heimendinger J, Patterson BH, et al. Fruit and vegetable intake in the United States: the baseline survey of the Five A Day for Better Health Program. *American Journal of Health Promotion* 1995;**9**(5):352-60
12. Giskes K, van Lenthe F, Kamphuis C, et al. Household and food shopping environments: do they play a role in socioeconomic inequalities in fruit and vegetable consumption? A multilevel study among Dutch adults. *Journal of epidemiology and community health* 2009;**63**(2):113-20
13. Laraia BA, Siega-Riz AM, Kaufman JS, et al. Proximity of supermarkets is positively associated with diet quality index for pregnancy. *Preventive medicine* 2004;**39**(5):869-75
14. Moore LV, Roux AVD, Nettleton JA, et al. Associations of the Local Food Environment with Diet Quality—A Comparison of Assessments based on Surveys and Geographic Information Systems The Multi-Ethnic Study of Atherosclerosis. *American Journal of Epidemiology* 2008;**167**(8):917-24
15. Morland K, Wing S, Roux AD. The contextual effect of the local food environment on residents' diets: the atherosclerosis risk in communities study. *Journal Information* 2002;**92**(11)
16. Rasmussen M, Krølner R, Klepp K-I, et al. Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: quantitative studies. *International Journal of Behavioral Nutrition and Physical Activity* 2006;**3**(1):22
17. Rose D, Richards R. Food store access and household fruit and vegetable use among participants in the US Food Stamp Program. *Public health nutrition* 2004;**7**(08):1081-88

18. Wrigley N, Warm D, Margetts B, et al. Assessing the impact of improved retail access on diet in 'a food desert': a preliminary report. *Urban Studies* 2002;**39**(11):2061-82

19. Zenk SN, Schulz AJ, Hollis-Neely T, et al. Fruit and vegetable intake in African Americans: income and store characteristics. *American journal of preventive medicine* 2005;**29**(1):1-9

20. Gustafson AA, Sharkey J, Samuel-Hodge CD, et al. Perceived and objective measures of the food store environment and the association with weight and diet among low-income women in North Carolina. *Public Health Nutrition* 2011;**14**(6):1032

21. Timperio A, Ball K, Roberts R, et al. Children's fruit and vegetable intake: associations with the neighbourhood food environment. *Preventive medicine* 2008;**46**(4):331-35

22. Pearson T, Russell J, Campbell MJ, et al. Do 'food deserts' influence fruit and vegetable consumption?—A cross-sectional study. *Appetite* 2005;**45**(2):195-97

23. Pearce J, Hiscock R, Blakely T, et al. The contextual effects of neighbourhood access to supermarkets and convenience stores on individual fruit and vegetable consumption. *Journal of Epidemiology and Community Health* 2008;**62**(3):198-201

24. Chum A, O'Campo P. Contextual determinants of cardiovascular diseases: Overcoming the residential trap by accounting for non-residential context and duration of exposure. *Health & Place* 2013;**24**:73-79

25. Canada S. General Social Survey, Cycle 24: Time Stress and Well-Being, 2010: Public Use Microdata Files. December 2011 ed. Ottawa, Ontario: Statistics Canada, 2011.

26. Kerr J, Frank L, Sallis JF, et al. Predictors of trips to food destinations. *Int J Behav Nutr Phys Act* 2012;**9**(1):58

27. Zenk SN, Schulz AJ, Matthews SA, et al. Activity space environment and dietary and physical activity behaviors: a pilot study. *Health & place* 2011;**17**(5):1150-61

28. Thornton LE, Lamb KE, Ball K. Employment status, residential and workplace food environments: Associations with women's eating behaviours. *Health & place* 2013;**24**:80-89

29. Pérez CE. Fruit and vegetable consumption. *Health Reports* 2002;**13**(3):23-31

30. Health Statistics Division Statistics Canada. Canadian Community Health Survey - Annual Component, 2010 2011.

31. Riediger ND, Moghadasian MH. Patterns of fruit and vegetable consumption and the influence of sex, age and socio-demographic factors among Canadian elderly. *Journal of the American College of Nutrition* 2008;**27**(2):306-13

32. Riediger ND, Shoostari S, Moghadasian MH. The influence of sociodemographic factors on patterns of fruit and vegetable consumption in Canadian adolescents. *Journal of the American Dietetic Association* 2007;**107**(9):1511-18

33. Canada H. Canada's Food Guide: Government of Canada, 2011.

34. Toronto Public Health. Toronto Healthy Environments Inspection System City of Toronto, 2012.

35. Bowman SA, Gortmaker SL, Ebbeling CB, et al. Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics* 2004;**113**(1):112-18

36. Mason KE, Bentley RJ, Kavanagh AM. Fruit and vegetable purchasing and the relative density of healthy and unhealthy food stores: evidence from an Australian multilevel study. *Journal of epidemiology and community health* 2013;**67**(3):231-36

37. Charreire H, Casey R, Salze P, et al. Measuring the food environment using geographical information systems: a methodological review. *Public Health Nutrition* 2010;**13**(11):1773

38. Schimpl M, Moore C, Lederer C, et al. Association between Walking Speed and Age in Healthy, Free-Living Individuals Using Mobile Accelerometry—A Cross-Sectional Study. *PLoS One* 2011;**6**(8):e23299

39. Gilliland JA, Rangel CY, Healy MA, et al. Linking Childhood Obesity to the Built Environment: A Multi-level Analysis of Home and School Neighbourhood Factors Associated With Body Mass Index. *Canadian Journal of Public Health* 2012;**103**
40. Smoyer-Tomic KE, Spence JC, Raine KD, et al. The association between neighborhood socioeconomic status and exposure to supermarkets and fast food outlets. *Health & place* 2008;**14**(4):740-54
41. Apparicio P, Cloutier M-S, Shearmur R. The case of Montreal's missing food deserts: evaluation of accessibility to food supermarkets. *International journal of health geographics* 2007;**6**(1):4
42. Raudenbush SW, Bryk AS. *Hierarchical linear models: Applications and data analysis methods*. 2nd ed: Sage, 2002.
43. Miller TM, Abdel-Maksoud MF, Crane LA, et al. Effects of social approval bias on self-reported fruit and vegetable consumption: a randomized controlled trial. *Nutr J* 2008;**7**(1):18
44. Stark JH, Neckerman K, Lovasi GS, et al. Neighbourhood food environments and body mass index among New York City adults. *Journal of epidemiology and community health* 2013;**67**(9):736-42

Figure 1: Mean duration (in minutes) at home on an average weekday by age and income groups (Canadian General Social Survey 2010) weighted N=28,075,610

Figure 2: Example of a network buffer for a 10 and 15 minute walking radius

CHECKLIST

We've ensured that all checklist items below have been included in our study.

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported
Objectives	3	State specific objectives, including any prespecified hypotheses
Methods		
Study design	4	Present key elements of study design early in the paper
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants (b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	Describe any efforts to address potential sources of bias
Study size	10	Explain how the study size was arrived at
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) Cohort study—If applicable, explain how loss to follow-up was addressed Case-control study—If applicable, explain how matching of cases and controls was addressed Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses

Continued on next page

For peer review only



Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) Cohort study—Summarise follow-up time (eg, average and total amount)
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time Case-control study—Report numbers in each exposure category, or summary measures of exposure Cross-sectional study—Report numbers of outcome events or summary measures
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses

Discussion


Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results

Other information


Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based
---------	----	---

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).



220x176mm (300 x 300 DPI)



116x90mm (300 x 300 DPI)

**BMJ Open**

# The effect of food environments on fruit and vegetable intake as modified by time spent at home: a cross-sectional study

Antony Chum, Eddie Farrell, Tyler Vaivada, Anna Labetski, Arianne Bohnert, Inthuja Selvaratnam, Kristian Larsen, Theresa Pinter and Patricia O'Campo

*BMJ Open* 2015 5:

doi: 10.1136/bmjopen-2014-006200

---

Updated information and services can be found at:  
<http://bmjopen.bmj.com/content/5/6/e006200>

---

*These include:*

## References

This article cites 39 articles, 9 of which you can access for free at:  
<http://bmjopen.bmj.com/content/5/6/e006200#BIBL>

## Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (2058)  
[Public health](#) (2151)  
[Research methods](#) (592)

---

## Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>