

**Supplemental Table 1: Prior knowledge and birth preparation among mothers who experienced non-cephalic birth**

		<b>n</b>	<b>%</b>
Number of infants reported as presenting in non-cephalic position		145	---
Knew prior to labor starting		42	29.0
		<b>n</b>	<b>% among those who knew</b>
Found out by*	Mother-self exam	8	19.0
	Traditional healer	0	0
	Traditional birth attendant	5	11.9
	Auxiliary nurse midwives	4	9.5
	Doctor or nurse	3	7.1
	Ultrasound	25	60.0
Preparations made based on knowledge*	No preparation	14	33.3
	Consulted facility/doctor	10	23.8
	Planned to deliver at hospital/facility	18	42.9
	Made financial arrangements	7	16.7
	Had trained clinical personnel try to flip the baby	3	7.1

\*Could provide multiple responses

**Supplemental Table 2: Prior knowledge and birth preparation among mothers who experienced multiple birth**

		<b>n</b>	<b>%</b>
Number of mothers reporting a multiple birth		55	---
Knew prior to labor starting		20	36.4
		<b>n</b>	<b>% among those who knew</b>
Found out through physical assessment by*	self	3	15
	neighbor	1	5
	traditional healer	0	0
	TBA	0	0
	ANM	1	5
	doctor or nurse	0	0
	Ultrasound exam	16	80
Preparations made based on knowledge*	No preparation	7	35
	Consulted facility/doctor	7	35
	Planned to delivery at a hospital/facility	11	55
	Made financial arrangements	4	20

\*Could provide multiple responses