

BMJ Open The prevalence of acne in Mainland China: a systematic review and meta-analysis

Danhui Li,¹ Qiang Chen,² Yi Liu,³ Tingting Liu,⁴ Wenhui Tang,⁵ Shengjie Li⁶

To cite: Li D, Chen Q, Liu Y, *et al*. The prevalence of acne in Mainland China: a systematic review and meta-analysis. *BMJ Open* 2017;**7**:e015354. doi:10.1136/bmjopen-2016-015354

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2016-015354>).

DL, QC and YL contributed equally.

Received 3 December 2016
Revised 13 March 2017
Accepted 15 March 2017



CrossMark

¹Key Laboratory of Environment and Genes Related to Diseases, Ministry of Education, Medical School, Xi'an Jiaotong University, Xi'an, China

²School of Public Health, Taishan Medical University, Tai'an, China

³Department of Dermatology, Affiliated Hospital of Jining Medical University, Jining, China

⁴The School Hospital of Taishan Medical University, Taishan Medical University, Tai'an, China

⁵Department of Biomedical Engineering, Maternal and Child Health Care of Laiwu City in Shandong Province, Laiwu, China

⁶Department of Clinical Laboratory, Eye & ENT Hospital, Shanghai Medical College, Fudan University, Shanghai, China

Correspondence to

Shengjie Li;
lshengjie6363020@163.com

ABSTRACT

Introduction Acne, a very common skin disease, can result in psychological distress and sustain impairment in quality of life. Data on the prevalence of acne and the differences in gender, region and age are limited. The aim of this review is to estimate the prevalence of acne in Mainland China comprehensively and to quantify its association with gender, region and age.

Methods We searched electronic databases with predetermined search terms to identify relevant studies published between 1 January 1996 and 30 September 2016. We pointed out repeated results using Note Express software and evaluated the studies for inclusion. Two independent reviewers extracted the data, followed with statistical analyses using Comprehensive Meta-Analysis software version 2.0. A random effects model was adopted to calculate the overall pooled prevalence and to merge categories, including gender (males and females), region (Northern China and Southern China) and age (primary and secondary students: 7–17 years old; undergraduates: 18–23 years old; overall: no limits of age) for subgroup analyses. Logistic meta-regression analysis was used to clarify the associations between acne and the predictors age, gender and region using OR and their associated 95% CI.

Results 25 relevant studies were included in this meta-analysis. The overall pooled prevalence rates of acne were 39.2% (95% CI 0.310 to 0.479). The prevalence rates in different age groups were 10.2% overall (95% CI 0.059 to 0.171), 50.2% for primary and secondary students (95% CI 0.451 to 0.554), and 44.5% for undergraduates (95% CI 0.358 to 0.534); by gender, the prevalence rates were 35.7% for females (95% CI 0.275 to 0.448) and 39.7% for males (95% CI 0.317 to 0.482); and by region, the prevalence rates were 34.2% for Northern China (95% CI 0.242 to 0.458) and 46.3% for Southern China (95% CI 0.374 to 0.555). The associations between acne and the predictors age, gender and region were statistically significant.

Conclusions In Mainland China, primary and secondary students exhibited higher prevalence rates than undergraduate students; males had higher prevalence rates of acne than females; and the prevalence rates of acne in Southern China was higher than Northern China.

INTRODUCTION

Acne, a very common skin disease among adolescents, is the fourth most common

Strengths and limitations of this study

- To our knowledge, this was the first systemic review evaluating the overall acne prevalence in Mainland China and analysing its associations with age, gender and region.
- This systematic review, covering 12 different provinces, was composed of 25 studies, with 80 000 people over a 41-year period.
- Some limitations should be considered: the diagnostic criteria for acne differed among the included studies; the age of the survey population could not be subdivided. Moreover due to heterogeneity among papers, the samples changed from variable to variable.

reason for patients aged 11–21 years to visit a doctor in the USA.¹ Acne is estimated to affect 9.4% of the world's population with the highest prevalence in adolescents.² Acne vulgaris-associated disease burden exhibits global distribution and has continued to grow in prevalence over time within this population.³ In addition, a group of increasing epidemiological data suggests that acne also affects a considerable number of adults,⁴ and women are more frequently affected by adult acne than men.⁵ The incidence of acne is different from various countries and ethnic groups. The prevalence of acne is reported to be low in developing countries of Africa.^{6,7} In Northern Tanzania, the prevalence of acne is reported to be 0.1%,⁸ whereas the prevalence of acne is 3.9% in the German population aged between 16 and 70 years.⁹ However, a comprehensive systemic review of the prevalence of acne in the Chinese population is still lacking. Several studies have shown that the estimated prevalence rate of acne varies from 8.1% to 85.1% in China^{10–12} depending on the region, subjects' ages and nationality studied.

Acne is a chronic inflammatory disease of the pilosebaceous unit resulting from androgen-induced increased sebum production, altered keratinisation, inflammation

and bacterial colonisation of hair follicles by *Propionibacterium acnes*.¹³ Acne vulgaris alters the normal skin physiology, impairing stratum corneum and causing transepidermal water loss.¹⁴ The clinical features of acne include papules, pustules, cysts, comedos and nodules. Acne occurs primarily on the face, neck and upper trunk and can lead to scar formation if treated improperly. Acne scarring is a frequent complication of acne, and patients often lack effective and safe methods for managing this condition.¹⁵ The resulting scars may negatively impact on an affected person's psychosocial and physical well-being. It has been reported that acne can result in psychological distress and have profound effects on the patients' self-esteem, which may lead to anxiety, depression, diminished self-confidence and communication difficulties.^{16 17}

Many large population-based studies have been conducted to estimate the prevalence rates of acne in regional populations, but a comprehensive statistical analysis of the prevalence of Chinese acne has not emerged so far. China, being a vast region, comprises of 34 province-level administrative regions, with 56 ethnic groups and a population of 1.3 billion people. Therefore, a country-wide, population-based study of the prevalence rates of acne in China was needed. In this review, we examined the prevalence of acne in Mainland China systematically and analysed the effects of gender, region and age on acne.

METHODS

Search strategy

Studies were aggregated from six databases between 1 January 1996 and 30 September 2016, including PubMed, EMBASE, Web of Science, the China National Knowledge Infrastructure, the VIP Database for Chinese Technical Periodicals and the Wan Fang Database for Chinese Periodicals. The following search terms were used to retrieve relevant studies: (acne), (prevalence or incidence) AND (China or Chinese). The combination of acne, prevalence, incidence, China and Chinese were used in varying combinations to identify relevant literature. Search strategies were customised to suit each database. The search strategy is presented in online supplementary table 1. In addition, a manual search was performed by checking the reference lists of eligible articles and relevant reviews.

INCLUSION CRITERIA

Studies that met all of the following criteria were included in the meta-analysis:

1. The study was a population-based survey.
2. The study evaluated the prevalence/incidence of acne.
3. The investigation involved random sampling or cluster sampling.
4. The sample size was >300.

A flow chart illustrating the article search process is presented in [figure 1](#).

Data management

Any duplicate studies were removed by Note Express. Two reviewers (QC and DHL) independently evaluated the title and abstract of all studies identified through the search against the inclusion and exclusion criteria. The full text of all eligible studies were then retrieved. Any disagreement were resolved by a third reviewer. Excluded studies and the reasons for exclusion were recorded.

DATA EXTRACTION

Customised data information was extracted independently by two investigators. The following information was collected for each study: (1) the name of the first author; (2) the location of the investigation area; (3) the year in which the investigation occurred; (4) the age range of the subjects; (5) the sample size (male, female and overall); (6) the prevalence rates obtained for males, females and overall; (7) the method used to sample subjects; (8) the criteria used to diagnose acne; and (9) the response rate.

Risk of bias (quality) assessment

Four key criteria^{18 19} were used by two independent investigators (YL and DHL) to estimate study quality: (1) the sampling scheme (random or consecutive); (2) whether the study included an adequate description of the characteristics of the study population; (3) whether a clear definition of the prevalence rate was provided; and (4) whether the response rate exceeded 75%. The scoring rule used was the following, for each quality item, 'clear or adequate' was scored as 1 point, whereas 'no' was scored as 0 point. The study was considered to be of adequate quality if the quality score was greater than or equal to 3. The two reviewers carefully assessed the included studies independently and agreed on the final grading. An additional reviewer was consulted should there be any uncertainty or disagreement.

Ethics and dissemination

Since primary data were not collected, formal ethical approval was not required. The results will be disseminated through peer-reviewed publications, conference presentations and the media.

STATISTICAL ANALYSIS

The statistical analyses were performed using Comprehensive Meta-Analysis software V.2.0 (Biostat, Englewood Cliffs, New Jersey, USA; <https://www.meta-analysis.com>). We calculated the pooled prevalence estimates as proportions and 95% CIs for the subgroup categories weighted by the sample sizes of the individual studies and then pooled these data to derive an overall proportion and its associated 95% CI. The heterogeneity of the pooled prevalence was estimated using the χ^2 -based Q statistic. Meanwhile, I^2 metrics were calculated to quantify heterogeneity. A random effects model was used if heterogeneity was



observed ($p < 0.05$); otherwise, a fixed effects model was applied. All studies were classified into one of two groups (North versus South) according to the geographical regions in which the studies were conducted. Similarly, the studies were classified into three groups based on the age of the participants in the samples, overall (no limits of age), undergraduate (18–23 years old) and primary and secondary students ('p and s', 7–17 years old). Subjects were also classified as male and female. The associations of age, gender and region with the prevalence of acne were evaluated using a logistic meta-regression model. The associations between age, gender, region and risk of acne were expressed as ORs and 95% CIs. A value of $p < 0.05$ was considered statistically significant.

Analysis of the publication bias

To examine the authenticity of data, the Egger test and Funnel plots were produced using the Comprehensive Meta-Analysis software V.2.0. No publication bias exists if the studies arranged symmetrically around the central line with a p value of > 0.05 .

RESULTS

Characteristics of the studies

Using the initial search strategy, 166 studies were identified. One hundred and forty-one studies were subsequently excluded (figure 1). A total of 25 studies^{10–12 20–41} involving 83 008 Chinese individuals were included in this meta-analysis. Detailed characteristics of each included study are presented in table 1. The overall quality of all included studies was found to be adequate (table 2).

The overall prevalence rates of acne in Mainland China

A total of 25 studies involving 83 008 Chinese people were included in this meta-analysis. However, there was significant heterogeneity in this meta-analysis ($I^2 = 99.797\%$, $Q = 11823.369$, $p < 0.001$). Thus, a random effects model was selected for the analysis. The overall pooled prevalence of acne was 39.2% (95% CI 0.310 to 0.479; figure 2).

The prevalence rates of acne in Mainland China by gender

A total of 24 studies involving 40 712 males and 41 907 females were included in this subgroup meta-analysis.




Figure 1 Flow-chart illustrating the article search process. First, we obtained 166 records identified through database searching. No additional records identified through other sources. Second, 89 records remained after duplicates were removed. Third, 58 studies were excluded after records screening. Then the remainder 31 studies were full-text articles assessed for eligibility and 6 studies were excluded. Finally 25 studies were included in the quantitative synthesis (meta-analysis).

Table 1 Characteristics of the populations examined in studies reporting the prevalence of acne in Mainland China

Name	Region	Year	Age range	Sample size	Prevalence (%)		
					Male % (n)	Female % (n)	Total (%)
Zhang Feng ²⁷	Dong guan	2013	Undergraduate	669	40.37 (379)	34.48 (290)	37.82(253)
Liu Lijuan ³⁵	Shi jiazhuang	2014	Undergraduate	742	42.11 (418)	36.11 (324)	39.49(293)
Liu Qing ³⁶	Zi bo	2013	12–49	1455	9.4 (504)	11.5 (951)	8.32(121)
Li Shengjie ³⁴	Tai an	2010	17–24	1416	39.85 (532)	46.61 (884)	44.07(624)
Zhang Zhiyong ⁴⁰	Han dan	2010	Undergraduate	1582	59.71 (834)	50.00 (748)	55.12(872)
Cai Luanduan ²⁵	Guang dong	2010	9–18	2015	49.90 (986)	59.86 (1029)	54.99(1108)
Zhang Jian ²⁰	Yi chun	2011	18–26	448	45.16 (217)	32.03 (231)	38.39(172)
Wang Renli ⁴¹	Si chuan	2008–2009	All	10,503	19.3 (4319)	16.9 (6184)	18.1(1901)
Pei Guangde ³⁷	He nan	2008	All	1547	8.80 (742)	7.86 (805)	8.39(129)
Chen Ying ²¹	Guang zhou	2008	Undergraduate	2252	34.7 (1253)	38.4 (999)	36.4(819)
Zhang Hong ³⁹	Jiang men	2006–2008	17–18	12 450	55.9 (7134)	46.02 (5136)	51.83(6453)
Deng Bin ³¹	Hai nan	2005	16–23	3500	34.8 (1990)	32.4 (1510)	33.8(1183)
Cui Jianping ³⁰	Xing tai	2007	10–18	2891	43.14 (1370)	35.44 (1521)	39.1(1130)
Li Min ²²	Fu jian	2007	15–23	1484	69.5 (836)	63.6 (648)	66.9(992)
Feng Jieying ²⁴	Guang zhou	2004–2005	11–19	1561	51.95 (743)	60.76 (818)	56.57(883)
Gao Ai ³²	Guang zhou	2004	12–20	2552	34.2 (1151)	31.6 (1305)	32.8(837)
Wu Tieqiang ²⁶	Zhu hai	2006	10–18	3200	54.9 (1790)	51.6 (1410)	53.5(17182)
Sun Zhenxiu ²³	Nan jing	2005	16–23	2100	37.91 (1000)	22 (1100)	30.33(637)
Liu Zhaorui ²⁸	Bei jing	2003	16–17	4933	67.1 (2364)	57.3 (2569)	62.0(3058)
Zhang Aihua ³⁸	Tai an, He ze	1996	14–23	1510	38.24 (829)	23.06 (581)	20.99(317)
Fu Rui ²⁹	Ji ning	2012	11–17	2560	45.40 (1343)	47.80 (1177)	46.51(1190)
Li Bing ³³	Shan xi	2012	12–17	741	55.8 (335)	49.2 (406)	52.2(387)
Shen Yiwei ¹⁰	China	2012	All	17 345	10.4 (7858)	6.1 (9487)	8.1(1405)
Law MP ¹¹	Hong Kong	2010	Undergraduate	389	N/A	N/A	85.1(331)
Wu TQ ¹²	Guang dong	2007	10–18	3163	51.3 (1785)	58.6 (1378)	53.5(1692)

(n), the number of participants, N/A, not available.

However, there was significant heterogeneity among both the males ($I^2=99.582\%$, $Q=5508.959$, $p<0.001$) and females ($I^2=99.614\%$, $Q=5957.125$, $p<0.001$) in this meta-analysis. Thus, a random effects model was selected for the analysis. Males (39.7%, 95% CI 0.317 to 0.482) exhibited a higher prevalence of acne than females (35.7%, 95% CI 0.275 to 0.448, $Z=3.903$, $p<0.001$) in the subgroup analysis (figure 3).

The prevalence rates of acne in Mainland China by region

A total of 24 studies involving 65 663 Chinese were included in this subgroup meta-analysis. Ten of these studies were conducted in Northern China (19 377 Chinese), and 14 studies were conducted in Southern China (46 286 Chinese). However, significant heterogeneity was found in both the North ($I^2=99.573\%$, $Q=2109.204$, $p<0.001$) and the South ($I^2=99.689\%$, $Q=4176.791$, $p<0.001$) subgroups in this meta-analysis. Thus, a random effects model was selected for the analysis. The South (46.3%, 95% CI 0.374 to 0.555) had a

higher prevalence of acne than the North (34.2%, 95% CI 0.242 to 0.458, $Z=2.498$, $p=0.012$) in the subgroup analysis (figure 4).

The prevalence rates of acne in Mainland China by age

A total of 25 studies, consisting of four studies that examined the overall population, 10 p and s studies, and 11 undergraduate studies, were included in this subgroup meta-analysis. However, significant heterogeneity was found overall ($I^2=99.534\%$, $Q=643.149$, $p<0.001$), for the p and s subgroup ($I^2=98.860\%$, $Q=789.719$, $p<0.001$) and for the undergraduates ($I^2=99.130\%$, $Q=1149.658$, $p<0.001$) in this meta-analysis. Thus, a random effects model was selected for the analysis. The prevalence of acne overall ages was 10.2% (95% CI 0.059 to 0.171). The p and s students (50.2%, 95% CI 0.451 to 0.554) had a higher prevalence of acne than did the undergraduates (44.5%, 95% CI 0.358 to 0.534, $Z=2.411$, $p=0.016$) in the subgroup analysis (figure 5).

**Table 2** Methodological quality of the studies reporting the prevalence of acne in Mainland China

Name	Sampling scheme	Population characteristics	Prevalence definition	Diagnostic criteria	Response rate %	Score
Zhang Feng ²⁷	Random stratified sample	Adequate	Clear	Clear	95	4
Liu Lijuan ³⁵	Random stratified sample	Adequate	Clear	Clear	N/A	3
Liu Qing ³⁶	Cluster sampling	Adequate	Clear	Clear	48.5	3
Li Shengjie ³⁴	Random stratified sample	Adequate	Clear	Clear	100	4
Zhang Zhiyong ⁴⁰	Cluster sampling	Adequate	Clear	Clear	97.75	4
Cai Luanduan ²⁵	Random stratified sample	Adequate	Clear	Clear	98.05	4
Zhang Jian ²⁰	Random stratified sample	Adequate	Clear	Clear	N/A	3
Wang Renli ⁴¹	Random stratified sample	Adequate	Clear	Clear	N/A	3
Pei Guangde ³⁷	Random stratified sample	Adequate	Clear	Clear	53.4	3
Chen Ying ²¹	Random stratified sample	Adequate	Clear	Clear	90.8	4
Zhang Hong ³⁹	Cross-sectional	Adequate	Clear	Clear	N/A	3
Deng Bin ³¹	Random stratified sample	Adequate	Clear	Clear	N/A	3
Cui Jianping ³⁰	Random stratified sample	Adequate	Clear	Clear	98.1	4
Li Min ²²	Random stratified sample	Adequate	Clear	Clear	N/A	3
Feng Jieying ²⁴	Random stratified sample	aAdequate	Clear	Clear	98.24	4
Gao Aili ³²	Cross-sectional	Adequate	Clear	Clear	96.2	4
Wu Tieqiang ²⁶	Cross-sectional	Adequate	Clear	Clear	98.84	4
Sun Zhenxiu ²³	Random stratified sample	Adequate	Clear	Clear	N/A	3
Liu Zhaorui ²⁸	Random stratified sample	Adequate	Clear	Clear	N/A	3
Zhang Aihua ³⁸	Random stratified sample	Adequate	Clear	Clear	N/A	3
Fu Rui ²⁹	Cross-sectional	Adequate	Clear	Clear	99.06	4
Li Bing ³³	Random stratified sample	Adequate	Clear	Clear	95.2	4
Shen Yiwei ¹⁰	Community-based study	Adequate	Clear	Clear	86.84	4
Law MP ¹¹	Cross-sectional	Adequate	No	Clear	99.3	4
Wu TQ ¹²	Cross-sectional	Adequate	Clear	Clear	98.8	4

N/A, not available.

Logistic meta-regression analysis of the associations between age, gender and region and the prevalence of acne

The associations between acne and the predictors age, gender and region were statistically significant (table 3) as measured by the corresponding ORs and their associated 95% CIs. The OR of acne prevalence was 1.217 (95% CI 0.109 to 14.681, $p=0.024$) between male and female subgroups. Suggesting that Chinese males might be more susceptible to acne than females. The OR between Southern China and Northern China was 1.184 (95% CI 0.002 to 3.833, $p=0.028$) for acne. The geographical factors of Southern China might be a risk factor for acne. The OR between p and s students and undergraduate was 3.127 (95% CI 0.001 to 3.838, $p=0.012$) for acne; therefore, age might have a vital role in the development of acne.

Analysis of the publication bias

There were no obvious asymmetries in the Funnel plots and the p -value exceeded 0.05 for the following groups: the overall pooled studies ($t=0.030$, $p=0.976$, 95% CI -0.215 to 0.222) (see online supplementary

figure 1), male subgroup ($t=0.346$, $p=0.733$, 95% CI -0.185 to 0.132) (see online supplementary figure 2), female subgroup ($t=0.143$, $p=0.887$, 95% CI -0.164 to 0.188) (see online supplementary figure 3), subgroup of all age layers ($t=0.501$, $p=0.666$, 95% CI -0.831 to 0.658) (see online supplementary figure 4), subgroup of p and s students ($t=0.580$, $p=0.578$, 95% CI -0.246 to 0.147) (see online supplementary figure 5), subgroup of undergraduates ($t=1.061$, $p=0.316$, 95% CI -0.121 to 0.335) (see online supplementary figure 6) and Southern China subgroup ($t=0.441$, $p=0.667$, 95% CI -0.194 to 0.293) (see online supplementary figure 7). Only the studies of the Northern China subgroup had publication bias ($t=3.369$, $p=0.01$, 95% CI -0.520 to -0.974) (see online supplementary figure 8).

DISCUSSION

This meta-analysis, based on 83008 subjects, derived from 25 studies, covering 12 provinces and municipalities in China, enables us to assess reliable prevalence rates of acne at the national level. Our results showed




Figure 2 Forest plot of the overall prevalence rates of acne. A total of 25 studies were included in this meta-analysis. Through analysing in a random effects model, the overall pooled prevalence of acne was calculated as 39.2% (95% CI 0.310 to 0.479).

that the overall pooled prevalence of acne was 39.2%. Other studies have reported a range of acne prevalence from different countries, such as 3.9% in the German population aged between 16 and 70 years⁹ and 61.5% in the Portugal population between the ages of 20 and 60 years old.⁴² The differences in age range, ethnic background, regions and pollution might explain the difference in the incidence of acne. Skin is the outermost barrier and various air pollutants such as oxides, particulate matter, ultraviolet (UV) radiation, polycyclic aromatic hydrocarbons and the ozone can all affect the skin. Puri⁴³ reported that air pollutants can cause damage to the skin by inducing oxidative stress, and prolonged or repetitive exposure to high levels of these pollutants may have profound negative effects on the skin. The various degrees of air pollution in different countries over the years have had different effects on the human skin. Males (39.7%) exhibited a higher prevalence of acne than females (35.7%, $Z=3.903$, $p<0.001$) in the subgroup

analysis of gender. Meanwhile, the South (46.3%) had a higher prevalence of acne than the North (34.2%, $Z=2.498$, $p=0.012$) in the regional subgroup analysis. Moreover, the primary and secondary students (50.2%) had a higher prevalence of acne than undergraduates (44.5%, $Z=2.411$, $p=0.016$) in the subgroup analysis of age. There was heterogeneity among the 25 included studies in this meta-analysis ($I^2=99.797\%$, $Q=11823.369$, $p<0.001$), and the following reasons might explain the phenomenon: (1) Differences existed in the design of studies, such as non-unified age range, collection of data and research objects, explaining the deviation of the prevalence of acne. (2) Evidence for seasonality was observed, with lower lipid production and reduced barrier function during the winter.⁴⁴ A correlation between the incidence of acne and skin surface lipid³⁴⁻⁴⁵ has been observed before, thus the included articles may have been conducted in different seasons,^{26 29 35 40} which resulted in different incidences of acne.




Figure 3 Forest plot of the prevalence rates of acne according to gender. A total of 24 studies were included in this subgroup meta-analysis. Through analysing in a random effects model, the prevalence rates of acne were 39.7% (95% CI 0.317 to 0.482) in males and 35.7% (95% CI 0.275 to 0.448) in females. Males (39.7%) exhibited a higher prevalence of acne than did females (35.7%) in the subgroup analysis ($Z=3.903$, $p<0.001$).

The age-related subgroup prevalence rates were also calculated in this study. The prevalence of primary and secondary students was 50.2%, which was consistent with the results of Wei *et al*⁴⁶ (51.3%). This result was similar to those reported by Karciauskiene *et al*⁴⁷ results among schoolchildren aged 7–19 years, which was 55.4% in Lithuania. However, these results are lower than a study conducted in Brazil, where a prevalence rate of 96% was found in adolescents aged 10–17⁴⁸. This difference may also be attributed to the different age range, regions and ethnic backgrounds of the study subjects. Our results suggest that primary and secondary students (50.2%) had a higher

prevalence of acne than undergraduates (44.5%, $Z=2.411$, $p=0.016$), which is in agreement with the findings of Shen *et al*.¹⁰ Moreover, using Logistic meta-regression analysis, we found that the OR between primary and secondary school students and undergraduate was 3.127 (95% CI 0.001 to 3.838, $p=0.012$) for acne. Vilar *et al*⁴⁹ and Yentzer *et al*⁵⁰ also reported that primary and secondary students (89.3%) had a higher prevalence of acne than the undergraduate subjects (61.9%) in the US population. Choi *et al*⁵¹ reported that in patients with late onset acne, the number of comedones and total number of acne lesions and the proportions of comedones were significantly less than in patients with




Figure 4 Forest plot of the prevalence rates of acne according to region. A total of 24 studies were included in this subgroup meta-analysis. Ten of these studies were conducted in Northern China and 14 studies were conducted in Southern China. Through analysing in a random effects model, the prevalence rates of acne in South was 46.3% (95% CI 0.374 to 0.555) and 34.2% in North (95% CI 0.242 to 0.458). The South (46.3%) had a higher prevalence of acne than the North (34.2%) in the subgroup analysis ($Z=2.498$, $p=0.012$).

early onset acne. The age of onset had a negative correlation with the number of comedones and the proportion of comedones in the T-zone and the entire face, namely, acne. The clinical differences in acne based on age appeared to be mainly a result of age of onset and not from the progression of the acne.⁵²⁻⁵³

In the present study, males (39.7%) had a 1.112 times higher prevalence rate of acne than did females (35.7%, $Z=3.903$, $p<0.001$). This result is in agreement with the findings of Schafer *et al*⁵⁴ that acne was more in men (29.9%) than women (23.7%) in the city of Hamburg (Germany). Adityan and Thappa⁵⁵ showed that the male to female ratio was 1.25:1 in South India. In Auckland, Lello *et al*⁵⁶ reported that males (91%) were more susceptible to acne compared with females (79%) and severe and moderately severe acne were significantly more common in males ($OR=2.6$, 95% CI $1.73 < OR < 3.9$) as well. Furthermore, moderate and severe acne (27.1% and 0.41% of patients, respectively) were significantly ($p<0.01$) common in males (36.2% and 0.9%, respectively) than females (21.2% and 0.2%, respectively) in Egypt.⁵⁷ This indicated that males had a higher risk of occurrence and development of acne than females. (1) This difference might be related to lifestyle. It is well known to us that males accounted for a larger proportion than females in drinking and smoking.⁵⁸⁻⁶⁰ (2) Males also tend to pay less attention to skin health than females in China. Inappropriate personal hygiene (use of abrasive soaps, harsh detergents and excessive scrubbing) can contribute

to the pathological process of acne.⁶¹⁻⁶² (3) Androgen levels elevation is greater in males than females during adolescence, and increased androgen levels is a risk factor of acne.⁶³ The interplay of growth hormone, insulin and insulin-like growth factor-1 signalling during puberty may also have a causal role in the pathogenesis of acne by influencing adrenal and gonadal androgen metabolism.⁶⁴ Adult women in different age categories have a lower prevalence of acne than adolescence. But recent research has shown that acne is affecting an increasing number of adults, particularly females. Adult female acne should be considered as a specific acne subtype distinct from adolescent acne, and a review on the topical and oral treatment options suitable for treating acne in adult females⁵ is expected in the future.

The prevalence rates of acne were also different in different regions. The prevalence of acne among individuals in Southern China (46.3%) was higher than those in Northern China (34.2%, $Z=2.498$, $p=0.012$). This is consistent with Subramaniyan,⁶⁵ who considered that the prevalence of acne was different between the East of the Indian mainland and its southern part. This difference may be due to several factors. (1) *UV radiation*. Enhanced sebum excretion, colonisation of the pilosebaceous duct with *P. acnes* and resultant inflammation were thought to play a critical role in the pathogenesis of acne.⁶⁶ The function of the sebaceous gland as an endocrine skin organ, which is mainly composed of sebocytes, has an important role in the occurrence and development of




Figure 5 Forest plot of the prevalence rates of acne according to age. All=overall age; p and s=primary and secondary students. A total of 25 studies, consisting of four studies that examined the overall population, 10 p and s studies and 11 undergraduate studies were included in this subgroup meta-analysis. Through analysing in a random effects model, the prevalence of acne over all ages was 10.2% (95% CI 0.059 to 0.171). The prevalence of acne in the primary and secondary students was 50.2% (95% CI 0.451 to 0.554) and 44.5% (95% CI 0.358 to 0.534) in the undergraduates. The primary and secondary students (50.2%) had a higher prevalence of acne than the undergraduates (44.5%) in the subgroup analysis ($Z=2.411$, $p=0.016$).

acne.⁶⁷ Excessive sebum production and its abnormal lipid ingredients from the sebaceous gland contributed to the formation of primary acne lesions.⁶⁸ Sebocytes also produce inflammatory cytokines and they have a vital

effect on the formation and aggravation of acne lesions.⁶⁹ Southern China has longer hours of sunshine than Northern China. Solar radiation contains UV radiation, which is an external environmental factor causing many skin diseases. UV radiation can affect the glands through direct or indirect pathways.⁷⁰ It is widely accepted that sebaceous gland hyperplasia and increased sebum secretion occur after irradiation of UV-B. The expression of inflammatory cytokines, especially interleukin (IL) 1b, IL-6, IL-8 and tumor necrosis factor (TNF)- α are significantly increased in cultured sebocytes after treatment with UV-B.⁷¹ Meanwhile, correlational research showed that time-distinct gene induction of TNF- α , IL-1b and matrix metalloproteinase (MMP)-1 in cultured HaCaT cells may be involved in UV-induced cellular responses.⁷² Although, mRNA levels do not always correspond to protein level in vivo, mRNA levels of inflammatory cytokines such as IL-1, IL-6 and TNF- α are also identified to be increased through studies using whole mouse skin and human skin exposed to UV-B.^{73 74} Furthermore, it has been known that epithelial keratinocytes contain a functional peroxisome proliferator-activated receptor gamma (PPAR)-gamma system, and this system is a target for UV-B radiation.⁷⁵ The synthesis of free fatty acids by PPAR-gamma stimulates the production

Table 3 Adjusted OR for acne obtained by logistic meta-regression analysis

Factor	Number of studies	OR (95% CI)	p Value
Age of range			
Undergraduate	11	1.00	
Primary and secondary school students	10	3.127 (0.001 to 3.838)	0.012
Gender			
Female	24	1.00	
Male	24	1.217 (0.109 to 14.681)	0.024
Location			
North	10	1.00	
South	14	1.184 (0.002 to 3.833)	0.028

of pro-inflammatory cytokines, such as IL-1b and TNF- α , in sebaceous glands.⁷¹ (2) *Humidity*. There is greater humidity in southern parts of China than Northern China. Subramaniyan⁶⁵ reported that the prevalence of dermatoses, including acne, is much higher in humid regions than arid regions. (3) *Diet*. Chili is a popular food for southern residents compared with the northern residents and is consumed in greater quantities than northern residents. Spicy food has been identified as a risk factor for the development of acne.³³ In simple terms, both environmental factors (climate and humidity) and diet may lead to the different prevalence rates of acne between Northern and Southern China.

This systematic review is likely limited by the different diagnostic criteria for acne among the included studies. Due to the enormous number of Chinese people, it was difficult to implement an unitive diagnostic criteria. In addition, the age of the survey population could not be subdivided, which made further detailed analysis of age groups impossible. More research are needed on the national prevalence of acne to provide better baseline data and to monitor the effect of acne over time in China.

CONCLUSION

The overall pooled prevalence rate of acne was 39.2% in Mainland China. Primary and secondary students exhibited higher prevalence rates than undergraduate students. Due to differences in lifestyle, skincare routines and androgen levels, males showed higher prevalence rates of acne than females. Possible etiological factors for the difference in prevalence rates of acne between Southern China and Northern China may be due to the varying UV radiation, humidity and dietary habits. The evidence generated from this paper may prove beneficial in terms of understanding the age and regional distribution and prevalence rates of acne among the Chinese population, which may help in identifying target prevention and treatment strategies for this cohort of patients.

Contributors SJL and DHL conceived the study and participated in drafting the final manuscript. QC and YL analysed the data and completed the final draft of the manuscript. TTL and WHT prepared all the figures. All authors reviewed the manuscript.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Ziv A, Boulet JR, Slap GB. Utilization of physician offices by adolescents in the united states. *Pediatrics* 1999;104:35–42.
- Tan JK, Bhate K. A global perspective on the epidemiology of acne. *Br J Dermatol* 2015;172 Suppl 1:3–12.
- Lynn DD, Umari T, Dunnick CA, et al. The epidemiology of acne vulgaris in late adolescence. *Adolesc Health Med Ther* 2016;7:13–25.
- Su P, Chen Wee Aw D, Lee SH, et al. Beliefs, perceptions and psychosocial impact of acne amongst singaporean students in tertiary institutions. *J Dtsch Dermatol Ges* 2015;13:227–33.
- Dréno B. Treatment of adult female acne: a new challenge. *J Eur Acad Dermatol Venereol* 2015;29 Suppl 5:14–19.
- Hogewoning AA, Koelemij I, Amoah AS, et al. Prevalence and risk factors of inflammatory acne vulgaris in rural and urban ghanaiian schoolchildren. *Br J Dermatol* 2009;161:475–7.
- Cordain L, Lindeberg S, Hurtado M, et al. Acne vulgaris: a disease of Western civilization. *Arch Dermatol* 2002;138:1584–90.
- Gibbs S. Skin disease and socioeconomic conditions in rural africa: tanzania. *Int J Dermatol* 1996;35:633–9.
- Augustin M, Herberger K, Hintzen S, et al. Prevalence of skin lesions and need for treatment in a cohort of 90 880 workers. *Br J Dermatol* 2011;165:865–73.
- Shen Y, Wang T, Zhou C, et al. Prevalence of acne vulgaris in Chinese adolescents and adults: a community-based study of 17,345 subjects in six cities. *Acta Derm Venereol* 2012;92:40–4.
- Law MP, Chuh AA, Lee A, et al. Acne prevalence and beyond: acne disability and its predictive factors among Chinese late adolescents in Hong Kong. *Clin Exp Dermatol* 2010;35:16–21.
- Wu TQ, Mei SQ, Zhang JX, et al. Prevalence and risk factors of facial acne vulgaris among chinese adolescents. *Int J Adolesc Med Health* 2007;19:407–12.
- Williams HC, Dellavalle RP, Garner S. Acne vulgaris. *The Lancet* 2012;379:361–72.
- McCarty M. Evaluation and management of refractory acne vulgaris in adolescent and adult men. *Dermatol Clin* 2016;34:203–6.
- Abdel Hay R, Shalaby K, Zaher H, et al. Interventions for acne scars. *Cochrane Database Syst Rev* 2016;4:CD011946.
- Lauermaann FT, Almeida HL, Duquia RP, et al. Acne scars in 18-year-old male adolescents: a population-based study of prevalence and associated factors. *An Bras Dermatol* 2016;91:291–5.
- Mulder MM, Sigurdsson V, van Zuuren EJ, et al. Psychosocial impact of acne vulgaris. evaluation of the relation between a change in clinical acne severity and psychosocial state. *Dermatology* 2001;203:124–30.
- Cheng JW, Cheng SW, Ma XY, et al. The prevalence of primary Glaucoma in mainland China: a systematic review and meta-analysis. *J Glaucoma* 2013;22:301–6.
- Say L, Donner A, Gülmezoglu AM, et al. The prevalence of stillbirths: a systematic review. *Reprod Health* 2006;3:1.
- Zhang J. Prevalence of acne in college students and the analysis of psychological condition of the patient. *J Yichun Coll* 2011;33:89–90.
- Ying C, Wu J, Liu Y, et al. The prevalence and related factors of acne in Guangzhou college students. *South Chin J Dermatol-Venereology* 2009;16:131–2.
- Li M, Candong L, Donghua C. Epidemiology investigation of acne in adolescents of different ages. *J Gansu Coll Tradit Chin Med* 2008;25:27–9.
- Zhenxiu S. Investigation of the incidence of acne in college students. *Chin J Leprosy Skin Dis* 2005;21:994–5.
- Jieying F, Ruiqiang F. Epidemiology of acne of 1561 middle school students in Guangzhou. *South Chin J Dermatol-Venereol* 2008;15:49–51.
- Luanduan C, Dinan Z, Yang L, et al. Epidemiological investigations of 2015 secondary students of acne. *Chin Mod Doctor* 2011;49:6–7.
- Wu T, Mei S, Zhang J, et al. Prevalence and risk factors of acne vulgaris in adolescents 10–18 years old. *Int J Dermatol Venereol* 2006;32:201–4.
- Zhang F, Liu J, Lin X, et al. Analysis of prevalence and awareness of acne of college students in Dongguan City. *Med Soc* 2014;27:68–70.
- Liu Z, Huang Y, Zhang H, et al. Prevalence of the knowledge and attitude behavior on acne of students grade 2 in senior high school in Beijing. *Chin J Dermatol* 2003;36:519–20.
- Rui F, Furen Z. Epidemiological investigation of acne vulgaris in Jining area junior high school students. *Chin J Leprosy Skin Dis* 2014;30:214–5.
- Jianping C, Dapeng D, Yun S. Prevalence of acne among primary and middle school students in urban and rural area of Xingtai city. *Chin J School Health* 2008;29:995–6.
- Bin D, Zeng H, Yongjiang D, et al. Survey of acne vulgaris in college students in Hainan Province and logistic analysis of risk factors. *Chin Trop Med* 2008;8:1867–8.
- Gao A, Zhang H, Zeng H, et al. The prevalence and risk factors analysis of adolescent acne in Guangzhou City tianhe district. *Chin J Leprosy Skin Dis* 2007;23:1052–3.



33. Bing L, Cangzhen H, Yanjie W. Investigation and analysis of risk factors for the occurrence of acne among middle school students in Weinan City. *Chin Health Care Nutri* 2012;22:2444–5.
34. Shengjie L, Zhang Y, Zheng G, *et al.* Investigation and analysis of sick status and influencing factors on the undergraduates facial acne vulgaris in Taian. *Chin J Dermatovenereol* 2012;26:625–7.
35. Liu L, Shiheng Song LC, *et al.* Prevalence and awareness of risk factors of acne among college students in Shijiazhuang City. *Chin J Dermatovenereol* 2014;28:171–2.
36. Liu Q, Gao X, Liu X, *et al.* An epidemiological survey on acne in zibo district, shandong province. *J Prac Dermatol* 2013;16:149–51.
37. Pei G, Du J, Huang Y, *et al.* An epidemiological survey on acne in Jiaozuo district of henan province. *Chin J Dermatovenereol* 2010;24:1129–31.
38. Aihua Z, Qingjuan G, Wanfa R, *et al.* Investigation on incidence rate of acne in youth. *Hebei Medicine* 1996;2:78–9.
39. Hong Z, Xiaobing H, Lichun F. The situation and strategies of prevention about acne of the Jiangmen middle-school students. *J Prac Dermatol* 2009;2:14–16.
40. Zhiyong Z, Ziyin L, Hui L, *et al.* Investigation of acne about related factors of college students in Handan City. *J Hainan Med Univ* 2011;17:1718–20.
41. Wang R, Ma A, Zhang X, *et al.* A survey about the prevalence of acne in Liangshan district of sichuan province. *Chin J Dermatol* 2010;43:875–7.
42. Semedo D, Ladeiro F, Ruivo M, *et al.* Adult acne: prevalence and portrayal in primary healthcare patients, in the greater Porto Area, Portugal. *Acta Med Port* 2016;29:507–13.
43. Puri P, Nandar SK, Kathuria S, *et al.* Effects of air pollution on the skin: a review. *Indian J Dermatol Venereol Leprol* 2017;7:1–9.
44. Meyer K, Pappas A, Dunn K, *et al.* Evaluation of seasonal changes in facial skin with and without acne. *J Drugs Dermatol* 2015;14:593–601.
45. Ikaraocha CI, Ikaraocha CI, Taylor GO, *et al.* Pattern of skin surface lipids in some south-western nigerians with acne vulgaris. *West Afr J Med* 2004;23:65–8.
46. Wei B, Pang Y, Zhu H, *et al.* The epidemiology of adolescent acne in north east China. *J Eur Acad Dermatol Venereol* 2010;24:953–7.
47. Karciauskiene J, Valiukeviciene S, Stang A, *et al.* Beliefs, perceptions, and treatment modalities of acne among schoolchildren in Lithuania: a cross-sectional study. *Int J Dermatol* 2015;54:e70–e78.
48. Bagatin E, Timpano DL, Guadanhim LR, *et al.* Acne vulgaris: prevalence and clinical forms in adolescents from são paulo, Brazil. *An Bras Dermatol* 2014;89:428–35.
49. Vilar GN, Santos LA, Sobral Filho JF. Quality of life, self-esteem and psychosocial factors in adolescents with acne vulgaris. *An Bras Dermatol* 2015;90:622–9.
50. Yentzer BA, Hick J, Reese EL, *et al.* Acne vulgaris in the United States, a descriptive epidemiology. *Cutis* 2010;86:94–9.
51. Choi CW, Lee DH, Kim HS, *et al.* The clinical features of late onset acne compared with early onset acne in women. *J Eur Acad Dermatol Venereol* 2011;25:454–61.
52. Williams C, Layton AM. Persistent acne in women : implications for the patient and for therapy. *Am J Clin Dermatol* 2006;7:281–90.
53. Marks R. Acne and its management beyond the age of 35 years. *Am J Clin Dermatol* 2004;5:459–62.
54. Schäfer T, Nienhaus A, Vieluf D, *et al.* Epidemiology of acne in the general population: the risk of smoking. *Br J Dermatol* 2001;145:100–4.
55. Adityan B, Thappa DM. Profile of acne vulgaris--a hospital-based study from south India. *Indian J Dermatol Venereol Leprol* 2009;75:272–8.
56. Lello J, Pearl A, Arroll B, *et al.* Prevalence of acne vulgaris in Auckland senior high school students. *N Z Med J* 1995;108:287–9.
57. El-Khateeb EA, Khafagy NH, Abd Elaziz KM, *et al.* Acne vulgaris: prevalence, beliefs, patients' attitudes, severity and impact on quality of life in Egypt. *Public Health* 2014;128:576–8.
58. Kegler MC, Hua X, Solomon M, *et al.* Factors associated with support for smoke-free policies among government workers in six chinese cities: a cross-sectional study. *BMC Public Health* 2014;14:1130.
59. Yang MJ. The chinese drinking problem: a review of the literature and its implication in a cross-cultural study. *Kaohsiung J Med Sci* 2002;18:543–50.
60. Yang X, Lu X, Wang L, *et al.* Common variants at 12q24 are associated with drinking behavior in han chinese. *Am J Clin Nutr* 2013;97:545–51.
61. Kilkeny M, Merlin K, Plunkett A, *et al.* The prevalence of common skin conditions in australian school students: 3. acne vulgaris. *Br J Dermatol* 1998;139:840–5.
62. Alexis AF, Lamb A. Concomitant therapy for acne in patients with skin of color: a case-based approach. *Dermatol Nurs* 2009;21:1.
63. Henze C, Hinney B, Wuttke W. Incidence of increased androgen levels in patients suffering from acne. *Dermatology* 1998;196:53–4.
64. Kumari R, Thappa DM. Role of insulin resistance and diet in acne. *Indian J Dermatol Venereol Leprol* 2013;79:291–9.
65. Subramaniyan R. Pattern of dermatoses among nicobarese in a community health camp at Nancowry, andaman and Nicobar islands. *Indian J Dermatol* 2016;61:187–9.
66. Cunliffe WJ. The sebaceous gland and acne-40 years on. *Dermatology* 1998;196:9–15.
67. Gollnick HP, Dreno B. Pathophysiology and management of acne. *J Eur Acad Dermatol Venereol* 2015;29 Suppl 4:1–2.
68. Zouboulis CC. Acne and sebaceous gland function. *Clin Dermatol* 2004;22:360–6.
69. Zouboulis CC, Adjaye J, Akamatsu H, *et al.* Human skin stem cells and the ageing process. *Exp Gerontol* 2008;43:986–97.
70. Roberts JE, Light RJE. Light and immunomodulation. *Ann N Y Acad Sci* 2000;917:435–45.
71. Lee WJ, Park KH, Sohn MY, *et al.* Ultraviolet B irradiation increases the expression of inflammatory cytokines in cultured sebocytes. *J Dermatol* 2013;40:993–7.
72. Skiba B, Neill B, Piva TJ. Gene expression profiles of TNF-alpha, TACE, Furin, IL-1beta and matrilysin in UVA- and UVB-irradiated HaCat cells. *Photodermatol Photoimmunol Photomed* 2005;21:173–82.
73. Murphy GM, Dowd PM, Hudspith BN, *et al.* Local increase in interleukin-1-like activity following UVB irradiation of human skin in vivo. *Photodermatol* 1989;6:268–74.
74. Scordi IA, Vincek V. Timecourse study of UVB-induced cytokine induction in whole mouse skin. *Photodermatol Photoimmunol Photomed* 2000;16:67–73.
75. Zhang Q, Southall MD, Mezsick SM, *et al.* Epidermal peroxisome proliferator-activated receptor gamma as a target for Ultraviolet B radiation. *J Biol Chem* 2005;280:73–9.

BMJ Open

The prevalence of acne in Mainland China: a systematic review and meta-analysis

Danhui Li, Qiang Chen, Yi Liu, Tingting Liu, Wenhui Tang and Shengjie Li

BMJ Open 2017 7:

doi: 10.1136/bmjopen-2016-015354

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/7/4/e015354>

These include:

References

This article cites 75 articles, 3 of which you can access for free at:
<http://bmjopen.bmj.com/content/7/4/e015354#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>