

Supplementary Table A

Comparison of reproductive and perinatal outcomes in the 1st pregnancy (live birth & full term) in women who had induced abortion, livebirth or miscarriage in the 2nd pregnancy

Outcome of 1st pregnancy	Outcome of second pregnancy			Crude Relative Risk (99% CI) ¹	
	Induced abortion N=30527	Live birth N=125855	Miscarriage N=22404	Induced abortion vs Live birth	Induced abortion vs Miscarriage
Live birth					
Pre-eclampsia	349 (1.1)	1447 (1.2)	325 (1.5)	0.99 (0.85, 1.16)	0.79 (0.65, 0.96)
Placenta previa	128 (0.4)	409 (0.3)	80 (0.4)	1.29 (0.99, 1.67)	1.17 (0.81, 1.69)
Abruptio placenta	84 (0.3)	262 (0.2)	57 (0.3)	1.32 (0.96, 1.82)	1.08 (0.70, 1.68)
Induction of labour ³	8064 (26.4)	33225 (26.4)	6103 (27.2)	1.00 (0.97, 1.03)	0.97 (0.93, 1.01)
Low birth weight <2500 ⁴	972 (3.2)	3727 (3.0)	626 (2.8)	1.08 (0.98, 1.18)	1.14 (1.00, 1.30)

Values are n (%) unless otherwise specified