



Supplemental material


Appendix to: **Experiences of patients with multimorbidity with primary care and the association with patient activation: a cross-sectional study in Germany**

Analysis – Diagrams for the Linear model



Residuen:





Results of the ordinal logistic regression analysis

	B (SE)	95 % CI	p
Age	-0.32 (0.02)	-0.08 to -0.02	0.239
Gender	-0.17 (0.37)	-0.92 to 0.56	0.637
Did your primary physician ask you about your preferences in your treatment?	0.27 (0.39)	-0.49 to 1.04	0.484
Have you agreed on treatment goals with your primary physician?	0.63 (0.39)	-0.14 to 1.47	0.111
<i>Do you feel as involved in decisions about your treatment as you would like to be?*</i>	0.89 (0.41)	0.08 to 1.70	0.030*
Have you been offered participation in a patient training programme after your diagnosis?	0.44 (0.46)	-0.47 to 1.35	0.347
Have you been offered the opportunity to participate in a support group after your diagnosis?	-0.71 (0.66)	-2.03 to 0.59	0.283

<i>Have you received a written (self-management) plan about what you can do to improve your health?*</i>	-1.21 (0.48)	-2.16 to -0.25	0.013*
Do you have a medication plan?	1.36 (0.73)	-0.07 to 2.80	0.062
In your opinion is this medication plan up-to-date?	-1.23 (0.63)	-2.47 to 0.00	0.051
Has your primary physician reviewed your medication with you in the last 12 months?	0.42 (0.44)	-0.44 to 1.30	0.340
Has your primary physician explained to you how and when you should take the medication?	-0.81 (0.53)	-1.86 to 0.23	0.127
Has your primary physician discussed with you how you cope with the burden of the chronic disease?	-0.69 (0.44)	-1.56 to 0.17	0.116

Note: * significant associations with patient activation level