
Vestibular syndromes, diagnosis and diagnostic errors in dizzy patients presenting to the emergency department. (Comolli et al.)

## **APPENDIX**

## Figure S1



Frequencies of vestibular syndromes, diagnosis and misdiagnosis rates in a tertiary emergency department (Comolli et al.)

Table S1: Cross table diagnoses emergency department (ED) vs. Follow-up

Bold cases represent the 29/662 (4.4%) cases where a dangerous diagnosis was found during follow-up but not during ED workup

		Diagnoses follow up														ı							
		Stroke / Minor Stroke	BPPV	Vestibular Deficit (e.g. Vestibular Neuritis)	TIA	Dysautonomia	Vestibular migraine	Menière's disease	PPPD	Tumor	Trauma	Medical side effects	Heart disease	Labyrinthitis	Infectious disease	Metabolic disorders	Neurodegenerative disease	Acoustic neuroma	Others	unknown	unknown etiology central vestibular syndrome	Total Diagnoses ED¹)	Change of diagnoses at follow- up <sup>2)</sup>
Diagnoses ED	Stroke / Minor Stroke	114	0	4	5	1	0	0	0	0	0	0	0	0	0	0	0	0	1	4	3	132	13.6%
	BPPV	2	20	3	0	0	0	0	0	0	0	0	0	1	0	0	0	0	2	2	0	28	28.6%
	Vestibular Deficit (e.g. Vestibular Neuritis)	2	0	75	1	0	1	2	0	0	0	0	0	1	0	0	1	1	2	6	0	89	15.7%
	TIA	7	1	0	34	0	0	0	0	0	0	1	0	0	0	1	0	0	0	4	1	49	30.6%
	Dysautonomia	0	0	0	0	6	0	0	0	0	0	1	2	0	0	0	0	0	2	0	0	9	33.3%
	Vestibular migraine	0	0	0	0	0	9	0	1	0	0	0	0	0	0	0	0	0	2	1	0	12	25.0%
	Menière's disease	0	0	3	0	0	0	11	0	0	0	0	0	1	0	0	0	0	1	0	0	15	26.7%
	PPPD	0	0	0	0	0	1	0	5	0	0	0	0	0	0	0	0	0	0	0	0	5	0.0%
	Tumor	0	0	0	0	0	0	0	0	13	0	0	0	0	0	0	0	0	0	1	0	14	7.1%
	Trauma	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	3	0.0%
	Medical side effects	1	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	3	33.3%
	Heart disease	0	0	0	0	0	0	0	0	0	0	0	5	0	0	0	0	0	0	0	0	5	0.0%
	Labyrinthitis	0	0	1	0	0	0	0	0	0	0	0	0	5	3	0	0	1	0	0	0	7	28.6%
	Infectious disease	0	0	3	0	0	0	0	0	0	0	0	0	1	3	0	0	0	0	0	0	6	50.0%
	Metabolic disorders	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0.0%
	Neurodegenerative disease	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	3	33.3%
	Acoustic neuroma	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	4	0	0	0	4	0.0%
	Others	0	1	0	0	2	0	0	0	1	0	0	0	0	0	0	0	0	38	1	0	42	9.5%
	unknown	14	6	35	9	8	22	8	6	1	0	1	1	1	1	3	0	0	28	104	4	249	58.2%
1)	Total Diagnoses Follow up1)	138	28	122		17	32	21	11	15	3	4	8	9	5	5	3	5	73	122	8	662	

<sup>1)</sup>The fields "total" refer to the number of the corresponding diagnosis. Since several diagnoses are possible, the columns and rows do not add up.

<sup>&</sup>lt;sup>2)</sup> The rate of changes of diagnoses at follow-up is calculated as follows: 100 \* (1 - correct diagnoses (grey fields) / total diagnoses ED) BPPV = benign paroxysmal positional vertigo; TIA= transient ischemic attack; PPPD = persistent postural-perceptual dizziness