

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

What are patients' preferences for revision surgery after periprosthetic joint infection? A discrete choice experiment

| | |
|-------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2019-031645 |
| Article Type: | Research |
| Date Submitted by the Author: | 13-May-2019 |
| Complete List of Authors: | Carroll, Fran; University of Bristol, Population Health Sciences, Bristol Medical School Goberman-Hill, Rachael; University of Bristol, Translational Health Sciences, Bristol Medical School Strange, Simon; University of Bristol, Translational Health Sciences, Bristol Medical School Blom, AW; University of Bristol, Translational Health Sciences, Bristol Medical School Moore, Andrew; University of Bristol, Translational Health Sciences, Bristol Medical School |
| Keywords: | Discrete Choice Experiment, Hip < ORTHOPAEDIC & TRAUMA SURGERY, periprosthetic infection, ORTHOPAEDIC & TRAUMA SURGERY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3 What are patients' preferences for revision surgery after periprosthetic joint infection? A discrete
4
5 choice experiment
6
7

8 **Authors:**
9

10 F E Carroll
11
12

13 R Goberman-Hill
14
15

16 S Strange
17
18

19 A W Blom
20
21

22 A J Moore
23
24

25 Department of Translational Health Sciences, Bristol Medical School, University of Bristol.
26
27

28 Corresponding author: Dr Andrew J Moore. Translational Health Sciences (Southmead),
29
30 Musculoskeletal Research Unit, Translational Health Sciences, Bristol Medical School, University of
31
32 Bristol, Learning and Research Building, Level 1, Southmead Hospital, Bristol, BS10 5NB, UK. Email:
33
34 a.j.moore@bristol.ac.uk. Telephone: 0117 414 7877
35
36

37
38 Word count: 3076
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: Understanding patients' preferences for treatment is crucial to provision of good care and shared decisions, especially when more than one treatment option exists for a given condition.

One such condition is infection of the area around the prosthesis after hip replacement, which affects between 0.4 and 3% of patients. There is more than one treatment option for this major complication and our study aimed to assess the value that patients place on aspects of revision surgery for periprosthetic hip infection.

Design: We identified four attributes of revision surgery for periprosthetic hip infection. Using a discrete choice experiment (DCE) we measured the value placed on each attribute by 57 people who had undergone either one- or two-stage revision surgery.

Setting: The DCE was conducted with participants from 9 National Health Service hospitals in the United Kingdom.

Participants: Adults who had undergone revision surgery for periprosthetic hip infection (n=57).

Results: Overall, the strongest preference was for a surgical option that resulted in no restrictions on engaging in valued activities after a new hip is fitted. Less valued, but still important attributes, included a shorter time taken from the start of treatment to return to normal activities, few or no side effects from antibiotics, and having only one operation.

Conclusions: Results highlight that people who have had revision surgery for periprosthetic hip infection most value aspects of care that affect their ability to engage in normal everyday activities.

These were the most important characteristics in decisions about revision surgery.

Keywords

Discrete choice experiment; hip; periprosthetic infection; orthopaedic surgery

Strengths and limitations of this study

- This is the first study to quantify the value that patients place on different aspects of revision surgery for prosthetic hip joint infection.
- Using a rigorous process involving empirical qualitative research we identified four attributes for the DCE.
- It was feasible for participants to complete the questionnaires, meaning the results were as expected with patients selecting a combination of options reflecting their preferred choice.
- While, the sample size of 57 participants sufficiently powered the analyses presented here, it is too small to conduct any subgroup analyses.
- The majority of the study sample were white, male and educated, so results may not reflect the preferences of the wider surgical population.

Introduction

For people with osteoarthritis, hip replacement is a common procedure that aims to improve function and reduce pain. In 2016 over 100,000 hip replacement procedures were conducted in the UK. With an ageing population, rates are predicted to rise ¹. Although successful for many people, approximately 0.4 -1% of patients who have undergone primary hip replacement ^{2,3} and 2-3% of patients undergoing revision hip replacement ⁴ develop deep periprosthetic joint infection (PJI) severe enough to warrant surgical revision. Patients with PJI find it devastating. Symptoms include severe pain, inflammation, discharge from the surgical wound, fever, nausea, malaise, reduction in or loss of function, dislocation, and if left untreated can lead to disability and/or death ⁵.

PJI is extremely challenging to treat. Treatment can include 'debridement, antibiotic treatment and implant retention' (DAIR), used in 7.6% of cases, or more commonly major revision surgery, which involves removal of the infected prosthesis, radical debridement of infected tissue and

1
2
3 reimplantation of a new prosthesis with subsequent antibiotic treatment ⁴. Revision surgery can be
4
5 provided in a single operation (one-stage) or as a staged operation (two-stage) where the infected
6
7 implant is removed and the patient is left without an implant (with or without a temporary spacer)
8
9 while receiving antibiotic treatment. In the staged process reimplantation of the new implant is
10
11 delayed, commonly for up to 6 months, but in some instances over 12 months later ^{5,6}. There is no
12
13 clear evidence that either a one or two-stage strategy is superior in eradicating infection ^{7,8}, but
14
15 qualitative research has shown that two-stage revision places a greater burden on patients and
16
17 families than one-stage revision. This burden is due to the extended period of immobility in between
18
19 operations, complications associated with the period of immobility, and deep psychological distress,
20
21 with some patients reporting depression and suicidal thoughts ⁶.
22
23
24
25

26 Surgeons' decisions about which type of revision surgery is most appropriate for an individual
27
28 patient take into account many factors. Our previous work indicates that decisions are based on a
29
30 combination of a surgeon's own training and clinical experience of different techniques; the
31
32 availability of hospital infrastructure such as microbiology services to quickly identify the infecting
33
34 organism; characteristics of the infecting organism and duration of infection; patient characteristics
35
36 such as age, comorbidities, frailty, and extent of damaged tissue; and published evidence of revision
37
38 techniques, or reports by senior colleagues. Surgeons also considered patients' preferences for
39
40 surgery, although this could often involve the choice between long-term suppressive antibiotics or
41
42 surgery ⁹. There is no quantitative evidence that characterises patients' preferences for one-stage or
43
44 two-stage revision surgery in this context and this is an important area of work to be investigated.
45
46 The aim of this study is to assess the surgical preferences of patients who underwent revision
47
48 surgery for prosthetic hip joint infection.
49
50
51
52
53
54
55
56

57 This study is part of a larger programme of research which aims to improve outcomes for patients
58
59 after periprosthetic joint infection (NIHR PGfAR: RP-PG-1210-12005). The programme includes a
60

1
2
3 randomised clinical trial comparing one-stage and two-stage revision surgery for prosthetic hip joint
4
5 infection¹⁰ within which this study was embedded.
6
7
8
9

10 11 Methods:

12 13 Discrete choice experiments (DCEs)

14
15
16
17
18 We undertook a discrete choice experiment (DCE) to quantify the surgical preferences of patients
19
20 who underwent revision surgery for prosthetic hip joint infection. The DCE was embedded within a
21
22 randomised clinical trial that evaluated clinical and cost effectiveness of revision surgery for
23
24 prosthetic hip joint infection and compared one-stage revision surgery with two-stage revision
25
26 surgery. Patients who were taking part in the trial were eligible for the DCE. Ethical approval for the
27
28 study was granted by NRES Committee South West – Frenchay on 31 December 2014 (14/SW/1166).
29
30 All participants provided written informed consent to take part in the DCE study.
31
32
33
34
35

36 DCEs are an established method in health services research and have been used to explore a range
37
38 of health-related services and treatments¹¹⁻¹³. DCEs involve asking respondents to choose between
39
40 hypothetical scenarios which describe goods or services, where a service may mean an intervention
41
42 or an approach to care. The method aims to establish what attributes of that service influence their
43
44 decision-making and to what extent. This enables quantification of the marginal impact of these
45
46 attributes. Scenarios within a DCE describe the service of interest (in this case, revision joint
47
48 replacement) using the same set of attributes, but at different levels in each scenario. Choices
49
50 between scenarios, or whether to accept or reject a scenario, are used to estimate the influence and
51
52 value of the different attribute levels.
53
54
55
56
57
58
59
60

Questionnaire development

When a patient faces revision surgery for prosthetic hip infection, a one-stage or a two-stage operation is required, and the decision about which is undertaken is made largely at the discretion of the surgeon, taking into account patient preferences for treatment. The DCE study was designed to engage patients in their preferences for the features associated with these two surgical options, given neither is currently known to be clinically superior in terms of patient outcomes.

Qualitative methods are recommended for DCE attribute development because they enable conceptual development of attributes directly from people's experiences and so better reflect the issues that are likely to matter most to people when making a decision¹⁴. For this study we developed attributes from our previous qualitative study which explored impact of prosthetic joint infection and its treatment on patients and their recovery process; one-to-one interviews were conducted with 19 patients with PJI, focusing on the impact of PJI and surgical treatment. The interviews were audio-recorded, transcribed and the qualitative data set was analysed thematically⁶. Levels of attributes were assigned by refining the language used to convey the meaning of the attributes, and particularly where some quantification of an attribute was mentioned by participants during qualitative interview, as illustrated in Table 1¹⁴.

Table 1 describes the attribute and level selection based on the earlier qualitative work, including illustrative quotations and rationale.

The final questionnaire comprised a single scenario task in which respondents were asked to imagine this was the first time that they had been offered the two surgical options presented and were asked to consider them carefully before selecting the one (from a pair) they would prefer.

1
2
3 With a $4 \times 4 \times 2 \times 2$ design (two four-level and two two-level attributes), a total of 64 different
4 combinations of attribute levels (profiles) is possible. This converts to 32 pairs of profiles, which as a
5 'full factorial' was considered to be too large for participants to complete¹⁵. An orthogonal main
6 effects plan was therefore used to reduce the number of choice sets to 16¹⁶. Each profile was
7 presented with its pair, and participants were required to select which option they preferred (see
8 Figure 1).
9
10
11
12
13
14
15

16 17 18 19 Patient and Public Involvement

20
21
22 The questionnaire was piloted and refined in collaboration with 5 patient and public involvement
23 representatives. At an initial meeting of representatives, group members were involved in
24 questionnaire development and suggested improvements to its formatting to aid readability, clearer
25 phrasing of the questions to avoid ambiguity, and shortening of the instruction leaflet for clarity. At a
26 subsequent meeting, members of the group completed the questionnaire and fed back on their
27 experience. They completed the questionnaire without assistance and felt the instructions were
28 clear but suggested that key points in the questions should be highlighted and that a contact
29 telephone number should be added to enable participants to seek assistance if needed. For
30 participants in the study, a summary of findings will be sent to those who indicated that they wished
31 to be informed.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 49 Participants

50
51
52 Patients who were recruited into the INFORM randomised controlled trial (ISRCTN10956306)
53 received the discrete choice task after completing the 18-month primary outcome measure. The
54 questionnaire was either posted to participants or completed in person with the assistance of a
55 research nurse during a hospital clinic visit.
56
57
58
59
60

Data analysis

Sample size calculations for DCEs are challenging due to dependence on the true parameter values estimated in the choice model¹⁷. However, reliable statistical analysis has been demonstrated with sample sizes of 40–120 respondents and combined with the rarity of periprosthetic joint infection (1% of those undergoing hip replacement), a sample size above 50 participants for the DCE was deemed adequate to obtain sufficient data for analysis and interpretation^{18, 19}.

Paper questionnaires were distributed to participants as part of the follow-up data collection in the INFORM trial. Questionnaires were returned to the study team between January 2017 and November 2018 and data entered onto a Microsoft Excel spreadsheet. DCE data were effects coded and analysed using STATA SE 15^{20, 21}. The influence of the four attributes on patient choices was analysed using a conditional logit model. As attribute levels are effects coded, the mean of all coefficients is 0 across each attribute. The coefficients indicate the strength of preference for each level, where more positive values indicate a greater likelihood that the patient would select a surgical option where that particular attribute level is present.

Results:

Participants

Of the 80 discrete choice questionnaires provided to trial participants, 57 were returned fully complete (71%) from patients from 9 trial sites. Data from 6 questionnaires was not used in analysis as they were partially or totally incomplete. Responding participants had a mean age of 70 (range 51 to 90), 21 (37%) were female, 26 (46%) had undergone 1-stage revision, 14 (25%) lived alone and 41 (72%) were retired from work. Table 2 provides further demographic information.

Table 2 – Demographic and Clinical characteristics of the DCE Respondents

| Characteristic | Participants (N=57) |
|---------------------------------------|---------------------|
| Age – Yrs [mean] (range) | 70 (51-90) |
| Gender [number] (%) | |
| Male | 36 (63) |
| Female | 21 (37) |
| Ethnicity [number] (%) | |
| White | 55(96) |
| Black | 1 (2) |
| Mixed | 1 (2) |
| Marital Status [number] (%) | |
| Married / Partner | 42 (74) |
| Divorced / Separated / Widowed | 12 (21) |
| Single | 3 (5) |
| Living Arrangements [number] (%) | |
| With Partner / somebody else | 43 (75) |
| Alone | 14 (25) |
| Schooling/Education [number] (%) | |
| Left at normal school leaving age | 35 (61) |
| Left after normal school leaving age | 15 (26) |
| Left before normal school leaving age | 7 (12) |
| Work Situation [number] (%) | |
| Retired | 41 (72) |
| Working / Sick leave | 14 (25) |

| | |
|--|---------|
| Unemployed | 2 (4) |
| Surgery Received for Prosthetic Hip Joint Infection [number] (%) | |
| 2-stage revision | 31 (54) |
| 1-stage revision | 26 (46) |

Discrete choice experiment

Table 3 shows the regression coefficients and the results from the 57 patients who had fully completed the discrete choice questionnaire.

Table 3 – discrete choice task results from conditional logistic regression

| <i>Attribute</i> | <i>Level</i> | <i>Coefficient</i> | <i>Standard error</i> | <i>95% confidence interval</i> | <i>P value</i> |
|---|-----------------------|--------------------|-----------------------|--------------------------------|----------------|
| Ability to engage in valued activities after new hip is fitted. | Can do everything* | 0.70 | | | |
| | Can do most things | 0.49 | .08 | .33 to .64 | <0.001 |
| | Cannot do most things | -0.39 | .07 | -.53 to -.24 | <0.001 |
| | Cannot do anything | -0.80 | .13 | -1.05 to -0.55 | <0.001 |
| Antibiotic side effects | Don't affect me much* | 0.22 | | | |

| | | | | | |
|---|------------------|-------|-----|--------------|--------|
| | Affects me a lot | -0.22 | .05 | -.33 to -.12 | <0.001 |
| | | | | | |
| Number of operations | 1* | 0.20 | | | |
| | 2 | -0.20 | .07 | -.35 to -.06 | <0.001 |
| | | | | | |
| Time taken after surgical treatment starts, to return to normal activities. | 3m* | 0.20 | | | |
| | 6m | 0.31 | .09 | .14 to .48 | <0.001 |
| | 12m | -0.06 | .05 | -.15 to .04 | 0.22 |
| | 18m | -0.45 | .10 | -.64 to -.26 | <0.001 |
| | | | | | |
| <i>*Indicates reference category within attribute</i> | | | | | |

Analysis indicates that participants had the strongest preference for a surgical option that resulted in the least restrictions on engaging in valued activities after the new hip is fitted, illustrated by the largest positive coefficient. Other less valued but important preferences were for a surgical strategy that would result in a shorter time after surgical treatment starts to return to normal activities, few or no side effects from antibiotics, and only one operation. The results also suggest that the least restrictions on engaging in valued activities, and the shortest time taken to return to normal activity are the individual attributes most valued by patients in this sample. This is indicated by the larger spread of coefficients (i.e. more 'value' is placed on changes in these attributes). The most acceptable option was a time period of between 3 to 6 months to return to normal activity, however

1
2
3 there is no clear preference up to 12 months, although 18 months appeared to be significantly
4
5 disfavoured.
6
7
8
9

10 Discussion:

11
12
13 This study aimed to investigate and understand patients' preferences for aspects of revision surgery
14
15 for periprosthetic joint infection. Four relevant attributes were identified through earlier qualitative
16
17 work, and quantitatively, patients in this study most value the ability to engage in valued activities
18
19 and the time taken to return to normal daily activities. This reflects the findings of our previous
20
21 qualitative work which shows that although both revision strategies impacted greatly on patients
22
23 and their families, patients receiving two-stage revision surgery experience particularly long periods
24
25 of immobility and social isolation. This was often followed by a protracted recovery period, which
26
27 could leave patients much less able than before their primary operation, and some patients
28
29 experienced profoundly negative psychological effects associated with physical suffering, loss of
30
31 dignity and independence⁶. It appears that for patients in our sample, 3-6 months to return to
32
33 normal activity was preferable, although there was no significant difference up to 12 months, but 18
34
35 months was disfavoured. This suggests that the acceptable margin of recovery for patients is up to
36
37 12 months after their receipt of a new hip joint.
38
39
40
41
42
43

44 Discrete choice methodology can be challenging for participants because the format of questions is
45
46 different to standard surveys and items can seem repetitive. We collected feedback from the first 11
47
48 participants who completed the questionnaire. We found that those participants who were
49
50 supported by a research nurse when completing the questionnaire were more likely to complete it
51
52 and return it, compared with those who received the questionnaire by post and completed it alone.
53
54 Participant feedback suggested that the questionnaire was difficult to complete, as the scenarios
55
56 were similar and appeared to be repetitive. To address this, we amended the questionnaire format
57
58 and instructions and offered participants support either face-to-face or by phone with one of the
59
60

1
2
3 study research nurses. Nurses were then able to answer queries about the questionnaire and offer
4 support if needed. The results suggest the group completed the questionnaire in a rational and
5
6 logical way, meaning the results were as expected with patients selecting an optimal combination of
7
8 options as their preferred choice. This is an important methodological finding because although our
9
10 study demonstrates the feasibility of the DCE method with this population, others conducting similar
11
12 studies with older, ill populations could consider in advance the need for professional support in the
13
14 completion of discrete choice questionnaires.
15
16
17

18
19
20
21 Participants were all individuals who participated in a clinical trial and had already undergone
22
23 revision surgery for periprosthetic joint infection. This meant that the choices that participants were
24
25 asked to make in the questionnaire were based on scenarios unlikely to reflect their real-life
26
27 experiences as in reality such choices would not be available to them, since decisions about surgical
28
29 strategies are based on a wider variety of clinical, surgeon, patient and organisational factors⁹. Also,
30
31 patients were not being faced with these decisions at the time of questionnaire completion as we
32
33 decided that it would be unethical to ask patients awaiting treatment to complete a DCE about
34
35 surgical options in a hypothetical context, at a time when they may be particularly vulnerable. To do
36
37 so would not meet an ethical standard of protection from harm as it might mean that patients were
38
39 inadvertently led to believe that there were more or different options available to them than were
40
41 clinically indicated at that time.
42
43
44

45
46
47 We also found that some participants reported difficulty separating their own recent personal
48
49 experience of revision surgery from the hypothetical scenarios presented in the questionnaire, as
50
51 they found it hard to imagine receiving a treatment option that differed from the one that they had
52
53 received. In terms of methodology, many DCE's are conducted with participants who already have
54
55 some experience of the treatment attributes under investigation^{12, 18, 19, 22}. In our study previous
56
57 experience meant that participants had some appreciation of the attributes being tested. While, the
58
59
60

1
2
3 sample size of 57 participants provided sufficient data for the analyses presented here, the sample
4
5 size was too small to conduct any subgroup analysis to identify whether preferences would differ
6
7 between participants who had received one or two stage revision. Similarly, patients who are older,
8
9 still working or live alone may have had stronger preferences for a one-stage operation than those
10
11 who are younger, retired and have support at home to cope with a two-stage procedure. The
12
13 majority of the study sample were also white, male and educated, which means that results may not
14
15 reflect the preferences of the wider surgical population. Further research could explore preferences
16
17 in a more diverse population.
18
19

20
21
22
23 This work has provided an initial and important first step in understanding patients' preferences for
24
25 characteristics associated with revision surgery for periprosthetic infection. It is important that
26
27 orthopaedic healthcare professionals discuss these attributes with patients when discussing options
28
29 for surgical and antibiotic treatment for periprosthetic infection.
30
31

32 33 34 35 Implications

36
37 Results of this study offer insight into the preferences of patients for revision surgery and provide
38
39 valuable information to surgeons from all disciplines. Although factors affecting patient preferences
40
41 for surgery differ to those valued by medical professionals, consideration should be given to such
42
43 factors in order to aid shared decision-making where clinical equipoise between options exists.
44
45

46
47
48 Previous research using discrete choice approaches has explored patient preferences in a surgical
49
50 context. This has included examination of preferences for surgical versus non-surgical interventions
51
52 in a range of conditions, such as oesophageal cancer and ulcerative colitis ^{22, 23}. Some research has
53
54 explored preferences for conditions in which there are two surgical options, including for ectopic
55
56 pregnancy, vaginal wall prolapse, and osteoarthritis ^{24, 25}. Across all of these conditions, evidence
57
58 suggests that patients choose options that reduce the need for further surgery or operations, have a
59
60

1
2
3 shorter recovery time, lower risk of symptom recurrence and improve ability to preserve existing
4 joint motion in the case of osteoarthritis. Findings from the current study are similar in that patients
5 prefer a surgical option that reduces the number of operations, recovery time and the side effects of
6 antibiotics. However, in the setting of infected joint replacement, patients placed highest value on
7 restoration of function. This was more important to the patients in our study than the number of
8 operations they would have to undergo. Although our study focused on the preferences between
9 one and two-stage surgery surgeons may need to consider these preferences during shared
10 decision-making about all options for revision surgery for prosthetic hip infection, including the role
11 of debridement with retention of implants (DAIR). Although DAIR is only efficacious in approximately
12 60% of cases, it is associated with a quicker return to valued activities and improved joint function ²⁶.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 Conclusions

29
30 Our results show that the most valued characteristics in decisions about revision surgery for
31 prosthetic hip infection were the ability to engage in valued activities and time taken to return to
32 normal activity. This builds on the findings of our previous qualitative work which shows that
33 although both revision strategies impact greatly on patients and their families' everyday lives,
34 patients receiving two-stage revision surgery experience particularly long periods of immobility and
35 social isolation ⁶. The desire to return to everyday activities should be taken into account when
36 surgeons are discussing options with patients, particularly when there is equipoise from a surgical
37 perspective about the options available, and when the decision is 'preference sensitive'.
38
39
40
41
42
43
44
45
46
47
48

49 Funding

50
51 This study is funded by the National Institute for Health Research Programme Grants for Applied
52 Research (NIHR PGfAR) programme (grant number: RP-PG-1210-12005) and supported by the NIHR
53 Comprehensive Clinical Research Network (CRN). This study was also supported by the NIHR
54 Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the
55
56
57
58
59
60

1
2
3 University of Bristol. The views expressed are those of the authors and not necessarily those of the
4
5 NIHR or the Department of Health and Social Care.
6
7

8 9 Author Contributions

10
11 FEC: Contributed to design, analysis and interpretation of data, and drafting the manuscript. RG-H:
12
13 Contributed to conception, design, interpretation of data, drafting and revising the manuscript. SS:
14
15 Contributed to design, data collection and revising the manuscript. AWB: Contributed to the
16
17 conception, design, interpretation of data and revising the manuscript. AJM: Contributed to the
18
19 design, interpretation of data, drafting and revising the manuscript. All authors gave final approval
20
21 of the submitted manuscript.
22
23
24
25

26 27 Competing Interests

28
29 None declared
30
31

32 33 Data Availability

34
35 Anonymised research data will be made available at the University of Bristol Research Data
36
37 Repository (data.bris) beginning 1 year after publication of the results. Access will be
38
39 granted to bona fide researchers after the University of Bristol Data Access Committee has
40
41 approved their request. Please contact data-bris@bristol.ac.uk in the first instance.
42
43
44
45

46 47 Acknowledgements

48
49 The authors thank the participants who gave their time to complete a discrete choice questionnaire.
50
51 We also thank Dr Erik Lenguerrand for providing demographic data for the study sample and advice
52
53 on presentation of statistical results. We would also like to thank the study administration and
54
55 management team: Makita Werrett and Beverley Evanson for their invaluable support. The authors
56
57 would also like to thank the INFORM Research Advisory Group (I-RAG) and the Patient Experience
58
59
60

1
2
3 Partnership in Research (PEP-R) at the University of Bristol for their contributions to the refinement
4
5 of the questionnaire.
6
7
8
9
10

11 References

- 12
13
14 1. National Joint Registry. National Joint Registry 14th Annual Report 2017: National Joint
15 Registry for England, Wales, Northern Ireland and the Isle of Man; 2017 [Available from:
16 www.njrreports.org.uk.
17
- 18 2. Blom AW, Taylor AH, Pattison G, Whitehouse S, Bannister GC. Infection after total hip
19 arthroplasty. The Avon experience. The Journal of bone and joint surgery British volume.
20 2003;85(7):956-9.
- 21 3. Phillips JE, Crane TP, Noy M, Elliott TS, Grimer RJ. The incidence of deep prosthetic infections
22 in a specialist orthopaedic hospital: a 15-year prospective survey. The Journal of bone and joint
23 surgery British volume. 2006;88(7):943-8.
- 24 4. Lenguerrand E, Whitehouse MR, Beswick AD, Jones SA, Porter ML, Blom AW. Revision for
25 prosthetic joint infection following hip arthroplasty: Evidence from the National Joint Registry. Bone
26 & joint research. 2017;6(6):391-8.
- 27 5. Zimmerli W, Trampuz A, Ochsner PE. Prosthetic-joint infections. The New England journal of
28 medicine. 2004;351(16):1645-54.
- 29 6. Moore AJ, Blom AW, Whitehouse MR, Goberman-Hill R. Deep prosthetic joint infection: a
30 qualitative study of the impact on patients and their experiences of revision surgery. BMJ open.
31 2015;5(12):e009495.
- 32 7. Kunutsor SK, Whitehouse MR, Blom AW, Beswick AD. Re-Infection Outcomes following One-
33 and Two-Stage Surgical Revision of Infected Hip Prosthesis: A Systematic Review and Meta-Analysis.
34 PloS one. 2015;10(9):e0139166.
- 35 8. Kunutsor SK, Whitehouse MR, Blom AW, Board T, Kay P, Wroblewski BM, et al. One- and
36 two-stage surgical revision of peri-prosthetic joint infection of the hip: a pooled individual
37 participant data analysis of 44 cohort studies. European journal of epidemiology. 2018;33(10):933-
38 46.
- 39 9. Moore AJ, Blom AW, Whitehouse MR, Goberman-Hill R. Managing uncertainty - a
40 qualitative study of surgeons' decision-making for one-stage and two-stage revision surgery for
41 prosthetic hip joint infection. BMC musculoskeletal disorders. 2017;18(1):154.
- 42 10. Strange S, Whitehouse MR, Beswick AD, Board T, Burston A, Burston B, et al. One-stage or
43 two-stage revision surgery for prosthetic hip joint infection--the INFORM trial: a study protocol for a
44 randomised controlled trial. Trials. 2016;17:90.
- 45 11. Louviere J, Street DJ, Burgess L. A 20+ years' retrospective on choice experiments.
46 Marketing research and modeling: Progress and prospects: Springer; 2004. p. 201-14.
- 47 12. de Bekker-Grob EW, Hofman R, Donkers B, van Ballegooijen M, Helmerhorst TJ, Raat H, et al.
48 Girls' preferences for HPV vaccination: a discrete choice experiment. Vaccine. 2010;28(41):6692-7.
- 49 13. Liede A, Mansfield CA, Metcalfe KA, Price MA, Snyder C, Lynch HT, et al. Preferences for
50 breast cancer risk reduction among BRCA1/BRCA2 mutation carriers: a discrete-choice experiment.
51 Breast cancer research and treatment. 2017;165(2):433-44.
- 52 14. Coast J, Al-Janabi H, Sutton EJ, Horrocks SA, Vosper AJ, Swancutt DR, et al. Using qualitative
53 methods for attribute development for discrete choice experiments: issues and recommendations.
54 Health economics. 2012;21(6):730-41.
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
15. Marshall D, Bridges JF, Hauber B, Cameron R, Donnalley L, Fyie K, et al. Conjoint Analysis Applications in Health - How are Studies being Designed and Reported?: An Update on Current Practice in the Published Literature between 2005 and 2008. *The patient*. 2010;3(4):249-56.
 16. Hedayat AS, Sloane NJA, Stufken J. *Orthogonal arrays: theory and applications*: Springer Science & Business Media; 2012.
 17. Lancsar E, Louviere J. Conducting discrete choice experiments to inform healthcare decision making: a user's guide. *PharmacoEconomics*. 2008;26(8):661-77.
 18. Salkeld G, Solomon M, Butow P, Short L. Discrete-choice experiment to measure patient preferences for the surgical management of colorectal cancer. *The British journal of surgery*. 2005;92(6):742-7.
 19. de Bekker-Grob EW, Essink-Bot ML, Meering WJ, Koes BW, Steyerberg EW. Preferences of GPs and patients for preventive osteoporosis drug treatment: a discrete-choice experiment. *PharmacoEconomics*. 2009;27(3):211-9.
 20. Bech M, Gyrd-Hansen D. Effects coding in discrete choice experiments. *Health economics*. 2005;14(10):1079-83.
 21. StataCorp. *Stata Statistical Software: Release 15*. College Station, TX: StataCorp LLC; 2017.
 22. Noordman BJ, de Bekker-Grob EW, Coene P, van der Harst E, Lagarde SM, Shapiro J, et al. Patients' preferences for treatment after neoadjuvant chemoradiotherapy for oesophageal cancer. *The British journal of surgery*. 2018;105(12):1630-8.
 23. Bewtra M, Kilambi V, Fairchild AO, Siegel CA, Lewis JD, Johnson FR. Patient preferences for surgical versus medical therapy for ulcerative colitis. *Inflammatory bowel diseases*. 2014;20(1):103-14.
 24. van Mello NM, Mol F, Opmeer BC, de Bekker-Grob EW, Essink-Bot ML, Ankum WM, et al. Salpingotomy or salpingectomy in tubal ectopic pregnancy: what do women prefer? *Reproductive biomedicine online*. 2010;21(5):687-93.
 25. Notten KJ, Essers BA, Weemhoff M, Rutten AG, Donners JJ, van Gestel I, et al. Do patients prefer mesh or anterior colporrhaphy for primary correction of anterior vaginal wall prolapse: a labelled discrete choice experiment. *BJOG : an international journal of obstetrics and gynaecology*. 2015;122(6):873-80.
 26. Kunutsor SK, Beswick AD, Whitehouse MR, Wylde V, Blom AW. Debridement, antibiotics and implant retention for periprosthetic joint infections: A systematic review and meta-analysis of treatment outcomes. *The Journal of infection*. 2018;77(6):479-88.

48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 – qualitative support for attributes included in the discrete choice questionnaire

| Attribute | Evidence of attribute inclusion, with pseudonym and surgery type | Levels | Rationale for levels |
|----------------------|--|----------------|-------------------------------|
| Number of operations | "There's no way I want two more big operations now at | 1) 1 operation | Two types of revision surgery |

| | | | |
|--|--|--|---|
| | <p>my time of life. You do it all or not at all..."I said there was no way I wanted two ops" (Harriet, 1-stage)</p> <p>"Of course, emotionally, you want it over and done with as soon as possible...but ultimately that has to be done in the correct way. There's the tortoise and hare situation. There's absolutely no point in rushing ahead if ultimately it's going to fail." (Maggie, 2-stage)</p> | <p>2) 2 operations</p> | <p>are currently provided in healthcare, and involve either one or two operations</p> |
| <p>Ability to engage in valued activities after new hip is fitted.</p> | <p>"Fourteen months without a hip joint so it meant that I couldn't drive a car, I couldn't do anything that I'd been used to doing, playing golf or doing anything. Well, I gave up</p> | <p>1) Can do everything</p> <p>2) Can do most things</p> <p>3) Cannot do most things</p> | <p>Following revision surgery, the ability to engage in valued activities can be</p> |

| | | |
|--|------------------------------|--|
| <p>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60</p> <p>golf actually after the first revision” (Don, 2-stage)</p> <p>“But when I for example went to, on holiday recently and I had serious problems getting into the bath to stand in the shower. Because I, I couldn’t get in. And in a wet – on a wet surface and that, I’m very conscious of not falling in. I can’t afford to fall. So, pain I’ve got none, stiffness none.</p> <p>Physical function limitations, and that’s one of them.” (Rory, 1-stage)</p> <p>“My aim has always been to get back on my feet as soon as I can, and to walk as good as I can, and that’s a big disappointment. I’m</p> | <p>4) Cannot do anything</p> | <p>reduced in a major, or somewhat more minor way. These levels capture variation in ability identified by patients.</p> |
|--|------------------------------|--|

| | | | |
|--|--|--|--|
| | <p>not where I think I should have been” (Robert, 2-stage)</p> | | |
| <p>Time taken after surgical treatment starts, to return to normal activities.</p> | <p>“I didn’t want to go 14 weeks with effectively one leg. What was worse was not knowing that I had to endure all of those weeks not knowing that I was ever going to get another hip joint back.” (Maggie, 2-stage)</p> <p>“If I had known how hard it was going to be for her to walk in that interim six months, and if there seemed to have been a reasonable, or a good possibility that the infection would be nuked in a one stage, then that might have been a better</p> | <ol style="list-style-type: none"> 1) 3 months 2) 6 months 3) 12 months 4) 18 months | <p>These time intervals demonstrate best approximations and a reflection of the need to ensure normal expectations of time taken for soft tissue recovery as expected by surgeons. 18 months is the maximum end point that surgeons would suggest for recovery time.</p> |

| | | | |
|-------------------------|---|---------------------|---|
| | <p>outcome for her” (Amelia, 2-stage)</p> <p>“I would’ve thought that if you wanted to go back to work...you wouldn’t be very happy [having a 2-stage operation] because you wouldn’t be able to do nothing for those six weeks again. Then you’d have to go all through it again after three months of having it done. Six weeks is only a month and a half, and then in another month and a half you’re having it all done again.” (Jim, 1-stage)</p> | | |
| Antibiotic side effects | <p>“...the nightmare on heavy antibiotics, toiletry wise. Now I’ve had to move into a, another bed. My wife and I are married 50 plus</p> | 1) Affects me a lot | Antibiotics are an essential method of attempting to ensure the |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

| | | |
|--|--------------------------------|--|
| <p>years, and I have to have my own room because I'm getting up in the night." (Rory, 1-stage)</p> <p>"...I stayed on antibiotics then for, for ever...and after the first [week of antibiotics] I was just dying. I just wanted to lie on the floor and die. I felt so sick. So ill." (Lottie, 2-stage)</p> <p>"I felt fine. When I was going to see the surgeon... They'd say, 'How are you today?' I'd say, 'I feel fine, fit! [laughs] I feel well in myself, I eat well.'" (Ray, 2-stage)</p> | <p>2) Don't affect me much</p> | <p>periprosthetic joint infection is treated and subsequently clear. For some patients, the impact of these antibiotics is significant, while for others there are less severe side effects.</p> |
|--|--------------------------------|--|

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

| OPTION H1 | OR | OPTION H2 |
|--|----|--|
| I take antibiotics and the side effects don't affect me much <i>and</i> I have two operations <i>and</i> After my new hip is fitted, I can do most of the things that I want to do <i>and</i> After my surgical treatment starts, it takes 6 months to get back to the things that I normally do | OR | I take antibiotics and the side effects affect me a lot <i>and</i> I have one operation <i>and</i> After my new hip is fitted, I cannot do most of the things that I want to do <i>and</i> After my surgical treatment starts, it takes 12 months to get back to the things that I normally do |
| Please only tick ONE box: | | |
| OPTION H1 <input type="checkbox"/> | | OPTION H2 <input type="checkbox"/> |

Figure 1 - example profile
338x190mm (96 x 96 DPI)

BMJ Open

What are patients' preferences for revision surgery after periprosthetic joint infection? A discrete choice experiment

| | |
|---------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2019-031645.R1 |
| Article Type: | Original research |
| Date Submitted by the Author: | 30-Sep-2019 |
| Complete List of Authors: | Carroll, Fran; University of Bristol, Population Health Sciences, Bristol Medical School Goberman-Hill, Rachael; University of Bristol, Translational Health Sciences, Bristol Medical School Strange, Simon; University of Bristol, Translational Health Sciences, Bristol Medical School Blom, AW; University of Bristol, Translational Health Sciences, Bristol Medical School Moore, Andrew; University of Bristol, Translational Health Sciences, Bristol Medical School |
| Primary Subject Heading: | Surgery |
| Secondary Subject Heading: | Health services research, Research methods |
| Keywords: | Discrete Choice Experiment, Hip < ORTHOPAEDIC & TRAUMA SURGERY, periprosthetic infection, ORTHOPAEDIC & TRAUMA SURGERY |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3 1 What are patients' preferences for revision surgery after periprosthetic joint infection? A discrete
4
5 2 choice experiment
6
7

8
9 3 **Authors:**

10
11
12 4 F E Carroll

13
14
15 5 R Gooberman-Hill

16
17
18 6 S Strange

19
20
21 7 A W Blom

22
23
24 8 A J Moore

25
26
27 9 Department of Translational Health Sciences, Bristol Medical School, University of Bristol.

28
29
30 10 Corresponding author: Dr Andrew J Moore. Translational Health Sciences (Southmead),
31
32 11 Musculoskeletal Research Unit, Translational Health Sciences, Bristol Medical School, University of
33
34 12 Bristol, Learning and Research Building, Level 1, Southmead Hospital, Bristol, BS10 5NB, UK. Email:
35
36 13 a.j.moore@bristol.ac.uk. Telephone: 0117 414 7877
37
38

39 14 Word count: 3088
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Abstract

2 Objectives: Understanding patients' preferences for treatment is crucial to provision of good care
3 and shared decisions, especially when more than one treatment option exists for a given condition.

4 One such condition is infection of the area around the prosthesis after hip replacement, which
5 affects between 0.4 and 3% of patients. There is more than one treatment option for this major
6 complication and our study aimed to assess the value that patients place on aspects of revision
7 surgery for periprosthetic hip infection.

8 Design: We identified four attributes of revision surgery for periprosthetic hip infection. Using a
9 discrete choice experiment (DCE) we measured the value placed on each attribute by 57 people who
10 had undergone either one- or two-stage revision surgery for infection.

11 Setting: The DCE was conducted with participants from 9 National Health Service hospitals in the
12 United Kingdom.

13 Participants: Adults who had undergone revision surgery for periprosthetic hip infection (n=57).

14 Results: Overall, the strongest preference was for a surgical option that resulted in no restrictions on
15 engaging in valued activities after a new hip is fitted ($\beta=0.7$). Less valued, but still important
16 attributes, included a shorter time taken from the start of treatment to return to normal activities (6
17 months; $\beta=0.3$), few or no side effects from antibiotics ($\beta=0.2$), and having only one operation
18 ($\beta=0.2$).

19 Conclusions: Results highlight that people who have had revision surgery for periprosthetic hip
20 infection most value aspects of care that affect their ability to engage in normal everyday activities.

21 These were the most important characteristics in decisions about revision surgery.

22

1 Keywords

2 *Discrete choice experiment; hip; periprosthetic infection; orthopaedic surgery*

3 Strengths and limitations of this study

- 4 • This is the first study to quantify the value that patients place on different aspects of revision
5 surgery for prosthetic hip joint infection.
- 6 • Using a rigorous process involving empirical qualitative research we identified four attributes
7 for the DCE.
- 8 • It was feasible for participants to complete the questionnaires, meaning the results were as
9 expected with patients selecting a combination of options reflecting their preferred choice.
- 10 • While, the sample size of 57 participants sufficiently powered the analyses presented here, it
11 is too small to conduct any subgroup analyses.
- 12 • The majority of the study sample were white, male and educated, so results may not reflect
13 the preferences of the wider surgical population.

15 Introduction

16 For people with osteoarthritis, hip replacement is a common procedure that aims to improve
17 function and reduce pain. In 2016 over 100,000 hip replacement procedures were conducted in the
18 UK. With an ageing population, rates are predicted to rise ¹. Although successful for many people,
19 approximately 0.4 -1% of patients who have undergone primary hip replacement ^{2,3} and 2-3% of
20 patients undergoing revision hip replacement ⁴ develop deep periprosthetic joint infection (PJI)
21 severe enough to warrant surgical revision. Patients with PJI find it devastating. Symptoms include

1
2
3 1 severe pain, inflammation, discharge from the surgical wound, fever, nausea, malaise, reduction in
4
5 2 or loss of function, dislocation, and if left untreated can lead to disability and/or death ⁵.
6
7
8 3 PJI is extremely challenging to treat. Treatment can include ‘debridement, antibiotic treatment and
9
10 4 implant retention’ (DAIR), used in 7.6% of cases, or more commonly major revision surgery, which
11
12 5 involves removal of the infected prosthesis, radical debridement of infected tissue and
13
14 6 reimplantation (“fitting”) of a new prosthesis with subsequent antibiotic treatment ⁴. Revision
15
16 7 surgery can be provided in a single operation (one-stage) or as a staged operation (two-stage) where
17
18 8 the infected implant is removed and the patient is left without an implant (with or without a
19
20 9 temporary spacer) while receiving antibiotic treatment. In the staged process reimplantation of the
21
22 10 new implant is delayed, commonly for up to 6 months, but in some instances over 12 months later ⁵.
23
24 11 ⁶. There is no clear evidence that either a one or two-stage strategy is superior in eradicating
25
26 12 infection ^{7,8}, but qualitative research has shown that two-stage revision places a greater burden on
27
28 13 patients and families than one-stage revision. This burden is due to the extended period of
29
30 14 immobility in between operations, complications associated with the period of immobility, and deep
31
32 15 psychological distress, with some patients reporting depression and suicidal thoughts ⁶.
33
34
35 16 Surgeons’ decisions about which type of revision surgery is most appropriate for an individual
36
37
38 17 patient take into account many factors. Our previous work indicates that decisions are based on a
39
40 18 combination of a surgeon’s own training and clinical experience of different techniques; the
41
42 19 availability of hospital infrastructure such as microbiology services to quickly identify the infecting
43
44 20 organism; characteristics of the infecting organism and duration of infection; patient characteristics
45
46 21 such as age, comorbidities, frailty, and extent of damaged tissue; and published evidence of revision
47
48 22 techniques, or reports by senior colleagues. Surgeons also considered patients’ preferences for
49
50 23 surgery, although this could often involve the choice between long-term suppressive antibiotics or
51
52 24 surgery ⁹. There is no quantitative evidence that characterises patients’ preferences for one-stage or
53
54 25 two-stage revision surgery in this context and this is an important area of work to be investigated.
55
56
57
58
59
60

1
2
3 1 The aim of this study is to assess the surgical preferences of patients who underwent revision
4
5 2 surgery for prosthetic hip joint infection.
6
7

8 3
9
10
11 4 This study is part of a larger programme of research which aims to improve outcomes for patients
12
13 5 after periprosthetic joint infection (NIHR PGfAR: RP-PG-1210-12005). The programme includes a
14
15 6 randomised clinical trial comparing one-stage and two-stage revision surgery for prosthetic hip joint
16
17 7 infection¹⁰ within which this study was embedded.
18
19
20
21 8

22 23 24 9 **Methods:**

25 26 27 28 29 10 **Discrete choice experiments (DCEs)**

30
31
32
33 11 We undertook a discrete choice experiment (DCE) to quantify the surgical preferences of patients
34
35 12 who underwent revision surgery for prosthetic hip joint infection. The DCE was embedded within a
36
37 13 randomised clinical trial that evaluated clinical and cost effectiveness of revision surgery for
38
39 14 prosthetic hip joint infection and compared one-stage revision surgery with two-stage revision
40
41 15 surgery. Patients who were taking part in the trial were eligible for the DCE. Ethical approval for the
42
43 16 study was granted by NRES Committee South West – Frenchay on 31 December 2014 (14/SW/1166).
44
45 17 All participants provided written informed consent to take part in the DCE study.
46
47
48
49 18

50
51 19 DCEs are an established method in health services research and have been used to explore a range
52
53 20 of health-related services and treatments¹¹⁻¹⁴. DCEs involve asking respondents to choose between
54
55 21 hypothetical scenarios which describe goods or services, where a 'service' may mean an intervention
56
57 22 or an approach to care. The method aims to establish what attributes of that service influence their
58
59 23 decision-making and to what extent. This enables quantification of the marginal impact of these

1
2
3 1 attributes. Scenarios within a DCE describe the service of interest (in this case, revision joint
4
5 2 replacement) using the same set of attributes, but at different levels in each scenario. Choices
6
7 3 between scenarios, or whether to accept or reject a scenario, are used to estimate the influence and
8
9 4 value of the different attribute levels.
10
11
12 5
13
14

15 6 Questionnaire development

16
17
18
19 7 When a patient faces revision surgery for prosthetic hip infection, a one-stage or a two-stage
20
21 8 operation is required, and the decision about which is undertaken is made largely at the discretion
22
23 9 of the surgeon, taking into account patient preferences for treatment. The DCE study was designed
24
25 10 to engage patients in their preferences for the features associated with these two surgical options,
26
27 11 given neither is currently known to be clinically superior in terms of patient outcomes.
28
29
30 12
31
32 13 Qualitative methods are recommended for DCE attribute development because they enable
33
34 14 conceptual development of attributes directly from people's experiences and so better reflect the
35
36 15 issues that are likely to matter most to people when making a decision ¹⁵. For this study we
37
38 16 developed attributes from our previous qualitative study which explored impact of prosthetic joint
39
40 17 infection and its treatment on patients and their recovery process; one-to-one interviews were
41
42 18 conducted with 19 patients with PJI, focusing on the impact of PJI and surgical treatment. The
43
44 19 interviews were audio-recorded, transcribed and the qualitative data set was analysed thematically
45
46 20 ⁶. Levels of attributes were assigned by refining the language used to convey the meaning of the
47
48 21 attributes, and particularly where some quantification of an attribute was mentioned by participants
49
50 22 during qualitative interview, as illustrated in Table 1 ¹⁵.
51
52
53 23
54
55 24 Table 1 describes the attribute and level selection based on the earlier qualitative work, including
56
57 25 illustrative quotations and rationale.
58
59
60

1
2
3 1
4
5 2 The final questionnaire comprised a single scenario task in which respondents were asked to imagine
6
7 3 this was the first time that they had been offered the two surgical options presented and were asked
8
9 4 to consider them carefully before selecting the one (from a pair) they would prefer.
10
11
12 5

13
14 6 With a $4 \times 4 \times 2 \times 2$ design (two four-level and two two-level attributes), a total of 64 different
15
16 7 combinations of attribute levels (profiles) is possible. This converts to 32 pairs of profiles, which as a
17
18 8 'full factorial' was considered to be too large for participants to complete ¹⁶. An orthogonal main
19
20 9 effects plan was therefore used to reduce the number of choice sets to 16 ¹⁷. Each profile was
21
22 10 presented with its pair, and participants were required to select which option they preferred (see
23
24 11 Figure 1).
25
26
27
28 12

30 13 Patient and Public Involvement

31
32
33
34 14 The questionnaire was piloted and refined in collaboration with 5 patient and public involvement
35
36 15 representatives. At an initial meeting of representatives, group members were involved in
37
38 16 questionnaire development and suggested improvements to its formatting to aid readability, clearer
39
40 17 phrasing of the questions to avoid ambiguity, and shortening of the instruction leaflet for clarity. At a
41
42 18 subsequent meeting, members of the group completed the questionnaire and fed back on their
43
44 19 experience. They completed the questionnaire without assistance and felt the instructions were
45
46 20 clear but suggested that key points in the questions should be highlighted and that a contact
47
48 21 telephone number should be added to enable participants to seek assistance if needed. For
49
50 22 participants in the study, a summary of findings will be sent to those who indicated that they wished
51
52 23 to be informed.
53
54
55
56
57
58
59
60

1
2
3
4 1
5
6
78 2 **Participants**
9

10
11
12 3 Patients who were recruited into the INFORM randomised controlled trial (ISRCTN10956306)
13
14 4 received the discrete choice task after completing the 18-month primary outcome measure. The
15
16 5 questionnaire was either posted to participants or completed in person with the assistance of a
17
18 6 research nurse during a hospital clinic visit.
19
20
21 7

22
23
24 8 **Data analysis**
25
26
27

28 9 Sample size calculations for DCEs are challenging due to dependence on the true parameter values
29
30 10 estimated in the choice model ¹⁸. However, reliable statistical analysis has been demonstrated with
31
32 11 sample sizes of 40–120 respondents and combined with the rarity of periprosthetic joint infection
33
34 12 (1% of those undergoing hip replacement), a sample size above 50 participants for the DCE was
35
36 13 deemed adequate to obtain sufficient data for exploratory analysis and interpretation ^{19, 20}.
37
38
39 14

40
41 15 Paper questionnaires were distributed to participants as part of the follow-up data collection in the
42
43 16 INFORM trial. Questionnaires were returned to the study team between January 2017 and
44
45 17 November 2018 and data entered onto a Microsoft Excel spreadsheet. DCE data were effects coded
46
47 18 and analysed using STATA SE 15 ^{21, 22}. The influence of the four attributes on patient choices was
48
49 19 analysed using a conditional logit model. As attribute levels are effects coded, the mean of all
50
51 20 coefficients is 0 across each attribute. The effects coded preference weights (coefficients) produced
52
53 21 by the conditional logit model are estimated relative to the mean effect of the attribute, with the p
54
55 22 value indicating the statistical significance of “the difference between the estimated preference
56
57 23 weight and the mean effect of the attribute” ²³.
58
59
60

1

2 **Results:**3 **Participants**

4 Of the 80 discrete choice questionnaires provided to trial participants, 57 were returned fully
 5 complete (71%) from patients from 9 trial sites. Data from 6 questionnaires was not used in analysis
 6 as they were partially or totally incomplete. Responding participants had a mean age of 70 (range 51
 7 to 90), 21 (37%) were female, 26 (46%) had undergone 1-stage revision, 14 (25%) lived alone and 41
 8 (72%) were retired from work. Table 2 provides further demographic information.

9

10 *Table 2 – Demographic and Clinical characteristics of the DCE Respondents*

| Characteristic | Participants (N=57) |
|--------------------------------|---------------------|
| Age – Yrs [mean] (range) | 70 (51-90) |
| Gender [number] (%) | |
| Male | 36 (63) |
| Female | 21 (37) |
| Ethnicity [number] (%) | |
| White | 55(96) |
| Black | 1 (2) |
| Mixed | 1 (2) |
| Marital Status [number] (%) | |
| Married / Partner | 42 (74) |
| Divorced / Separated / Widowed | 12 (21) |
| Single | 3 (5) |

| | |
|--|---------|
| Living Arrangements [number] (%) | |
| With Partner / somebody else | 43 (75) |
| Alone | 14 (25) |
| Schooling/Education [number] (%) | |
| Left at normal school leaving age | 35 (61) |
| Left after normal school leaving age | 15 (26) |
| Left before normal school leaving age | 7 (12) |
| Work Situation [number] (%) | |
| Retired | 41 (72) |
| Working / Sick leave | 14 (25) |
| Unemployed | 2 (4) |
| Surgery Received for Prosthetic Hip Joint Infection [number] (%) | |
| 2-stage revision | 31 (54) |
| 1-stage revision | 26 (46) |

1

2 Discrete choice experiment

3 Table 3 shows the regression coefficients and the results from the 57 patients who had fully
4 completed the discrete choice questionnaire.

5

6 Table 3 – discrete choice task results from conditional logistic regression

| <i>Attribute</i> | <i>Level</i> | <i>Coefficient</i> | <i>Standard error</i> | <i>95% confidence interval</i> | <i>P value</i> |
|------------------|--------------|--------------------|---------------------------|--|----------------|
| | | | | | |

58

59

60

| | | | | | | |
|--|---|-----------------------|-------|-----|----------------|--------|
| 1 2 3 4 5 6 7 8 9 | Ability to engage in valued activities after new hip is fitted. | Can do everything* | 0.70 | | | |
| 10 11 12 13 14 | | Can do most things | 0.49 | .08 | .33 to .64 | <0.001 |
| 15 16 17 18 | | Cannot do most things | -0.39 | .07 | -.53 to -.24 | <0.001 |
| 19 20 21 22 23 24 25 | | Cannot do anything | -0.80 | .13 | -1.05 to -0.55 | <0.001 |
| 26 27 28 29 | Antibiotic side effects | Don't affect me much* | 0.22 | | | |
| 30 31 32 33 34 35 36 | | Affects me a lot | -0.22 | .05 | -.33 to -.12 | <0.001 |
| 37 38 39 40 41 42 43 | Number of operations | 1* | 0.20 | | | |
| 44 45 46 47 48 49 50 51 52 | | 2 | -0.20 | .07 | -.35 to -.06 | <0.001 |
| 53 54 55 56 57 58 59 60 | Time taken after surgical treatment starts, to return to normal activities. | 3months* | 0.20 | | | |
| | | 6months | 0.31 | .09 | .14 to .48 | <0.001 |
| | | 12months | -0.06 | .05 | -.15 to .04 | 0.22 |
| | | 18months | -0.45 | .10 | -.64 to -.26 | <0.001 |

| | | | | | |
|---|--|--|--|--|--|
| | | | | | |
| <i>*Indicates reference category within attribute</i> | | | | | |

1

2

3 Analysis indicates that participants had the strongest preference for a surgical option that resulted in
 4 the least restrictions on engaging in valued activities after the new hip is fitted, illustrated by the
 5 largest preference weight. Other less valued but important preferences were for a surgical strategy
 6 that would result in a shorter time after surgical treatment starts to return to normal activities, few
 7 or no side effects from antibiotics, and only one operation. The results also suggest that the least
 8 restrictions on engaging in valued activities, and the shortest time taken to return to normal activity
 9 are the individual attributes most valued by patients in this sample. This is indicated by the larger
 10 spread of coefficients (i.e. more 'value' is placed on changes in these attributes). The most
 11 acceptable option was a time period of between 3 to 6 months to return to normal activity, however
 12 there is no clear preference up to 12 months, although 18 months appeared to be significantly
 13 disfavoured.

14

15 Discussion:

16 This study aimed to investigate and understand patients' preferences for aspects of revision surgery
 17 for periprosthetic joint infection. Four relevant attributes were identified through earlier qualitative
 18 work, and quantitatively, patients in this study most value the ability to engage in valued activities
 19 and the time taken to return to normal daily activities. This reflects the findings of our previous
 20 qualitative work which shows that although both revision strategies impacted greatly on patients
 21 and their families, patients receiving two-stage revision surgery experience particularly long periods
 22 of immobility and social isolation. This was often followed by a protracted recovery period, which
 23 could leave patients much less able than before their primary operation, and some patients

1
2
3 1 experienced profoundly negative psychological effects associated with physical suffering, loss of
4
5 2 dignity and independence ⁶. It appears that for patients in our sample, 3-6 months to return to
6
7 3 normal activity was preferable, although there was no significant difference up to 12 months, but 18
8
9 4 months was disfavoured. This suggests that the acceptable margin of recovery for patients is up to
10
11 5 12 months after their receipt of a new hip joint.
12
13
14 6

15
16 7 Discrete choice methodology can be challenging for participants because the format of questions is
17
18 8 different to standard surveys and items can seem repetitive. We collected feedback from the first 11
19
20 9 participants who completed the questionnaire. We found that those participants who were
21
22 10 supported by a research nurse when completing the questionnaire were more likely to complete it
23
24 11 and return it, compared with those who received the questionnaire by post and completed it alone.
25
26 12 Participant feedback suggested that the questionnaire was difficult to complete, as the scenarios
27
28 13 were similar and appeared to be repetitive. To address this, we amended the questionnaire format
29
30 14 and instructions and offered participants support either face-to-face or by phone with one of the
31
32 15 study research nurses. Nurses were then able to answer queries about the questionnaire and offer
33
34 16 support if needed. The results suggest the group completed the questionnaire in a rational and
35
36 17 logical way, meaning the results were as expected with patients selecting an optimal combination of
37
38 18 options as their preferred choice. This is an important methodological finding because although our
39
40 19 study demonstrates the feasibility of the DCE method with this population, others conducting similar
41
42 20 studies with older, ill populations could consider in advance the need for professional support in the
43
44 21 completion of discrete choice questionnaires.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

22
23 23 Participants were all individuals who participated in a clinical trial and had already undergone
24
25 24 revision surgery for periprosthetic joint infection. This meant that the choices that participants were
26
27 25 asked to make in the questionnaire were based on scenarios unlikely to reflect their real-life
28
29 26 experiences as in reality such choices would not be available to them, since decisions about surgical

1 strategies are based on a wider variety of clinical, surgeon, patient and organisational factors⁹. Also,
2 patients were not being faced with these decisions at the time of questionnaire completion as we
3 decided that it would be unethical to ask patients awaiting treatment to complete a DCE about
4 surgical options in a hypothetical context, at a time when they may be particularly vulnerable. To do
5 so would not meet an ethical standard of protection from harm as it might mean that patients were
6 inadvertently led to believe that there were more or different options available to them than were
7 clinically indicated at that time.

8
9 We also found that some participants reported difficulty separating their own recent personal
10 experience of revision surgery from the hypothetical scenarios presented in the questionnaire, as
11 they found it hard to imagine receiving a treatment option that differed from the one that they had
12 received. In terms of methodology, many DCE's are conducted with participants who already have
13 some experience of the treatment attributes under investigation^{12, 19, 20, 24}. In our study previous
14 experience meant that participants had some appreciation of the attributes being tested. While, the
15 sample size of 57 participants provided sufficient data for the analyses presented here, the sample
16 size was too small to conduct any subgroup analysis to identify whether preferences would differ
17 between participants who had received one or two stage revision. Similarly, patients who are older,
18 still working or live alone may have had stronger preferences for a one-stage operation than those
19 who are younger, retired and have support at home to cope with a two-stage procedure. The
20 majority of the study sample were also white, male and educated, which means that results may not
21 reflect the preferences of the wider surgical population. Further research could explore preferences
22 in a more diverse population.

23
24 This work has provided an initial and important first step in understanding patients' preferences for
25 characteristics associated with revision surgery for periprosthetic infection. It is important that

1 orthopaedic healthcare professionals discuss these attributes with patients when discussing options
2 for surgical and antibiotic treatment for periprosthetic infection.

4 Implications

5 Results of this study offer insight into the preferences of patients for revision surgery and provide
6 valuable information to surgeons from all disciplines. Although factors affecting patient preferences
7 for surgery differ to those valued by medical professionals, consideration should be given to such
8 factors in order to aid shared decision-making where clinical equipoise between options exists.

9
10 Previous research using discrete choice approaches has explored patient preferences in a surgical
11 context. This has included examination of preferences for surgical versus non-surgical interventions
12 in a range of conditions, such as oesophageal cancer and ulcerative colitis ^{24, 25}. Some research has
13 explored preferences for conditions in which there are two surgical options, including for ectopic
14 pregnancy, vaginal wall prolapse, and osteoarthritis ^{26, 27}. Across all of these conditions, evidence
15 suggests that patients choose options that reduce the need for further surgery or operations, have a
16 shorter recovery time, lower risk of symptom recurrence and improve ability to preserve existing
17 joint motion in the case of osteoarthritis. Findings from the current study are similar in that patients
18 prefer a surgical option that reduces the number of operations, recovery time and the side effects of
19 antibiotics. However, in the setting of infected joint replacement, patients placed highest value on
20 restoration of function. This was more important to the patients in our study than the number of
21 operations they would have to undergo. Although our study focused on the preferences between
22 one and two-stage surgery surgeons may need to consider these preferences during shared
23 decision-making about all options for revision surgery for prosthetic hip infection, including the role
24 of debridement with retention of implants (DAIR). Although DAIR is only efficacious in approximately
25 60% of cases, it is associated with a quicker return to valued activities and improved joint function ²⁸.

1
2
3 1
4
5

6 2 Conclusions

7
8
9 3 Our results show that the most valued characteristics in decisions about revision surgery for
10
11 4 prosthetic hip infection were the ability to engage in valued activities and time taken to return to
12
13 5 normal activity. This builds on the findings of our previous qualitative work which shows that
14
15 6 although both revision strategies impact greatly on patients and their families' everyday lives,
16
17 7 patients receiving two-stage revision surgery experience particularly long periods of immobility and
18
19 8 social isolation⁶. The desire to return to everyday activities should be taken into account when
20
21 9 surgeons are discussing options with patients, particularly when there is equipoise from a surgical
22
23 10 perspective about the options available, and when the decision is 'preference sensitive'.
24
25
26
27

28 11 Funding

29
30
31
32 12 This study is funded by the National Institute for Health Research Programme Grants for Applied
33
34 13 Research (NIHR PGfAR) programme (grant number: RP-PG-1210-12005) and supported by the NIHR
35
36 14 Comprehensive Clinical Research Network (CRN). This study was also supported by the NIHR
37
38 15 Biomedical Research Centre at the University Hospitals Bristol NHS Foundation Trust and the
39
40 16 University of Bristol. The views expressed are those of the authors and not necessarily those of the
41
42 17 NIHR or the Department of Health and Social Care.
43
44
45
46
47

48 18 Author Contributions

49
50
51 19 FEC: Contributed to design, analysis and interpretation of data, drafting and revising the manuscript.
52
53 20 RG-H: Contributed to conception, design, interpretation of data, drafting and revising the
54
55 21 manuscript. SS: Contributed to design, data collection and revising the manuscript. AWB:
56
57 22 Contributed to the conception, design, interpretation of data and revising the manuscript. AJM:
58
59
60

1
2
3 1 Contributed to the design, interpretation of data, drafting and revising the manuscript. All authors
4
5 2 gave final approval of the submitted manuscript.
6
7
8

9 3 Competing Interests

10
11
12
13 4 None declared
14
15
16

17 5 Data Availability

18
19
20
21 6 Anonymised research data will be made available at the University of Bristol Research Data
22
23 7 Repository (data.bris) beginning 1 year after publication of the results. Access will be
24
25 8 granted to bona fide researchers after the University of Bristol Data Access Committee has
26
27 9 approved their request. Please contact data-bris@bristol.ac.uk in the first instance.
28
29
30
31

32 10 Acknowledgements

33
34
35
36 11 The authors thank the participants who gave their time to complete a discrete choice questionnaire.
37
38 12 We also thank Dr Erik Lenguerand for providing demographic data for the study sample and advice
39
40 13 on presentation of statistical results. We would also like to thank the study administration and
41
42 14 management team: Makita Werrett and Beverley Evanson for their invaluable support. The authors
43
44 15 would also like to thank the INFORM Research Advisory Group (I-RAG) and the Patient Experience
45
46 16 Partnership in Research (PEP-R) at the University of Bristol for their contributions to the refinement
47
48 17 of the questionnaire.
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 – qualitative support for attributes included in the discrete choice questionnaire

| Attribute | Evidence of attribute inclusion, with pseudonym and surgery type | Levels | Rationale for levels |
|----------------------|--|--|---|
| Number of operations | <p>“There’s no way I want two more big operations now at my time of life. You do it all or not at all...” “I said there was no way I wanted two ops” (Harriet, 1-stage)</p> <p>“Of course, emotionally, you want it over and done with as soon as possible...but ultimately that has to be done in the correct way. There’s the tortoise and hare situation. There’s absolutely no point in rushing ahead if</p> | <p>1) 1 operation</p> <p>2) 2 operations</p> | <p>Two types of revision surgery are currently provided in healthcare, and involve either one or two operations</p> |

| | | | |
|--|--|--|--|
| | <p>ultimately it's going to fail.” (Maggie, 2-stage)</p> | | |
| <p>Ability to engage in valued activities after new hip is fitted.</p> | <p>“Fourteen months without a hip joint so it meant that I couldn't drive a car, I couldn't do anything that I'd been used to doing, playing golf or doing anything. Well, I gave up golf actually after the first revision” (Don, 2-stage)</p> <p>“But when I for example went to, on holiday recently and I had serious problems getting into the bath to stand in the shower. Because I, I couldn't get in. And in a wet – on a wet surface and that, I'm very conscious of not falling in. I can't afford to fall. So, pain I've got</p> | <ol style="list-style-type: none"> 1) Can do everything 2) Can do most things 3) Cannot do most things 4) Cannot do anything | <p>Following revision surgery, the ability to engage in valued activities can be reduced in a major, or somewhat more minor way. These levels capture variation in ability identified by patients.</p> |

| | | | |
|--|--|--|--|
| | <p>none, stiffness none.</p> <p>Physical function limitations, and that's one of them." (Rory, 1-stage)</p> <p>"My aim has always been to get back on my feet as soon as I can, and to walk as good as I can, and that's a big disappointment. I'm not where I think I should have been" (Robert, 2-stage)</p> | | |
| <p>Time taken after surgical treatment starts, to return to normal activities.</p> | <p>"I didn't want to go 14 weeks with effectively one leg. What was worse was not knowing that I had to endure all of those weeks not knowing that I was ever going to get another hip joint back." (Maggie, 2-stage)</p> | <ol style="list-style-type: none"> 1) 3 months 2) 6 months 3) 12 months 4) 18 months | <p>These time intervals demonstrate best approximations and a reflection of the need to ensure normal expectations of time taken for soft tissue recovery as</p> |

| | | | |
|---|--|--|--|
| <p>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60</p> | <p>“If I had known how hard it was going to be for her to walk in that interim six months, and if there seemed to have been a reasonable, or a good possibility that the infection would be nuked in a one stage, then that might have been a better outcome for her” (Amelia, 2-stage)</p> <p>“I would’ve thought that if you wanted to go back to work...you wouldn’t be very happy [having a 2-stage operation] because you wouldn’t be able to do nothing for those six weeks again. Then you’d have to go all through it again after three months of having it done. Six weeks is only a month and a half, and then</p> | | <p>expected by surgeons. 18 months is the maximum end point that surgeons would suggest for recovery time.</p> |
|---|--|--|--|

| | | | |
|-------------------------|---|---|--|
| | <p>in another month and a half you're having it all done again." (Jim, 1-stage)</p> | | |
| Antibiotic side effects | <p>"...the nightmare on heavy antibiotics, toiletry wise. Now I've had to move into a, another bed. My wife and I are married 50 plus years, and I have to have my own room because I'm getting up in the night." (Rory, 1-stage)</p> <p>"...I stayed on antibiotics then for, for ever...and after the first [week of antibiotics] I was just dying. I just wanted to lie on the floor and die. I felt so sick. So ill." (Lottie, 2-stage)</p> <p>"I felt fine. When I was going to see the surgeon...</p> | <p>1) Affects me a lot</p> <p>2) Don't affect me much</p> | <p>Antibiotics are an essential method of attempting to ensure the periprosthetic joint infection is treated and subsequently clear. For some patients, the impact of these antibiotics is significant, while for others there are less severe side effects.</p> |

| | | | |
|--|---|--|--|
| | They'd say, 'How are you today?' I'd say, 'I feel fine, fit! [laughs] I feel well in myself, I eat well.'" (Ray, 2-stage) | | |
|--|---|--|--|

References

1. National Joint Registry. National Joint Registry 14th Annual Report 2017: National Joint Registry for England, Wales, Northern Ireland and the Isle of Man; 2017 [Available from: www.njrreports.org.uk.
2. Blom AW, Taylor AH, Pattison G, Whitehouse S, Bannister GC. Infection after total hip arthroplasty. The Avon experience. The Journal of bone and joint surgery British volume. 2003;85(7):956-9.
3. Phillips JE, Crane TP, Noy M, Elliott TS, Grimer RJ. The incidence of deep prosthetic infections in a specialist orthopaedic hospital: a 15-year prospective survey. The Journal of bone and joint surgery British volume. 2006;88(7):943-8.
4. Lenguerrand E, Whitehouse MR, Beswick AD, Jones SA, Porter ML, Blom AW. Revision for prosthetic joint infection following hip arthroplasty: Evidence from the National Joint Registry. Bone & joint research. 2017;6(6):391-8.
5. Zimmerli W, Trampuz A, Ochsner PE. Prosthetic-joint infections. The New England journal of medicine. 2004;351(16):1645-54.
6. Moore AJ, Blom AW, Whitehouse MR, Goberman-Hill R. Deep prosthetic joint infection: a qualitative study of the impact on patients and their experiences of revision surgery. BMJ open. 2015;5(12):e009495.
7. Kunutsor SK, Whitehouse MR, Blom AW, Beswick AD. Re-Infection Outcomes following One- and Two-Stage Surgical Revision of Infected Hip Prosthesis: A Systematic Review and Meta-Analysis. PloS one. 2015;10(9):e0139166.
8. Kunutsor SK, Whitehouse MR, Blom AW, Board T, Kay P, Wroblewski BM, et al. One- and two-stage surgical revision of peri-prosthetic joint infection of the hip: a pooled individual participant data analysis of 44 cohort studies. European journal of epidemiology. 2018;33(10):933-46.
9. Moore AJ, Blom AW, Whitehouse MR, Goberman-Hill R. Managing uncertainty - a qualitative study of surgeons' decision-making for one-stage and two-stage revision surgery for prosthetic hip joint infection. BMC musculoskeletal disorders. 2017;18(1):154.
10. Strange S, Whitehouse MR, Beswick AD, Board T, Burston A, Burston B, et al. One-stage or two-stage revision surgery for prosthetic hip joint infection--the INFORM trial: a study protocol for a randomised controlled trial. Trials. 2016;17:90.
11. Louviere J, Street DJ, Burgess L. A 20+ years' retrospective on choice experiments. Marketing research and modeling: Progress and prospects: Springer; 2004. p. 201-14.

12. de Bekker-Grob EW, Hofman R, Donkers B, van Ballegooijen M, Helmerhorst TJ, Raat H, et al. Girls' preferences for HPV vaccination: a discrete choice experiment. *Vaccine*. 2010;28(41):6692-7.
13. Liede A, Mansfield CA, Metcalfe KA, Price MA, Snyder C, Lynch HT, et al. Preferences for breast cancer risk reduction among BRCA1/BRCA2 mutation carriers: a discrete-choice experiment. *Breast cancer research and treatment*. 2017;165(2):433-44.
14. van Dijk JD, Groothuis-Oudshoorn CG, Marshall DA, MJ IJ. An Empirical Comparison of Discrete Choice Experiment and Best-Worst Scaling to Estimate Stakeholders' Risk Tolerance for Hip Replacement Surgery. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2016;19(4):316-22.
15. Coast J, Al-Janabi H, Sutton EJ, Horrocks SA, Vosper AJ, Swancutt DR, et al. Using qualitative methods for attribute development for discrete choice experiments: issues and recommendations. *Health economics*. 2012;21(6):730-41.
16. Marshall D, Bridges JF, Hauber B, Cameron R, Donnalley L, Fyie K, et al. Conjoint Analysis Applications in Health - How are Studies being Designed and Reported?: An Update on Current Practice in the Published Literature between 2005 and 2008. *The patient*. 2010;3(4):249-56.
17. Hedayat AS, Sloane NJA, Stufken J. *Orthogonal arrays: theory and applications*: Springer Science & Business Media; 2012.
18. Lancsar E, Louviere J. Conducting discrete choice experiments to inform healthcare decision making: a user's guide. *PharmacoEconomics*. 2008;26(8):661-77.
19. Salkeld G, Solomon M, Butow P, Short L. Discrete-choice experiment to measure patient preferences for the surgical management of colorectal cancer. *The British journal of surgery*. 2005;92(6):742-7.
20. de Bekker-Grob EW, Essink-Bot ML, Meerding WJ, Koes BW, Steyerberg EW. Preferences of GPs and patients for preventive osteoporosis drug treatment: a discrete-choice experiment. *PharmacoEconomics*. 2009;27(3):211-9.
21. Bech M, Gyrd-Hansen D. Effects coding in discrete choice experiments. *Health economics*. 2005;14(10):1079-83.
22. StataCorp. *Stata Statistical Software: Release 15*. College Station, TX: StataCorp LLC; 2017.
23. Hauber AB, Gonzalez JM, Groothuis-Oudshoorn CG, Prior T, Marshall DA, Cunningham C, et al. Statistical Methods for the Analysis of Discrete Choice Experiments: A Report of the ISPOR Conjoint Analysis Good Research Practices Task Force. *Value in health : the journal of the International Society for Pharmacoeconomics and Outcomes Research*. 2016;19(4):300-15.
24. Noordman BJ, de Bekker-Grob EW, Coene P, van der Harst E, Lagarde SM, Shapiro J, et al. Patients' preferences for treatment after neoadjuvant chemoradiotherapy for oesophageal cancer. *The British journal of surgery*. 2018;105(12):1630-8.
25. Bewtra M, Kilambi V, Fairchild AO, Siegel CA, Lewis JD, Johnson FR. Patient preferences for surgical versus medical therapy for ulcerative colitis. *Inflammatory bowel diseases*. 2014;20(1):103-14.
26. van Mello NM, Mol F, Opmeer BC, de Bekker-Grob EW, Essink-Bot ML, Ankum WM, et al. Salpingotomy or salpingectomy in tubal ectopic pregnancy: what do women prefer? *Reproductive biomedicine online*. 2010;21(5):687-93.
27. Notten KJ, Essers BA, Weemhoff M, Rutten AG, Donners JJ, van Gestel I, et al. Do patients prefer mesh or anterior colporrhaphy for primary correction of anterior vaginal wall prolapse: a labelled discrete choice experiment. *BJOG : an international journal of obstetrics and gynaecology*. 2015;122(6):873-80.
28. Kunutsor SK, Beswick AD, Whitehouse MR, Wylde V, Blom AW. Debridement, antibiotics and implant retention for periprosthetic joint infections: A systematic review and meta-analysis of treatment outcomes. *The Journal of infection*. 2018;77(6):479-88.

1
2
3
4
5
6 Legend: Figure 1. Example profile of DCE
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

| OPTION H1 | OR | OPTION H2 |
|--|----|--|
| I take antibiotics and the side effects don't affect me much <i>and</i> I have two operations <i>and</i> After my new hip is fitted, I can do most of the things that I want to do <i>and</i> After my surgical treatment starts, it takes 6 months to get back to the things that I normally do | OR | I take antibiotics and the side effects affect me a lot <i>and</i> I have one operation <i>and</i> After my new hip is fitted, I cannot do most of the things that I want to do <i>and</i> After my surgical treatment starts, it takes 12 months to get back to the things that I normally do |
| Please only tick ONE box: | | |
| OPTION H1 <input type="checkbox"/> | | OPTION H2 <input type="checkbox"/> |

Figure 1. Example profile of DCE
160x90mm (300 x 300 DPI)