

## Correction: *The premature closure of ROMPA clinical trial: mortality reduction in septic shock by plasma adsorption*

Giménez-Esparza C, Portillo-Requena C, Colomina-Climent F, *et al.* The premature closure of ROMPA clinical trial: mortality reduction in septic shock by plasma adsorption. *BMJ Open* 2019;9:e030139. doi: 10.1136/bmjopen-2019-030139.

This article was previously published with an error.

The following statement in the discussion section is incorrect: 'In addition, as in our study, the COMPACT 2 group used a randomisation system based on prognostic scores,<sup>6</sup> which means that the groups will not be similar until the end of recruitment (which is reason to introduce the propensity score in our results).' The COMPACT 2 group used a blocked randomization schedule (randomly permuting blocks of four and six), with stratification according to site and the presence of septic shock at admission.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

*BMJ Open* 2020;10:e030139corr1. doi:10.1136/bmjopen-2019-030139corr1

