

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Characteristics of older adults with back pain associated with choice of first primary care provider: The Back Complaints in the Elders – Norway (BACE-N) study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053229
Article Type:	Original research
Date Submitted by the Author:	08-May-2021
Complete List of Authors:	Vigdal, Ørjan; Oslo Metropolitan University, Department of Physiotherapy Storheim, Kjersti; Oslo University Hospital; Oslo Metropolitan University, Department of Physiotherapy Munk Killingmo, Rikke ; Oslo Metropolitan University Faculty of Health Sciences, Department of Physiotherapy Småstuen, Milada; Oslo Metropolitan University, Department of Nursing and Health Promotion Grotle, Margreth; Oslo Metropolitan University, Department of Physiotherapy; Oslo universitetssykehus Ullevål, Research and Communication Unit for Musculoskeletal Health
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, PRIMARY CARE, GERIATRIC MEDICINE

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Article title:**

Characteristics of older adults with back pain associated with choice of first primary care provider:  
The Back Complaints in the Elders – Norway (BACE-N) study

**Authors:**

Ørjan Nesse Vigdal<sup>1</sup>

Kjersti Storheim<sup>2,1</sup>

Rikke Munk Killingmo<sup>1</sup>

Milada C. Småstuen<sup>1</sup>

Margreth Grotle<sup>1,2</sup>

**Author affiliations:**

<sup>1</sup>Department of Physiotherapy, Faculty of Health Science, OsloMet - Oslo Metropolitan University, Oslo, Norway

<sup>2</sup>Research and Communication Unit for Musculoskeletal Health (FORMI), Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway

**Corresponding author:**

Ørjan Nesse Vigdal

E-mail address: orvig@oslomet.no

Mailing address: OsloMet – Oslo Metropolitan University, Post box 4, St.Olavs plass, 0130 Oslo, Norway

Telephone number: +47 414 59 314

**Manuscript number of pages (excluding title page, including tables): 20**

**Word count: 3975**

**Total number of tables: 2**

**Supplementary files: 1**

## Abstract

**Background:** Studies of older adults with back pain in primary care have found significant national differences in patient characteristics. There is a lack of knowledge of whether characteristics of older back pain patients differ according to their choice of first primary care provider.

**Objectives:** To describe characteristics of older adults with back pain in primary care, and to assess associations between patient characteristics and type of first primary care provider (general practitioner (GP), physiotherapist (PT) or chiropractor).

**Methods:** This cross-sectional study included patients aged  $\geq 55$  years seeking Norwegian primary care with a new episode of back pain. Patient characteristics were collected through questionnaires and a clinical examination, covering the following domains: sociodemographic, general health, current and previous back pain, psychological and clinical factors. Associations between patient characteristics and visiting a GP or PT compared to a chiropractor were assessed with multiple multinomial regression analyses.

**Results:** We included 452 patients: 127 first visited a GP, 130 first visited a PT and 195 first visited a chiropractor. Median (IQR) age was 66 (59-72) years. Median (IQR) back-related disability (Roland-Morris Disability Questionnaire, 0-24) was 9 (5-13). Recurring episodes were common, 301 (67%) patients had monthly or yearly recurrences. Patients with worse back-related disability, longer duration of symptoms, lower expectations for full recovery and worse physical performance measured with the Back Performance scale had higher odds of visiting a GP or PT compared to a chiropractor.

**Conclusion:** Older back pain patients in primary care had moderate to severe levels of back-related disability, and most had recurring episodes. Our results suggest that older adult's choice of first primary care provider was associated with important patient characteristics, which highlights the need for caution with generalizations of study results across primary care populations.

**Trial registration number:** ClinicalTrials identifier: NCT04261309

## Data availability statement

Data not available.

**Keywords:** Back pain, older adults, primary care, characteristics, care-seeking behaviour

## Article summary

### Strengths and limitations of this study

- This is the first study to compare characteristics of older adults with back pain visiting a GP, physiotherapist or chiropractor.
- This study provides a thorough comprehensive overview of older adults with back pain, and thus contributes with important knowledge in a research field with few previous studies
- It was not possible to obtain data on eligible patients that were not invited or declined to participate in the study. This might reduce external validity.

For peer review only

## Introduction

Back pain is the number one cause of years lived with disability globally, with an estimated point prevalence of 11.9% [1, 2]. Older adults have historically been under-represented in back pain research [3, 4], but have recently received increased attention [5, 6]. Although the prevalence of pathoanatomical findings on diagnostic imaging increases with age [7-9], the prevalence of serious pathology, such as vertebral fractures and neuropathic pain, in older back pain patients in primary care is low. Studies have reported a prevalence of 6% and 2-11%, respectively [10, 11]. Moreover, studies in primary care have found significant national differences in the characteristics and burden of back pain in older adults [12, 13]. This highlights the importance of caution when generalizing results from studies from one setting to another.

Most patients seeking healthcare for back pain are treated in primary care [14]. In Norway, back pain is the reason for 10%, 27% and 86% of the visits to general practitioners (GP), physiotherapists (PT) and chiropractors, respectively [15]. One study suggests that choice of first primary care provider has consequences for future healthcare consumption [16]. To optimize decision making regarding treatment, research and health policies, detailed knowledge of patient populations is required. Most of the previous studies exploring patient populations seeking primary care have compared GP and chiropractic populations, showing that patients seeking care from a GP have a higher overall burden of back pain compared to chiropractic patients [17-24]. Only a few studies include PT populations [25-28]. These studies suggest that patients seeking care from PTs are older and have more disability than those seeking care from chiropractors [25, 26, 28]. To the best of our knowledge, only one study has been performed in an exclusively older population [27]. This study found that older women seeking care from GPs reported worse back pain and worse health-related quality of life than older women visiting a PT or a chiropractor [27]. The study only included women between 59-64 years of age, and it is not clear if the results are also generalizable to men or adults over 65 years of age. Further, they did not examine back-related disability or other back pain factors, sociodemographic factors, psychological factors or clinical factors. Thus, there is still a considerable lack of knowledge regarding whether characteristics of older back pain patients differ according to their choice of first primary care provider.

Therefore, the aims of this study were 1) to describe the characteristics of patients  $\geq 55$  years of age seeking primary care for a new episode of back pain in terms of sociodemographic, general health, current back pain and back pain history, psychological and clinical characteristics, and 2) to assess if patient characteristics are associated with type of first primary care provider (GP, PT or chiropractor).

## Methods

### Design and setting

This cross-sectional study presents baseline data from the Back Complaints in the Elders – Norway (BACE-N) study, a prospective observational cohort study in Norwegian primary care. The BACE-N study is a part of the international BACE consortium, with research groups from Brazil, the Netherlands and Australia [6]. The BACE-N study protocol has been registered in ClinicalTrials.gov (Identifier NCT04261309). The study was classified as a quality assessment study by the Norwegian Regional Committee for Medical Research Ethics (reference no. 2014/1634/REK vest) and was approved by the Norwegian Social Science Data Service in 2015 (reference no. 42149).

### Participants and recruitment procedure

Eligible patients were  $\geq 55$  years of age, seeking primary care from a GP, PT or chiropractor in primary care for a new episode of back pain. Back pain was defined as pain located in the region from the top of the scapula to the sacrum, with or without radiating leg pain. A new episode was defined as not having received healthcare for the same complaint in the last six months. Patients were excluded if they had difficulties completing the questionnaire due to language barriers, or if they had difficulties completing the clinical examination (for example wheelchair-bound patients). Participants received care as usual.

Patients were recruited from GPs, PTs, and chiropractors in urban and rural parts of Norway between April 2015 and February 2020, either during or immediately after the consultation. The primary care providers were instructed to invite consecutive patients. To facilitate the recruitment process, media advertisements were also used. Eligible patients received oral and written information about the study. The final screening for eligibility and inclusion to the study was performed by the researchers. All included patients signed an informed consent form before enrolment in the study. The baseline measurements, consisting of questionnaires and a clinical examination, were collected as soon after the first primary care consultation as possible.

### Measurements

#### *Sociodemographic variables*

Information regarding age, sex, marital status, employment status and educational level were collected.

#### *General health variables*



1  
2  
3 Health-related quality of life (HR-QoL) was measured using the Short-Form Health Survey 36-item  
4 (SF-36) physical and mental summary measures (range 0-100, higher score indicates better HR-QoL)  
5 [29]. Alcohol consumption was measured using the 3-item Alcohol Use Disorder Identification Test  
6 consumption questions (AUDIT-C) (range 0-12, higher score indicates higher alcohol consumption)  
7 [30]. Hazardous alcohol consumption was defined as an AUDIT-C score of  $\geq 3/12$  for women and  
8  $\geq 4/12$  for men [31, 32]. Smoking status (current smoker, previous smoker, non-smoker) was  
9 collected. The number of comorbidities was measured using the Self-Administered Comorbidity  
10 Questionnaire (SCQ) [33]. The SCQ has 13 pre-defined comorbidities and two optional comorbidities.  
11 Item 12, "back pain", was replaced with a third optional comorbidity. Widespread pain was  
12 measured using the pain drawing from McGill Pain Questionnaire and the revised criteria from Wolfe  
13 et al. for widespread pain [34, 35]. The number of falls during the last six weeks was collected, and  
14 falls self-efficacy was measured using the Falls Efficacy Scale-International (FES-I) (range 16-64,  
15 higher score indicated lower falls efficacy) [36].

#### 26 *Current back pain and back pain history*

27 Back pain location (thoracic or lumbar, or both) was collected. Average back pain severity last week  
28 was measured using the Numeric Rating Scale (NRS) (range 0-10, higher score indicates higher back  
29 pain severity) [37]. Back-related disability was measured with the 24-item Roland-Morris Disability  
30 Questionnaire (RMDQ) (range 0-24, higher score indicated more back-related disability) [38]. Back  
31 pain duration was measured in days and categorized into "<6 weeks", "6 weeks to 3 months", and  
32 ">3 months". Frequency of previous back pain episodes (monthly, yearly, every 1-5 years, every five  
33 years, once) was collected. Sleep problems attributable to back pain were measured using item 5i  
34 from the Pittsburgh Sleep Quality Index (PSQI) [39], and dichotomized to "weekly/less than weekly".  
35 Morning stiffness was measured with item six from Knee injury and Osteoarthritis Outcome Score  
36 (KOOS) [40], where we replaced the word "knee" with "back".

#### 46 *Psychological variables*

47 Kinesiophobia was measured using the Fear-Avoidance Beliefs Questionnaire-Physical Activity  
48 subscale (FABQ-PA) (range 0-24, higher score indicates higher levels of kinesiophobia) [41]. Signs of  
49 depression were measured with the Center for Epidemiological Studies-Depression questionnaire  
50 (CES-D) (range 0-60, higher score indicates more signs of depression) [42]. Pain catastrophizing was  
51 measured using the Pain Catastrophizing Scale (PCS) (range 0-52, higher score indicates more pain  
52 catastrophizing) [43]. Beliefs and attitudes towards back pain was measured using the Back Beliefs  
53 Questionnaire (BBQ) (range 9-45, higher score indicates more positive beliefs) [44]. Start Back  
54 Screening Tool was used to assess prognostic risk profiles [45]. Expectations of recovery from back  
55  
56  
57  
58  
59  
60

1  
2  
3 pain within the next 3 months was assessed with a five-point scale, with the categories “Fully  
4 recovered”, “Much better”, “No difference”, “Much worse”, and “Worse than ever”.  
5  
6  
7

### 8 *Clinical variables*

9  
10 Pain with active movements was assessed for forward flexion, lateral flexion and rotation of the  
11 back. Physical performance with focus on trunk mobility was assessed with the 6-item Back  
12 Performance Scale (BPS) (range 0-18, higher score indicates worse trunk mobility performance) [46].  
13 Walking function was assessed with the Timed-Up-and-Go (TUG) [47]. Signs of radiculopathy was  
14 measured using a clinical diagnostic model that summarizes five items: Subjective sensory changes (1  
15 point), radiating pain below the knee (2 points), leg pain worse than back pain (2 points), positive  
16 neural tension test (3 points) and neurological deficit of myotome, dermatome or reflexes in the  
17 lower limb (2 points) [48]. A score of  $\geq 5/10$  has been shown to indicate >80% probability of  
18 radiculopathy [48]. Twelve red flags were assessed: Cancer, first episode of back pain, constant pain,  
19 unexplained weight loss, systemically unwell, fever, urinary retention or loss of bladder control, age  
20  $\geq 75$  years, trauma cause of back pain, osteoporosis, cortisone use and severe morning stiffness.  
21  
22  
23  
24  
25  
26  
27  
28  
29

### 30 **Statistical analyses**

31 All analyses were performed using the IBM SPSS Statistics version 26 for Windows (IBM Corporation,  
32 Armonk, NY, USA). To handle missing data, five multiple imputation datasets with 10 iterations were  
33 created using regression estimation, and the pooled estimates are presented in this study. Patient  
34 characteristics were described with counts and percentages for categorical variables, mean and  
35 standard deviation (SD) for normally distributed continuous variables and median and interquartile  
36 range (IQR) for continuous variables with a skewed distribution. Mann-Whitney U-test was used to  
37 assess differences in days between first primary care contact and inclusion to the study between  
38 primary care practitioners, and between those recruited from primary care and those recruited from  
39 media advertisements. Multinomial regression was used to assess the strength of the associations  
40 between patient characteristics and patient’s choice of first primary care provider. First primary care  
41 provider (GP, PT or chiropractor) was the dependent variable. The chiropractic group was the largest,  
42 and therefore chosen as the reference group. Patient characteristics were organized into five blocks,  
43 for which we created separate models: i) Sociodemographic ii) general health iii) current back pain  
44 episode and back pain history iv) psychological variables and v) clinical variables. All variables in the  
45 block were simultaneously included in the model, without univariate pre-testing. The strength of  
46 associations is expressed as odds ratios (OR) with 95% confidence intervals (CI). We considered our  
47 study as exploratory, so no correction for multiple testing was performed [49]. P-values <0.05 were  
48 thus considered statistically significant. All tests were two-sided.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### *Assessment of generalizability*

Because of economic and practical reasons, we were unable to collect data on eligible participants that declined to participate or for other reasons were not invited. Therefore, we performed a descriptive comparison of the BACE-N on age, sex, nationality, educational level, work status, marital status, BMI, alcohol use, HR-QoL, depression and walking distance with individual data from a subsample from the study “The Norwegian study on life course, ageing and generation (NORLAG)” [50, 51]. This study used a random sampling strategy in the general population and included 11028 participants. The subsample (NORLAG MSK) consisted of 794 participants collected in 2017. The participants of the subsample were  $\geq 55$  years of age and had at least one musculoskeletal complaint.

### *Sensitivity analyses*

We performed two sensitivity analyses: 1) To assess possible bias introduced by the multiple imputation procedure, the multiple multinomial regression analyses were performed on complete case data. 2) Because PT services became available through direct access in Norway from 01.01.2018, characteristics of PT patients recruited before and after 01.01.2018 were compared using individual sample t-tests or Mann-Whitney U-tests for continuous variables, and chi-square tests for categorical variables. Results from the sensitivity analyses are available in supplementary material S1 and S2.

### *Sample size consideration*

Sample size was considered for the BACE-N study as a whole, with the following criteria: Having sufficient statistical power for up to 14 variables in a multivariate logistic regression analysis using the “10 events per variable” rule [52], with an outcome prevalence of 40%, and allowing for a dropout-rate of 20%. This yielded a preferred sample size of 450 participants. As the multinomial regression models in this study includes a maximum of 8 independent variables, we expect the sample size to be sufficient.

### **Patient and public involvement**

Patient representatives were part of the scientific board of the study and involved in designing and establishing BACE-N. Results will be disseminated to the recruiting primary care providers and the participating patients in an annual newsletter.

### **Results**

1  
2  
3 A total of 452 patients were included in the study, 127 first visited a GP, 130 first visited a PT and 195  
4 first visited a chiropractor. Eighteen patients were included from media advertisements. Median  
5 (IQR) number of days from first primary care contact to inclusion in the study was 7 (2-17) days.  
6  
7 Duration from first primary care contact to inclusion in the study was significantly shorter for  
8 chiropractic patients compared to GP patients ( $p<0.01$ ) and PT patients ( $p<0.01$ ). There was no  
9 difference in duration from first primary care contact to inclusion between those recruited directly  
10 from primary care practices, and those recruited through media advertisements.  
11  
12  
13  
14  
15

### 16 *Patient characteristics*

17 Missingness ranged from 0.0 to 16.8% for the variables, and total missingness was 4.4% across all  
18 values. Consult table 1 for details regarding patient characteristics. The median age of the patients  
19 was 66, around half of the patients were women, were in paid work, and had university-level  
20 education. Half of the patients had a hazardous alcohol consumption level, and nearly 60% of them  
21 were either current or previous smokers. One in six patients had experienced a fall during the last six  
22 weeks. Half of the patients had one or more comorbidities.  
23  
24  
25  
26  
27  
28  
29

30 Most patients reported moderate levels of back pain and moderate to severe levels of back-related  
31 disability with a median (IQR) RMDQ-score of 9 (5-13). Almost 60% of the patients experienced  
32 monthly or yearly recurrences of back pain. Over 40% experienced weekly sleep problems  
33 attributable to back pain, and 70% experienced moderate to extreme morning stiffness. Two thirds  
34 of the patients had a low-risk profile according to the SBT, and only 6.6% had a high-risk profile.  
35 Expectations of recovery were generally high, with three out of four expecting to be much better or  
36 fully recovered within three months.  
37  
38  
39  
40  
41  
42

### 43 *Associations between patient characteristics and type of first primary care provider*

44 Table 2 presents the associations from the multinomial regression analyses. Patients with higher  
45 back-related disability, longer duration of symptoms, worse physical performance, probable  
46 radiculopathy, poorer HR-QoL and lower expectations of being fully recovered within the next three  
47 months were more likely to visit a GP compared to a chiropractor. Patients with widespread pain  
48 were more likely to visit a chiropractor than a GP. The characteristics strongest associated with  
49 choosing a GP versus a chiropractor were duration of symptoms, widespread pain and expectation of  
50 being fully recovered.  
51  
52  
53  
54  
55  
56  
57

58 Patients that were older, had a longer duration of symptoms, higher back-related disability,  
59 moderate morning stiffness, higher levels of pain catastrophizing, physical performance, lower  
60

1  
2  
3 expectations of being fully recovered within the next three months were more likely to visit a PT  
4 compared to a chiropractor. Patients in the SBT medium risk group were more likely to visit a  
5 chiropractor compared to a PT. The characteristics strongest associated with choosing a PT versus a  
6 chiropractor were duration of symptoms and expectation of being fully recovered.  
7  
8  
9

10  
11 Gender, education level, marital status, employment status, comorbidities, back pain severity, sleep  
12 problems, kinesiophobia, depressive signs, back beliefs, red flags, pain on active range of motion and  
13 Timed Up and Go-scores were not associated with type of primary care provider.  
14  
15  
16

### 17 18 *Assessment of generalizability*

19  
20 The BACE-N study sample had more men (48% versus 36.3% in NORLAG MSK), more participants with  
21 high educational level (44% versus 28.6% in NORLAG MSK), more participants currently in paid work  
22 (45.3% 31.6% in NORLAG MSK), and more participants living with a partner (76.8% versus 62.2% in  
23 NORLAG MSK). Age, nationality, alcohol consumption, BMI, depressive signs, HR-QoL and walking  
24 distance were similar between BACE-N and NORLAG MSK.  
25  
26  
27  
28  
29  
30

### 31 **Discussion**

32  
33 This study showed that nearly all older patients with back pain had experienced back pain previously,  
34 and for most patients this episode was the latest of a series of annually or monthly recurring  
35 episodes. This is in accordance with several studies on back pain trajectories, where episodic or  
36 fluctuating pain was shown to be common both in the short and long term [53-56]. Further, patients  
37 with more severe back-related disability and other symptoms and signs were overall more likely to  
38 visit a GP or a physiotherapist than a chiropractor. Contrary to this finding, patients with widespread  
39 pain were more likely to choose a chiropractor over a GP. This is the first study to assess associations  
40 of a broad range of patient characteristics and choice of first primary care provider in an older  
41 population.  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 The burden of back pain and psychological profile were comparable between younger Norwegian  
52 back pain cohorts and the older BACE-N sample [57, 58]. The characteristics of the included patients  
53 in this study was largely comparable to the BACE-study from the Netherlands [12, 59], with a few  
54 exceptions. Both in our total study sample and our GP subsample, a larger proportion of patients had  
55 paid work, fewer experienced their first episode of back pain, and they reported lower levels of  
56 kinesiophobia and pain catastrophizing compared to the Dutch study sample. When comparing our  
57 results to the Brazilian BACE-study [12, 60], the Brazilian study had a higher proportion of women.  
58  
59  
60

1  
2  
3 Further, our study sample had more patients in paid work, more patients with hazardous alcohol  
4 consumption patterns, more smokers, the patients had fewer comorbidities, lower levels of back-  
5 related disability and back pain severity, kinesiophobia, depression signs, and pain catastrophizing  
6 compared to the Brazilian BACE-sample. These differences between populations within the BACE  
7 consortium might be explained in part by minor differences in recruitment strategies in the different  
8 countries [12] or differences in how primary care is organized in the different countries. In the  
9 Netherlands, patients were recruited exclusively from a GP setting [59], whereas in Brazil patients  
10 were recruited from primary care centres or health centres specialized in geriatrics [60]. Another  
11 possible explanation may be cultural differences in the expression and interpretation of and coping  
12 with pain [61].  
13  
14  
15  
16  
17  
18  
19  
20

21  
22 In line with previous research on healthcare utilization for back pain in younger populations [18, 20-  
23 22, 24-27], our results suggest that patients with “less complex” characteristics were more likely to  
24 visit a chiropractor compared to a GP or a physiotherapist. Unsurprisingly, studies using bivariate  
25 analyses [17, 19, 22, 24, 27, 28] to compare the provider groups find more significant associations or  
26 differences than studies using multivariate analyses [18, 20, 21, 25, 26]. However, regardless of  
27 statistical approach, these studies suggest that patients who seek chiropractic care have an overall  
28 lower burden of back pain compared to patients seeking GP or PT care [17-22, 24]. One notable  
29 exception is the study of Eklund et al. [23], which found that Swedish chiropractic patients had more  
30 pain and worse psychological and behavioural characteristics compared to a sample of sick-listed  
31 primary care (specific provider unknown) patients at high risk for chronicity. Our finding showing that  
32 patients with widespread pain were more likely to choose a chiropractor over a GP was contrary to  
33 the general pattern of chiropractic patients being less “complex.” To the best of our knowledge, no  
34 previous studies have compared prevalence of widespread pain in the two populations, but one  
35 study showed that GP patients had more musculoskeletal comorbidities [24], possibly implying more  
36 widespread pain. Two previous studies found an association between higher age and odds of seeking  
37 care from a physiotherapist compared to a chiropractor [25, 26], in line with our results.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Many of the patient characteristics associated with choice of primary care provider in this study have  
51 previously been found to be significant prognostic factors for the persistent back-related disability  
52 and back pain in older people. For example, duration of back pain and expectation of improvement  
53 [62-67], and higher levels of back-related disability [63-68], are consistently reported as significant  
54 prognostic factors for a poor outcome of a back pain episode. A few studies in older people have  
55 found that single symptoms of neurological involvement such as leg pain below the knee, and the  
56 diagnosis of spinal stenosis were prognostic factors for the outcome of a back pain episode [62, 65].  
57  
58  
59  
60

1  
2  
3 We combined single symptoms of neurological involvement into a compound measure, but it is likely  
4 that older patients with radiculopathy have worse outcomes than those without radiculopathy.  
5  
6 Although slightly different from widespread pain, the presence of multi-site pain has also in some  
7  
8 studies been found to be a prognostic factor for the outcome of back pain in older adults [65, 69].  
9  
10 The impact of pain catastrophizing on the clinical course of back pain is less clear in older adults [64,  
11  
12 67] compared to younger populations [70], but it is not unreasonable to believe that pain  
13  
14 catastrophizing may be a prognostic factor for back pain in older adults. Thus, the associations  
15  
16 between potential prognostic factors and choice of first primary care provider imply that we can  
17  
18 expect the clinical course of patients in the three primary care groups to be different.

19  
20 The results of this study need to be viewed with consideration of some limitations. We instructed the  
21  
22 recruiting primary care providers to invite consecutive patients, but because of obvious time  
23  
24 constraints in clinical practice we could not ask them to keep record of how many declined to  
25  
26 participate, nor of eligible patients that were not invited. This recruitment strategy increases the risk  
27  
28 of selection bias, and thus could reduce the external validity of the study. To compensate for this  
29  
30 limitation, we compared the BACE-N sample with the NORLAG MSK subsample. The characteristics of  
31  
32 the two samples were largely comparable, but BACE-N has more men, more participants with higher  
33  
34 education, more in paid work, and more living with their partner. Sex and education level have  
35  
36 previously been shown to be associated with back pain severity and back-related disability in older  
37  
38 adults [12, 13]. Thus, it may be possible that the levels of back pain and back-related disability  
39  
40 presented in this study are slightly underestimated. The NORLAG MSK subsample is sampled from  
41  
42 the general population, which may not be representative of those who seek care. However, the most  
43  
44 important determinants of care-seeking for back pain seems to be pain severity and disability levels  
45  
46 [71]. We therefore believe the assessment to be justified.

47  
48 Another limitation may be the analysis strategy. We chose to keep the variables in the five blocks to  
49  
50 provide a broad assessment of the differences in case-mix in the three primary care settings. To limit  
51  
52 the number of statistical tests performed, univariate pre-testing and testing a “final model” across  
53  
54 blocks were avoided. Furthermore, a different organization of the variables, for example strictly  
55  
56 adhering to the biopsychosocial model [72] or Andersen’s behavioural model of health services use  
57  
58 [73], may have yielded slightly different results. However, our results are largely supported by  
59  
60 previous studies, so the potential differences because of analysis strategy or variable organization  
may be negligible. A third limitation is that we were unable to examine some possibly important  
determinants for healthcare use, such as access to different providers, patient’s familiarity with  
providers, the patient’s economic situation and social network referrals [73-75]. Including these

1  
2  
3 factors would have given an even broader overview of associations between individual and  
4 contextual characteristics and choice of primary care provider and suggest that future research focus  
5 on examining the contextual and social factors associated with healthcare service use.  
6  
7  
8  
9

## 10 **Conclusion**

11 We found that nearly all older adults with back pain seeking primary care had experienced back pain  
12 previously, and recurring episodes were common. In general, patients with more severe back-related  
13 disability and other clinical symptoms and signs were more likely to visit a GP or a physiotherapist  
14 than a chiropractor. Our results suggest that important patient characteristics are associated with  
15 older adult's choice of primary care providers due to back pain, which may affect the clinical course  
16 of back pain for these patients. The findings highlight the need for caution with generalization of  
17 study results across primary care populations. This is an important consideration for healthcare  
18 providers, for the development and implementation of clinical practice guidelines, and for regulators  
19 when developing primary care pathways for back pain. Further research is needed in assessing if the  
20 choice of primary care provider affects future care pathways and the clinical course of back pain in  
21 older adults.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

## 33 **Acknowledgments**

34 The authors thank all study participants for their significant contribution. Further, we would like to  
35 thank all the recruiting healthcare providers: Kerstin Ulrich, Lise Lothe, Tim Raven, Andreas Hoff  
36 Nordvik, Christoffer Børsheim, Steinar Forshei, Mette Brekke, Daniel Major, Jan Harald Lønn, Agnes  
37 Mordt, Andrea Kolstad, Mathias Svanevik, Alexander Diesen, Lars Gullestad, Ida Svalstuen, Joakim  
38 Ordahl, Mona Øversveen, Jørgen Kongtorp, Are Hansen, Geir Haram, Palwinder Singh, Svein Erik  
39 Sandelien, Bent Ulseth, Harald Nordheim, Ola Sand, Ragnhild Perstølen, Jorun Salater, Anna Allen-  
40 Unhammer, Morten Nilsen, Haakon Lilleeng Asmyhr, Philip Wilkens, Ane Klevberg, Eli Magnesen,  
41 Aleksander Killingmo, Bjørn Tore Bjørkedal, Stina Lund, Daniel Ekeberg, Berte Marie Enger, Johan  
42 Edvard Tellum, Morten de la Cruz, Bård Kvam, Marte Paulsen, Astrid Figger, Christian Mayer, Trond  
43 Magne Aasberg, Thea Tømmervåg, Jørgen Øyen, Håvard Nordås, Tore Viste Ollestad, Olav Aase,  
44 Renate Meier, Bjørn Røe, Jørgen Øyen, Jack Johnson, Carl-Erik Høgquist, Ingrid Hystad, Mats  
45 Thorbeck, Elisabeth Barø, Kaja McCormick, Lars Martin Fischer, Martin Haagensen, Cathrine  
46 Rosslund, Marianne Storberget, Nora Helk, Grete Bråten, Annecken Lister Haugen, Hege Herstad,  
47 Cathrine Natland, Frøydis Blaker Åsbø, Ola Klaastad, Thorleif Henning Monsen, Geir Wiik, Jørn  
48 Christian Halvorsen, Tonje Høgdahl Mysen, Tine Tandberg, Eir Marie Bergan, Rune Solheim, Ole  
49 Kristoffer Larsen. We thank the research assistants for assisting with clinical examinations and  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 providing baseline questionnaires to study participants. We thank the BACE-N scientific board, and  
4 the BACE Consortium for important input with designing the study.  
5  
6  
7

8 **Author contributions:**

9  
10 ØNV: Study design, data collection, data analyses, manuscript draft. KS: Study design, data  
11 interpretation, critical revision. RMK: Data collection, data interpretation, critical revision. MCS:  
12 Statistical advisor, data interpretation, critical revision. MG: Principal investigator, study design, data  
13 interpretation, critical revision.  
14  
15  
16

17  
18 **Funding:**

19  
20 This work was funded by The Norwegian Fund for Post-Graduate Training in Physiotherapy (grant  
21 number 90749). The funding body was not involved in designing the study, data collection, analyses  
22 or interpretation of data, nor in writing the manuscript.  
23  
24  
25

26  
27 **Competing interests**

28 The authors declare no competing interests.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

1. Vos, T., et al., *Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010*. The Lancet, 2012. **380**(9859): p. 2163-2196.
2. Hoy, D., et al., *A systematic review of the global prevalence of low back pain*. Arthritis & Rheumatism, 2012. **64**(6): p. 2028-2037.
3. Bressler, H.B., et al., *The Prevalence of Low Back Pain in the Elderly: A Systematic Review of the Literature*. Spine, 1999. **24**(17): p. 1813-1819.
4. Paeck, T., et al., *Are older adults missing from low back pain clinical trials? A systematic review and meta-analysis*. Arthritis Care Res (Hoboken), 2014. **66**(8): p. 1220-6.
5. Jarvik, J.G., et al., *Study protocol: the back pain outcomes using longitudinal data (BOLD) registry*. BMC musculoskeletal disorders, 2012. **13**(1): p. 64.
6. Scheele, J., et al., *Back complaints in the elders (BACE); design of cohort studies in primary care: an international consortium*. BMC Musculoskelet Disord, 2011. **12**: p. 193.
7. Kalichman, L., et al., *Spinal stenosis prevalence and association with symptoms: the Framingham Study*. The Spine Journal, 2009. **9**(7): p. 545-550.
8. Van Der Klift, M., et al., *The incidence of vertebral fractures in men and women: the Rotterdam Study*. Journal of bone and mineral research, 2002. **17**(6): p. 1051-1056.
9. de Schepper, E.I., et al., *The association between lumbar disc degeneration and low back pain: the influence of age, gender, and individual radiographic features*. Spine, 2010. **35**(5): p. 531-536.
10. Enthoven, W.T., et al., *Prevalence and "Red Flags" Regarding Specified Causes of Back Pain in Older Adults Presenting in General Practice*. Physical Therapy, 2016. **96**(3): p. 305-312.
11. Enthoven, W.T., et al., *Back Complaints in Older Adults: Prevalence of Neuropathic Pain and Its Characteristics*. Pain Medicine, 2013. **14**(11): p. 1664-1672.
12. Jesus-Moraleida, F.R., et al., *Back Complaints in the Elders in Brazil and the Netherlands: a cross-sectional comparison*. Age and ageing, 2017. **46**(3): p. 476-481.
13. Jarvik, J.G., et al., *Back pain in seniors: the back pain outcomes using longitudinal data (BOLD) cohort baseline data*. BMC Musculoskeletal Disorders, 2014. **15**(1): p. 1-23.
14. Kinge, J.M., et al., *Musculoskeletal disorders in Norway: prevalence of chronicity and use of primary and specialist health care services*. BMC Musculoskeletal Disorders, 2015. **16**(1): p. 1-9.
15. Werner, E.L. and A. Indahl, *Kunnskap, praksis og holdninger til rygglidelser hos leger, fysioterapeuter og kiropraktorer*. Tidsskr Nor Laegeforen, 2005. **125**(13): p. 1794–7.
16. Fritz, J.M., J. Kim, and J. Dorius, *Importance of the type of provider seen to begin health care for a new episode low back pain: associations with future utilization and costs*. Journal of evaluation in clinical practice, 2016. **22**(2): p. 247-252.
17. Carey, T.S., et al., *Acute Severe Low Back Pain: A Population-based Study of Prevalence and Care-seeking*. Spine, 1996. **21**(3): p. 339-344.
18. Hurwitz, E.L. and H. Morgenstern, *The effects of comorbidity and other factors on medical versus chiropractic care for back problems*. Spine, 1997. **22**(19): p. 2254-2263.
19. Nyiendo, J., et al., *Patient characteristics and physicians' practice activities for patients with chronic low back pain: A practice-based study of primary care and chiropractic physicians*. 2001. **24**(2): p. 92-100.
20. Côté, P., J.D. Cassidy, and L. Carroll, *The treatment of neck and low back pain: who seeks care? who goes where?* Medical care, 2001: p. 956-967.
21. Sharma, R., M. Haas, and M. Stano, *Patient Attitudes, Insurance, and Other Determinants of Self-Referral to Medical and Chiropractic Physicians*. American Journal of Public Health, 2003. **93**(12): p. 2111-2117.

- 1  
2  
3 22. Hestbaek, L., et al., *Low back pain in primary care: a description of 1250 patients with low*  
4 *back pain in danish general and chiropractic practice.* International journal of family  
5 medicine, 2014. **2014**.
- 6 23. Eklund, A., et al., *Psychological and behavioral differences between low back pain*  
7 *populations: a comparative analysis of chiropractic, primary and secondary care patients.*  
8 BMC Musculoskeletal Disorders, 2015. **16**(1).
- 9 24. Hartvigsen, L., et al., *Baseline Characteristics May Help Indicate the Best Choice of Health*  
10 *Care Provider for Back Pain Patients in Primary Care: Results From a Prospective Cohort*  
11 *Study.* Journal of Manipulative and Physiological Therapeutics, 2020. **43**(1): p. 13-23.
- 12 25. Chevan, J. and D.L. Riddle, *Factors associated with care seeking from physicians, physical*  
13 *therapists, or chiropractors by persons with spinal pain: a population-based study.* journal of  
14 orthopaedic & sports physical therapy, 2011. **41**(7): p. 467-476.
- 15 26. Blanchette, M.-A., et al., *Workers' characteristics associated with the type of healthcare*  
16 *provider first seen for occupational back pain.* BMC Musculoskeletal Disorders, 2016. **17**(1).
- 17 27. Sibbritt, D., et al., *Severity of back pain may influence choice and order of practitioner*  
18 *consultations across conventional, allied and complementary health care: a cross-sectional*  
19 *study of 1851 mid-age Australian women.* BMC Musculoskeletal Disorders, 2016. **17**(1).
- 20 28. Walker, B.F., R. Muller, and W.D. Grant, *Low back pain in Australian adults. Health provider*  
21 *utilization and care seeking.* Journal of manipulative and physiological therapeutics, 2004.  
22 **27**(5): p. 327-335.
- 23 29. Loge, J.H. and S. Kaasa, *Short Form 36 (SF-36) health survey: normative data from the general*  
24 *Norwegian population.* Scandinavian Journal of Social Medicine, 1998. **26**(4): p. 250-258.
- 25 30. Bush, K., *The AUDIT Alcohol Consumption Questions (AUDIT-C). An Effective Brief Screening*  
26 *Test for Problem Drinking.* Archives of Internal Medicine, 1998. **158**(16): p. 1789.
- 27 31. Bradley, K.A., et al., *AUDIT-C as a Brief Screen for Alcohol Misuse in Primary Care.* Alcoholism:  
28 Clinical and Experimental Research, 2007. **31**(7): p. 1208-1217.
- 29 32. Norwegian National Advisory Unit on Concurrent Substance Abuse and Mental Health  
30 Disorders. *AUDIT-C.* 23.09.2020]; Available from: [https://rop.no/kartleggingsverktøey/audit-](https://rop.no/kartleggingsverktøey/audit-c/)  
31 [c/](https://rop.no/kartleggingsverktøey/audit-c/).
- 32 33. Sangha, O., et al., *The self-administered comorbidity questionnaire: A new method to assess*  
33 *comorbidity for clinical and health services research.* Arthritis & Rheumatism, 2003. **49**(2): p.  
34 156-163.
- 35 34. Wolfe, F., et al., *Revised chronic widespread pain criteria: development from and integration*  
36 *with fibromyalgia criteria.* Scandinavian Journal of Pain, 2019. **20**(1): p. 77-86.
- 37 35. Melzack, R., *The McGill Pain Questionnaire: Major properties and scoring methods.* Pain,  
38 1975. **1**(3): p. 277-299.
- 39 36. Helbostad, J.L., et al., *Validation of the Falls Efficacy Scale-International in fall-prone older*  
40 *persons.* Age and Ageing, 2010. **39**(2): p. 259-259.
- 41 37. Von Korff, M., M.P. Jensen, and P. Karoly, *Assessing Global Pain Severity by Self-Report in*  
42 *Clinical and Health Services Research.* Spine, 2000. **25**(24): p. 3140-3151.
- 43 38. Grotle, M., J. Brox, and N. Vollestad, *Cross-cultural adaptation of the Norwegian versions of*  
44 *the Roland-Morris Disability Questionnaire and the Oswestry Disability Index.* Journal of  
45 Rehabilitation Medicine, 2003. **35**(5): p. 241-247.
- 46 39. Buysse, D.J., et al., *The Pittsburgh Sleep Quality Index: a new instrument for psychiatric*  
47 *practice and research.* Psychiatry research, 1989. **28**(2): p. 193-213.
- 48 40. Roos, E.M., et al., *Knee Injury and Osteoarthritis Outcome Score (KOOS)—Development of a*  
49 *Self-Administered Outcome Measure.* Journal of Orthopaedic & Sports Physical Therapy,  
50 1998. **28**(2): p. 88-96.
- 51 41. Grotle, M., J.I. Brox, and N.K. Vøllestad, *Reliability, validity and responsiveness of the fear-*  
52 *avoidance beliefs questionnaire: methodological aspects of the Norwegian version.* Journal of  
53 rehabilitation medicine, 2006. **38**(6): p. 346-353.
- 54  
55  
56  
57  
58  
59  
60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
42. Radloff, L.S., *The CES-D scale: A self report depression scale for research in the general population*. Applied Psychological Measurements, 1977. **1**(3): p. 385-401.
43. Fernandes, L., et al., *Cross-cultural adaptation and validation of the Norwegian pain catastrophizing scale in patients with low back pain*. BMC musculoskeletal disorders, 2012. **13**(1): p. 111.
44. Tingulstad, A., et al., *Back beliefs among elderly seeking health care due to back pain; psychometric properties of the Norwegian version of the back beliefs questionnaire*. BMC Musculoskeletal Disorders, 2019. **20**(1).
45. Hill, J.C., et al., *A primary care back pain screening tool: Identifying patient subgroups for initial treatment*. Arthritis & Rheumatism, 2008. **59**(5): p. 632-641.
46. Strand, L.I., R. Moe-Nilssen, and A.E. Ljunggren, *Back Performance Scale for the Assessment of Mobility-Related Activities in People With Back Pain*. Physical Therapy, 2002. **82**(12): p. 1213-1223.
47. Podsiadlo, D. and S. Richardson, *The Timed "Up & Go": A Test of Basic Functional Mobility for Frail Elderly Persons*. Journal of the American Geriatrics Society, 1991. **39**(2): p. 142-148.
48. Stynes, S., et al., *Clinical diagnostic model for sciatica developed in primary care patients with low back-related leg pain*. PloS one, 2018. **13**(4).
49. Bender, R. and S. Lange, *Adjusting for multiple testing—when and how?* Journal of Clinical Epidemiology, 2001. **54**(4): p. 343-349.
50. Slagsvold, B., et al., *Life-course, ageing and generations in Norway:the NorLAG study*. Norsk Epidemiologi, 2012. **22**(2).
51. Torsteinsen, A. and A. Holmøy, *Den norske studien av livsløp, aldring og generasjon—tredje runde (NorLAG3). Dokumentasjonsrapport*. 2019.
52. Moons, K.G.M., et al., *Prognosis and prognostic research: what, why, and how?* BMJ, 2009. **338**: p. b375.
53. Dunn, K.M., K. Jordan, and P.R. Croft, *Characterizing the course of low back pain: a latent class analysis*. American journal of epidemiology, 2006. **163**(8): p. 754-761.
54. Dunn, K.M., P. Campbell, and K.P. Jordan, *Long-term trajectories of back pain: cohort study with 7-year follow-up*. BMJ open, 2013. **3**(12): p. e003838.
55. Kongsted, A., et al., *Patients with low back pain had distinct clinical course patterns that were typically neither complete recovery nor constant pain. A latent class analysis of longitudinal data*. The Spine Journal: Official Journal Of The North American Spine Society, 2015. **15**(5): p. 885-894.
56. Kongsted, A., et al., *What have we learned from ten years of trajectory research in low back pain?* BMC Musculoskeletal Disorders, 2016. **17**: p. 1-11.
57. Nordstoga, A.L., et al., *Improvement in Work Ability, Psychological Distress and Pain Sites in Relation to Low Back Pain Prognosis: A Longitudinal Observational Study in Primary Care*. Spine, 2019. **44**(7): p. E423-E429.
58. Grotle, M., et al., *Prognostic factors in first-time care seekers due to acute low back pain*. European Journal of Pain, 2007. **11**(3): p. 290-298.
59. Scheele, J., et al., *Characteristics of older patients with back pain in general practice: BACE cohort study*. Eur J Pain, 2013. **18**(2): p. 279-87.
60. Jesus-Moraleida, F.R.D., et al., *The Brazilian Back Complaints in the Elders (Brazilian BACE) study: characteristics of Brazilian older adults with a new episode of low back pain*. Brazilian Journal of Physical Therapy, 2017.
61. Peacock, S. and S. Patel, *Cultural influences on pain*. Reviews in pain, 2008. **1**(2): p. 6-9.
62. Scheele, J., et al., *Course and prognosis of older back pain patients in general practice: a prospective cohort study*. Pain, 2013. **154**(6): p. 951-7.
63. Deyo, R.A., et al., *Trajectories of symptoms and function in older adults with low back disorders*. Spine (Phila Pa 1976), 2015. **40**(17): p. 1352-62.
64. Enthoven, W.T., et al., *Defining trajectories in older adults with back pain presenting in general practice*. Age and ageing, 2016. **45**(6): p. 878-883.

- 1  
2  
3 65. Rundell, S.D., et al., *Predictors of Persistent Disability and Back Pain in Older Adults with a*  
4 *New Episode of Care for Back Pain*. Pain Med, 2017. **18**(6): p. 1049-1062.
- 5 66. Jarvik, J.G., et al., *Long-term outcomes of a large, prospective observational cohort of older*  
6 *adults with back pain*. Spine J, 2018. **18**(9): p. 1540-1551.
- 7 67. van der Gaag, W.H., et al., *Developing clinical prediction models for non-recovery in older*  
8 *patients seeking care for back pain: the BACE prospective cohort study*. PAIN, 2020. **Articles**  
9 **in Press**.
- 10 68. Van Den Berg, R., et al., *Clinical and radiographic features of spinal osteoarthritis predict*  
11 *long-term persistence and severity of back pain in older adults*. Annals of Physical and  
12 Rehabilitation Medicine, 2020.
- 13 69. Rundell, S.D., et al., *Multisite Pain Is Associated with Long-term Patient-Reported Outcomes*  
14 *in Older Adults with Persistent Back Pain*. Pain Med, 2019.
- 15 70. Wertli, M.M., et al., *Catastrophizing—a prognostic factor for outcome in patients with low*  
16 *back pain: a systematic review*. The Spine Journal, 2014. **14**(11): p. 2639-2657.
- 17 71. Ferreira, M., et al., *Factors defining care-seeking in low back pain - A meta-analysis of*  
18 *population based surveys*. European Journal of Pain, 2010. **14**(7): p. 747.e1-747.e7.
- 19 72. Waddell, G., *1987 Volvo award in clinical sciences. A new clinical model for the treatment of*  
20 *low-back pain*. Spine, 1987. **12**(7): p. 632-644.
- 21 73. Andersen, R.M., *National health surveys and the behavioral model of health services use*.  
22 Medical care, 2008. **46**(7): p. 647-653.
- 23 74. Davis, M.A., et al., *Regional Supply of Chiropractic Care and Visits to Primary Care Physicians*  
24 *for Back and Neck Pain*. The Journal of the American Board of Family Medicine, 2015. **28**(4):  
25 p. 481-490.
- 26 75. Kirby, E.R., et al., *A qualitative study of influences on older women's practitioner choices for*  
27 *back pain care*. BMC Health Services Research, 2014. **14**(1): p. 131.
- 28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Tables:

Table 1: Baseline characteristics of study participants\*

	Missing, n (%)	Total (n=452)	GP (n=127)	PT (n=130)	Chiro (n=195)
<b>Sociodemographic variables</b>					
Age, median (IQR)	0 (0.0)	66 (59-72)	67 (60-73)	68 (63-74)	63 (58-71)
Female, n (%)	0 (0.0)	235 (52.0)	74 (58.3)	70 (53.8)	89 (46.1)
Marital status	19 (4.2)				
Married or living with partner, n (%)		347 (76.8)	90 (70.1)	98 (74.6)	158 (81.0)
Employment status	5 (1.1)				
Currently in paid work, n (%)		212 (45.3)	57 (43.3)	49 (31.5)	106 (55.9)
Educational level, n (%)	20 (4.4)				
Low (elementary + high school)		253 (56.0)	72 (56.7)	70 (55.1)	110 (56.4)
High (university level)		199 (44.0)	55 (43.3)	60 (44.9)	85 (43.6)
<b>General health variables</b>					
Health-related quality of life (SF-36 0-100)	41 (9.1)				
Mental sumscore, mean (SD)		52.5 (10.0)	50.5 (11.5)	53.4 (10.0)	53.2 (8.8)
Physical sumscore, mean (SD)		41.4 (8.4)	40.0 (7.9)	40.6 (8.0)	42.8 (8.9)
Hazardous alcohol consumption (AUDIT-C <sup>A</sup> ), n (%)	59 (13.1)	228 (50.4)	65 (51.1)	65 (50.0)	98 (50.2)
Smoking status, n (%)	22 (4.9)				
Current smoker		63 (13.9)	21 (16.5)	13 (10.0)	28 (14.3)
Previous		203 (44.9)	59 (46.4)	60 (46.2)	84 (43.1)
Never		186 (41.2)	47 (37.0)	57 (43.8)	83 (42.6)
Number of comorbidities (SCQ 0-15), median (IQR)	18 (4.0)	1 (1-2)	1 (0-2)	2 (1-2)	1 (1-5)
BMI, mean (SD)	14 (3.1)	27.6 (4.7)	27.6 (4.5)	27.5 (4.7)	27.7 (4.8)
Fall last 6 weeks, n (%)	24 (5.3)	73 (16.1)	13 (10.2)	24 (18.4)	35 (18.2)
Falls self-efficacy (FESI 16-64), mean (SD)	48 (10.6)	21.8 (6.0)	22.4 (6.3)	22.2 (6.1)	21.1 (5.7)
Widespread pain, n (%)	16 (3.5)	33 (7.3)	5 (4.0)	7 (5.3)	21 (10.8)
<b>Current back pain and back pain history variables</b>					
Previous back pain, n (%)	58 (12.8)				
Monthly		127 (28.1)	42 (33.1)	46 (35.4)	40 (20.5)
Every year		174 (38.5)	45 (35.4)	44 (33.8)	86 (44.1)
Every 1-5 years		90 (19.9)	26 (20.5)	19 (14.6)	45 (23.1)
Every five years		45 (10.0)	10 (7.9)	16 (12.3)	20 (10.3)
Only once		15 (3.3)	4 (3.1)	6 (4.6)	4 (2.1)
Back pain location of current episode, n (%)	11 (2.4)				
Thoracic only		19 (4.2)	4 (3.1)	7 (5.4)	8 (4.1)
Lumbar only		382 (84.5)	106 (83.5)	109 (83.8)	167 (85.6)
Both		51 (11.3)	17 (13.4)	14 (10.8)	20 (10.3)
Duration of current episode, n (%)	76 (16.8)				
0-6 weeks		297 (65.7)	74 (58.3)	67 (51.5)	156 (80.0)
6 weeks to 3 months		59 (13.1)	22 (17.3)	21 (16.2)	16 (8.2)
3 months or over		96 (21.2)	31 (24.4)	42 (32.3)	23 (11.8)
Back pain severity (NRS 0-10), mean (SD)	31 (6.9)	5.4 (2.3)	5.7 (2.2)	5.1 (2.3)	5.4 (2.4)
Back-related disability (RMDQ 0-24), median (IQR)	45 (10.0)	9 (5-13)	10 (6-14)	9 (6-13)	8 (3-13)
Sleep problems due to back pain, n (%)	24 (5.3)				
Weekly		189 (41.8)	60 (47.2)	49 (37.7)	80 (41.0)
Less than weekly		263 (58.2)	67 (52.8)	81 (62.3)	115 (59.0)
Morning stiffness, n (%)	26 (5.8)				
Significant or extreme		178 (39.3)	47 (37.0)	51 (39.2)	81 (41.5)
Moderate		144 (31.9)	44 (34.6)	48 (36.9)	51 (26.2)
Some or none		130 (28.8)	36 (28.3)	31 (23.9)	63 (32.3)
<b>Psychological variables</b>					
Kinesiophobia (FABQ-PA 0-24), median (IQR)	18 (4.0)	10 (5-14)	11 (6-14)	10 (5-15)	9 (3-13)
Depression (CES-D 0-60), median (IQR)	57 (12.6)	8 (4-15)	10 (4-17)	8.5 (4-15)	7 (4-13)
Pain catastrophizing (PCS 0-52), median (IQR)	35 (7.7)	10 (4-16)	11 (5-18)	12 (5-18)	7 (3-14)
Back beliefs (BBQ 9-45), mean (SD)	57 (12.6)	29.8 (7.0)	28.0 (6.9)	29.3 (7.2)	31.3 (6.7)
Expectations for back pain next 3 months, n (%)	19 (4.2)				

1					
2					
3	Fully recovered	115 (25.4)	19 (15.0)	24 (18.5)	72 (36.9)
4	Much better	226 (50.0)	66 (52.0)	71 (54.6)	89 (45.6)
5	No change or worse	111 (24.6)	42 (33.0)	35 (26.9)	33 (16.9)
6	Start Back Screening Tool risk profiles, n (%)	31 (6.9)			
7	Low	297 (65.7)	72 (56.7)	92 (70.8)	133 (68.2)
8	Medium	125 (27.7)	38 (29.9)	32 (24.6)	55 (28.2)
9	High	30 (6.6)	16 (12.6)	6 (4.6)	8 (4.1)
10	<b>Clinical variables</b>				
11	Physical performance (BPS 0-18), median (IQR)	20 (4.4)	5 (2-8)	7 (3-9)	5 (3-8)
12	Timed up and go, mean seconds (SD)	7 (1.5)	8.0 (2.5)	8.2 (3.0)	8.3 (2.3)
13	Positive diagnostic rule for radiculopathy, n (%)	38 (8.4)	99 (22.0)	37 (29.1)	31 (15.9)
14	Number of red flags (0-12), median (IQR)	50 (11.0)	1 (0-2)	1 (0-2)	1 (0-1)
15	Pain on active range of motion, n (%)	9 (2.0)	295 (65.3)	86 (67.7)	88 (67.7)
16	GP: General practitioner; PT: Physiotherapist; Chiro: Chiropractor; IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.				
17	* The presented characteristics are pooled estimates based on multiple imputation procedures				
18	^ AUDIT-C scores of $\geq 3/12$ for women and $\geq 4/12$ indicates hazardous alcohol consumption				
19					
20					
21					
22					
23					
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
51					
52					
53					
54					
55					
56					
57					
58					
59					
60					

**Table 2:** Multinomial regression analyses; multivariate associations between patient characteristics and choice of healthcare provider (dependent variable) \*

	GP (n=127)	p-value	PT (n=130)	p-value
	Odds ratio (95% CI)		Odds ratio (95% CI)	
<b>Block i) Sociodemographic variables</b>				
Age	1.03 (0.99, 1.07)	0.11	1.04 (1.00, 1.08)	<b>0.03</b>
Gender				
Female	1.53 (0.96, 2.45)	0.07	1.33 (0.83, 2.12)	0.24
Male (ref.)	1.00		1.00	
Marital status				
Married/cohabiting	0.67 (0.38, 1.19)	0.17	0.90 (0.51, 1.61)	0.73
Not married/cohabiting (ref.)	1.00		1.00	
Educational level				
Higher education	1.02 (0.64, 1.62)	0.94	1.08 (0.68, 1.73)	0.73
Lower education (ref.)	1.00		1.00	
Employment status				
Currently in paid work	0.86 (0.46, 1.62)	0.64	0.55 (0.30, 1.01)	0.05
No paid work (ref.)	1.00		1.00	
<b>Block ii) General health variables</b>				
Hazardous alcohol intake (AUDIT-C)				
Yes	1.20 (0.73, 1.97)	0.47	1.08 (0.64, 1.81)	0.77
No (ref.)	1.00		1.00	
Smoking status				
Yes	1.18 (0.56, 2.46)	0.67	0.64 (0.28, 1.48)	0.29
Previously	1.31 (0.77, 2.23)	0.32	1.11 (0.67, 1.83)	0.70
No (ref.)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.96 (0.93, 1.00)	<b>0.03</b>	0.98 (0.95, 1.01)	0.19
Mental component	0.97 (0.95, 1.00)	<b>0.02</b>	1.01 (0.98, 1.03)	0.73
BMI	0.98 (0.93, 1.04)	0.53	0.97 (0.92, 1.02)	0.28
Comorbidities (SCQ, 0-15)	1.07 (0.86, 1.33)	0.53	1.15 (0.95, 1.40)	0.17
Widespread pain				
Yes	0.22 (0.06, 0.81)	<b>0.02</b>	0.46 (0.18, 1.16)	0.10
No (ref.)	1.00		1.00	
Falls self-efficacy (FES-I, 16-64)	1.00 (0.95, 1.05)	0.98	1.03 (0.95, 1.05)	0.32
<b>Block iii) Current back pain and back pain history variables</b>				
Back pain severity (NRS, 0-10)	1.02 (0.91, 1.14)	0.77	0.90 (0.80, 1.01)	0.08
Back-related disability (RMDQ, 0-24)	1.06 (1.00, 1.12)	<b>0.04</b>	1.07 (1.01, 1.13)	<b>0.02</b>
Duration				
Over 3 months	2.92 (1.28, 6.66)	<b>0.01</b>	4.57 (1.99, 10.50)	<b>&lt;0.01</b>
6 weeks to 3 months	3.03 (1.27, 4.97)	<b>0.02</b>	3.17 (1.28, 7.84)	<b>0.01</b>
0-6 weeks (ref.)	1.00		1.00	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Morning stiffness				
Significant or extreme	0.76 (0.41, 1.42)	0.39	1.21 (0.64, 2.30)	0.55
Moderate	1.37 (0.74, 2.56)	0.32	2.03 (1.08, 3.81)	<b>0.03</b>
A little or none (ref.)	1.00		1.00	
Sleep problems attributable to back pain				
Weekly	1.09 (0.63, 1.89)	0.76	0.75 (0.41, 1.35)	0.33
Less than weekly (ref.)	1.00		1.00	
Previous back pain frequency				
Yearly	1.11 (0.65, 1.92)	0.70	1.00 (0.59, 1.69)	0.99
Not yearly (ref.)	1.00		1.00	
<b>Block iv) Psychological variables</b>				
Fear-avoidance (FABQ-PA, 0-24)	1.02 (0.97, 1.07)	0.38	1.03 (0.98, 1.08)	0.20
Pain catastrophizing (PCS, 0-52)	1.03 (0.99, 1.07)	0.10	1.06 (1.02, 1.10)	<b>&lt;0.01</b>
Depression symptoms (CESD, 0-60)	0.99 (0.95, 1.02)	0.44	0.99 (0.96, 1.03)	0.68
Back beliefs (BBQ, 9-45)	0.97 (0.93, 1.01)	0.22	0.99 (0.95, 1.03)	0.67
Expectation for back pain in 3 months				
Recovered	0.27 (0.13, 0.57)	<b>&lt;0.01</b>	0.38 (0.19, 0.79)	<b>0.01</b>
Much better	0.65 (0.35, 1.21)	0.18	0.85 (0.46, 1.56)	0.60
No change or worse (ref.)	1.00		1.00	
Start Back Screening tool risk category				
High risk	1.65 (0.52, 5.24)	0.40	0.29 (0.08, 1.10)	0.07
Medium risk	0.97 (0.52, 1.80)	0.92	<b>0.52 (0.28, 0.97)</b>	<b>0.04</b>
Low risk (ref.)	1.00		1.00	
<b>Block v) Clinical variables</b>				
Number of red flags (0-12)	1.25 (0.99, 1.58)	0.06	1.19 (0.96, 1.48)	0.12
Diagnostic tool for radiculopathy				
Positive	1.94 (1.08, 3.47)	<b>0.03</b>	1.52 (0.85, 2.73)	0.16
Negative (ref.)	1.00		1.00	
Pain on active range of motion				
Yes	0.95 (0.57, 1.58)	0.85	1.09 (0.67, 1.80)	0.72
No (ref.)	1.00		1.00	
Trunk mobility performance (BPS, 0-18)	1.16 (1.08, 1.24)	<b>&lt;0.01</b>	1.07 (1.00, 1.15)	<b>0.04</b>
Timed Up and Go, mean seconds	0.93 (0.83, 1.04)	0.20	1.00 (0.90, 1.11)	0.93

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

\* The multinomial regression analyses are based on pooled estimates from multiple regression analyses

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.

The chiropractic group (n=195) was the reference dependent variable.

Models were built block-wise within the five blocks: i) sociodemographic ii) general health iii) current episode and back pain history iv) psychological and v) clinical. All variables were included simultaneously.



**SUPPLEMENTARY FILE**

Table of Contents

Sensitivity analyses S1, complete case analyses: ..... 2

Sensitivity analyses S2, physiotherapy patients:..... 4

Assessment of generalizability S3: ..... 6

For peer review only

Sensitivity analyses S1, complete case analyses:

**Methods:**

Multiple multinomial regression. One model per variable block.

The total number of available cases per category will vary with number of missing for each block, and is thus shown for each block.

**Results:**

See Table S1 for details. No substantial changes in point estimates were detected in the multinomial regression analyses when comparing complete cases analyses to the pooled imputed estimates.

There were, however, some changes in p-values. In the complete case analyses, age and being in the SBT medium risk group were not significantly associated with choosing a PT compared to a chiropractor. Further, in the complete case analyses, having more red flags were significantly associated with choosing a GP compared to a chiropractor.

**Table S1:** Complete case analyses of multiple multinomial regression analyses. Chiropractic group is the reference group.

<b>Block i) Sociodemographic factors. Chiropractor n=181</b>				
	GP (n=113) Odds ratio (95% CI)	p-value	Physio (n=108) Odds ratio (95% CI)	p-value
Age	1.03 (0.99, 1.08)	0.11	1.03 (0.99, 1.07)	0.14
Gender				
Female	1.33 (0.81, 2.17)	0.26	1.40 (0.85, 2.33)	0.19
Male (ref)	1.00		1.00	
Marital status				
Married/cohabiting	0.66 (0.37, 1.19)	0.17	0.92 (0.49, 1.72)	0.79
Not married/cohabiting (ref)	1.00		1.00	
Educational level				
Higher education	1.02 (0.63, 1.65)	0.95	1.08 (0.66, 1.77)	0.77
Lower education (ref)	1.00		1.00	
Employment status				
Currently in paid work	0.96 (0.50, 1.86)	0.91	0.53 (0.27, 1.03)	0.06
No paid work (ref)	1.00		1.00	
<b>Block ii) General health factors. Chiropractor n=155</b>				
	GP (n=92) Odds ratio (95% CI)	p-value	Physio (n=89) Odds ratio (95% CI)	p-value
Hazardous alcohol intake (AUDIT-C)				
Yes	1.23 (0.70, 2.15)	0.48	1.67 (0.95, 2.92)	0.07
No (ref)	1.00		1.00	
Smoking				
Yes	1.37 (0.57, 3.26)	0.48	0.63 (0.22, 1.76)	0.37
Previously	1.47 (0.82, 2.66)	0.20	1.43 (0.81, 2.54)	0.22
No (ref)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.96 (0.92, 0.99)	<b>0.03</b>	0.97 (0.94, 1.01)	0.96
Mental component	0.95 (0.92, 0.98)	<b>0.002</b>	1.00 (0.94, 1.07)	0.99
BMI	0.99 (0.93, 1.06)	0.81	1.00 (0.94, 1.07)	0.99
Comorbidities (SCQ, 0-15)	1.02 (0.81, 1.29)	0.88	1.12 (0.89, 1.41)	0.33
Widespread pain				
Yes	0.16 (0.03, 0.79)	<b>0.03</b>	0.50 (0.15, 1.67)	0.26
No (ref)				
Falls self-efficacy (FESI, 16-64)	0.99 (0.93, 1.05)	0.73	0.99 (0.93, 1.06)	0.77
<b>Block iii) Current episode and back pain history. Chiropractor n=134</b>				
	GP (n=80) Odds ratio (95% CI)	p-value	Physio (n=92) Odds ratio (95% CI)	p-value

Back pain severity (NRS, 0-10)	1.06 (0.91, 1.22)	0.49	0.94 (0.82, 1.08)	0.40
Back-related disability (RMDQ, 0-24)	1.06 (0.99, 1.13)	0.12	1.06 (0.99, 1.13)	0.11
Duration				
Over 3 months	5.49 (2.34, 12.85)	<b>0.000</b>	9.00 (4.03, 20.13)	<b>0.000</b>
6 weeks to 3 months	4.92 (1.92, 12.61)	<b>0.001</b>	4.90 (1.91, 12.56)	<b>0.001</b>
0-6 weeks (ref)	1.00		1.00	
Morning stiffness				
Significant or extreme	1.02 (0.47, 2.24)	0.96	1.23 (0.57, 2.67)	0.60
Moderate	1.92 (0.88, 4.22)	0.10	2.36 (1.09, 5.14)	<b>0.03</b>
A little or none (ref)	1.00		1.00	
Sleep problems attributable to back pain				
- Weekly	0.82 (0.42, 1.62)	0.57	0.77 (0.39, 1.52)	0.45
- Less than weekly (ref)	1.00		1.00	
Previous back pain frequency				
- Yearly	1.06 (0.57, 1.96)	0.57	1.07 (0.58, 1.99)	0.82
- Not yearly (ref)	1.00		1.00	
<b>Block iv) Psychological factors. Chiropractor n=155</b>				
	GP (n=96)		Physio (n=94)	
	Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-value
Fear-avoidance (FABQ-PA, 0-24)	1.03 (0.98, 1.09)	0.25	1.03 (0.98, 1.09)	0.24
Pain catastrophizing (PCS, 0-52)	1.01 (0.96, 1.05)	0.82	1.06 (1.02, 1.11)	<b>0.01</b>
Depression symptoms (CESD, 0-60)	1.00 (0.95, 1.04)	0.88	0.99 (0.95, 1.04)	0.70
Back beliefs (BBQ, 9-45)	0.97 (0.92, 1.02)	0.23	0.99 (0.94, 1.04)	0.61
Expectation for back pain in 3 months				
Recovered	0.24 (0.11, 0.54)	<b>0.01</b>	0.43 (0.19, 0.96)	<b>0.04</b>
Much better	0.57 (0.29, 1.13)	0.11	0.83 (0.41, 1.67)	0.60
No change or worse(ref)	1.00		1.00	
Start Back Screening tool				
High risk	3.41 (0.69, 16.94)	0.13	0.21 (0.03, 1.72)	0.14
Medium risk	1.26 (0.65, 2.46)	0.50	0.51 (0.25, 1.03)	0.06
Low risk (ref)	1.00		1.00	
<b>Block v) Clinical variables. Chiropractor n=159</b>				
	GP (n=105)		Physio (n=110)	
	Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-value
Number of red flags (0-12)	1.30 (1.02, 1.67)	<b>0.04</b>	1.26 (0.99, 1.60)	0.06
Nerve involvement diagnostic tool				
Positive	2.34 (1.27, 4.31)	<b>0.01</b>	1.70 (0.93, 3.14)	0.09
Negative (ref)	1.00		1.00	
Pain on active range of motion				
Yes	0.76 (0.43, 1.32)	0.33	0.92 (0.54, 1.58)	0.77
No (ref)	1.00		1.00	
Trunk mobility performance (BPS, 0-18)	1.19 (1.10, 1.29)	<b>0.000</b>	1.10 (1.02, 1.19)	<b>0.01</b>
Timed up and go, mean seconds	0.90 (0.79, 1.03)	0.14	1.00 (0.88, 1.13)	0.94

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale - International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire - Physical Activity subscale; CES-D: Center for Epidemiological Studies Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.

## Sensitivity analyses S2, physiotherapy patients:

Analyses of differences between physiotherapy patients recruited before 01.01.2018 and after 01.01.2018. After 01.01.2018, there was direct access to physiotherapy in Norway, which potentially could change the population characteristics.

### Methods:

- Univariate analyses corresponding to measurement level and distribution: Chi square test or  $\chi^2$  for categorical variables, individual sample t-test for normally distributed continuous variables, Mann Whitney U-test for continuous variables with a skewed distribution
- We used the pooled estimates from multiple imputation that were used in the article table 1 and 2

### Results:

See Table S2 for details. We found statistically significant differences between PT patients recruited before and after 01.01.2018 on the BBQ and BPS. PT patients recruited before 01.01.2018 held significantly more optimistic beliefs about back pain, with a mean (SD) BBQ score of 30.3 (6.8) for patients recruited before 01.01.2018 compared to 27.3 (7.5) for patients recruited after 01.01.2018 ( $p=0.03$ ). PT patients recruited before 01.01.2018 had significantly better trunk mobility performance, with a median (IQR) of 5 (2-7) for patients recruited before 01.01.2018 compared to 7 (4-9.75) for patients recruited after 01.01.2018 ( $p=0.003$ ).

**Table S2:** Univariate analyses of differences between physiotherapy patients recruited before and after 01.01.2018.

	Physio before (n=90)	Physio after (n=40)	p-value
Age, median (IQR)	68 (62.75, 73)	68.5 (61.5, 76)	0.323
Sex female, n (%)	53 (58.9)	17 (42.5)	0.084
Married or living with partner, n (%)	69 (76.7)	29 (72.5)	0.580
Paid work, n (%)	30 (33.3)	12 (30.0)	0.606
Education level			0.317
- Low (elementary+high school)	51 (56.7)	19 (47.5)	
- High (university+ uni 4+)	39 (43.3)	21 (52.5)	
Health-related quality of life			
- Mental sumscore, median (IQR)	56.29 (51.01, 60.99)	54.63 (47.35, 60.37)	0.396
- Physical sumscore, mean (SD)	40.61 (7.91)	40.67 (8.30)	0.969
Hazardous alcohol consumption, n (%)	44 (48.9)	21 (52.5)	0.786
Smoking status			0.202
- Current smoker	9 (10)	4 (10)	
- Previous	46 (51.1)	14 (35)	
- Never	35 (38.9)	22 (55)	
Number of comorbidities, median (IQR)	2 (1, 2.25)	1 (1, 2)	0.235
BMI, median (IQR)	26.60 (24.41, 30.47)	26.37 (24.60, 29.27)	0.913

1				
2				
3	Fall last 6 weeks, n (%)	18 (20)	7 (17.5)	0.623
4	Falls self-efficacy, median (IQR)	20 (18, 23.35)	22.5 (17, 26.9)	0.424
5	Widespread pain, n (%)	5 (5.6)	2 (5.0)	0.880
6	Previous back pain, n (%)			0.479
7	- Monthly	35 (38.9)	11 (27.5)	
8	- Every year	30 (33.3)	14 (35.0)	
9	- Every 1-5 years	13 (14.4)	5 (12.5)	
10	- Every five years	8 (8.9)	8 (20.0)	
11	- Only once	4 (4.4)	2 (5.0)	
12	Duration of current episode, n (%)			0.538
13	- 0-6 weeks	30 (33.3)	11 (27.5)	
14	- 6 weeks to 3 months	17 (18.9)	11 (27.5)	
15	- 3 months or over	43 (47.8)	19 (47.5)	
16	Back pain, mean (SD)	5.22 (2.53)	4.69 (1.87)	0.208
17	Back-related disability, RMDQ, median (IQR)	8 (6, 13)	9.5 (4.25, 14)	0.808
18	Sleep problems due to back pain, n (%)			0.374
19	- Weekly	36 (40)	13 (32.5)	
20	- Less than weekly	54 (60)	27 (67.5)	
21	Morning stiffness, n (%)			0.753
22	- Significant or extreme	35 (38.9)	16 (40)	
23	- Moderate	35 (38.9)	13 (32.5)	
24	- Some or none	20 (22.2)	11 (27.5)	
25	Walking distance, n (%)			0.285
26	- More than 3km	40 (44.4)	16 (40.0)	
27	- 200m to 3km	41 (45.6)	16 (40.0)	
28	- Less than 200m	9 (10)	8 (20.0)	
29	Kinesiophobia (FABQ-PA), median (IQR)	10 (5, 15)	10.5 (5, 14)	0.842
30	Depression (CES-D), median (IQR)	8 (3.75, 14)	9.5 (5.25, 17.3)	0.305
31	Pain catastrophizing (PCS), median (IQR)	12 (5.3, 17)	11 (4, 19.6)	0.872
32	Back beliefs (BBQ), mean (SD)	30.3 (6.8)	27.3 (7.5)	<b>0.03</b>
33	Expectations for back pain next 3 months			0.821
34	- Fully recovered	17 (18.9)	7 (17.5)	
35	- Much better	50 (55.5)	21 (52.5)	
36	- No change or worse	23 (25.5)	12 (30.0)	
37	SBT risk profiles			0.163
38	- Low	68 (75.5)	24 (60)	
39	- Medium	18 (20)	14 (35)	
40	- High	4 (4.4)	2 (5)	
41	Back performance scale, median (range)	5 (2, 7)	7 (4, 9.75)	<b>0.003</b>
42	Timed up and go, median (IQR)	7.99 (6.66, 9.18)	7.42 (6.64, 9.86)	0.655
43	Probable nerve root involvement, n (%)	20 (22.2)	13 (32.5)	0.194
44	Number of red flags, median (range)	1 (0, 2)	1 (0, 2)	0.815
45	Pain on active range of motion, n (%)	61 (67.8)	27 (67.5)	0.905

IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire Physical Activity subscale; CES-D: Center for Epidemiological Studies Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
AUDIT-C scores of  $\geq 12$  for women and  $\geq 12$  indicates hazardous alcohol consumption



STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	4
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	5
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	5
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5-7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5-7
Bias	9	Describe any efforts to address potential sources of bias	8
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7
		(b) Describe any methods used to examine subgroups and interactions	NA
		(c) Explain how missing data were addressed	7
		(d) If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	8
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9
		(b) Give reasons for non-participation at each stage	-
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	9, 19-20
		(b) Indicate number of participants with missing data for each variable of interest	9, 19-20
Outcome data	15*	Report numbers of outcome events or summary measures	9
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9-10, 20-21

		(b) Report category boundaries when continuous variables were categorized	6-7
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10, suppl.
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	10
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	10-12
Generalisability	21	Discuss the generalisability (external validity) of the study results	12
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).



# BMJ Open

## Characteristics of older adults with back pain associated with choice of first primary care provider: a cross-sectional analysis from the BACE-N cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053229.R1
Article Type:	Original research
Date Submitted by the Author:	18-Jul-2021
Complete List of Authors:	Vigdal, Ørjan; Oslo Metropolitan University, Department of Physiotherapy Storheim, Kjersti; Oslo University Hospital; Oslo Metropolitan University, Department of Physiotherapy Munk Killingmo, Rikke ; Oslo Metropolitan University Faculty of Health Sciences, Department of Physiotherapy Småstuen, Milada; Oslo Metropolitan University, Department of Physiotherapy Grotle, Margreth; Oslo Metropolitan University, Department of Physiotherapy; Oslo universitetssykehus Ulleval, Research and Communication Unit for Musculoskeletal Health
<b>Primary Subject Heading</b>:	Health services research
Secondary Subject Heading:	Rehabilitation medicine, Geriatric medicine, General practice / Family practice
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, PRIMARY CARE, GERIATRIC MEDICINE, REHABILITATION MEDICINE

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Article title:**

Characteristics of older adults with back pain associated with choice of first primary care provider: a cross-sectional analysis from the BACE-N cohort study

**Authors:**

Ørjan Nesse Vigdal<sup>1</sup>

Kjersti Storheim<sup>2,1</sup>

Rikke Munk Killingmo<sup>1</sup>

Milada C. Småstuen<sup>1</sup>

Margreth Grotle<sup>1,2</sup>

**Author affiliations:**

<sup>1</sup>Department of Physiotherapy, Faculty of Health Science, OsloMet - Oslo Metropolitan University, Oslo, Norway

<sup>2</sup>Research and Communication Unit for Musculoskeletal Health (FORMI), Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway

**Corresponding author:**

Ørjan Nesse Vigdal

E-mail address: orvig@oslomet.no

Mailing address: OsloMet – Oslo Metropolitan University, Post box 4, St.Olavs plass, 0130 Oslo, Norway

Telephone number: +47 414 59 314

**Manuscript number of pages (excluding title page, including tables): 21**

**Word count: 4303**

**Total number of tables: 2**

**Supplementary files: 1**

## Abstract

**Objectives:** To describe characteristics of older adults with back pain in primary care, and to assess associations between patient characteristics and type of first primary care provider (general practitioner (GP), physiotherapist (PT) or chiropractor).

**Design:** Cross-sectional analysis from the BACE-N cohort study.

**Setting:** Norwegian GP, PT and chiropractic primary care centres.

**Participants:** Patients aged  $\geq 55$  years seeking Norwegian primary care with a new episode of back pain were invited to participate. Between April 2015 and February 2020, we included 452 patients: 127 first visited a GP, 130 first visited a PT and 195 first visited a chiropractor.

**Primary and secondary outcome measures:** For the first objective, the outcome measure was descriptive statistics of patient characteristics, covering the following domains: sociodemographic, general health, current and previous back pain, psychological and clinical factors. For the second objective, first primary care provider was the outcome measure. Associations between patient characteristics and visiting a GP or PT compared to a chiropractor were assessed with multiple multinomial regression analyses.

**Results:** Median (IQR) age was 66 (59-72) years. Levels of back-related disability was moderate to severe, with a median (IQR) Roland-Morris Disability Questionnaire (range 0-24) score of 9 (5-13). Recurring episodes were common, 301 (67%) patients had monthly or yearly recurrences. Patients with worse back-related disability, longer duration of symptoms, lower expectations for full recovery and worse physical performance measured with the Back Performance Scale had higher odds of visiting a GP or PT compared to a chiropractor ( $p < 0.05$ ).

**Conclusion:** Older back pain patients in primary care had moderate to severe levels of back-related disability, and most had recurring episodes. Our results suggest that older adult's choice of first primary care provider was associated with important patient characteristics, which highlights the need for caution with generalizations of study results across primary care populations.

**Trial registration number:** ClinicalTrials identifier: NCT04261309

## Data availability statement

Data not available.

1  
2  
3 **Keywords:** Back pain, older adults, primary care, characteristics, care-seeking behaviour  
4  
5  
6

7 **Article summary**

8  
9 **Strengths and limitations of this study**

- 10  
11 • This is the first study to compare characteristics of older adults with back pain visiting a GP,  
12 physiotherapist or chiropractor.  
13 • This study provides a thorough comprehensive overview of older adults with back pain, and thus  
14 contributes with important knowledge in a research field with few previous studies  
15 • It was not possible to obtain data on eligible patients that were not invited or declined to  
16 participate in the study. This might reduce external validity.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## Introduction

Back pain is the number one cause of years lived with disability globally, with an estimated point prevalence of 11.9% [1, 2]. Older adults have historically been under-represented in back pain research [3, 4], but have recently received increased attention [5, 6]. Although the prevalence of pathoanatomical findings on diagnostic imaging increases with age [7-9], the prevalence of serious pathology, such as vertebral fractures and neuropathic pain in older back pain patients in primary care is low. Studies have reported a prevalence of 6% and 2-11%, respectively [10, 11]. Moreover, studies in primary care have found significant national differences in the characteristics and burden of back pain in older adults [12, 13]. This highlights the importance of caution when generalizing results from studies from one setting to another.

Most patients seeking healthcare for back pain are treated in primary care [14]. In Norway, back pain is the reason for 10%, 27% and 86% of the visits to general practitioners (GP), physiotherapists (PT) and chiropractors, respectively [15]. Some studies suggest that choice of first primary care provider has consequences for future healthcare consumption, including imaging and opioid use [16, 17]. To optimize decision making regarding treatment, research and health policies, detailed knowledge of patient populations is required. Most of the previous studies exploring patient populations seeking primary care have compared GP and chiropractic populations, showing that patients seeking care from a GP have a higher overall burden of back pain compared to chiropractic patients [18-25]. Only a few studies include PT populations [26-29]. These studies suggest that patients seeking care from PTs are older and have more disability than those seeking care from chiropractors [26, 27, 29]. To the best of our knowledge, only one study has been performed in an exclusively older population [28]. This study found that older women seeking care from GPs reported worse back pain and worse health-related quality of life than older women visiting a PT or a chiropractor [28]. The study only included women between 59-64 years of age, and it is not clear if the results are also generalizable to men or adults over 65 years of age. Further, they did not examine back-related disability or other back pain factors, sociodemographic factors, psychological factors or clinical factors. Thus, there is still a considerable lack of knowledge regarding whether characteristics of older back pain patients differ according to their choice of first primary care provider.

Therefore, the aims of this study were 1) to describe the characteristics of patients  $\geq 55$  years of age seeking primary care for a new episode of back pain in terms of sociodemographic, general health, current back pain and back pain history, psychological and clinical characteristics, and 2) to assess if patient characteristics are associated with type of first primary care provider (GP, PT or chiropractor).

## Methods

### Design and setting

This cross-sectional study presents baseline data from the Back Complaints in the Elders – Norway (BACE-N) study, a prospective observational cohort study in Norwegian primary care. The BACE-N study is a part of the international BACE consortium, with research groups from Brazil, the Netherlands and Australia [6]. The BACE-N study protocol has been registered in ClinicalTrials.gov (Identifier NCT04261309). The study was classified as a quality assessment study by the Norwegian Regional Committee for Medical Research Ethics (reference no. 2014/1634/REK vest) and was approved by the Norwegian Social Science Data Service in 2015 (reference no. 42149).

Norwegian primary care is organized by the municipalities and financed through the National Insurance Scheme, the municipalities, and patient co-payment [30]. There is direct access to GPs, PTs (from 2018) and chiropractors [30]. Patient co-payment rates vary between healthcare providers, with chiropractors generally having the highest co-payment cost [30]. Treatments provided usually differ between the healthcare providers. For example, patients visiting a GP is more likely to receive pharmacological therapy, patients visiting a PT is more likely to receive exercise therapy, and patients visiting chiropractors are more likely to receive manipulation therapy [15].

### Participants and recruitment procedure

Eligible patients were  $\geq 55$  years of age, seeking primary care from a GP, PT or chiropractor in primary care for a new episode of back pain. Back pain was defined as pain located in the region from the top of the scapula to the sacrum, with or without radiating leg pain. A new episode was defined as not having received healthcare for the same complaint in the last six months. Patients were excluded if they had difficulties completing the questionnaire due to language barriers, or if they had difficulties completing the clinical examination (for example wheelchair-bound patients). Participants received care as usual.

Patients were recruited from GPs, PTs, and chiropractors in urban and rural parts of Norway between April 2015 and February 2020, either during or immediately after the consultation. The primary care providers were instructed to invite consecutive patients. To facilitate the recruitment process, media advertisements were also used. Eligible patients received oral and written information about the study. The final screening for eligibility and inclusion to the study was performed by the researchers. All included patients signed an informed consent form before enrolment in the study. The baseline

1  
2  
3 measurements, consisting of questionnaires and a clinical examination, were collected as soon after  
4 the first primary care consultation as possible.  
5  
6  
7

## 8 **Measurements**

### 9 *Sociodemographic variables*

10 Information regarding age, sex, marital status, employment status and educational level were  
11 collected.  
12  
13  
14

### 15 *General health variables*

16 Health-related quality of life (HR-QoL) was measured using the Short-Form Health Survey 36-item  
17 (SF-36) physical and mental summary measures (range 0-100, higher score indicates better HR-QoL)  
18 [31]. Alcohol consumption was measured using the 3-item Alcohol Use Disorder Identification Test  
19 consumption questions (AUDIT-C) (range 0-12, higher score indicates higher alcohol consumption)  
20 [32]. Hazardous alcohol consumption was defined as an AUDIT-C score of  $\geq 3/12$  for women and  
21  $\geq 4/12$  for men [33, 34]. Smoking status (current smoker, previous smoker, non-smoker) was  
22 collected. The number of comorbidities was measured using the Self-Administered Comorbidity  
23 Questionnaire (SCQ) [35]. The SCQ has 13 pre-defined comorbidities and two optional comorbidities.  
24 Item 12, "back pain", was replaced with a third optional comorbidity. Widespread pain was  
25 measured using the pain drawing from McGill Pain Questionnaire and the revised criteria from Wolfe  
26 et al. for widespread pain [36, 37]. The number of falls during the last six weeks was collected, and  
27 falls self-efficacy was measured using the Falls Efficacy Scale-International (FES-I) (range 16-64,  
28 higher score indicated lower falls efficacy) [38].  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

### 42 *Current back pain and back pain history*

43 Back pain location (thoracic or lumbar, or both) was collected. Average back pain severity last week  
44 was measured using the Numeric Rating Scale (NRS) (range 0-10, higher score indicates higher back  
45 pain severity) [39]. Back-related disability was measured with the 24-item Roland-Morris Disability  
46 Questionnaire (RMDQ) (range 0-24, higher score indicated more back-related disability) [40]. Back  
47 pain duration was measured in days and categorized into "<6 weeks", "6 weeks to 3 months", and  
48 ">3 months". Frequency of previous back pain episodes (monthly, yearly, every 1-5 years, every five  
49 years, once) was collected. Sleep problems attributable to back pain were measured using item 5i  
50 from the Pittsburgh Sleep Quality Index (PSQI) [41], and dichotomized to "weekly/less than weekly".  
51 Morning stiffness was measured with item six from Knee injury and Osteoarthritis Outcome Score  
52 (KOOS) [42], where we replaced the word "knee" with "back".  
53  
54  
55  
56  
57  
58  
59  
60



### *Psychological variables*

Kinesiophobia was measured using the Fear-Avoidance Beliefs Questionnaire-Physical Activity subscale (FABQ-PA) (range 0-24, higher score indicates higher levels of kinesiophobia) [43]. Signs of depression were measured with the Center for Epidemiological Studies-Depression questionnaire (CES-D) (range 0-60, higher score indicates more signs of depression) [44]. Pain catastrophizing was measured using the Pain Catastrophizing Scale (PCS) (range 0-52, higher score indicates more pain catastrophizing) [45]. Beliefs and attitudes towards back pain was measured using the Back Beliefs Questionnaire (BBQ) (range 9-45, higher score indicates more positive beliefs) [46]. Start Back Screening Tool (SBT) was used to assess prognostic risk profiles [47]. Expectations of recovery from back pain within the next 3 months was assessed with a five-point scale, with the categories “Fully recovered”, “Much better”, “No difference”, “Much worse”, and “Worse than ever”.

### *Clinical variables*

Pain with active movements was assessed for forward flexion, lateral flexion and rotation of the back. Physical performance with focus on trunk mobility was assessed with the 6-item Back Performance Scale (BPS) (range 0-18, higher score indicates worse trunk mobility performance) [48]. Walking function was assessed with the Timed-Up-and-Go (TUG) [49]. Signs of radiculopathy was measured using a clinical diagnostic model that summarizes five items: Subjective sensory changes (1 point), radiating pain below the knee (2 points), leg pain worse than back pain (2 points), positive neural tension test (3 points) and neurological deficit of myotome, dermatome or reflexes in the lower limb (2 points) [50]. A score of  $\geq 5/10$  has been shown to indicate  $>80\%$  probability of radiculopathy [50]. Twelve red flags were assessed: Cancer, first episode of back pain, constant pain, unexplained weight loss, systemically unwell, fever, urinary retention or loss of bladder control, age  $\geq 75$  years, trauma cause of back pain, osteoporosis, cortisone use and severe morning stiffness.

### **Statistical analyses**

All analyses were performed using the IBM SPSS Statistics version 26 for Windows (IBM Corporation, Armonk, NY, USA). To handle missing data, five multiple imputation datasets with 10 iterations were created using regression estimation, and the pooled estimates are presented in this study. Patient characteristics were described with counts and percentages for categorical variables, mean and standard deviation (SD) for normally distributed continuous variables and median and interquartile range (IQR) for continuous variables with a skewed distribution. Mann-Whitney U-test was used to assess differences in days between first primary care contact and inclusion to the study between primary care practitioners, and between those recruited from primary care and those recruited from media advertisements. Multinomial regression was used to assess the strength of the associations

1  
2  
3 between patient characteristics and patient's choice of first primary care provider. First primary care  
4 provider (GP, PT or chiropractor) was the dependent variable. The chiropractic group was the largest,  
5 and therefore chosen as the reference group. Patient characteristics were organized into five blocks,  
6 for which we created separate models: i) Sociodemographic ii) general health iii) current back pain  
7 episode and back pain history iv) psychological variables and v) clinical variables. All variables in the  
8 block were simultaneously included in the model, without univariate pre-testing. The strength of  
9 associations is expressed as odds ratios (OR) with 95% confidence intervals (CI). We considered our  
10 study as exploratory, so no correction for multiple testing was performed [51]. P-values <0.05 were  
11 thus considered statistically significant. All tests were two-sided.  
12  
13  
14  
15  
16  
17  
18  
19

### 20 *Assessment of generalizability*

21 Because of economic and practical reasons, we were unable to collect data on eligible participants  
22 that declined to participate or for other reasons were not invited. Therefore, we performed a  
23 descriptive comparison of the BACE-N on age, sex, nationality, educational level, work status, marital  
24 status, BMI, alcohol use, HR-QoL, depression and walking distance with individual data from a  
25 subsample from the study "The Norwegian study on life course, ageing and generation (NORLAG)"  
26 [52, 53]. This study used a random sampling strategy in the general population and included 11028  
27 participants. The subsample (NORLAG MSK) consisted of 794 participants collected in 2017. The  
28 participants of the subsample were ≥55 years of age and had at least one musculoskeletal complaint.  
29  
30  
31  
32  
33  
34  
35

### 36 *Sensitivity analyses*

37 We performed three sensitivity analyses: 1) To assess possible bias introduced by the multiple  
38 imputation procedure, the multiple multinomial regression analyses were performed on complete  
39 case data. We included a bootstrapping approach to assess the robustness of the coefficients. 2)  
40 Because PT services became available through direct access in Norway from 01.01.2018,  
41 characteristics of PT patients recruited before and after 01.01.2018 were compared using individual  
42 sample t-tests or Mann-Whitney U-tests for continuous variables, and chi-square tests for categorical  
43 variables. 3) We performed the multiple multinomial regression analyses in the subgroup with low  
44 back pain only. Results from the sensitivity analyses are available in supplementary material S1  
45 through S3.  
46  
47  
48  
49  
50  
51  
52  
53

### 54 *Sample size consideration*

55 Sample size was considered for the BACE-N study as a whole, with the following criteria: Having  
56 sufficient statistical power for up to 14 variables in a multivariate logistic regression analysis using  
57 the "10 events per variable" rule [54], with an outcome prevalence of 40%, and allowing for a  
58  
59  
60

1  
2  
3 dropout-rate of 20%. This yielded a preferred sample size of 450 participants. As the multinomial  
4 regression models in this study includes a maximum of 8 independent variables, we expect the  
5 sample size to be sufficient.  
6  
7  
8  
9

### 10 **Patient and public involvement**

11 Patient representatives were part of the scientific board of the study and involved in designing and  
12 establishing BACE-N. Results will be disseminated to the recruiting primary care providers and the  
13 participating patients in an annual newsletter.  
14  
15  
16  
17  
18  
19

### 20 **Results**

21 A total of 452 patients were included in the study, 127 first visited a GP, 130 first visited a PT and 195  
22 first visited a chiropractor. Eighteen patients were included from media advertisements. Median  
23 (IQR) number of days from first primary care contact to inclusion in the study was 13 (3-21) days for  
24 GP patients, 9 (3-21) for physiotherapy patients and 5 (1-13) for chiropractic patients. The duration  
25 was significantly shorter for chiropractic patients compared to GP patients ( $p<0.01$ ) and PT patients  
26 ( $p<0.01$ ). There was no statistically significant difference in duration from first primary care contact  
27 to inclusion between those recruited directly from primary care practices (median (IQR) 7 (2-15)  
28 days), and those recruited through media advertisements (median (IQR) 16 (1-28) days) ( $p=0.315$ ).  
29  
30  
31  
32  
33  
34  
35  
36  
37

#### 38 *Patient characteristics*

39 Missingness ranged from 0.0 to 16.8% for the variables, and total missingness was 4.4% across all  
40 values. Rates of missingness was similarly distributed across the primary care provider groups.  
41 Consult table 1 for details regarding patient characteristics. The median age of the patients was 66,  
42 around half of the patients were women, were in paid work, and had university-level education. Half  
43 of the patients had a hazardous alcohol consumption level, and nearly 60% of them were either  
44 current or previous smokers. One in six patients had experienced a fall during the last six weeks. Half  
45 of the patients had one or more comorbidities.  
46  
47  
48  
49  
50  
51

52 Most patients reported moderate levels of back pain and moderate to severe levels of back-related  
53 disability with a median (IQR) RMDQ-score of 9 (5-13). Almost 70% of the patients experienced  
54 monthly or yearly recurrences of back pain. Over 40% experienced weekly sleep problems  
55 attributable to back pain, and 70% experienced moderate to extreme morning stiffness. Two thirds  
56 of the patients had a low-risk profile according to the SBT, and only 6.6% had a high-risk profile.  
57  
58  
59  
60

1  
2  
3 Expectations of recovery were generally high, with three out of four expecting to be much better or  
4 fully recovered within three months.  
5  
6  
7

#### 8 *Associations between patient characteristics and type of first primary care provider*

9

10 Table 2 presents the associations from the multinomial regression analyses. Patients with higher  
11 back-related disability, longer duration of symptoms, worse physical performance, probable  
12 radiculopathy, poorer HR-QoL and lower expectations of being fully recovered within the next three  
13 months were more likely to visit a GP compared to a chiropractor. Patients with widespread pain  
14 were more likely to visit a chiropractor than a GP. The characteristics strongest associated with  
15 choosing a GP versus a chiropractor were duration of symptoms, widespread pain and expectation of  
16 being fully recovered.  
17  
18  
19  
20  
21  
22

23 Patients that were older, had a longer duration of symptoms, higher back-related disability,  
24 moderate morning stiffness, higher levels of pain catastrophizing, worse physical performance, lower  
25 expectations of being fully recovered within the next three months were more likely to visit a PT  
26 compared to a chiropractor. Patients in the SBT medium or high risk group were more likely to visit a  
27 chiropractor compared to a PT. The characteristics strongest associated with choosing a PT versus a  
28 chiropractor were duration of symptoms and expectation of being fully recovered.  
29  
30  
31  
32  
33  
34

35 Gender, education level, marital status, employment status, comorbidities, back pain severity, sleep  
36 problems, kinesiophobia, depressive signs, back beliefs, red flags, pain on active range of motion and  
37 Timed Up and Go-scores were not associated with type of primary care provider.  
38  
39  
40  
41

#### 42 *Assessment of generalizability*

43

44 The BACE-N study sample had more men (48% versus 36.3% in NORLAG MSK), more participants with  
45 high educational level (44% versus 28.6% in NORLAG MSK), more participants currently in paid work  
46 (45.3% versus 31.6% in NORLAG MSK), and more participants living with a partner (76.8% versus  
47 62.2% in NORLAG MSK). Age, nationality, alcohol consumption, BMI, depressive signs, HR-QoL and  
48 walking distance were similar for BACE-N and NORLAG MSK. See supplementary material S4 for  
49 further details.  
50  
51  
52  
53  
54  
55  
56

## 57 **Discussion**

58 This study showed that nearly all older patients with back pain had experienced back pain previously,  
59 and for most patients this episode was the latest of a series of annually or monthly recurring  
60

1  
2  
3 episodes. This is in accordance with several studies on back pain trajectories, where episodic or  
4 fluctuating pain was shown to be common both in the short and long term [55-58]. Further, patients  
5 with more severe back-related disability and other symptoms and signs were overall more likely to  
6 visit a GP or a physiotherapist than a chiropractor. Contrary to this finding, patients with widespread  
7 pain were more likely to choose a chiropractor over a GP. This is the first study to assess associations  
8 of a broad range of patient characteristics and choice of first primary care provider in an older  
9 population. Older adults have previously been under-represented in back pain studies [3, 4], and the  
10 evidence underlying treatment decisions in this age group may have been over-reliant on studies  
11 performed in younger populations. Thus, this study provides evidence to improve knowledge about  
12 older adults with back pain. This may prove important for clinical guideline development and  
13 informing stakeholders aiming to improve quality of care for older adults with back pain.  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 The burden of back pain and psychological profile were comparable between younger Norwegian  
24 back pain cohorts and the older BACE-N sample [59, 60]. The characteristics of the included patients  
25 in this study was largely comparable to the BACE-study from the Netherlands [12, 61], with a few  
26 exceptions. Both in our total study sample and our GP subsample, a larger proportion of patients had  
27 paid work, fewer experienced their first episode of back pain, and they reported lower levels of  
28 kinesiophobia and pain catastrophizing compared to the Dutch study sample. When comparing our  
29 results to the Brazilian BACE-study [12, 62], the Brazilian study had a higher proportion of women.  
30 Further, our study sample had more patients in paid work, more patients with hazardous alcohol  
31 consumption patterns, more smokers, the patients had fewer comorbidities, lower levels of back-  
32 related disability and back pain severity, kinesiophobia, depression signs, and pain catastrophizing  
33 compared to the Brazilian BACE-sample. These differences between populations within the BACE  
34 consortium might be explained in part by minor differences in recruitment strategies in the different  
35 countries [12] or differences in how primary care is organized in the different countries. In the  
36 Netherlands, patients were recruited exclusively from a GP setting [61], whereas in Brazil patients  
37 were recruited from primary care centres or health centres specialized in geriatrics [62]. Another  
38 possible explanation may be cultural differences in the expression and interpretation of and coping  
39 with pain [63].  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 In line with previous research on healthcare utilization for back pain in younger populations [19, 21-  
54 23, 25-28], our results suggest that patients with “less complex” characteristics were more likely to  
55 visit a chiropractor compared to a GP or a physiotherapist. Unsurprisingly, studies using bivariate  
56 analyses [18, 20, 23, 25, 28, 29] to compare the provider groups find more significant associations or  
57 differences than studies using multivariate analyses [19, 21, 22, 26, 27]. However, regardless of  
58  
59  
60

1  
2  
3 statistical approach, these studies suggest that patients who seek chiropractic care have an overall  
4 lower burden of back pain compared to patients seeking GP or PT care [18-23, 25]. One notable  
5 exception is the study of Eklund et al. [24], which found that Swedish chiropractic patients had more  
6 pain and worse psychological and behavioural characteristics compared to a sample of sick-listed  
7 primary care (specific provider unknown) patients at high risk for chronicity. Our finding showing that  
8 patients with widespread pain were more likely to choose a chiropractor over a GP was contrary to  
9 the general pattern of chiropractic patients being less “complex.” To the best of our knowledge, no  
10 previous studies have compared prevalence of widespread pain in the two populations, but one  
11 study showed that GP patients had more musculoskeletal comorbidities [25], possibly implying more  
12 widespread pain. Two previous studies found an association between higher age and odds of seeking  
13 care from a physiotherapist compared to a chiropractor [26, 27], in line with our results.  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 Many of the patient characteristics associated with choice of primary care provider in this study have  
24 previously been found to be significant prognostic factors for the persistent back-related disability  
25 and back pain in older people. For example, duration of back pain and expectation of improvement  
26 [64-69], and higher levels of back-related disability [65-70], are consistently reported as significant  
27 prognostic factors for a poor outcome of a back pain episode. A few studies in older people have  
28 found that single symptoms of neurological involvement such as leg pain below the knee, and the  
29 diagnosis of spinal stenosis were prognostic factors for the outcome of a back pain episode [64, 67].  
30 We combined single symptoms of neurological involvement into a compound measure, but it is likely  
31 that older patients with radiculopathy have worse outcomes than those without radiculopathy.  
32 Although slightly different from widespread pain, the presence of multi-site pain has also in some  
33 studies been found to be a prognostic factor for the outcome of back pain in older adults [67, 71].  
34 The impact of pain catastrophizing on the clinical course of back pain is less clear in older adults [66,  
35 69] compared to younger populations [72], but it is not unreasonable to believe that pain  
36 catastrophizing may be a prognostic factor for back pain in older adults. Thus, the associations  
37 between potential prognostic factors and choice of first primary care provider imply that we can  
38 expect the clinical course of patients in the three primary care groups to be different. Further, they  
39 imply that caution should be exercised when generalizing across primary care populations.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 The results of this study need to be viewed with consideration of some limitations. We instructed the  
54 recruiting primary care providers to invite consecutive patients, but because of obvious time  
55 constraints in clinical practice we could not ask them to keep record of how many declined to  
56 participate, nor of eligible patients that were not invited. This recruitment strategy increases the risk  
57 of selection bias, and thus could reduce the external validity of the study. To compensate for this  
58  
59  
60

1  
2  
3 limitation, we compared the BACE-N sample with the NORLAG MSK subsample. The characteristics of  
4 the two samples were largely comparable, but BACE-N has more men, more participants with higher  
5 education, more in paid work, and more living with their partner. Sex and education level have  
6 previously been shown to be associated with back pain severity and back-related disability in older  
7 adults [12, 13]. Thus, it may be possible that the levels of back pain and back-related disability  
8 presented in this study are slightly underestimated. The NORLAG MSK subsample is sampled from  
9 the general population, which may not be representative of those who seek care. However, the most  
10 important determinants of care-seeking for back pain seems to be pain severity and disability levels  
11 [73]. We therefore believe the assessment to be justified.  
12  
13  
14  
15  
16  
17  
18  
19

20 Another limitation may be the analysis strategy. We chose to keep the variables in the five blocks to  
21 provide a broad assessment of the differences in case-mix in the three primary care settings. To limit  
22 the number of statistical tests performed, univariate pre-testing and testing a “final model” across  
23 blocks were avoided. Furthermore, a different organization of the variables, for example strictly  
24 adhering to the biopsychosocial model [74] or Andersen’s behavioural model of health services use  
25 [75], may have yielded slightly different results. However, our results are largely supported by  
26 previous studies, so the potential differences because of analysis strategy or variable organization  
27 may be negligible. A third limitation is that we were unable to examine some possibly important  
28 determinants for healthcare use, such as access to different providers, patient’s familiarity with  
29 providers, the patient’s economic situation and social network referrals [75-77]. These factors may  
30 be the most important determinants in driving the patient’s choice of first primary care provider, and  
31 including these factors would have given an even broader overview of associations between  
32 individual and contextual characteristics and choice of primary care provider. We suggest that future  
33 research focus on examining the contextual and social factors associated with healthcare service use.  
34 Finally, generalization of our results to other healthcare systems may be limited. Different healthcare  
35 systems may have different access to care, different payment schemes and different professional  
36 training and responsibilities for the healthcare providers, all of which may impact health services  
37 utilization and consequently the patient characteristics associated with choosing different primary  
38 care providers [75, 78, 79].  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 **Conclusion**

55 We found that nearly all older adults with back pain seeking primary care had experienced back pain  
56 previously, and recurring episodes were common. In general, patients with more severe back-related  
57 disability and other clinical symptoms and signs were more likely to visit a GP or a physiotherapist  
58 than a chiropractor. Our results suggest that important patient characteristics are associated with  
59  
60

1  
2  
3 older adult's choice of primary care providers due to back pain, which may affect the clinical course  
4 of back pain for these patients. The findings highlight the need for caution with generalization of  
5 study results across primary care populations. This is an important consideration for healthcare  
6 providers, for the development and implementation of clinical practice guidelines, and for regulators  
7 when developing primary care pathways for back pain. Further research is needed in assessing if the  
8 choice of primary care provider affects future care pathways and the clinical course of back pain in  
9 older adults.  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **Acknowledgments**

19  
20 The authors thank all study participants for their significant contribution. Further, we would like to  
21 thank all the recruiting healthcare providers: Kerstin Ulrich, Lise Lothe, Tim Raven, Andreas Hoff  
22 Nordvik, Christoffer Børsheim, Steinar Forshei, Mette Brekke, Daniel Major, Jan Harald Lønn, Agnes  
23 Mordt, Andrea Kolstad, Mathias Svanevik, Alexander Diesen, Lars Gullestad, Ida Svalstuen, Joakim  
24 Ordahl, Mona Øversveen, Jørgen Kongtorp, Are Hansen, Geir Haram, Palwinder Singh, Svein Erik  
25 Sandelien, Bent Ulseth, Harald Nordheim, Ola Sand, Ragnhild Perstølen, Jorun Salater, Anna Allen-  
26 Unhammer, Morten Nilsen, Haakon Lilleeng Asmyhr, Philip Wilkens, Ane Klevberg, Eli Magnesen,  
27 Aleksander Killingmo, Bjørn Tore Bjørkedal, Stina Lund, Daniel Ekeberg, Berte Marie Enger, Johan  
28 Edvard Tellum, Morten de la Cruz, Bård Kvam, Marte Paulsen, Astrid Figger, Christian Mayer, Trond  
29 Magne Aasberg, Thea Tømmervåg, Jørgen Øyen, Håvard Nordås, Tore Viste Ollestad, Olav Aase,  
30 Renate Meier, Bjørn Røe, Jørgen Øyen, Jack Johnson, Carl-Erik Høgquist, Ingrid Hystad, Mats  
31 Thorbeck, Elisabeth Barø, Kaja McCormick, Lars Martin Fischer, Martin Haagenen, Cathrine  
32 Rosslund, Marianne Storberget, Nora Helk, Grete Bråten, Annecken Lister Haugen, Hege Herstad,  
33 Cathrine Natland, Frøydis Blaker Åsbø, Ola Klaastad, Thorleif Henning Monsen, Geir Wiik, Jørn  
34 Christian Halvorsen, Tonje Høgdahl Mysen, Tine Tandberg, Eir Marie Bergan, Rune Solheim, Ole  
35 Kristoffer Larsen. We thank the research assistants for assisting with clinical examinations and  
36 providing baseline questionnaires to study participants. We thank the BACE-N scientific board, and  
37 the BACE Consortium for important input with designing the study.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 **Author contributions:**

52 ØNV: Study design, data collection, data analyses, manuscript draft. KS: Study design, data  
53 interpretation, critical revision. RMK: Data collection, data interpretation, critical revision. MCS:  
54 Statistical advisor, data interpretation, critical revision. MG: Principal investigator, study design, data  
55 interpretation, critical revision.  
56  
57  
58  
59  
60



**Funding:**

This work was funded by The Norwegian Fund for Post-Graduate Training in Physiotherapy (grant number 90749). The funding body was not involved in designing the study, data collection, analyses or interpretation of data, nor in writing the manuscript.

**Competing interests**

The authors declare no competing interests.

For peer review only

## REFERENCES

1. Vos, T., et al., *Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010*. The Lancet, 2012. **380**(9859): p. 2163-2196.
2. Hoy, D., et al., *A systematic review of the global prevalence of low back pain*. Arthritis & Rheumatism, 2012. **64**(6): p. 2028-2037.
3. Bressler, H.B., et al., *The Prevalence of Low Back Pain in the Elderly: A Systematic Review of the Literature*. Spine, 1999. **24**(17): p. 1813-1819.
4. Paeck, T., et al., *Are older adults missing from low back pain clinical trials? A systematic review and meta-analysis*. Arthritis Care Res (Hoboken), 2014. **66**(8): p. 1220-6.
5. Jarvik, J.G., et al., *Study protocol: the back pain outcomes using longitudinal data (BOLD) registry*. BMC musculoskeletal disorders, 2012. **13**(1): p. 64.
6. Scheele, J., et al., *Back complaints in the elders (BACE); design of cohort studies in primary care: an international consortium*. BMC Musculoskelet Disord, 2011. **12**: p. 193.
7. Kalichman, L., et al., *Spinal stenosis prevalence and association with symptoms: the Framingham Study*. The Spine Journal, 2009. **9**(7): p. 545-550.
8. Van Der Klift, M., et al., *The incidence of vertebral fractures in men and women: the Rotterdam Study*. Journal of bone and mineral research, 2002. **17**(6): p. 1051-1056.
9. de Schepper, E.I., et al., *The association between lumbar disc degeneration and low back pain: the influence of age, gender, and individual radiographic features*. Spine, 2010. **35**(5): p. 531-536.
10. Enthoven, W.T., et al., *Prevalence and "Red Flags" Regarding Specified Causes of Back Pain in Older Adults Presenting in General Practice*. Physical Therapy, 2016. **96**(3): p. 305-312.
11. Enthoven, W.T., et al., *Back Complaints in Older Adults: Prevalence of Neuropathic Pain and Its Characteristics*. Pain Medicine, 2013. **14**(11): p. 1664-1672.
12. Jesus-Moraleida, F.R., et al., *Back Complaints in the Elders in Brazil and the Netherlands: a cross-sectional comparison*. Age and ageing, 2017. **46**(3): p. 476-481.
13. Jarvik, J.G., et al., *Back pain in seniors: the back pain outcomes using longitudinal data (BOLD) cohort baseline data*. BMC Musculoskeletal Disorders, 2014. **15**(1): p. 1-23.
14. Kinge, J.M., et al., *Musculoskeletal disorders in Norway: prevalence of chronicity and use of primary and specialist health care services*. BMC Musculoskeletal Disorders, 2015. **16**(1): p. 1-9.
15. Werner, E.L. and A. Indahl, *Kunnskap, praksis og holdninger til rygglidelser hos leger, fysioterapeuter og kiropraktorer*. Tidsskr Nor Laegeforen, 2005. **125**(13): p. 1794–7.
16. Fritz, J.M., J. Kim, and J. Dorius, *Importance of the type of provider seen to begin health care for a new episode low back pain: associations with future utilization and costs*. Journal of evaluation in clinical practice, 2016. **22**(2): p. 247-252.
17. Kazis, L.E., et al., *Observational retrospective study of the association of initial healthcare provider for new-onset low back pain with early and long-term opioid use*. BMJ Open, 2019. **9**(9): p. e028633.
18. Carey, T.S., et al., *Acute Severe Low Back Pain: A Population-based Study of Prevalence and Care-seeking*. Spine, 1996. **21**(3): p. 339-344.
19. Hurwitz, E.L. and H. Morgenstern, *The effects of comorbidity and other factors on medical versus chiropractic care for back problems*. Spine, 1997. **22**(19): p. 2254-2263.
20. Nyiendo, J., et al., *Patient characteristics and physicians' practice activities for patients with chronic low back pain: A practice-based study of primary care and chiropractic physicians*. 2001. **24**(2): p. 92-100.
21. Côté, P., J.D. Cassidy, and L. Carroll, *The treatment of neck and low back pain: who seeks care? who goes where?* Medical care, 2001: p. 956-967.

- 1
- 2
- 3 22. Sharma, R., M. Haas, and M. Stano, *Patient Attitudes, Insurance, and Other Determinants of*
- 4 *Self-Referral to Medical and Chiropractic Physicians*. American Journal of Public Health, 2003.
- 5 **93**(12): p. 2111-2117.
- 6
- 7 23. Hestbaek, L., et al., *Low back pain in primary care: a description of 1250 patients with low*
- 8 *back pain in danish general and chiropractic practice*. International journal of family
- 9 *medicine*, 2014. **2014**.
- 10
- 11 24. Eklund, A., et al., *Psychological and behavioral differences between low back pain*
- 12 *populations: a comparative analysis of chiropractic, primary and secondary care patients*.
- 13 *BMC Musculoskeletal Disorders*, 2015. **16**(1).
- 14
- 15 25. Hartvigsen, L., et al., *Baseline Characteristics May Help Indicate the Best Choice of Health*
- 16 *Care Provider for Back Pain Patients in Primary Care: Results From a Prospective Cohort*
- 17 *Study*. Journal of Manipulative and Physiological Therapeutics, 2020. **43**(1): p. 13-23.
- 18
- 19 26. Chevan, J. and D.L. Riddle, *Factors associated with care seeking from physicians, physical*
- 20 *therapists, or chiropractors by persons with spinal pain: a population-based study*. journal of
- 21 *orthopaedic & sports physical therapy*, 2011. **41**(7): p. 467-476.
- 22
- 23 27. Blanchette, M.-A., et al., *Workers' characteristics associated with the type of healthcare*
- 24 *provider first seen for occupational back pain*. BMC Musculoskeletal Disorders, 2016. **17**(1).
- 25
- 26 28. Sibbritt, D., et al., *Severity of back pain may influence choice and order of practitioner*
- 27 *consultations across conventional, allied and complementary health care: a cross-sectional*
- 28 *study of 1851 mid-age Australian women*. BMC Musculoskeletal Disorders, 2016. **17**(1).
- 29
- 30 29. Walker, B.F., R. Muller, and W.D. Grant, *Low back pain in Australian adults. Health provider*
- 31 *utilization and care seeking*. Journal of manipulative and physiological therapeutics, 2004.
- 32 **27**(5): p. 327-335.
- 33
- 34 30. Sperre Saunes, I., et al., *Norway: Health system review*. 2020.
- 35
- 36 31. Loge, J.H. and S. Kaasa, *Short Form 36 (SF-36) health survey: normative data from the general*
- 37 *Norwegian population*. Scandinavian Journal of Social Medicine, 1998. **26**(4): p. 250-258.
- 38
- 39 32. Bush, K., *The AUDIT Alcohol Consumption Questions (AUDIT-C). An Effective Brief Screening*
- 40 *Test for Problem Drinking*. Archives of Internal Medicine, 1998. **158**(16): p. 1789.
- 41
- 42 33. Bradley, K.A., et al., *AUDIT-C as a Brief Screen for Alcohol Misuse in Primary Care*. Alcoholism:
- 43 *Clinical and Experimental Research*, 2007. **31**(7): p. 1208-1217.
- 44
- 45 34. Norwegian National Advisory Unit on Concurrent Substance Abuse and Mental Health
- 46 *Disorders. AUDIT-C*. 23.09.2020]; Available from: [https://rop.no/kartleggingsverktoey/audit-](https://rop.no/kartleggingsverktoey/audit-c/)
- 47 [c/](https://rop.no/kartleggingsverktoey/audit-c/).
- 48
- 49 35. Sangha, O., et al., *The self-administered comorbidity questionnaire: A new method to assess*
- 50 *comorbidity for clinical and health services research*. Arthritis & Rheumatism, 2003. **49**(2): p.
- 51 156-163.
- 52
- 53 36. Wolfe, F., et al., *Revised chronic widespread pain criteria: development from and integration*
- 54 *with fibromyalgia criteria*. Scandinavian Journal of Pain, 2019. **20**(1): p. 77-86.
- 55
- 56 37. Melzack, R., *The McGill Pain Questionnaire: Major properties and scoring methods*. Pain,
- 57 1975. **1**(3): p. 277-299.
- 58
- 59 38. Helbostad, J.L., et al., *Validation of the Falls Efficacy Scale-International in fall-prone older*
- 60 *persons*. Age and Ageing, 2010. **39**(2): p. 259-259.
- 61
- 62 39. Von Korff, M., M.P. Jensen, and P. Karoly, *Assessing Global Pain Severity by Self-Report in*
- 63 *Clinical and Health Services Research*. Spine, 2000. **25**(24): p. 3140-3151.
- 64
- 65 40. Grotle, M., J. Brox, and N. Vollestad, *Cross-cultural adaptation of the Norwegian versions of*
- 66 *the Roland-Morris Disability Questionnaire and the Oswestry Disability Index*. Journal of
- 67 *Rehabilitation Medicine*, 2003. **35**(5): p. 241-247.
- 68
- 69 41. Buysse, D.J., et al., *The Pittsburgh Sleep Quality Index: a new instrument for psychiatric*
- 70 *practice and research*. Psychiatry research, 1989. **28**(2): p. 193-213.
- 71
- 72 42. Roos, E.M., et al., *Knee Injury and Osteoarthritis Outcome Score (KOOS)—Development of a*
- 73 *Self-Administered Outcome Measure*. Journal of Orthopaedic & Sports Physical Therapy,
- 74 1998. **28**(2): p. 88-96.

- 1  
2  
3 43. Grotle, M., J.I. Brox, and N.K. Vøllestad, *Reliability, validity and responsiveness of the fear-avoidance beliefs questionnaire: methodological aspects of the Norwegian version*. Journal of rehabilitation medicine, 2006. **38**(6): p. 346-353.
- 4  
5  
6 44. Radloff, L.S., *The CES-D scale: A self report depression scale for research in the general population*. Applied Psychological Measurements, 1977. **1**(3): p. 385-401.
- 7  
8 45. Fernandes, L., et al., *Cross-cultural adaptation and validation of the Norwegian pain catastrophizing scale in patients with low back pain*. BMC musculoskeletal disorders, 2012. **13**(1): p. 111.
- 9  
10  
11 46. Tingulstad, A., et al., *Back beliefs among elderly seeking health care due to back pain; psychometric properties of the Norwegian version of the back beliefs questionnaire*. BMC Musculoskeletal Disorders, 2019. **20**(1).
- 12  
13  
14 47. Hill, J.C., et al., *A primary care back pain screening tool: Identifying patient subgroups for initial treatment*. Arthritis & Rheumatism, 2008. **59**(5): p. 632-641.
- 15  
16  
17 48. Strand, L.I., R. Moe-Nilssen, and A.E. Ljunggren, *Back Performance Scale for the Assessment of Mobility-Related Activities in People With Back Pain*. Physical Therapy, 2002. **82**(12): p. 1213-1223.
- 18  
19  
20 49. Podsiadlo, D. and S. Richardson, *The Timed "Up & Go": A Test of Basic Functional Mobility for Frail Elderly Persons*. Journal of the American Geriatrics Society, 1991. **39**(2): p. 142-148.
- 21  
22  
23 50. Stynes, S., et al., *Clinical diagnostic model for sciatica developed in primary care patients with low back-related leg pain*. PloS one, 2018. **13**(4).
- 24  
25  
26 51. Bender, R. and S. Lange, *Adjusting for multiple testing—when and how?* Journal of Clinical Epidemiology, 2001. **54**(4): p. 343-349.
- 27  
28 52. Slagsvold, B., et al., *Life-course, ageing and generations in Norway:the NorLAG study*. Norsk Epidemiologi, 2012. **22**(2).
- 29  
30 53. Torsteinsen, A. and A. Holmøy, *Den norske studien av livsløp, aldring og generasjon—tredje runde (NorLAG3). Dokumentasjonsrapport*. 2019.
- 31  
32 54. Moons, K.G.M., et al., *Prognosis and prognostic research: what, why, and how?* BMJ, 2009. **338**: p. b375.
- 33  
34 55. Dunn, K.M., K. Jordan, and P.R. Croft, *Characterizing the course of low back pain: a latent class analysis*. American journal of epidemiology, 2006. **163**(8): p. 754-761.
- 35  
36 56. Dunn, K.M., P. Campbell, and K.P. Jordan, *Long-term trajectories of back pain: cohort study with 7-year follow-up*. BMJ open, 2013. **3**(12): p. e003838.
- 37  
38 57. Kongsted, A., et al., *Patients with low back pain had distinct clinical course patterns that were typically neither complete recovery nor constant pain. A latent class analysis of longitudinal data*. The Spine Journal: Official Journal Of The North American Spine Society, 2015. **15**(5): p. 885-894.
- 39  
40  
41 58. Kongsted, A., et al., *What have we learned from ten years of trajectory research in low back pain?* BMC Musculoskeletal Disorders, 2016. **17**: p. 1-11.
- 42  
43  
44 59. Nordstoga, A.L., et al., *Improvement in Work Ability, Psychological Distress and Pain Sites in Relation to Low Back Pain Prognosis: A Longitudinal Observational Study in Primary Care*. Spine, 2019. **44**(7): p. E423-E429.
- 45  
46  
47 60. Grotle, M., et al., *Prognostic factors in first-time care seekers due to acute low back pain*. European Journal of Pain, 2007. **11**(3): p. 290-298.
- 48  
49  
50 61. Scheele, J., et al., *Characteristics of older patients with back pain in general practice: BACE cohort study*. Eur J Pain, 2013. **18**(2): p. 279-87.
- 51  
52  
53 62. Jesus-Moraleida, F.R.D., et al., *The Brazilian Back Complaints in the Elders (Brazilian BACE) study: characteristics of Brazilian older adults with a new episode of low back pain*. Brazilian Journal of Physical Therapy, 2017.
- 54  
55  
56 63. Peacock, S. and S. Patel, *Cultural influences on pain*. Reviews in pain, 2008. **1**(2): p. 6-9.
- 57  
58 64. Scheele, J., et al., *Course and prognosis of older back pain patients in general practice: a prospective cohort study*. Pain, 2013. **154**(6): p. 951-7.
- 59  
60

- 1  
2  
3 65. Deyo, R.A., et al., *Trajectories of symptoms and function in older adults with low back disorders*. *Spine (Phila Pa 1976)*, 2015. **40**(17): p. 1352-62.
- 4  
5 66. Enthoven, W.T., et al., *Defining trajectories in older adults with back pain presenting in general practice*. *Age and ageing*, 2016. **45**(6): p. 878-883.
- 6  
7 67. Rundell, S.D., et al., *Predictors of Persistent Disability and Back Pain in Older Adults with a New Episode of Care for Back Pain*. *Pain Med*, 2017. **18**(6): p. 1049-1062.
- 8  
9 68. Jarvik, J.G., et al., *Long-term outcomes of a large, prospective observational cohort of older adults with back pain*. *Spine J*, 2018. **18**(9): p. 1540-1551.
- 10  
11 69. van der Gaag, W.H., et al., *Developing clinical prediction models for non-recovery in older patients seeking care for back pain: the BACE prospective cohort study*. *PAIN*, 2021. **162**(6): p. 1632-1640.
- 12  
13  
14 70. Van Den Berg, R., et al., *Clinical and radiographic features of spinal osteoarthritis predict long-term persistence and severity of back pain in older adults*. *Annals of Physical and Rehabilitation Medicine*, 2020.
- 15  
16 71. Rundell, S.D., et al., *Multisite Pain Is Associated with Long-term Patient-Reported Outcomes in Older Adults with Persistent Back Pain*. *Pain Med*, 2019.
- 17  
18 72. Wertli, M.M., et al., *Catastrophizing—a prognostic factor for outcome in patients with low back pain: a systematic review*. *The Spine Journal*, 2014. **14**(11): p. 2639-2657.
- 19  
20 73. Ferreira, M., et al., *Factors defining care-seeking in low back pain - A meta-analysis of population based surveys*. *European Journal of Pain*, 2010. **14**(7): p. 747.e1-747.e7.
- 21  
22 74. Waddell, G., *1987 Volvo award in clinical sciences. A new clinical model for the treatment of low-back pain*. *Spine*, 1987. **12**(7): p. 632-644.
- 23  
24 75. Andersen, R.M., *National health surveys and the behavioral model of health services use*. *Medical care*, 2008. **46**(7): p. 647-653.
- 25  
26 76. Davis, M.A., et al., *Regional Supply of Chiropractic Care and Visits to Primary Care Physicians for Back and Neck Pain*. *The Journal of the American Board of Family Medicine*, 2015. **28**(4): p. 481-490.
- 27  
28 77. Kirby, E.R., et al., *A qualitative study of influences on older women's practitioner choices for back pain care*. *BMC Health Services Research*, 2014. **14**(1): p. 131.
- 29  
30 78. Garrity, B., et al., *Unrestricted Direct Access to Physical Therapist Services Is Associated With Lower Health Care Utilization and Costs in Patients With New-Onset Low Back Pain*. *Physical Therapy*, 2019.
- 31  
32 79. Carey, K., et al., *Health insurance design and conservative therapy for low back pain*. *Am J Manag Care*, 2019. **25**(6): p. e182-e187.
- 33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1:** Baseline characteristics of study participants\*

	Missing, n (%)	Total (n=452)	GP (n=127)	PT (n=130)	Chiro (n=195)
<b>Sociodemographic variables</b>					
Age, median (IQR)	0 (0.0)	66 (59-72)	67 (60-73)	68 (63-74)	63 (58-71)
Female, n (%)	0 (0.0)	235 (52.0)	74 (58.3)	70 (53.8)	89 (46.1)
Marital status	19 (4.2)				
Married or living with partner, n (%)		347 (76.8)	90 (70.1)	98 (74.6)	158 (81.0)
Employment status	5 (1.1)				
Currently in paid work, n (%)		212 (45.3)	57 (43.3)	49 (31.5)	106 (55.9)
Educational level, n (%)	20 (4.4)				
Low (elementary + high school)		253 (56.0)	72 (56.7)	70 (55.1)	110 (56.4)
High (university level)		199 (44.0)	55 (43.3)	60 (44.9)	85 (43.6)
<b>General health variables</b>					
Health-related quality of life (SF-36 0-100)	41 (9.1)				
Mental sumscore, mean (SD)		52.5 (10.0)	50.5 (11.5)	53.4 (10.0)	53.2 (8.8)
Physical sumscore, mean (SD)		41.4 (8.4)	40.0 (7.9)	40.6 (8.0)	42.8 (8.9)
Hazardous alcohol consumption (AUDIT-C <sup>a</sup> ), n (%)	59 (13.1)	228 (50.4)	65 (51.1)	65 (50.0)	98 (50.2)
Smoking status, n (%)	22 (4.9)				
Current smoker		63 (13.9)	21 (16.5)	13 (10.0)	28 (14.3)
Previous		203 (44.9)	59 (46.4)	60 (46.2)	84 (43.1)
Never		186 (41.2)	47 (37.0)	57 (43.8)	83 (42.6)
Number of comorbidities (SCQ 0-15), median (IQR)	18 (4.0)	1 (1-2)	1 (0-2)	2 (1-2)	1 (1-5)
BMI, mean (SD)	14 (3.1)	27.6 (4.7)	27.6 (4.5)	27.5 (4.7)	27.7 (4.8)
Fall last 6 weeks, n (%)	24 (5.3)	73 (16.1)	13 (10.2)	24 (18.4)	35 (18.2)
Falls self-efficacy (FESI 16-64), mean (SD)	48 (10.6)	21.8 (6.0)	22.4 (6.3)	22.2 (6.1)	21.1 (5.7)
Widespread pain, n (%)	16 (3.5)	33 (7.3)	5 (4.0)	7 (5.3)	21 (10.8)
<b>Current back pain and back pain history variables</b>					
Previous back pain, n (%)	58 (12.8)				
Monthly		127 (28.1)	42 (33.1)	46 (35.4)	40 (20.5)
Every year		174 (38.5)	45 (35.4)	44 (33.8)	86 (44.1)
Every 1-5 years		90 (19.9)	26 (20.5)	19 (14.6)	45 (23.1)
Every five years		45 (10.0)	10 (7.9)	16 (12.3)	20 (10.3)
Only once		15 (3.3)	4 (3.1)	6 (4.6)	4 (2.1)
Back pain location of current episode, n (%)	11 (2.4)				
Thoracic only		19 (4.2)	4 (3.1)	7 (5.4)	8 (4.1)
Lumbar only		382 (84.5)	106 (83.5)	109 (83.8)	167 (85.6)
Both		51 (11.3)	17 (13.4)	14 (10.8)	20 (10.3)
Duration of current episode, n (%)	76 (16.8)				
0-6 weeks		297 (65.7)	74 (58.3)	67 (51.5)	156 (80.0)
6 weeks to 3 months		59 (13.1)	22 (17.3)	21 (16.2)	16 (8.2)
3 months or over		96 (21.2)	31 (24.4)	42 (32.3)	23 (11.8)
Back pain severity (NRS 0-10), mean (SD)	31 (6.9)	5.4 (2.3)	5.7 (2.2)	5.1 (2.3)	5.4 (2.4)
Back-related disability (RMDQ 0-24), median (IQR)	45 (10.0)	9 (5-13)	10 (6-14)	9 (6-13)	8 (3-13)
Sleep problems due to back pain, n (%)	24 (5.3)				
Weekly		189 (41.8)	60 (47.2)	49 (37.7)	80 (41.0)
Less than weekly		263 (58.2)	67 (52.8)	81 (62.3)	115 (59.0)
Morning stiffness, n (%)	26 (5.8)				
Significant or extreme		178 (39.3)	47 (37.0)	51 (39.2)	81 (41.5)
Moderate		144 (31.9)	44 (34.6)	48 (36.9)	51 (26.2)
Some or none		130 (28.8)	36 (28.3)	31 (23.9)	63 (32.3)
<b>Psychological variables</b>					
Kinesiophobia (FABQ-PA 0-24), median (IQR)	18 (4.0)	10 (5-14)	11 (6-14)	10 (5-15)	9 (3-13)
Depression (CES-D 0-60), median (IQR)	57 (12.6)	8 (4-15)	10 (4-17)	8.5 (4-15)	7 (4-13)
Pain catastrophizing (PCS 0-52), median (IQR)	35 (7.7)	10 (4-16)	11 (5-18)	12 (5-18)	7 (3-14)
Back beliefs (BBQ 9-45), mean (SD)	57 (12.6)	29.8 (7.0)	28.0 (6.9)	29.3 (7.2)	31.3 (6.7)
Expectations for back pain next 3 months, n (%)	19 (4.2)				
Fully recovered		115 (25.4)	19 (15.0)	24 (18.5)	72 (36.9)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Much better		226 (50.0)	66 (52.0)	71 (54.6)	89 (45.6)
No change or worse		111 (24.6)	42 (33.0)	35 (26.9)	33 (16.9)
Start Back Screening Tool risk profiles, n (%)	31 (6.9)				
Low		297 (65.7)	72 (56.7)	92 (70.8)	133 (68.2)
Medium		125 (27.7)	38 (29.9)	32 (24.6)	55 (28.2)
High		30 (6.6)	16 (12.6)	6 (4.6)	8 (4.1)
<b>Clinical variables</b>					
Physical performance (BPS 0-18), median (IQR)	20 (4.4)	5 (2-8)	7 (3-9)	5 (3-8)	4 (1-7)
Timed up and go, mean seconds (SD)	7 (1.5)	8.0 (2.5)	8.2 (3.0)	8.3 (2.3)	7.8 (2.2)
Positive diagnostic rule for radiculopathy, n (%)	38 (8.4)	99 (22.0)	37 (29.1)	31 (23.8)	31 (15.9)
Number of red flags (0-12), median (IQR)	50 (11.0)	1 (0-2)	1 (0-2)	1 (0-2)	1 (0-1)
Pain on active range of motion, n (%)	9 (2.0)	295 (65.3)	86 (67.7)	88 (67.7)	120 (61.5)

GP: General practitioner; PT: Physiotherapist; Chiro: Chiropractor; IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
\* The presented characteristics are pooled estimates based on multiple imputation procedures  
^ AUDIT-C scores of ≥3/12 for women and ≥4/12 indicates hazardous alcohol consumption

**Table 2:** Multinomial regression analyses; multivariate associations between patient characteristics and choice of healthcare provider (dependent variable) \*

	GP (n=127)	p-value	PT (n=130)	p-value
	Odds ratio (95% CI)		Odds ratio (95% CI)	
<b>Block i) Sociodemographic variables</b>				
Age	1.03 (0.99-1.07)	0.11	1.04 (1.00-1.08)	<b>0.03</b>
Gender				
Female	1.53 (0.96-2.45)	0.07	1.33 (0.83-2.12)	0.24
Male (ref.)	1.00		1.00	
Marital status				
Married/cohabiting	0.67 (0.38-1.19)	0.17	0.90 (0.51-1.61)	0.73
Not married/cohabiting (ref.)	1.00		1.00	
Educational level				
Higher education	1.02 (0.64-1.62)	0.94	1.08 (0.68-1.73)	0.73
Lower education (ref.)	1.00		1.00	
Employment status				
Currently in paid work	0.86 (0.46-1.62)	0.64	0.55 (0.30-1.01)	0.05
No paid work (ref.)	1.00		1.00	
<b>Block ii) General health variables</b>				
Hazardous alcohol intake (AUDIT-C)				
Yes	1.20 (0.73-1.97)	0.47	1.08 (0.64-1.81)	0.77
No (ref.)	1.00		1.00	
Smoking status				
Yes	1.18 (0.56-2.46)	0.67	0.64 (0.28-1.48)	0.29
Previously	1.31 (0.77-2.23)	0.32	1.11 (0.67-1.83)	0.70
No (ref.)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.96 (0.93-1.00)	<b>0.03</b>	0.98 (0.95-1.01)	0.19
Mental component	0.97 (0.95-1.00)	<b>0.02</b>	1.01 (0.98-1.03)	0.73
BMI	0.98 (0.93-1.04)	0.53	0.97 (0.92-1.02)	0.28
Comorbidities (SCQ, 0-15)	1.07 (0.86-1.33)	0.53	1.15 (0.95-1.40)	0.17
Widespread pain				
Yes	0.22 (0.06-0.81)	<b>0.02</b>	0.46 (0.18-1.16)	0.10
No (ref.)	1.00		1.00	
Falls self-efficacy (FES-I, 16-64)	1.00 (0.95-1.05)	0.98	1.03 (0.95-1.05)	0.32
<b>Block iii) Current back pain and back pain history variables</b>				
Back pain severity (NRS, 0-10)	1.02 (0.91-1.14)	0.77	0.90 (0.80-1.01)	0.08
Back-related disability (RMDQ, 0-24)	1.06 (1.00-1.12)	<b>0.04</b>	1.07 (1.01-1.13)	<b>0.02</b>
Duration				
Over 3 months	2.92 (1.28-6.66)	<b>0.01</b>	4.57 (1.99-10.50)	<b>&lt;0.01</b>

6 weeks to 3 months	3.03 (1.27-4.97)	<b>0.02</b>	3.17 (1.28-7.84)	<b>0.01</b>
0-6 weeks (ref.)	1.00		1.00	
Morning stiffness				
Significant or extreme	0.76 (0.41-1.42)	0.39	1.21 (0.64-2.30)	0.55
Moderate	1.37 (0.74-2.56)	0.32	2.03 (1.08-3.81)	<b>0.03</b>
A little or none (ref.)	1.00		1.00	
Sleep problems attributable to back pain				
Weekly	1.09 (0.63-1.89)	0.76	0.75 (0.41-1.35)	0.33
Less than weekly (ref.)	1.00		1.00	
Previous back pain frequency				
Yearly	1.11 (0.65-1.92)	0.70	1.00 (0.59-1.69)	0.99
Not yearly (ref.)	1.00		1.00	
<b>Block iv) Psychological variables</b>				
Fear-avoidance (FABQ-PA, 0-24)	1.02 (0.98-1.07)	0.32	1.03 (0.98-1.08)	0.22
Pain catastrophizing (PCS, 0-52)	1.04 (1.00-1.07)	0.05	1.06 (1.02-1.10)	<b>&lt;0.01</b>
Depression symptoms (CESD, 0-60)	0.99 (0.95-1.03)	0.53	0.99 (0.96-1.03)	0.61
Back beliefs (BBQ, 9-45)	0.97 (0.93-1.02)	0.23	0.99 (0.95-1.03)	0.67
Expectation for back pain in 3 months				
Recovered	0.26 (0.12-0.56)	<b>&lt;0.01</b>	0.39 (0.19-0.79)	<b>0.01</b>
Much better	0.65 (0.35-1.19)	0.16	0.85 (0.46-1.58)	0.61
No change or worse (ref.)	1.00		1.00	
Start Back Screening tool risk category				
Medium + high risk	1.02 (0.55-1.87)	0.95	0.49 (0.26-0.92)	<b>0.03</b>
Low risk (ref.)	1.00		1.00	
<b>Block v) Clinical variables</b>				
Number of red flags (0-12)	1.25 (0.99-1.58)	0.06	1.19 (0.96-1.48)	0.12
Diagnostic tool for radiculopathy				
Positive	1.94 (1.08-3.47)	<b>0.03</b>	1.52 (0.85-2.73)	0.16
Negative (ref.)	1.00		1.00	
Pain on active range of motion				
Yes	0.95 (0.57-1.58)	0.85	1.09 (0.67-1.80)	0.72
No (ref.)	1.00		1.00	
Trunk mobility performance (BPS, 0-18)	1.16 (1.08-1.24)	<b>&lt;0.01</b>	1.07 (1.00-1.15)	<b>0.04</b>
Timed Up and Go, mean seconds	0.93 (0.83-1.04)	0.20	1.00 (0.90-1.11)	0.93

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale - International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire - Physical Activity subscale; CES-D: Center for Epidemiological Studies - Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

\* The multinomial regression analyses are based on pooled estimates from multiple regression analyses

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.

The chiropractic group (n=195) was the reference dependent variable.

Models were built block-wise within the five blocks: i) sociodemographic ii) general health iii) current episode and back pain history iv) psychological and v) clinical. All variables were included simultaneously.



**SUPPLEMENTARY FILE**

Table of Contents

Sensitivity analyses S1, complete case analyses: ..... 2

Sensitivity analyses S2, physiotherapy patients:..... 5

Sensitivity analyses S3, only low back pain patients: ..... 7

Assessment of generalizability S4: ..... 9

For peer review only

## Sensitivity analyses S1, complete case analyses:

### Methods:

Multiple multinomial regression. One model per variable block. The total number of available cases per category will vary with number of missing for each block, and is thus shown for each block.

Additionally, bootstrapping was performed for n=1000 bootstrapping samples. The average bootstrapping odds ratios and their corresponding bias-corrected accelerated 95% confidence intervals (BCa 95% CI) are provided. Because of few observations in the Start Back Screening Tool high risk group, we chose to combine this group with the medium risk group.

### Results:

See Table S1 for details. No substantial changes in point estimates were detected in the multinomial regression analyses when comparing complete cases analyses to the pooled imputed estimates. There were, however, some changes in p-values. In the complete case analyses, age and being in the SBT medium risk group were not significantly associated with choosing a PT compared to a chiropractor. Further, in the complete case analyses, having more red flags were significantly associated with choosing a GP compared to a chiropractor. As can be seen from the bootstrapping procedure, odds ratios and BCa 95% CIs were stable for all variables, except for the SBT high risk group. Here, the BCa 95% CIs indicate that the odds ratios cannot be trusted for this specific variable.

**Table S1:** Complete case analyses of multiple multinomial regression analyses. Chiropractic group is the reference group.

<b>Block i) Sociodemographic factors. Chiropractor n=181</b>						
	GP (n=113) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Physio (n=108) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Age	1.03 (0.99, 1.08)	0.11	1.03 (0.99, 1.08)	1.03 (0.99, 1.07)	0.14	1.03 (0.98, 1.08)
Gender						
Female	1.33 (0.81, 2.17)	0.26	1.33 (0.80, 2.09)	1.40 (0.85, 2.33)	0.19	1.40 (0.82, 2.27)
Male (ref)	1.00		1.00	1.00		1.00
Marital status						
Married/cohabiting	0.66 (0.37, 1.19)	0.17	0.66 (0.36, 1.26)	0.92 (0.49, 1.72)	0.79	0.92 (0.48, 1.68)
Not married/cohabiting (ref)	1.00		1.00	1.00		1.00
Educational level						
Higher education	1.02 (0.63, 1.65)	0.95	1.02 (0.60, 1.70)	1.08 (0.66, 1.77)	0.77	1.08 (0.65, 1.79)
Lower education (ref)	1.00		1.00	1.00		1.00
Employment status						
Currently in paid work	0.96 (0.50, 1.86)	0.91	0.96 (0.50, 1.78)	0.53 (0.27, 1.03)	0.06	0.53 (0.26, 1.00)
No paid work (ref)	1.00		1.00	1.00		1.00
<b>Block ii) General health factors. Chiropractor n=155</b>						
	GP (n=92) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Physio (n=89) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Hazardous alcohol intake (AUDIT-C)						
Yes	1.23 (0.70, 2.15)	0.48	1.23 (0.67, 2.15)	1.67 (0.95, 2.92)	0.07	1.67 (0.96, 3.12)
No (ref)	1.00		1.00	1.00		1.00
Smoking						
Yes	1.37 (0.57, 3.26)	0.48	1.37 (0.44, 3.89)	0.63 (0.22, 1.76)	0.37	0.63 (0.16, 1.64)
Previously	1.47 (0.82, 2.66)	0.20	1.47 (0.82, 3.16)	1.43 (0.81, 2.54)	0.22	1.43 (0.80, 2.78)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

No (ref)	1.00		1.00	1.00		1.00
Health-related quality of life (SF-36, 0-100)						
Physical component	0.96 (0.92, 0.99)	<b>0.03</b>	0.96 (0.92, 0.99)	0.97 (0.94, 1.01)	0.96	0.97 (0.94, 1.01)
Mental component	0.95 (0.92, 0.98)	<b>0.002</b>	0.95 (0.92, 0.98)	1.00 (0.94, 1.07)	0.99	1.00 (0.97, 1.04)
BMI	0.99 (0.93, 1.06)	0.81	0.99 (0.92, 1.06)	1.00 (0.94, 1.07)	0.99	1.00 (0.92, 1.09)
Comorbidities (SCQ, 0-15)	1.02 (0.81, 1.29)	0.88	1.02 (0.76, 1.31)	1.12 (0.89, 1.41)	0.33	1.12 (0.90, 1.42)
Widespread pain						
Yes	0.16 (0.03, 0.79)	<b>0.03</b>	0.16 (0.07, 0.38)	0.50 (0.15, 1.67)	0.26	0.49 (0.10, 1.41)
No (ref)	1.00		1.00	1.00		1.00
Falls self-efficacy (FESI, 16-64)	1.00 (0.93, 1.05)	0.73	0.99 (0.92, 1.07)	0.99 (0.93, 1.06)	0.77	0.99 (0.92, 1.07)

**Block iii) Current episode and back pain history. Chiropractor n=134**

	GP (n=80)			Physio (n=92)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Back pain severity (NRS, 0-10)	1.06 (0.91, 1.22)	0.49	1.05 (0.90, 1.22)	0.94 (0.82, 1.08)	0.40	0.94 (0.82, 1.10)
Back-related disability (RMDQ, 0-24)	1.06 (0.99, 1.13)	0.12	1.06 (0.97, 1.14)	1.06 (0.99, 1.13)	0.11	1.06 (0.97, 1.15)
Duration						
Over 3 months	5.49 (2.34, 12.85)	<b>&lt;0.001</b>	5.49(1.93, 22.47)	9.00 (4.03, 20.13)	<b>&lt;0.001</b>	9.00 (3.53, 32.69)
6 weeks to 3 months	4.92 (1.92, 12.61)	<b>0.001</b>	4.92 (1.78, 36.27)	4.90 (1.91, 12.56)	<b>0.001</b>	4.90 (1.62, 18.99)
0-6 weeks (ref)	1.00		1.00	1.00		1.00
Morning stiffness						
Significant or extreme	1.02 (0.47, 2.24)	0.96	1.02 (0.54, 2.80)	1.23 (0.57, 2.67)	0.60	1.23 (0.56, 2.75)
Moderate	1.92 (0.88, 4.22)	0.10	1.93 (0.69, 5.55)	2.36 (1.09, 5.14)	<b>0.03</b>	2.36 (1.07, 5.75)
A little or none (ref)	1.00		1.00	1.00		1.00
Sleep problems attributable to back pain						
Weekly	0.82 (0.42, 1.62)	0.57	0.82 (0.33, 1.61)	0.77 (0.39, 1.52)	0.45	0.77 (0.38, 1.47)
Less than weekly (ref)	1.00		1.00	1.00		1.00
Previous back pain frequency						
Yearly	1.06 (0.57, 1.96)	0.57	1.06 (0.49, 2.14)	1.07 (0.58, 1.99)	0.82	1.07 (0.51, 2.17)
Not yearly (ref)	1.00		1.00	1.00		1.00

**Block iv) Psychological factors. Chiropractor n=155**

	GP (n=96)			Physio (n=94)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Fear-avoidance (FABQ-PA, 0-24)	1.03 (0.98, 1.09)	0.22	1.03 (0.98, 1.10)	1.03 (0.98, 1.08)	0.24	1.03 (0.98, 1.09)
Pain catastrophizing (PCS, 0-52)	1.02 (0.97, 1.06)	0.45	1.02 (0.97, 1.07)	1.05 (1.01, 1.10)	<b>0.02</b>	1.05 (1.01, 1.11)
Depression symptoms (CESD, 0-60)	1.00 (0.96, 1.04)	0.97	1.00 (0.95, 1.04)	0.99 (0.95, 1.04)	0.66	0.99 (0.94, 1.04)
Back beliefs (BBQ, 9-45)	0.97 (0.92, 1.02)	0.21	0.97 (0.92, 1.01)	0.99 (0.94, 1.04)	0.64	0.99 (0.94, 1.04)
Expectation for back pain in 3 months						
Recovered	0.24 (0.10, 0.54)	<b>0.01</b>	0.24 (0.09, 0.48)	0.43 (0.19, 0.95)	<b>0.04</b>	0.43 (0.19, 0.93)
Much better	0.57 (0.29, 1.12)	0.10	0.57 (0.28, 1.10)	0.83 (0.41, 1.68)	0.61	0.83 (0.40, 1.91)
No change or worse(ref)	1.00		1.00	1.00		1.00
Start Back Screening tool						
Medium+high risk	1.31 (0.68, 2.53)	0.42	1.31 (0.62, 2.83)	0.49 (0.24, 1.00)	0.05	0.49 (0.26, 0.86)
Low risk (ref)	1.00		1.00	1.00		1.00

**Block v) Clinical variables. Chiropractor n=159**

	GP (n=105)			Physio (n=110)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Number of red flags (0-12)	1.30 (1.02, 1.67)	<b>0.04</b>	1.30 (0.97, 1.75)	1.26 (0.99, 1.60)	0.06	1.26 (0.98, 1.59)
Nerve involvement diagnostic tool						
Positive	2.34 (1.27, 4.31)	<b>0.01</b>	2.34 (1.23, 4.69)	1.70 (0.93, 3.14)	0.09	1.70 (0.91, 3.33)
Negative (ref)	1.00		1.00	1.00		1.00
Pain on active range of motion						
Yes	0.76 (0.43, 1.32)	0.33	0.76 (0.44, 1.28)	0.92 (0.54, 1.58)	0.77	0.92 (0.53, 1.64)
No (ref)	1.00		1.00	1.00		1.00
Physical performance (BPS, 0-18)	1.19 (1.10, 1.29)	<b>&lt;0.001</b>	1.19 (1.11, 1.32)	1.10 (1.02, 1.19)	<b>0.01</b>	1.10 (1.02, 1.64)
Timed up and go, mean seconds	0.90 (0.79, 1.03)	0.14	0.90 (0.80, 1.02)	1.00 (0.88, 1.13)	0.94	1.00 (0.87, 1.13)

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire;

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.  
\*OR (BCa 95% CI) is average odds ratios from 1000 bootstrapping samples, including bias-corrected accelerated 95% confidence intervals.

For peer review only

## Sensitivity analyses S2, physiotherapy patients:

Analyses of differences between physiotherapy patients recruited before 01.01.2018 and after 01.01.2018. After 01.01.2018, there was direct access to physiotherapy in Norway, which potentially could change the population characteristics.

### Methods:

- Univariate analyses corresponding to measurement level and distribution: Chi square test or Fischer's exact test for categorical variables, individual sample t-test for normally distributed continuous variables, Mann Whitney U-test for continuous variables with a skewed distribution
- We used the pooled estimates from multiple imputation that were used in the article table 1 and 2

### Results:

See Table S2 for details. We found statistically significant differences between PT patients recruited before and after 01.01.2018 on the BBQ and BPS. PT patients recruited before 01.01.2018 held significantly more optimistic beliefs about back pain, with a mean (SD) BBQ score of 30.3 (6.8) for patients recruited before 01.01.2018 compared to 27.3 (7.5) for patients recruited after 01.01.2018 ( $p=0.03$ ). PT patients recruited before 01.01.2018 had significantly better trunk mobility performance, with a median (IQR) of 5 (2-7) for patients recruited before 01.01.2018 compared to 7 (4-9.75) for patients recruited after 01.01.2018 ( $p=0.003$ ).

**Table S2:** Univariate analyses of differences between physiotherapy patients recruited before and after 01.01.2018.

	Physio before (n=90)	Physio after (n=40)	p-value
Age, median (IQR)	68 (62.75, 73)	68.5 (61.5, 76)	0.323
Sex female, n (%)	53 (58.9)	17 (42.5)	0.084
Married or living with partner, n (%)	69 (76.7)	29 (72.5)	0.580
Paid work, n (%)	30 (33.3)	12 (30.0)	0.606
Education level			0.317
- Low (elementary+high school)	51 (56.7)	19 (47.5)	
- High (university+ uni 4+)	39 (43.3)	21 (52.5)	
Health-related quality of life			
- Mental sumscore, median (IQR)	56.29 (51.01, 60.99)	54.63 (47.35, 60.37)	0.396
- Physical sumscore, mean (SD)	40.61 (7.91)	40.67 (8.30)	0.969
Hazardous alcohol consumption, n (%)	44 (48.9)	21 (52.5)	0.786
Smoking status			0.202
- Current smoker	9 (10)	4 (10)	
- Previous	46 (51.1)	14 (35)	
- Never	35 (38.9)	22 (55)	
Number of comorbidities, median (IQR)	2 (1, 2.25)	1 (1, 2)	0.235
BMI, median (IQR)	26.60 (24.41, 30.47)	26.37 (24.60, 29.27)	0.913

1				
2				
3	Fall last 6 weeks, n (%)	18 (20)	7 (17.5)	0.623
4	Falls self-efficacy, median (IQR)	20 (18, 23.35)	22.5 (17, 26.9)	0.424
5	Widespread pain, n (%)	5 (5.6)	2 (5.0)	0.880
6	Previous back pain, n (%)			0.479
7	- Monthly	35 (38.9)	11 (27.5)	
8	- Every year	30 (33.3)	14 (35.0)	
9	- Every 1-5 years	13 (14.4)	5 (12.5)	
10	- Every five years	8 (8.9)	8 (20.0)	
11	- Only once	4 (4.4)	2 (5.0)	
12	Duration of current episode, n (%)			0.538
13	- 0-6 weeks	30 (33.3)	11 (27.5)	
14	- 6 weeks to 3 months	17 (18.9)	11 (27.5)	
15	- 3 months or over	43 (47.8)	19 (47.5)	
16	Back pain, mean (SD)	5.22 (2.53)	4.69 (1.87)	0.208
17	Back-related disability, RMDQ, median (IQR)	8 (6, 13)	9.5 (4.25, 14)	0.808
18	Sleep problems due to back pain, n (%)			0.374
19	- Weekly	36 (40)	13 (32.5)	
20	- Less than weekly	54 (60)	27 (67.5)	
21	Morning stiffness, n (%)			0.753
22	- Significant or extreme	35 (38.9)	16 (40)	
23	- Moderate	35 (38.9)	13 (32.5)	
24	- Some or none	20 (22.2)	11 (27.5)	
25	Walking distance, n (%)			0.285
26	- More than 3km	40 (44.4)	16 (40.0)	
27	- 200m to 3km	41 (45.6)	16 (40.0)	
28	- Less than 200m	9 (10)	8 (20.0)	
29	Kinesiophobia (FABQ-PA), median (IQR)	10 (5, 15)	10.5 (5, 14)	0.842
30	Depression (CES-D), median (IQR)	8 (3.75, 14)	9.5 (5.25, 17.3)	0.305
31	Pain catastrophizing (PCS), median (IQR)	12 (5.3, 17)	11 (4, 19.6)	0.872
32	Back beliefs (BBQ), mean (SD)	30.3 (6.8)	27.3 (7.5)	<b>0.03</b>
33	Expectations for back pain next 3 months			0.821
34	- Fully recovered	17 (18.9)	7 (17.5)	
35	- Much better	50 (55.5)	21 (52.5)	
36	- No change or worse	23 (25.5)	12 (30.0)	
37	SBT risk profiles			0.163
38	- Low	68 (75.5)	24 (60)	
39	- Medium	18 (20)	14 (35)	
40	- High	4 (4.4)	2 (5)	
41	Physical performance (BPS), median (range)	5 (2, 7)	7 (4, 9.75)	<b>0.003</b>
42	Timed up and go, median (IQR)	7.99 (6.66, 9.18)	7.42 (6.64, 9.86)	0.655
43	Probable nerve root involvement, n (%)	20 (22.2)	13 (32.5)	0.194
44	Number of red flags, median (range)	1 (0, 2)	1 (0, 2)	0.815
45	Pain on active range of motion, n (%)	61 (67.8)	27 (67.5)	0.905

IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
AUDIT-C scores of  $\geq 3/12$  for women and  $\geq 4/12$  indicates hazardous alcohol consumption

## Sensitivity analyses S3, only low back pain patients:

### Methods:

Multiple multinomial regression. One model per variable block. The chiropractic group is the reference group. For these analyses, 382 patients were available; 106 GP patients, 109 physiotherapy patients, and 167 chiropractic patients.

### Results:

See Table S3 for details. Overall, there were very few substantial changes in point estimates and p-values compared to the analyses of all included patients in the article main body. SF-36 physical component summary score was no longer significantly associated with first visiting a GP. Having widespread pain was significantly associated with visiting a physiotherapist compared to a chiropractor. Although point estimates for back-related disability was identical, it was no longer significantly associated with visiting a GP or a physiotherapist. For the Start Back Screening Tool, medium risk category was no longer significantly associated with visiting a chiropractor compared to a physiotherapist, but high risk was significant. Having a positive diagnostic rule for radiculopathy was significantly associated with visiting a physiotherapist compared to a chiropractor.

**Table S3:** Subgroup analyses of the multinomial regression analyses for patients with low back pain only. Chiropractic group (n=167) is the reference group.

<b>Block i) Sociodemographic factors.</b>				
	GP (n=106)		Physio (n=109)	
	Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-value
Age	1.03 (0.99, 1.07)	0.23	1.04 (1.00, 1.08)	<b>0.05</b>
Gender				
Female	1.43 (0.86, 2.37)	0.17	1.31 (0.78, 2.19)	0.31
Male (ref)	1.00		1.00	
Marital status				
Married/cohabiting	0.58 (0.30, 1.09)	0.09	0.73 (0.38, 1.40)	0.34
Not married/cohabiting (ref)	1.00		1.00	
Educational level				
Higher education	0.97 (0.58, 1.61)	0.91	1.18 (0.71, 1.96)	0.52
Lower education (ref)	1.00		1.00	
Employment status				
Currently in paid work	0.79 (0.40, 1.55)	0.49	0.63 (0.31, 1.28)	0.20
No paid work (ref)	1.00			
<b>Block ii) General health factors.</b>				
Hazardous alcohol intake (AUDIT-C)				
Yes	1.19 (0.69, 2.05)	0.54	1.18 (0.69, 2.01)	0.54
No (ref)	1.00		1.00	
Smoking				
Yes	1.42 (0.64, 3.19)	0.39	0.64 (0.24, 1.71)	0.37
Previously	1.37 (0.75, 2.47)	0.30	1.02 (0.59, 1.77)	0.95
No (ref)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.97 (0.93, 1.00)	0.08	0.98 (0.94, 1.01)	0.20
Mental component	0.97 (0.94, 1.00)	<b>0.04</b>	1.00 (0.97, 1.03)	0.96
BMI	0.99 (0.93, 1.05)	0.76	0.96 (0.90, 1.02)	0.23
Comorbidities (SCQ, 0-15)	1.13 (0.90, 1.42)	0.29	1.18 (0.96, 1.47)	0.12
Widespread pain				
Yes	0.16 (0.04, 0.65)	<b>0.01</b>	0.30 (0.09, 0.99)	<b>0.05</b>
No (ref)	1.00		1.00	
Falls self-efficacy (FESI, 16-64)	1.00 (0.95, 1.05)	0.99	1.01 (0.95, 1.06)	0.85

<b>Block iii) Current episode and back pain history.</b>					
Back pain severity (NRS, 0-10)	0.98 (0.86, 1.11)	0.73	0.89 (0.78, 1.01)	0.07	
Back-related disability (RMDQ, 0-24)	1.06 (1.00, 1.13)	0.05	1.06 (1.00, 1.13)	0.06	
Duration					
Over 3 months	3.54 (1.42, 8.80)	<b>&lt;0.01</b>	3.85 (1.69, 8.77)	<b>&lt;0.01</b>	
6 weeks to 3 months	3.40 (1.12, 10.37)	<b>0.03</b>	3.25 (1.16, 9.09)	<b>0.03</b>	
0-6 weeks (ref)	1.00		1.00		
Morning stiffness					
Significant or extreme	0.79 (0.39, 1.60)	0.51	1.35 (0.68, 2.67)	0.39	
Moderate	1.63 (0.82, 3.24)	0.16	2.02 (1.02, 4.03)	<b>0.05</b>	
A little or none (ref)	1.00		1.00		
Sleep problems attributable to back pain					
- Weekly	1.13 (0.60, 2.14)	0.70	0.66 (0.34, 1.26)	0.20	
- Less than weekly (ref)	1.00		1.00		
Previous back pain frequency					
- Yearly	1.03 (0.57, 1.87)	0.93	1.04 (0.59, 1.83)	0.88	
- Not yearly (ref)	1.00		1.00		
<b>Block iv) Psychological factors.</b>					
Fear-avoidance (FABQ-PA, 0-24)	1.00 (0.95, 1.05)	0.97	1.03 (0.98, 1.08)	0.31	
Pain catastrophizing (PCS, 0-52)	1.03 (0.99, 1.07)	0.20	1.06 (1.02, 1.10)	<b>&lt;0.01</b>	
Depression symptoms (CESD, 0-60)	0.97 (0.94, 1.03)	0.50	0.99 (0.95, 1.03)	0.70	
Back beliefs (BBQ, 9-45)	0.96 (0.92, 1.01)	0.12	0.99 (0.94, 1.04)	0.63	
Expectation for back pain in 3 months					
Recovered	0.21 (0.09, 0.49)	<b>&lt;0.01</b>	0.34 (0.16, 0.73)	<b>&lt;0.01</b>	
Much better	0.60 (0.31, 1.16)	0.13	0.71 (0.36, 1.39)	0.31	
No change or worse(ref)	1.00		1.00		
Start Back Screening tool					
High risk	1.82 (0.55, 6.05)	0.33	0.19 (0.04, 0.90)	<b>0.04</b>	
Medium risk	1.03 (0.52, 2.06)	0.92	0.59 (0.30, 1.17)	0.13	
Low risk (ref)	1.00		1.00		
<b>Block v) Clinical variables.</b>					
Number of red flags (0-12)	1.28 (0.98, 1.68)	0.07	1.16 (0.90, 1.50)	0.24	
Diagnostic rule for radiculopathy					
Positive	2.32 (1.24, 4.34)	<b>&lt;0.01</b>	1.89 (1.00, 3.57)	<b>0.05</b>	
Negative (ref)	1.00		1.00		
Pain on active range of motion					
Yes	0.88 (0.50, 1.53)	0.64	1.06 (0.62, 1.80)	0.84	
No (ref)	1.00				
Physical performance (BPS, 0-18)	1.19 (1.10, 1.28)	<b>0.03</b>	1.09 (1.01, 1.17)	<b>0.03</b>	
Timed up and go, mean seconds	0.89 (0.78, 1.01)	0.06	0.97 (0.86, 1.09)	0.56	

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale - International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire - Physical Activity subscale; CES-D: Center for Epidemiological Studies - Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.



## Assessment of generalizability S4:

**Table S4:** Descriptive comparison of NORLAG sample and NORLAG 2017 musculoskeletal (MSK) subsample with BACE-N sample.

	NORLAG 2017 subsample MSK conditions <sup>^</sup> (n=794)	BACE-N (n=452)
Age, median (IQR, range)	66 (60-74, 50-93)	66 (59-72, 55-89)
Gender female, n (%)	506 (63.7)	235 (52)
Mother tongue Norwegian (n=432), n (valid %)		412 (95.4)
Country of origin Norway, n (%)	728 (91.7)	
Educational level, n (%)		
- Low (elementary + high school)	566 (71.4)	253 (56.0)
- High (university level)	227 (28.6)	199 (44.0)
In paid work, n (%)	251 (31.6)	205 (45.3)
Living with partner, n (%)	494 (62.2)	347 (76.8)
BMI, mean (SD)	26.3 (4.4)	27.6 (4.7)
How many alcoholic units do you normally drink? <sup>~</sup> n (valid %)		
- 1-2		
- 3-4		
- 5-6	183 (70.1)	289 (63.9)
- 7-9	62 (23.8)	136 (30.1)
- 10 or more	10 (3.8)	22 (4.8)
	1 (0.4)	2 (0.4)
	5 (1.9)	3 (0.7)
How often have you drunk alcohol until you felt intoxicated? (n=433) n, (valid %)		
- Once per week	12 (2.8)	
- 2-3 times per week	3 (0.7)	
- 2-3 times per month	18 (4.2)	
- Once per month	37 (8.5)	
- Rarely	235 (54.3)	
- Never	128 (29.6)	
How often do you drink 6 alcoholic units or more?		
- Almost daily		1 (0.2)
- Some days per week		3 (0.7)
- Some days per month		41 (9.1)
- Rarely		194 (42.9)
- Never		213 (47.1)
CES-D (IQR, range)	8 (4-14, 0-38)	8 (4-15, 0-46)
HR-QoL, physical summary score*, mean (SD)	37.5 (11.3)	41.4 (8.4)
HR-QoL, mental summary score*, mean (SD)	54.7 (8.2)	52.5 (10.0)
Walking distance		
- Cannot walk	13 (1.7)	
- A few steps	22 (2.8)	
- 10-100 m	59 (7.6)	
- 100-500m	57 (7.3)	
- 500m-1km	82 (10.5)	
- 1-5km	235 (30.1)	
- 5km+	313 (40.1)	
Walking distance		
- Less than 15m		20 (0.7)
- 15m-200m		310 (11.5)
- 200m-3km		1130 (42.1)
- 3km+		1218 (45.3)

IQR: Interquartile range; SD; Standard deviation; BMI: Body mass index; CES-D: Center for Epidemiological Studies – Depression questionnaire; HR-QoL: Health-related quality of life

<sup>^</sup>The subsample was collected in 2017 and consisted of participants aged 55 years or older, with at least one musculoskeletal condition

<sup>~</sup> In NORLAG, this variable is continuously, as “number of alcoholic drinks usually drunk per time you drink alcohol”. In BACE-N, it is the AUDIT-C question 2, a categorical question with 5 categories: 1-2, 3-4, 5-6, 7-9 and 10 or more.

\*NORLAG used Short Form Health Survey-12, BACE-N used Short Form Health Survey-36

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	4
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	5
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	5
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5-7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5-7
Bias	9	Describe any efforts to address potential sources of bias	8
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7
		(b) Describe any methods used to examine subgroups and interactions	NA
		(c) Explain how missing data were addressed	7
		(d) If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	8
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9
		(b) Give reasons for non-participation at each stage	-
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	9, 19-20
		(b) Indicate number of participants with missing data for each variable of interest	9, 19-20
Outcome data	15*	Report numbers of outcome events or summary measures	9
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9-10, 20-21

		(b) Report category boundaries when continuous variables were categorized	6-7
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10, suppl.
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	10
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	10-12
Generalisability	21	Discuss the generalisability (external validity) of the study results	12
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Characteristics of older adults with back pain associated with choice of first primary care provider: a cross-sectional analysis from the BACE-N cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-053229.R2
Article Type:	Original research
Date Submitted by the Author:	01-Sep-2021
Complete List of Authors:	Vigdal, Ørjan; Oslo Metropolitan University, Department of Physiotherapy Storheim, Kjersti; Oslo University Hospital; Oslo Metropolitan University, Department of Physiotherapy Munk Killingmo, Rikke ; Oslo Metropolitan University Faculty of Health Sciences, Department of Physiotherapy Småstuen, Milada; Oslo Metropolitan University, Department of Physiotherapy Grotle, Margreth; Oslo Metropolitan University, Department of Physiotherapy; Oslo universitetssykehus Ullevål, Research and Communication Unit for Musculoskeletal Health
<b>Primary Subject Heading</b>:	Health services research
Secondary Subject Heading:	Rehabilitation medicine, Geriatric medicine, General practice / Family practice
Keywords:	Back pain < ORTHOPAEDIC & TRAUMA SURGERY, PRIMARY CARE, GERIATRIC MEDICINE, REHABILITATION MEDICINE

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Article title:**

Characteristics of older adults with back pain associated with choice of first primary care provider: a cross-sectional analysis from the BACE-N cohort study

**Authors:**

Ørjan Nesse Vigdal<sup>1</sup>

Kjersti Storheim<sup>2,1</sup>

Rikke Munk Killingmo<sup>1</sup>

Milada C. Småstuen<sup>1</sup>

Margreth Grotle<sup>1,2</sup>

**Author affiliations:**

<sup>1</sup>Department of Physiotherapy, Faculty of Health Science, OsloMet - Oslo Metropolitan University, Oslo, Norway

<sup>2</sup>Research and Communication Unit for Musculoskeletal Health (FORMI), Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway

**Corresponding author:**

Ørjan Nesse Vigdal

E-mail address: orvig@oslomet.no

Mailing address: OsloMet – Oslo Metropolitan University, Post box 4, St.Olavs plass, 0130 Oslo, Norway

Telephone number: +47 414 59 314

**Manuscript number of pages (excluding title page, including tables): 21**

**Word count: 4320**

**Total number of tables: 2**

**Supplementary files: 1**

## Abstract

**Objectives:** To describe characteristics of older adults with back pain in primary care, and to assess associations between patient characteristics and type of first primary care provider (general practitioner (GP), physiotherapist (PT) or chiropractor).

**Design:** Cross-sectional analysis from the BACE-N cohort study.

**Setting:** Norwegian GP, PT and chiropractic primary care centres.

**Participants:** Patients aged  $\geq 55$  years seeking Norwegian primary care with a new episode of back pain were invited to participate. Between April 2015 and February 2020, we included 452 patients: 127 first visited a GP, 130 first visited a PT and 195 first visited a chiropractor.

**Primary and secondary outcome measures:** For the first objective, the outcome measure was descriptive statistics of patient characteristics, covering the following domains: sociodemographic, general health, current and previous back pain, psychological and clinical factors. For the second objective, first primary care provider was the outcome measure. Associations between patient characteristics and visiting a GP or PT compared to a chiropractor were assessed with multiple multinomial regression analyses.

**Results:** Median (IQR) age was 66 (59-72) years. Levels of back-related disability was moderate to severe, with a median (IQR) Roland-Morris Disability Questionnaire (range 0-24) score of 9 (5-13). Recurring episodes were common, 301 (67%) patients had monthly or yearly recurrences. Patients with worse back-related disability, longer duration of symptoms, lower expectations for full recovery and worse physical performance measured with the Back Performance Scale had higher odds of visiting a GP or PT compared to a chiropractor ( $p < 0.05$ ).

**Conclusion:** Older back pain patients in primary care had moderate to severe levels of back-related disability, and most had recurring episodes. Our results suggest that older adult's choice of first primary care provider was associated with important patient characteristics, which highlights the need for caution with generalizations of study results across primary care populations.

**Trial registration number:** ClinicalTrials identifier: NCT04261309

## Data availability statement

Data not available.

1  
2  
3 **Keywords:** Back pain, older adults, primary care, characteristics, care-seeking behaviour  
4  
5  
6

7 **Article summary**

8 **Strengths and limitations of this study**

- 9  
10  
11 • We used descriptive statistics to provide a thorough presentation of characteristics of older  
12 people seeking primary care for a new episode of back pain.  
13 • This study utilized multivariate, multinomial regression analyses to provide a comprehensive  
14 overview of associations between patient characteristics and choice of first healthcare provider.  
15 • It was not possible to obtain data on eligible patients that were not invited or declined to  
16 participate in the study, which might reduce external validity.  
17 • Due to differences in primary care organization between countries, readers are advised to  
18 exercise caution with generalizations of results to other healthcare systems.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Introduction

Back pain is the number one cause of years lived with disability globally, with an estimated point prevalence of 11.9% [1, 2]. Older adults have historically been under-represented in back pain research [3, 4], but have recently received increased attention [5, 6]. Although the prevalence of pathoanatomical findings on diagnostic imaging increases with age [7-9], the prevalence of serious pathology, such as vertebral fractures and neuropathic pain in older back pain patients in primary care is low. Studies have reported a prevalence of 6% and 2-11%, respectively [10, 11]. Moreover, studies in primary care have found significant national differences in the characteristics and burden of back pain in older adults [12, 13]. This highlights the importance of caution when generalizing results from studies from one setting to another.

Most patients seeking healthcare for back pain are treated in primary care [14]. In Norway, back pain is the reason for 10%, 27% and 86% of the visits to general practitioners (GP), physiotherapists (PT) and chiropractors, respectively [15]. Some studies suggest that choice of first primary care provider has consequences for future healthcare consumption, including imaging and opioid use [16, 17]. To optimize decision making regarding treatment, research and health policies, detailed knowledge of patient populations is required. Most of the previous studies exploring patient populations seeking primary care have compared GP and chiropractic populations, showing that patients seeking care from a GP have a higher overall burden of back pain compared to chiropractic patients [18-25]. Only a few studies include PT populations [26-29]. These studies suggest that patients seeking care from PTs are older and have more disability than those seeking care from chiropractors [26, 27, 29]. To the best of our knowledge, only one study has been performed in an exclusively older population [28]. This study found that older women seeking care from GPs reported worse back pain and worse health-related quality of life than older women visiting a PT or a chiropractor [28]. The study only included women between 59-64 years of age, and it is not clear if the results are also generalizable to men or adults over 65 years of age. Further, they did not examine back-related disability or other back pain factors, sociodemographic factors, psychological factors or clinical factors. Thus, there is still a considerable lack of knowledge regarding whether characteristics of older back pain patients differ according to their choice of first primary care provider.

Therefore, the aims of this study were 1) to describe the characteristics of patients  $\geq 55$  years of age seeking primary care for a new episode of back pain in terms of sociodemographic, general health, current back pain and back pain history, psychological and clinical characteristics, and 2) to assess if patient characteristics are associated with type of first primary care provider (GP, PT or chiropractor).

## Methods

### Design and setting

This cross-sectional study presents baseline data from the Back Complaints in the Elders – Norway (BACE-N) study, a prospective observational cohort study in Norwegian primary care. The BACE-N study is a part of the international BACE consortium, with research groups from Brazil, the Netherlands and Australia [6]. The BACE-N study protocol has been registered in ClinicalTrials.gov (Identifier NCT04261309). The study was classified as a quality assessment study by the Norwegian Regional Committee for Medical Research Ethics (reference no. 2014/1634/REK vest) and was approved by the Norwegian Social Science Data Service in 2015 (reference no. 42149).

Norwegian primary care is organized by the municipalities and financed through the National Insurance Scheme, the municipalities, and patient co-payment [30]. There is direct access to GPs, PTs (from 2018) and chiropractors [30]. Patient co-payment rates vary between healthcare providers, with chiropractors generally having the highest co-payment cost [30]. Treatments provided usually differ between the healthcare providers. For example, patients visiting a GP are more likely to receive pharmacological therapy, patients visiting a PT are more likely to receive exercise therapy, and patients visiting chiropractors are more likely to receive manipulation therapy [15].

### Participants and recruitment procedure

Eligible patients were  $\geq 55$  years of age, seeking primary care from a GP, PT or chiropractor in primary care for a new episode of back pain. Back pain was defined as pain located in the region from the top of the scapula to the sacrum, with or without radiating leg pain. A new episode was defined as not having received healthcare for the same complaint in the last six months. Patients were excluded if they had difficulties completing the questionnaire due to language barriers, or if they had difficulties completing the clinical examination (for example wheelchair-bound patients). Participants received care as usual.

Patients were recruited from GPs, PTs, and chiropractors in urban and rural parts of Norway between April 2015 and February 2020, either during or immediately after the consultation. The primary care providers were instructed to invite consecutive patients. To facilitate the recruitment process, media advertisements were also used. Eligible patients received oral and written information about the study. The final screening for eligibility and inclusion to the study was performed by the researchers. All included patients signed an informed consent form before enrolment in the study. The baseline

1  
2  
3 measurements, consisting of questionnaires and a clinical examination, were collected as soon after  
4 the first primary care consultation as possible.  
5  
6  
7

## 8 **Measurements**

### 9 *Sociodemographic variables*

10 Information regarding age, sex, marital status, employment status and educational level were  
11 collected.  
12  
13  
14

### 15 *General health variables*

16 Health-related quality of life (HR-QoL) was measured using the Short-Form Health Survey 36-item  
17 (SF-36) physical and mental summary measures (standardized with a mean of 50 and a standard  
18 deviation of 10 according to a general US population with higher scores denoting better health) [31].  
19 Alcohol consumption was measured using the 3-item Alcohol Use Disorder Identification Test  
20 consumption questions (AUDIT-C) (range 0-12, higher score indicates higher alcohol consumption)  
21 [32]. Hazardous alcohol consumption was defined as an AUDIT-C score of  $\geq 3/12$  for women and  
22  $\geq 4/12$  for men [33, 34]. Smoking status (current smoker, previous smoker, non-smoker) was  
23 collected. The number of comorbidities was measured using the Self-Administered Comorbidity  
24 Questionnaire (SCQ) [35]. The SCQ has 13 pre-defined comorbidities and two optional comorbidities.  
25 Item 12, "back pain", was replaced with a third optional comorbidity. Widespread pain was  
26 measured using the pain drawing from McGill Pain Questionnaire and the revised criteria from Wolfe  
27 et al. for widespread pain [36, 37]. The number of falls during the last six weeks was collected, and  
28 falls self-efficacy was measured using the Falls Efficacy Scale-International (FES-I) (range 16-64,  
29 higher score indicated lower falls efficacy) [38].  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 *Current back pain and back pain history*

44 Back pain location (thoracic or lumbar, or both) was collected. Average back pain severity last week  
45 was measured using the Numeric Rating Scale (NRS) (range 0-10, higher score indicates higher back  
46 pain severity) [39]. Back-related disability was measured with the 24-item Roland-Morris Disability  
47 Questionnaire (RMDQ) (range 0-24, higher score indicated more back-related disability) [40]. Back  
48 pain duration was measured in days and categorized into "<6 weeks", "6 weeks to 3 months", and  
49 ">3 months". Frequency of previous back pain episodes (monthly, yearly, every 1-5 years, every five  
50 years, once) was collected. Sleep problems attributable to back pain were measured using item 5i  
51 from the Pittsburgh Sleep Quality Index (PSQI) [41], and dichotomized to "weekly/less than weekly".  
52 Morning stiffness was measured with item six from Knee injury and Osteoarthritis Outcome Score  
53 (KOOS) [42], where we replaced the word "knee" with "back".  
54  
55  
56  
57  
58  
59  
60

### *Psychological variables*

Kinesiophobia was measured using the Fear-Avoidance Beliefs Questionnaire-Physical Activity subscale (FABQ-PA) (range 0-24, higher score indicates higher levels of kinesiophobia) [43]. Signs of depression were measured with the Center for Epidemiological Studies-Depression questionnaire (CES-D) (range 0-60, higher score indicates more signs of depression) [44]. Pain catastrophizing was measured using the Pain Catastrophizing Scale (PCS) (range 0-52, higher score indicates more pain catastrophizing) [45]. Beliefs and attitudes towards back pain was measured using the Back Beliefs Questionnaire (BBQ) (range 9-45, higher score indicates more positive beliefs) [46]. Start Back Screening Tool (SBT) was used to assess prognostic risk profiles [47]. Expectations of recovery from back pain within the next 3 months was assessed with a five-point scale, with the categories “Fully recovered”, “Much better”, “No difference”, “Much worse”, and “Worse than ever”.

### *Clinical variables*

Pain with active movements was assessed for forward flexion, lateral flexion and rotation of the back. Physical performance with focus on trunk mobility was assessed with the 6-item Back Performance Scale (BPS) (range 0-18, higher score indicates worse trunk mobility performance) [48]. Walking function was assessed with the Timed-Up-and-Go (TUG) [49]. Signs of radiculopathy was measured using a clinical diagnostic model that summarizes five items: Subjective sensory changes (1 point), radiating pain below the knee (2 points), leg pain worse than back pain (2 points), positive neural tension test (3 points) and neurological deficit of myotome, dermatome or reflexes in the lower limb (2 points) [50]. A score of  $\geq 5/10$  has been shown to indicate  $>80\%$  probability of radiculopathy [50]. Twelve red flags were assessed: Cancer, first episode of back pain, constant pain, unexplained weight loss, systemically unwell, fever, urinary retention or loss of bladder control, age  $\geq 75$  years, trauma cause of back pain, osteoporosis, cortisone use and severe morning stiffness.

### **Statistical analyses**

All analyses were performed using the IBM SPSS Statistics version 26 for Windows (IBM Corporation, Armonk, NY, USA). To handle missing data, five multiple imputation datasets with 10 iterations were created using regression estimation, and the pooled estimates are presented in this study. Patient characteristics were described with counts and percentages for categorical variables, mean and standard deviation (SD) for normally distributed continuous variables and median and interquartile range (IQR) for continuous variables with a skewed distribution. Mann-Whitney U-test was used to assess differences in days between first primary care contact and inclusion to the study between primary care practitioners, and between those recruited from primary care and those recruited from

1  
2  
3 media advertisements. Multinomial regression was used to assess the strength of the associations  
4 between patient characteristics and patient's choice of first primary care provider. First primary care  
5 provider (GP, PT or chiropractor) was the dependent variable. The chiropractic group was the largest,  
6 and therefore chosen as the reference group. Patient characteristics were organized into five blocks,  
7 for which we created separate models: i) Sociodemographic ii) general health iii) current back pain  
8 episode and back pain history iv) psychological variables and v) clinical variables. All variables in the  
9 block were simultaneously included in the model, without univariate pre-testing. The strength of  
10 associations is expressed as odds ratios (OR) with 95% confidence intervals (CI). We considered our  
11 study as exploratory, so no correction for multiple testing was performed [51]. P-values <0.05 were  
12 thus considered statistically significant. All tests were two-sided.  
13  
14  
15  
16  
17  
18  
19  
20  
21

### 22 *Assessment of generalizability*

23 Because of economic and practical reasons, we were unable to collect data on eligible participants  
24 that declined to participate or for other reasons were not invited. Therefore, we performed a  
25 descriptive comparison of the BACE-N on age, sex, nationality, educational level, work status, marital  
26 status, BMI, alcohol use, HR-QoL, depression and walking distance with individual data from a  
27 subsample from the study "The Norwegian study on life course, ageing and generation (NORLAG)"  
28 [52, 53]. This study used a random sampling strategy in the general population and included 11028  
29 participants. The subsample (NORLAG MSK) consisted of 794 participants collected in 2017. The  
30 participants of the subsample were  $\geq 55$  years of age and had at least one musculoskeletal complaint.  
31  
32  
33  
34  
35  
36  
37

### 38 *Sensitivity analyses*

39 We performed three sensitivity analyses: 1) To assess possible bias introduced by the multiple  
40 imputation procedure, the multiple multinomial regression analyses were performed on complete  
41 case data. We included a bootstrapping approach to assess the robustness of the coefficients. 2)  
42 Because PT services became available through direct access in Norway from 01.01.2018,  
43 characteristics of PT patients recruited before and after 01.01.2018 were compared using individual  
44 sample t-tests or Mann-Whitney U-tests for continuous variables, and chi-square tests for categorical  
45 variables. 3) We performed the multiple multinomial regression analyses in the subgroup with low  
46 back pain only. Results from the sensitivity analyses are available in supplementary material S1  
47 through S3.  
48  
49  
50  
51  
52  
53  
54  
55  
56

### 57 *Sample size consideration*

58 Sample size was considered for the BACE-N study as a whole, with the following criteria: Having  
59 sufficient statistical power for up to 14 variables in a multivariate logistic regression analysis using  
60

1  
2  
3 the “10 events per variable” rule [54], with an outcome prevalence of 40%, and allowing for a  
4 dropout-rate of 20%. This yielded a preferred sample size of 450 participants. As the multinomial  
5 regression models in this study includes a maximum of 8 independent variables, we expect the  
6 sample size to be sufficient.  
7  
8  
9

### 10 11 **Patient and public involvement**

12 Patient representatives were part of the scientific board of the study and involved in designing and  
13 establishing BACE-N. Results will be disseminated to the recruiting primary care providers and the  
14 participating patients in an annual newsletter.  
15  
16  
17  
18  
19  
20  
21

### 22 **Results**

23 A total of 452 patients were included in the study, 127 first visited a GP, 130 first visited a PT and 195  
24 first visited a chiropractor. Eighteen patients were included from media advertisements. Median  
25 (IQR) number of days from first primary care contact to inclusion in the study was 13 (3-21) days for  
26 GP patients, 9 (3-21) for physiotherapy patients and 5 (1-13) for chiropractic patients. The duration  
27 was significantly shorter for chiropractic patients compared to GP patients ( $p<0.01$ ) and PT patients  
28 ( $p<0.01$ ). There was no statistically significant difference in duration from first primary care contact  
29 to inclusion between those recruited directly from primary care practices (median (IQR) 7 (2-15)  
30 days), and those recruited through media advertisements (median (IQR) 16 (1-28) days) ( $p=0.315$ ).  
31  
32  
33  
34  
35  
36  
37  
38

#### 39 *Patient characteristics*

40 Missingness ranged from 0.0 to 16.8% for the variables, and total missingness was 4.4% across all  
41 values. Rates of missingness was similarly distributed across the primary care provider groups.  
42 Consult table 1 for details regarding patient characteristics. The median age of the patients was 66,  
43 around half of the patients were women, were in paid work, and had university-level education. Half  
44 of the patients had a hazardous alcohol consumption level, and nearly 60% of them were either  
45 current or previous smokers. One in six patients had experienced a fall during the last six weeks. Half  
46 of the patients had one or more comorbidities.  
47  
48  
49  
50  
51  
52

53 Most patients reported moderate levels of back pain and moderate to severe levels of back-related  
54 disability with a median (IQR) RMDQ-score of 9 (5-13). Almost 70% of the patients experienced  
55 monthly or yearly recurrences of back pain. Over 40% experienced weekly sleep problems  
56 attributable to back pain, and 70% experienced moderate to extreme morning stiffness. Two thirds  
57 of the patients had a low-risk profile according to the SBT, and only 6.6% had a high-risk profile.  
58  
59  
60

1  
2  
3 Expectations of recovery were generally high, with three out of four expecting to be much better or  
4 fully recovered within three months.  
5  
6  
7

#### 8 *Associations between patient characteristics and type of first primary care provider*

9

10 Table 2 presents the associations from the multinomial regression analyses. Patients with higher  
11 back-related disability, longer duration of symptoms, worse physical performance, probable  
12 radiculopathy, poorer HR-QoL and lower expectations of being fully recovered within the next three  
13 months were more likely to visit a GP compared to a chiropractor. Patients with widespread pain  
14 were more likely to visit a chiropractor than a GP. The characteristics strongest associated with  
15 choosing a GP versus a chiropractor were duration of symptoms, widespread pain and expectation of  
16 being fully recovered.  
17  
18  
19  
20  
21  
22

23 Patients that were older, had a longer duration of symptoms, higher back-related disability,  
24 moderate morning stiffness, higher levels of pain catastrophizing, worse physical performance, lower  
25 expectations of being fully recovered within the next three months were more likely to visit a PT  
26 compared to a chiropractor. Patients in the SBT medium or high risk group were more likely to visit a  
27 chiropractor compared to a PT. The characteristics strongest associated with choosing a PT versus a  
28 chiropractor were duration of symptoms and expectation of being fully recovered.  
29  
30  
31  
32  
33  
34

35 Gender, education level, marital status, employment status, comorbidities, back pain severity, sleep  
36 problems, kinesiophobia, depressive signs, back beliefs, red flags, pain on active range of motion and  
37 Timed Up and Go-scores were not associated with type of primary care provider.  
38  
39  
40  
41

#### 42 *Assessment of generalizability*

43

44 The BACE-N study sample had more men (48% versus 36.3% in NORLAG MSK), more participants with  
45 high educational level (44% versus 28.6% in NORLAG MSK), more participants currently in paid work  
46 (45.3% versus 31.6% in NORLAG MSK), and more participants living with a partner (76.8% versus  
47 62.2% in NORLAG MSK). Age, nationality, alcohol consumption, BMI, depressive signs, HR-QoL and  
48 walking distance were similar for BACE-N and NORLAG MSK. See supplementary material S4 for  
49 further details.  
50  
51  
52  
53  
54  
55  
56

## 57 **Discussion**

58 This study showed that nearly all older patients with back pain had experienced back pain previously,  
59 and for most patients this episode was the latest of a series of annually or monthly recurring  
60

1  
2  
3 episodes. This is in accordance with several studies on back pain trajectories, where episodic or  
4 fluctuating pain was shown to be common both in the short and long term [55-58]. Further, patients  
5 with more severe back-related disability and other symptoms and signs were overall more likely to  
6 visit a GP or a physiotherapist than a chiropractor. Contrary to this finding, patients with widespread  
7 pain were more likely to choose a chiropractor over a GP. This is the first study to assess associations  
8 of a broad range of patient characteristics and choice of first primary care provider in an older  
9 population. Older adults have previously been under-represented in back pain studies [3, 4], and the  
10 evidence underlying treatment decisions in this age group may have been over-reliant on studies  
11 performed in younger populations. Thus, this study provides evidence to improve knowledge about  
12 older adults with back pain. This may prove important for clinical guideline development and  
13 informing stakeholders aiming to improve quality of care for older adults with back pain.  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 The burden of back pain and psychological profile were comparable between younger Norwegian  
24 back pain cohorts and the older BACE-N sample [59, 60]. The characteristics of the included patients  
25 in this study was largely comparable to the BACE-study from the Netherlands [12, 61], with a few  
26 exceptions. Both in our total study sample and our GP subsample, a larger proportion of patients had  
27 paid work, fewer experienced their first episode of back pain, and they reported lower levels of  
28 kinesiophobia and pain catastrophizing compared to the Dutch study sample. When comparing our  
29 results to the Brazilian BACE-study [12, 62], the Brazilian study had a higher proportion of women.  
30 Further, our study sample had more patients in paid work, more patients with hazardous alcohol  
31 consumption patterns, more smokers, the patients had fewer comorbidities, lower levels of back-  
32 related disability and back pain severity, kinesiophobia, depression signs, and pain catastrophizing  
33 compared to the Brazilian BACE-sample. These differences between populations within the BACE  
34 consortium might be explained in part by minor differences in recruitment strategies in the different  
35 countries [12] or differences in how primary care is organized in the different countries. In the  
36 Netherlands, patients were recruited exclusively from a GP setting [61], whereas in Brazil patients  
37 were recruited from primary care centres or health centres specialized in geriatrics [62]. Another  
38 possible explanation may be cultural differences in the expression and interpretation of and coping  
39 with pain [63].  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 In line with previous research on healthcare utilization for back pain in younger populations [19, 21-  
54 23, 25-28], our results suggest that patients with “less complex” characteristics were more likely to  
55 visit a chiropractor compared to a GP or a physiotherapist. Unsurprisingly, studies using bivariate  
56 analyses [18, 20, 23, 25, 28, 29] to compare the provider groups find more significant associations or  
57 differences than studies using multivariate analyses [19, 21, 22, 26, 27]. However, regardless of  
58  
59  
60



1  
2  
3 statistical approach, these studies suggest that patients who seek chiropractic care have an overall  
4 lower burden of back pain compared to patients seeking GP or PT care [18-23, 25]. One notable  
5 exception is the study of Eklund et al. [24], which found that Swedish chiropractic patients had more  
6 pain and worse psychological and behavioural characteristics compared to a sample of sick-listed  
7 primary care (specific provider unknown) patients at high risk for chronicity. Our finding showing that  
8 patients with widespread pain were more likely to choose a chiropractor over a GP was contrary to  
9 the general pattern of chiropractic patients being less “complex.” To the best of our knowledge, no  
10 previous studies have compared prevalence of widespread pain in the two populations, but one  
11 study showed that GP patients had more musculoskeletal comorbidities [25], possibly implying more  
12 widespread pain. Two previous studies found an association between higher age and odds of seeking  
13 care from a physiotherapist compared to a chiropractor [26, 27], in line with our results.  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 Many of the patient characteristics associated with choice of primary care provider in this study have  
24 previously been found to be significant prognostic factors for the persistent back-related disability  
25 and back pain in older people. For example, duration of back pain and expectation of improvement  
26 [64-69], and higher levels of back-related disability [65-70], are consistently reported as significant  
27 prognostic factors for a poor outcome of a back pain episode. A few studies in older people have  
28 found that single symptoms of neurological involvement such as leg pain below the knee, and the  
29 diagnosis of spinal stenosis were prognostic factors for the outcome of a back pain episode [64, 67].  
30 We combined single symptoms of neurological involvement into a compound measure, but it is likely  
31 that older patients with radiculopathy have worse outcomes than those without radiculopathy.  
32 Although slightly different from widespread pain, the presence of multi-site pain has also in some  
33 studies been found to be a prognostic factor for the outcome of back pain in older adults [67, 71].  
34 The impact of pain catastrophizing on the clinical course of back pain is less clear in older adults [66,  
35 69] compared to younger populations [72], but it is not unreasonable to believe that pain  
36 catastrophizing may be a prognostic factor for back pain in older adults. Thus, the associations  
37 between potential prognostic factors and choice of first primary care provider imply that we can  
38 expect the clinical course of patients in the three primary care groups to be different. Further, they  
39 imply that caution should be exercised when generalizing across primary care populations.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 The results of this study need to be viewed with consideration of some limitations. We instructed the  
54 recruiting primary care providers to invite consecutive patients, but because of obvious time  
55 constraints in clinical practice we could not ask them to keep record of how many declined to  
56 participate, nor of eligible patients that were not invited. This recruitment strategy increases the risk  
57 of selection bias, and thus could reduce the external validity of the study. To compensate for this  
58  
59  
60

1  
2  
3 limitation, we compared the BACE-N sample with the NORLAG MSK subsample. The characteristics of  
4 the two samples were largely comparable, but BACE-N has more men, more participants with higher  
5 education, more in paid work, and more living with their partner. Sex and education level have  
6 previously been shown to be associated with back pain severity and back-related disability in older  
7 adults [12, 13]. Thus, it may be possible that the levels of back pain and back-related disability  
8 presented in this study are slightly underestimated. The NORLAG MSK subsample is sampled from  
9 the general population, which may not be representative of those who seek care. However, the most  
10 important determinants of care-seeking for back pain seems to be pain severity and disability levels  
11 [73]. We therefore believe the assessment to be justified.  
12  
13  
14  
15  
16  
17  
18  
19

20 Another limitation may be the analysis strategy. We chose to keep the variables in the five blocks to  
21 provide a broad assessment of the differences in case-mix in the three primary care settings. To limit  
22 the number of statistical tests performed, univariate pre-testing and testing a “final model” across  
23 blocks were avoided. Furthermore, a different organization of the variables, for example strictly  
24 adhering to the biopsychosocial model [74] or Andersen’s behavioural model of health services use  
25 [75], may have yielded slightly different results. However, our results are largely supported by  
26 previous studies, so the potential differences because of analysis strategy or variable organization  
27 may be negligible. A third limitation is that we were unable to examine some possibly important  
28 determinants for healthcare use, such as access to different providers, patient’s familiarity with  
29 providers, the patient’s economic situation and social network referrals [75-77]. These factors may  
30 be the most important determinants in driving the patient’s choice of first primary care provider, and  
31 including these factors would have given an even broader overview of associations between  
32 individual and contextual characteristics and choice of primary care provider. We suggest that future  
33 research focus on examining the contextual and social factors associated with healthcare service use.  
34 Finally, generalization of our results to other healthcare systems may be limited. Different healthcare  
35 systems may have different access to care, different payment schemes and different professional  
36 training and responsibilities for the healthcare providers, all of which may impact health services  
37 utilization and consequently the patient characteristics associated with choosing different primary  
38 care providers [75, 78, 79].  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 **Conclusion**

55 We found that nearly all older adults with back pain seeking primary care had experienced back pain  
56 previously, and recurring episodes were common. In general, patients with more severe back-related  
57 disability and other clinical symptoms and signs were more likely to visit a GP or a physiotherapist  
58 than a chiropractor. Our results suggest that important patient characteristics are associated with  
59  
60

1  
2  
3 older adult's choice of primary care providers due to back pain, which may affect the clinical course  
4 of back pain for these patients. The findings highlight the need for caution with generalization of  
5 study results across primary care populations. This is an important consideration for healthcare  
6 providers, for the development and implementation of clinical practice guidelines, and for regulators  
7 when developing primary care pathways for back pain. Further research is needed in assessing if the  
8 choice of primary care provider affects future care pathways and the clinical course of back pain in  
9 older adults.  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **Acknowledgments**

19  
20 The authors thank all study participants for their significant contribution. Further, we would like to  
21 thank all the recruiting healthcare providers: Kerstin Ulrich, Lise Lothe, Tim Raven, Andreas Hoff  
22 Nordvik, Christoffer Børsheim, Steinar Forshei, Mette Brekke, Daniel Major, Jan Harald Lønn, Agnes  
23 Mordt, Andrea Kolstad, Mathias Svanevik, Alexander Diesen, Lars Gullestad, Ida Svalstuen, Joakim  
24 Ordahl, Mona Øversveen, Jørgen Kongtorp, Are Hansen, Geir Haram, Palwinder Singh, Svein Erik  
25 Sandelien, Bent Ulseth, Harald Nordheim, Ola Sand, Ragnhild Perstølen, Jorun Salater, Anna Allen-  
26 Unhammer, Morten Nilsen, Haakon Lilleeng Asmyhr, Philip Wilkens, Ane Klevberg, Eli Magnesen,  
27 Aleksander Killingmo, Bjørn Tore Bjørkedal, Stina Lund, Daniel Ekeberg, Berte Marie Enger, Johan  
28 Edvard Tellum, Morten de la Cruz, Bård Kvam, Marte Paulsen, Astrid Figger, Christian Mayer, Trond  
29 Magne Aasberg, Thea Tømmervåg, Jørgen Øyen, Håvard Nordås, Tore Viste Ollestad, Olav Aase,  
30 Renate Meier, Bjørn Røe, Jørgen Øyen, Jack Johnson, Carl-Erik Høgquist, Ingrid Hystad, Mats  
31 Thorbeck, Elisabeth Barø, Kaja McCormick, Lars Martin Fischer, Martin Haagenen, Cathrine  
32 Rosslund, Marianne Storberget, Nora Helk, Grete Bråten, Annecken Lister Haugen, Hege Herstad,  
33 Cathrine Natland, Frøydis Blaker Åsbø, Ola Klaastad, Thorleif Henning Monsen, Geir Wiik, Jørn  
34 Christian Halvorsen, Tonje Høgdahl Mysen, Tine Tandberg, Eir Marie Bergan, Rune Solheim, Ole  
35 Kristoffer Larsen. We thank the research assistants for assisting with clinical examinations and  
36 providing baseline questionnaires to study participants. We thank the BACE-N scientific board, and  
37 the BACE Consortium for important input with designing the study.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 **Author contributions:**

52 ØNV: Study design, data collection, data analyses, manuscript draft. KS: Study design, data  
53 interpretation, critical revision. RMK: Data collection, data interpretation, critical revision. MCS:  
54 Statistical advisor, data interpretation, critical revision. MG: Principal investigator, study design, data  
55 interpretation, critical revision.  
56  
57  
58  
59  
60

**Funding:**

This work was funded by The Norwegian Fund for Post-Graduate Training in Physiotherapy (grant number 90749). The funding body was not involved in designing the study, data collection, analyses or interpretation of data, nor in writing the manuscript.

**Competing interests**

The authors declare no competing interests.

For peer review only

## REFERENCES

1. Vos, T., et al., *Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010*. The Lancet, 2012. **380**(9859): p. 2163-2196.
2. Hoy, D., et al., *A systematic review of the global prevalence of low back pain*. Arthritis & Rheumatism, 2012. **64**(6): p. 2028-2037.
3. Bressler, H.B., et al., *The Prevalence of Low Back Pain in the Elderly: A Systematic Review of the Literature*. Spine, 1999. **24**(17): p. 1813-1819.
4. Paeck, T., et al., *Are older adults missing from low back pain clinical trials? A systematic review and meta-analysis*. Arthritis Care Res (Hoboken), 2014. **66**(8): p. 1220-6.
5. Jarvik, J.G., et al., *Study protocol: the back pain outcomes using longitudinal data (BOLD) registry*. BMC musculoskeletal disorders, 2012. **13**(1): p. 64.
6. Scheele, J., et al., *Back complaints in the elders (BACE); design of cohort studies in primary care: an international consortium*. BMC Musculoskelet Disord, 2011. **12**: p. 193.
7. Kalichman, L., et al., *Spinal stenosis prevalence and association with symptoms: the Framingham Study*. The Spine Journal, 2009. **9**(7): p. 545-550.
8. Van Der Klift, M., et al., *The incidence of vertebral fractures in men and women: the Rotterdam Study*. Journal of bone and mineral research, 2002. **17**(6): p. 1051-1056.
9. de Schepper, E.I., et al., *The association between lumbar disc degeneration and low back pain: the influence of age, gender, and individual radiographic features*. Spine, 2010. **35**(5): p. 531-536.
10. Enthoven, W.T., et al., *Prevalence and "Red Flags" Regarding Specified Causes of Back Pain in Older Adults Presenting in General Practice*. Physical Therapy, 2016. **96**(3): p. 305-312.
11. Enthoven, W.T., et al., *Back Complaints in Older Adults: Prevalence of Neuropathic Pain and Its Characteristics*. Pain Medicine, 2013. **14**(11): p. 1664-1672.
12. Jesus-Moraleida, F.R., et al., *Back Complaints in the Elders in Brazil and the Netherlands: a cross-sectional comparison*. Age and ageing, 2017. **46**(3): p. 476-481.
13. Jarvik, J.G., et al., *Back pain in seniors: the back pain outcomes using longitudinal data (BOLD) cohort baseline data*. BMC Musculoskeletal Disorders, 2014. **15**(1): p. 1-23.
14. Kinge, J.M., et al., *Musculoskeletal disorders in Norway: prevalence of chronicity and use of primary and specialist health care services*. BMC Musculoskeletal Disorders, 2015. **16**(1): p. 1-9.
15. Werner, E.L. and A. Indahl, *Kunnskap, praksis og holdninger til rygglidelser hos leger, fysioterapeuter og kiropraktorer*. Tidsskr Nor Laegeforen, 2005. **125**(13): p. 1794–7.
16. Fritz, J.M., J. Kim, and J. Dorius, *Importance of the type of provider seen to begin health care for a new episode low back pain: associations with future utilization and costs*. Journal of evaluation in clinical practice, 2016. **22**(2): p. 247-252.
17. Kazis, L.E., et al., *Observational retrospective study of the association of initial healthcare provider for new-onset low back pain with early and long-term opioid use*. BMJ Open, 2019. **9**(9): p. e028633.
18. Carey, T.S., et al., *Acute Severe Low Back Pain: A Population-based Study of Prevalence and Care-seeking*. Spine, 1996. **21**(3): p. 339-344.
19. Hurwitz, E.L. and H. Morgenstern, *The effects of comorbidity and other factors on medical versus chiropractic care for back problems*. Spine, 1997. **22**(19): p. 2254-2263.
20. Nyiendo, J., et al., *Patient characteristics and physicians' practice activities for patients with chronic low back pain: A practice-based study of primary care and chiropractic physicians*. 2001. **24**(2): p. 92-100.
21. Côté, P., J.D. Cassidy, and L. Carroll, *The treatment of neck and low back pain: who seeks care? who goes where?* Medical care, 2001: p. 956-967.

22. Sharma, R., M. Haas, and M. Stano, *Patient Attitudes, Insurance, and Other Determinants of Self-Referral to Medical and Chiropractic Physicians*. American Journal of Public Health, 2003. **93**(12): p. 2111-2117.
23. Hestbaek, L., et al., *Low back pain in primary care: a description of 1250 patients with low back pain in danish general and chiropractic practice*. International journal of family medicine, 2014. **2014**.
24. Eklund, A., et al., *Psychological and behavioral differences between low back pain populations: a comparative analysis of chiropractic, primary and secondary care patients*. BMC Musculoskeletal Disorders, 2015. **16**(1).
25. Hartvigsen, L., et al., *Baseline Characteristics May Help Indicate the Best Choice of Health Care Provider for Back Pain Patients in Primary Care: Results From a Prospective Cohort Study*. Journal of Manipulative and Physiological Therapeutics, 2020. **43**(1): p. 13-23.
26. Chevan, J. and D.L. Riddle, *Factors associated with care seeking from physicians, physical therapists, or chiropractors by persons with spinal pain: a population-based study*. journal of orthopaedic & sports physical therapy, 2011. **41**(7): p. 467-476.
27. Blanchette, M.-A., et al., *Workers' characteristics associated with the type of healthcare provider first seen for occupational back pain*. BMC Musculoskeletal Disorders, 2016. **17**(1).
28. Sibbritt, D., et al., *Severity of back pain may influence choice and order of practitioner consultations across conventional, allied and complementary health care: a cross-sectional study of 1851 mid-age Australian women*. BMC Musculoskeletal Disorders, 2016. **17**(1).
29. Walker, B.F., R. Muller, and W.D. Grant, *Low back pain in Australian adults. Health provider utilization and care seeking*. Journal of manipulative and physiological therapeutics, 2004. **27**(5): p. 327-335.
30. Sperre Saunes, I., et al., *Norway: Health system review*. 2020.
31. Loge, J.H. and S. Kaasa, *Short Form 36 (SF-36) health survey: normative data from the general Norwegian population*. Scandinavian Journal of Social Medicine, 1998. **26**(4): p. 250-258.
32. Bush, K., *The AUDIT Alcohol Consumption Questions (AUDIT-C). An Effective Brief Screening Test for Problem Drinking*. Archives of Internal Medicine, 1998. **158**(16): p. 1789.
33. Bradley, K.A., et al., *AUDIT-C as a Brief Screen for Alcohol Misuse in Primary Care*. Alcoholism: Clinical and Experimental Research, 2007. **31**(7): p. 1208-1217.
34. Norwegian National Advisory Unit on Concurrent Substance Abuse and Mental Health Disorders. *AUDIT-C*. 23.09.2020]; Available from: <https://rop.no/kartleggingsverktoey/audit-c/>.
35. Sangha, O., et al., *The self-administered comorbidity questionnaire: A new method to assess comorbidity for clinical and health services research*. Arthritis & Rheumatism, 2003. **49**(2): p. 156-163.
36. Wolfe, F., et al., *Revised chronic widespread pain criteria: development from and integration with fibromyalgia criteria*. Scandinavian Journal of Pain, 2019. **20**(1): p. 77-86.
37. Melzack, R., *The McGill Pain Questionnaire: Major properties and scoring methods*. Pain, 1975. **1**(3): p. 277-299.
38. Helbostad, J.L., et al., *Validation of the Falls Efficacy Scale-International in fall-prone older persons*. Age and Ageing, 2010. **39**(2): p. 259-259.
39. Von Korff, M., M.P. Jensen, and P. Karoly, *Assessing Global Pain Severity by Self-Report in Clinical and Health Services Research*. Spine, 2000. **25**(24): p. 3140-3151.
40. Grotle, M., J. Brox, and N. Vollestad, *Cross-cultural adaptation of the Norwegian versions of the Roland-Morris Disability Questionnaire and the Oswestry Disability Index*. Journal of Rehabilitation Medicine, 2003. **35**(5): p. 241-247.
41. Buysse, D.J., et al., *The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research*. Psychiatry research, 1989. **28**(2): p. 193-213.
42. Roos, E.M., et al., *Knee Injury and Osteoarthritis Outcome Score (KOOS)—Development of a Self-Administered Outcome Measure*. Journal of Orthopaedic & Sports Physical Therapy, 1998. **28**(2): p. 88-96.

- 1  
2  
3 43. Grotle, M., J.I. Brox, and N.K. Vøllestad, *Reliability, validity and responsiveness of the fear-avoidance beliefs questionnaire: methodological aspects of the Norwegian version*. Journal of rehabilitation medicine, 2006. **38**(6): p. 346-353.
- 4  
5  
6 44. Radloff, L.S., *The CES-D scale: A self report depression scale for research in the general population*. Applied Psychological Measurements, 1977. **1**(3): p. 385-401.
- 7  
8 45. Fernandes, L., et al., *Cross-cultural adaptation and validation of the Norwegian pain catastrophizing scale in patients with low back pain*. BMC musculoskeletal disorders, 2012. **13**(1): p. 111.
- 9  
10  
11 46. Tingulstad, A., et al., *Back beliefs among elderly seeking health care due to back pain; psychometric properties of the Norwegian version of the back beliefs questionnaire*. BMC Musculoskeletal Disorders, 2019. **20**(1).
- 12  
13  
14 47. Hill, J.C., et al., *A primary care back pain screening tool: Identifying patient subgroups for initial treatment*. Arthritis & Rheumatism, 2008. **59**(5): p. 632-641.
- 15  
16  
17 48. Strand, L.I., R. Moe-Nilssen, and A.E. Ljunggren, *Back Performance Scale for the Assessment of Mobility-Related Activities in People With Back Pain*. Physical Therapy, 2002. **82**(12): p. 1213-1223.
- 18  
19  
20 49. Podsiadlo, D. and S. Richardson, *The Timed "Up & Go": A Test of Basic Functional Mobility for Frail Elderly Persons*. Journal of the American Geriatrics Society, 1991. **39**(2): p. 142-148.
- 21  
22  
23 50. Stynes, S., et al., *Clinical diagnostic model for sciatica developed in primary care patients with low back-related leg pain*. PloS one, 2018. **13**(4).
- 24  
25  
26 51. Bender, R. and S. Lange, *Adjusting for multiple testing—when and how?* Journal of Clinical Epidemiology, 2001. **54**(4): p. 343-349.
- 27  
28  
29 52. Slagsvold, B., et al., *Life-course, ageing and generations in Norway:the NorLAG study*. Norsk Epidemiologi, 2012. **22**(2).
- 30  
31  
32 53. Torsteinsen, A. and A. Holmøy, *Den norske studien av livsløp, aldring og generasjon—tredje runde (NorLAG3). Dokumentasjonsrapport*. 2019.
- 33  
34  
35 54. Moons, K.G.M., et al., *Prognosis and prognostic research: what, why, and how?* BMJ, 2009. **338**: p. b375.
- 36  
37  
38 55. Dunn, K.M., K. Jordan, and P.R. Croft, *Characterizing the course of low back pain: a latent class analysis*. American journal of epidemiology, 2006. **163**(8): p. 754-761.
- 39  
40  
41 56. Dunn, K.M., P. Campbell, and K.P. Jordan, *Long-term trajectories of back pain: cohort study with 7-year follow-up*. BMJ open, 2013. **3**(12): p. e003838.
- 42  
43  
44 57. Kongsted, A., et al., *Patients with low back pain had distinct clinical course patterns that were typically neither complete recovery nor constant pain. A latent class analysis of longitudinal data*. The Spine Journal: Official Journal Of The North American Spine Society, 2015. **15**(5): p. 885-894.
- 45  
46  
47 58. Kongsted, A., et al., *What have we learned from ten years of trajectory research in low back pain?* BMC Musculoskeletal Disorders, 2016. **17**: p. 1-11.
- 48  
49  
50 59. Nordstoga, A.L., et al., *Improvement in Work Ability, Psychological Distress and Pain Sites in Relation to Low Back Pain Prognosis: A Longitudinal Observational Study in Primary Care*. Spine, 2019. **44**(7): p. E423-E429.
- 51  
52  
53 60. Grotle, M., et al., *Prognostic factors in first-time care seekers due to acute low back pain*. European Journal of Pain, 2007. **11**(3): p. 290-298.
- 54  
55  
56 61. Scheele, J., et al., *Characteristics of older patients with back pain in general practice: BACE cohort study*. Eur J Pain, 2013. **18**(2): p. 279-87.
- 57  
58  
59 62. Jesus-Moraleida, F.R.D., et al., *The Brazilian Back Complaints in the Elders (Brazilian BACE) study: characteristics of Brazilian older adults with a new episode of low back pain*. Brazilian Journal of Physical Therapy, 2017.
- 60  
61  
62 63. Peacock, S. and S. Patel, *Cultural influences on pain*. Reviews in pain, 2008. **1**(2): p. 6-9.
- 63  
64  
65 64. Scheele, J., et al., *Course and prognosis of older back pain patients in general practice: a prospective cohort study*. Pain, 2013. **154**(6): p. 951-7.

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
65. Deyo, R.A., et al., *Trajectories of symptoms and function in older adults with low back disorders*. *Spine (Phila Pa 1976)*, 2015. **40**(17): p. 1352-62.
66. Enthoven, W.T., et al., *Defining trajectories in older adults with back pain presenting in general practice*. *Age and ageing*, 2016. **45**(6): p. 878-883.
67. Rundell, S.D., et al., *Predictors of Persistent Disability and Back Pain in Older Adults with a New Episode of Care for Back Pain*. *Pain Med*, 2017. **18**(6): p. 1049-1062.
68. Jarvik, J.G., et al., *Long-term outcomes of a large, prospective observational cohort of older adults with back pain*. *Spine J*, 2018. **18**(9): p. 1540-1551.
69. van der Gaag, W.H., et al., *Developing clinical prediction models for non-recovery in older patients seeking care for back pain: the BACE prospective cohort study*. *PAIN*, 2021. **162**(6): p. 1632-1640.
70. Van Den Berg, R., et al., *Clinical and radiographic features of spinal osteoarthritis predict long-term persistence and severity of back pain in older adults*. *Annals of Physical and Rehabilitation Medicine*, 2020.
71. Rundell, S.D., et al., *Multisite Pain Is Associated with Long-term Patient-Reported Outcomes in Older Adults with Persistent Back Pain*. *Pain Med*, 2019.
72. Wertli, M.M., et al., *Catastrophizing—a prognostic factor for outcome in patients with low back pain: a systematic review*. *The Spine Journal*, 2014. **14**(11): p. 2639-2657.
73. Ferreira, M., et al., *Factors defining care-seeking in low back pain - A meta-analysis of population based surveys*. *European Journal of Pain*, 2010. **14**(7): p. 747.e1-747.e7.
74. Waddell, G., *1987 Volvo award in clinical sciences. A new clinical model for the treatment of low-back pain*. *Spine*, 1987. **12**(7): p. 632-644.
75. Andersen, R.M., *National health surveys and the behavioral model of health services use*. *Medical care*, 2008. **46**(7): p. 647-653.
76. Davis, M.A., et al., *Regional Supply of Chiropractic Care and Visits to Primary Care Physicians for Back and Neck Pain*. *The Journal of the American Board of Family Medicine*, 2015. **28**(4): p. 481-490.
77. Kirby, E.R., et al., *A qualitative study of influences on older women's practitioner choices for back pain care*. *BMC Health Services Research*, 2014. **14**(1): p. 131.
78. Garrity, B., et al., *Unrestricted Direct Access to Physical Therapist Services Is Associated With Lower Health Care Utilization and Costs in Patients With New-Onset Low Back Pain*. *Physical Therapy*, 2019.
79. Carey, K., et al., *Health insurance design and conservative therapy for low back pain*. *Am J Manag Care*, 2019. **25**(6): p. e182-e187.



**Table 1:** Baseline characteristics of study participants\*

	Missing, n (%)	Total (n=452)	GP (n=127)	PT (n=130)	Chiro (n=195)
<b>Sociodemographic variables</b>					
Age, median (IQR)	0 (0.0)	66 (59-72)	67 (60-73)	68 (63-74)	63 (58-71)
Female, n (%)	0 (0.0)	235 (52.0)	74 (58.3)	70 (53.8)	89 (46.1)
Marital status	19 (4.2)				
Married or living with partner, n (%)		347 (76.8)	90 (70.1)	98 (74.6)	158 (81.0)
Employment status	5 (1.1)				
Currently in paid work, n (%)		212 (45.3)	57 (43.3)	49 (31.5)	106 (55.9)
Educational level, n (%)	20 (4.4)				
Low (elementary + high school)		253 (56.0)	72 (56.7)	70 (55.1)	110 (56.4)
High (university level)		199 (44.0)	55 (43.3)	60 (44.9)	85 (43.6)
<b>General health variables</b>					
Health-related quality of life (SF-36 0-100)	41 (9.1)				
Mental sumscore, mean (SD)		52.5 (10.0)	50.5 (11.5)	53.4 (10.0)	53.2 (8.8)
Physical sumscore, mean (SD)		41.4 (8.4)	40.0 (7.9)	40.6 (8.0)	42.8 (8.9)
Hazardous alcohol consumption (AUDIT-C <sup>a</sup> ), n (%)	59 (13.1)	228 (50.4)	65 (51.1)	65 (50.0)	98 (50.2)
Smoking status, n (%)	22 (4.9)				
Current smoker		63 (13.9)	21 (16.5)	13 (10.0)	28 (14.3)
Previous		203 (44.9)	59 (46.4)	60 (46.2)	84 (43.1)
Never		186 (41.2)	47 (37.0)	57 (43.8)	83 (42.6)
Number of comorbidities (SCQ 0-15), median (IQR)	18 (4.0)	1 (1-2)	1 (0-2)	2 (1-2)	1 (1-5)
BMI, mean (SD)	14 (3.1)	27.6 (4.7)	27.6 (4.5)	27.5 (4.7)	27.7 (4.8)
Fall last 6 weeks, n (%)	24 (5.3)	73 (16.1)	13 (10.2)	24 (18.4)	35 (18.2)
Falls self-efficacy (FESI 16-64), mean (SD)	48 (10.6)	21.8 (6.0)	22.4 (6.3)	22.2 (6.1)	21.1 (5.7)
Widespread pain, n (%)	16 (3.5)	33 (7.3)	5 (4.0)	7 (5.3)	21 (10.8)
<b>Current back pain and back pain history variables</b>					
Previous back pain, n (%)	58 (12.8)				
Monthly		127 (28.1)	42 (33.1)	46 (35.4)	40 (20.5)
Every year		174 (38.5)	45 (35.4)	44 (33.8)	86 (44.1)
Every 1-5 years		90 (19.9)	26 (20.5)	19 (14.6)	45 (23.1)
Every five years		45 (10.0)	10 (7.9)	16 (12.3)	20 (10.3)
Only once		15 (3.3)	4 (3.1)	6 (4.6)	4 (2.1)
Back pain location of current episode, n (%)	11 (2.4)				
Thoracic only		19 (4.2)	4 (3.1)	7 (5.4)	8 (4.1)
Lumbar only		382 (84.5)	106 (83.5)	109 (83.8)	167 (85.6)
Both		51 (11.3)	17 (13.4)	14 (10.8)	20 (10.3)
Duration of current episode, n (%)	76 (16.8)				
0-6 weeks		297 (65.7)	74 (58.3)	67 (51.5)	156 (80.0)
6 weeks to 3 months		59 (13.1)	22 (17.3)	21 (16.2)	16 (8.2)
3 months or over		96 (21.2)	31 (24.4)	42 (32.3)	23 (11.8)
Back pain severity (NRS 0-10), mean (SD)	31 (6.9)	5.4 (2.3)	5.7 (2.2)	5.1 (2.3)	5.4 (2.4)
Back-related disability (RMDQ 0-24), median (IQR)	45 (10.0)	9 (5-13)	10 (6-14)	9 (6-13)	8 (3-13)
Sleep problems due to back pain, n (%)	24 (5.3)				
Weekly		189 (41.8)	60 (47.2)	49 (37.7)	80 (41.0)
Less than weekly		263 (58.2)	67 (52.8)	81 (62.3)	115 (59.0)
Morning stiffness, n (%)	26 (5.8)				
Significant or extreme		178 (39.3)	47 (37.0)	51 (39.2)	81 (41.5)
Moderate		144 (31.9)	44 (34.6)	48 (36.9)	51 (26.2)
Some or none		130 (28.8)	36 (28.3)	31 (23.9)	63 (32.3)
<b>Psychological variables</b>					
Kinesiophobia (FABQ-PA 0-24), median (IQR)	18 (4.0)	10 (5-14)	11 (6-14)	10 (5-15)	9 (3-13)
Depression (CES-D 0-60), median (IQR)	57 (12.6)	8 (4-15)	10 (4-17)	8.5 (4-15)	7 (4-13)
Pain catastrophizing (PCS 0-52), median (IQR)	35 (7.7)	10 (4-16)	11 (5-18)	12 (5-18)	7 (3-14)
Back beliefs (BBQ 9-45), mean (SD)	57 (12.6)	29.8 (7.0)	28.0 (6.9)	29.3 (7.2)	31.3 (6.7)
Expectations for back pain next 3 months, n (%)	19 (4.2)				
Fully recovered		115 (25.4)	19 (15.0)	24 (18.5)	72 (36.9)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Much better		226 (50.0)	66 (52.0)	71 (54.6)	89 (45.6)
No change or worse		111 (24.6)	42 (33.0)	35 (26.9)	33 (16.9)
Start Back Screening Tool risk profiles, n (%)	31 (6.9)				
Low		297 (65.7)	72 (56.7)	92 (70.8)	133 (68.2)
Medium		125 (27.7)	38 (29.9)	32 (24.6)	55 (28.2)
High		30 (6.6)	16 (12.6)	6 (4.6)	8 (4.1)
<b>Clinical variables</b>					
Physical performance (BPS 0-18), median (IQR)	20 (4.4)	5 (2-8)	7 (3-9)	5 (3-8)	4 (1-7)
Timed up and go, mean seconds (SD)	7 (1.5)	8.0 (2.5)	8.2 (3.0)	8.3 (2.3)	7.8 (2.2)
Positive diagnostic rule for radiculopathy, n (%)	38 (8.4)	99 (22.0)	37 (29.1)	31 (23.8)	31 (15.9)
Number of red flags (0-12), median (IQR)	50 (11.0)	1 (0-2)	1 (0-2)	1 (0-2)	1 (0-1)
Pain on active range of motion, n (%)	9 (2.0)	295 (65.3)	86 (67.7)	88 (67.7)	120 (61.5)

GP: General practitioner; PT: Physiotherapist; Chiro: Chiropractor; IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
\* The presented characteristics are pooled estimates based on multiple imputation procedures  
^ AUDIT-C scores of ≥3/12 for women and ≥4/12 indicates hazardous alcohol consumption

**Table 2:** Multinomial regression analyses; multivariate associations between patient characteristics and choice of healthcare provider (dependent variable) \*

	GP (n=127) Odds ratio (95% CI)	p-value	PT (n=130) Odds ratio (95% CI)	p-value
<b>Block i) Sociodemographic variables</b>				
Age	1.03 (0.99-1.07)	0.11	1.04 (1.00-1.08)	<b>0.03</b>
Gender				
Female	1.53 (0.96-2.45)	0.07	1.33 (0.83-2.12)	0.24
Male (ref.)	1.00		1.00	
Marital status				
Married/cohabiting	0.67 (0.38-1.19)	<b>0.17</b>	0.90 (0.51-1.61)	0.73
Not married/cohabiting (ref.)	1.00		1.00	
Educational level				
Higher education	1.02 (0.64-1.62)	0.94	1.08 (0.68-1.73)	0.73
Lower education (ref.)	1.00		1.00	
Employment status				
Currently in paid work	0.86 (0.46-1.62)	0.64	0.55 (0.30-1.01)	0.05
No paid work (ref.)	1.00		1.00	
<b>Block ii) General health variables</b>				
Hazardous alcohol intake (AUDIT-C)				
Yes	1.20 (0.73-1.97)	0.47	1.08 (0.64-1.81)	0.77
No (ref.)	1.00		1.00	
Smoking status				
Yes	1.18 (0.56-2.46)	0.67	0.64 (0.28-1.48)	0.29
Previously	1.31 (0.77-2.23)	0.32	1.11 (0.67-1.83)	0.70
No (ref.)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.96 (0.93-1.00)	<b>0.03</b>	0.98 (0.95-1.01)	0.19
Mental component	0.97 (0.95-1.00)	<b>0.02</b>	1.01 (0.98-1.03)	0.73
BMI	0.98 (0.93-1.04)	0.53	0.97 (0.92-1.02)	0.28
Comorbidities (SCQ, 0-15)	1.07 (0.86-1.33)	0.53	1.15 (0.95-1.40)	0.17
Widespread pain				
Yes	0.22 (0.06-0.81)	<b>0.02</b>	0.46 (0.18-1.16)	0.10
No (ref.)	1.00		1.00	
Falls self-efficacy (FES-I, 16-64)	1.00 (0.95-1.05)	0.98	1.03 (0.95-1.05)	0.32
<b>Block iii) Current back pain and back pain history variables</b>				
Back pain severity (NRS, 0-10)	1.02 (0.91-1.14)	0.77	0.90 (0.80-1.01)	0.08
Back-related disability (RMDQ, 0-24)	1.06 (1.00-1.12)	<b>0.04</b>	1.07 (1.01-1.13)	<b>0.02</b>
Duration				
Over 3 months	2.92 (1.28-6.66)	<b>0.01</b>	4.57 (1.99-10.50)	<b>&lt;0.01</b>

6 weeks to 3 months	3.03 (1.27-4.97)	<b>0.02</b>	3.17 (1.28-7.84)	<b>0.01</b>
0-6 weeks (ref.)	1.00		1.00	
Morning stiffness				
Significant or extreme	0.76 (0.41-1.42)	0.39	1.21 (0.64-2.30)	0.55
Moderate	1.37 (0.74-2.56)	0.32	2.03 (1.08-3.81)	<b>0.03</b>
A little or none (ref.)	1.00		1.00	
Sleep problems attributable to back pain				
Weekly	1.09 (0.63-1.89)	0.76	0.75 (0.41-1.35)	0.33
Less than weekly (ref.)	1.00		1.00	
Previous back pain frequency				
Yearly	1.11 (0.65-1.92)	0.70	1.00 (0.59-1.69)	0.99
Not yearly (ref.)	1.00		1.00	
<b>Block iv) Psychological variables</b>				
Fear-avoidance (FABQ-PA, 0-24)	1.02 (0.98-1.07)	0.32	1.03 (0.98-1.08)	0.22
Pain catastrophizing (PCS, 0-52)	1.04 (1.00-1.07)	0.05	1.06 (1.02-1.10)	<b>&lt;0.01</b>
Depression symptoms (CESD, 0-60)	0.99 (0.95-1.03)	0.53	0.99 (0.96-1.03)	0.61
Back beliefs (BBQ, 9-45)	0.97 (0.93-1.02)	0.23	0.99 (0.95-1.03)	0.67
Expectation for back pain in 3 months				
Recovered	0.26 (0.12-0.56)	<b>&lt;0.01</b>	0.39 (0.19-0.79)	<b>0.01</b>
Much better	0.65 (0.35-1.19)	0.16	0.85 (0.46-1.58)	0.61
No change or worse (ref.)	1.00		1.00	
Start Back Screening tool risk category				
Medium + high risk	1.02 (0.55-1.87)	0.95	0.49 (0.26-0.92)	<b>0.03</b>
Low risk (ref.)	1.00		1.00	
<b>Block v) Clinical variables</b>				
Number of red flags (0-12)	1.25 (0.99-1.58)	0.06	1.19 (0.96-1.48)	0.12
Diagnostic tool for radiculopathy				
Positive	1.94 (1.08-3.47)	<b>0.03</b>	1.52 (0.85-2.73)	0.16
Negative (ref.)	1.00		1.00	
Pain on active range of motion				
Yes	0.95 (0.57-1.58)	0.85	1.09 (0.67-1.80)	0.72
No (ref.)	1.00		1.00	
Trunk mobility performance (BPS, 0-18)	1.16 (1.08-1.24)	<b>&lt;0.01</b>	1.07 (1.00-1.15)	<b>0.04</b>
Timed Up and Go, mean seconds	0.93 (0.83-1.04)	0.20	1.00 (0.90-1.11)	0.93

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale - International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire - Physical Activity subscale; CES-D: Center for Epidemiological Studies - Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

\* The multinomial regression analyses are based on pooled estimates from multiple regression analyses

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.

The chiropractic group (n=195) was the reference dependent variable.

Models were built block-wise within the five blocks: i) sociodemographic ii) general health iii) current episode and back pain history iv) psychological and v) clinical. All variables were included simultaneously.

**SUPPLEMENTARY FILE**

Table of Contents

Sensitivity analyses S1, complete case analyses: ..... 2

Sensitivity analyses S2, physiotherapy patients:..... 5

Sensitivity analyses S3, only low back pain patients: ..... 7

Assessment of generalizability S4: ..... 9

For peer review only

## Sensitivity analyses S1, complete case analyses:

### Methods:

Multiple multinomial regression. One model per variable block. The total number of available cases per category will vary with number of missing for each block, and is thus shown for each block.

Additionally, bootstrapping was performed for n=1000 bootstrapping samples. The average bootstrapping odds ratios and their corresponding bias-corrected accelerated 95% confidence intervals (BCa 95% CI) are provided. Because of few observations in the Start Back Screening Tool high risk group, we chose to combine this group with the medium risk group.

### Results:

See Table S1 for details. No substantial changes in point estimates were detected in the multinomial regression analyses when comparing complete cases analyses to the pooled imputed estimates. There were, however, some changes in p-values. In the complete case analyses, age and being in the SBT medium risk group were not significantly associated with choosing a PT compared to a chiropractor. Further, in the complete case analyses, having more red flags were significantly associated with choosing a GP compared to a chiropractor. As can be seen from the bootstrapping procedure, odds ratios and BCa 95% CIs were stable for all variables, except for the SBT high risk group. Here, the BCa 95% CIs indicate that the odds ratios cannot be trusted for this specific variable.

**Table S1:** Complete case analyses of multiple multinomial regression analyses. Chiropractic group is the reference group.

<b>Block i) Sociodemographic factors. Chiropractor n=181</b>						
	GP (n=113) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Physio (n=108) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Age	1.03 (0.99, 1.08)	0.11	1.03 (0.99, 1.08)	1.03 (0.99, 1.07)	0.14	1.03 (0.98, 1.08)
Gender						
Female	1.33 (0.81, 2.17)	0.26	1.33 (0.80, 2.09)	1.40 (0.85, 2.33)	0.19	1.40 (0.82, 2.27)
Male (ref)	1.00		1.00	1.00		1.00
Marital status						
Married/cohabiting	0.66 (0.37, 1.19)	0.17	0.66 (0.36, 1.26)	0.92 (0.49, 1.72)	0.79	0.92 (0.48, 1.68)
Not married/cohabiting (ref)	1.00		1.00	1.00		1.00
Educational level						
Higher education	1.02 (0.63, 1.65)	0.95	1.02 (0.60, 1.70)	1.08 (0.66, 1.77)	0.77	1.08 (0.65, 1.79)
Lower education (ref)	1.00		1.00	1.00		1.00
Employment status						
Currently in paid work	0.96 (0.50, 1.86)	0.91	0.96 (0.50, 1.78)	0.53 (0.27, 1.03)	0.06	0.53 (0.26, 1.00)
No paid work (ref)	1.00		1.00	1.00		1.00
<b>Block ii) General health factors. Chiropractor n=155</b>						
	GP (n=92) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Physio (n=89) Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Hazardous alcohol intake (AUDIT-C)						
Yes	1.23 (0.70, 2.15)	0.48	1.23 (0.67, 2.15)	1.67 (0.95, 2.92)	0.07	1.67 (0.96, 3.12)
No (ref)	1.00		1.00	1.00		1.00
Smoking						
Yes	1.37 (0.57, 3.26)	0.48	1.37 (0.44, 3.89)	0.63 (0.22, 1.76)	0.37	0.63 (0.16, 1.64)
Previously	1.47 (0.82, 2.66)	0.20	1.47 (0.82, 3.16)	1.43 (0.81, 2.54)	0.22	1.43 (0.80, 2.78)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

No (ref)	1.00		1.00	1.00		1.00
Health-related quality of life (SF-36, 0-100)						
Physical component	0.96 (0.92, 0.99)	<b>0.03</b>	0.96 (0.92, 0.99)	0.97 (0.94, 1.01)	0.96	0.97 (0.94, 1.01)
Mental component	0.95 (0.92, 0.98)	<b>0.002</b>	0.95 (0.92, 0.98)	1.00 (0.94, 1.07)	0.99	1.00 (0.97, 1.04)
BMI	0.99 (0.93, 1.06)	0.81	0.99 (0.92, 1.06)	1.00 (0.94, 1.07)	0.99	1.00 (0.92, 1.09)
Comorbidities (SCQ, 0-15)	1.02 (0.81, 1.29)	0.88	1.02 (0.76, 1.31)	1.12 (0.89, 1.41)	0.33	1.12 (0.90, 1.42)
Widespread pain						
Yes	0.16 (0.03, 0.79)	<b>0.03</b>	0.16 (0.07, 0.38)	0.50 (0.15, 1.67)	0.26	0.49 (0.10, 1.41)
No (ref)	1.00		1.00	1.00		1.00
Falls self-efficacy (FESI, 16-64)	1.00 (0.93, 1.05)	0.73	0.99 (0.92, 1.07)	0.99 (0.93, 1.06)	0.77	0.99 (0.92, 1.07)

**Block iii) Current episode and back pain history. Chiropractor n=134**

	GP (n=80)			Physio (n=92)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Back pain severity (NRS, 0-10)	1.06 (0.91, 1.22)	0.49	1.05 (0.90, 1.22)	0.94 (0.82, 1.08)	0.40	0.94 (0.82, 1.10)
Back-related disability (RMDQ, 0-24)	1.06 (0.99, 1.13)	0.12	1.06 (0.97, 1.14)	1.06 (0.99, 1.13)	0.11	1.06 (0.97, 1.15)
Duration						
Over 3 months	5.49 (2.34, 12.85)	<b>&lt;0.001</b>	5.49(1.93, 22.47)	9.00 (4.03, 20.13)	<b>&lt;0.001</b>	9.00 (3.53, 32.69)
6 weeks to 3 months	4.92 (1.92, 12.61)	<b>0.001</b>	4.92 (1.78, 36.27)	4.90 (1.91, 12.56)	<b>0.001</b>	4.90 (1.62, 18.99)
0-6 weeks (ref)	1.00		1.00	1.00		1.00
Morning stiffness						
Significant or extreme	1.02 (0.47, 2.24)	0.96	1.02 (0.54, 2.80)	1.23 (0.57, 2.67)	0.60	1.23 (0.56, 2.75)
Moderate	1.92 (0.88, 4.22)	0.10	1.93 (0.69, 5.55)	2.36 (1.09, 5.14)	<b>0.03</b>	2.36 (1.07, 5.75)
A little or none (ref)	1.00		1.00	1.00		1.00
Sleep problems attributable to back pain						
Weekly	0.82 (0.42, 1.62)	0.57	0.82 (0.33, 1.61)	0.77 (0.39, 1.52)	0.45	0.77 (0.38, 1.47)
Less than weekly (ref)	1.00		1.00	1.00		1.00
Previous back pain frequency						
Yearly	1.06 (0.57, 1.96)	0.57	1.06 (0.49, 2.14)	1.07 (0.58, 1.99)	0.82	1.07 (0.51, 2.17)
Not yearly (ref)	1.00		1.00	1.00		1.00

**Block iv) Psychological factors. Chiropractor n=155**

	GP (n=96)			Physio (n=94)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Fear-avoidance (FABQ-PA, 0-24)	1.03 (0.98, 1.09)	0.22	1.03 (0.98, 1.10)	1.03 (0.98, 1.08)	0.24	1.03 (0.98, 1.09)
Pain catastrophizing (PCS, 0-52)	1.02 (0.97, 1.06)	0.45	1.02 (0.97, 1.07)	1.05 (1.01, 1.10)	<b>0.02</b>	1.05 (1.01, 1.11)
Depression symptoms (CESD, 0-60)	1.00 (0.96, 1.04)	0.97	1.00 (0.95, 1.04)	0.99 (0.95, 1.04)	0.66	0.99 (0.94, 1.04)
Back beliefs (BBQ, 9-45)	0.97 (0.92, 1.02)	0.21	0.97 (0.92, 1.01)	0.99 (0.94, 1.04)	0.64	0.99 (0.94, 1.04)
Expectation for back pain in 3 months						
Recovered	0.24 (0.10, 0.54)	<b>0.01</b>	0.24 (0.09, 0.48)	0.43 (0.19, 0.95)	<b>0.04</b>	0.43 (0.19, 0.93)
Much better	0.57 (0.29, 1.12)	0.10	0.57 (0.28, 1.10)	0.83 (0.41, 1.68)	0.61	0.83 (0.40, 1.91)
No change or worse(ref)	1.00		1.00	1.00		1.00
Start Back Screening tool						
Medium+high risk	1.31 (0.68, 2.53)	0.42	1.31 (0.62, 2.83)	0.49 (0.24, 1.00)	0.05	0.49 (0.26, 0.86)
Low risk (ref)	1.00		1.00	1.00		1.00

**Block v) Clinical variables. Chiropractor n=159**

	GP (n=105)			Physio (n=110)		
	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*	Odds ratio (95% CI)	p-value	OR (BCa 95% CI)*
Number of red flags (0-12)	1.30 (1.02, 1.67)	<b>0.04</b>	1.30 (0.97, 1.75)	1.26 (0.99, 1.60)	0.06	1.26 (0.98, 1.59)
Nerve involvement diagnostic tool						
Positive	2.34 (1.27, 4.31)	<b>0.01</b>	2.34 (1.23, 4.69)	1.70 (0.93, 3.14)	0.09	1.70 (0.91, 3.33)
Negative (ref)	1.00		1.00	1.00		1.00
Pain on active range of motion						
Yes	0.76 (0.43, 1.32)	0.33	0.76 (0.44, 1.28)	0.92 (0.54, 1.58)	0.77	0.92 (0.53, 1.64)
No (ref)	1.00		1.00	1.00		1.00
Physical performance (BPS, 0-18)	1.19 (1.10, 1.29)	<b>&lt;0.001</b>	1.19 (1.11, 1.32)	1.10 (1.02, 1.19)	<b>0.01</b>	1.10 (1.02, 1.64)
Timed up and go, mean seconds	0.90 (0.79, 1.03)	0.14	0.90 (0.80, 1.02)	1.00 (0.88, 1.13)	0.94	1.00 (0.87, 1.13)

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire;

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

---

FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.  
\*OR (BCa 95% CI) is average odds ratios from 1000 bootstrapping samples, including bias-corrected accelerated 95% confidence intervals.

For peer review only

## Sensitivity analyses S2, physiotherapy patients:

Analyses of differences between physiotherapy patients recruited before 01.01.2018 and after 01.01.2018. After 01.01.2018, there was direct access to physiotherapy in Norway, which potentially could change the population characteristics.

### Methods:

- Univariate analyses corresponding to measurement level and distribution: Chi square test or Fischer's exact test for categorical variables, individual sample t-test for normally distributed continuous variables, Mann Whitney U-test for continuous variables with a skewed distribution
- We used the pooled estimates from multiple imputation that were used in the article table 1 and 2

### Results:

See Table S2 for details. We found statistically significant differences between PT patients recruited before and after 01.01.2018 on the BBQ and BPS. PT patients recruited before 01.01.2018 held significantly more optimistic beliefs about back pain, with a mean (SD) BBQ score of 30.3 (6.8) for patients recruited before 01.01.2018 compared to 27.3 (7.5) for patients recruited after 01.01.2018 ( $p=0.03$ ). PT patients recruited before 01.01.2018 had significantly better trunk mobility performance, with a median (IQR) of 5 (2-7) for patients recruited before 01.01.2018 compared to 7 (4-9.75) for patients recruited after 01.01.2018 ( $p=0.003$ ).

**Table S2:** Univariate analyses of differences between physiotherapy patients recruited before and after 01.01.2018.

	Physio before (n=90)	Physio after (n=40)	p-value
Age, median (IQR)	68 (62.75, 73)	68.5 (61.5, 76)	0.323
Sex female, n (%)	53 (58.9)	17 (42.5)	0.084
Married or living with partner, n (%)	69 (76.7)	29 (72.5)	0.580
Paid work, n (%)	30 (33.3)	12 (30.0)	0.606
Education level			0.317
- Low (elementary+high school)	51 (56.7)	19 (47.5)	
- High (university+ uni 4+)	39 (43.3)	21 (52.5)	
Health-related quality of life			
- Mental sumscore, median (IQR)	56.29 (51.01, 60.99)	54.63 (47.35, 60.37)	0.396
- Physical sumscore, mean (SD)	40.61 (7.91)	40.67 (8.30)	0.969
Hazardous alcohol consumption, n (%)	44 (48.9)	21 (52.5)	0.786
Smoking status			0.202
- Current smoker	9 (10)	4 (10)	
- Previous	46 (51.1)	14 (35)	
- Never	35 (38.9)	22 (55)	
Number of comorbidities, median (IQR)	2 (1, 2.25)	1 (1, 2)	0.235
BMI, median (IQR)	26.60 (24.41, 30.47)	26.37 (24.60, 29.27)	0.913



1				
2				
3	Fall last 6 weeks, n (%)	18 (20)	7 (17.5)	0.623
4	Falls self-efficacy, median (IQR)	20 (18, 23.35)	22.5 (17, 26.9)	0.424
5	Widespread pain, n (%)	5 (5.6)	2 (5.0)	0.880
6	Previous back pain, n (%)			0.479
7	- Monthly	35 (38.9)	11 (27.5)	
8	- Every year	30 (33.3)	14 (35.0)	
9	- Every 1-5 years	13 (14.4)	5 (12.5)	
10	- Every five years	8 (8.9)	8 (20.0)	
11	- Only once	4 (4.4)	2 (5.0)	
12	Duration of current episode, n (%)			0.538
13	- 0-6 weeks	30 (33.3)	11 (27.5)	
14	- 6 weeks to 3 months	17 (18.9)	11 (27.5)	
15	- 3 months or over	43 (47.8)	19 (47.5)	
16	Back pain, mean (SD)	5.22 (2.53)	4.69 (1.87)	0.208
17	Back-related disability, RMDQ, median (IQR)	8 (6, 13)	9.5 (4.25, 14)	0.808
18	Sleep problems due to back pain, n (%)			0.374
19	- Weekly	36 (40)	13 (32.5)	
20	- Less than weekly	54 (60)	27 (67.5)	
21	Morning stiffness, n (%)			0.753
22	- Significant or extreme	35 (38.9)	16 (40)	
23	- Moderate	35 (38.9)	13 (32.5)	
24	- Some or none	20 (22.2)	11 (27.5)	
25	Walking distance, n (%)			0.285
26	- More than 3km	40 (44.4)	16 (40.0)	
27	- 200m to 3km	41 (45.6)	16 (40.0)	
28	- Less than 200m	9 (10)	8 (20.0)	
29	Kinesiophobia (FABQ-PA), median (IQR)	10 (5, 15)	10.5 (5, 14)	0.842
30	Depression (CES-D), median (IQR)	8 (3.75, 14)	9.5 (5.25, 17.3)	0.305
31	Pain catastrophizing (PCS), median (IQR)	12 (5.3, 17)	11 (4, 19.6)	0.872
32	Back beliefs (BBQ), mean (SD)	30.3 (6.8)	27.3 (7.5)	<b>0.03</b>
33	Expectations for back pain next 3 months			0.821
34	- Fully recovered	17 (18.9)	7 (17.5)	
35	- Much better	50 (55.5)	21 (52.5)	
36	- No change or worse	23 (25.5)	12 (30.0)	
37	SBT risk profiles			0.163
38	- Low	68 (75.5)	24 (60)	
39	- Medium	18 (20)	14 (35)	
40	- High	4 (4.4)	2 (5)	
41	Physical performance (BPS), median (range)	5 (2, 7)	7 (4, 9.75)	<b>0.003</b>
42	Timed up and go, median (IQR)	7.99 (6.66, 9.18)	7.42 (6.64, 9.86)	0.655
43	Probable nerve root involvement, n (%)	20 (22.2)	13 (32.5)	0.194
44	Number of red flags, median (range)	1 (0, 2)	1 (0, 2)	0.815
45	Pain on active range of motion, n (%)	61 (67.8)	27 (67.5)	0.905

IQR: Interquartile range; SD: Standard deviation; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale – International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire – Physical Activity subscale; CES-D: Center for Epidemiological Studies – Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.  
AUDIT-C scores of  $\geq 3/12$  for women and  $\geq 4/12$  indicates hazardous alcohol consumption

## Sensitivity analyses S3, only low back pain patients:

### Methods:

Multiple multinomial regression. One model per variable block. The chiropractic group is the reference group. For these analyses, 382 patients were available; 106 GP patients, 109 physiotherapy patients, and 167 chiropractic patients.

### Results:

See Table S3 for details. Overall, there were very few substantial changes in point estimates and p-values compared to the analyses of all included patients in the article main body. SF-36 physical component summary score was no longer significantly associated with first visiting a GP. Having widespread pain was significantly associated with visiting a physiotherapist compared to a chiropractor. Although point estimates for back-related disability was identical, it was no longer significantly associated with visiting a GP or a physiotherapist. For the Start Back Screening Tool, medium risk category was no longer significantly associated with visiting a chiropractor compared to a physiotherapist, but high risk was significant. Having a positive diagnostic rule for radiculopathy was significantly associated with visiting a physiotherapist compared to a chiropractor.

**Table S3:** Subgroup analyses of the multinomial regression analyses for patients with low back pain only. Chiropractic group (n=167) is the reference group.

<b>Block i) Sociodemographic factors.</b>				
	GP (n=106)		Physio (n=109)	
	Odds ratio (95% CI)	p-value	Odds ratio (95% CI)	p-value
Age	1.03 (0.99, 1.07)	0.23	1.04 (1.00, 1.08)	<b>0.05</b>
Gender				
Female	1.43 (0.86, 2.37)	0.17	1.31 (0.78, 2.19)	0.31
Male (ref)	1.00		1.00	
Marital status				
Married/cohabiting	0.58 (0.30, 1.09)	0.09	0.73 (0.38, 1.40)	0.34
Not married/cohabiting (ref)	1.00		1.00	
Educational level				
Higher education	0.97 (0.58, 1.61)	0.91	1.18 (0.71, 1.96)	0.52
Lower education (ref)	1.00		1.00	
Employment status				
Currently in paid work	0.79 (0.40, 1.55)	0.49	0.63 (0.31, 1.28)	0.20
No paid work (ref)	1.00			
<b>Block ii) General health factors.</b>				
Hazardous alcohol intake (AUDIT-C)				
Yes	1.19 (0.69, 2.05)	0.54	1.18 (0.69, 2.01)	0.54
No (ref)	1.00		1.00	
Smoking				
Yes	1.42 (0.64, 3.19)	0.39	0.64 (0.24, 1.71)	0.37
Previously	1.37 (0.75, 2.47)	0.30	1.02 (0.59, 1.77)	0.95
No (ref)	1.00		1.00	
Health-related quality of life (SF-36, 0-100)				
Physical component	0.97 (0.93, 1.00)	0.08	0.98 (0.94, 1.01)	0.20
Mental component	0.97 (0.94, 1.00)	<b>0.04</b>	1.00 (0.97, 1.03)	0.96
BMI	0.99 (0.93, 1.05)	0.76	0.96 (0.90, 1.02)	0.23
Comorbidities (SCQ, 0-15)	1.13 (0.90, 1.42)	0.29	1.18 (0.96, 1.47)	0.12
Widespread pain				
Yes	0.16 (0.04, 0.65)	<b>0.01</b>	0.30 (0.09, 0.99)	<b>0.05</b>
No (ref)	1.00		1.00	
Falls self-efficacy (FESI, 16-64)	1.00 (0.95, 1.05)	0.99	1.01 (0.95, 1.06)	0.85

<b>Block iii) Current episode and back pain history.</b>					
Back pain severity (NRS, 0-10)	0.98 (0.86, 1.11)	0.73	0.89 (0.78, 1.01)	0.07	
Back-related disability (RMDQ, 0-24)	1.06 (1.00, 1.13)	0.05	1.06 (1.00, 1.13)	0.06	
Duration					
Over 3 months	3.54 (1.42, 8.80)	<b>&lt;0.01</b>	3.85 (1.69, 8.77)	<b>&lt;0.01</b>	
6 weeks to 3 months	3.40 (1.12, 10.37)	<b>0.03</b>	3.25 (1.16, 9.09)	<b>0.03</b>	
0-6 weeks (ref)	1.00		1.00		
Morning stiffness					
Significant or extreme	0.79 (0.39, 1.60)	0.51	1.35 (0.68, 2.67)	0.39	
Moderate	1.63 (0.82, 3.24)	0.16	2.02 (1.02, 4.03)	<b>0.05</b>	
A little or none (ref)	1.00		1.00		
Sleep problems attributable to back pain					
- Weekly	1.13 (0.60, 2.14)	0.70	0.66 (0.34, 1.26)	0.20	
- Less than weekly (ref)	1.00		1.00		
Previous back pain frequency					
- Yearly	1.03 (0.57, 1.87)	0.93	1.04 (0.59, 1.83)	0.88	
- Not yearly (ref)	1.00		1.00		
<b>Block iv) Psychological factors.</b>					
Fear-avoidance (FABQ-PA, 0-24)	1.00 (0.95, 1.05)	0.97	1.03 (0.98, 1.08)	0.31	
Pain catastrophizing (PCS, 0-52)	1.03 (0.99, 1.07)	0.20	1.06 (1.02, 1.10)	<b>&lt;0.01</b>	
Depression symptoms (CESD, 0-60)	0.97 (0.94, 1.03)	0.50	0.99 (0.95, 1.03)	0.70	
Back beliefs (BBQ, 9-45)	0.96 (0.92, 1.01)	0.12	0.99 (0.94, 1.04)	0.63	
Expectation for back pain in 3 months					
Recovered	0.21 (0.09, 0.49)	<b>&lt;0.01</b>	0.34 (0.16, 0.73)	<b>&lt;0.01</b>	
Much better	0.60 (0.31, 1.16)	0.13	0.71 (0.36, 1.39)	0.31	
No change or worse(ref)	1.00		1.00		
Start Back Screening tool					
High risk	1.82 (0.55, 6.05)	0.33	0.19 (0.04, 0.90)	<b>0.04</b>	
Medium risk	1.03 (0.52, 2.06)	0.92	0.59 (0.30, 1.17)	0.13	
Low risk (ref)	1.00		1.00		
<b>Block v) Clinical variables.</b>					
Number of red flags (0-12)	1.28 (0.98, 1.68)	0.07	1.16 (0.90, 1.50)	0.24	
Diagnostic rule for radiculopathy					
Positive	2.32 (1.24, 4.34)	<b>&lt;0.01</b>	1.89 (1.00, 3.57)	<b>0.05</b>	
Negative (ref)	1.00		1.00		
Pain on active range of motion					
Yes	0.88 (0.50, 1.53)	0.64	1.06 (0.62, 1.80)	0.84	
No (ref)	1.00				
Physical performance (BPS, 0-18)	1.19 (1.10, 1.28)	<b>0.03</b>	1.09 (1.01, 1.17)	<b>0.03</b>	
Timed up and go, mean seconds	0.89 (0.78, 1.01)	0.06	0.97 (0.86, 1.09)	0.56	

GP: General practitioner; PT: Physiotherapist; CI: Confidence interval; ref.: reference category; AUDIT-C: Alcohol Use Disorder Identification Test - Consumption questions; SF-36: Short Form Health Survey 36 Item; SCQ: Self-administered Comorbidity Questionnaire; FES-I: Falls Self-Efficacy Scale - International; NRS: Numeric Rating Scale; RMDQ: Roland-Morris Disability Questionnaire; FABQ-PA: Fear-Avoidance Beliefs Questionnaire - Physical Activity subscale; CES-D: Center for Epidemiological Studies - Depression; PCS: Pain Catastrophizing Scale; BBQ: Back Beliefs Questionnaire; BPS: Back Performance Scale.

The odds ratios for continuous variables represent the change in odds with a one-unit increase in the continuous variable.

## Assessment of generalizability S4:

**Table S4:** Descriptive comparison of NORLAG sample and NORLAG 2017 musculoskeletal (MSK) subsample with BACE-N sample.

	NORLAG 2017 subsample MSK conditions <sup>^</sup> (n=794)	BACE-N (n=452)
Age, median (IQR, range)	66 (60-74, 50-93)	66 (59-72, 55-89)
Gender female, n (%)	506 (63.7)	235 (52)
Mother tongue Norwegian (n=432), n (valid %)		412 (95.4)
Country of origin Norway, n (%)	728 (91.7)	
Educational level, n (%)		
- Low (elementary + high school)	566 (71.4)	253 (56.0)
- High (university level)	227 (28.6)	199 (44.0)
In paid work, n (%)	251 (31.6)	205 (45.3)
Living with partner, n (%)	494 (62.2)	347 (76.8)
BMI, mean (SD)	26.3 (4.4)	27.6 (4.7)
How many alcoholic units do you normally drink? <sup>~</sup> n (valid %)		
- 1-2		
- 3-4		
- 5-6	183 (70.1)	289 (63.9)
- 7-9	62 (23.8)	136 (30.1)
- 10 or more	10 (3.8)	22 (4.8)
	1 (0.4)	2 (0.4)
	5 (1.9)	3 (0.7)
How often have you drunk alcohol until you felt intoxicated? (n=433) n, (valid %)		
- Once per week	12 (2.8)	
- 2-3 times per week	3 (0.7)	
- 2-3 times per month	18 (4.2)	
- Once per month	37 (8.5)	
- Rarely	235 (54.3)	
- Never	128 (29.6)	
How often do you drink 6 alcoholic units or more?		
- Almost daily		1 (0.2)
- Some days per week		3 (0.7)
- Some days per month		41 (9.1)
- Rarely		194 (42.9)
- Never		213 (47.1)
CES-D (IQR, range)	8 (4-14, 0-38)	8 (4-15, 0-46)
HR-QoL, physical summary score*, mean (SD)	37.5 (11.3)	41.4 (8.4)
HR-QoL, mental summary score*, mean (SD)	54.7 (8.2)	52.5 (10.0)
Walking distance		
- Cannot walk	13 (1.7)	
- A few steps	22 (2.8)	
- 10-100 m	59 (7.6)	
- 100-500m	57 (7.3)	
- 500m-1km	82 (10.5)	
- 1-5km	235 (30.1)	
- 5km+	313 (40.1)	
Walking distance		
- Less than 15m		20 (0.7)
- 15m-200m		310 (11.5)
- 200m-3km		1130 (42.1)
- 3km+		1218 (45.3)

IQR: Interquartile range; SD; Standard deviation; BMI: Body mass index; CES-D: Center for Epidemiological Studies – Depression questionnaire; HR-QoL: Health-related quality of life

<sup>^</sup>The subsample was collected in 2017 and consisted of participants aged 55 years or older, with at least one musculoskeletal condition

<sup>~</sup> In NORLAG, this variable is continuously, as “number of alcoholic drinks usually drunk per time you drink alcohol”. In BACE-N, it is the AUDIT-C question 2, a categorical question with 5 categories: 1-2, 3-4, 5-6, 7-9 and 10 or more.

\*NORLAG used Short Form Health Survey-12, BACE-N used Short Form Health Survey-36

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	4
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	5
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	5
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5-7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5-7
Bias	9	Describe any efforts to address potential sources of bias	8
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7
		(b) Describe any methods used to examine subgroups and interactions	NA
		(c) Explain how missing data were addressed	7
		(d) If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	8
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9
		(b) Give reasons for non-participation at each stage	-
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	9, 19-20
		(b) Indicate number of participants with missing data for each variable of interest	9, 19-20
Outcome data	15*	Report numbers of outcome events or summary measures	9
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9-10, 20-21

		(b) Report category boundaries when continuous variables were categorized	6-7
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10, suppl.
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	10
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	10-12
Generalisability	21	Discuss the generalisability (external validity) of the study results	12
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).