

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Transcatheter aortic valve implantation outcomes in patients with sarcopenia: protocol for a systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2022-067461
Article Type:	Protocol
Date Submitted by the Author:	21-Aug-2022
Complete List of Authors:	Luo, Kai; Sichuan University, Department of Anesthesiology Yang, Lei; Sichuan University, Department of Anesthesiology; Chinese Academy of Medical Sciences, Department of Anesthesiology Li, Yu; Sichuan University, Department of Anesthesiology; Chinese Academy of Medical Sciences, Department of Anesthesiology
Keywords:	SURGERY, Anaesthesia in cardiology < ANAESTHETICS, Adverse events < THERAPEUTICS

SCHOLARONE™
Manuscripts

1
2
3
4 1 **Transcatheter aortic valve implantation outcomes in patients with sarcopenia: protocol**
5
6 2 **for a systematic review and meta-analysis**
7
8

9 3 Kai Luo¹, Lei Yang^{1,2}, Yu Li^{1,2}
10

11
12 4 ¹Department of Anesthesiology, West China Hospital, Sichuan University, No. 37, Guoxue
13
14 5 Xiang, Chengdu 610041, China
15
16

17 6 ²Department of Anesthesiology, West China Hospital, Sichuan University and The Research
18
19 7 Units of West China, Chinese Academy of Medical Sciences, Chengdu, China.
20
21

22
23 8 Corresponding author
24

25 9 Name: Yu Li
26
27

28 10 Address: Department of Anesthesiology, West China Hospital, Sichuan University, No. 37,
29
30

31 11 Guoxue Xiang, Chengdu 610041, China
32
33

34 12 E-mail: liyu1936@163.com
35

36 13 **Abstract**
37
38

39 14 **Introduction**
40
41

42 15 Sarcopenia, which represents a central biological substratum of frailty and increases the
43
44 16 incidence of adverse events and mortality after elective esophageal cancer surgery, gastrectomy,
45
46 17 and pancreatic surgery, has been recently suggested as a predictor of outcomes in patients
47
48 18 undergoing transcatheter aortic valve implantation (TAVI). However, the results of these
49
50 19 studies are variable, and therefore, we would like to perform a systematic review and meta-
51
52 20 analysis of the current literature to evaluate sarcopenia as a predictor of outcome post-TAVI.
53
54
55
56

57
58 21 **Methods and analysis**
59
60

1
2
3
4 1 Two investigators will search electronic databases independently, including PubMed,
5
6 2 EMBASE, Web of Science, Medline, and The Cochrane Library. In addition, four clinical trial
7
8 3 registries and the related references will be manually retrieved. Included studies will be assessed
9
10 4 for risk of bias according to Cochrane and certainty of evidence using the Grading of
11
12 5 Recommendations Assessment. Subgroup analyses and sensitivity analyses will be performed
13
14 6 if substantial heterogeneity is encountered. Review Manager (version 5.4) software will be used
15
16 7 to perform the whole data analysis. The results will be presented as forest plots, and
17
18 8 heterogeneity between the included studies will be assessed by Cochran's Q and I2 tests.
19
20
21
22
23
24

25 9 **Ethics and dissemination**

26
27
28 10 No ethical approval is needed, as we will recruit from previously published studies in which
29
30 11 informed consent was obtained. The results will be disseminated in a peer-reviewed journal
31
32 12 upon completion.
33
34

35
36 13 PROSPERO registration number CRD42022349525.
37
38

39 14 **Strengths and limitations of this study**

40
41
42 15 Transcatheter aortic valve implantation (TAVI) represents the current mainstay of
43
44 16 treatment for patients affected by severe aortic stenosis (AS). However, preoperative
45
46 17 risk stratification models are imperfect, and the effect of sarcopenia on postoperative
47
48 18 outcomes remains unclear.
49
50

51
52
53 19 To the best of our knowledge, this is the first systematic review and meta-analysis to
54
55 20 estimate the effect of sarcopenia in predicting outcomes post-TAVI.
56
57

58 21 This extensive systematic review and meta-analysis is an applied study based on clinical
59
60

1
2
3
4 1 practice that will provide high-level evidence to evaluate the impact of sarcopenia on
5
6 2 clinical outcomes in patients undergoing TAVI.
7
8

9 3 The gold standard for noninvasive muscle wasting assessment in clinical practice is an
10
11 4 abdominal computed tomography (CT) scan, and the assessment of sarcopenia may
12
13 5 interfere with the final data because its adoption in clinical practice remains limited by
14
15 6 a lack of standardized definition.
16
17

18
19
20 7 Inherent to all meta-analyses is the potential for heterogeneity among the selected
21
22 8 studies. Meanwhile, the measurement of sarcopenia is not uniform, and our analysis
23
24 9 may use several tools that were used to assess muscle areas with a different threshold
25
26 10 to define sarcopenic patients and may interfere with the outcome. However, sensitivity
27
28 11 and subgroup analysis can be used to resolve this disagreement.
29
30
31

32 33 34 12 **Introduction**

35
36 13 Transcatheter aortic valve implantation (TAVI) represents the current mainstay of treatment for
37
38 14 older adults affected by severe aortic stenosis (AS) and those with high or prohibitive surgical
39
40 15 risk[1, 2]. Several clinical, functional, and anatomical predictors of procedural success and
41
42 16 long-term outcomes have been identified[3-5]. To address new treatment targets and improve
43
44 17 current management strategies, sarcopenia has been recently suggested as a predictor of
45
46 18 outcomes in patients undergoing TAVI[6]. For TAVI planning and perioperative preparation,
47
48 19 a computed tomography (CT) scan is routinely performed and may cover the complete thorax
49
50 20 and abdomen. In addition to assessing preoperative cardiopulmonary function, a CT scan can
51
52 21 also be used to measure body composition.
53
54
55
56
57
58
59
60

1 Sarcopenia, which is defined as the loss of skeletal muscle mass and functioning due to aging[7,
2 8], represents a central biological substratum of frailty and reflects a state of declined functional
3 capacity and increased vulnerability to disease, disability, and death[9-11]. The psoas muscle
4 area (PMA) obtained from axial cuts on CT scan, which has been validated as a surrogate for
5 muscle mass[12], low muscle mass as a measure of sarcopenia is easily measured from
6 preoperative CT images. Sarcopenia is considered an independent marker of life quality and
7 prognosis and has been demonstrated to herald adverse outcomes across a range of clinical
8 conditions[9, 10, 13]. Sarcopenia has been observed to be a strong prognostic indicator for
9 perioperative complications and increased short- and long-term postoperative mortality and
10 morbidity[14-16], including cognitive impairment[17], mental disorders[18], acute kidney
11 injury[19, 20], and even survival[16, 21, 22]; meanwhile, the sarcopenia cohort was associated
12 with longer ICU stays and hospital stays[16]. However, the evidence comes mainly from
13 patients undergoing noncardiac surgery, with limited evidence and recommendations for
14 sarcopenia as a predictor of poor outcomes after TAVI.

15 Studies have suggested that sarcopenia is highly prevalent among patients undergoing TAVI,
16 negatively affects important outcomes[23] and is an independent predictor of outcomes in
17 patients undergoing TAVI. Therefore, we will perform further analysis to assess the impact of
18 sarcopenia on mortality and postoperative complications in patients following TAVI.

19 **Methods**

20 This protocol was registered on PROSPERO (CRD42022349525) and will be reported
21 following the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols

1 (PRISMA-P) statement guidelines[24]. The results of this systematic review and meta-
 2 analysis will be published in a specialist journal or presented at a conference. This protocol is
 3 based on existing studies without patients or the public being involved directly.

4 **Search strategy**

5 Articles will be retrieved from five databases, including PubMed, EMBASE, Web of Science,
 6 Medline, and The Cochrane Library, four clinical trial registries (Australia-new Zealand
 7 Clinical Trials Registry, ClinicalTrials.gov, International Registry of Standard Randomized
 8 Controlled Trial Numbers and Chinese Clinical Trial Register), and the related references will
 9 be manually retrieved. The following search algorithm and Medical Subject Headings (MeSH)
 10 terms will be used: [(sarcopenia) and (TAVI)]. A specific search strategy is presented in Table
 11 1 using PubMed as an example.

Table 1 Search strategy of PubMed

Search	Query
#1	"Sarcopenia"[MeSH Terms]
#2	"Sarcopenia"[All Fields] OR "Sarcopenias"[All Fields] OR "Muscle mass"[All Fields] OR "Muscle area"[All Fields]
#3	#1 OR #2
#4	"Transcatheter aortic valve implantation"[MeSH Terms]
#5	"Transcatheter aortic valve implantation"[All Fields] OR "Transcatheter aortic valve replacement"[All Fields] OR "TAVI"[All Fields] OR "TAVR"[All Fields]
#6	#4 OR #5

#7 **#3 AND #6**

1 Search results will be exported to an Endnote database. Two investigators (L K and Y L) will
2 each independently search the database and read the titles/abstracts of all the records, and full-
3 text articles will be retrieved if they are judged potentially eligible by an investigator. Both
4 investigators will evaluate the full texts of all the potentially eligible retrieved articles, and the
5 article will be excluded for an explicit reason. The two researchers will cross-check the data
6 after completion of data extraction. Irreconcilable disagreements between the two researchers
7 will be resolved by consensus with a third independent reviewer (L Y). The study flow is
8 presented in Fig 1.

9 Inclusion and exclusion criteria

10 The inclusion criteria

- 11 (1) adult patients aged 18 or older;
- 12 (2) randomized controlled trials or observational studies (cohort, cross-sectional, or
13 case-control studies) published as original articles that evaluated and compared the effect of
14 sarcopenia in patients undergoing TAVI;
- 15 (3) a reference group composed of nonsarcopenic patients;
- 16 (4) Sufficient quantitative data, such as contingency tables, ORs, relative risks, or hazard
17 ratios with 95% confidence intervals (CIs), were provided;

18 The exclusion criteria:

- 19 (1) unfinished or ongoing studies;
- 20 (2) Case reports, comments, animal studies, meta-analyses and systematic reviews.

1 **Outcomes**

2 The primary outcome is short-term mortality (30-day mortality). Secondary outcomes are
3 long-term mortality (> 30 days), length of ICU stay, need for ICU admission (the number of
4 patients in the sarcopenia or nonsarcopenia group needs ICU admission), length of hospital
5 stay and total complications.

6 **Data collection and bias assessment**

7 Two investigators will collect the following items: authors, publication year, age, BMI, ASA
8 classification, sample size, diagnostic criteria for sarcopenia, and characteristics of the
9 included population. If the outcome data were presented in the form of a graph and could not
10 be directly extracted, we will use a plot digitizer or contact the corresponding author.

11 Two investigators (L K and Y L) will independently use the Quality in Prognosis Studies
12 (QUIPS) critical assessment tool to assess the risk of bias for the included studies[25, 26].

13 The scale mainly includes study participation, study attrition, prognostic factor measurement,
14 outcome measurement, study confounding, and statistical analysis and reporting. Each
15 domain is assessed against criteria, thereby resulting in a rating of 'high', 'moderate', or 'low'
16 risk of bias. Any discrepancies in the assessment will be resolved by consensus or third-
17 author arbitration.

18 GRADE (Grading of Recommendations, Assessment, Development, and Evaluation)[27] will
19 be used to assess the quality of evidence for short- and long-term mortality, length of ICU
20 stay, need for ICU admission, length of hospital stay and complications. We will rate the
21 quality of the evidence as 'high', 'moderate', 'low', and 'very low' based on the risk of bias,

1 inconsistency, indirectness, imprecision, and other considerations.

2 **Statistical analysis**

3 Review Manager (version 5.4) software will be used to perform the whole data analysis. The
4 mean difference (MD) with 95% confidence intervals (CI) will be chosen to calculate
5 continuous variables, while risk ratios (RR) with the same CI will be employed to present
6 dichotomous outcomes. Continuous results will be presented as the mean and standard
7 deviation, and if the median is displayed, we will convert the median and interquartile range
8 to the mean and standard deviation using the statistical formula[28, 29]. Statistical
9 heterogeneity across studies will be evaluated with a corresponding *P* value derived from the
10 chi-square test and I^2 test, and $I^2 > 50\%$ or $P < 0.10$ is regarded as significantly heterogeneous.
11 If the included studies are homogeneous ($p \geq 0.10$ and $I^2 < 50\%$), we will use the fixed-effects
12 model to estimate the outcome data. A random-effect model will be used to aggregate the data
13 owing to significant heterogeneity, and we will identify any potential sources of heterogeneity
14 and conduct subgroup analysis. Subgroup analysis and meta-regression analysis will be
15 conducted to explore the possible sources of heterogeneity and inconsistency, such as
16 participants' mean age, diagnosis of sarcopenia and race. In addition, leave-one-out jackknife
17 sensitivity analysis will be conducted by removing one study at a time before repeating the
18 analysis to verify the robustness of the results.

19 **Subgroup and sensitivity analysis**

20 Subgroup analyses may be performed depending on the number of studies identified. The
21 level of heterogeneity across included studies and strength of evidence for heterogeneity will

1
2
3
4 1 be examined using the Cochrane Q and I² statistics with associated 95% CI, an I² of 50% and
5
6 2 above is considered a substantial level of heterogeneity. The random effect models will be
7
8
9 3 used because of potential heterogeneity between study variations in population, regions or
10
11
12 4 assessment methods of sarcopenia across studies, or the fixed effect model will be used if
13
14
15 5 tests of heterogeneity among studies are not significant. A range of sensitivity analyses may
16
17
18 6 be conducted to examine the methodological quality and potential sources of heterogeneity of
19
20
21 7 the included studies. Sources of variations may include tool for assessment of sarcopenia, age,
22
23
24 8 preoperative comorbidities and the level of care from different centers. These will be stratified
25
26
27 9 and separate sensitivity analyses conducted, leave-one-out jackknife sensitivity analysis will
28
29
30 10 be used to verify the robustness of the results. A minimum of two studies are generally
31
32
33 11 considered sufficient to perform a meta-analysis, if the level of heterogeneity is high between
34
35
36 12 studies and pooled analysis of the studies is not possible, a narrative summary of the outcome
37
38
39 13 of the selected studies will be presented in a systematic review and report the reasons for the
40
41
42 14 results.

15 **Patient and public involvement**

16 No patients or the public were directly involved in the design, writing or editing of this
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

18 **Discussion**

19 This is the first study to demonstrate that body composition analysis using pre-TAVI CT is
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 of sarcopenia on outcomes post-TAVI to provide important predictive information for general
2 medical management strategies, nursing goals, and rehabilitation expectations.

3 Our study will provide a reliable basis for clinical practice. As pre-TAVI CT scans are currently
4 routinely performed, they may provide important complementary information for sarcopenia
5 assessment. On the one hand, it highlights sarcopenia as a risk predictor in TAVI patients and
6 evaluates sarcopenia as an objective, readily available tool to be introduced in clinical practice.
7 On the other hand, sarcopenia as a risk predictor suggests a potential role for the correction of
8 sarcopenia to improve post-TAVI outcomes, such as protein supplementation and exercise
9 training.

10 Limitations exist in this meta-analysis. First, the measurement of muscle mass and the definition
11 of sarcopenia are not uniform and will inevitably have a negative impact on the outcomes. In
12 addition, more than 90% of all TAVI cases are currently performed via the transfemoral route.
13 To avoid protracted recovery associated with thoracic access, transapical vascular access and
14 subclavian, carotid, and transcarotid routes are usually favored. However, we can attempt to
15 solve this problem by subgroup analysis if sufficient studies are included. Meanwhile,
16 publication bias was unavoidable, as we selected published articles.

17 **Ethics statements**

18 Patient consent for publication: Not applicable.

19 **Funding statement**

20 This research received no specific grant from any funding agency in the public, commercial or
21 not-for-profit sectors.

1 Contributions:

2 L K registered the methods on the PROSPERO website and drafted the manuscript. The
 3 manuscript was subsequently revised for critical intellectual content by L K, Y L, and L Y.

4 Competing Interest statemen

5 No competing interest

7 References

- 8 1. Mack M J, M B Leon, V H Thourani, et al. *Transcatheter Aortic-Valve Replacement with a Balloon-Expandable Valve in Low-Risk Patients*. N Engl J Med, 2019; **380**(18): 1695-1705.
- 9 2. Leon M B, C R Smith, M J Mack, et al. *Transcatheter or Surgical Aortic-Valve Replacement in Intermediate-Risk Patients*. N Engl J Med, 2016; **374**(17): 1609-20.
- 10 3. Gallone G, A Depaoli, F D'Ascenzo, et al. *Impact of computed-tomography defined sarcopenia on outcomes of older adults undergoing transcatheter aortic valve implantation*. J Cardiovasc Comput Tomogr, 2022; **16**(3): 207-214.
- 11 4. Holmes D R, Jr., J M Brennan, et al. *Clinical outcomes at 1 year following transcatheter aortic valve replacement*. Jama, 2015; **313**(10): 1019-28.
- 12 5. Debonnaire P, L Fusini, R Wolterbeek, et al. *Value of the "TAVI2-SCORE" versus surgical risk scores for prediction of one year mortality in 511 patients who underwent transcatheter aortic valve implantation*. Am J Cardiol, 2015; **115**(2): 234-42.
- 13 6. Soud M, F Alahdab, G Ho, et al. *Usefulness of skeletal muscle area detected by computed tomography to predict mortality in patients undergoing transcatheter aortic valve replacement: a meta-analysis study*. Int J Cardiovasc Imaging, 2019; **35**(6):1141-1147.
- 14 7. Cruz-Jentoft A J, J P Baeyens, J M Bauer, et al. *Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People*. Age Ageing, 2010; **39**(4): 412-23.
- 15 8. Dodds R M.A A Sayer, *Sarcopenia, frailty and mortality: the evidence is growing*. Age Ageing, 2016; **45**(5): 570-1.
- 16 9. Cruz-Jentoft A J, G Bahat, J Bauer, et al. *Sarcopenia: revised European consensus on definition and diagnosis*. Age Ageing, 2019; **48**(1):16-31.
- 17 10. Jones K, A Gordon-Weeks, C Coleman.M Silva, *Radiologically Determined Sarcopenia Predicts Morbidity and Mortality Following Abdominal Surgery: A Systematic Review and Meta-Analysis*. World J Surg, 2017; **41**(9): 2266-2279.
- 18 11. Chang S F.P L Lin, *Systematic Literature Review and Meta-Analysis of the Association of Sarcopenia With Mortality*. Worldviews Evid Based Nurs, 2016; **13**(2):

- 153-62.
12. Boutin R D, L Yao, R J Canter, L Lenchik, *Sarcopenia: Current Concepts and Imaging Implications*. AJR Am J Roentgenol, 2015; **205**(3): W255-66.
 13. ter Maaten J M, K Damman, H L Hillege, et al. *Creatinine excretion rate, a marker of muscle mass, is related to clinical outcome in patients with chronic systolic heart failure*. Clin Res Cardiol, 2014; **103**(12): 976-83.
 14. Weerink L B M, A van der Hoorn, B L van Leeuwen, G H de Bock, *Low skeletal muscle mass and postoperative morbidity in surgical oncology: a systematic review and meta-analysis*. J Cachexia Sarcopenia Muscle, 2020; **11**(3): 636-649.
 15. Aleixo G F P, S S Shachar, K A Nyrop, et al. *Myosteatosis and prognosis in cancer: Systematic review and meta-analysis*. Crit Rev Oncol Hematol, 2020; **145**: 102839.
 16. Yang T R, K Luo, X Deng, et al. *Effect of sarcopenia in predicting postoperative mortality in emergency laparotomy: a systematic review and meta-analysis*. World J Emerg Surg, 2022; **17**(1): 36.
 17. Chang K V, T H Hsu, W T Wu, et al. *Association Between Sarcopenia and Cognitive Impairment: A Systematic Review and Meta-Analysis*. J Am Med Dir Assoc, 2016; **17**(12): 1164.e7-1164.e15.
 18. Chang K V, T H Hsu, W T Wu, et al. *Is sarcopenia associated with depression? A systematic review and meta-analysis of observational studies*. Age Ageing, 2017; **46**(5): 738-746.
 19. Bang J Y, I G Jun, J B Lee, et al. *Impact of Sarcopenia on Acute Kidney Injury after Infrarenal Abdominal Aortic Aneurysm Surgery: A Propensity Matching Analysis*. Nutrients, 2021. **13**(7).
 20. Sim J H, H M Kwon, I G Jun, S H Kim, K S Kim, Y J Moon, J G Song, G S Hwang, *Association of skeletal muscle index with postoperative acute kidney injury in living donor hepatectomy: A retrospective single-centre cohort study*. Liver Int, 2022; **42**(2): 425-434.
 21. Francomacaro L M, C Walker, K Jaap, et al. *Sarcopenia predicts poor outcomes in urgent exploratory laparotomy*. Am J Surg, 2018; **216**(6): 1107-1113.
 22. McQuade C, D O Kavanagh, C O'Brien, et al. *CT-determined sarcopenia as a predictor of post-operative outcomes in patients undergoing an emergency laparotomy*. Clin Imaging, 2021; **79**: 273-277.
 23. Bertschi D, C M Kiss, A W Schoenenberger, et al. *Sarcopenia in Patients Undergoing Transcatheter Aortic Valve Implantation (TAVI): A Systematic Review of the Literature*. J Nutr Health Aging, 2021; **25**(1): 64-70.
 24. Shamseer L, D Moher, M Clarke, D Ghersi, et al. *Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation*. Bmj, 2015; **350**: g7647.
 25. Hayden J A, P Côté, C Bombardier, *Evaluation of the quality of prognosis studies in systematic reviews*. Ann Intern Med, 2006; **144**(6): 427-37.
 26. Hayden J A, D A van der Windt, J L Cartwright, et al. *Assessing bias in studies of prognostic factors*. Ann Intern Med, 2013; **158**(4): 280-6.

- 1
2
3 1 27. Brozek J L, E A Akl, P Alonso-Coello, et al. *Grading quality of evidence and strength*
4 2 *of recommendations in clinical practice guidelines. Part 1 of 3. An overview of the*
5 3 *GRADE approach and grading quality of evidence about interventions.* Allergy, 2009;
6 4 **64**(5): 669-77.
7 5 28. Luo D, X Wan, J Liu.T Tong, *Optimally estimating the sample mean from the sample*
8 6 *size, median, mid-range, and/or mid-quartile range.* Stat Methods Med Res, 2018;
9 7 **27**(6): 1785-1805.
10 8 29. Wan X, W Wang, J Liu.T Tong, *Estimating the sample mean and standard deviation*
11 9 *from the sample size, median, range and/or interquartile range.* BMC Med Res
12 10 *Methodol*, 2014; **14**: 135.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

Figure 1 is provided without any warranty of any kind, either express or implied, the flow chart will be used to document the process of selecting articles for inclusion in this systematic review and meta-analysis.




Fig 1. Flowchart showing selection of articles for review



PRISMA 2020 Checklist

Section and Topic	Item #	Checklist item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review.	Page 1
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist.	Page 1
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	Page 1
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	Page 1
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	Page 6
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	Page 5
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	Page 5
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	Page 5 to Page 6
Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	Page 7
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	Page 7 to Page 8
	10b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	Page 7 to Page 8
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	Page 8
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	Page 7 to Page 8
Synthesis methods	13a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item #5)).	Page 8
	13b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	Page 8
	13c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	Page 2
	13d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	Page 7 to Page 8
	13e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	Page 8
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	Page 8
Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	No
Certainty	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	Page 7



PRISMA 2020 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Section and Topic	Item #	Checklist item	Location where item is reported
assessment			
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	Page 5 to Page 6
	16b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	Page 6
Study characteristics	17	Cite each included study and present its characteristics.	Not applicable
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	Not applicable
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	Not applicable
Results of syntheses	20a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	Not applicable
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	Not applicable
	20c	Present results of all investigations of possible causes of heterogeneity among study results.	Not applicable
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	Not applicable
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	Not applicable
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	Not applicable
DISCUSSION			
Discussion	23a	Provide a general interpretation of the results in the context of other evidence.	Not applicable
	23b	Discuss any limitations of the evidence included in the review.	Not applicable
	23c	Discuss any limitations of the review processes used.	Page 9
	23d	Discuss implications of the results for practice, policy, and future research.	Page 9
OTHER INFORMATION			
Registration and protocol	24a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	Page 2
	24b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	Page 5
	24c	Describe and explain any amendments to information provided at registration or in the protocol.	Page 5
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	Page 9
Competing interests	26	Declare any competing interests of review authors. For peer review only - http://bmjopen.bmj.com/site/about/guidelines.xhtml	Page 9



PRISMA 2020 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Section and Topic	Item #	Checklist item	Location where item is reported
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	Page 5

For peer review only

36/bmjopen-2022-067461 on 30 November 2022. Downloaded from <http://bmjopen.bmj.com/> on April 20, 2024 by guest. Protected by copyright.

PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification	1a	Identify the report as a protocol of a systematic review (✓)
Update	1b	If the protocol is for an update of a previous systematic review, identify as such (×)
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number (✓)
Authors:		
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author (✓)
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review (✓)
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments (×)
Support:		
Sources	5a	Indicate sources of financial or other support for the review (✓)
Sponsor	5b	Provide name for the review funder and/or sponsor (×)
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol (×)
INTRODUCTION		
Rationale	6	Describe the rationale for the review in the context of what is already known (✓)
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO) (✓)
METHODS		
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review (✓)
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage (✓)
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated (✓)

Study records:		
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review (✓)
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis) (✓)
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms done independently, in duplicate), any processes for obtaining and confirming data from investigators (✓)
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications (✓)
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale (✓)
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis (✓)
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised (✓)
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I^2 , Kendall's τ) (✓)
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) (✓)
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned (×)
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies) (✓)
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE) (✓)

✓: The items has been presented in the manuscript

×: Not applicable

BMJ Open

Sarcopenia as a predictor of outcome after transcatheter aortic valve implantation: protocol for systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2022-067461.R1
Article Type:	Protocol
Date Submitted by the Author:	04-Nov-2022
Complete List of Authors:	Luo, Kai; Sichuan University, Department of Anesthesiology Yang, Lei; Sichuan University, Department of Anesthesiology; Chinese Academy of Medical Sciences, Department of Anesthesiology Li, Yu; Sichuan University, Department of Anesthesiology; Chinese Academy of Medical Sciences, Department of Anesthesiology
Primary Subject Heading:	Surgery
Secondary Subject Heading:	Surgery, Anaesthesia
Keywords:	SURGERY, Anaesthesia in cardiology < ANAESTHETICS, Adverse events < THERAPEUTICS

SCHOLARONE™
Manuscripts

1
2
3
4 **Sarcopenia as a predictor of outcome after transcatheter aortic valve**
5
6 **implantation: protocol for systematic review and meta-analysis**
7
8

9 Kai Luo¹, Lei Yang^{1,2}, Yu Li^{1,2}
10

11 ¹Department of Anesthesiology, West China Hospital, Sichuan University, No. 37,
12
13 Guoxue Xiang, Chengdu 610041, China.
14

15
16 ²Department of Anesthesiology, West China Hospital, Sichuan University and The
17
18 Research Units of West China, Chinese Academy of Medical Sciences, Chengdu,
19
20
21
22 China.
23

24
25
26
27 Correspondence to:
28

29
30 Yu Li
31

32 Department of Anesthesiology, West China Hospital, Sichuan University, No. 37,
33

34
35 Guoxue Xiang, Chengdu 610041, China
36

37
38 liyu1936@163.com
39
40
41
42

43 **Abstract**
44

45 **Introduction**
46

47
48 Sarcopenia represents a central biological substratum of frailty, which increases the
49
50 incidence of adverse events and mortality after surgery for esophageal cancer,
51
52 gastrectomy, and pancreatic surgery. Recently, sarcopenia has been suggested as a
53
54 predictor of outcomes in patients undergoing transcatheter aortic valve implantation
55
56 (TAVI). However, since relevant data were variable, we aimed to perform a systematic
57
58
59
60

1
2
3
4 review and meta-analysis of the current literature to evaluate sarcopenia as a predictor
5
6 of post-TAVI outcomes.
7

8 9 **Methods and analysis**

10
11 Two investigators will conduct independent searches in PubMed, EMBASE, Web of
12
13 Science, MEDLINE, and the Cochrane Library, from database inception until October
14
15 2022. The search will not be limited by language or region. Eligible studies will include
16
17 reports investigating post-TAVI outcomes in patients with sarcopenia, who are
18
19 aged >18 years and diagnosed using a CT scan. The primary outcome is short-term
20
21 mortality (30-day mortality), while the secondary outcomes include long-term mortality
22
23 (>30 days), length of ICU stay, need for ICU admission (the number of patients in the
24
25 sarcopenia or non-sarcopenia group requiring ICU admission), length of hospital stay,
26
27 and overall complications. Included studies will be assessed for risk of bias according
28
29 to the Quality in Prognosis Studies (QUIPS) critical assessment tool and certainty of
30
31 evidence using the GRADE. The analysis will be done with Review Manager (version
32
33 5.4) software. If testing reveals little or no statistical heterogeneity, a fixed-effect model
34
35 will be used for data synthesis; otherwise, a random-effect model may be employed.
36
37 Upon encountering substantial heterogeneity, subgroup analysis and leave-one-out
38
39 jackknife sensitivity analysis will be used to verify the robustness of the results. The
40
41 obtained results will be presented as forest plots while Cochran's Q test and I² test will
42
43 be used to calculate the heterogeneity (>50% indicating strong heterogeneity).
44
45
46
47
48
49
50
51
52
53
54

55 56 **Ethics and dissemination**

57
58 No ethical approval is needed for this study since we will be using data from previously
59
60

published studies. The results will be disseminated in a peer-reviewed journal.

PROSPERO registration number: CRD42022349525.

Strengths and limitations of this study

- This systematic review and meta-analysis will synthesise evidence regarding the effect of sarcopenia on outcomes after transcatheter aortic valve implantation.
- We will strictly follow the guidelines for the conduct and reporting of systematic reviews and meta-analyses to minimize bias.
- We aim be as thorough as possible, including conducting sensitivity and subgroup analyses.
- We will use the Quality in Prognosis Studies (QUIPS) critical assessment tool to evaluate the methodological quality of included studies, while GRADE will be used to determine the certainty of the evidence.
- The included studies may have high heterogeneity in terms of methods and results, which could impact the robustness of the review findings.

Introduction

Transcatheter aortic valve implantation (TAVI) remains the current treatment for older adults affected by severe aortic stenosis (AS) and those with high or prohibitive surgical risk [1, 2]. Previously, several clinical, functional, and anatomical predictors were identified for procedural success and long-term outcomes [3–5]. To address new

1
2
3
4 treatment targets and improve current management strategies, sarcopenia was recently
5
6 suggested as a predictor of outcomes in patients undergoing TAVI [6].
7

8
9 Sarcopenia is highly prevalent among patients undergoing TAVI and is reported in a
10
11 vast majority of men (80%) compared to women, who exhibited approximately half of
12
13 the value (47%). Of the patients who underwent TAVI with a BMI <25, 73% had
14
15 sarcopenia, which was considered a predictor of mortality [7]. In patients undergoing
16
17 transcatheter aortic valve implantation (TAVI) for severe aortic stenosis, skeletal
18
19 muscle mass was investigated as a marker of sarcopenia. Although it is technically easy
20
21 and reproducible to measure Psoas Muscle Area (PMA) from CT, it is not readily
22
23 available in many clinics. Hence, a pre-procedural scan of the chest, abdomen, and
24
25 pelvis is routinely performed among TAVI candidates for interventional planning [3].
26
27
28
29

30
31 Sarcopenia is defined as the loss of skeletal muscle mass and functioning due to aging
32
33 [8, 9], which represents a central biological substratum of frailty reflecting a state of
34
35 declined functional capacity, and increased vulnerability to disease, disability, and
36
37 ultimately death [10–12]. The PMA obtained from axial cuts on a CT scan was
38
39 validated as a surrogate for muscle mass [13]. A low muscle mass is a measure of
40
41 sarcopenia, which can be easily determined through preoperative CT images.
42
43
44
45
46
47
48 Sarcopenia is demonstrated to herald adverse outcomes across a range of clinical
49
50 conditions and is considered an independent marker of life quality and disease
51
52 prognosis [10, 11, 14]. Sarcopenia was observed to be a strong prognostic indicator for
53
54 perioperative complications, which increased short- and long-term postoperative
55
56 mortality and morbidity [15–17], including cognitive impairment [18], mental disorders
57
58
59
60

1
2
3
4 [19], acute kidney injury [20, 21], and even survival [17, 22, 23]. Moreover, the
5
6 sarcopenia cohort was associated with longer ICU and hospital stays [17], whose
7
8 evidence was mainly observed in patients undergoing non-cardiac surgery. Yet, no full
9
10 consensus has been achieved on sarcopenia being a predictor of poor outcomes after
11
12 TAVI. Studies suggested that sarcopenia was associated with longer hospital stay (LOS)
13
14 and worse one-year health-related quality of life post-TAVI [24]. However, Guglielmo
15
16 et al. believed that sarcopenia overestimated additive prognostic value over current
17
18 post-TAVI mortality risk estimators; for example, it was shown that early safety,
19
20 clinical efficacy, and 30-day all-cause death remained unaffected by sarcopenia [3].
21
22 Sarcopenia can occur secondary to a systemic disease, especially one that invokes
23
24 inflammatory processes, e.g., malignancy or organ failure. Furthermore, sarcopenia can
25
26 develop as a result of inadequate intake of energy or protein, probably due to anorexia,
27
28 malabsorption, limited access to healthy foods, or limited ability to eat [10]. Michael
29
30 reported that patients who underwent TAVI with sarcopenia showed a negative effect
31
32 on important outcomes [25], increasing mortality or prolonging the length of hospital
33
34 stay. However, the conclusion stating that sarcopenia negatively affected the outcomes
35
36 of TAVI patients is cursory to be accepted since a few previous studies did not use the
37
38 revised guideline-based criteria for diagnosing the disease. Additionally, the qualitative
39
40 systematic review of the included studies was previously performed without a meta-
41
42 analysis. Therefore, we aim to perform further analysis to assess the impact of
43
44 sarcopenia on mortality and postoperative complications in patients after TAVI.
45
46
47
48
49
50
51
52
53
54
55
56
57

58 **Methods and analysis**

59
60

1
2
3
4 This study was registered on PROSPERO (CRD42022349525) and this protocol was
5
6 prepared according to the guidelines of the Preferred Reporting Items for Systematic
7
8 Review and Meta-Analysis Protocols (PRISMA-P) statement [26]. The results of this
9
10 systematic review and meta-analysis will be reported in accordance with PRISMA
11
12 guidance.
13
14
15

16 17 **Search strategy**

18
19 Articles will be retrieved from five databases, including PubMed, EMBASE, Web of
20
21 Science, MEDLINE, and the Cochrane Library, from database inception until October
22
23 2022. The search will not be limited by language or region. Relevant publications will
24
25 be searched manually for the reference lists of the collected studies, including grey
26
27 literature, e.g., conference articles. Studies will first be screened independently using
28
29 the title/abstract and then in full versions to determine suitability for their inclusion in
30
31 a review. During the search process, the main search terms will be combined with a free
32
33 word search. The following search algorithm and Medical Subject Headings (MeSH)
34
35 terms will be used: [(Sarcopenia) and (TAVI)]. Additional details regarding search
36
37 strategies are provided in the online supplemental appendix.
38
39
40
41
42
43
44

45 The obtained search results will be exported to the endnote database. Two investigators
46
47 (LK and YL) will each independently search the databases and screen the
48
49 titles/abstracts of all records. Full-text articles potentially judged as eligible by an
50
51 investigator will be retrieved and evaluated by both investigators. An article will be
52
53 excluded only for an explicit reason. After the completion of data extraction, both
54
55 researchers will cross-check the data. If there is any irreconcilable disagreement
56
57
58
59
60

1
2
3
4 between the two researchers, it will be resolved by consensus, with input from a third
5
6 independent reviewer (LY). The template study flowchart is presented in Figure 1.
7
8

9 **Inclusion and exclusion criteria**

10
11 The inclusion criteria will be as follows:

- 12
13
14 (1) adult patients aged 18 or older;
- 15
16
17 (2) original published articles, including randomized controlled trials (RCTs) or
18
19 observational studies (cohort, cross-sectional, or case-control studies) evaluating and
20
21 comparing the effect of sarcopenia on patients undergoing TAVI;
- 22
23
24 (3) definition of sarcopenia being supported by the Asian Working Group for
25
26 Sarcopenia (AWGS) [27] and the European Working Group for Sarcopenia in Older
27
28 People (EWGSOP1) [8] or EWGSOP2 [10].
- 29
30
31 (4) a reference group including non-sarcopenic patients;
- 32
33
34 (5) having sufficient quantitative data, including contingency tables, ORs, relative risks,
35
36 or hazard ratios with 95% confidence intervals (CIs).
37
38
39

40 The exclusion criteria will be as follows:

- 41
42
43 (1) unfinished or ongoing studies;
- 44
45
46 (2) case reports, comments, animal studies, meta-analyses, and systematic reviews.
47

48 **Outcomes**

49
50
51 The primary outcome of this study is short-term mortality (30-day mortality), while the
52
53 secondary outcomes will include long-term mortality (>30 days), length of ICU stay,
54
55 need for ICU admission (the number of patients in the sarcopenia or non-sarcopenia
56
57 group requiring ICU admission), length of hospital stay, and overall complications
58
59
60

1
2
3
4 (myocardial infarction, stroke, bleeding complications, acute kidney injury, vascular
5
6 complications, conduction disturbances, and arrhythmias).
7
8

9 **Data collection and bias assessment**

10
11 Both investigators will collect the following items from the included studies: authors,
12
13 publication year, age, BMI, ASA classification, sample size, diagnostic criteria for
14
15 sarcopenia, and characteristics of the included population. In case the presentation of
16
17 the outcome data is in the form of a graph and can not be directly extracted, a plot
18
19 digitizer will be used, or the corresponding author will be contacted.
20
21
22
23

24
25 Two investigators (LK and YL) will independently use the Quality in Prognosis Studies
26
27 (QUIPS) critical assessment tool and assess the bias risk in the included studies [28,
28
29 29]. The scale mainly includes study participation, study attrition, prognostic factor
30
31 measurement, outcome measurement, study confounding, and statistical analysis and
32
33 reporting. Each domain will be assessed against the criteria, resulting in a rating of
34
35 either 'high', 'moderate', or 'low' risk of bias. Any discrepancies in the assessment can
36
37 be resolved by consensus or third-author arbitration.
38
39
40
41

42
43 GRADE (Grading of Recommendations, Assessment, Development, and Evaluation)
44
45 [30] system will be used to assess the quality of evidence for short- and long-term
46
47 mortality, length of ICU stay, need for ICU admission, length of hospital stay, and other
48
49 complications. Based on the risk of bias, inconsistency, indirectness, imprecision, and
50
51 other considerations, the quality of evidence will be rated as 'high', 'moderate', 'low',
52
53 and 'very low'.
54
55
56
57

58 **Statistical analysis**

1
2
3
4 Review Manager (version 5.4) software will be used to perform the analysis.
5
6 Continuous variables will be calculated using the mean difference (MD) with 95%
7
8 confidence intervals (CI), while dichotomous outcomes, such as 30-day mortality, can
9
10 be presented using the risk ratios (RR) with the same CI. Continuous results will be
11
12 presented as the mean and standard deviation. If the median is displayed, the median
13
14 and interquartile range can be converted to mean and standard deviation using the
15
16 statistical formula [31, 32]. For each endpoint, the heterogeneity will be visualized in
17
18 forest plots and assessed using Cochran's Q test and I^2 statistics. Statistical
19
20 heterogeneity across studies will be evaluated using a corresponding *p-value* derived
21
22 from the chi-square and I^2 tests. A value of $I^2 > 50\%$ or $P < 0.10$ is considered
23
24 significantly heterogeneous. If the included studies are homogeneous ($P \geq 0.10$ and
25
26 $I^2 < 50\%$), a fixed-effect model will be used to estimate the outcome data. However, a
27
28 random-effect model will be used to aggregate data with significant heterogeneity,
29
30 wherein any potential sources of heterogeneity will be identified and analyzed using
31
32 subgroup analysis. Subgroup and meta-regression analyses will be conducted to explore
33
34 the possible sources of heterogeneity and inconsistency, including the assessment tool
35
36 for sarcopenia, age, preoperative comorbidities, and the level of care obtained from
37
38 different centers. Moreover, the robustness of the results will be verified using a leave-
39
40 one-out jackknife sensitivity analysis conducted by removing one study at a time before
41
42 repeating the analysis. If more than ten studies are included in the meta-analysis, a
43
44 funnel plot will be generated to assess publication bias. After synthesizing the data and
45
46 classifying it as outlined above, the final report will be written following the PRISMA
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 criteria.

5 6 **Subgroup and sensitivity analyses**

7
8
9 Subgroup analyses will be performed depending on the number of identified studies.
10
11 The level of heterogeneity across included studies and the strength of the supporting
12
13 evidence will be examined using the Cochrane Q and I² statistics with associated 95%
14
15 CI. An I² value of 50% and above is considered a substantial level of heterogeneity. A
16
17 random-effect model will be used if potential heterogeneity is seen among study
18
19 variations, including population, regions, or assessment methods of sarcopenia.
20
21 However, if tests of heterogeneity among studies are not found to be significant, a fixed-
22
23 effect model will be used. To examine the methodological quality and potential sources
24
25 of heterogeneity in the included studies, a range of sensitivity analyses may be
26
27 conducted. Sources of variations can include the assessment tool for sarcopenia, age,
28
29 preoperative comorbidities, and the level of care from different centers. These sources
30
31 will be stratified, and sensitivity analyses will be conducted separately. A leave-one-
32
33 out jackknife sensitivity analysis will be used to verify the robustness of the results.
34
35 Generally, a minimum of two studies are sufficient to perform a meta-analysis. If the
36
37 level of heterogeneity between studies is high with no possibility of performing a
38
39 pooled analysis, a narrative summary of the outcome will be presented in a systematic
40
41 review along with the reasons for the obtained results.
42
43
44
45
46
47
48
49
50
51

52 53 **Patient and public involvement**

54
55
56 None.

57 58 **Ethics and dissemination**

1
2
3
4 No ethical approval is needed for this review as we will use data from previously
5
6 published studies. Upon completion, the results will be disseminated in a peer-reviewed
7
8 journal.
9

10 11 **Discussion**

12
13
14 To the best of our knowledge, this will be the first systematic review and meta-analysis
15
16 to examine the effects of sarcopenia on TAVI outcomes. Since TAVI candidates are
17
18 elderly and exhibit high morbidity due to sarcopenia, it is becoming a growing problem
19
20 among the aging population. This study aims to investigate the impact of sarcopenia on
21
22 post-TAVI outcomes, providing important predictive information for general medical
23
24 management strategies, nursing goals, and rehabilitation expectations.
25
26
27
28

29
30 Our study findings could provide a reliable basis for clinical practice. Sarcopenia is
31
32 vulnerable to iatrogenic stressors and also prone to deterioration in the long term.
33
34 Therefore, it ultimately needs to be integrated into the complex decision-making of
35
36 choosing between the operational and conservative approaches in TAVI candidates [33].
37
38 Objectively, evidence has supported the simplification of the evaluation of sarcopenia
39
40 for clinical purposes. Currently, pre-TAVI CT scans are routinely performed, which
41
42 may provide important complementary information for sarcopenia assessment.
43
44
45
46
47 Sarcopenia is highlighted as a risk predictor in TAVI patients, evaluating it as an
48
49 objective and readily available tool to be introduced in clinical practice. However,
50
51 sarcopenia, as a risk predictor, can also suggest the correction of sarcopenia using
52
53 protein supplementation and exercise training, ultimately improving post-TAVI
54
55
56
57
58 outcomes.
59
60

1
2
3
4 Nevertheless, this meta-analysis protocol may include some limitations. First, the
5
6 measurement of muscle mass and the definition of sarcopenia may not be uniform
7
8
9 across studies, which may impact the outcomes. Second, although more than 90% of
10
11 all TAVI cases are currently performed via the transfemoral route, transapical vascular
12
13 access along with subclavian, carotid, and trans-carotid routes may be usually favored
14
15 to avoid protracted recovery associated with thoracic access. However, this problem
16
17 could potentially be addressed by subgroup analysis and including sufficient studies.
18
19
20 Third, since we will be selecting published articles, publication bias may be
21
22 unavoidable.
23
24
25
26
27
28
29

30 ** **
31
32
33
34

35 **Patient consent for publication**

36
37 Not applicable.
38
39

40 **Funding**

41
42 This research received no specific grant from any funding agency in public, commercial
43
44 or not-for-profit sectors.
45
46
47

48 **Contributors**

49
50 Preparation of the manuscript, and figures was performed by K.L., L.Y. and all authors
51
52 reviewed the final manuscript. Other specific contributions are as follows: Conception
53
54 and design: K.L., L.Y. Collection and assembly of studies: K.L., L.Y. Manuscript
55
56 writing: All authors. Drafting and revision of manuscript and Final approval of
57
58
59
60

manuscript: All authors.

Competing interests

No competing interests.

References

1. Mack M J, M B Leon, V H Thourani, R Makkar, S K Kodali, M Russo, S R Kapadia, S C Malaisrie, D J Cohen, P Pibarot, J Leipsic, R T Hahn, P Blanke, M R Williams, J M McCabe, D L Brown, V Babaliaros, S Goldman, W Y Szeto, P Genereux, A Pershad, S J Pocock, M C Alu, J G Webb. C R Smith, Transcatheter Aortic-Valve Replacement with a Balloon-Expandable Valve in Low-Risk Patients. *N Engl J Med*, 2019. **380**(18): p. 1695-1705.
2. Leon M B, C R Smith, M J Mack, R R Makkar, L G Svensson, S K Kodali, V H Thourani, E M Tuzcu, D C Miller, H C Herrmann, D Doshi, D J Cohen, A D Pichard, S Kapadia, T Dewey, V Babaliaros, W Y Szeto, M R Williams, D Kereiakes, A Zajarias, K L Greason, B K Whisenant, R W Hodson, J W Moses, A Trento, D L Brown, W F Fearon, P Pibarot, R T Hahn, W A Jaber, W N Anderson, M C Alu. J G Webb, Transcatheter or Surgical Aortic-Valve Replacement in Intermediate-Risk Patients. *N Engl J Med*, 2016. **374**(17): p. 1609-20.
3. Gallone G, A Depaoli, F D'Ascenzo, D Tore, L Allois, F Bruno, M Casale, F Atzeni, G De Lio, P P Bocchino, F Piroli, F Angelini, A Angelini, L Scudeler, F De Lio, A Andreis, S Salizzoni, M La Torre, F Conrotto, M Rinaldi, P Fonio. G M De Ferrari, Impact of computed-tomography defined sarcopenia on outcomes of older adults undergoing transcatheter aortic valve implantation. *J Cardiovasc Comput Tomogr*, 2022. **16**(3): p. 207-214.
4. Holmes D R, Jr., J M Brennan, J S Rumsfeld, D Dai, S M O'Brien, S Vemulapalli, F H Edwards, J Carroll, D Shahian, F Grover, E M Tuzcu, E D Peterson, R G Brindis. M J Mack, Clinical outcomes at 1 year following transcatheter aortic valve replacement. *Jama*, 2015. **313**(10): p. 1019-28.
5. Debonnaire P, L Fusini, R Wolterbeek, V Kamperidis, P van Rosendaal, F van der Kley, S Katsanos, E Joyce, G Tamborini, M Muratori, P Gripari, J J Bax, N A Marsan, M Pepi. V Delgado, Value of the "TAVI2-SCORE" versus surgical risk scores for prediction of one year mortality in 511 patients who underwent transcatheter aortic valve implantation. *Am J Cardiol*, 2015. **115**(2): p. 234-42.
6. Soud M, F Alahdab, G Ho, K O Kuku, M Cejudo-Tejeda, A Hideo-Kajita, P de Araujo Gonçalves, R C Teles, R Waksman. H M Garcia-Garcia, Usefulness of skeletal muscle area detected by computed tomography to predict mortality in patients undergoing transcatheter aortic valve replacement: a meta-analysis

- study. *Int J Cardiovasc Imaging*, 2019. **35**(6): p. 1141-1147.
7. Mok M, R Allende, J Leipsic, O A Altisent, M Del Trigo, F Campelo-Parada, R DeLarochelière, E Dumont, D Doyle, M Côté, M Freeman, J Webb.J Rodés-Cabau, Prognostic Value of Fat Mass and Skeletal Muscle Mass Determined by Computed Tomography in Patients Who Underwent Transcatheter Aortic Valve Implantation. *Am J Cardiol*, 2016. **117**(5): p. 828-33.
 8. Cruz-Jentoft A J, J P Baeyens, J M Bauer, Y Boirie, T Cederholm, F Landi, F C Martin, J P Michel, Y Rolland, S M Schneider, E Topinková, M Vandewoude.M Zamboni, Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*, 2010. **39**(4): p. 412-23.
 9. Dodds R M.A A Sayer, Sarcopenia, frailty and mortality: the evidence is growing. *Age Ageing*, 2016. **45**(5): p. 570-1.
 10. Cruz-Jentoft A J, G Bahat, J Bauer, Y Boirie, O Bruyère, T Cederholm, C Cooper, F Landi, Y Rolland, A A Sayer, S M Schneider, C C Sieber, E Topinkova, M Vandewoude, M Visser.M Zamboni, Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*, 2019. **48**(1): p. 16-31.
 11. Jones K, A Gordon-Weeks, C Coleman.M Silva, Radiologically Determined Sarcopenia Predicts Morbidity and Mortality Following Abdominal Surgery: A Systematic Review and Meta-Analysis. *World J Surg*, 2017. **41**(9): p. 2266-2279.
 12. Chang S F.P L Lin, Systematic Literature Review and Meta-Analysis of the Association of Sarcopenia With Mortality. *Worldviews Evid Based Nurs*, 2016. **13**(2): p. 153-62.
 13. Boutin R D, L Yao, R J Canter.L Lenchik, Sarcopenia: Current Concepts and Imaging Implications. *AJR Am J Roentgenol*, 2015. **205**(3): p. W255-66.
 14. ter Maaten J M, K Damman, H L Hillege, S J Bakker, S D Anker, G Navis.A A Voors, Creatinine excretion rate, a marker of muscle mass, is related to clinical outcome in patients with chronic systolic heart failure. *Clin Res Cardiol*, 2014. **103**(12): p. 976-83.
 15. Weerink L B M, A van der Hoorn, B L van Leeuwen.G H de Bock, Low skeletal muscle mass and postoperative morbidity in surgical oncology: a systematic review and meta-analysis. *J Cachexia Sarcopenia Muscle*, 2020. **11**(3): p. 636-649.
 16. Aleixo G F P, S S Shachar, K A Nyrop, H B Muss, L Malpica.G R Williams, Myosteatosis and prognosis in cancer: Systematic review and meta-analysis. *Crit Rev Oncol Hematol*, 2020. **145**: p. 102839.
 17. Yang T R, K Luo, X Deng, L Xu, R R Wang.P Ji, Effect of sarcopenia in predicting postoperative mortality in emergency laparotomy: a systematic review and meta-analysis. *World J Emerg Surg*, 2022. **17**(1): p. 36.
 18. Chang K V, T H Hsu, W T Wu, K C Huang.D S Han, Association Between Sarcopenia and Cognitive Impairment: A Systematic Review and Meta-


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Analysis. *J Am Med Dir Assoc*, 2016. **17**(12): p. 1164.e7-1164.e15.
19. Chang K V, T H Hsu, W T Wu, K C Huang, D S Han, Is sarcopenia associated with depression? A systematic review and meta-analysis of observational studies. *Age Ageing*, 2017. **46**(5): p. 738-746.
20. Bang J Y, I G Jun, J B Lee, Y S Ko, K W Kim, J H Jeong, S H Kim, J G Song, Impact of Sarcopenia on Acute Kidney Injury after Infrarenal Abdominal Aortic Aneurysm Surgery: A Propensity Matching Analysis. *Nutrients*, 2021. **13**(7).
21. Sim J H, H M Kwon, I G Jun, S H Kim, K S Kim, Y J Moon, J G Song, G S Hwang, Association of skeletal muscle index with postoperative acute kidney injury in living donor hepatectomy: A retrospective single-centre cohort study. *Liver Int*, 2022. **42**(2): p. 425-434.
22. Francomacaro L M, C Walker, K Jaap, J Dove, M Hunsinger, K Widom, D Torres, M Shabahang, J Blansfield, J Wild, Sarcopenia predicts poor outcomes in urgent exploratory laparotomy. *Am J Surg*, 2018. **216**(6): p. 1107-1113.
23. McQuade C, D O Kavanagh, C O'Brien, K Hunter, D Nally, C Hickie, E Ward, W C Torreggiani, CT-determined sarcopenia as a predictor of post-operative outcomes in patients undergoing an emergency laparotomy. *Clin Imaging*, 2021. **79**: p. 273-277.
24. Damluji A A, G Rodriguez, T Noel, L Davis, V Dahya, B Tehrani, K Epps, M Sherwood, E Sarin, J Walston, K Bandeen-Roche, J R Resar, T T Brown, G Gerstenblith, C M O'Connor, W Batchelor, Sarcopenia and health-related quality of life in older adults after transcatheter aortic valve replacement. *Am Heart J*, 2020. **224**: p. 171-181.
25. Bertschi D, C M Kiss, A W Schoenenberger, A E Stuck, R W Kressig, Sarcopenia in Patients Undergoing Transcatheter Aortic Valve Implantation (TAVI): A Systematic Review of the Literature. *J Nutr Health Aging*, 2021. **25**(1): p. 64-70.
26. Shamseer L, D Moher, M Clarke, D Ghera, A Liberati, M Petticrew, P Shekelle, L A Stewart, Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ*, 2015. **350**: p. g7647.
27. Chen L K, L K Liu, J Woo, P Assantachai, T W Auyeung, K S Bahyah, M Y Chou, L Y Chen, P S Hsu, O Krairit, J S Lee, W J Lee, Y Lee, C K Liang, P Limpawattana, C S Lin, L N Peng, S Satake, T Suzuki, C W Won, C H Wu, S N Wu, T Zhang, P Zeng, M Akishita, H Arai, Sarcopenia in Asia: consensus report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc*, 2014. **15**(2): p. 95-101.
28. Hayden J A, P Côté, C Bombardier, Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med*, 2006. **144**(6): p. 427-37.
29. Hayden J A, D A van der Windt, J L Cartwright, P Côté, C Bombardier, Assessing bias in studies of prognostic factors. *Ann Intern Med*, 2013. **158**(4): p. 280-6.
30. Brozek J L, E A Akl, P Alonso-Coello, D Lang, R Jaeschke, J W Williams, B

- 1
2
3 Phillips, M, Leigemann, A, Lethaby, J, Bousquet, G, H Guyatt, H, J Schünemann,
4 Grading quality of evidence and strength of recommendations in clinical
5 practice guidelines. Part 1 of 3. An overview of the GRADE approach and
6 grading quality of evidence about interventions. *Allergy*, 2009. **64**(5): p. 669-
7 77.
8
9
10 31. Luo D, X Wan, J Liu, T Tong, Optimally estimating the sample mean from the
11 sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med*
12 *Res*, 2018. **27**(6): p. 1785-1805.
13
14 32. Wan X, W Wang, J Liu, T Tong, Estimating the sample mean and standard
15 deviation from the sample size, median, range and/or interquartile range. *BMC*
16 *Med Res Methodol*, 2014. **14**: p. 135.
17
18 33. Andò G, G Basile, Sarcopenia: only one of the domains of frailty in patients
19 undergoing transcatheter aortic valve implantation. *J Cardiovasc Med*
20 *(Hagerstown)*, 2020. **21**(10): p. 787-789.
21
22

Figure title/legend

Figure 1. Flowchart

The flowchart will be used to document the process of selecting articles for
inclusion in the systematic review and meta-analysis.



45 **Fig 1.** Flowchart showing selection of articles for review

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Appendix

We will search the reference lists of all identified publications for additional studies. There is no restriction on the language of publication. The searches will be rerun prior to the final analyses and any further studies identified will be included

Search terms for PubMed

Search	Query
#1	"Sarcopenia"[MeSH Terms]
#2	"Sarcopenia"[Title/Abstract] OR "Sarcopenias"[Title/Abstract] OR "Muscle mass"[Title/Abstract] OR "Muscle area"[Title/Abstract]
#3	#1 OR #2
#4	"Transcatheter aortic valve implantation"[MeSH Terms] "Transcatheter aortic valve implantation"[Title/Abstract] OR "Transcatheter aortic valve replacement"[Title/Abstract] OR "TAVI"[Title/Abstract] OR "TAVR"[Title/Abstract]
#5	#4 OR #5
#6	#4 OR #5
#7	#3 AND #6

Search terms for MEDLINE

Search	Query
#1	Sarcopenia. ab, ti.
#2	Sarcopenias. ab, ti.
#3	Muscle mass. ab, ti.
#4	Muscle area. ab, ti.
#5	#1 OR #2 OR #3 OR #4
#6	Transcatheter aortic valve implantation. ab, ti.
#7	Transcatheter aortic valve replacement. ab, ti.
#8	TAVI. ab, ti.
#9	TAVR. ab, ti.
#10	#6 OR #7 OR #8 OR #9
#11	#5 AND #10

Search terms for the Cochrane Library (in Title, Abstract, Keyword)

Search	Query
#1	MeSH descriptor: [Sarcopenia] explode all trees
#2	Sarcopenia
#3	Sarcopenias
#4	MeSH descriptor: [Muscle mass] explode all trees
#5	Muscle mass
#6	Skeletal muscle mass
#7	MeSH descriptor: [Muscle area] explode all trees
#8	Muscle area

1	
2	
3	#9 Muscle areas
4	#10 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9
5	#11 MeSH descriptor: [Transcatheter aortic valve implantation] explode all trees
6	#12 Transcatheter aortic valve implantation
7	#13 Transcatheter aortic valve replacement
8	#14 TAVI
9	#15 TAVR
10	#16 #11 OR #12 OR #13 OR #14 OR #15
11	#17 #10 AND #16
12	
13	
14	
15	
16	
17	

Search terms for EMBASE

Search	Query
#1	'Sarcopenia'/exp
#2	'Sarcopenia': ab, ti
#3	'Sarcopenias': ab, ti
#4	'Muscle mass'/exp
#5	'Muscle mass': ab, ti.
#6	'Muscle area'/exp
#7	'Muscle area': ab, ti.
#8	'Muscle areas': ab, ti.
#9	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8
#10	'Transcatheter aortic valve implantation'/exp
#11	'Transcatheter aortic valve implantation': ab, ti.
#12	'Transcatheter aortic valve replacement': ab, ti.
#13	'TAVI': ab, ti.
#14	'TAVR': ab, ti.
#15	#10 OR #11 OR #12 OR #13 OR #14
#16	#9 AND #15

Search terms for Web of Science

Search	Query
#1	Sarcopenia (Topic) OR Sarcopenia (Title) OR Sarcopenia (Abstract) OR Sarcopenias (Topic) OR Sarcopenias (Title) OR Sarcopenias (Abstract)
#2	Muscle mass (Topic) OR Muscle mass (Title) OR Muscle mass (Abstract) OR Muscle area (Topic) OR Muscle area (Title) OR Muscle area (Abstract) OR Muscle areas (Topic) OR Muscle areas (Title) OR Muscle areas (Abstract)
#3	#1 OR #2
#4	Transcatheter aortic valve implantation (Topic) OR Transcatheter aortic valve implantation (Title) OR Transcatheter aortic valve implantation (Abstract)
#5	Transcatheter aortic valve implantation (Topic) OR Transcatheter aortic valve implantation (Title) OR Transcatheter aortic valve implantation (Abstract)
#6	TAVI (Topic) OR TAVI (Title) OR TAVI (Abstract)

1
2
3 #7 TAVR (Topic) OR TAVR (Title) OR TAVR (Abstract)
4 #8 #4 OR #5 OR #6 OR #7
5 #9 #3 AND #8
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item
ADMINISTRATIVE INFORMATION		
Title:		
Identification	1a	Identify the report as a protocol of a systematic review (page 1)
Update	1b	If the protocol is for an update of a previous systematic review, identify as such (No)
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number (page 3)
Authors:		
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author (page 1)
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review (page 12)
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments (No)
Support:		
Sources	5a	Indicate sources of financial or other support for the review (page 12)
Sponsor	5b	Provide name for the review funder and/or sponsor (No)
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol (No)
INTRODUCTION		
Rationale	6	Describe the rationale for the review in the context of what is already known (page 3 to 5)
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO) (page 6 to 7)
METHODS		
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review (page 6 to 7)
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage (page 6)
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated (page 6 and appendix)
Study records:		
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review (page 8)

1			
2			
3	Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis) (page 7 to 8)
4			
5	Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms done independently, in duplicate), any processes for obtaining and confirming data from investigators (page 6 and page 7)
6			
7	Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications (page 7)
8			
9	Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale (page 7)
10			
11	Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis (page 8)
12			
13	Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised (page 8 to 9)
14		15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I^2 , Kendall's τ) (page 8 to 9)
15		15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression) (page 9 to 10)
16		15d	If quantitative synthesis is not appropriate, describe the type of summary planned (page 10)
17			
18	Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies) (page 9)
19			
20	Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE) (page 8)
21			
22			
23			
24			
25			
26			
27			
28			
29			
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			