

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Exploring digital health interventions for pregnant women at high risk for preeclampsia and eclampsia in low-and-middle-income countries: a scoping review

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-056130
Article Type:	Original research
Date Submitted by the Author:	04-Aug-2021
Complete List of Authors:	Shahil Feroz, Anam; Aga Khan University, ; University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Afzal, Noreen; The Aga Khan University Medical College Pakistan, Dean's Office Seto, Emily; University of Toronto, Institute of Health Policy, Management and Evaluation; University Health Network, Centre for Global eHealth Innovation
Keywords:	PUBLIC HEALTH, Prenatal diagnosis < OBSTETRICS, Health informatics < BIOTECHNOLOGY & BIOINFORMATICS, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

## Title of Scoping Review

Exploring digital health interventions for pregnant women at high risk for preeclampsia and eclampsia in low-and-middle-income countries: a scoping review

a. Author and Co-author's name

1. Anam Shahil Feroz [anam.shahil@mail.utoroto.ca](mailto:anam.shahil@mail.utoroto.ca)
2. Noreen Afzal [noreen.afzal100@gmail.com](mailto:noreen.afzal100@gmail.com)
3. Emily Seto [Emily.seto@utoronto.ca](mailto:Emily.seto@utoronto.ca)

b. The name, postal address, e-mail, telephone, and fax numbers of the corresponding author.

Name: Anam shahil Feroz

Affiliations:

- Aga Khan University, Community Health Sciences Department
- University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Toronto, ON, CAN M5T 3M6

Email: [anam.shahil@mail.utoronto.ca](mailto:anam.shahil@mail.utoronto.ca)

Phone: +1647 679 6702

c. The full names, institutions, city, and country of all co-authors.

1. Anam Shahil Feroz

Affiliations:

- Aga Khan University, Community Health Sciences Department, Karachi, Sindh Pakistan
- University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Toronto, Ontario, Canada

2. Noreen Afzal

Affiliations:

- The Aga Khan University Medical College Pakistan, Dean's Office Karachi, Sindh Pakistan

3. Emily Seto

Affiliations:

- University of Toronto, Institute of Health Policy, Management and Evaluation Toronto, Ontario, Canada
- University Health Network, Centre for Global eHealth Innovation Toronto, Ontario, Canada

d. Up to five keywords or phrases suitable for use in an index (it is recommended to use MeSH terms).

Digital health interventions, pregnant women, preeclampsia, eclampsia, low-and-middle-income countries, predictive models, mHealth applications, devices

e. Word count - excluding title page, references, figures and tables: 5899

**Abstract**

**Objective:** To explore digital health interventions (DHIs) that have been used to support pregnant women at HRPE/E in LMICs

**Design:** Scoping review

**Data Source:** Five electronic databases were searched including EMBASE, MEDLINE, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and CINAHL, between January 1, 2000, and October 20, 2020, to identify the articles that described DHIs implemented to support pregnant women at HRPE/E.

**Results:** A total of 19 publications describing seven unique studies and nine different DHIs were included. Most studies were conducted in South Asia and Sub-Saharan Africa (n=16). Of nine unique DHIs, two served the purpose of predicting risk for adverse maternal health outcomes while seven DHIs focused on monitoring high-risk pregnant women for managing PE/E. Both of these purposes utilized mobile phone applications as interface to facilitate data collection, decision making, and communication between health workers and pregnant women. Seven articles described the use of more than one unique DHI. Data collection, prediction of adverse maternal outcomes, integrated diagnostic and clinical decision support, and personal health tracking were reported as the key functions of DHIs. The articles reported three major outcomes: 1) maternal health outcomes (n=4), 2) usability and acceptability (n=5), and 3) intervention feasibility and fidelity (n=7).

**Conclusion:** Although the current evidence base of DHIs shows some potential for the use of different DHIs to support pregnant women in early diagnosis of PE/E, more prospective experimental and longitudinal studies are needed prior to recommending the use of DHIs for pregnant women at HRPE/E in LMICs. To support pregnant women at HRPE, future research work should incorporate telemedicine to enable remote consultation between pregnant women and healthcare providers, consider a multidisciplinary team approach for designing DHIs, and enable the use of DHIs by pregnant women as end-users instead of healthcare providers as end-users.

**Strengths and limitations of this study**

- 1  
2  
3 1. First scoping review to explore the use of DHIs in LMICs to support pregnant women at  
4 HRPE/E.  
5
- 6  
7 2. The high heterogeneity of the DHIs and study outcomes limited the interpretation of the studies  
8 through quantitative analysis.  
9
- 10  
11 3. This review also only included peer-reviewed articles and papers published in the English  
12 Language.  
13
- 14  
15 4. The review did not include information that may have been found in other databases and sources  
16 (abstracts, reviews, conference proceedings, opinion papers, books).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

Approximately 16% of all maternal deaths in low-and-middle-income countries (LMICs) are attributable to preeclampsia/eclampsia (PE/E) [1]. High maternal mortality from PE/E results from: 1) lack of early identification and treatment of pregnant women, 2) difficulties in reaching treatment centers and, 3) poor health-seeking behaviors linked with low patient education [2]. To meet the United Nations Sustainable Developmental Goal target 3.1 of reducing the maternal mortality ratio to less than 70/100,000 live births by 2030, innovations are required to decrease PE/E-related mortality [3].

The most effective strategies to ensure early diagnosis and management of PE/E include self-monitoring of blood pressure, use of magnesium sulfate therapy, proteinuria determinations, and timely delivery [1]. International guidelines including the European Society of Hypertension, American Heart Association, National Institute for Health and Care Excellence, and American Society of Hypertension guidelines, recommend self-monitoring for PE symptoms and recording of blood pressure for pregnant women at high risk for preeclampsia and eclampsia (HRPE/E) because of their potential benefits such as effective control of blood pressure, early risk identification, and treatment, and cost-savings due to fewer hospital visits [4-6]. Self-monitoring also has a role in preventing conditions like white coat hypertension and masked hypertension in pregnant women at HRPE/E. The World Health Organization (WHO) suggests home blood pressure monitoring for pregnant women at HRPE/E to detect changes in blood pressure between antenatal visits and to ensure care continuity [7].

Digital health interventions (DHIs) are increasingly being used to support pregnant women at HRPE/E for remote monitoring of blood pressure and symptoms. To date, four reviews explored the use of digital tools for remote monitoring of pregnant women at HRPE/E. In 2020, Aquino et al. reported 16 unique, feasible, and cost-effective telemonitoring interventions to support pregnant women with hypertensive disorder of pregnancy [6]. However, the review mainly focused on telemonitoring interventions for remote blood pressure monitoring of pregnant women. The review also primarily identified studies from high-income countries like the UK, USA, and Belgium [6]. Lanssens et al. (2017) reported 14 studies from 1988 to 2010 that used telemonitoring interventions for pregnant women during the prenatal period[8]. This review, however, used a narrow time range and focused on telemonitoring solutions implemented in high-income countries for pregnant women at high risk for gestational diabetes and

1  
2  
3 preterm labor. In addition, the included studies had a high methodological risk of bias. When only  
4 studies with low risk of bias were considered, the added value of telemonitoring became less  
5 pronounced [8]. Rivera-Romero et al. (2018) captured only 11 studies conducted in high-income  
6 countries, on mobile health (mHealth) interventions for the hypertensive disorder of pregnancy[9]. The  
7 included studies showed positive results in the improvement of maternal health and acceptability of  
8 solutions, although most of the studies involved a small number of participants, and none were complete  
9 clinical studies [9]. Van den Heuvel et al. (2018) reported 12 studies on the use of telemonitoring and  
10 teleconsulting interventions to improve pregnancy care generally [10]. The review did not focus on the  
11 use of eHealth for the hypertensive disorder of pregnancy and generally included all aspects of perinatal  
12 care.

13  
14 These four reviews provided foundational information on the use of telemonitoring to support high-risk  
15 pregnant women in antepartum and postpartum period. However, quality evidence on the appropriate  
16 use of DHIs to support pregnant women at HRPE/E in LMIC is scarce. None of the reviews extensively  
17 documented the use of DHIs in LMICs for the early diagnosis and management of pregnant women at  
18 HRPE/E. This gap highlights the need to explore the potential role of DHIs to support pregnant women  
19 at HRPE/E in LMICs. This review aims to systematically explore the available literature on the use of  
20 DHIs to support early detection and management of PE/E in LMICs. Specifically, the research question  
21 for this scoping review is: What is known in the literature about DHIs that have been used to support  
22 pregnant women at HRPE/E in LMICs?

## 23 24 **Methods**

25  
26 The “Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping  
27 Reviews ” checklist was used to guide the design and reporting of this scoping review [11]  
28 (Supplementary file 1: Completed PRISMA-ScR Checklist). The review was registered in the Open  
29 Science Framework (OSF) - Center for Open Science on Oct 19, 2020 (Registration link:  
30 <https://osf.io/gncvj>). The review was guided by the methodological framework by Levac et al. [12] and  
31 Arksey et al. [13] to examine articles describing the use of digital health solutions to support early  
32 detection and management of PE/E in LMICs.



### ***Eligibility Criteria***

The review included studies that involved pregnant women at HRPE/E and implemented the digital health solutions to support early detection and management of PE/E in LMICs. For this scoping review, the DHIs included wearable devices, predictive models operationalized through clinical applications, health information technologies, health management systems, and other innovations related to mobile health, telehealth, and telemedicine that can guide diagnosis, monitoring, and treatment[14]. The review included only English-language studies, which were conducted in LMICs. The World Bank's (WB) 2020 country classification list was used to select LMICs with a Gross National Income (GNI) per capita between \$1,036 and \$4,045 [15]. The review primarily aimed to include original and primary research studies, including experimental studies (e.g., randomized controlled trials, quasi-experimental studies), observational studies (e.g., cohort, case-control, cross-sectional, qualitative studies), and study protocols. All types of reviews, meta-analyses, letters to editors, commentaries, viewpoints, news articles, abstracts, and books were excluded. Articles published between January 1, 2000, and October 20, 2020, were included, given that DHIs prior to 2000 would likely have little applicability for current implementation (Supplementary file 2: Eligibility Criteria).

### ***Information Sources and Search Strategy***

Five main electronic databases were searched including Excerpta Medica Database (EMBASE), Medical Literature Analysis and Retrieval System Online (MEDLINE), Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and Cumulated Index to Nursing and Allied Health Literature (CINAHL). A supplementary search was conducted using the first seven pages of Google Scholar to capture peer-reviewed literature on the use of DHIs to support pregnant women at HRPE. The reference lists of relevant systematic reviews and final included articles were also hand-searched to find pertinent studies. The search strategy was developed with the assistance of an expert librarian specializing in health services research. It included four main concepts of interest: target population (pregnant women), health condition (PE), intervention (digital health tools), and settings (LMICs). The search strategy included both keywords and subject headings such as MeSH, and Emtree (Supplementary file 3: Search strategy for the MEDLINE database).

### ***Selection Procedure***

Records from all the electronic databases were exported to Endnote software for screening purposes.

The primary reviewer (AS) developed a pre-defined screening form, and pilot testing was carried out using 10 randomly selected articles to ensure appropriate screening reliability among the two reviewers (AS and NA), which was found to be 90%. All articles were independently screened by the two reviewers to exclude those that did not fulfill the inclusion criteria. Two reviewers then met to review any discrepancies which were discussed until a consensus was reached.

The initial search found a total of 4,078 articles. After de-duplication, 3,389 titles and abstracts were screened by the two reviewers (AS and NA) to evaluate whether they met the eligibility criteria. Of these, 72 records were found to be eligible for full-text screening by the two reviewers. Finally, 19 articles were identified after the full-text screening that met the inclusion criteria for this review [16-34]. Fifty-three articles were excluded for the following reasons: (1) the study was not reported in the English language; (2) the publication did not talk about pregnant women at HRPE; (3) the research did not include any of the DHIs; (4) the publication was a conference abstract, review, editorial, commentary; or (5) the study implemented the DHIs for pregnant women at HRPE in high-income countries. The study selection procedure was recorded according to the “Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA)” flow diagram (Figure 1: PRISMA Flow Diagram for Database Search of Studies).

### ***Data Extraction***

A data abstraction form was designed collectively by the research team to determine appropriate variables such as study characteristics, type of DHIs, intervention description, and study outcomes (Supplementary file 4: Data Abstraction form). To ensure consistency in the data extraction process, the form was pilot tested using three randomly selected articles, which resulted in consistent data being abstracted by both reviewers. Both reviewers (AS, NA) independently completed the data extraction sheet for each of the 19 final articles. The data abstraction sheets of both the reviewers were compared to confirm that all major results were included in the scoping review. In the case of inconsistencies

1  
2  
3 between the data extraction sheets from the two reviewers, a third reviewer would have been invited to  
4  
5 make a final decision, but no inconsistencies were found.

### 6 7 ***Data Analysis***

8  
9 An inductive approach was used to thematically organize and summarize the results from the included  
10  
11 articles to explore our research question. The extracted results from each article were read several times  
12  
13 to identify frequent patterns, similarities, and differences on the use of DHIs to support pregnant women  
14  
15 at HRPE in LMICs. The identified emerging patterns were organized into five thematic groupings  
16  
17 including study characteristics, overview and appraisal of included studies, purpose of DHIs, users of  
18  
19 DHIs, and types of outcomes examined by the included studies. The first, and last author discussed the  
20  
21 results and agreed upon the final groupings of the results.  
22  
23  
24  
25

## 26 **Results**

### 27 28 ***Study Characteristics***

29  
30 A total of 19 publications describing 7 unique studies were included in this review. The included articles  
31  
32 were published between 2009 and 2020. Of these 19 articles, a total of 16 articles described studies that  
33  
34 were conducted in South Asia and Sub-Saharan Africa, one article described a study conducted in  
35  
36 Africa, Southern Asia, and the Middle East, and the remaining two articles described studies conducted  
37  
38 in unspecified resource-poor settings (LMICs) (Supplementary file 5: Overview of the included  
39  
40 articles).  
41  
42

43 The 19 articles were classified into three types of articles: observational studies (n=12), experimental  
44  
45 studies (n=4 including two RCTs), and protocol papers (n= 3). All included articles reported the use of  
46  
47 DHIs for antepartum women. The articles reported varying eligibility criteria for selecting high-risk  
48  
49 pregnant women for different DHIs. Some articles selected high-risk pregnant women based on the  
50  
51 National Institute for Health and Care Excellence (NICE) guidelines [35], specific age groups such as  
52  
53 pregnant women aged 15–49 years [22], while a few articles selected pregnant women based on their  
54  
55 residential area such as women living in study catchment area [23], permanent resident of the particular  
56  
57 area, or non-resident who delivered in the study area[18]. . Most DHIs collected blood pressure, heart  
58  
59 rate, and pulse oximetry, with some innovations collecting data on additional indicators such as  
60

1  
2  
3 demographic data, hemoglobin, urine dipstick test to detect proteinuria and glucose, other urinary  
4 markers, and PE symptoms. Only one article reported the use of international guideline (NICE clinical  
5 guideline 107) to determine blood pressure thresholds [28] (Supplementary file 6: DHIs  
6 characteristics).  
7  
8  
9

10  
11 Seven articles described the application of theoretical frameworks to guide the implementation and  
12 evaluation of digital health tools, including the technology acceptance model [25], diffusion of  
13 innovation model [26, 31], three delay model [26, 29], normalization process theory [23], medical  
14 research council framework [34], logic models [31, 34], realist evaluation theories [31], and cost-  
15 effectiveness models [22]. Two articles described the use of the LambdaNative framework for the  
16 development of the 'PIERS on the Move' (POTM) mHealth application [19, 24]. The remaining 10  
17 articles did not mention the use of theory or frameworks for the implementation of DHIs.  
18  
19  
20  
21  
22  
23  
24  
25

### 26 ***Overview of the Appraisal of Included Studies***

27  
28 A total of ten publications in this review reported research work of the monitoring component of PRE-  
29 EMPT (PE/E Monitoring, Prevention & Treatment) project by Peter von Dadelszen et al., University  
30 of British Columbia [17-19, 22-24, 29-31, 36]. The elements of the monitoring component include  
31 predictive models, Community Level Interventions for PE (CLIP) and integrated mHealth applications.  
32  
33 The PRE-EMPT initiative involved the work of the following research groups: CLIP Pakistan working  
34 group, CLIP India working group, CLIP trial collaborative group, and MiniPIERS and FullPIERS study  
35 working group. The PRE-EMPT project was funded through the Bill & Melinda Gates Foundation  
36 (\$25.9 million).  
37  
38  
39  
40  
41  
42  
43  
44

45 A total of four articles reported research work of CRADLE VSA trial led by Nathan et al., which aimed  
46 to evaluate the ability of the device to accurately detect abnormalities in women's vital signs during  
47 pregnancy [27, 28, 34, 37]. The remaining five publications reported five unique DHIs to support  
48 pregnant women at HRPE including the Congo Red Dot test [21], a hypothetical telemonitoring  
49 program [20], a new hypertension detector [32], an integrated diagnostic and clinical decision support  
50 system named 'bliss4midwives' [16], and a smart wristwatch (called the F1 smart wristwatch) for blood  
51 pressure monitoring of expectant mother [25].  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Following PRISMA-ScR guidelines, each of the above-mentioned included article was reviewed to  
4 identify emerging themes related to the use of DHIs to support pregnant women at HRPE in LMICs.  
5  
6 The key themes that emerged from the observational and experimental studies and protocol papers are  
7  
8 as follows: (1) purpose of DHIs including risk prediction and monitoring of high-risk pregnant women;  
9  
10  
11 (2) users of DHIs including healthcare providers (HCPs), caregivers, and pregnant women; (3) types of  
12  
13 outcomes examined in included studies including maternal and neonatal health outcomes, usability and  
14  
15 acceptability and intervention feasibility.  
16

### 17 ***Purpose of Digital Health Interventions***

18  
19  
20 This review reports nine unique DHIs from 19 included articles to support pregnant women at HRPE/E  
21  
22 in LMICs. These unique interventions are clustered into two main groups based on their purpose:  
23  
24 predicting risk of adverse maternal health outcomes (n=2) and monitoring high-risk pregnant women  
25  
26 to manage PE/E (n=7). Most articles (n=7) described the use of more than one unique DHI (Figure 2:  
27  
28 Classification of the Included Studies Based on the Purpose of Digital Health Interventions.).  
29

### 30 ***Predicting Risk of Adverse Maternal Health Outcomes***

31  
32  
33 Five observational studies and two RCTs described the use of two unique clinical predictive models  
34  
35 named fullPIERS [19] and miniPIERS [17-19, 24, 29-31] to facilitate the prediction of adverse maternal  
36  
37 outcomes occurring as a result of PE based on demographics, symptoms, clinical signs (including  
38  
39 SpO<sub>2</sub>), and laboratory tests. In order to implement these predictive models, the mobile application  
40  
41 'POTM' was developed as an interface to enable healthcare workers to easily determine the risk of  
42  
43 adverse maternal health outcomes. One article reported the use of both the miniPIERS and fullPIERS  
44  
45 predictive models [19], while six articles only reported the use of the miniPIERS model to predict  
46  
47 adverse health outcomes among pregnant women with PE/E in LMICs [17, 18, 24, 29-31]. Payne et al.  
48  
49 described the development process of the miniPIERS model to identify pregnant women at HRPE/E in  
50  
51 five LMICs using simple-to-measure indicators: personal demographics (gestational age); clinical signs  
52  
53 (blood pressure readings and proteinuria); and PE symptoms (headache, visual disturbances, chest pain,  
54  
55 dyspnea, vaginal bleeding, and abdominal pain) [29]. The fullPIERS model included additional  
56  
57 predictors such as SpO<sub>2</sub> and laboratory tests, to calculate a risk score for pregnant women.  
58  
59

### 60 ***Monitoring High-Risk Pregnant Women for Managing PE/E conditions***

1  
2  
3 The review identified seven unique DHIs for continuous monitoring high-risk pregnant women for  
4 managing PE/E including one diagnostic test named Congo Red Dot for monitoring misfolded protein  
5 in the preeclamptic urine [21], CLIP intervention for monitoring blood pressure among high-risk  
6 women through community health workers [17, 18, 22, 23, 30, 31], as well as five unique devices for  
7 monitoring blood pressure [16, 20, 25, 27, 28, 32, 34, 37]. The five unique devices for measuring blood  
8 pressure among high-risk pregnant women include the Microlife CRADLE VSA device [27, 28, 34,  
9 37], the Bliss4Midwives' device [16], a new hypertension detector device [32], hypothetical  
10 telemonitoring program [20] and the F1 smart wristwatch [25].

11  
12 The Congo Red Dot test was evaluated in a prospective experimental study design. The Congo Red  
13 Dot test requires minimal specialized equipment and enables minimally trained personnel to diagnose  
14 PE in resource-limited health care settings. The test was developed in 2016, based on the ability of  
15 constituents in preeclamptic urine to bind the amyloidophilic dye Congo Red. At the core of the test is  
16 the discovery that preeclamptic women eliminate misfolded proteins in their urine, a molecular feature  
17 that is proportional to disease severity [21].

18  
19 The CLIP intervention was implemented in Mozambique, Pakistan, India, and Nigeria as part of cluster  
20 randomized controlled trials (cRCTs) [17, 18, 22, 23, 30, 31]. The implementation of CLIP intervention  
21 involved scaling-up of existing community health workforce to provide community engagement and  
22 community health worker-led app-guided monitoring for high-risk pregnant women for hypertension.  
23 Community health workers were able to undertake all aspects of the app-guided visits, and  
24 approximately 10% of pregnant women were found to be hypertensive.

25  
26 As a first example of blood pressure measurement device, Nathan et al. assessed the accuracy of the  
27 Microlife 3AS1-2 blood pressure device in 2014 for use in pregnancy and PE in a low-resource setting  
28 [27]. The study recruited a total of 45 pregnant women, of whom 15 had PE, from Kimberley Hospital  
29 in South Africa. The study concluded that the device can be recommended for use in pregnancy,  
30 including PE as it fulfills the requirements stipulated by the WHO for an automated blood pressure  
31 device suitable for use in antenatal clinics and primary healthcare facilities of LMICs. The device has  
32 been extensively validated for accuracy, usability, and acceptability in low-resource settings [27]. The  
33 device calculates the pregnant woman's risk of hypovolaemic or septic shock and alerts frontline  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 healthcare workers about vital sign abnormalities through a traffic light early warning system display.  
4  
5 In 2018, a three-month mixed-methodology feasibility study was conducted to incorporate the  
6  
7 CRADLE VSA device into routine maternity care in 10 low-income sites [34]. Primary, secondary, and  
8  
9 tertiary facilities were allocated devices and training packages consisting of a short-animated film,  
10  
11 interactive sessions, booklet, and posters.  
12

13  
14 As a second example, a study conducted in Ghana used the Bliss4Midwives (B4M) device which  
15  
16 included infrared sensors to measure hemoglobin, a self-inflating cuff for blood pressure measurement,  
17  
18 and an automated reader for urinary protein and glucose through dipsticks. The device facilitated non-  
19  
20 invasive screening of PE and served as an integrated diagnostic and clinical decision support device for  
21  
22 PE [16]. The third example of a device for blood pressure monitoring was a new hypertension detector,  
23  
24 developed by Thakor et al., which was compared in an observational study with other traditional devices  
25  
26 for use in developing countries to support pregnant women at HRPE/E[32]. The new device was found  
27  
28 to be more accurate and easy-to-use than CRADLE VSA and other devices, due to the reduced number  
29  
30 of steps required for use [32]. As a fourth example of a device for blood pressure monitoring was a  
31  
32 hypothetical telemonitoring program [20], which was described in a qualitative study protocol. The  
33  
34 study intended to explore the perspectives, needs, and preferences of a telemonitoring program for  
35  
36 pregnant women at HRPE in a tertiary health facility of Karachi, to inform future implementation.  
37  
38

39  
40 Finally, one prospective experimental study used a wearable device called the F1 smart wristwatch that  
41  
42 included an integrated chip for sensing blood pressure readings and displaying real-time data on the  
43  
44 screen. The smartwatch on the expectant mother's wrist takes blood pressure readings and transfers  
45  
46 them by Bluetooth to their phone at regular intervals to facilitate personal health tracking. The caregiver  
47  
48 can access the expectant mother's records, as well as receive alerts on blood pressure readings [25].  
49

50  
51 Both of these purposes utilized mobile phone applications as an interface to facilitate data collection,  
52  
53 decision making, and communication between health workers and pregnant women. The majority of  
54  
55 these studies used the POTM application [17-24, 30, 31, 33] to facilitate the collection of relevant  
56  
57 clinical data during antenatal visits. The application was used by community health workers in India,  
58  
59 Pakistan, Nigeria, and Mozambique, as part of a CLIP cluster RCT [17, 30]. The POTM platform  
60  
combined two interventions, which were the miniPIERS model and a Phone Oximeter to accurately



1  
2  
3 predict the risk score for pregnant women at HRPE/E in LMICs. The application generated a risk  
4 estimate which enabled community health workers and other healthcare providers (HCPs) to stratify  
5 high-risk pregnant women, escalate care, and make referrals to the facility. In addition, Jonas et al. study  
6 used a mobile application for administrating CRD test for monitoring misfolded protein in the  
7 preeclamptic urine [21]. Finally, the Feroz et al. study protocol described a hypothetical mobile-based  
8 telemonitoring program which would serve as a communication aid between nurses and high-risk  
9 pregnant women[20, 38].

### 17 ***Users of Digital Health Interventions***

18 Most articles involved HCPs (n=17) as the targeted primary users of the DHIs, while only two articles  
19 had pregnant women and caregivers as the primary users of the DHI [20, 25]. The articles described  
20 various healthcare workers as the users of the DHIs, including mid-level HCPs, community-based  
21 HCPs, lady health supervisors, semi-literate volunteers, community health nurses, lady health workers,  
22 midwives, and accredited social health activists. Sixteen articles included information on the training  
23 of patients and HCPs on how to use the DHI, interpret physiological metrics, and take actionable  
24 measures for critical results [16, 18, 19, 21-24, 26-34]. The HCPs received advanced training to  
25 enhance their assessment skills and ability to facilitate the overall management of pregnant women at  
26 HRPE/E. Three articles did not specify the training component for either HCPs or patients [17, 20, 25].

### 38 ***Type of Outcomes Examined***

39 The included articles (n=19) reported on three major outcomes: 1) maternal and neonatal health  
40 outcomes (n=4), 2) usability and acceptability (n=5), and 3) intervention feasibility (n=7)  
41 (Supplementary file 7: Outcomes of DHIs).

### 46 ***Maternal and Neonatal Health Outcomes***

47 Four articles examining maternal and neonatal health outcomes were observational studies (n=3) and  
48 RCTs (n=2) [17, 18, 28, 30]. Maternal health outcomes included magnesium sulfate use, hospital  
49 admissions, CCU admissions, birth preparedness, complication readiness, facility delivery attended by  
50 skilled birth attendants, and adverse maternal outcomes such as an increase in kidney injury, maternal  
51 morbidity, and mortality. Both RCTs reported non-significant findings regarding maternal morbidity  
52 and mortality for participants in the DHI arm [17, 30]. Neonatal health outcomes included stillbirths,  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 fetal and neonatal morbidity, and mortality. Only one of the two RCTs reported a reduction in stillbirths  
4 in the DHI group; however, no impact on neonatal morbidity or mortality was reported for participants  
5 in the DHI group [30].  
6  
7  
8

### 9 Usability and Acceptability

10  
11 Five articles reported on the usability and acceptability of DHIs in LMICs [19, 24-26, 32]. The articles  
12 mentioned pregnant women, caregivers, and HCPs' experience of use of DHIs in LMICs. Usability  
13 outcomes included: trust in technology, ease of use, content richness, perceived usefulness, and user  
14 satisfaction. Musyoka et al. (2019) study found that a 24-hour ambulatory blood pressure monitoring  
15 system had shown great potential for actual adoption in healthcare systems in developing countries,  
16 given its simplicity and affordability [25]. The study found that content richness had a slightly positive  
17 linear effect on perceived ease of use, while there is a slightly negative relationship between content  
18 richness and perceived usefulness [25]. Lim et al. used the computer systems usability questionnaire to  
19 assess the usability of the POTM mHealth application [24]. Nurses and midwives who participated in  
20 the study rated the usability high for the integration of these technologies and thought it would help  
21 their fieldwork. The study found that usability issues were often related to navigation of the app and  
22 phone features such as scroll wheels, touch screen use, etc.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

### 37 Intervention Feasibility and Fidelity

38  
39 Most articles (n=7) reported on the feasibility and fidelity of DHIs for pregnant women at HRPE/E in  
40 LMICs in order to provide evidence on the evaluation of DHIs for replication and scale-up of  
41 successful DHIs [16, 21, 23, 27, 29, 31, 34]. Study outcomes included: fidelity and accuracy of the  
42 CRADLE VSA device, MiniPIERS model development and validation, understanding of enabling and  
43 impeding factors for CLIP trial implementation, experiences of pregnant women with B4M  
44 intervention, and cost-effectiveness of the Congo Red Dot test. One mixed-methods study reported high  
45 fidelity of the implementation of the CRADLE VSA device, with improved HCPs ability to make  
46 clinical decisions, escalate care, and make immediate referrals in case of emergency [34]. The study by  
47 Khowaja et al. (2016) reported factors associated with the feasibility of the CLIP trial implementation  
48 including community mobilization, institutional support, system integration, knowledge gaps, lack of  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 trained personnel, cultural myths and misconceptions, poor health service quality, and high cost of care  
4  
5 [23].  
6  
7  
8

## 9 **Discussion**

### 10 ***Principal Findings***

11  
12 This review summarizes evidence on the existing DHIs to support pregnant women at HRPE/E in  
13 LMICs. Given that most articles (11 out of 19) were published between 2015 and 2020, the novelty of  
14 DHIs use to support pregnant women with HRPE/E was indicated. Only nine unique DHIs were  
15 identified in this review from 19 included articles, reflecting the limited understanding and use of DHIs  
16 to support pregnant women in LMICs. Most included articles used observational and exploratory  
17 research methods to study DHIs. This suggested the need for concerted efforts to learn from small  
18 innovation projects and deployments as outlined in WHO guide on monitoring and evaluation of DHIs  
19 [39]. Most articles in this review did not report information on the blood pressure thresholds, which  
20 limited our understanding of standardized blood pressure thresholds used in LMICs. The explicit  
21 reporting of standardized blood pressure thresholds could help in designing effective clinical decision  
22 support systems for monitoring pregnant women in LMICs [40].  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

### 37 ***Implementation Barriers and Strategies for DHIs***

38  
39 The Microlife CRADLE VSA blood pressure monitoring device has been extensively validated for use  
40 in LMICs for pregnant women [27, 28, 34, 37]. However, HCPs faced several barriers during the  
41 implementation of CRADLE VSA device including lack of supportive supervision for device use, high  
42 staff turnover, and poor availability of the device, poor battery life of device, misleading displays,  
43 broken hand pump, tubing and broken charging ports [34]. Nathan et al. and Vousden et al. suggested  
44 a range of implementation strategies to address known barriers, prior to scale-up, including recognizing  
45 designated device champions who can provide in-depth local training and support for device use,  
46 emphasizing the importance of a device training package (short animated film, interactive sessions,  
47 booklet, and posters), updating training materials to explain the traffic light alert system, providing  
48 chargers in addition to the USB cable, and ensuring an adequate supply of VSA devices [28, 34]. Lim  
49 et al. study mentioned that the general unfamiliarness of using touch screen smart phones was reported  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 as the major barrier faced during the implementation of POTM application[24]. Abejirinde et al. study  
4 trained users on the technical and operational functions of the device to address technical and procedural  
5 issues including software freezes, slow response time, and low user dexterity with operating the device  
6  
7 were two main factors that contributed to delays[16].  
8  
9

### 11 ***Research Gaps and Suggestions for Future Research***

12 *Enabling the use of DHIs by pregnant women as end-users instead of HCPs as end-users:* Most articles  
13 in this review targeted DHIs at HCPs who have less formal training and education, as opposed to studies  
14 conducted in high-income countries where DHIs have been targeted at family physicians and clinicians  
15 who have specialized medical training [6]. This review identified only one study that targeted DHI at  
16 pregnant women for personal health tracking [25]; however, DHIs implemented in high-income  
17 countries are often targeted for use by pregnant women to improve maternal health behaviors and  
18 maternal-fetal health outcomes [41]. Given the increasing cell phone penetration in LMICs [42], there  
19 is an opportunity to use mobile phone technology to target DHIs at the patient level (pregnant women)  
20 to encourage personal health tracking. Yet, health informatics researchers should consider issues of  
21 technological literacy, user characteristics (age, gender, computer skills, experience), cultural factors,  
22 and socioeconomic status when designing and implementing DHIs in the LMIC context [43]. None of  
23 the studies delivered targeted client instructions via a digital platform, in response to abnormal blood  
24 pressure readings or signs and symptoms of PE. In high-income countries, some digital health platforms  
25 have delivered manual or automated targeted instructions to the pregnant women to provide information  
26 about medications, referrals, and diet [44]. LMICs can learn from the experiences of high-income  
27 countries for developing context-specific digital platforms that can facilitate targeted client  
28 communication between providers and pregnant women. Evidence suggests that the targeted client  
29 communication for transmission of health information, health event alerts and reminders, and diagnostic  
30 results have shown positive impacts on health behaviors and health outcomes in high-income countries  
31 [45].  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

54  
55 *Using Multidisciplinary Team Approach for Designing DHIs:* None of the DHIs used a  
56 multidisciplinary team approach for monitoring of pregnant women for PE/E. Blandford et al. suggest  
57 that DHIs should involve collaboration between different cadres of HCPs across all levels of the health  
58  
59  
60

1  
2  
3 system, to achieve the full potential of digital intervention [46]. For instance, a nurse or midwife at a  
4 primary level could communicate about a pregnant women's health condition to a clinician at a  
5 secondary institution to seek recommendations for managing pregnant women at HRPE/E. Murray et  
6 al. suggest that high-quality research in the digital health field requires fertile multidisciplinary  
7 collaborations that draw on insights and experience from multiple fields, including clinical medicine,  
8 health services research, behavioral science, education, engineering, and computer science[47]. Thus,  
9 research aimed at designing and evaluating DHIs to support pregnant women at HRPE/E should draw  
10 insights from collaborators belonging to diverse disciplines including obstetricians and gynecologists,  
11 telemedicine experts, knowledge users, HCPs (nurses, doctors), public health specialists, maternal  
12 health specialist, health services researchers, as well as patient partners.

23  
24 *Exploring Telemedicine Use to Enable Remote Consultation Between Pregnant Women and Healthcare*

25  
26 *Providers:* Most articles used DHIs for the prediction of adverse maternal outcomes, data collection  
27 and decision aid, diagnostic and clinical decision support, and personal health tracking. There is a lack  
28 of evidence on using DHIs for referral coordination, teleconsultation between pregnant women and  
29 HCPs, communication between the HCP and their supervisor, and HCPs' training. Telemedicine has  
30 been extensively used in high-income countries for providing a range of obstetrical services such as  
31 using videoconference to replace in-person visits, implementing at-home monitoring, enabling  
32 consultation with remote specialists, earlier postpartum follow up visits, and access to lactation  
33 consultants [48]. This evidence shows the potential of using telemedicine for pregnant women at  
34 HRPE/E in LMICs to enable remote monitoring and remote consultation between pregnant women and  
35 providers.

36  
37 *Monitoring and Evaluating the Implementation and Effectiveness of DHIs:* Most articles reported on  
38 intervention feasibility, usability, and acceptability outcomes. Two RCTs reported non-significant  
39 findings for maternal morbidity, mortality, and neonatal deaths [17, 30] with only one RCT that reported  
40 a significant difference in stillbirth rate in DHIs group [30]. This suggests the need of conducting more  
41 experimental studies such as RCTs to evaluate the efficacy and effectiveness of diverse DHIs to improve  
42 maternal and child health outcomes. In the review, only one study protocol described the methodology  
43 to conduct an economic evaluation of the CLIP package in South Asian and African countries [22].  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 This shows the paucity of evidence on the economic impact of DHIs to support pregnant women with  
4 PE/E. Ramsey et al. recommend that future clinical trials should incorporate cost-analysis of DHIs as  
5 there is mounting evidence on embedding economic evaluations within clinical trials to build a robust  
6 cost-effectiveness model that has high internal validity and timeliness [49]. The articles included in this  
7 review did not extensively identify facilitators and challenges encountered during the implementation  
8 of DHIs for pregnant women with PE/E in LMICs, unlike many studies conducted in high-income  
9 countries [6]. This review identified only a few facilitators: easy to use technology, trust in technology,  
10 and availability of diagnostic service at the point of care. This indicates the need to examine and report  
11 on enablers and barriers faced when employing DHIs for pregnant women at HRPE/E across the stages  
12 of design, development, implementation, and evaluation.

13  
14 In summary, this scoping review suggests four recommendations for future research: 1) enable the use  
15 of DHIs by pregnant women as end-users to encourage personal health tracking including  
16 individualized patient instructions; 2) consider a multidisciplinary team approach when designing DHIs  
17 for pregnant women at HRPE/E; 3) explore the potential of using telemedicine in LMICs to enable  
18 remote consultation between pregnant women and health provider; 4) conduct further studies including  
19 prospective longitudinal and experimental studies to establish the implementation effectiveness and  
20 efficacy of DHIs to support pregnant women at HRPE; exploratory studies to identify barriers and  
21 enablers associated with the development, implementation, and evaluation of DHIs; and economic  
22 evaluations of DHIs within large clinical trials to identify cost-effective DHIs.

### 23 24 25 **Conclusion**

26 The current evidence base is sparse but shows some potential for the use of different DHIs to support  
27 pregnant women in early diagnosis of PE/E through predicting the risk for adverse maternal health  
28 outcomes and monitoring high-risk pregnant women for PE/E through devices and other DHIs . Limited  
29 evidence exists on types, benefits, cost-effectiveness, and outcomes of DHIs. The weak evidence may  
30 impede the adoption of these promising technologies in community and healthcare settings to support  
31 pregnant women at HRPE/E in LMICs. Future research work should target DHIs at the pregnant women  
32 level to promote personal health tracking with targeted instructions for pregnant women, consider a  
33 multidisciplinary team approach for designing DHIs, explore the role of telemedicine to enable remote  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

ANAM SHAHIL

19

consultation between pregnant women and healthcare providers, and evaluate the implementation and effectiveness of DHIs.

**Acknowledgements**

None

**Conflicts of Interest**

Authors declare no competing interests.

**Contributors**

ASF and ES authors conceptualized and designed the study. ASF screened the articles and performed data extraction, synthesized the data, and drafted the manuscript. NA independently performed screening of articles. ES served as the senior author including participating in the data analysis and providing critical feedback on the manuscript. All authors read and approved the final manuscript.

**Funding**

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

**Data sharing statement**

No additional data available

**Patient consent for publication**

Not required.

**Ethics Statement**

Not required

**Abbreviations**

CLIP: Community-Level Interventions for Preeclampsia

CRD: Congo Red Dot

DHI: Digital Health Intervention

PRE-EMPT: Preeclampsia, Eclampsia Monitoring, Prevention & Treatment

GNI: Gross National Income

HCP: Healthcare Provider

HRPE: High Risk for Preeclampsia

LMIC: Low Middle Income Country

ANAM SHAHIL

NICE: National Institute for Health and Care Excellence

OSF: Open Science Framework

PE/E: Preeclampsia/Eclampsia

PIERS: Preeclampsia Integrated Estimate of RiSk

POTM: PIERS on the Move

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-analyses

RCT: Randomized Controlled Trials

TM: Telemonitoring

VSA: Vital Sign Alert

WHO: World Health Organization



## References

1. Goldenberg, R.L., et al., *Reducing maternal mortality from preeclampsia and eclampsia in low-resource countries—what should work?* 2015. **94**(2): p. 148-155.
2. Khowaja, A.R., et al., *Community's perceptions of pre-eclampsia and eclampsia in Sindh Pakistan: a qualitative study.* 2016. **13**(1): p. 39-44.
3. Howden-Chapman, P., et al., *SDG 3: Ensure healthy lives and promote wellbeing for all at all ages.* 2017: p. 81-126.
4. Van Den Heuvel, J., et al., *Home-based telemonitoring versus hospital admission in high risk pregnancies: a qualitative study on women's experiences.* 2020. **20**(1): p. 1-9.
5. van den Heuvel, J.F., et al., *SAFE@ HOME—Feasibility study of a telemonitoring platform combining blood pressure and preeclampsia symptoms in pregnancy care.* 2019. **240**: p. 226-231.
6. Aquino, M., et al., *Exploring the Use of Telemonitoring for Patients at High Risk for Hypertensive Disorders of Pregnancy in the Antepartum and Postpartum Periods: Scoping Review.* 2020. **8**(4): p. e15095.
7. WHO, *WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience.* Geneva, Switzerland. 2016.
8. Lanssens, D., et al., *Effectiveness of telemonitoring in obstetrics: scoping review.* 2017. **19**(9): p. e327.
9. Rivera-Romero, O., et al., *Mobile health solutions for hypertensive disorders in pregnancy: scoping literature review.* 2018. **6**(5): p. e130.
10. van den Heuvel, J.F., et al., *eHealth as the Next-Generation Perinatal Care: An Overview of the Literature.* J Med Internet Res, 2018. **20**(6): p. e202.
11. de Oliveira, B.G.R.B., *Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist.* 2015.
12. Levac, D., H. Colquhoun, and K.K.J.I.s. O'Brien, *Scoping studies: advancing the methodology.* 2010. **5**(1): p. 69.
13. O'Malley, L.J.I.J.S.R.M., *Scoping studies: towards a methodological framework AU-Arksey, Hilary.* 2005. **8**: p. 19-32.
14. Guo, C., et al., *Challenges for the evaluation of digital health solutions—A call for innovative evidence generation approaches.* npj Digital Medicine, 2020. **3**(1): p. 1-14.
15. Serajuddin, U. and N. Hamadeh, *New World Bank country classifications by income level: 2020–2021.* World Bank Blogs, 2020.
16. Abejirinde, I.O., et al., *Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana.* BMC Pregnancy Childbirth, 2018. **18**(1): p. 209.
17. Bellad, M.B., et al., *Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial.* Pregnancy Hypertens, 2020. **21**: p. 166-175.
18. Bellad, M.B., et al., *Maternal and newborn health in Karnataka state, India: the community level interventions for pre-eclampsia (CLIP) Trial's baseline study results.* 2017. **12**(1): p. e0166623.
19. Dunsmuir, D.T., et al., *Development of mHealth applications for pre-eclampsia triage.* IEEE J Biomed Health Inform, 2014. **18**(6): p. 1857-64.
20. Feroz, A., S. Saleem, and E. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol.* Reprod Health, 2020. **17**(1): p. 135.
21. Jonas, S.M., et al., *Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources.* J Am Med Inform Assoc, 2016. **23**(1): p. 166-73.




- 1
- 2
- 3
- 4 22. Khowaja, A.R., et al., *Economic evaluation of Community Level Interventions for Pre-eclampsia (CLIP) in South Asian and African countries: a study protocol*. Implement Sci, 2015. **10**: p. 76.
- 5
- 6
- 7 23. Khowaja, A.R., et al., *The feasibility of community level interventions for pre-eclampsia in South Asia and Sub-Saharan Africa: a mixed-methods design*. Reprod Health, 2016. **13 Suppl 1**: p. 56.
- 8
- 9
- 10 24. Lim, J., et al., *Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage*. JMIR Mhealth Uhealth, 2015. **3(2)**: p. e37.
- 11
- 12 25. Musyoka, F.M., M.M. Thiga, and G.M. Muketha, *A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care*. Informatics in Medicine Unlocked, 2019. **16**.
- 13
- 14 26. Nathan, H.L., et al., *Evaluation of a novel device for the management of high blood pressure and shock in pregnancy in low-resource settings: study protocol for a stepped-wedge cluster-randomised controlled trial (CRADLE-3 trial)*. 2018. **19(1)**: p. 1-10.
- 15
- 16 27. Nathan, H.L., et al., *An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2*. Blood Press Monit, 2015. **20(1)**: p. 52-5.
- 17
- 18 28. Nathan, H.L., et al., *Early warning system hypertension thresholds to predict adverse outcomes in pre-eclampsia: A prospective cohort study*. Pregnancy Hypertens, 2018. **12**: p. 183-188.
- 19
- 20 29. Payne, B.A., et al., *A risk prediction model for the assessment and triage of women with hypertensive disorders of pregnancy in low-resourced settings: the miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) multi-country prospective cohort study*. PLoS Med, 2014. **11(1)**: p. e1001589.
- 21
- 22 30. Qureshi, R.N., et al., *Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial*. Pregnancy Hypertens, 2020. **22**: p. 109-118.
- 23
- 24 31. Sharma, S., et al., *A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria*. 2017. **17(1)**: p. 238.
- 25
- 26 32. Thakor, N., et al., *Hypertension Detector for Developing Countries*. 2010, Department of Biomedical Engineering, Johns Hopkins University.
- 27
- 28 33. von Dadelszen, P., et al., *The PRECISE (PREgnancy Care Integrating translational Science, Everywhere) Network's first protocol: deep phenotyping in three sub-Saharan African countries*. Reprod Health, 2020. **17(Suppl 1)**: p. 51.
- 29
- 30 34. Vousden, N., et al., *Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial*. BMC Pregnancy Childbirth, 2018. **18(1)**: p. 115.
- 31
- 32 35. Feroz, A., S. Saleem, and E.J.R.H. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol*. 2020. **17(1)**: p. 1-7.
- 33
- 34 36. von Dadelszen, P., et al., *The Community-Level Interventions for Pre-eclampsia (CLIP) cluster randomised trials in Mozambique, Pakistan, and India: an individual participant-level meta-analysis*. The Lancet, 2020. **396(10250)**: p. 553-563.
- 35
- 36 37. Nathan, H.L., et al., *The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings*. Reprod Health, 2018. **15(1)**: p. 5.
- 37
- 38 38. Feroz, A., S. Saleem, and E. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol*. Reproductive health, 2020. **17(1)**: p. 1-7.
- 39
- 40 39. Organization, W.H., *Monitoring and evaluating digital health interventions: a practical guide to conducting research and assessment*. 2016.
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3
- 4 40. Nee, O. and A.J.D.S.S. Hein, Advances in, Book, editor Ger Devlin, *Clinical decision support*
- 5 *with guidelines and Bayesian networks*. 2010: p. 117-137.
- 6 41. Hussain, T., P. Smith, and L.M. Yee, *Mobile Phone–Based Behavioral Interventions in*
- 7 *Pregnancy to Promote Maternal and Fetal Health in High-Income Countries: Systematic*
- 8 *Review*. JMIR mHealth and uHealth, 2020. **8**(5): p. e15111.
- 9 42. Muir, S., *To what extent is mHealth an effective Intervention in LMICs?* 2019.
- 10 43. Feroz, A., M.M. Kadir, and S.J.G.h.a. Saleem, *Health systems readiness for adopting mhealth*
- 11 *interventions for addressing non-communicable diseases in low-and middle-income*
- 12 *countries: a current debate*. 2018. **11**(1): p. 1496887.
- 13 44. Haddad, S.M., R.T. Souza, and J.G. Cecatti, *Mobile technology in health (mHealth) and*
- 14 *antenatal care—searching for apps and available solutions: a systematic review*. International
- 15 *journal of medical informatics*, 2019. **127**: p. 1-8.
- 16 45. Organization, W.H., *WHO guideline: recommendations on digital interventions for health*
- 17 *system strengthening: web supplement 2: summary of findings and GRADE tables*. 2019,
- 18 *World Health Organization*.
- 19 46. Blandford, A., et al., *Seven lessons for interdisciplinary research on interactive digital health*
- 20 *interventions*. 2018. **4**: p. 2055207618770325.
- 21 47. Murray, E., et al., *Evaluating digital health interventions: key questions and approaches*.
- 22 2016, Elsevier.
- 23 48. Weigel, G., B. Frederiksen, and U. Ranji, *Telemedicine and pregnancy care*. The Henry J Kaiser
- 24 *Family Foundation*. February, 2020.
- 25 49. Ramsey, S., et al., *Good research practices for cost-effectiveness analysis alongside clinical*
- 26 *trials: the ISPOR RCT-CEA Task Force report*. 2005. **8**(5): p. 521-533.
- 27
- 28
- 29
- 30
- 31
- 32


### Figure Legends

Figure 1: PRISMA Flow Diagram for Database Search of Studies

Figure 2: Classification of the Included Studies Based on the Purpose of Digital Health Interventions.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47



136/bmjopen-2021-056630 on 8 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

## Appendix I: Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
<b>TITLE</b>			
Title	1	Identify the report as a scoping review.	1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	1
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	2&3
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	3
<b>METHODS</b>			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	3
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	3&4
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	4
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	4
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	4&5
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	5
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	Appendix IV
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	5-6



SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	5
<b>RESULTS</b>			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	4&5
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	Appendix V-VII
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	5-6
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	7-10/ Appendix V-VII
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	7-10
<b>DISCUSSION</b>			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	10-12
Limitations	20	Discuss the limitations of the scoping review process.	12
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	12
<b>FUNDING</b>			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	NA

JBI = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.

\* Where *sources of evidence* (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.

† A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with *information sources* (see first footnote).

‡ The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JBI guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.

§ The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

From: Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med.* 2018;169:467–473. doi: 10.7326/M18-0850.





## Appendix II-Eligibility Criteria

### Question 1: Does this study include humans?

- a) If yes, **INCLUDE**
- b) **EXCLUDE** animal studies/models, non-humans or vertebrae studies

### Question 2: Is the primary language of the study English?

#### Is the primary language of the study English?

- a) If yes, **INCLUDE**
- b) **EXCLUDE** if study is listed as described in a non-English language

### Question 3: Is the article classified as one of the following?

- a) **INCLUDE** all types of study designs including, observational studies, experimental studies, qualitative studies, study protocols, grey literature.
- b) **EXCLUDE:** systematic reviews, meta-analysis, letter to editors, scoping reviews, commentaries, news articles

### Question 4: Does this study examine care provided to pregnant women with Preeclampsia/eclampsia (PE/E)/or at high risk for PE/E (HRPE/E)?

- a) If yes, **INCLUDE**
- b) **EXCLUDE** if study does not focus on care provided to pregnant women with PE/E or at HRPE/E

### Question 5: Does this study examine digital technologies to support pregnant women with preeclampsia/eclampsia or at HRPE/E

- a) **INCLUDE** studies that are focused on use of digital technologies to support pregnant women with PE/E or at HRPE/E. Digital technologies may include:
  - Telephone communication
  - Video communication
  - Text messaging (asynchronous)
  - Email messaging (asynchronous)
  - Portals, apps, and other applications
  - Remote monitoring
  - Devices
  - Predictive models

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
- Provider-provider communication through one of the above modalities
    - Synonyms: digital health, virtual care, virtual visits, eVisits, telehealth, telemedicine, eConsultation, mobile health, mHealth, teleconsultation, teleconference, telecommunications, tele\* (e.g., telepsychiatry, teledermatology, etc), videoconferencing, video visits, phone, telephone, electronic consultation, online consultation, e-mail, text messaging, asynchronous messaging, secure messaging, direct messaging, messaging
  - b) **INCLUDE** studies focused on using digital technologies for early diagnosis, screening, and management of pregnant women with PE/E or HRPE/E.
  - c) **INCLUDE** studies that used digital technologies to support pregnant women with PE/E or at HRPE/E
  - d) **EXCLUDE** studies focused on digital health interventions that do not explicitly focus on pregnant women with PE/E or HRPE/E

#### Question 7: Is this study based on low-and-middle-income contexts?

See list of countries by income classification here: <https://data.worldbank.org/country/XN>

- a) If yes, **INCLUDE**
- b) **INCLUDE** if study focused on high and middle income together.
- c) **EXCLUDE** if based on only high-income country context



**Appendix III -Medical Literature Analysis and Retrieval System Online search strategy**

1. Pregnant Women/
2. exp pregnancy/
3. (pregnan\* adj3 ("at risk" or "at-risk" or "high risk" or "high-risk")).tw,kw.
4. exp Eclampsia/
5. exp Pre-Eclampsia/
6. (Pre Eclampsia or preeclampsia or pre-eclampsia or pre eclampsia or eclampsia or gestosis or proteinuria or toxemia\*).tw,kw.
7. or/1-6
8. Telemedicine/
9. Medical informatics/
10. Digital health.mp.
11. mHealth app.mp.
12. predictive model.mp.
13. CLIP.mp.
14. informatics/
15. exp Telecommunications/
16. Monitoring, Ambulatory/
17. exp Telemetry/
18. Monitoring, Physiologic/
19. exp Computer Communication Networks/
20. Mobile Applications/
21. Smartphone/
22. Cell Phone/
23. (tele-monitor\* or telemonitor\* or telemed\* or tele-med\* or teleinterpret\* or tele-interpret\* or telecomm\* or tele-comm\* or telemetry).tw,kw.
24. (mhealth\* or m-health\* or ehealth\* or e-health\* or telehealth\* or tele-health\*).tw,kw.
25. (mobile adj3 (health\* or technolog\* or app\* or solution\* or phone\* or communicat\*)).tw,kw.
26. (remote\* adj3 (transmi\* or transfer\* or tele\* or monitor\* or consult\* or follow-up or program\* or connect\* or web-base\* or "web base\*" or term)).tw,kw.
27. (monitor\* adj3 (home or remote or distan\* or ambulatory or tele\* or online or on-line or "on line" or phone or digital\* or Skype or electronic\* or implant\* or wireless\* or web-base\* or "web base\*")).tw,kw.
28. (interven\* adj3 (remote\* or distan\* or tele\* or online or on-line or "on line" or phone\* or digital\* or Skype or electronic\* or wireless\*)).tw,kw.
29. (smartphone\* or "smart phone\*" or bluetooth\* or Internet\* or phone\* or text messag\*).tw,kw.
30. ((app or apps or application\*) adj3 (mobile or electronic or software)).tw,kw.
31. ((digital\* or electronic\* or online\* or on-line\* or "on line" or Internet) adj3 (health\* or solution\* or transmit\* or transmiss\* or transfer\* or device\* or connect\*)).tw,kw.
32. (broadband adj3 (device\* or capab\*)).tw,kw.
33. (multi-media\* or multimedia\*).tw,kw.
34. (self monitor\* or self-monitor\*).tw,kw.
35. or/8-34
36. 7 and 35
37. developing countries/
38. low-and-middle-income countries.mp.
39. LMICs
40. Honduras/
41. Angola/
42. Papua New Guinea/
43. Algeria/
44. India/
45. Philippines/

- 1  
2  
3 46. Bangladesh/  
4 47. Kenya/  
5 48. Sao Tome and Principe.mp.  
6 49. Benin/  
7 50. Kiribati.mp.  
8 51. Senegal/  
9 52. Bhutan/  
10 53. Kyrgyzstan/  
11 54. Solomon Islands.mp.  
12 55. Bolivia/  
13 56. Laos/  
14 57. Sri Lanka/  
15 58. Cabo Verde/  
16 59. Lesotho/  
17 60. Tanzania/  
18 61. Cambodia/  
19 62. Mauritania/  
20 63. Timor-Leste/  
21 64. Cameroon/  
22 65. Micronesia/  
23 66. Tunisia/  
24 67. Comoros/  
25 68. Moldova/  
26 69. Ukraine/  
27 70. "Democratic Republic of the Congo"/  
28 71. Mongolia/  
29 72. Uzbekistan  
30 73. Cote d'Ivoire/  
31 74. Morocco/  
32 75. Vanuatu/  
33 76. Djibouti/  
34 77. Myanmar/  
35 78. Vietnam/  
36 79. Egypt/  
37 80. Nepal/  
38 81. West Bank and Gaza.mp.  
39 82. El Salvador/  
40 83. Nicaragua/  
41 84. Zambia/  
42 85. Eswatini/  
43 86. Nigeria/  
44 87. Zimbabwe/  
45 88. Ghana/  
46 89. Pakistan/  
47  
48 90. (Angola or Honduras or Papua New Guinea or Algeria or India or Philippines or Bangladesh or Kenya or Sao  
49 Tome and Principe or Benin or Kiribati or Senegal or Bhutan or Kyrgyz Republic or Solomon Islands or Bolivia or  
50 Lao PDR or Sri Lanka or Cabo Verde or Lesotho or Tanzania or Cambodia or Mauritania or Timor-Leste or  
51 Cameroon or Micronesia or Tunisia or Comoros or Moldova or Ukraine or Democratic Republic of the Congo or  
52 Mongolia or Uzbekistan or Cote d'Ivoire or Morocco or Vanuatu or Djibouti or Myanmar or Vietnam or Egypt  
53 or Nepal or West Bank and Gaza or El Salvador or Nicaragua or Zambia or Eswatini or Nigeria or Zimbabwe or  
54 Ghana or Pakistan).tw,kw.  
55 91. or/37-90  
56 92. 36 and 91  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 93. exp animals/ not humans.sh.
- 94. 92 not 93
- 95. remove duplicates from 94

For peer review only

## Appendix IV- Data abstraction form

1. Author, year
2. Journal
3. Study design (observational, experimental, protocol paper)
4. Study setting/country (LMICs)
5. Study population/Health Condition (PW at HRPE/E)
6. Study Objective
7. Number of participants (Sample)
8. Study period
  - Duration of intervention
  - Duration of data collection
9. Digital health intervention (DHI) used
  - Type of DHI (Predicative model, mHealth applications, devices)
  - Intervention validation (Yes/No)
  - Targeted primary user of intervention (health care provider/pregnant women)
  - User training on use of digital health intervention (Yes/No)
  - Function of digital health intervention
10. Study outcomes
  - Maternal and fetal health outcomes
  - Intervention feasibility/usability/fidelity/acceptability
11. Framework/model used
12. Study limitations
13. Comments

## Appendix V-Overview of included articles

Reference	Year	Study Title	Type of Study Design	Objective	Setting	N	Health Condition	Purpose of Digital Health Intervention
Musyoka et al. [25]	2019	A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care	Prospective experimental study	The study sought to implement a 24-hour ambulatory blood pressure monitoring solution for preeclampsia management, using a smartwatch in conjunction with a mobile and cloud-based application.	Kenya	N=30	preeclampsia	Monitoring
Lim et al. [24]	2015	Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage	Observational	The aim of this study was to assess the usability of PIERS on the Move PotM (with mid-level health workers) for iteratively refining the system.	South Africa	N=37	preeclampsia	Predicting
Vousden et al. [34]	2018	Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial	Observational	Prior to the CRADLE 3 trial start, a mixed-methodology feasibility study was undertaken to finalise the intervention and implementation processes which were guided by the Expert Recommendations for Implementing Change (ERIC) project	Zimbabwe, Ethiopia, India	Number of HCP trained=204	Preeclampsia, eclampsia and shock	Monitoring
Nathan et al. [26]	2018	The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings	Observational	This qualitative study aimed to determine the usability, feasibility and acceptability of the CRADLE VSA among a variety of users and in diverse socio-economic settings, considering these five clusters of influence. This will inform future device modifications and successful dissemination of the CRADLE VSA for routine use.	India, Mozambique, Nigeria and South Africa	N=205	Preeclampsia and shock	Monitoring
Feroz et al. [20]	2020	Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a	Protocol paper	The study aims to explore the perspectives, preferences, and needs of telemonitoring (TM) for pregnant women at HRPE in Karachi, to inform future implementation strategies.	Pakistan	N=30	Preeclampsia	Monitoring

		tertiary health facility of Karachi: a qualitative study protocol						
Dunsmuir et al. [19]	2014	Development of mHealth Applications for Pre-Eclampsia Triage	Observational	This paper describes the design process of two versions of the POTM application, the original version application referred to as POTM), and a simplified, community-based version for the Community Level Interventions for Pre-eclampsia cluster randomized controlled trial (application referred to as CLIP POTM),	Nigeria, Mozambique, Pakistan, and India	Projected +30,000 pregnant women  500 community HCPs	Preeclampsia	Predicting
Jonas et al. [21]	2016	Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources	Prospective experimental study design	The study proposes an innovative mobile health (mHealth) solution that enables the quantification of the congo red dot test as a batch laboratory test, with minimal cost and equipment.	Resource poor settings	N=273	preeclampsia	Monitoring
Thakor et al. [32]	2009	Hypertension Detector for Developing Countries	Observational	A prototype of a low-cost device engineered specifically for semi-literate volunteers in developing countries has been created.	Africa, Southern Asia, and the Middle East	-	Preeclampsia	Monitoring
Nathan et al. [27]	2015	An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2	Observational	The study aims to assess the accuracy of the Microlife 3AS1-2 blood pressure device in pregnancy and pre-eclampsia in a low-resource setting.	South Africa	N=45	Preeclampsia	Monitoring
Nathan et al. [28]	2018	Early warning system hypertension thresholds to predict adverse outcomes	Observational	The study aims to evaluate the association between blood pressure (BP) measurements and adverse outcomes in women with pre-eclampsia.	South Africa	N= 1547	Preeclampsia	Monitoring

		in pre-eclampsia: A prospective cohort study							
1									
2	Payne et al. [29]	2014	A Risk Prediction Model for the Assessment and Triage of Women with Hypertensive Disorders of Pregnancy in Low-Resourced Settings: The miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) Multi-country Prospective Cohort Study	Observational	The objective of the miniPIERS study was to develop and validate a simplified clinical prediction model for adverse maternal outcomes among women with HDP for use in community and primary health care facilities in LMICs.	LMICs	N= 2,133	Preeclampsia	Predicting
3									
4									
5									
6									
7									
8									
9									
10									
11									
12									
13									
14									
15									
16	Bellad et al. [17]	2020	Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial	Experimental study (RCT)	The objective of the Community-Level Interventions for reeclampsia (CLIP) India cluster randomised controlled trial (cRCT) was to test the hypothesis that implementing community-level, evidence-based care focused on pregnancy hypertension would reduce all-cause maternal, fetal and newborn mortality and major morbidity, without causing harm	India	N=14,783 pregnancies	Preeclampsia	Monitoring and Predicting
17									
18									
19									
20									
21									
22									
23									
24	Qureshi et al. [30]	2020	Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial	Experimental study (RCT)	The aim of the Community-Level Interventions for Pre-eclampsia (CLIP) cluster randomised controlled trial (cRCT) in Sindh Province, Pakistan was to reduce maternal and perinatal mortality and major morbidity by 20% or more in intervention (vs. control) clusters, through a community-level intervention to address triage, (initial) treatment, and transport (to facility) of women with pregnancy hypertension.	Pakistan	N= 35,974 women	Preeclampsia	Monitoring and Predicting
25									
26									
27									
28									
29									
30									
31									
32									
33	Khowaja et al [22]	2015	Economic evaluation of Community Level Interventions for Pre-eclampsia (CLIP) in South Asian and African countries: a study protocol	Protocol paper	The study aims to conduct an economic evaluation alongside of the CLIP Trial, to inform decision makers not only of clinical outcomes but the cost required to obtain those outcomes.	Nigeria, Mozambique, Pakistan, and India	N= 154,000	Preeclampsia	Monitoring
34									
35									
36									
37									
38									
39									
40	Khowaja et al [23]	2016	The feasibility of community level	Observational study	The study aimed to describe the health system, identify community and individual barriers and facilitators that influence care of pregnant women	Nigeria, Mozambique,	N= 337 (health facilities)	Preeclampsia	Monitoring
41									
42									
43									
44									
45									
46									

		interventions for pre-eclampsia in South Asia and Sub-Saharan Africa: a mixed-methods design		in the community, in preparation for the conduct of a community-based cluster randomized trial	Pakistan, and India	N= 100 (IDIs) N= 123 (FGDs)		
Von Dadelzen et al. [33]	2020	The PRECISE (PREgnancy Care Integrating translational Science, Everywhere) Network's first protocol: deep phenotyping in three sub-Saharan African countries	Protocol paper	This paper describes the protocol that underpins the clinical research activity of the Network, so that the investigators, and broader global health community, can have access to 'deep phenotyping' of women as they advance through pregnancy to the end of the puerperium.	Gambia Kenya Mozambique	N= 600 (each country)	Preeclampsia, and eclampsia	Monitoring
Abejirinde et al [16]	2018	Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana	Observational	This paper therefore explores the experiences of women exposed to the B4M device, to answer the research questions: i) How did women experience the use of Bliss4Midwives during their routine antenatal care consultations? ii) What influence did Bliss4Midwives have on woman-provider relationships and on ANC service utilization?	Ghana	N=30	preeclampsia, gestational diabetes and anaemia	Monitoring
Bellad et al [18]	2017	Maternal and Newborn Health in Karnataka State, India: The Community Level Interventions for Pre-Eclampsia (CLIP) Trial's Baseline Study Results	Observational	To describe baseline demographics and health outcomes prior to initiation of the CLIP trial and to improve knowledge of population-level health, in particular of maternal and neonatal outcomes related to hypertensive disorders of pregnancy, in northern districts the state of Karnataka, India.	India	N= 5,469	Hypertension disorders of pregnancy, preeclampsia	Monitoring and Predicting
Sharma et al [31]	2017	A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria	Observational	To evaluate implementation processes of the complex CLIP intervention, assess mechanisms of impact and identify emerging unintended causal pathways.	Nigeria	N= 32,785	preeclampsia	Monitoring and Predicting



Appendix VI- Digital Health Intervention Characteristics

Reference	Digital health intervention	Validated	Intervention use for	Technological component(s)	Targeted primary user	User training
Musyoka et al. [25]	24-hour ambulatory blood pressure monitoring system	Validated	Blood pressure data collection	F1 smart wristwatch Blood Pressure Monitoring Mobile Application Cloud Data center Caregiver's smartphone	Expectant mother and the caregiver	Not specified
Lim et al. [24]	Pre-eclampsia Integrated Estimate of RiSk (PIERS) on the Move (PotM)	Not specified	Demographics (gestational age at presentation), clinical signs (blood pressure, SPO2 and dipstick proteinuria), and symptoms (chest pain or dyspnoea, headache or visual disturbances, vaginal bleeding with abdominal pain)	mHealth platform	Mid-level health workers	Yes
Vousden et al. [34]	CRADLE (Community blood pressure monitoring in Rural Africa & Asia: Detection of	Validated	Measures blood pressure, pulse and calculates the mothers risk of shock  A traffic light Early Warning System display alerts users to	Microlife CRADLE VSA device	Healthcare providers	Yes

136/bmjopen-2021-056130 on 8 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

	underlying pre-Eclampsia and shock) Vital Sign Alert		abnormalities in the vital signs results.			
Nathan et al. [26]	Microlife® CRADLE (Community blood pressure monitoring in Rural Africa & Asia: Detection of underlying pre-Eclampsia and shock) Vital Signs Alert (VSA)	Validated	Device accurately measures blood pressure and pulse. Traffic lights within the device help healthcare workers identify women who need additional treatment for these conditions	Microlife® CRADLE VSA device	Healthcare providers	Yes
Feroz et al. [20]	Hypothetical elemonitoring program	Not specified	Blood pressure measurement	-	Pregnant women and caregiver	-
Dunsmuir et al. [19]	MiniPIERS AND FullPIERS models  Two versions of the POTM application, 1) Original version (application referred to as POTM), 2)Simplified, community-based version for the Community Level Interventions for Pre-eclampsia cluster randomized controlled trial (application referred to as CLIP	Not specified	Mean BP, SpO2, gestational age, proteinuria, symptoms.	Smartphone, mobile health applications (POTM/CLIP POTM), Research electronic data capture server	Community-based health care providers	Yes

	POTM)					
Jonas et al. [21]	Smartphone-based diagnostic test (Congo Red Dot) for preeclampsia	Validated	urine markers	mHealth solution for administering the Congo Red Dot (CRD) test	Modestly trained personnel	Yes
Thakor et al. [32]	New device (Hypertension Detector for Developing Countries), intraarterial, sphygmomanometers, assorted automatic blood pressure devices, and proteinuria measurement	Not specified	Blood pressure measurement	-	Semi-literate volunteers with minimal training	Yes
Nathan et al. [27]	Microlife 3AS1-2 blood pressure device	Validated	Measures blood pressure	Device	Staff with minimal training	Yes
Nathan et al. [28]	CRADLE Vital Signs Alert (VSA)	Validated	Measures BP and pulse to facilitate prompt recognition of abnormalities in vital signs	Device Traffic light early warning system	Healthcare providers	yes
Payne et al. [29]	miniPIERS risk prediction model	Validated	miniPIERS (measures demographics, symptoms and signs).	Mobile health application	Mid-level health workers	yes
Bellad et al. [17]	CLIP intervention package included miniPIERS model, PIERS On the Move (POM) tool, and	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	mobile-based CLIP POM mobile health application (app),	Community health workers	yes

	Microlife BP 3AS1-2 device			central REDCap server		
Qureshi et al. [30]	CLIP intervention package included miniPIERS model, Microlife BP 3AS1-2 device and PIERS On the Move (POM) mobile health (mHealth) application	Validated	BP measurement and pulse oximetry	POM mHealth application	Lady health workers	Yes
Khowaja et al [22]	CLIP intervention package PIERS On the Move (POM) mobile health (mHealth) application	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	POM mHealth application	Community-based health care providers	Yes
Khowaja et al [23]	CLIP intervention package PIERS On the Move (POM) mobile health (mHealth) application	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	POM mHealth application	Community-based health care providers	Yes
Von Dodelszen et al. [33]	CRADLE BP device, pulse oximetry and TraCer platform, POM mHealth application	Validated	CRADLE VSA semi-automated and validated BP device will be used for all clinical measurements of blood pressure (BP) in the study pulse oximetry  POM platform to provide time-of-disease	POM mHealth application	Healthcare providers	Yes

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

			risk estimates to hypertensive pregnant women using PIERS models			
Abejirinde et al [16]	Bliss4Midwives' (B4M)	Not specified	non-invasive device for measuring haemoglobin via infrared sensors mounted on a finger clip; a self-inflating blood pressure cuff; and an automated reader for urinary protein and glucose through dipsticks.	Data from all diagnostic devices are automatically or manually linked to an android tablet equipped with decision support algorithms	Midwives and community health nurses	Yes
Bellad et al [18]	Community Level Interventions for Pre-Eclampsia (CLIP) Package	Not specified	Measuring blood pressure	mHealth platform	Community-based health activists ASHAs	Not specified
Sharma et al [31]	Community Level Interventions for Pre-Eclampsia (CLIP) Package	Not specified	Blood pressure measurement	PIERS On the Move (POM) mHealth tool, Microlife VSA blood pressure device	Community health workers	Yes

## Appendix VIII- Outcomes of digital health interventions

Reference	Study Title	Study outcomes pertaining to digital health intervention use	Framework/model used
<b>Maternal and fetal health outcomes (4 studies)</b>			
Nathan et al. (2018) [28]	Early warning system hypertension thresholds to predict adverse outcomes in pre-eclampsia: A prospective cohort study	Of 1547 women with pre-eclampsia, 33.0% of women triggered a red light on admission and 78.6% at their highest BP. Severe hypertension and adverse outcomes were common across yellow and red categories. Comparing admission red to yellow lights, there was a significant increase in kidney injury (OR 1.74, CI 1.31–2.33, test p=.003), magnesium sulfate use (OR 3.40, CI 2.24–5.18, p < .001) and CCU admission (OR 1.50, CI 1.18–1.91, p < .001), but not for maternal death, eclampsia, extended perinatal death or preterm delivery.	No framework described
Bellad et al. (2020) [17]	Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial	The primary outcome did not differ between intervention and control arms (adjusted odds ratio (aOR) 0.92 [95% confidence interval 0.74, 1.15]; p = 0.47; intraclass correlation coefficient 0.013). There was no intervention-related safety concerns following administration of either methyldopa or MgSO <sub>4</sub> , and 401 facility referrals. Compared with intervention arm women without CLIP contacts, those with ≥8 contacts suffered fewer stillbirths (aOR 0.19 [0.10, 0.35]; p < 0.001), at the probable expense of survivable neonatal morbidity (aOR 1.39 [0.97, 1.99]; p = 0.072).	No framework described
Qureshi et al. (2020) [30]	Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial.	The primary outcome did not differ between intervention (26.6%) and control (21.9%) clusters (adjusted odds ratio, aOR, 1.20 [95% confidence interval 0.84- 1.72]; p = 0.31). There was reduction in stillbirths (0.89 [0.81-0.99]; p = 0.03), but no impact on maternal death (1.08 [0.69, 1.71]; p = 0.74) or morbidity (1.12 [0.57, 2.16]; p = 0.77); early (0.99 [0.82-1.09]; p = 0.46) or late neonatal deaths (1.23 [0.97-1.55]; p = 0.09); or neonatal morbidity (1.22 [0.77, 1.96]; p = 0.40). Improvements in outcome rates were observed with 4–7 (p = 0.015) and ≥8 (p < 0.001) (vs. 0) CLIP contacts.	No framework described
Bellad et al. (2017) [18]	Maternal and newborn health in Karnataka state, India: the community level interventions for pre-eclampsia (CLIP) Trial's baseline study results	A majority of the women reported institutional deliveries (96.0%), largely attended by skilled birth attendants. The maternal mortality ratio of 104 (per 100,000 livebirths) was observed during this study, neonatal mortality ratio was 25 per 1,000 livebirths, and perinatal mortality ratio was 50 per 1,000 livebirths. Despite a high number of institutional deliveries, rates of stillbirth were 2.86%.	No framework described
<b>Usability and acceptability (5 studies)</b>			

Musyoka et al. (2019) [25]	A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care. Informatics in Medicine Unlocked.	Content richness has a slightly positive linear effect on Perceived Ease of Use, while there is a slightly negative relationship between Content Richness and Perceived Usefulness. Overall, the 24-hour ambulatory blood pressure monitoring system has shown great potential for actual adoption in healthcare systems in developing countries, given its simplicity and affordability.	Technology Acceptance Model
Lim et al. (2015) [24]	Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage.	Overall, users felt the app was usable using the Computer Systems Usability Questionnaire; median (range) values for Study 1 = 2 (1-6) and Study 2 = 1 (1-7). Usability problems were often related to mobile phone features (eg, scroll wheels, touch screen use).	LambdaNative framework for app development
Nathan et al. (2018) [26]	The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings.	Most HCWs perceived the CRADLE device to be easy to use and accurate. The traffic lights early warning system was unanimously reported positively, giving HCWs, Pregnant women and families understanding of vital signs and confidence with decision-making. Some described manual inflation as tiring, particularly when measuring vital signs in obese and hypertensive women (n=4) and a few South African HCWs distrusted the device's accuracy (n =7).	Diffusion of innovation model Three delay model
Thakor et al. (2010) [32]	Hypertension Detector for Developing Countries.	The study developed a prototype of a low-cost device engineered specifically for semi-literate volunteers in developing countries. Preliminary testing has shown reliable hypertension detection and plans have been made for field testing in rural communities this August 2010 in Nepal.	No framework described
Dunsmuir et al (2014) [19]	Development of mHealth applications for pre-eclampsia triage. IEEE J Biomed Health Inform.	The paper outlines the POTM application development process. The paper concludes that the successful development of an mHealth tool, must consider the user and the setting in which it is deployed. CLIP POTM began with a single specification document, but study discovered differing requests from the different countries with their cultural differences, leading to modified application versions for each country.	LambdaNative Framework for developing application
<b>Intervention Feasibility and Fidelity (7 studies)</b>			
Vousden et al (2018) [34]	Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial	Intervention was implemented with high fidelity (85% of HCP trained, n=204). Results indicated a good understanding of device use with 75% of participants scoring >75% (n=97; 90% of those distributed). Interviews with HCPs reported that the intervention improved capacity to make clinical decisions, escalate care and make appropriate referrals.	Medical Research Council framework and logic model
Khowaja et al (2016) [23]	The feasibility of community level interventions for pre-eclampsia in South Asia and	The study highlight enabling factors including need for community mobilization, awareness raising programs, institutional support, community safety nets for	Normalization process theory

	Sub-Saharan Africa: a mixed-methods design.	emergency funds, and system integration. Whereas, impeding factors included delays in care seeking, knowledge gaps, lack of trained human resource, cultural myths and misconceptions, high cost of care, and poor health service quality.	
Abejirinde et al (2018) [16]	Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana.	Pregnant women generally valued the availability of diagnostic services at the point-of-care. The intervention made women feel listened to and cared for. Process outcomes of the B4M encounter also showed that it was perceived as improving the skills and knowledge of the health workers which facilitated trust in diagnostic recommendations and was therefore believed to motivate referral compliance.	No framework described
Sharma et al (2017) [31]	A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria.	This paper offers robust measures of the process indicators, external validity of conclusions about effectiveness can best be complemented by efficacy studies using a RCT. The methodology allows to examine the internal validity of the efficacy of the intervention by assessing the implementation (quantity and quality) of what is delivered.	Logic model, Diffusions of innovations and realist evaluation theories
Nathan et al (2015) [27]	An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2	The Microlife 3AS1-2 device achieved an overall B/A grade in pregnancy (including pre-eclampsia), passing the British Hypertension Society protocol requirements and achieving the International Organization for Standardization standard with a mean difference and SD of $-3.8 \pm 7.3$ and $-1.5 \pm 6.2$ mmHg for systolic and diastolic pressures, respectively. The device can be recommended for use in pregnancy, including preeclampsia. Also, it fulfils the requirements of WHO for an automated blood pressure device suitable for use in a low-resource setting.	No framework described
Payne et al (2014) [29]	A risk prediction model for the assessment and triage of women with hypertensive disorders of pregnancy in low-resourced settings: the miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) multi-country prospective cohort study.	The miniPIERS model was well-calibrated and had an area under the receiver operating characteristic curve (AUC ROC) of 0.768 (95% CI 0.735–0.801) with an average optimism of 0.037. External validation AUC ROC was 0.713 (95% CI 0.668–0.768). A predicted probability $\geq 25\%$ to define a positive test classified women with 85.5% accuracy. The miniPIERS model shows reasonable ability to identify women at increased risk of adverse maternal outcomes associated with the hypertensive disorders of pregnancy	Three delay model
Jonas et al. (2016) [21]	Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources.	The results suggests that combining smartphone-based image analysis with molecular-specific disease features represents a cost-effective application of mHealth that has the potential to fill gaps in access to health care solutions that are critical to reducing adverse events related to PE in resource-poor settings	No framework described



For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
<b>TITLE</b>			
Title	1	Identify the report as a scoping review.	1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	1
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	2&3
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	3
<b>METHODS</b>			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	3
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	3&4
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	4
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	4
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	4&5
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	5
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	Appendix IV
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	5-6



SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	5
<b>RESULTS</b>			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	4&5
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	Appendix V-VII
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	5-6
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	7-10/ Appendix V-VII
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	7-10
<b>DISCUSSION</b>			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	10-12
Limitations	20	Discuss the limitations of the scoping review process.	12
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	12
<b>FUNDING</b>			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	NA

JBI = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.

\* Where *sources of evidence* (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.

† A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with *information sources* (see first footnote).

‡ The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JBI guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.

§ The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

From: Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med.* 2018;169:467–473. doi: [10.7326/M18-0850](https://doi.org/10.7326/M18-0850).



# BMJ Open

## Exploring digital health interventions for pregnant women at high risk for preeclampsia and eclampsia in low-and-middle-income countries: a scoping review

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-056130.R1
Article Type:	Original research
Date Submitted by the Author:	06-Dec-2021
Complete List of Authors:	Shahil Feroz, Anam; Aga Khan University, Community Health Sciences; University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Afzal, Noreen; The Aga Khan University Medical College Pakistan, Dean's Office Seto, Emily; University of Toronto, Institute of Health Policy, Management and Evaluation; University Health Network, Centre for Global eHealth Innovation
<b>Primary Subject Heading</b>:	Health informatics
Secondary Subject Heading:	Reproductive medicine, Public health, Research methods
Keywords:	PUBLIC HEALTH, Prenatal diagnosis < OBSTETRICS, Health informatics < BIOTECHNOLOGY & BIOINFORMATICS, Telemedicine < BIOTECHNOLOGY & BIOINFORMATICS

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

## Title of Scoping Review

Exploring digital health interventions for pregnant women at high risk for preeclampsia and eclampsia in low-and-middle-income countries: a scoping review

a. Author and Co-author's name

1. Anam Shahil Feroz [anam.shahil@mail.utoroto.ca](mailto:anam.shahil@mail.utoroto.ca)
2. Noreen Afzal [noreen.afzal100@gmail.com](mailto:noreen.afzal100@gmail.com)
3. Emily Seto [Emily.seto@utoronto.ca](mailto:Emily.seto@utoronto.ca)

b. The name, postal address, e-mail, telephone, and fax numbers of the corresponding author.

Name: Anam shahil Feroz

Affiliations:

- Aga Khan University, Community Health Sciences Department
- University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Toronto, ON, CAN M5T 3M6

Email: [anam.shahil@mail.utoronto.ca](mailto:anam.shahil@mail.utoronto.ca)

Phone: +1647 679 6702

c. The full names, institutions, city, and country of all co-authors.

1. Anam Shahil Feroz

Affiliations:

- Aga Khan University, Community Health Sciences Department, Karachi, Sindh Pakistan
- University of Toronto Institute of Health Policy Management and Evaluation, Dalla Lana School of Public Health Toronto, Ontario, Canada

2. Noreen Afzal

Affiliations:

- The Aga Khan University Medical College Pakistan, Dean's Office Karachi, Sindh Pakistan

3. Emily Seto

Affiliations:

- University of Toronto, Institute of Health Policy, Management and Evaluation Toronto, Ontario, Canada
- University Health Network, Centre for Global eHealth Innovation Toronto, Ontario, Canada

d. Up to five keywords or phrases suitable for use in an index (it is recommended to use MeSH terms).

Digital health interventions, pregnant women, preeclampsia, eclampsia, low-and-middle-income countries, predictive models, mHealth applications, devices

e. Word count - excluding title page, references, figures and tables: 6247

## Abstract

**Objective:** To explore digital health interventions that have been used to support pregnant women at high risk for preeclampsia/ eclampsia (HRPE/E) in low-middle-income countries (LMICs).

**Design:** Scoping review

**Data Source:** EMBASE, MEDLINE, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and CINAHL were searched between January 1, 2000, and October 20, 2020.

**Eligibility criteria:** The review included original research studies that were published in English, involved pregnant women at HRPE/E, and implemented digital health interventions for PE/E in LMICs.

**Data extraction and synthesis:** Two reviewers independently completed the data extraction for each of the 19 final articles. An inductive approach was used to thematically organize and summarize the results from the included articles.

**Results:** A total of 19 publications describing seven unique studies and nine different digital health interventions were included. Most studies were conducted in South Asia and Sub-Saharan Africa (n=16). Of nine unique digital health interventions, two served the purpose of predicting risk for adverse maternal health outcomes while seven focused on monitoring high-risk pregnant women for PE/E. Both of these purposes utilized mobile phone applications as interface to facilitate data collection, decision making, and communication between health workers and pregnant women. The review identified key functions of interventions including data collection, prediction of adverse maternal outcomes, integrated diagnostic and clinical decision support, and personal health tracking.. The review reported three major outcomes: maternal health outcomes including maternal and neonatal morbidity and mortality (n=4); usability and acceptability including ease-of-use, and perceived usefulness, (n=5);and intervention feasibility and fidelity including accuracy of device, and intervention implementation (n=7).

**Conclusion:** Although the current evidence base shows some potential for the use of digital health interventions for PE/E, more prospective experimental and longitudinal studies are needed prior to recommending the use of digital health interventions for PE/E.

## Strengths and limitations of this study



- 1  
2  
3 1. First scoping review to explore the use of digital health interventions (DHIs) in low-middle-  
4 income countries (LMICs) to support pregnant women at high risk for preeclampsia/eclampsia  
5 (HRPE/E).  
6  
7
- 8  
9 2. The scoping review has identified several gaps in the area of DHIs use for PE/E in LMICs  
10 which can be explored through future research.  
11  
12
- 13 3. The high heterogeneity of the DHIs and study outcomes limited the interpretation of the studies  
14 through quantitative analysis.  
15  
16
- 17 4. This review only included peer-reviewed articles and papers published in the English  
18 Language.  
19  
20
- 21 5. The review did not include information that may have been found in other databases and sources  
22 (abstracts, reviews, conference proceedings, opinion papers, books).  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Introduction

Approximately 16% of all maternal deaths in low-and-middle-income countries (LMICs) are attributable to preeclampsia/eclampsia (PE/E) [1]. High maternal mortality from PE/E results from: 1) lack of early identification and treatment of pregnant women, 2) difficulties in reaching treatment centers and, 3) poor health-seeking behaviors linked with low patient education [2]. To meet the United Nations Sustainable Developmental Goal target 3.1 of reducing the maternal mortality ratio to less than 70/100,000 live births by 2030, innovations are required to decrease PE/E-related mortality [3].

The most effective strategies to ensure early diagnosis and management of PE/E include self-monitoring of blood pressure, use of magnesium sulfate therapy, proteinuria determinations, and timely delivery [1]. International guidelines including the European Society of Hypertension, American Heart Association, National Institute for Health and Care Excellence, and American Society of Hypertension guidelines, recommend self-monitoring for PE symptoms and recording of blood pressure for pregnant women at high risk for preeclampsia and eclampsia (HRPE/E) because of their potential benefits such as effective control of blood pressure, early risk identification, and treatment, and cost-savings due to fewer hospital visits [4-6]. Self-monitoring also has a role in preventing conditions like white coat hypertension and masked hypertension in pregnant women at HRPE/E. The World Health Organization (WHO) suggests home blood pressure monitoring for pregnant women at HRPE/E to detect changes in blood pressure between antenatal visits and to ensure care continuity [7].

Digital health interventions (DHIs) are increasingly being used to support pregnant women at HRPE/E for remote monitoring of blood pressure and symptoms. To date, four reviews explored the use of digital tools for remote monitoring of pregnant women at HRPE/E. In 2020, Aquino et al. reported 16 unique, feasible, and cost-effective telemonitoring interventions to support pregnant women with hypertensive disorder of pregnancy [6]. However, the review mainly focused on telemonitoring interventions for remote blood pressure monitoring of pregnant women. The review also primarily identified studies from high-income countries like the UK, USA, and Belgium [6]. Lanssens et al. (2017) reported 14 studies from 1988 to 2010 that used telemonitoring interventions for pregnant women during the prenatal period[8]. This review, however, used a narrow time range and focused on telemonitoring solutions implemented in high-income countries for pregnant women at high risk for gestational diabetes and

1  
2  
3 preterm labor. In addition, the included studies had a high methodological risk of bias. When only  
4 studies with low risk of bias were considered, the added value of telemonitoring became less  
5 pronounced [8]. Rivera-Romero et al. (2018) captured only 11 studies conducted in high-income  
6 countries, on mobile health (mHealth) interventions for the hypertensive disorder of pregnancy[9]. The  
7 included studies showed positive results in the improvement of maternal health and acceptability of  
8 solutions, although most of the studies involved a small number of participants, and none were complete  
9 clinical studies [9]. Van den Heuvel et al. (2018) reported 12 studies on the use of telemonitoring and  
10 teleconsulting interventions to improve pregnancy care generally [10]. The review did not focus on the  
11 use of eHealth for the hypertensive disorder of pregnancy and generally included all aspects of perinatal  
12 care.  
13

14  
15 These four reviews provided foundational information on the use of telemonitoring to support high-risk  
16 pregnant women in antepartum and postpartum period. However, quality evidence on the appropriate  
17 use of DHIs to support pregnant women at HRPE/E in LMIC is scarce. None of the reviews extensively  
18 documented the use of DHIs in LMICs for the early diagnosis and management of pregnant women at  
19 HRPE/E. This gap highlights the need to explore the potential role of DHIs to support pregnant women  
20 at HRPE/E in LMICs. This review aims to systematically explore the available literature on the use of  
21 DHIs to support early detection and management of PE/E in LMICs.  
22  
23

## 24 **Methods**

25  
26 The “Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping  
27 Reviews” (PRISMA-ScR) checklist was used to guide the design and reporting of this scoping review  
28 [11]. . The review was registered in the Open Science Framework (OSF) - Center for Open Science on  
29 Oct 19, 2020 (Registration link: <https://osf.io/gncvj>). The review was guided by the methodological  
30 framework by Levac et al. [12] and Arksey et al. [13] to examine articles describing the use of digital  
31 health solutions to support early detection and management of PE/E in LMICs.  
32  
33

## 34 ***Identifying Research Question***

35  
36 The main research question for this scoping review is: What is known in the literature about DHIs that  
37 have been used to support pregnant women at HRPE/E in LMICs?  
38  
39

1  
2  
3 Our study has used the broad population, concept and context (PCC) framework recommended by the  
4 Joanna Briggs Institute for Scoping Reviews. The operationalization of PCC framework for our scoping  
5 review include: population (pregnant women at HRPE/E), concept (DHIs), and context (LMICs).  
6  
7  
8

### 9 ***Eligibility Criteria***

10  
11 The review included studies that involved pregnant women at HRPE/E and implemented the digital  
12 health solutions to support early detection and management of PE/E in LMICs. For this scoping review,  
13 the DHIs included wearable devices, predictive models operationalized through clinical applications,  
14 health information technologies, health management systems, and other innovations related to mobile  
15 health, telehealth, and telemedicine that can guide diagnosis, monitoring, and treatment[14] . The  
16 review included only English-language studies, which were conducted in LMICs. The World Bank's  
17 (WB) 2020 country classification list was used to select LMICs with a Gross National Income (GNI)  
18 per capita between \$1,036 and \$4,045 [15]. The review primarily aimed to include original and primary  
19 research studies, including experimental studies (e.g., randomized controlled trials, quasi-experimental  
20 studies), observational studies (e.g., cohort, case-control, cross-sectional, qualitative studies), and study  
21 protocols. All types of reviews, meta-analyses, letters to editors, commentaries, viewpoints, news  
22 articles, abstracts, and books were excluded. Articles published between January 1, 2000, and October  
23 20, 2020, were included, given that DHIs prior to 2000 would likely have little applicability for current  
24 implementation (Supplementary file 1: Eligibility Criteria).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

### 41 ***Information Sources and Search Strategy***

42  
43 Five main electronic databases were searched including Excerpta Medica Database (EMBASE),  
44 Medical Literature Analysis and Retrieval System Online (MEDLINE), Cochrane Central Register of  
45 Controlled Trials, Cochrane Database of Systematic Reviews, and Cumulated Index to Nursing and  
46 Allied Health Literature (CINAHL). A supplementary search was conducted using the first seven pages  
47 of Google Scholar to capture peer-reviewed literature on the use of DHIs to support pregnant women  
48 at HRPE. The reference lists of relevant systematic reviews and final included articles were also hand-  
49 searched to find pertinent studies. The search strategy was developed with the assistance of an expert  
50 librarian specializing in health services research. It included four main concepts of interest: target  
51 population (pregnant women), health condition (PE), intervention (digital health tools), and settings  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (LMICs). The search strategy included both keywords and subject headings such as MeSH, and Emtree  
4  
5 (Supplementary file 2: Search strategy).  
6  
7  
8

### 9 ***Selection Procedure***

10  
11 Records from all the electronic databases were exported to Endnote software for screening purposes.  
12  
13 The primary reviewer (AS) developed a pre-defined screening form, and pilot testing was carried out  
14  
15 using 10 randomly selected articles to ensure appropriate screening reliability among the two reviewers  
16  
17 (AS and NA), which was found to be 90%. All articles were independently screened by the two  
18  
19 reviewers to exclude those that did not fulfill the inclusion criteria. Two reviewers then met to review  
20  
21 any discrepancies which were discussed until a consensus was reached.  
22  
23

24  
25 The initial search found a total of 4,078 articles. After de-duplication, 3,389 titles and abstracts were  
26  
27 screened by the two reviewers (AS and NA) to evaluate whether they met the eligibility criteria. Of  
28  
29 these, 72 records were found to be eligible for full-text screening by the two reviewers. Finally, 19  
30  
31 articles were identified after the full-text screening that met the inclusion criteria for this review [16-  
32  
33 34]. Fifty-three articles were excluded for the following reasons: (1) the study was not reported in the  
34  
35 English language; (2) the publication did not talk about pregnant women at HRPE; (3) the research did  
36  
37 not include any of the DHIs; (4) the publication was a conference abstract, review, editorial,  
38  
39 commentary; or (5) the study implemented the DHIs for pregnant women at HRPE in high-income  
40  
41 countries. The study selection procedure was recorded according to the PRISMA-ScR flow diagram  
42  
43 (Figure 1: PRISMA-ScR Flow Diagram for Database Search of Studies).  
44

### 45 ***Data Extraction***

46  
47 A data abstraction form was designed collectively by the research team to determine appropriate  
48  
49 variables such as study characteristics, type of DHIs, intervention description, and study outcomes  
50  
51 (Supplementary file 3: Data Abstraction form). To ensure consistency in the data extraction process,  
52  
53 the form was pilot tested using three randomly selected articles, which resulted in consistent data being  
54  
55 abstracted by both reviewers. Both reviewers (AS, NA) independently completed the data extraction  
56  
57 sheet for each of the 19 final articles. The data abstraction sheets of both the reviewers were compared  
58  
59 to confirm that all major results were included in the scoping review. In the case of inconsistencies  
60

1  
2  
3 between the data extraction sheets from the two reviewers, a third reviewer would have been invited to  
4  
5 make a final decision, but no inconsistencies were found.

### 6 7 ***Data Analysis***

8  
9 An inductive approach was used to thematically organize and summarize the results from the included  
10  
11 articles to explore our research question. The extracted results from each article were read several times  
12  
13 to identify frequent patterns, similarities, and differences on the use of DHIs to support pregnant women  
14  
15 at HRPE in LMICs. The identified emerging patterns were organized into five thematic groupings  
16  
17 including study characteristics, overview and appraisal of included studies, purpose of DHIs, users of  
18  
19 DHIs, and types of outcomes examined by the included studies. The first, and last author discussed the  
20  
21 results and agreed upon the final groupings of the results.  
22  
23

### 24 25 ***Patient and Public Involvement***

26  
27 No patients or members of the public were involved in the protocol design and conduct of the scoping  
28  
29 review.

### 30 31 **Results**

#### 32 33 ***Study Characteristics***

34  
35 A total of 19 publications describing 7 unique studies were included in this review. The included articles  
36  
37 were published between 2009 and 2020. Of these 19 articles, a total of 16 articles described studies that  
38  
39 were conducted in South Asia and Sub-Saharan Africa, one article described a study conducted in  
40  
41 Africa, Southern Asia, and the Middle East, and the remaining two articles described studies conducted  
42  
43 in unspecified resource-poor settings (LMICs) (Supplementary file 4: Overview of the included  
44  
45 articles).

46  
47 The 19 articles were classified into three types of articles: observational studies (n=12), experimental  
48  
49 studies (n=4 including two RCTs), and protocol papers (n= 3). All included articles reported the use of  
50  
51 DHIs for antepartum women. The articles reported varying eligibility criteria for selecting high-risk  
52  
53 pregnant women for different DHIs. Some articles selected high-risk pregnant women based on the  
54  
55 National Institute for Health and Care Excellence (NICE) guidelines [35], specific age groups such as  
56  
57 pregnant women aged 15–49 years [22], while a few articles selected pregnant women based on their  
58  
59 residential area such as women living in study catchment area [23], permanent resident of the particular  
60

1  
2  
3 area, or non-resident who delivered in the study area[18]. . Most DHIs collected blood pressure, heart  
4 rate, and pulse oximetry, with some innovations collecting data on additional indicators such as  
5 demographic data, hemoglobin, urine dipstick test to detect proteinuria and glucose, other urinary  
6 markers, and PE symptoms. Only one article reported the use of international guideline (NICE clinical  
7 guideline 107) to determine blood pressure thresholds [28] (Supplementary file 5: DHIs  
8 characteristics).

9  
10  
11  
12  
13  
14  
15  
16 Seven articles described the application of theoretical frameworks to guide the implementation and  
17 evaluation of digital health tools, including the technology acceptance model [25], diffusion of  
18 innovation model [26, 31], three delay model [26, 29], normalization process theory [23], medical  
19 research council framework [34], logic models [31, 34], realist evaluation theories [31], and cost-  
20 effectiveness models [22]. Two articles described the use of the LambdaNative framework for the  
21 development of the ‘PIERS on the Move’ (POTM) mHealth application [19, 24]. The remaining 10  
22 articles did not mention the use of theory or frameworks for the implementation of DHIs.

### 23 24 25 26 27 28 29 30 31 ***Overview of the Appraisal of Included Studies***

32  
33 A total of ten publications in this review reported research work of the monitoring component of PRE-  
34 EMPT (PE/E Monitoring, Prevention & Treatment) project by Peter von Dadelszen et al., University  
35 of British Columbia [17-19, 22-24, 29-31, 36]. The elements of the monitoring component include  
36 predictive models, Community Level Interventions for PE (CLIP) and integrated mHealth applications.  
37 The PRE-EMPT initiative involved the work of the following research groups: CLIP Pakistan working  
38 group, CLIP India working group, CLIP trial collaborative group, and MiniPIERS and FullPIERS study  
39 working group. The PRE-EMPT project was funded through the Bill & Melinda Gates Foundation  
40 (\$25.9 million).

41  
42  
43  
44  
45  
46  
47  
48  
49 A total of four articles reported research work of CRADLE vital sign alert (VSA) trial led by Nathan et  
50 al., which aimed to evaluate the ability of the device to accurately detect abnormalities in women’s vital  
51 signs during pregnancy [27, 28, 34, 37]. The remaining five publications reported five unique DHIs to  
52 support pregnant women at HRPE including the Congo Red Dot test [21], a hypothetical telemonitoring  
53 program [20], a new hypertension detector [32], an integrated diagnostic and clinical decision support  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 system named 'bliss4midwives'[16], and a smart wristwatch (called the F1 smart wristwatch) for blood  
4 pressure monitoring of expectant mother [25].  
5

6  
7 Following PRISMA-ScR guidelines, each of the above-mentioned included article was reviewed to  
8 identify emerging themes related to the use of DHIs to support pregnant women at HRPE in LMICs.  
9

10  
11 The key themes that emerged from the observational and experimental studies and protocol papers are  
12 as follows: (1) purpose of DHIs including risk prediction and monitoring of high-risk pregnant women;  
13  
14 (2) users of DHIs including healthcare providers (HCPs), caregivers, and pregnant women; (3) types of  
15  
16 outcomes examined in included studies including maternal and neonatal health outcomes, usability and  
17  
18 acceptability and intervention feasibility.  
19  
20

### 21 22 ***Purpose of Digital Health Interventions***

23  
24 This review reports nine unique DHIs from 19 included articles to support pregnant women at HRPE/E  
25  
26 in LMICs. These unique interventions are clustered into two main groups based on their purpose:  
27  
28 predicting risk of adverse maternal health outcomes (n=2) and monitoring high-risk pregnant women  
29  
30 to manage PE/E (n=7). Most articles (n=7) described the use of more than one unique DHI (Figure 2:  
31  
32 Classification of the Included Studies Based on the Purpose of Digital Health Interventions.).  
33  
34

### 35 ***Predicting Risk of Adverse Maternal Health Outcomes***

36  
37 Five observational studies and two RCTs described the use of two unique clinical predictive models  
38  
39 named fullPIERS [19] and miniPIERS [17-19, 24, 29-31] to facilitate the prediction of adverse maternal  
40  
41 outcomes occurring as a result of PE based on demographics, symptoms, clinical signs (including  
42  
43 SpO<sub>2</sub>), and laboratory tests. In order to implement these predictive models, the mobile application  
44  
45 'POTM' was developed as an interface to enable healthcare workers to easily determine the risk of  
46  
47 adverse maternal health outcomes. One article reported the use of both the miniPIERS and fullPIERS  
48  
49 predictive models [19], while six articles only reported the use of the miniPIERS model to predict  
50  
51 adverse health outcomes among pregnant women with PE/E in LMICs [17, 18, 24, 29-31]. Payne et al.  
52  
53 described the development process of the miniPIERS model to identify pregnant women at HRPE/E in  
54  
55 five LMICs using simple-to-measure indicators: personal demographics (gestational age); clinical signs  
56  
57 (blood pressure readings and proteinuria); and PE symptoms (headache, visual disturbances, chest pain,  
58  
59  
60

1  
2  
3 dyspnea, vaginal bleeding, and abdominal pain) [29]. The fullPIERS model included additional  
4 predictors such as SpO2 and laboratory tests, to calculate a risk score for pregnant women.  
5  
6

### 7 Monitoring High-Risk Pregnant Women for Managing PE/E conditions

9 The review identified seven unique DHIs for continuous monitoring high-risk pregnant women for  
10 managing PE/E including one diagnostic test named Congo Red Dot for monitoring misfolded protein  
11 in the preeclamptic urine [21], CLIP intervention for monitoring blood pressure among high-risk  
12 women through community health workers [17, 18, 22, 23, 30, 31], as well as five unique devices for  
13 monitoring blood pressure [16, 20, 25, 27, 28, 32, 34, 37]. The five unique devices for measuring blood  
14 pressure among high-risk pregnant women include the Microlife CRADLE VSA device [27, 28, 34,  
15 37], the Bliss4Midwives' device [16], a new hypertension detector device [32], hypothetical  
16 telemonitoring program [20] and the F1 smart wristwatch [25].  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 The Congo Red Dot test was evaluated in a prospective experimental study design. The Congo Red  
27 Dot test requires minimal specialized equipment and enables minimally trained personnel to diagnose  
28 PE in resource-limited health care settings. The test was developed in 2016, based on the ability of  
29 constituents in preeclamptic urine to bind the amyloidophilic dye Congo Red. At the core of the test is  
30 the discovery that preeclamptic women eliminate misfolded proteins in their urine, a molecular feature  
31 that is proportional to disease severity [21].  
32  
33  
34  
35  
36  
37  
38

39 The CLIP intervention was implemented in Mozambique, Pakistan, India, and Nigeria as part of cluster  
40 randomized controlled trials (cRCTs) [17, 18, 22, 23, 30, 31]. The implementation of CLIP intervention  
41 involved scaling-up of existing community health workforce to provide community engagement and  
42 community health worker-led app-guided monitoring for high-risk pregnant women for hypertension.  
43 Community health workers were able to undertake all aspects of the app-guided visits, and  
44 approximately 10% of pregnant women were found to be hypertensive.  
45  
46  
47  
48  
49  
50

51 As a first example of blood pressure measurement device, Nathan et al. assessed the accuracy of the  
52 Microlife 3AS1-2 blood pressure device in 2014 for use in pregnancy and PE in a low-resource setting  
53 [27]. The study recruited a total of 45 pregnant women, of whom 15 had PE, from Kimberley Hospital  
54 in South Africa. The study concluded that the device can be recommended for use in pregnancy,  
55 including PE as it fulfills the requirements stipulated by the WHO for an automated blood pressure  
56  
57  
58  
59  
60

1  
2  
3 device suitable for use in antenatal clinics and primary healthcare facilities of LMICs. The device has  
4 been extensively validated for accuracy, usability, and acceptability in low-resource settings [27]. The  
5 device calculates the pregnant woman's risk of hypovolaemic or septic shock and alerts frontline  
6 healthcare workers about vital sign abnormalities through a traffic light early warning system display.  
7  
8 In 2018, a three-month mixed-methodology feasibility study was conducted to incorporate the  
9 CRADLE VSA device into routine maternity care in 10 low-income sites [34]. Primary, secondary, and  
10 tertiary facilities were allocated devices and training packages consisting of a short-animated film,  
11 interactive sessions, booklet, and posters.  
12  
13

14  
15 As a second example, a study conducted in Ghana used the Bliss4Midwives (B4M) device which  
16 included infrared sensors to measure hemoglobin, a self-inflating cuff for blood pressure measurement,  
17 and an automated reader for urinary protein and glucose through dipsticks. The device facilitated non-  
18 invasive screening of PE and served as an integrated diagnostic and clinical decision support device for  
19 PE [16]. The third example of a device for blood pressure monitoring was a new hypertension detector,  
20 developed by Thakor et al., which was compared in an observational study with other traditional devices  
21 for use in developing countries to support pregnant women at HRPE/E[32]. The new device was found  
22 to be more accurate and easy-to-use than CRADLE VSA and other devices, due to the reduced number  
23 of steps required for use [32]. As a fourth example of a device for blood pressure monitoring was a  
24 hypothetical telemonitoring program [20], which was described in a qualitative study protocol. The  
25 study intended to explore the perspectives, needs, and preferences of a telemonitoring program for  
26 pregnant women at HRPE in a tertiary health facility of Karachi, to inform future implementation.  
27  
28

29  
30 Finally, one prospective experimental study used a wearable device called the F1 smart wristwatch that  
31 included an integrated chip for sensing blood pressure readings and displaying real-time data on the  
32 screen. The smartwatch on the expectant mother's wrist takes blood pressure readings and transfers  
33 them by Bluetooth to their phone at regular intervals to facilitate personal health tracking. The caregiver  
34 can access the expectant mother's records, as well as receive alerts on blood pressure readings [25].  
35  
36

37  
38 Both of these purposes utilized mobile phone applications as an interface to facilitate data collection,  
39 decision making, and communication between health workers and pregnant women. The majority of  
40 these studies used the POTM application [17-24, 30, 31, 33] to facilitate the collection of relevant  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 clinical data during antenatal visits. The application was used by community health workers in India,  
4 Pakistan, Nigeria, and Mozambique, as part of a CLIP cluster RCT [17, 30]. The POTM platform  
5 combined two interventions, which were the miniPIERS model and a Phone Oximeter to accurately  
6 predict the risk score for pregnant women at HRPE/E in LMICs. The application generated a risk  
7 estimate which enabled community health workers and other healthcare providers (HCPs) to stratify  
8 high-risk pregnant women, escalate care, and make referrals to the facility. In addition, Jonas et al. study  
9 used a mobile application for administrating CRD test for monitoring misfolded protein in the  
10 preeclamptic urine [21]. Finally, the Feroz et al. study protocol described a hypothetical mobile-based  
11 telemonitoring program which would serve as a communication aid between nurses and high-risk  
12 pregnant women[20, 38].

### 23 ***Users of Digital Health Interventions***

24 Most articles involved HCPs (n=17) as the targeted primary users of the DHIs, while only two articles  
25 had pregnant women and caregivers as the primary users of the DHI [20, 25]. The articles described  
26 various healthcare workers as the users of the DHIs, including mid-level HCPs, community-based  
27 HCPs, lady health supervisors, semi-literate volunteers, community health nurses, lady health workers,  
28 midwives, and accredited social health activists. Sixteen articles included information on the training  
29 of patients and HCPs on how to use the DHI, interpret physiological metrics, and take actionable  
30 measures for critical results [16, 18, 19, 21-24, 26-34]. The HCPs received advanced training to  
31 enhance their assessment skills and ability to facilitate the overall management of pregnant women at  
32 HRPE/E. Three articles did not specify the training component for either HCPs or patients [17, 20, 25].

### 33 ***Type of Outcomes Examined***

34 The included articles (n=19) reported on three major outcomes: 1) maternal and neonatal health  
35 outcomes (n=4), 2) usability and acceptability (n=5), and 3) intervention feasibility (n=7)  
36 (Supplementary file 6: Outcomes of DHIs).

### 37 ***Maternal and Neonatal Health Outcomes***

38 Four articles examining maternal and neonatal health outcomes were observational studies (n=2) and  
39 RCTs (n=2) [17, 18, 28, 30]. Maternal health outcomes included magnesium sulfate use, hospital  
40 admissions, CCU admissions, birth preparedness, complication readiness, facility delivery attended by  
41

1  
2  
3 skilled birth attendants, and adverse maternal outcomes such as an increase in kidney injury, maternal  
4 morbidity, and mortality. For example, Nathan et al.'s observational study evaluated the association  
5 between blood pressure measurements and adverse outcomes in women with PE using CRADLE VSA  
6 traffic light early warning system. The study demonstrated that the risk of maternal death, eclampsia,  
7 and perinatal death was similar across the women who triggered a yellow or red light on the CRADLE  
8 VSA. However, the risk of kidney injury, maternal use of magnesium sulfate, maternal CCU admission  
9 and preterm delivery, was greater for those who triggered a red light, compared to a yellow light. [28].  
10  
11 The two RCTs reported non-significant findings regarding maternal morbidity and mortality for  
12 participants in the DHI arm [17, 30]. Neonatal health outcomes included stillbirths, fetal and neonatal  
13 morbidity, and mortality. Only one of the two RCTs reported a reduction in stillbirths (0·89 [0·81-  
14 0·99];  $p = 0·03$ ) in the DHI group; however, no impact on neonatal morbidity or mortality was reported  
15 for participants in the DHI group [30].

### 16 Usability and Acceptability

17  
18 Five articles reported on the usability and acceptability of DHIs in LMICs [19, 24-26, 32]. The articles  
19 mentioned pregnant women, caregivers, and HCPs' experience of use of DHIs in LMICs. Usability  
20 outcomes included: trust in technology, ease of use, content richness, perceived usefulness, and user  
21 satisfaction. For instance, Musyoka et al.'s (2019) study found that a 24-hour ambulatory blood pressure  
22 monitoring system has a great potential for actual adoption in healthcare systems in developing  
23 countries, given its simplicity and affordability [25]. The study found that content richness had a slightly  
24 positive linear effect on perceived ease of use, while there is a slightly negative relationship between  
25 content richness and perceived usefulness [25]. Lim et al. used the computer systems usability  
26 questionnaire to assess the usability of the POTM mHealth application [24]. Nurses and midwives who  
27 participated in the study rated the usability high for the integration of these technologies and thought it  
28 would help their fieldwork. The study found that usability issues were often related to navigation of the  
29 app and phone features such as scroll wheels, touch screen use, etc. In a study by Nathan et al., most  
30 HCWs perceived the CRADLE device to be easy to use; however, some described manual inflation as  
31 tiring, particularly when measuring vital signs in obese and hypertensive women [26]. Dunsmuir et al.'s  
32 study reported on the usability of CLIP POTM application; the CLIP trial received requests from  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 different countries for modifications in POTM to consider different user needs and cultural differences  
4 leading to modified application versions for each country [19]. *Intervention Feasibility and Fidelity*  
5  
6 Most articles (n=7) reported on the feasibility and fidelity of DHIs for pregnant women at HRPE/E in  
7  
8 LMICs in order to provide evidence on the evaluation of DHIs for replication and scale-up of  
9  
10 successful DHIs [16, 21, 23, 27, 29, 31, 34]. Study outcomes included: fidelity and accuracy of the  
11  
12 CRADLE VSA device, MiniPIERS model development and validation, understanding of enabling and  
13  
14 impeding factors for CLIP trial implementation, experiences of pregnant women with B4M  
15  
16 intervention, and cost-effectiveness of the Congo Red Dot test. For example, Payne et al.'s study  
17  
18 informed that miniPIERS model has a reasonable ability to identify women at increased risk of adverse  
19  
20 maternal outcomes associated with the hypertensive disorders of pregnancy [29]. Nathan et al.'s another  
21  
22 study assessed the accuracy of Microlife 3AS1-2 blood pressure device for accuracy for use in  
23  
24 pregnancy in LMICs. The authors concluded that the device can be recommended for use in pregnancy,  
25  
26 including PE as it meets the standards stipulated by the WHO for automated blood pressure devices  
27  
28 suitable for low-resource settings [27]. One mixed-methods study reported high fidelity of the  
29  
30 implementation of the CRADLE VSA device, with improved HCPs ability to make clinical decisions,  
31  
32 escalate care, and make immediate referrals in case of emergency [34]. The study by Khowaja et al.  
33  
34 (2016) reported factors associated with the feasibility of the CLIP trial implementation including  
35  
36 community mobilization, institutional support, system integration, knowledge gaps, lack of trained  
37  
38 personnel, cultural myths and misconceptions, poor health service quality, and high cost of care [23].  
39  
40  
41  
42  
43  
44

## 45 Discussion

### 46 *Principal Findings*

47  
48 This review summarizes evidence on the existing DHIs to support pregnant women at HRPE/E in  
49  
50 LMICs. Given that most articles (11 out of 19) were published between 2015 and 2020, the novelty of  
51  
52 DHIs use to support pregnant women with HRPE/E was indicated. Only nine unique DHIs were  
53  
54 identified in this review from 19 included articles, reflecting the limited understanding and use of DHIs  
55  
56 to support pregnant women in LMICs. Most included articles used observational and exploratory  
57  
58 research methods to study DHIs. This suggested the need for concerted efforts to learn from small  
59  
60



1  
2  
3 innovation projects and deployments as outlined in WHO guide on monitoring and evaluation of DHIs  
4 [39]. Most articles in this review did not report information on the blood pressure thresholds, which  
5 limited our understanding of standardized blood pressure thresholds used in LMICs. The explicit  
6 reporting of standardized blood pressure thresholds could help in designing effective clinical decision  
7 support systems for monitoring pregnant women in LMICs [40].  
8  
9  
10  
11  
12

### 13 ***Implementation Barriers and Strategies for DHIs***

14  
15 The Microlife CRADLE VSA blood pressure monitoring device has been extensively validated for use  
16 in LMICs for pregnant women [27, 28, 34, 37]. However, HCPs faced several barriers during the  
17 implementation of CRADLE VSA device including lack of supportive supervision for device use, high  
18 staff turnover, and poor availability of the device, poor battery life of device, misleading displays,  
19 broken hand pump, tubing and broken charging ports [34]. Nathan et al. and Vousden et al. suggested  
20 a range of implementation strategies to address known barriers, prior to scale-up, including recognizing  
21 designated device champions who can provide in-depth local training and support for device use,  
22 emphasizing the importance of a device training package (short animated film, interactive sessions,  
23 booklet, and posters), updating training materials to explain the traffic light alert system, providing  
24 chargers in addition to the USB cable, and ensuring an adequate supply of VSA devices [28, 34]. Lim  
25 et al. study mentioned that the general unfamiliarness of using touch screen smart phones was reported  
26 as the major barrier faced during the implementation of POTM application[24]. Abejirinde et al. study  
27 trained users on the technical and operational functions of the device to address technical and procedural  
28 issues including software freezes, slow response time, and low user dexterity with operating the device  
29 were two main factors that contributed to delays[16].  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

### 47 ***Research Gaps and Suggestions for Future Research***

48  
49 *Enabling the use of DHIs by pregnant women as end-users instead of HCPs as end-users:* Most articles  
50 in this review targeted DHIs at HCPs who have less formal training and education, as opposed to studies  
51 conducted in high-income countries where DHIs have been targeted at family physicians and clinicians  
52 who have specialized medical training [6]. This review identified only one study that targeted DHI at  
53 pregnant women for personal health tracking [25]; however, DHIs implemented in high-income  
54 countries are often targeted for use by pregnant women to improve maternal health behaviors and  
55  
56  
57  
58  
59  
60



1  
2  
3 maternal-fetal health outcomes [41]. Given the increasing cell phone penetration in LMICs [42], there  
4 is an opportunity to use mobile phone technology to target DHIs at the patient level (pregnant women)  
5 to encourage personal health tracking. Yet, health informatics researchers should consider issues of  
6 technological literacy, user characteristics (age, gender, computer skills, experience), cultural factors,  
7 and socioeconomic status when designing and implementing DHIs in the LMIC context [43]. None of  
8 the studies delivered targeted client instructions via a digital platform, in response to abnormal blood  
9 pressure readings or signs and symptoms of PE. In high-income countries, some digital health platforms  
10 have delivered manual or automated targeted instructions to the pregnant women to provide information  
11 about medications, referrals, and diet [44]. LMICs can learn from the experiences of high-income  
12 countries for developing context-specific digital platforms that can facilitate targeted client  
13 communication between providers and pregnant women. Evidence suggests that the targeted client  
14 communication for transmission of health information, health event alerts and reminders, and diagnostic  
15 results have shown positive impacts on health behaviors and health outcomes in high-income countries  
16 [45].

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33 Using Multidisciplinary Team Approach for Designing DHIs: None of the DHIs used a  
34 multidisciplinary team approach for monitoring of pregnant women for PE/E. Blandford et al. suggest  
35 that DHIs should involve collaboration between different cadres of HCPs across all levels of the health  
36 system, to achieve the full potential of digital intervention [46]. For instance, a nurse or midwife at a  
37 primary level could communicate about a pregnant women's health condition to a clinician at a  
38 secondary institution to seek recommendations for managing pregnant women at HRPE/E. Murray et  
39 al. suggest that high-quality research in the digital health field requires fertile multidisciplinary  
40 collaborations that draw on insights and experience from multiple fields, including clinical medicine,  
41 health services research, behavioral science, education, engineering, and computer science[47]. Thus,  
42 research aimed at designing and evaluating DHIs to support pregnant women at HRPE/E should draw  
43 insights from collaborators belonging to diverse disciplines including obstetricians and gynecologists,  
44 telemedicine experts, knowledge users, HCPs (nurses, doctors), public health specialists, maternal  
45 health specialist, health services researchers, as well as patient partners.

1  
2  
3 *Exploring Telemedicine Use to Enable Remote Consultation Between Pregnant Women and Healthcare*

4  
5 *Providers:* Most articles used DHIs for the prediction of adverse maternal outcomes, data collection  
6 and decision aid, diagnostic and clinical decision support, and personal health tracking. There is a lack  
7 of evidence on using DHIs for referral coordination, teleconsultation between pregnant women and  
8 HCPs, communication between the HCP and their supervisor, and HCPs' training. Telemedicine has  
9 been extensively used in high-income countries for providing a range of obstetrical services such as  
10 using videoconference to replace in-person visits, implementing at-home monitoring, enabling  
11 consultation with remote specialists, earlier postpartum follow up visits, and access to lactation  
12 consultants [48]. This evidence shows the potential of using telemedicine for pregnant women at  
13 HRPE/E in LMICs to enable remote monitoring and remote consultation between pregnant women and  
14 providers.  
15

16  
17 *Monitoring and Evaluating the Implementation and Effectiveness of DHIs:* Most articles reported on  
18 intervention feasibility, usability, and acceptability outcomes. Two RCTs reported non-significant  
19 findings for maternal morbidity, mortality, and neonatal deaths [17, 30] with only one RCT that reported  
20 a significant difference in stillbirth rate in DHIs group [30]. This suggests the need of conducting more  
21 experimental studies such as RCTs to evaluate the efficacy and effectiveness of diverse DHIs to improve  
22 maternal and child health outcomes. In the review, only one study protocol described the methodology  
23 to conduct an economic evaluation of the CLIP package in South Asian and African countries [22].  
24 This shows the paucity of evidence on the economic impact of DHIs to support pregnant women with  
25 PE/E. Ramsey et al. recommend that future clinical trials should incorporate cost-analysis of DHIs as  
26 there is mounting evidence on embedding economic evaluations within clinical trials to build a robust  
27 cost-effectiveness model that has high internal validity and timeliness [49]. The articles included in this  
28 review did not extensively identify facilitators and challenges encountered during the implementation  
29 of DHIs for pregnant women with PE/E in LMICs, unlike many studies conducted in high-income  
30 countries [6]. This review identified only a few facilitators: easy to use technology, trust in technology,  
31 and availability of diagnostic service at the point of care. This indicates the need to examine and report  
32 on enablers and barriers faced when employing DHIs for pregnant women at HRPE/E across the stages  
33 of design, development, implementation, and evaluation.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In summary, this scoping review suggests four recommendations for future research: 1) enable the use  
4 of DHIs by pregnant women as end-users to encourage personal health tracking including  
5 individualized patient instructions; 2) consider a multidisciplinary team approach when designing DHIs  
6 for pregnant women at HRPE/E; 3) explore the potential of using telemedicine in LMICs to enable  
7 remote consultation between pregnant women and health provider; 4) conduct further studies including  
8 prospective longitudinal and experimental studies to establish the implementation effectiveness and  
9 efficacy of DHIs to support pregnant women at HRPE; exploratory studies to identify barriers and  
10 enablers associated with the development, implementation, and evaluation of DHIs; and economic  
11 evaluations of DHIs within large clinical trials to identify cost-effective DHIs.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

## 22 **Conclusion**

23  
24 The current evidence base is sparse but shows some potential for the use of different DHIs to support  
25 pregnant women in early diagnosis of PE/E through predicting the risk for adverse maternal health  
26 outcomes and monitoring high-risk pregnant women for PE/E through devices and other DHIs . Limited  
27 evidence exists on types, benefits, cost-effectiveness, and outcomes of DHIs. The weak evidence may  
28 impede the adoption of these promising technologies in community and healthcare settings to support  
29 pregnant women at HRPE/E in LMICs. Future research work should target DHIs at the pregnant women  
30 level to promote personal health tracking with targeted instructions for pregnant women, consider a  
31 multidisciplinary team approach for designing DHIs, explore the role of telemedicine to enable remote  
32 consultation between pregnant women and healthcare providers, and evaluate the implementation and  
33 effectiveness of DHIs.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

## 45 **Acknowledgements**

46 None

## 47 **Conflicts of Interest**

48 Authors declare no competing interests.

## 49 **Contributors**

50 ASF and ES authors conceptualized and designed the study. ASF screened the articles and performed  
51 data extraction, synthesized the data, and drafted the manuscript. NA independently performed  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 screening of articles. ES served as the senior author including participating in the data analysis and  
4 providing critical feedback on the manuscript. All authors read and approved the final manuscript.  
5  
6

### 7 **Funding**

8  
9 This research received no specific grant from any funding agency in the public, commercial or not-for-  
10 profit sectors  
11  
12

13 **Data availability statement** No additional data available  
14

### 15 **Patient consent for publication**

16  
17 Not required.  
18  
19

### 20 **Ethics Statement**

21  
22 Not required  
23  
24

### 25 **Abbreviations**

26 CLIP: Community-Level Interventions for Preeclampsia  
27

28 CRD: Congo Red Dot  
29

30 DHI: Digital Health Intervention  
31

32 PRE-EMPT: Preeclampsia, Eclampsia Monitoring, Prevention & Treatment  
33

34 GNI: Gross National Income  
35

36 HCP: Healthcare Provider  
37

38 HRPE: High Risk for Preeclampsia  
39

40 LMIC: Low Middle Income Country  
41

42 NICE: National Institute for Health and Care Excellence  
43

44 OSF: Open Science Framework  
45

46 PE/E: Preeclampsia/Eclampsia  
47

48 PIERS: Preeclampsia Integrated Estimate of RiSk  
49

50 POTM: PIERS on the Move  
51

52 PRISMA-ScR: Preferred Reporting Items for Systematic Reviews and Meta-analyses-Scoping Review  
53

54 RCT: Randomized Controlled Trials  
55

56 TM: Telemonitoring  
57

58 VSA: Vital Sign Alert  
59

60 WHO: World Health Organization

## References

1. Goldenberg, R.L., et al., *Reducing maternal mortality from preeclampsia and eclampsia in low-resource countries—what should work?* 2015. **94**(2): p. 148-155.
2. Khowaja, A.R., et al., *Community's perceptions of pre-eclampsia and eclampsia in Sindh Pakistan: a qualitative study.* 2016. **13**(1): p. 39-44.
3. Howden-Chapman, P., et al., *SDG 3: Ensure healthy lives and promote wellbeing for all at all ages.* 2017: p. 81-126.
4. Van Den Heuvel, J., et al., *Home-based telemonitoring versus hospital admission in high risk pregnancies: a qualitative study on women's experiences.* 2020. **20**(1): p. 1-9.
5. van den Heuvel, J.F., et al., *SAFE@ HOME—Feasibility study of a telemonitoring platform combining blood pressure and preeclampsia symptoms in pregnancy care.* 2019. **240**: p. 226-231.
6. Aquino, M., et al., *Exploring the Use of Telemonitoring for Patients at High Risk for Hypertensive Disorders of Pregnancy in the Antepartum and Postpartum Periods: Scoping Review.* 2020. **8**(4): p. e15095.
7. WHO, *WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience.* Geneva, Switzerland. 2016.
8. Lanssens, D., et al., *Effectiveness of telemonitoring in obstetrics: scoping review.* 2017. **19**(9): p. e327.
9. Rivera-Romero, O., et al., *Mobile health solutions for hypertensive disorders in pregnancy: scoping literature review.* 2018. **6**(5): p. e130.
10. van den Heuvel, J.F., et al., *eHealth as the Next-Generation Perinatal Care: An Overview of the Literature.* J Med Internet Res, 2018. **20**(6): p. e202.
11. de Oliveira, B.G.R.B., *Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist.* 2015.
12. Levac, D., H. Colquhoun, and K.K.J.I.s. O'Brien, *Scoping studies: advancing the methodology.* 2010. **5**(1): p. 69.
13. O'Malley, L.J.I.J.S.R.M., *Scoping studies: towards a methodological framework AU-Arksey, Hilary.* 2005. **8**: p. 19-32.
14. Guo, C., et al., *Challenges for the evaluation of digital health solutions—A call for innovative evidence generation approaches.* npj Digital Medicine, 2020. **3**(1): p. 1-14.
15. Serajuddin, U. and N. Hamadeh, *New World Bank country classifications by income level: 2020–2021.* World Bank Blogs, 2020.
16. Abejirinde, I.O., et al., *Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana.* BMC Pregnancy Childbirth, 2018. **18**(1): p. 209.
17. Bellad, M.B., et al., *Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial.* Pregnancy Hypertens, 2020. **21**: p. 166-175.
18. Bellad, M.B., et al., *Maternal and newborn health in Karnataka state, India: the community level interventions for pre-eclampsia (CLIP) Trial's baseline study results.* 2017. **12**(1): p. e0166623.
19. Dunsmuir, D.T., et al., *Development of mHealth applications for pre-eclampsia triage.* IEEE J Biomed Health Inform, 2014. **18**(6): p. 1857-64.
20. Feroz, A., S. Saleem, and E. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol.* Reprod Health, 2020. **17**(1): p. 135.
21. Jonas, S.M., et al., *Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources.* J Am Med Inform Assoc, 2016. **23**(1): p. 166-73.

- 1
- 2
- 3
- 4 22. Khowaja, A.R., et al., *Economic evaluation of Community Level Interventions for Pre-eclampsia (CLIP) in South Asian and African countries: a study protocol*. Implement Sci, 2015. **10**: p. 76.
- 5
- 6
- 7 23. Khowaja, A.R., et al., *The feasibility of community level interventions for pre-eclampsia in South Asia and Sub-Saharan Africa: a mixed-methods design*. Reprod Health, 2016. **13 Suppl 1**: p. 56.
- 8
- 9
- 10 24. Lim, J., et al., *Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage*. JMIR Mhealth Uhealth, 2015. **3**(2): p. e37.
- 11
- 12 25. Musyoka, F.M., M.M. Thiga, and G.M. Muketha, *A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care*. Informatics in Medicine Unlocked, 2019. **16**.
- 13
- 14 26. Nathan, H.L., et al., *The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings*. Reprod Health, 2018. **15**(1): p. 5.
- 15
- 16 27. Nathan, H.L., et al., *An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2*. Blood Press Monit, 2015. **20**(1): p. 52-5.
- 17
- 18 28. Nathan, H.L., et al., *Early warning system hypertension thresholds to predict adverse outcomes in pre-eclampsia: A prospective cohort study*. Pregnancy Hypertens, 2018. **12**: p. 183-188.
- 19
- 20 29. Payne, B.A., et al., *A risk prediction model for the assessment and triage of women with hypertensive disorders of pregnancy in low-resourced settings: the miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) multi-country prospective cohort study*. PLoS Med, 2014. **11**(1): p. e1001589.
- 21
- 22 30. Qureshi, R.N., et al., *Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial*. Pregnancy Hypertens, 2020. **22**: p. 109-118.
- 23
- 24 31. Sharma, S., et al., *A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria*. 2017. **17**(1): p. 238.
- 25
- 26 32. Thakor, N., et al., *Hypertension Detector for Developing Countries*. 2010, Department of Biomedical Engineering, Johns Hopkins University.
- 27
- 28 33. von Dadelszen, P., et al., *The PRECISE (PREgnancy Care Integrating translational Science, Everywhere) Network's first protocol: deep phenotyping in three sub-Saharan African countries*. Reprod Health, 2020. **17**(Suppl 1): p. 51.
- 29
- 30 34. Vousden, N., et al., *Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial*. BMC Pregnancy Childbirth, 2018. **18**(1): p. 115.
- 31
- 32 35. Feroz, A., S. Saleem, and E.J.R.H. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol*. 2020. **17**(1): p. 1-7.
- 33
- 34 36. von Dadelszen, P., et al., *The Community-Level Interventions for Pre-eclampsia (CLIP) cluster randomised trials in Mozambique, Pakistan, and India: an individual participant-level meta-analysis*. The Lancet, 2020. **396**(10250): p. 553-563.
- 35
- 36 37. Nathan, H.L., et al., *Evaluation of a novel device for the management of high blood pressure and shock in pregnancy in low-resource settings: study protocol for a stepped-wedge cluster-randomised controlled trial (CRADLE-3 trial)*. 2018. **19**(1): p. 1-10.
- 37
- 38 38. Feroz, A., S. Saleem, and E. Seto, *Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a tertiary health facility of Karachi: a qualitative study protocol*. Reproductive health, 2020. **17**(1): p. 1-7.
- 39
- 40 39. Organization, W.H., *Monitoring and evaluating digital health interventions: a practical guide to conducting research and assessment*. 2016.
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60



- 1
- 2
- 3
- 4 40. Nee, O. and A.J.D.S.S. Hein, Advances in, Book, editor Ger Devlin, *Clinical decision support*
- 5 *with guidelines and Bayesian networks*. 2010: p. 117-137.
- 6 41. Hussain, T., P. Smith, and L.M. Yee, *Mobile Phone–Based Behavioral Interventions in*
- 7 *Pregnancy to Promote Maternal and Fetal Health in High-Income Countries: Systematic*
- 8 *Review*. JMIR mHealth and uHealth, 2020. **8**(5): p. e15111.
- 9 42. Muir, S., *To what extent is mHealth an effective Intervention in LMICs?* 2019.
- 10 43. Feroz, A., M.M. Kadir, and S.J.G.h.a. Saleem, *Health systems readiness for adopting mhealth*
- 11 *interventions for addressing non-communicable diseases in low-and middle-income*
- 12 *countries: a current debate*. 2018. **11**(1): p. 1496887.
- 13 44. Haddad, S.M., R.T. Souza, and J.G. Cecatti, *Mobile technology in health (mHealth) and*
- 14 *antenatal care—searching for apps and available solutions: a systematic review*. International
- 15 *journal of medical informatics*, 2019. **127**: p. 1-8.
- 16 45. Organization, W.H., *WHO guideline: recommendations on digital interventions for health*
- 17 *system strengthening: web supplement 2: summary of findings and GRADE tables*. 2019,
- 18 *World Health Organization*.
- 19 46. Blandford, A., et al., *Seven lessons for interdisciplinary research on interactive digital health*
- 20 *interventions*. 2018. **4**: p. 2055207618770325.
- 21 47. Murray, E., et al., *Evaluating digital health interventions: key questions and approaches*.
- 22 2016, Elsevier.
- 23 48. Weigel, G., B. Frederiksen, and U. Ranji, *Telemedicine and pregnancy care*. The Henry J Kaiser
- 24 *Family Foundation*. February, 2020.
- 25 49. Ramsey, S., et al., *Good research practices for cost-effectiveness analysis alongside clinical*
- 26 *trials: the ISPOR RCT-CEA Task Force report*. 2005. **8**(5): p. 521-533.
- 27
- 28
- 29
- 30
- 31

## 32 Figure Legends

33 Figure 1: PRISMA-ScR Flow Diagram for Database Search of Studies

34 Figure 2: Classification of the Included Studies Based on the Purpose of Digital Health Interventions.

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56


57

58


59

60





1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47



136/bmjopen-2021-056630 on 8 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

## Supplementary File 1 - Eligibility Criteria

**Question 1: Does this study include humans?**

- a) If yes, **INCLUDE**
- b) **EXCLUDE** animal studies/models, non-humans or vertebrae studies

**Question 2: Is the primary language of the study English?****Is the primary language of the study English?**

- a) If yes, **INCLUDE**
- b) **EXCLUDE** if study is listed as described in a non-English language

**Question 3: Is the article classified as one of the following?**

- a) **INCLUDE** all types of study designs including, observational studies, experimental studies, qualitative studies, study protocols, grey literature.
- b) **EXCLUDE:** systematic reviews, meta-analysis, letter to editors, scoping reviews, commentaries, news articles

**Question 4: Does this study examine care provided to pregnant women with Preeclampsia/eclampsia (PE/E)/or at high risk for PE/E (HRPE/E)?**

- a) If yes, **INCLUDE**
- b) **EXCLUDE** if study does not focus on care provided to pregnant women with PE/E or at HRPE/E

**Question 5: Does this study examine digital technologies to support pregnant women with preeclampsia/eclampsia or at HRPE/E**

- a) **INCLUDE** studies that are focused on use of digital technologies to support pregnant women with PE/E or at HRPE/E. Digital technologies may include:
  - Telephone communication
  - Video communication
  - Text messaging (asynchronous)
  - Email messaging (asynchronous)
  - Portals, apps, and other applications
  - Remote monitoring

- Devices
- Predictive models
- Provider-provider communication through one of the above modalities
  - Synonyms: digital health, virtual care, virtual visits, eVisits, telehealth, telemedicine, eConsultation, mobile health, mHealth, teleconsultation, teleconference, telecommunications, tele\* (e.g., telepsychiatry, teledermatology, etc), videoconferencing, video visits, phone, telephone, electronic consultation, online consultation, e-mail, text messaging, asynchronous messaging, secure messaging, direct messaging, messaging

- b) **INCLUDE** studies focused on using digital technologies for early diagnosis, screening, and management of pregnant women with PE/E or HRPE/E.
- c) **INCLUDE** studies that used digital technologies to support pregnant women with PE/E or at HRPE/E
- d) **EXCLUDE** studies focused on digital health interventions that do not explicitly focus on pregnant women with PE/E or HRPE/E

### Question 7: Is this study based on low-and-middle-income contexts?

See list of countries by income classification here: <https://data.worldbank.org/country/XN>

- a) If yes, **INCLUDE**
- b) **INCLUDE** if study focused on high and middle income together.
- c) **EXCLUDE** if based on only high-income country context

## Supplementary File 2 - Search Strategy

1. Pregnant Women/
2. exp pregnancy/
3. (pregnan\* adj3 ("at risk" or "at-risk" or "high risk" or "high-risk")).tw,kw.
4. exp Eclampsia/
5. exp Pre-Eclampsia/
6. (Pre Eclampsia or preeclampsia or pre-eclampsia or pre eclampsia or eclampsia or gestosis or proteinuria or toxemia\*).tw,kw.
7. or/1-6
8. Telemedicine/
9. Medical informatics/
10. Digital health.mp.
11. mHealth app.mp.
12. predictive model.mp.
13. CLIP.mp.
14. informatics/
15. exp Telecommunications/
16. Monitoring, Ambulatory/
17. exp Telemetry/
18. Monitoring, Physiologic/
19. exp Computer Communication Networks/
20. Mobile Applications/
21. Smartphone/
22. Cell Phone/
23. (tele-monitor\* or telemonitor\* or telemed\* or tele-med\* or teleinterpret\* or tele-interpret\* or telecomm\* or tele-comm\* or telemetry).tw,kw.
24. (mhealth\* or m-health\* or ehealth\* or e-health\* or telehealth\* or tele-health\*).tw,kw.
25. (mobile adj3 (health\* or technolog\* or app\* or solution\* or phone\* or communicat\*)).tw,kw.
26. (remote\* adj3 (transmi\* or transfer\* or tele\* or monitor\* or consult\* or follow-up or program\* or connect\* or web-base\* or "web base\*" or term)).tw,kw.
27. (monitor\* adj3 (home or remote or distan\* or ambulatory or tele\* or online or on-line or "on line" or phone or digital\* or Skype or electronic\* or implant\* or wireless\* or web-base\* or "web base\*")).tw,kw.
28. (interven\* adj3 (remote\* or distan\* or tele\* or online or on-line or "on line" or phone\* or digital\* or Skype or electronic\* or wireless\*)).tw,kw.
29. (smartphone\* or "smart phone\*" or bluetooth\* or Internet\* or phone\* or text messag\*).tw,kw.
30. ((app or apps or application\*) adj3 (mobile or electronic or software)).tw,kw.
31. ((digital\* or electronic\* or online\* or on-line\* or "on line" or Internet) adj3 (health\* or solution\* or transmit\* or transmiss\* or transfer\* or device\* or connect\*)).tw,kw.
32. (broadband adj3 (device\* or capab\*)).tw,kw.
33. (multi-media\* or multimedia\*).tw,kw.
34. (self monitor\* or self-monitor\*).tw,kw.
35. or/8-34
36. 7 and 35
37. developing countries/
38. low-and-middle-income countries.mp.
39. LMICs
40. Honduras/
41. Angola/
42. Papua New Guinea/
43. Algeria/
44. India/
45. Philippines/

- 1
- 2
- 3 46. Bangladesh/
- 4 47. Kenya/
- 5 48. Sao Tome and Principe.mp.
- 6 49. Benin/
- 7 50. Kiribati.mp.
- 8 51. Senegal/
- 9 52. Bhutan/
- 10 53. Kyrgyzstan/
- 11 54. Solomon Islands.mp.
- 12 55. Bolivia/
- 13 56. Laos/
- 14 57. Sri Lanka/
- 15 58. Cabo Verde/
- 16 59. Lesotho/
- 17 60. Tanzania/
- 18 61. Cambodia/
- 19 62. Mauritania/
- 20 63. Timor-Leste/
- 21 64. Cameroon/
- 22 65. Micronesia/
- 23 66. Tunisia/
- 24 67. Comoros/
- 25 68. Moldova/
- 26 69. Ukraine/
- 27 70. "Democratic Republic of the Congo"/
- 28 71. Mongolia/
- 29 72. Uzbekistan
- 30 73. Cote d'Ivoire/
- 31 74. Morocco/
- 32 75. Vanuatu/
- 33 76. Djibouti/
- 34 77. Myanmar/
- 35 78. Vietnam/
- 36 79. Egypt/
- 37 80. Nepal/
- 38 81. West Bank and Gaza.mp.
- 39 82. El Salvador/
- 40 83. Nicaragua/
- 41 84. Zambia/
- 42 85. Eswatini/
- 43 86. Nigeria/
- 44 87. Zimbabwe/
- 45 88. Ghana/
- 46 89. Pakistan/
- 47
- 48 90. (Angola or Honduras or Papua New Guinea or Algeria or India or Philippines or Bangladesh or Kenya or Sao
- 49 Tome and Principe or Benin or Kiribati or Senegal or Bhutan or Kyrgyz Republic or Solomon Islands or Bolivia or
- 50 Lao PDR or Sri Lanka or Cabo Verde or Lesotho or Tanzania or Cambodia or Mauritania or Timor-Leste or
- 51 Cameroon or Micronesia or Tunisia or Comoros or Moldova or Ukraine or Democratic Republic of the Congo or
- 52 Mongolia or Uzbekistan or Cote d'Ivoire or Morocco or Vanuatu or Djibouti or Myanmar or Vietnam or Egypt
- 53 or Nepal or West Bank and Gaza or El Salvador or Nicaragua or Zambia or Eswatini or Nigeria or Zimbabwe or
- 54 Ghana or Pakistan).tw,kw.
- 55 91. or/37-90
- 56 92. 36 and 91
- 57
- 58
- 59
- 60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 93. exp animals/ not humans.sh.
- 94. 92 not 93
- 95. remove duplicates from 94

For peer review only



## Supplementary File 3 - Data Abstraction Form

1. Author, year
2. Journal
3. Study design (observational, experimental, protocol paper)
4. Study setting/country (LMICs)
5. Study population/Health Condition (PW at HRPE/E)
6. Study Objective
7. Number of participants (Sample)
8. Study period
  - Duration of intervention
  - Duration of data collection
9. Digital health intervention (DHI) used
  - Type of DHI (Predictive model, mHealth applications, devices)
  - Intervention validation (Yes/No)
  - Targeted primary user of intervention (health care provider/pregnant women)
  - User training on use of digital health intervention (Yes/No)
  - Function of digital health intervention
10. Study outcomes
  - Maternal and fetal health outcomes
  - Intervention feasibility/usability/fidelity/acceptability
11. Framework/model used
12. Study limitations
13. Comments

## Supplementary File 4 - Overview of the Included Articles

Reference	Year	Study Title	Type of Study Design	Objective	Setting	N	Health Condition	Purpose of Digital Health Intervention
Musyoka et al. [25]	2019	A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care	Prospective experimental study	The study sought to implement a 24-hour ambulatory blood pressure monitoring solution for preeclampsia management, using a smartwatch in conjunction with a mobile and cloud-based application.	Kenya	N=30	preeclampsia	Monitoring
Lim et al. [24]	2015	Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage	Observational	The aim of this study was to assess the usability of PIERS on the Move PotM (with mid-level health workers) for iteratively refining the system.	South Africa	N=37	preeclampsia	Predicting
Vousden et al. [34]	2018	Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial	Observational	Prior to the CRADLE 3 trial start, a mixed-methodology feasibility study was undertaken to finalise the intervention and implementation processes which were guided by the Expert Recommendations for Implementing Change (ERIC) project	Zimbabwe, Ethiopia, India	Number of HCP trained=204	Preeclampsia, eclampsia and shock	Monitoring
Nathan et al. [26]	2018	The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings	Observational	This qualitative study aimed to determine the usability, feasibility and acceptability of the CRADLE VSA among a variety of users and in diverse socio-economic settings, considering these five clusters of influence. This will inform future device modifications and successful dissemination of the CRADLE VSA for routine use.	India, Mozambique, Nigeria and South Africa	N=205	Preeclampsia and shock	Monitoring
Feroz et al. [20]	2020	Exploring perspectives, preferences and needs of a telemonitoring program for women at high risk for preeclampsia in a	Protocol paper	The study aims to explore the perspectives, preferences, and needs of telemonitoring (TM) for pregnant women at HRPE in Karachi, to inform future implementation strategies.	Pakistan	N=30	Preeclampsia	Monitoring

		tertiary health facility of Karachi: a qualitative study protocol						
Dunsmuir et al. [19]	2014	Development of mHealth Applications for Pre-Eclampsia Triage	Observational	This paper describes the design process of two versions of the POTM application, the original version application referred to as POTM), and a simplified, community-based version for the Community Level Interventions for Pre-eclampsia cluster randomized controlled trial (application referred to as CLIP POTM),	Nigeria, Mozambique, Pakistan, and India	Projected +30,000 pregnant women  500 community HCPs	Preeclampsia	Predicting
Jonas et al. [21]	2016	Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources	Prospective experimental study design	The study proposes an innovative mobile health (mHealth) solution that enables the quantification of the congo red dot test as a batch laboratory test, with minimal cost and equipment.	Resource poor settings	N=273	preeclampsia	Monitoring
Thakor et al. [32]	2009	Hypertension Detector for Developing Countries	Observational	A prototype of a low-cost device engineered specifically for semi-literate volunteers in developing countries has been created.	Africa, Southern Asia, and the Middle East	-	Preeclampsia	Monitoring
Nathan et al. [27]	2015	An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2	Observational	The study aims to assess the accuracy of the Microlife 3AS1-2 blood pressure device in pregnancy and pre-eclampsia in a low-resource setting.	South Africa	N=45	Preeclampsia	Monitoring
Nathan et al. [28]	2018	Early warning system hypertension thresholds to predict adverse outcomes	Observational	The study aims to evaluate the association between blood pressure (BP) measurements and adverse outcomes in women with pre-eclampsia.	South Africa	N= 1547	Preeclampsia	Monitoring

		in pre-eclampsia: A prospective cohort study						
Payne et al. [29]	2014	A Risk Prediction Model for the Assessment and Triage of Women with Hypertensive Disorders of Pregnancy in Low-Resourced Settings: The miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) Multi-country Prospective Cohort Study	Observational	The objective of the miniPIERS study was to develop and validate a simplified clinical prediction model for adverse maternal outcomes among women with HDP for use in community and primary health care facilities in LMICs.	LMICs	N= 2,133	Preeclampsia	Predicting
Bellad et al. [17]	2020	Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial	Experimental study (RCT)	The objective of the Community-Level Interventions for reeclampsia (CLIP) India cluster randomised controlled trial (cRCT) was to test the hypothesis that implementing community-level, evidence-based care focused on pregnancy hypertension would reduce all-cause maternal, fetal and newborn mortality and major morbidity, without causing harm	India	N=14,783 pregnancies	Preeclampsia	Monitoring and Predicting
Qureshi et al. [30]	2020	Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial	Experimental study (RCT)	The aim of the Community-Level Interventions for Pre-eclampsia (CLIP) cluster randomised controlled trial (cRCT) in Sindh Province, Pakistan was to reduce maternal and perinatal mortality and major morbidity by 20% or more in intervention (vs. control) clusters, through a community-level intervention to address triage, (initial) treatment, and transport (to facility) of women with pregnancy hypertension.	Pakistan	N= 35,974 women	Preeclampsia	Monitoring and Predicting
Khowaja et al [22]	2015	Economic evaluation of Community Level Interventions for Pre-eclampsia (CLIP) in South Asian and African countries: a study protocol	Protocol paper	The study aims to conduct an economic evaluation alongside of the CLIP Trial, to inform decision makers not only of clinical outcomes but the cost required to obtain those outcomes.	Nigeria, Mozambique, Pakistan, and India	N= 154,000	Preeclampsia	Monitoring
Khowaja et al [23]	2016	The feasibility of community level	Observational study	The study aimed to describe the health system, identify community and individual barriers and facilitators that influence care of pregnant women	Nigeria, Mozambique,	N= 337 (health facilities)	Preeclampsia	Monitoring

		interventions for pre-eclampsia in South Asia and Sub-Saharan Africa: a mixed-methods design		in the community, in preparation for the conduct of a community-based cluster randomized trial	Pakistan, and India	N= 100 (IDIs) N= 123 (FGDs)		
Von Dadelzen et al. [33]	2020	The PRECISE (PREgnancy Care Integrating translational Science, Everywhere) Network's first protocol: deep phenotyping in three sub-Saharan African countries	Protocol paper	This paper describes the protocol that underpins the clinical research activity of the Network, so that the investigators, and broader global health community, can have access to 'deep phenotyping' of women as they advance through pregnancy to the end of the puerperium.	Gambia Kenya Mozambique	N= 600 (each country)	Preeclampsia, and eclampsia	Monitoring
Abejirinde et al [16]	2018	Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana	Observational	This paper therefore explores the experiences of women exposed to the B4M device, to answer the research questions: i) How did women experience the use of Bliss4Midwives during their routine antenatal care consultations? ii) What influence did Bliss4Midwives have on woman-provider relationships and on ANC service utilization?	Ghana	N=30	preeclampsia, gestational diabetes and anaemia	Monitoring
Bellad et al [18]	2017	Maternal and Newborn Health in Karnataka State, India: The Community Level Interventions for Pre-Eclampsia (CLIP) Trial's Baseline Study Results	Observational	To describe baseline demographics and health outcomes prior to initiation of the CLIP trial and to improve knowledge of population-level health, in particular of maternal and neonatal outcomes related to hypertensive disorders of pregnancy, in northern districts the state of Karnataka, India.	India	N= 5,469	Hypertension disorders of pregnancy, preeclampsia	Monitoring and Predicting
Sharma et al [31]	2017	A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria	Observational	To evaluate implementation processes of the complex CLIP intervention, assess mechanisms of impact and identify emerging unintended causal pathways.	Nigeria	N= 32,785	preeclampsia	Monitoring and Predicting

Supplementary File 5- Digital Health Intervention Characteristics

Reference	Digital health intervention	Validated	Intervention use for	Technological component(s)	Targeted primary user	User training
Musyoka et al. [25]	24-hour ambulatory blood pressure monitoring system	Validated	Blood pressure data collection	F1 smart wristwatch Blood Pressure Monitoring Mobile Application Cloud Data center Caregiver's smartphone	Expectant mother and the caregiver	Not specified
Lim et al. [24]	Pre-eclampsia Integrated Estimate of RiSk (PIERS) on the Move (PotM)	Not specified	Demographics (gestational age at presentation), clinical signs (blood pressure, SPO2 and dipstick proteinuria), and symptoms (chest pain or dyspnoea, headache or visual disturbances, vaginal bleeding with abdominal pain)	mHealth platform	Mid-level health workers	Yes
Vousden et al. [34]	CRADLE (Community blood pressure monitoring in Rural Africa & Asia: Detection of	Validated	Measures blood pressure, pulse and calculates the mothers risk of shock  A traffic light Early Warning System display alerts users to	Microlife CRADLE VSA device	Healthcare providers	Yes

	underlying pre-Eclampsia and shock) Vital Sign Alert		abnormalities in the vital signs results.			
Nathan et al. [26]	Microlife® CRADLE (Community blood pressure monitoring in Rural Africa & Asia: Detection of underlying pre-Eclampsia and shock) Vital Signs Alert (VSA)	Validated	Device accurately measures blood pressure and pulse. Traffic lights within the device help healthcare workers identify women who need additional treatment for these conditions	Microlife® CRADLE VSA device	Healthcare providers	Yes
Feroz et al. [20]	Hypothetical elemonitoring program	Not specified	Blood pressure measurement	-	Pregnant women and caregiver	-
Dunsmuir et al. [19]	MiniPIERS AND FullPIERS models  Two versions of the POTM application, 1) Original version (application referred to as POTM), 2)Simplified, community-based version for the Community Level Interventions for Pre-eclampsia cluster randomized controlled trial (application referred to as CLIP	Not specified	Mean BP, SpO2, gestational age, proteinuria, symptoms.	Smartphone, mobile health applications (POTM/CLIP POTM), Research electronic data capture server	Community-based health care providers	Yes



	POTM)					
Jonas et al. [21]	Smartphone-based diagnostic test (Congo Red Dot) for preeclampsia	Validated	urine markers	mHealth solution for administering the Congo Red Dot (CRD) test	Modestly trained personnel	Yes
Thakor et al. [32]	New device (Hypertension Detector for Developing Countries), intraarterial, sphygmomanometers, assorted automatic blood pressure devices, and proteinuria measurement	Not specified	Blood pressure measurement	-	Semi-literate volunteers with minimal training	Yes
Nathan et al. [27]	Microlife 3AS1-2 blood pressure device	Validated	Measures blood pressure	Device	Staff with minimal training	Yes
Nathan et al. [28]	CRADLE Vital Signs Alert (VSA)	Validated	Measures BP and pulse to facilitate prompt recognition of abnormalities in vital signs	Device Traffic light early warning system	Healthcare providers	yes
Payne et al. [29]	miniPIERS risk prediction model	Validated	miniPIERS (measures demographics, symptoms and signs).	Mobile health application	Mid-level health workers	yes
Bellad et al. [17]	CLIP intervention package included miniPIERS model, PIERS On the Move (POM) tool, and	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	mobile-based CLIP POM mobile health application (app),	Community health workers	yes

136/bmjopen-2021-056130 on 8 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

	Microlife BP 3AS1-2 device			central REDCap server		
Qureshi et al. [30]	CLIP intervention package included miniPIERS model, Microlife BP 3AS1-2 device and PIERS On the Move (POM) mobile health (mHealth) application	Validated	BP measurement and pulse oximetry	POM mHealth application	Lady health workers	Yes
Khowaja et al [22]	CLIP intervention package PIERS On the Move (POM) mobile health (mHealth) application	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	POM mHealth application	Community-based health care providers	Yes
Khowaja et al [23]	CLIP intervention package PIERS On the Move (POM) mobile health (mHealth) application	Validated	Measure BP, pre-eclampsia symptoms and dipstick proteinuria	POM mHealth application	Community-based health care providers	Yes
Von Dodelszen et al. [33]	CRADLE BP device, pulse oximetry and TraCer platform, POM mHealth application	Validated	CRADLE VSA semi-automated and validated BP device will be used for all clinical measurements of blood pressure (BP) in the study pulse oximetry  POM platform to provide time-of-disease	POM mHealth application	Healthcare providers	Yes

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

			risk estimates to hypertensive pregnant women using PIERS models			
Abejirinde et al [16]	Bliss4Midwives' (B4M)	Not specified	non-invasive device for measuring haemoglobin via infrared sensors mounted on a finger clip; a self-inflating blood pressure cuff; and an automated reader for urinary protein and glucose through dipsticks.	Data from all diagnostic devices are automatically or manually linked to an android tablet equipped with decision support algorithms	Midwives and community health nurses	Yes
Bellad et al [18]	Community Level Interventions for Pre-Eclampsia (CLIP) Package	Not specified	Measuring blood pressure	mHealth platform	Community-based health activists ASHAs	Not specified
Sharma et al [31]	Community Level Interventions for Pre-Eclampsia (CLIP) Package	Not specified	Blood pressure measurement	PIERS On the Move (POM) mHealth tool, Microlife VSA blood pressure device	Community health workers	Yes

## Supplementary File 6 - Outcomes of Digital Health Interventions

Reference	Study Title	Study outcomes pertaining to digital health intervention use	Framework/model used
<b>Maternal and fetal health outcomes (4 studies)</b>			
Nathan et al. (2018) [28]	Early warning system hypertension thresholds to predict adverse outcomes in pre-eclampsia: A prospective cohort study	Of 1547 women with pre-eclampsia, 33.0% of women triggered a red light on admission and 78.6% at their highest BP. Severe hypertension and adverse outcomes were common across yellow and red categories. Comparing admission red to yellow lights, there was a significant increase in kidney injury (OR 1.74, CI 1.31–2.33, test p=.003), magnesium sulfate use (OR 3.40, CI 2.24–5.18, p < .001) and CCU admission (OR 1.50, CI 1.18–1.91, p < .001), but not for maternal death, eclampsia, extended perinatal death or preterm delivery.	No framework described
Bellad et al. (2020) [17]	Community level interventions for pre-eclampsia (CLIP) in India: A cluster randomised controlled trial	The primary outcome did not differ between intervention and control arms (adjusted odds ratio (aOR) 0.92 [95% confidence interval 0.74, 1.15]; p = 0.47; intraclass correlation coefficient 0.013). There was no intervention-related safety concerns following administration of either methyldopa or MgSO <sub>4</sub> , and 401 facility referrals. Compared with intervention arm women without CLIP contacts, those with ≥8 contacts suffered fewer stillbirths (aOR 0.19 [0.10, 0.35]; p < 0.001), at the probable expense of survivable neonatal morbidity (aOR 1.39 [0.97, 1.99]; p = 0.072).	No framework described
Qureshi et al. (2020) [30]	Community-level interventions for pre-eclampsia (CLIP) in Pakistan: A cluster randomised controlled trial.	The primary outcome did not differ between intervention (26.6%) and control (21.9%) clusters (adjusted odds ratio, aOR, 1.20 [95% confidence interval 0.84- 1.72]; p = 0.31). There was reduction in stillbirths (0.89 [0.81-0.99]; p = 0.03), but no impact on maternal death (1.08 [0.69, 1.71]; p = 0.74) or morbidity (1.12 [0.57, 2.16]; p = 0.77); early (0.99 [0.82-1.09]; p = 0.46) or late neonatal deaths (1.23 [0.97-1.55]; p = 0.09); or neonatal morbidity (1.22 [0.77, 1.96]; p = 0.40). Improvements in outcome rates were observed with 4–7 (p = 0.015) and ≥8 (p < 0.001) (vs. 0) CLIP contacts.	No framework described
Bellad et al. (2017) [18]	Maternal and newborn health in Karnataka state, India: the community level interventions for pre-eclampsia (CLIP) Trial's baseline study results	A majority of the women reported institutional deliveries (96.0%), largely attended by skilled birth attendants. The maternal mortality ratio of 104 (per 100,000 livebirths) was observed during this study, neonatal mortality ratio was 25 per 1,000 livebirths, and perinatal mortality ratio was 50 per 1,000 livebirths. Despite a high number of institutional deliveries, rates of stillbirth were 2.86%.	No framework described
<b>Usability and acceptability (5 studies)</b>			

136/bmjopen-2024-058730 on 8 February 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

Musyoka et al. (2019) [25]	A 24-hour ambulatory blood pressure monitoring system for preeclampsia management in antenatal care. Informatics in Medicine Unlocked.	Content richness has a slightly positive linear effect on Perceived Ease of Use, while there is a slightly negative relationship between Content Richness and Perceived Usefulness. Overall, the 24-hour ambulatory blood pressure monitoring system has shown great potential for actual adoption in healthcare systems in developing countries, given its simplicity and affordability.	Technology Acceptance Model
Lim at al. (2015) [24]	Usability and Feasibility of PIERS on the Move: An mHealth App for Pre-Eclampsia Triage.	Overall, users felt the app was usable using the Computer Systems Usability Questionnaire; median (range) values for Study 1 = 2 (1-6) and Study 2 = 1 (1-7). Usability problems were often related to mobile phone features (eg, scroll wheels, touch screen use).	LambdaNative framework for app development
Nathan et al. (2018) [26]	The CRADLE vital signs alert: qualitative evaluation of a novel device designed for use in pregnancy by healthcare workers in low-resource settings.	Most HCWs perceived the CRADLE device to be easy to use and accurate. The traffic lights early warning system was unanimously reported positively, giving HCWs, Pregnant women and families understanding of vital signs and confidence with decision-making. Some described manual inflation as tiring, particularly when measuring vital signs in obese and hypertensive women (n=4) and a few South African HCWs distrusted the device's accuracy (n =7).	Diffusion of innovation model Three delay model
Thakor et al. (2010) [32]	Hypertension Detector for Developing Countries.	The study developed a prototype of a low-cost device engineered specifically for semi-literate volunteers in developing countries. Preliminary testing has shown reliable hypertension detection and plans have been made for field testing in rural communities this August 2010 in Nepal.	No framework described
Dunsmuir et al (2014) [19]	Development of mHealth applications for pre-eclampsia triage. IEEE J Biomed Health Inform.	The paper outlines the POTM application development process. The paper concludes that the successful development of an mHealth tool, must consider the user and the setting in which it is deployed. CLIP POTM began with a single specification document, but study discovered differing requests from the different countries with their cultural differences, leading to modified application versions for each country.	LambdaNative Framework for developing application
<b>Intervention Feasibility and Fidelity (7 studies)</b>			
Vousden et al (2018) [34]	Evaluation of a novel vital sign device to reduce maternal mortality and morbidity in low-resource settings: a mixed method feasibility study for the CRADLE-3 trial	Intervention was implemented with high fidelity (85% of HCP trained, n=204). Results indicated a good understanding of device use with 75% of participants scoring >75% (n=97; 90% of those distributed). Interviews with HCPs reported that the intervention improved capacity to make clinical decisions, escalate care and make appropriate referrals.	Medical Research Council framework and logic model
Khowaja et al (2016) [23]	The feasibility of community level interventions for pre-eclampsia in South Asia and	The study highlight enabling factors including need for community mobilization, awareness raising programs, institutional support, community safety nets for	Normalization process theory

	Sub-Saharan Africa: a mixed-methods design.	emergency funds, and system integration. Whereas, impeding factors included delays in care seeking, knowledge gaps, lack of trained human resource, cultural myths and misconceptions, high cost of care, and poor health service quality.	
Abejirinde et al (2018) [16]	Pregnant women's experiences with an integrated diagnostic and decision support device for antenatal care in Ghana.	Pregnant women generally valued the availability of diagnostic services at the point-of-care. The intervention made women feel listened to and cared for. Process outcomes of the B4M encounter also showed that it was perceived as improving the skills and knowledge of the health workers which facilitated trust in diagnostic recommendations and was therefore believed to motivate referral compliance.	No framework described
Sharma et al (2017) [31]	A process evaluation plan for assessing a complex community-based maternal health intervention in Ogun State, Nigeria.	This paper offers robust measures of the process indicators, external validity of conclusions about effectiveness can best be complemented by efficacy studies using a RCT. The methodology allows to examine the internal validity of the efficacy of the intervention by assessing the implementation (quantity and quality) of what is delivered.	Logic model, Diffusions of innovations and realist evaluation theories
Nathan et al (2015) [27]	An accurate semiautomated oscillometric blood pressure device for use in pregnancy (including pre-eclampsia) in a low-income and middle-income country population: the Microlife 3AS1-2	The Microlife 3AS1-2 device achieved an overall B/A grade in pregnancy (including pre-eclampsia), passing the British Hypertension Society protocol requirements and achieving the International Organization for Standardization standard with a mean difference and SD of $-3.8 \pm 7.3$ and $-1.5 \pm 6.2$ mmHg for systolic and diastolic pressures, respectively. The device can be recommended for use in pregnancy, including preeclampsia. Also, it fulfils the requirements of WHO for an automated blood pressure device suitable for use in a low-resource setting.	No framework described
Payne et al (2014) [29]	A risk prediction model for the assessment and triage of women with hypertensive disorders of pregnancy in low-resourced settings: the miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) multi-country prospective cohort study.	The miniPIERS model was well-calibrated and had an area under the receiver operating characteristic curve (AUC ROC) of 0.768 (95% CI 0.735–0.801) with an average optimism of 0.037. External validation AUC ROC was 0.713 (95% CI 0.668–0.768). A predicted probability $\geq 25\%$ to define a positive test classified women with 85.5% accuracy. The miniPIERS model shows reasonable ability to identify women at increased risk of adverse maternal outcomes associated with the hypertensive disorders of pregnancy	Three delay model
Jonas et al. (2016) [21]	Smartphone-based diagnostic for preeclampsia: an mHealth solution for administering the Congo Red Dot (CRD) test in settings with limited resources.	The results suggests that combining smartphone-based image analysis with molecular-specific disease features represents a cost-effective application of mHealth that has the potential to fill gaps in access to health care solutions that are critical to reducing adverse events related to PE in resource-poor settings	No framework described

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47



## Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist

SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
<b>TITLE</b>			
Title	1	Identify the report as a scoping review.	1
<b>ABSTRACT</b>			
Structured summary	2	Provide a structured summary that includes (as applicable): background, objectives, eligibility criteria, sources of evidence, charting methods, results, and conclusions that relate to the review questions and objectives.	1
<b>INTRODUCTION</b>			
Rationale	3	Describe the rationale for the review in the context of what is already known. Explain why the review questions/objectives lend themselves to a scoping review approach.	2&3
Objectives	4	Provide an explicit statement of the questions and objectives being addressed with reference to their key elements (e.g., population or participants, concepts, and context) or other relevant key elements used to conceptualize the review questions and/or objectives.	3
<b>METHODS</b>			
Protocol and registration	5	Indicate whether a review protocol exists; state if and where it can be accessed (e.g., a Web address); and if available, provide registration information, including the registration number.	3
Eligibility criteria	6	Specify characteristics of the sources of evidence used as eligibility criteria (e.g., years considered, language, and publication status), and provide a rationale.	3&4
Information sources*	7	Describe all information sources in the search (e.g., databases with dates of coverage and contact with authors to identify additional sources), as well as the date the most recent search was executed.	4
Search	8	Present the full electronic search strategy for at least 1 database, including any limits used, such that it could be repeated.	4
Selection of sources of evidence†	9	State the process for selecting sources of evidence (i.e., screening and eligibility) included in the scoping review.	4&5
Data charting process‡	10	Describe the methods of charting data from the included sources of evidence (e.g., calibrated forms or forms that have been tested by the team before their use, and whether data charting was done independently or in duplicate) and any processes for obtaining and confirming data from investigators.	5
Data items	11	List and define all variables for which data were sought and any assumptions and simplifications made.	Appendix IV
Critical appraisal of individual sources of evidence§	12	If done, provide a rationale for conducting a critical appraisal of included sources of evidence; describe the methods used and how this information was used in any data synthesis (if appropriate).	5-6



SECTION	ITEM	PRISMA-ScR CHECKLIST ITEM	REPORTED ON PAGE #
Synthesis of results	13	Describe the methods of handling and summarizing the data that were charted.	5
<b>RESULTS</b>			
Selection of sources of evidence	14	Give numbers of sources of evidence screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally using a flow diagram.	4&5
Characteristics of sources of evidence	15	For each source of evidence, present characteristics for which data were charted and provide the citations.	Appendix V-VII
Critical appraisal within sources of evidence	16	If done, present data on critical appraisal of included sources of evidence (see item 12).	5-6
Results of individual sources of evidence	17	For each included source of evidence, present the relevant data that were charted that relate to the review questions and objectives.	7-10/ Appendix V-VII
Synthesis of results	18	Summarize and/or present the charting results as they relate to the review questions and objectives.	7-10
<b>DISCUSSION</b>			
Summary of evidence	19	Summarize the main results (including an overview of concepts, themes, and types of evidence available), link to the review questions and objectives, and consider the relevance to key groups.	10-12
Limitations	20	Discuss the limitations of the scoping review process.	12
Conclusions	21	Provide a general interpretation of the results with respect to the review questions and objectives, as well as potential implications and/or next steps.	12
<b>FUNDING</b>			
Funding	22	Describe sources of funding for the included sources of evidence, as well as sources of funding for the scoping review. Describe the role of the funders of the scoping review.	NA

JBI = Joanna Briggs Institute; PRISMA-ScR = Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews.

\* Where *sources of evidence* (see second footnote) are compiled from, such as bibliographic databases, social media platforms, and Web sites.

† A more inclusive/heterogeneous term used to account for the different types of evidence or data sources (e.g., quantitative and/or qualitative research, expert opinion, and policy documents) that may be eligible in a scoping review as opposed to only studies. This is not to be confused with *information sources* (see first footnote).

‡ The frameworks by Arksey and O'Malley (6) and Levac and colleagues (7) and the JBI guidance (4, 5) refer to the process of data extraction in a scoping review as data charting.

§ The process of systematically examining research evidence to assess its validity, results, and relevance before using it to inform a decision. This term is used for items 12 and 19 instead of "risk of bias" (which is more applicable to systematic reviews of interventions) to include and acknowledge the various sources of evidence that may be used in a scoping review (e.g., quantitative and/or qualitative research, expert opinion, and policy document).

From: Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med.* 2018;169:467–473. doi: [10.7326/M18-0850](https://doi.org/10.7326/M18-0850).

