

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

The role of educational settings in the transmission chain of SARS-CoV-2 in 2020: a systematic review

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058308
Article Type:	Original research
Date Submitted by the Author:	14-Oct-2021
Complete List of Authors:	Vardavas, Constantine; University of Crete; Harvard University, Department of Oral Health Policy and Epidemiology Nikitara, Katerina; University of Crete, Mathioudakis, Alexander; The University of Manchester, Immunity and Respiratory Medicine; Manchester Academic Health Science Centre Boon, Michelle; Glasgow Caledonian University, WISE Centre for Economic Justice Phalkey, Revati; Public Health England; University of Nottingham Leonardi-Bee, Jo; University of Nottingham, Division of Epidemiology and Public Health, School of Medicine Pharris, Anastasia; European Centre for Disease Prevention and Control Deogan, Charlotte; European Centre for Disease Prevention and Control Suk, Jonathan; European Centre for Disease Prevention and Control
Keywords:	COVID-19, Public health < INFECTIOUS DISEASES, Community child health < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1 The role of educational settings in the transmission chain of SARS- 2 CoV-2 in 2020: a systematic review

3 **Authors:** Constantine Vardavas^{1,2}, Katerina Nikitara¹, Alexander Mathioudakis^{3,4}, Michele
4 Hilton-Boon⁵, Revati Phalkey^{6,7}, Jo Leonardi-Bee⁷, Anastasia Pharris⁸, Charlotte Deogan⁸,
5 Jonathan E. Suk^{8*}

7 Affiliations

8 1: School of Medicine, University of Crete, Greece

9 2: Department of Oral Health Policy and Epidemiology, Harvard School of Dental Medicine, Harvard
10 University, Boston, MA, USA

11 3: Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The
12 University of Manchester, Manchester, UK.

13 4: North West Lung Centre, Wythenshawe Hospital, Manchester University NHS Foundation Trust,
14 Manchester Academic Health Science Centre, Manchester, UK.

15 5: WISE Centre for Economic Justice, Glasgow Caledonian University

16 6: Climate Change and Health Group, Public Health England, United Kingdom

17 7: Division of Epidemiology and Public Health, School of Medicine, University of Nottingham, UK

18 8: European Centre for Disease Prevention and Control, Solna, Sweden

19
20
21 ***: Corresponding author: Jonathan E. Suk, jonathan.suk@ecdc.europa.eu, +46858601633. Gustav III:s
22 Boulevard, 169 73 Solna, Sweden.**

23 Word counting: 3216 words

24 Keywords: SARS-CoV-2, COVID-19, pandemic, children, schools, household

27 **Abstract (240 words)**

28 **Objectives:** School closures have been used as a core non pharmaceutical intervention during
29 the COVID-19 pandemic. This review aims at identifying the role of children in COVID-19
30 transmission in educational settings.

31 **Methods:** This systematic literature review assessed studies published between December
32 2019 and April 1, 2021 in Medline and Embase, which included studies that assessed
33 educational settings from approximately January 2020 to January 2021. The inclusion criteria
34 were based on the PCC framework (P-Population, C-Concept, C-Context). The study
35 *Population* was restricted to people 1-17 years old (excluding neonatal transmission), the
36 *Concept* was to assess child-to-child and child-to-adult transmission, while the *Context* was to
37 assess specifically educational setting transmission clusters.

38 **Results:** Fifteen studies met inclusion criteria, ranging from daycare centers to high schools
39 and summer camps, while eight studies assessed the re-opening of schools in the 2020-2021
40 school year. In principle although there is sufficient evidence that children can both be
41 infected by and transmit SARS-CoV-2 in school settings, the SAR remain relatively low -when
42 NPI measures are implemented in parallel. Moreover, although the evidence was limited
43 there was an indication that younger children may have a lower SAR than adolescents.

44 **Conclusions:** Transmission in educational settings in 2020 was minimal -when NPI measures
45 were implemented in parallel. However, with an upsurge of cases related to variants of
46 concern, continuous surveillance and assessment of the evidence is warranted to ensure the
47 maximum protection of the health of students and the educational workforce, while also
48 minimising the numerous negative impacts that school closures may have on children.

49 **Strengths and limitations of this study**

- 50 • This study provides a rapid review of the peer-reviewed literature pertaining to SARS-CoV-
51 2 transmission by children within educational settings.
- 52 • The review reflects the status quo of the previous school years (January 2020 - January
53 2021) due to the lag time between study implementation, peer review and publication.
- 54 • The included studies represent child-to-child transmission within the context of previous
55 SARS-CoV-2 strains and are not directly applicable to newer variants.

1
2
3 56 **MAIN TEXT**
4

5
6 57 **INTRODUCTION**
7

8 58 One of the more perplexing and controversial dimensions during the first year of the COVID-
9 59 19 pandemic surrounded the role of children in the transmission. Are they drivers of the
10 60 pandemic, or are they merely innocent bystanders, affected in myriad ways by school closures
11 61 and other physical distancing measures while not being generally at-risk of COVID-19
12 62 themselves?
13

14
15
16
17
18 63 Epidemiologic indicators of SARS-CoV-2 infection in children provide a complex picture
19 64 regarding their potential role in the transmission chain. Systematic reviews have concluded
20 65 that children and adolescents have lower susceptibility to SARS-CoV-2 infection [1, 2].
21
22 66 However, when infected and symptomatic, children may shed viral RNA in similar quantities
23 67 to adults [3], and that younger children (under 5 years) with mild to moderate symptoms may
24 68 shed even more virus than older children and adults [4]. While the proportion of
25 69 asymptomatic SARS-CoV-2 infections among children in the general population is uncertain,
26 70 initial data had indicated that 16% of paediatric cases in Europe in the first phase of the
27 71 pandemic were classified as asymptomatic [5], while up to 90% of paediatric cases in China
28 72 were deemed to be asymptomatic, mild, or moderate [6]. Moreover, it is possible that
29 73 children are less often asymptomatic carriers than adults: a study of non-COVID-19-related
30 74 hospitalizations in Milan identified 1% of children and 9% of adults as asymptomatic carriers
31 75 of SARS-CoV-2 [7]. Meanwhile, while children are overall noted to have lower rates of severe
32 76 COVID-19 cases [8], there is evidence of differing transmission dynamics between younger vs.
33 77 older children [2]. There is evidence that when index cases, younger children, such as under
34 78 10 years of age, lead to lower secondary attack rates than older children and adult [9, 10].
35
36
37
38
39
40
41
42
43
44
45
46
47

48 79 Important potential sources of evidence surrounding the role of children in the COVID-19
49 80 pandemic come from studies situated in the community, household, healthcare or
50 81 educational settings. Transmission of SARS-COV-2 has thus far been documented to be higher
51 82 in household settings than in other community settings – including schools – a finding which
52 83 may be potentially attributable to the individual, behavioural and contextual factors of
53 84 households vs. other settings, as has been suggested elsewhere [9].
54
55
56
57
58
59
60

1
2
3 86 Although, at the time of writing, the more transmissible Delta variant of concern is driving
4
5 87 SARS-CoV-2 transmission (*ref to add ECDC RRA 16-pending*) there is currently a gap in
6
7 88 published studies looking at the transmission of Delta in school settings. However, as
8
9 89 decisions currently need to be taken to ensure high levels of preparedness in school settings
10
11 90 [11], the literature published thus far may have important insights to guide decision-making
12
13 91 around school closures and re-openings, as well support decision making for mitigation
14
15 92 measures in educational settings. This systematic literature review was conducted to assess
16
17 93 child-to-child and child-to-adult SARS-CoV-2 transmission within educational settings and to
18
19 94 calculate where possible the secondary attack rate (SAR) when the child is the index case.

20 95 **METHODS**

21 96 ***Search Strategy***

22
23
24 97 This systematic literature review is reported in accordance with the Preferred Reporting Items
25
26 98 for Systematic Reviews and Meta-Analysis (PRISMA) guidelines [12]. Relevant studies
27
28 99 published between December 2019 and April 1, 2021 were identified by searching Medline
29
30 100 and Embase. The following set of inclusion criteria were used to determine eligibility of the
31
32 101 studies, which is based on the PCC framework (P-Population, C-Concept, C-Context). The
33
34 102 study *Population* was restricted to people 1-17 years old (excluding neonatal transmission
35
36 103 [13]), the *Concept* was to assess child-to-child and child-to-adult transmission when the child
37
38 104 is the index case, while the *Context* was to assess specifically educational setting transmission
39
40 105 clusters. Subject heading terms and free text words relating to the Population, Concept and
41
42 106 Context terms as identified in the inclusion criteria were used to develop a comprehensive
43
44 107 list of terms for the search strategy, from which this specific review was based. We included
45
46 108 all studies of quantitative research, while, opinion pieces, commentaries, case reports and
47
48 109 editorials were excluded. Mathematical modelling and simulation studies were also excluded.
49
50 110 We additionally screened reference lists of the included articles to identify further relevant
51
52 111 studies. The search was limited to the English language.

53 112 54 113 ***Study selection***

55 114 Initially, a pilot training screening process was used where 100 identical articles were
56
57 115 screened for their eligibility independently by two reviewers to ensure consistency in
58
59 116 screening. As a high measure of inter-rater agreement was achieved between the two
60

1
2
3 117 reviewers during the pilot assessment (percentage agreement >90% and/ or Cohen's Kappa
4 118 >0.81), the remaining titles were randomly allocated to the two reviewers and screened for
5
6 119 eligibility independently by them. After an initial selection of the titles, each reviewer
7
8 120 assessed each other's selected studies. The retrieved articles were then independently
9
10 121 double-screened by two reviewers based on the full text of the articles.
11
12

13 122 **Data extraction**

14
15 123 The data extraction template was piloted independently by the two reviewers on a random
16
17 124 sample of two included studies to enable an assessment of consistency in data extraction and
18
19 125 to identify where amendments needed to be made to the template. The remaining studies
20
21 126 were then data extracted independently by two reviewers, and the results were double
22
23 127 checked across the original manuscript by a third reviewer.
24

25 128 **Data synthesis**

26
27 129 Characteristics of the included studies were presented in tabulated form detailing the study
28
29 130 design, geographical location of the study, sample size, characteristics of the populations
30
31 131 considered, setting, context, parallel implemented Non Pharmaceutical Interventions (NPI),
32
33 132 and the findings of the study. Depending on the level of information available, infection SAR
34
35 133 were noted. A narrative synthesis approach was applied to look systematically at the data and
36
37 134 to describe each study categorized by the study design. Patterns in the data were identified
38
39 135 through tabulation of results, and an inductive approach was taken to translate the data to
40
41 136 identify areas of commonality between studies.
42

43 137 **Patient and Public Involvement statement**

44
45 138 Patients or the public were not involved in the design, or conduct, or reporting, or
46
47 139 dissemination plans of our research.
48

49 140 **RESULTS**

50 51 141 **Study selection and description**

52
53 142 A total of 5,406 studies were identified according to the specified selection criteria from
54
55 143 Medline and Embase. After the removal of duplicates, 5,233 were screened by title/abstract,
56
57 144 out of which 333 were assessed via full text, and 15 studies subsequently included in this
58
59 145 review. The PRISMA flowchart showing the flow of study selection is presented in **Figure 1**.
60

1
2
3 146 Fifteen published studies were identified to address child-to-child and/or child-to-adult
4
5 147 transmission of SARS-CoV-2. Timeframes of data collection within these studies ranged
6
7 148 between January 2020 and January 2021. Studies from 11 countries were included (United
8
9 149 States, South Korea, Israel, Germany, Italy, Ireland, France, Singapore, Australia, Norway, and
10
11 150 England). A full detailed overview of the published studies is provided in **Table 1**.

13 151 ***Studies assessing outbreaks in Educational Settings***

15 152 Heavey et al. [14] conducted a case study in order to explore the role of transmission among
16
17 153 children in the school setting in the Republic of Ireland, before school closure. Three pediatric
18
19 154 index cases of COVID-19 with a history of school attendance were detected with 895 contacts.
20
21 155 Child-to-adult transmission or child-to-child transmission was not reported in this study.
22
23 156 Similarly Danis et al. [15] presented the contact tracing results of a nine-year-old child in
24
25 157 France, who visited 3 different schools the first days of symptom appearance. There was no
26
27 158 evidence of secondary transmission in any of the school contacts. Moreover, Yung et al.
28
29 159 traced three COVID-19 cases (2 pediatric and 1 adult) in three different educational settings,
30
31 160 and the results were negative, as were the tracing of close contacts of a preschool case in S
32
33 161 Korea [16]. Gold et al, in early 2020 had also indicated the possibility of educators playing a
34
35 162 role in school transmission as identified through the assessment of a transmission cluster in
36
37 163 primary (elementary) schools in the US [17] while Lopez et al assessed three COVID-19
38
39 164 outbreaks in child care facilities in Utah, during April 1–July 10, 2020 and noted that SARS-
40
41 165 CoV-2 Infections among young children acquired in child care settings were transmitted to
42
43 166 their household members [18].

44 167 One study from New South Wales, Australia presented an overview of COVID-19 cases and
45
46 168 transmission in schools. In a total number of 15 schools and 10 Early Childhood Educational
47
48 169 and Care Settings, 27 index cases were identified, among which 12 were children and 15 staff
49
50 170 members. Secondary transmission was noted only in four out of 25 educational settings,
51
52 171 where 2 children and 1 adult secondary cases were detected after the tracing of 752 contacts
53
54 172 [19].

55 173 ***Studies assessing the re-opening of schools and summer camps***

57 174 Eight studies reported on the regional evidence after the re-opening of schools. A school
58
59 175 outbreak in Israel after reopening of schools in May 2020 was described by Stein-Zamir et al.

1
2
3 176 The outbreak assessment was initiated by two pediatric COVID-19 cases that were not
4
5 177 epidemiologically related. The results showed that 153/1161 students and 25/151 staff
6
7 178 members tested positive for COVID-19. However, this outbreak was attributed to crowded
8
9 179 classes, combined with the exemption of facemasks and the use of air-condition due to an
10
11 180 extreme heatwave [20]. On the contrary, a study by Link-Gelles et al., in Rhode Island, USA.
12
13 181 among 666 child care programs that reopened on 1 June, 2020 after a 3-month closure
14
15 182 revealed 52 confirmed and probable cases (33 confirmed cases), of which 30 were among
16
17 183 children and 22 among adults. Secondary transmission for 10 cases was noted in only 4/666
18
19 184 childcare programs, which was attributed to class distancing, the use of face masks for adults,
20
21 185 universal symptom screening daily and disinfection [21]. The regional reopening of schools in
22
23 186 Germany in May 2020 was assessed by Ehrhardt et al., who noted that child-to-child
24
25 187 transmission in schools/childcare facilities appeared very uncommon, with an estimated six
26
27 188 of the identified 137 cases that had attended school to have led to a secondary transmission
28
29 189 overall to 11 additional pupils [22]. While two additional studies from S Korea by Yoon et al.,
30
31 190 indicated that upon the return of children to school in May-June 2020, no indication of
32
33 191 secondary transmission was noted in kindergarten children, middle school or high schools,
34
35 192 while in primary school only two cases of secondary transmission was noted [23, 24]. The
36
37 193 reopening of schools in September 2020 in Italy was not associated with elevated SAR, which
38
39 194 reached 3.8% overall, 0% in preschool, 0.38% in primary and 6.46% in secondary schools,
40
41 195 however these percentages included both adult and child cases [25]. Brandal et al., assessed
42
43 196 the transmission of COVID-19 in school settings in Norway between August-November 2020
44
45 197 and identified minimal child-to-child (0.9%, 2/234) and child-to-adult (1.7%, 1/58)
46
47 198 transmission [26].

48
49 199 Summer educational camps are presented separately, as close proximity between students is
50
51 200 not only noted within school hours but throughout the day and night due to additional extra
52
53 201 curriculum activities and close sleeping proximity. Two studies assessed secondary
54
55 202 transmission within summer educational camps, with striking differences. Pray et al identified
56
57 203 a rapid transmission of SARS-CoV-2 at an overnight retreat where adolescents and young
58
59 204 adults aged 14–24 years had prolonged contact and shared sleeping quarters, where one
60
205 index case/child led to the infection of 76% of attendees [27]. On the contrary Blaisdell in four
206
overnight camps noted no indication of secondary transmission following the isolation of the

207 paediatric index case and quarantine of their cohort, indicating the importance of the
208 implementation of NPI to reduce COVID-19 transmission [28].

209 ***Secondary attack rates of COVID-19 transmission in educational settings***

210 **Table 2** presents the SAR extracted from the studies, ranging from 0 to 76%, depending on
211 the setting, the timeframe and the implementation of NPI. With the exception of the study
212 by Pray et al., [27] within the context of summer camps in which a high transmission rate
213 (76%) was noted, in all studies within the context of school settings, the reported SARs were
214 minimal. Age differentiations were noted, for instance in the study by Larosa et al., across 36
215 schools in northern Italy, who identified an overall SAR of 3.2%, reaching 6.6% in middle and
216 high schools and 0.38% in primary schools.

217 **DISCUSSION**

218 This study provides a rapid review of the peer-reviewed literature pertaining to SARS-CoV-2
219 transmission by children within educational settings, a topic which is a crucial input to
220 assessments of the role of school settings in COVID-19 transmission. The literature appraised
221 in this review provides sufficient evidence that children can both be infected by and transmit
222 SARS-CoV-2 in school settings, however the SAR remained relatively low within the studies
223 assessed by our review, reflecting primarily schools in 2020. Our results with regards to
224 educational settings are in line with population based studies published after the cut-off of
225 this review, in which SARS-CoV-2 outbreaks were uncommon in educational settings [29] in
226 England [30], Canada [31] and in Utah, [32], Missouri [33] and New Jersey, USA [34], during
227 similar periods.

228 During the first waves of the COVID-19 pandemic, the vast uncertainty surrounding the
229 epidemiology of SARS-CoV-2 led many countries globally to include school closure
230 concomitant with other NPIs for reducing COVID-19 transmission. Within our review there
231 were limited cases in the assessed studies in which a child index case was responsible for
232 extensive secondary transmission in schools, with the notable exception of an outbreak in
233 Israel (which was associated with dense spacing, lack of the use of facemasks and closed
234 spaces with poor ventilation) and secondary transmission within summer educational camps,
235 where prolonged exposure between case-contact pairs is expected [27]. This finding is
236 supported by data from a large population based study assessing transmission dynamics that

1
2
3 237 identified that patterns of enhanced transmission risk in similar age pairs were strongest
4
5 238 among children aged 0 to 14 years [2].
6

7
8 239 On the contrary, evidence from studies that note a very small number of cases after school
9
10 240 reopenings the authors attribute to the strict NPIs implemented including the use of face
11
12 241 masks, physical distancing, screening for symptoms and classroom disinfection. Close
13
14 242 proximity between students was linked to elevated transmission rates in both school settings
15
16 243 and educational camps [20, 27], while adult educators have also been noted to play a role in
17
18 244 school transmission [17].

19 245 Modelling studies using various assumptions of infectivity from the first 3-4 months of the
20
21 246 pandemic [35-41], have previously assessed the role of school closure, and overall indicated
22
23 247 that school closure is associated with a reduction in the number of cases, hospitalisations and
24
25 248 ICU admissions, with the effect of school closure dependent on the transmission rate, and the
26
27 249 duration of school closure. Within this context age is noted to be a crucial aspect, as recent
28
29 250 modelling studies from the Netherlands indicated that contact restrictions within the age
30
31 251 group of 10-20 years old caused a slightly more significant reduction in R_e compared to 5-10
32
33 252 years old [54]. The same study also assessed the impact of reducing school contacts in
34
35 253 pandemic progression and showed that if complete school closure were implemented after
36
37 254 the summer holidays, R would be reduced by 10%, however, if school closure was enacted in
38
39 255 November, after implementing a partial lockdown since August, it could further decrease R
40
41 256 by 16%. Another recent European study that assessed school closure, based on the population
42
43 257 of two large cities of Norway, Oslo and Tromsø, indicating that a controlled and gradual school
44
45 258 re-opening would only have a slight increase in the reproduction number of less than 0.25,
46
47 259 and probably in the range between 0.10 and 0.14, which would not substantially affect the
48
49 260 infection rates [55].

50
51 261 While school closure may reduce SARS-CoV-2 transmission as noted above, the societal and
52
53 262 economic impacts of prolonged school closure are noteworthy, as they may impact the
54
55 263 availability of the healthcare workforce [37, 42] and may also have negative effects on
56
57 264 children through the interruption of the educational learning, social isolation, increased
58
59 265 exposure to domestic violence, and rise in dropout rates [43]. Furthermore, the impact of
60
266 school closures has been noted to impact significantly also special education [44], while
267
research performed within the context of the COVID-19 pandemic has identified that

268 contextual factors of particular relevance during school closures had negative impacts on
269 student wellbeing [45]. In light of the above, policy makers need to be aware of the
270 cost/benefit in each setting when considering school closures as a NPI [46].

271 Transmission of SARS-COV-2 has been noted to be higher in household settings than other
272 community settings, including schools, a finding which may be potentially attributable to the
273 individual, behavioural and contextual factors of the household vs. other settings, which may
274 support transmission dynamics [47]. Direct evidence showing children as a source of
275 transmission is scarce and largely based on small studies or studies investigating few
276 paediatric cases, however the results presented here concur with other and previous
277 systematic reviews that have summarised the evidence on the role of children in SARS-CoV-2
278 transmission [48-50].

279 There are important limitations to this study that may impact the direct implications for
280 decision-making. As we assessed peer-reviewed evidence published in two biomedical
281 databases, it inherently reflects the status quo of the interim of the previous school years
282 (January 2020 - January 2021) due to the lag time between study implementation, peer
283 review and publication. A further limitation of this report refers to the fact that these studies
284 represent child-to-child transmission within the context of previous SARS-CoV-2 strains and
285 are not directly applicable to newer and more transmissible variants, such as the SARS-CoV-2
286 Delta (B.1.617.2) variant of concern. Finally, the included studies reflect a broad geographical
287 and temporal range and are limited in comparability due to varying factors such as:
288 background levels of community SARS-CoV-2 transmission; enrolment strategies and varying
289 NPI policies which in turn depends highly on the geographical region and the socioeconomic
290 context, while accountability to government and political stability were found to exert
291 influence [51]. Hence in light of the above, supporting educators and parents in the
292 implementation of NPIs is important as population based studies have indicated that adults
293 concerned about the impact of COVID-19 on their children's education were more likely to
294 practice personal protective measures and social distancing [52].

295 **CONCLUSIONS**

296 The findings presented here provide an assessment of the published peer-reviewed evidence
297 on transmission in educational settings during 2020, in which transmission was minimal -

1
2
3 298 when NPI measures were implemented in parallel. However, with an upsurge of cases related
4
5 299 to new variants of concern, notably Delta, continuous surveillance and assessment of the
6
7 300 evidence is warranted to ensure the maximum protection of the health of students and the
8
9 301 educational workforce, while also minimising the numerous negative impacts that school
10
11 302 closures may have on children. Where schools remain open, in-school NPI measures should
12
13 303 be continually refined according to new knowledge according to the epidemiologic context,
14
15 304 taking into account levels of community SARS-CoV-2 transmission, information on the
16
17 305 severity of the Delta variant, and vaccination coverage levels among eligible students, which
18
19 306 includes children over 12 in many jurisdictions, at the time of writing [53].
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

307 **Acknowledgments**

308 We would like to thank Katerina Papathanasaki, Chrysa Chatzopoulou, Konstantinos
309 Skouloudakis and Ioanna Lagou for their assistance in data archiving and report preparation.
310 We acknowledge the bravery and dedication of educational professionals everywhere during
311 these most challenging times.

312

313 **Funding**

314 This report was produced under a service contract 6-ECD.11297, within Framework contract
315 ECDC/2019/001 Lot 1B, with the European Centre for Disease Prevention and Control (ECDC),
316 acting under the mandate from the European Commission. The information and views set out
317 in this piece of work are those of the authors and do not necessarily reflect the official opinion
318 of the Commission/Agency. The Commission/Agency do not guarantee the accuracy of the
319 data included in this analysis. Neither the Commission/Agency nor any person acting on the
320 Commission's/Agency's behalf may be held responsible for the use which may be made of the
321 information contained therein.

322 **Conflicts of interest/Competing interests**

323 None to report.

324 **Availability of data and material**

325 Not applicable

326 **Ethics statement**

327 For the purposes of this review publicly accessible documents were used as evidence, and,
328 hence, no ethics approval was required.

329 **Contributors**

330 CV, JL-B, RP and JES designed the study. KN, MHB and AM undertook the literature review
331 and extracted the data. JL-B and RP developed the search code. KN, MHB and AM analysed
332 and interpreted the data. AP and CD participated in data evaluation and interpretation along
333 with CV, JL-B, RP, JES, KN, MHB, and AM. CV wrote the first draft of the manuscript with input
334 from all authors. All authors reviewed and revised subsequent drafts.

TABLES

Table 1. Studies assessing SARS-CoV-2 transmission in educational settings, reported secondary cases and parallel non pharmaceutical interventions.

Lead Author, Year	Country	Timeframe	Age Range*	Setting	No of symptomatic pediatric index cases	No of asymptomatic pediatric index cases	Secondary cases in the school settings ¹	Parallel non pharmaceutical interventions
Child care settings								
Lopez et al., 2020 (18)	USA, Utah	April –July 2020	0.2-16	3 childcare facilities	0 child (3 adults)		Transmission was documented from 12 secondary pediatric cases (3 asymptomatic) to at least 12/46 non-facility contacts (confirmed or probable cases)	Quarantine for 14 days of cases + contacts; in 2 facilities: daily screening and staff members were using masks
Yoon et al., 2021 (24)	South Korea	February – March 2020	4	1 childcare center	1 (information about symptoms not reported)		0/190	Adult staff wore masks, but mask wearing by children were not consistent. After the index case-patient was identified, the center was closed. All potentially exposed persons were quarantined at home for 14 days.
Combined childcare-school settings								
Heavey et al., 2020 (14)	Ireland	March 2020	10-15	Schools	2	1	0/822 school contacts 0/73 other contacts	Exposure before school closure. Schools closed, contacts were quarantined
Danis et al., 2020 (15)	France	January to February 2020	9	3 schools	1	0	0/86 school contacts	Not reported

							1/6 hospitalised contacts	
Yung et al., 2020 (54)	Singapore	February to March 2020	2-8-15	3 schools	2	0	0/42 symptomatic contacts	Contacts were quarantined. Targeted measures at the school level
Macartney et al., 2020 (19)	Australia, NSW	25 January to 10 April 2020	<18	15 schools and 10 childcare settings	12 (information about symptoms not reported)		3/752 (3: 2 children & 1 adult)	Contacts were quarantined
Stein-Zamir et al., 2020 (20)	Israel	May 2020	12-18	1 high school	2	0	178/1312 (178 children & 25 staff)	Closed spaces with poor ventilation, high temperatures, crowded spaces and close contact with no masks
Link-Gelles et al., 2020 (21)	USA, RI	June – July 2020	<18	666 educational settings	33 confirmed and 19 probable cases in 29 settings		17 cases in 4/666 educational settings with.	Class distancing, the use of face masks for adults, universal symptom screening daily and disinfection
Ehrhardt et al., 2020 (22)	Germany, Baden-Württemberg	May – August 2020	<18	Schools and childcare facilities	137 (information about symptoms not reported)		11/>2300, estimation of 1 secondary case per roughly 25 infectious school days	Masks, social distancing, hygiene, ventilation, smaller class sizes, cancelled activities, exclusion of sick children
Brandal et al., 2020 (26)	Norway, Oslo and Viken counties	August – November 2020	5-13	Primary schools	13 (information about symptoms not reported)		3/292 (3: 2 children & 1 adult)	National guideline-based infection control measures, i.e. hygiene, physical distancing, symptomatic children to stay at home. Masks not worn in schools
Gold et al., 2021 (17)	USA, Georgia	December 2020 – January 2021	5-13	8 primary schools	1 (information about symptoms not reported)		5/contacts traced not reported	Physical distancing and masks; imperfect compliance noted

2021-08-30 08:08 on 5 April 2022. Downloaded from <http://bmjopen.bmj.com/> on April 22, 2024 by guest. Protected by copyright.

Larosa et al., 2020 (25)	Italy, Reggio Emilia	September – October 2020	<18	8 preschools, 10 primary 18 secondary	43	0	17/1,198 (17 children & 0 adults)	Mandatory surgical masks for children except when seated and not speaking; physical distancing measures
Yoon et al., 2020 (23)	South Korea	Up to July 2020	<18	6 preschools 13 primary, 6 secondary, 14 high schools	44 (information about symptoms not reported)		2/≥ 13,100	School closure continued until 6/4/2020. Social distancing strategies and mask wearing when schools opened with rigorous contact tracing and rapid testing on any suspected cases.
Summer Camps								
Pray et al., 2020 (27)	USA, Wisconsin	July – August 2020	14-24	1 overnight camp	1	0	115/151 confirmed or probable cases	Documentation of a negative prearrival RT-PCR result, 7-day prearrival quarantine, and outdoor programming
Blaisdell et al., 2020 (28)	USA, Maine	June – August 2020	7-18	4 overnight camps	0	1	No secondary transmission identified	Prearrival quarantine, pre- and postarrival testing and symptom screening, cohorting, use of face coverings, physical distancing, enhanced hygiene measures, cleaning and disinfecting, and maximal outdoor programming

1: Measured from the date of last contact; 2: Probable cases; *Except when the age refers to only 1 pediatric case and age range is 1-14

Table 2. Studies that assessed the secondary attack rate (SAR), when children are the index case within educational settings.

Lead Author, Year	Country	Timeframe	SAR
Heavey et al., 2020 (14)	Ireland	March 2020	0

Danis et al., 2020 (15)	France	January to February 2020	School: 0/86, Community: 0/80, Hospitalised: 1/6
Yung et al., 2020 (54)	Singapore	February to March 2020	0/42
Macartney et al., 2020 (19)	Australia, NSW	25 January to 9 April, 2020	All settings, all child case to child contacts 0.3% (2/649) All settings, all child case to staff member contacts 1.0% (1/103), Child close contacts 28.0% (7/25)
Stein-Zamir et al., 2020 (20)	Israel	May 2020	178 / 1312
Heavey et al., 2020 (14)	USA, Rhode Island	1 June- 31 July, 2020	n/a
Pray et al., 2020 (27)	United States, Wisconsin	July-August 2020	115/151 (76%)
Blaisdell et al., 2020 (28)	United States, Maine	June-August 2020	0
Lopez et al., 2020 (18)	USA, Utah	April-July 2020	n/a
Ehrhardt et al., 2020 (22)	Germany, Baden-Württemberg	25 May - 5 August 2020	estimation of one secondary case per roughly 25 infectious school days
Brandal et al., 2020 (26)	Norway, Oslo and Viken counties	28 August - 11 November 2020	child 2/234 (0.9%), adult 1/58 (1.7%)
Gold et al., 2021 (17)	United States, Georgia	1 December 2020 - 22 January 2021	n/a
Larosa et al., 2020 (25)	Italy	1 September -15 October 2020	38/994 (3.82%) overall 0.38% in primary schools (1/266) 6.46% in secondary schools (37/572)
Yoon et al., 2021 (24)	South Korea	27/2-16/3/2020	0
Yoon et al., 2020 (23)	Korea	up to 31/7/2020	2/≥ 13,100

6/bmjopen-2021-058308 on 5 April 2022. Downloaded from <http://bmjopen.bmj.com/> on April 22, 2024 by guest. Protected by copyright.

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURES

Figure 1. PRISMA Flowchart of study selection included in the rapid review.

For peer review only

References (imported via Endnote)


1. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, et al. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults. *JAMA Pediatrics*. 2021;175(2):143.
2. Laxminarayan R, Wahl B, Dudala SR, Gopal K, Mohan BC, Neelima S, et al. Epidemiology and transmission dynamics of COVID-19 in two Indian states. *Science*. 2020/11/07 ed2020. p. 691-7.
3. Jones TC, Mühlemann B, Veith T, Biele G, Zuchowski M, Hofmann J, et al. An analysis of SARS-CoV-2 viral load by patient age. medRxiv2020. p. 2020.06.08.20125484.
4. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). *JAMA Pediatr*. 2020;174(9):902-3.
5. Götzinger F, Santiago-García B, Noguera-Julián A, Lanasa M, Lancella L, Calò Carducci FI, et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Health*. 2020;4(9):653-61.
6. Dong QQ, Qiu LR, Cheng LM, Shu SN, Chen Y, Zhao Y, et al. Kindergartens Reopening in the Period of Regular Epidemic Prevention and Control, Beneficial or Harmful? *Curr Med Sci*. 2020;40(5):817-21.
7. Milani GP, Bottino I, Rocchi A, Marchisio P, Elli S, Agostoni C, et al. Frequency of Children vs Adults Carrying Severe Acute Respiratory Syndrome Coronavirus 2 Asymptomatically. *JAMA Pediatr*. 2021;175(2):193-4.
8. Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr*. 2020;109(6):1088-95.
9. Goldstein E, Lipsitch M. Temporal rise in the proportion of younger adults and older adolescents among coronavirus disease (COVID-19) cases following the introduction of physical distancing measures, Germany, March to April 2020. *Euro Surveill*. 2020;25(17).
10. Cho EY, Choi EH, Kim JH. Interpreting Transmissibility of COVID-19 in Children. *Emerg Infect Dis*. 2020;26(12):3106-7.
11. Control ECDC. COVID-19 in children and the role of school settings in transmission - second update. 8 July 2021.
12. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
13. Antonakou A. The latest update on the effects of COVID-19 infection in pregnancy. *Eur J Midwifery*. 2020;4:12.
14. Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary transmission of COVID-19 from children attending school in Ireland, 2020. *Eurosurveillance* 2020.
15. Danis K, Epaulard O, Bénet T, Gaymard A, Campoy S, Botelho-Nevers E, et al. Cluster of Coronavirus Disease 2019 (COVID-19) in the French Alps, February 2020. *Clinical Infectious Diseases* 2020. p. 825-32.
16. Yung CF, Kam K-Q, Nadua KD, Chong CY, Tan NWH, Li J, et al. Novel Coronavirus 2019 Transmission Risk in Educational Settings. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2021. p. 1055-8.
17. Gold JA GJ, Kimball A, et al. Infection Among Elementary School Educators and Students in One School District — Georgia, December 2020–January 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:289–2922021.

18. Lopez A, Hill M, Antezano J, Vilven D, Rutner T, Bogdanow L, et al. Transmission Dynamics of COVID-19 Outbreaks Associated with Child Care Facilities - Salt Lake City, Utah, April-July 2020. *MMWR Morbidity and mortality weekly report* 2020. p. 1319-23.
19. Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *The Lancet Child & Adolescent Health: Elsevier BV*; 2020. p. 807-16.
20. Stein-Zamir C, Abramson N, Shoob H, Libal E, Bitan M, Cardash T, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Eurosurveillance: European Centre for Disease Control and Prevention (ECDC)*; 2020.
21. Link-Gelles R, DellaGrotta AL, Molina C, Clyne A, Campagna K, Lanzieri TM, et al. Limited Secondary Transmission of SARS-CoV-2 in Child Care Programs - Rhode Island, June 1-July 31, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(34):1170-2.
22. Ehrhardt J, Ekinci A, Krehl H, Meincke M, Finci I, Klein J, et al. Transmission of SARS-CoV-2 in children aged 0 to 19 years in childcare facilities and schools after their reopening in May 2020, Baden-Wurttemberg, Germany. *Euro Surveill.* 2020/09/12 ed2020.
23. Yoon Y, Kim KR, Park H, Kim S, Kim YJ. Stepwise School Opening and an Impact on the Epidemiology of COVID-19 in the Children. *J Korean Med Sci.* 2020/12/02 ed2020. p. e414.
24. Yoon Y, Choi GJ, Kim JY, Kim KR, Park H, Chun JK, et al. Childcare Exposure to Severe Acute Respiratory Syndrome Coronavirus 2 for 4-Year-Old Presymptomatic Child, South Korea. *Emerg Infect Dis.* 2020/12/01 ed2021. p. 341-7.
25. Larosa E, Djuric O, Cassinadri M, Cilloni S, Bisaccia E, Vicentini M, et al. Secondary transmission of COVID-19 in preschool and school settings in northern Italy after their reopening in September 2020: a population-based study. *Euro Surveill.* 2020/12/12 ed2020.
26. Brandal LT, Ofitserova TS, Meijerink H, Rykkvin R, Lund HM, Hungnes O, et al. Minimal transmission of SARS-CoV-2 from paediatric COVID-19 cases in primary schools, Norway, August to November 2020. *Euro Surveill.* 2021/01/09 ed2021.
27. Pray I, Gibbons-Burgener S, Rosenberg A, Cole D, Borenstein S, Bateman A, et al. COVID-19 Outbreak at an Overnight Summer School Retreat — Wisconsin, July–August 2020. *MMWR Morbidity and Mortality Weekly Report* 2020. p. 1600-4.
28. Blaisdell L, Cohn W, Pavell J, Rubin D, Vergales J. Preventing and mitigating SARS-CoV-2 transmission - Four Overnight Camps, Maine, June-August 2020. *MMWR Morbidity and mortality weekly report* 2020. p. 1216-20.
29. Ladhani SN, Baawuah F, Beckmann J, Okike IO, Ahmad S, Garstang J, et al. SARS-CoV-2 infection and transmission in primary schools in England in June-December, 2020 (sKIDs): an active, prospective surveillance study. *Lancet Child Adolesc Health.* 2021;5(6):417-27.
30. Ismail SA, Saliba V, Lopez Bernal J, Ramsay ME, Ladhani SN. SARS-CoV-2 infection and transmission in educational settings: a prospective, cross-sectional analysis of infection clusters and outbreaks in England. *Lancet Infect Dis.* 2021;21(3):344-53.
31. Bark D, Dhillon N, St-Jean M, Kinniburgh B, McKee G, Choi A. SARS-CoV-2 transmission in kindergarten to grade 12 schools in the Vancouver Coastal Health region: a descriptive epidemiologic study. *CMAJ Open.* 2021;9(3):E810-e7.
32. Hershov RB, Wu K, Lewis NM, Milne AT, Currie D, Smith AR, et al. Low SARS-CoV-2 Transmission in Elementary Schools - Salt Lake County, Utah, December 3, 2020-January 31, 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70(12):442-8.
33. Dawson P, Worrell MC, Malone S, Tinker SC, Fritz S, Maricque B, et al. Pilot Investigation of SARS-CoV-2 Secondary Transmission in Kindergarten Through Grade 12 Schools Implementing Mitigation Strategies - St. Louis County and City of Springfield, Missouri, December 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70(12):449-55.

- 1
2
3 34. Volpp KG, Kraut BH, Ghosh S, Neatherlin J. Minimal SARS-CoV-2 Transmission After
4 Implementation of a Comprehensive Mitigation Strategy at a School - New Jersey, August 20-
5 November 27, 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70(11):377-81.
- 6 35. Abdollahi E, Haworth-Brockman M, Keynan Y, Langley JM, Moghadas SM. Simulating the
7 effect of school closure during COVID-19 outbreaks in Ontario, Canada. *BMC medicine.* 2020. p.
8 230.
- 9 36. Chin ET, Huynh BQ, Lo NC, Hastie T, Basu S. Projected geographic disparities in healthcare
10 worker absenteeism from COVID-19 school closures and the economic feasibility of child care
11 subsidies: a simulation study. *BMC medicine.* 2020. p. 218.
- 12 37. Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care
13 workforce and net mortality: a modelling study. *Lancet Public Health.* 2020;5(5):e271-e8.
- 14 38. Zhang J, Litvinova M, Liang Y, Wang Y, Wang W, Zhao S, et al. Changes in contact patterns
15 shape the dynamics of the COVID-19 outbreak in China. *Science (New York, NY)*2020. p. 1481-6.
- 16 39. Koo JR, Cook AR, Park M, Sun Y, Sun H, Lim JT, et al. Interventions to mitigate early spread
17 of SARS-CoV-2 in Singapore: a modelling study. *The Lancet Infectious diseases.* 2020. p. 678-88.
- 18 40. Kim S, Kim YJ, Peck KR, Jung E. School Opening Delay Effect on Transmission Dynamics of
19 Coronavirus Disease 2019 in Korea: Based on Mathematical Modeling and Simulation Study. *J*
20 *Korean Med Sci.* 2020/04/04 ed2020. p. e143.
- 21 41. Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al. The effect of control
22 strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a
23 modelling study. *The Lancet Public Health.* 2020. p. e261-e70.
- 24 42. Sadique MZ, Adams EJ, Edmunds WJ. Estimating the costs of school closure for mitigating
25 an influenza pandemic. *BMC Public Health.* 2008;8:135.
- 26 43. Adverse consequences of school closures: UNESCO; [Available from:
27 <https://en.unesco.org/covid19/educationresponse/consequences>.
28
- 29 44. Yakut AD. Educators' experiences in special education institutions during the COVID-19
30 outbreak. *J Res Spec Educ Needs.* 2021.
- 31 45. Mansfield KL, Newby D, Sonesson E, Vaci N, Jindra C, Geulayov G, et al. COVID-19 partial
32 school closures and mental health problems: A cross-sectional survey of 11,000 adolescents to
33 determine those most at risk. *JCPP Adv.* 2021;1(2):e12021.
- 34 46. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and
35 management practices during coronavirus outbreaks including COVID-19: a rapid systematic
36 review. *Lancet Child Adolesc Health.* 2020;4(5):397-404.
- 37 47. Merckx J, Labrecque JA, Kaufman JS. Transmission of SARS-CoV-2 by Children. *Dtsch*
38 *Arztebl Int.* 2020;117(33-34):553-60.
- 39 48. Li X, Xu W, Dozier M, He Y, Kirolos A, Theodoratou E. The role of children in transmission
40 of SARS-CoV-2: A rapid review. *J Glob Health.* 2020;10(1):011101.
- 41 49. Suk JE, Vardavas C, Nikitara K, Phalkey R, Leonardi-Bee J, Pharris A, et al. The role of
42 children in the transmission chain of SARS-CoV-2: a systematic review and update of current
43 evidence. medRxiv. 2020:2020.11.06.20227264.
- 44 50. Wiedenmann M, Goutaki M, Keiser O, Stringhini S, Tanner M, Low N. The role of children
45 and adolescents in the SARS-CoV-2 pandemic: a rapid review. *Swiss Med Wkly.* 2021;151:w30058.
- 46 51. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, et al. Ranking the
47 effectiveness of worldwide COVID-19 government interventions. *Nat Hum Behav.*
48 2020;4(12):1303-12.
- 49 52. Vardavas CI, Odani S, Nikitara K, El Banhawi H, Kyriakos CN, Taylor L, et al. Perceptions and
50 practice of personal protective behaviors to prevent COVID-19 transmission in the G7 nations.
51 *Population Medicine.* 2020;2(June).
- 52
53
54
55
56
57
58
59
60

- 1
2
3 53. Control ECfDPa. Overview of the implementation of COVID-19 vaccination strategies and
4 vaccine deployment plans in the EU/EEA 23 September 2021.
5 54. Yung CF, Kam KQ, Chong CY, Nadua KD, Li J, Tan NWH, et al. Household Transmission of
6 Severe Acute Respiratory Syndrome Coronavirus 2 from Adults to Children. *J Pediatr.* 2020/07/08
7 ed2020. p. 249-51.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only



BMJ Open

Transmission of SARS-CoV-2 in educational settings in 2020: a review

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058308.R1
Article Type:	Original research
Date Submitted by the Author:	13-Jan-2022
Complete List of Authors:	Vardavas, Constantine; University of Crete; Harvard University, Department of Oral Health Policy and Epidemiology Nikitara, Katerina; University of Crete, Mathioudakis, Alexander; The University of Manchester, Immunity and Respiratory Medicine; Manchester Academic Health Science Centre Boon, Michelle; Glasgow Caledonian University, WISE Centre for Economic Justice Phalkey, Revati; Public Health England; University of Nottingham Leonardi-Bee, Jo; University of Nottingham, Division of Epidemiology and Public Health, School of Medicine Pharris, Anastasia; European Centre for Disease Prevention and Control Deogan, Charlotte; European Centre for Disease Prevention and Control Suk, Jonathan; European Centre for Disease Prevention and Control
Primary Subject Heading:	Infectious diseases
Secondary Subject Heading:	Public health, Epidemiology, Infectious diseases, Health policy, Global health
Keywords:	COVID-19, Public health < INFECTIOUS DISEASES, Community child health < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1 Transmission of SARS-CoV-2 in educational settings in 2020: a 2 review

3 **Authors:** Constantine Vardavas^{1,2}, Katerina Nikitara¹, Alexander G. Mathioudakis^{3,4}, Michele
4 Hilton-Boon⁵, Revati Phalkey^{6,7}, Jo Leonardi-Bee⁷, Anastasia Pharris⁸, Charlotte Deogan⁸,
5 Jonathan E. Suk^{8*}

7 Affiliations

8 1: School of Medicine, University of Crete, Greece

9 2: Department of Oral Health Policy and Epidemiology, Harvard School of Dental Medicine, Harvard
10 University, Boston, MA, USA

11 3: Division of Infection, Immunity and Respiratory Medicine, School of Biological Sciences, The
12 University of Manchester, Manchester, UK.

13 4: North West Lung Centre, Wythenshawe Hospital, Manchester University NHS Foundation Trust,
14 Manchester Academic Health Science Centre, Manchester, UK.

15 5: WISE Centre for Economic Justice, Glasgow Caledonian University

16 6: Climate Change and Health Group, Public Health England, United Kingdom

17 7: Division of Epidemiology and Public Health, School of Medicine, University of Nottingham, UK

18 8: European Centre for Disease Prevention and Control, Solna, Sweden

19
20
21 ***: Corresponding author: Jonathan E. Suk, jonathan.suk@ecdc.europa.eu, +46858601633. Gustav III:s
22 Boulevard, 169 73 Solna, Sweden.**

23 Word counting: 3216 words

24 Keywords: SARS-CoV-2, COVID-19, pandemic, children, schools, household

27 **Abstract (240 words)**

28 **Objectives:** School closures have been used as a core non pharmaceutical intervention (NPI)
29 during the COVID-19 pandemic. This review aims at identifying SARS-CoV-2 transmission in
30 educational settings during the first waves of the pandemic.

31 **Methods:** This literature review assessed studies published between December 2019 and
32 April 1, 2021 in Medline and Embase, which included studies that assessed educational
33 settings from approximately January 2020 to January 2021. The inclusion criteria were based
34 on the PCC framework (P-Population, C-Concept, C-Context). The study *Population* was
35 restricted to people 1-17 years old (excluding neonatal transmission), the *Concept* was to
36 assess child-to-child and child-to-adult transmission, while the *Context* was to assess
37 specifically educational setting transmission.

38 **Results:** Fifteen studies met inclusion criteria, ranging from daycare centers to high schools
39 and summer camps, while eight studies assessed the re-opening of schools in the 2020-2021
40 school year. In principle although there is sufficient evidence that children can both be
41 infected by and transmit SARS-CoV-2 in school settings, the SAR remain relatively low -when
42 NPI measures are implemented in parallel. Moreover, although the evidence was limited
43 there was an indication that younger children may have a lower SAR than adolescents.

44 **Conclusions:** Transmission in educational settings in 2020 was minimal -when NPI measures
45 were implemented in parallel. However, with an upsurge of cases related to variants of
46 concern, continuous surveillance and assessment of the evidence is warranted to ensure the
47 maximum protection of the health of students and the educational workforce, while also
48 minimising the numerous negative impacts that school closures may have on children.

49 **Strengths and limitations of this study**

- 50 • This review followed a systematic search approach.
- 51 • The included studies of this review have heterogenous methodologies and a meta-
52 analysis could not be performed.
- 53 • The search represents peer reviewed literature that included previous variants of interest
54 and do not cover the Delta or Omicron.

1
2
3 56 **MAIN TEXT**
4

5
6 57 **INTRODUCTION**
7

8 58 One of the more perplexing and controversial dimensions during the first year of the COVID-
9 59 19 pandemic surrounded the role of children in the transmission.

10
11
12 60 Epidemiologic indicators of SARS-CoV-2 infection in children provide a complex picture
13 61 regarding their potential role in the transmission chain. Systematic reviews have concluded
14 62 that children and adolescents have lower susceptibility to SARS-CoV-2 infection (1, 2).
15 63 However, when infected and symptomatic, children may shed viral RNA in similar quantities
16 64 to adults (3), and that younger children (under 5 years) with mild to moderate symptoms may
17 65 shed even more virus than older children and adults (4). While the proportion of
18 66 asymptomatic SARS-CoV-2 infections among children in the general population is uncertain,
19 67 initial data had indicated that 16% of paediatric cases in Europe in the first phase of the
20 68 pandemic were classified as asymptomatic (5), while up to 90% of paediatric cases in China
21 69 were deemed to be asymptomatic, mild, or moderate (6). Moreover, it is possible that
22 70 children are less often asymptomatic carriers than adults: a study of non-COVID-19-related
23 71 hospitalizations in Milan identified 1% of children and 9% of adults as asymptomatic carriers
24 72 of SARS-CoV-2 (7). Meanwhile, while children are overall noted to have lower rates of severe
25 73 COVID-19 cases (8), there is evidence of differing transmission dynamics between younger vs.
26 74 older children (2). There is evidence that when index cases, younger children, such as under
27 75 10 years of age, lead to lower secondary attack rates than older children and adult (9, 10).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 76 Important potential sources of evidence surrounding the role of children in the COVID-19
57 77 pandemic come from studies situated in the community, household, healthcare or
58 78 educational settings. Transmission of SARS-COV-2 has thus far been documented to be higher
59 79 in household settings than in other community settings – including schools – a finding which
60 80 may be potentially attributable to the individual, behavioural and contextual factors of
81 81 households vs. other settings, as has been suggested elsewhere (9).
82
83
84

83 Although, at the time of writing, the more transmissible Delta and potentially Omicron
84 84 variants is driving SARS-CoV-2 transmission there is currently a gap in published studies

1
2
3 85 looking at the transmission of COVID-19 during the first waves in school settings. However, as
4
5 86 decisions currently need to be taken to ensure high levels of preparedness in school settings
6
7 87 (11), the literature published thus far may have important insights to guide decision-making
8
9 88 around school closures and re-openings, as well support decision making for mitigation
10
11 89 measures in educational settings. With the above in mind this literature review was
12
13 90 conducted to assess child-to-child and child-to-adult SARS-CoV-2 transmission within
14
15 91 educational settings during the first wave of the pandemic and to calculate where possible
16
17 92 the secondary attack rate (SAR) when the child is the index case.

18 93 **METHODS**

19 94 ***Search Strategy***

20
21
22 95 This literature review is reported in accordance with the Preferred Reporting Items for
23
24 96 Systematic Reviews and Meta-Analysis (PRISMA) guidelines (12). Relevant studies published
25
26 97 between December 2019 and April 1, 2021 were identified by searching Medline and Embase.
27
28 98 The following set of inclusion criteria were used to determine eligibility of the studies, which
29
30 99 is based on the PCC framework (P-Population, C-Concept, C-Context). The study *Population*
31
32 100 was restricted to people 1-17 years old (excluding neonatal transmission (13)), the *Concept*
33
34 101 was to assess child-to-child and child-to-adult transmission when the child is the index case,
35
36 102 while the *Context* was to assess specifically educational setting transmission clusters. Subject
37
38 103 heading terms and free text words relating to the Population, Concept and Context terms as
39
40 104 identified in the inclusion criteria were used to develop a comprehensive list of terms for the
41
42 105 search strategy (so as to ensure we would not lose information), from which this specific
43
44 106 review on educational settings was based. We included all studies of quantitative research,
45
46 107 while, opinion pieces, commentaries, case reports and editorials were excluded.
47
48 108 Mathematical modelling and simulation studies were also excluded. We additionally screened
49
50 109 reference lists of the included articles to identify further relevant studies. The search was
51
52 110 limited to the English language. The search terms of the review are presented in the
53
54 111 Supplementary file.

55 112 56 113 ***Study selection***

57 114 Initially, a pilot training screening process was used where 100 identical articles were
58
59 115 screened for their eligibility independently by two reviewers to ensure consistency in
60

1
2
3 116 screening. As a high measure of inter-rater agreement was achieved between the two
4
5 117 reviewers during the pilot assessment (percentage agreement >90% and/ or Cohen's Kappa
6
7 118 >0.81), the remaining titles were randomly allocated to the two reviewers and screened for
8
9 119 eligibility independently by them. After an initial selection of the titles, each reviewer
10
11 120 assessed each other's selected studies. The retrieved articles were then independently
12
13 121 double-screened by two reviewers based on the full text of the articles.

15 122 ***Data extraction***

17 123 The data extraction template was piloted independently by the two reviewers on a random
18
19 124 sample of two included studies to enable an assessment of consistency in data extraction and
20
21 125 to identify where amendments needed to be made to the template. The remaining studies
22
23 126 were then data extracted independently by two reviewers, and the results were double
24
25 127 checked across the original manuscript by a third reviewer.

27 128 ***Data synthesis***

29 129 Characteristics of the included studies were presented in tabulated form detailing the study
30
31 130 design, geographical location of the study, sample size, characteristics of the populations
32
33 131 considered, setting, context, parallel implemented Non Pharmaceutical Interventions (NPI),
34
35 132 and the findings of the study. Depending on the level of information available, infection SAR
36
37 133 were noted as defined in each included study. A narrative synthesis approach was applied to
38
39 134 look systematically at the data and to describe each study categorized by the study design.
40
41 135 Patterns in the data were identified through tabulation of results, and an inductive approach
42
43 136 was taken to translate the data to identify areas of commonality between studies.

44 137 ***Patient and Public Involvement statement***

46 138 Patients or the public were not involved in the design, or conduct, or reporting, or
47
48 139 dissemination plans of our research.

51 140 **RESULTS**

53 141 ***Study selection and description***

55 142 A total of 5,406 studies were identified according to the specified selection criteria from
56
57 143 Medline and Embase. After the removal of duplicates, 5,233 were screened by title/abstract,
58
59
60

1
2
3 144 out of which 333 were assessed via full text, and 15 studies subsequently included in this
4
5 145 review. The PRISMA flowchart showing the flow of study selection is presented in **Figure 1**.

6
7 146 Fifteen published studies were identified to report child-to-child and/or child-to-adult
8
9 147 transmission of SARS-CoV-2. Timeframes of data collection within these studies ranged
10
11 148 between January 2020 and January 2021. Studies from 11 countries were included (United
12
13 149 States, South Korea, Israel, Germany, Italy, Ireland, France, Singapore, Australia, Norway, and
14
15 150 England). A full detailed overview of the published studies is provided in **Table 1**.

17 151 ***Studies assessing outbreaks in Educational Settings***

18
19 152 Heavey et al. (14) conducted a case study in order to explore the role of transmission among
20
21 153 children in the school setting in the Republic of Ireland, before school closure. Three pediatric
22
23 154 index cases of COVID-19 with a history of school attendance were detected with 895 contacts.
24
25 155 Child-to-adult transmission or child-to-child transmission was not reported in this study.
26
27 156 Similarly Danis et al. (15) presented the contact tracing results of a nine-year-old child in
28
29 157 France, who visited 3 different schools the first days of symptom appearance. There was no
30
31 158 evidence of secondary transmission in any of the school contacts. Moreover, Yung et al.
32
33 159 traced three COVID-19 cases (2 pediatric and 1 adult) in three different educational settings,
34
35 160 and the results were negative, as were the tracing of close contacts of a preschool case in S
36
37 161 Korea (16). Gold et al, in early 2020 had also indicated the possibility of educators playing a
38
39 162 role in school transmission as identified through the assessment of transmission clusters in
40
41 163 primary (elementary) schools in Georgia, US. More specific, in four clusters the index case was
42
43 164 an educator, while a student was the index case in one cluster.(17). Also,Lopez et al assessed
44
45 165 three COVID-19 outbreaks in child care facilities in Utah, during April 1–July 10, 2020 and
46
47 166 noted that SARS-CoV-2 infections among young children acquired in child care settings were
48
49 167 transmitted to their household members (18).

50
51 168 One study from New South Wales, Australia presented an overview of COVID-19 cases and
52
53 169 transmission in schools. In a total number of 15 schools and 10 Early Childhood Educational
54
55 170 and Care Settings, 27 index cases were identified, among which 12 were children and 15 staff
56
57 171 members. Secondary transmission was noted only in four out of 25 educational settings, (19).

58 172 ***Studies assessing the re-opening of schools and summer camps***

1
2
3 173 Eight studies reported on the regional evidence after the re-opening of schools. A school
4 174 outbreak in Israel after reopening of schools in May 2020 was described by Stein-Zamir et al.
5 175 The outbreak assessment was initiated by two pediatric COVID-19 cases that were not
6 176 epidemiologically related. The results showed that 153/1161 students and 25/151 staff
7 177 members tested positive for COVID-19 (20). On the contrary, a study by Link-Gelles et al., in
8 178 Rhode Island, USA. among 666 child care programs revealed 52 confirmed and probable cases
9 179 (33 confirmed cases), of which 30 were among children and 22 among adults. Secondary
10 180 transmission for 10 cases was noted in only 4/666 childcare programs(21). The regional
11 181 reopening of schools in Germany in May 2020 was assessed by Ehrhardt et al., who noted that
12 182 child-to-child transmission in schools/childcare facilities appeared very uncommon, with an
13 183 estimated six of the identified 137 cases that had attended school to have led to a secondary
14 184 transmission overall to 11 additional pupils (22). While two additional studies from S Korea
15 185 by Yoon et al., indicated that upon the return of children to school in May-June 2020, no
16 186 indication of secondary transmission was noted in kindergarten children, middle school or
17 187 high schools, while in primary school only two cases of secondary transmission was noted (23,
18 188 24). The reopening of schools in September 2020 in Italy was not associated with elevated
19 189 SAR, which reached 3.8% overall, 0% in preschool, 0.38% in primary and 6.46% in secondary
20 190 schools, however these percentages included both adult and child cases (25). Brandal et al.,
21 191 assessed the transmission of COVID-19 in school settings in Norway between August-
22 192 November 2020 and identified minimal child-to-child (0.9%, 2/234) and child-to-adult (1.7%,
23 193 1/58) transmission (26).

24 194 Summer educational camps are presented separately, as close proximity between students is
25 195 not only noted within school hours but throughout the day and night due to additional extra
26 196 curriculum activities and close sleeping proximity. Two studies assessed secondary
27 197 transmission within summer educational camps, with striking differences. Pray et al identified
28 198 a rapid transmission of SARS-CoV-2 at an overnight retreat where adolescents and young
29 199 adults aged 14–24 years had prolonged contact and shared sleeping quarters, where one
30 200 index case/child led to the infection of 76% of attendees (27). On the contrary Blaisdell in four
31 201 overnight camps noted no indication of secondary transmission following the isolation of the
32 202 paediatric index case and quarantine of their cohort, indicating the importance of the
33 203 implementation of NPI to reduce COVID-19 transmission (28).

204 ***Secondary attack rates of COVID-19 transmission in educational settings***

205 **Table 2** presents the SAR extracted from the studies, ranging from 0 to 76%, depending on
206 the setting, the timeframe and the implementation of NPI. With the exception of the study
207 by Pray et al., (27) within the context of summer camps in which a high transmission rate
208 (76%) was noted, in all studies within the context of school settings, the reported SARs were
209 minimal. Age differentiations were noted, for instance in the study by Larosa et al., across 36
210 schools in northern Italy, who identified an overall SAR of 3.2%, reaching 6.6% in middle and
211 high schools and 0.38% in primary schools (25).

212 **DISCUSSION**

213 This study provides a rapid review of the peer-reviewed literature pertaining to SARS-CoV-2
214 transmission by children within educational settings, a topic which is a crucial input to
215 assessments of the role of school settings in COVID-19 transmission. The literature appraised
216 in this review provides sufficient evidence that children can both be infected by and transmit
217 SARS-CoV-2 in school settings, however the SAR remained relatively low within the studies
218 assessed by our review, reflecting primarily schools in 2020. Our results with regards to
219 educational settings are in line with population based studies published after the cut-off of
220 this review, in which SARS-CoV-2 outbreaks were uncommon in educational settings (29) in
221 England (30), Canada (31) and in Utah, (32), Missouri (33) and New Jersey, USA (34), North
222 Carolina (35) during similar periods.

223 During the first waves of the COVID-19 pandemic, the vast uncertainty surrounding the
224 epidemiology of SARS-CoV-2 led many countries globally to include school closure
225 concomitant with other NPIs for reducing COVID-19 transmission. Within our review there
226 were limited cases in the assessed studies in which a child index case was responsible for
227 extensive secondary transmission in schools, with the notable exception of an outbreak in
228 Israel (which was associated with dense spacing, lack of the use of facemasks and closed
229 spaces with poor ventilation) and secondary transmission within summer educational camps,
230 where prolonged exposure between case-contact pairs is expected (27). This finding is
231 supported by data from a large population-based study assessing transmission dynamics that
232 identified that patterns of enhanced transmission risk in similar age pairs were strongest
233 among children aged 0 to 14 years (2).

1
2
3 234 On the contrary, evidence from studies that note a very small number of cases after school
4
5 235 reopening (36, 37) the authors attribute to the strict NPIs implemented including the use of
6
7 236 face masks, physical distancing, screening for symptoms and classroom disinfection (21).
8
9 237 Close proximity between students was linked to elevated transmission rates in both school
10
11 238 settings and educational camps (20, 27), while adult educators have also been noted to play
12
13 239 a role in school transmission (17).

14
15 240 Modelling studies using various assumptions of infectivity from the first 3-4 months of the
16
17 241 pandemic (38-44), have previously assessed the role of school closure, and overall indicated
18
19 242 that school closure is associated with a reduction in the number of cases, hospitalisations and
20
21 243 ICU admissions, with the effect of school closure dependent on the transmission rate, and the
22
23 244 duration of school closure. Within this context, age is noted to be a crucial aspect, as recent
24
25 245 modelling studies from the Netherlands indicated that contact restrictions within the age
26
27 246 group of 10-20 years old caused a slightly more significant reduction in R_e compared to 5-10
28
29 247 years old [54]. Another European study that assessed school closure, based on the population
30
31 248 of two large cities of Norway, Oslo and Tromsø, indicated that a controlled and gradual school
32
33 249 re-opening would only have a slight increase in the reproduction number of less than 0.25,
34
35 250 and probably in the range between 0.10 and 0.14, which would not substantially affect the
36
37 251 infection rates [55]. Interestingly, a rapid review conducted by Viner et al (45), underlined
38
39 252 that while modelling studies support the closure of educational institutions as part of the
40
41 253 social distancing measures that need to be implemented, the only study examining school
42
43 254 closures exclusively found relatively marginal impact, by reasonably assuming increased levels
44
45 255 of household and community as a result. However, a recent review that included only
46
47 256 empirical studies, conducted by Mendez-Brito et al., (46) indicated that school closure,
48
49 257 followed by workplace and entertainment venue closure, as well as bans of public events
50
51 258 were the most effective NPIs, concluding that an early response and a combination of specific
52
53 259 social distancing measures are of crucial importance for the reduction of COVID-19 cases and
54
55 260 deaths.

56
57 261 While school closure may reduce SARS-CoV-2 transmission, the societal and economic
58
59 262 impacts of prolonged school closure are noteworthy, as they may impact the availability of
60
263 the healthcare workforce (40, 47) and may also have negative effects on children through the
264
interruption of the educational learning, social isolation, increased exposure to domestic

1
2
3 265 violence, and rise in dropout rates (48). Furthermore, the impact of school closures has been
4
5 266 noted to impact significantly also special education (49), while research performed within the
6
7 267 context of the COVID-19 pandemic has identified that contextual factors of particular
8
9 268 relevance during school closures had negative impacts on student wellbeing (50). In light of
10
11 269 the above, policy makers need to be aware of the cost/benefit in each setting when
12
13 270 considering school closures as a NPI (45).

14
15 271 Transmission of SARS-COV-2 has been noted to be higher in household settings than other
16
17 272 community settings, including schools, a finding which may be potentially attributable to the
18
19 273 individual, behavioural and contextual factors of the household vs. other settings, which may
20
21 274 support transmission dynamics (51). Direct evidence showing children as a source of
22
23 275 transmission is scarce and largely based on small studies or studies investigating few
24
25 276 paediatric cases, however the results presented here concur with other and previous
26
27 277 systematic reviews that have summarised the evidence on the role of children in SARS-CoV-2
28
29 278 transmission (52-54).

30
31 279 There are important limitations to this study that may impact the direct implications for
32
33 280 decision-making. As we assessed peer-reviewed evidence published in two biomedical
34
35 281 databases, it inherently reflects the status quo of the interim of the previous school years
36
37 282 (January 2020 - January 2021) due to the lag time between study implementation, peer
38
39 283 review and publication. A further limitation of this report refers to the fact that these studies
40
41 284 represent child-to-child transmission within the context of previous SARS-CoV-2 strains and
42
43 285 are not directly applicable to newer and more transmissible variants, such as the SARS-CoV-2
44
45 286 Delta (B.1.617.2) variant of concern or the more recent Omicron variant. Finally, the included
46
47 287 studies reflect a broad geographical and temporal range and are limited in comparability due
48
49 288 to varying factors such as: background levels of community SARS-CoV-2 transmission;
50
51 289 enrolment strategies and varying NPI policies which in turn depends highly on the
52
53 290 geographical region and the socioeconomic context, while accountability to government and
54
55 291 political stability were found to exert influence (55). Hence in light of the above, supporting
56
57 292 educators and parents in the implementation of NPIs is important as population based studies
58
59 293 have indicated that adults concerned about the impact of COVID-19 on their children's
60
294 education were more likely to practice personal protective measures and social distancing
295 (56).

1
2
3 296
4
56 297 **CONCLUSIONS**
7

8 298 The findings presented here provide an assessment of the published peer-reviewed evidence
9
10 299 on transmission in educational settings during 2020, in which transmission was minimal -
11
12 300 when NPI measures were implemented in parallel. However, with an upsurge of cases related
13
14 301 to new variants of concern, notably Delta and Omicron, continuous surveillance and
15
16 302 assessment of the evidence is warranted to ensure the maximum protection of the health of
17
18 303 students and the educational workforce, while also minimising the numerous negative
19
20 304 impacts that school closures may have on children. Where schools remain open, in-school NPI
21
22 305 measures should be continually refined according to new knowledge according to the
23
24 306 epidemiologic context, taking into account levels of community SARS-CoV-2 transmission,
25
26 307 information on the severity of the Delta and Omicron variants, and vaccination coverage
27
28 308 levels among eligible students, which includes children over 12 in many jurisdictions, at the
29
30 309 time of writing (57). Finally, future studies should focus more on identifying SARS-CoV-2
31
32 310 variants and on providing specific definitions about cases and contacts, while more detailed
33
34 311 information on the contact tracing strategies and the implemented NPIs would reduce the
35
36 312 limitations.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

313 **Acknowledgments**

314 We would like to thank Katerina Papathanasaki, Chrysa Chatzopoulou, Konstantinos
315 Skouloudakis and Ioanna Lagou for their assistance in data archiving and report preparation.
316 We acknowledge the bravery and dedication of educational professionals everywhere during
317 these most challenging times.

318 **Funding**

319 This report was produced under a service contract 6-ECD.11297, within Framework contract
320 ECDC/2019/001 Lot 1B, with the European Centre for Disease Prevention and Control (ECDC),
321 acting under the mandate from the European Commission. The information and views set out
322 in this piece of work are those of the authors and do not necessarily reflect the official opinion
323 of the Commission/Agency. The Commission/Agency do not guarantee the accuracy of the
324 data included in this analysis. Neither the Commission/Agency nor any person acting on the
325 Commission's/Agency's behalf may be held responsible for the use which may be made of the
326 information contained therein.

327 **Conflicts of interest/Competing interests**

328 None to report.

329 **Availability of data and material**

330 No data are available.

331 **Ethics statement**

332 For the purposes of this review publicly accessible documents were used as evidence, and,
333 hence, no ethics approval was required.

334 **Contributors**

335 CV, JL-B, RP and JES designed the study. KN, MHB and AM undertook the literature review
336 and extracted the data. JL-B and RP developed the search code. KN, MHB and AM analysed
337 and interpreted the data. AP and CD participated in data evaluation and interpretation along
338 with CV, JL-B, RP, JES, KN, MHB, and AM. CV wrote the first draft of the manuscript with input
339 from all authors. All authors reviewed and revised subsequent drafts.

TABLES

Table 1. Studies assessing SARS-CoV-2 transmission in educational settings, reported secondary cases and parallel non pharmaceutical interventions, until January 2021.

Lead Author, Year	Country	Timeframe	Age Range*	Setting	No of symptomatic pediatric index cases	No of asymptomatic pediatric index cases	Secondary cases in the school settings ¹	Parallel non pharmaceutical interventions in the community setting
Child care settings								
Lopez et al., 2020 (18)	USA, Utah	April –July 2020	0.2-16	3 childcare facilities (3 clusters)	0 child (3 adults)		Transmission was documented from 12 secondary pediatric cases (3 asymptomatic) to at least 12/46 non-facility contacts (confirmed or probable cases)	Quarantine for 14 days of cases + contacts; in 2 facilities: daily screening and staff members were using masks
Yoon et al., 2021 (24)	South Korea	February – March 2020	4	1 childcare center	1 (information about symptoms not reported)		0/190	Adult staff wore masks, but mask wearing by children were not consistent. After the index case-patient was identified, the center was closed. All potentially exposed persons were quarantined at home for 14 days.
Combined childcare-school settings								
Heavey et al., 2020 (14)	Ireland	March 2020	10-15	Schools	2	1	0/822 school contacts 0/73 other contacts	Exposure before school closure. Schools closed, contacts were quarantined
Danis et al., 2020 (15)	France	January to February 2020	9	3 schools	1	0	0/86 school contacts	Not reported

							1/6 hospitalised contacts	
Yung et al., 2020 (58)	Singapore	February to March 2020	2.8-15	3 schools	2	0	0/42 symptomatic contacts	Contacts were quarantined. Targeted measures at the school level
Macartney et al., 2020 (19)	Australia, NSW	25 January to 10 April 2020	<18	15 schools and 10 childcare settings (3 clusters)	12 (information about symptoms not reported)		3/752 (3: 2 children & 1 adult)	Contacts were quarantined
Stein-Zamir et al., 2020 (20)	Israel	May 2020	12-18	1 high school (1 cluster)	2	0	178/1312 (178: 153 children & 25 staff)	Closed spaces with poor ventilation, high temperatures, crowded spaces and close contact with no masks
Link-Gelles et al., 2020 (21)	USA, RI	June – July 2020	<18	666 educational settings (4 clusters)	33 confirmed and 19 probable cases in 29 settings		17 cases in 4/666 educational settings with.	Class distancing, the use of face masks for adults, universal symptom screening daily and disinfection
Ehrhardt et al., 2020 (22)	Germany, Baden-Württemberg	May – August 2020	<18	Schools and childcare facilities (11 clusters)	137 (information about symptoms not reported)		11/>2300, estimation of 1 secondary case per roughly 25 infectious school days	Masks, social distancing, hygiene, ventilation, smaller class sizes, cancelled activities, exclusion of sick children
Brandal et al., 2020 (26)	Norway, Oslo and Viken counties	August – November 2020	5-13	Primary schools (2 clusters)	13 (information about symptoms not reported)		3/292 (3: 2 children & 1 adult)	National guideline-based infection control measures, i.e. hygiene, physical distancing, symptomatic children to stay at home. Masks not worn in schools

http://bmjopen-2021-028308 on 5 April 2022. Downloaded from http://bmjopen.bmj.com/ on April 22, 2024 by guest. Protected by copyright.

Gold et al., 2021 (17)	USA, Georgia	December 2020 – January 2021	5-13	8 primary schools (nine clusters)	1 (information about symptoms not reported)		5/contacts traced not reported	Physical distancing and masks; imperfect compliance noted
Larosa et al., 2020 (25)	Italy, Reggio Emilia	September – October 2020	<18	8 preschools, 10 primary 18 secondary (nine clusters)	43	0	17/1198 (17 children & 0 adults)	Mandatory surgical masks for children except when seated and not speaking; physical distancing measures
Yoon et al., 2020 (23)	South Korea	Up to July 2020	<18	6 preschools 13 primary, 6 secondary, 14 high schools (2 clusters)	44 (information about symptoms not reported)		2/≥ 13,100	School closure continued until 6/4/2020. Social distancing strategies and mask wearing when schools opened with rigorous contact tracing and rapid testing on any suspected cases.
Summer Camps								
Pray et al., 2020 (27)	USA, Wisconsin	July – August 2020	14-24	1 overnight camp	1	0	115/151 confirmed or probable cases	Documentation of a negative prearrival RT-PCR result, 7-day prearrival quarantine, and outdoor programming
Blaisdell et al., 2020 (28)	USA, Maine	June – August 2020	7-18	4 overnight camps	0	1	No secondary transmission identified	Prearrival quarantine, pre- and postarrival testing and symptom screening, cohorting, use of face coverings, physical distancing, enhanced hygiene measures, cleaning and disinfecting, and maximal outdoor programming

1: Measured from the date of last contact; 2: Probable cases; *Except when the age refers to only 1 pediatric case and age range is a

Table 2. Studies that assessed the secondary attack rate (SAR)¹, when children are the index case within educational settings.

Lead Author, Year	Country	Timeframe	SAR
Heavey et al., 2020 (14)	Ireland	March 2020	0
Danis et al., 2020 (15)	France	January to February 2020	School: 0/86, Community: 0/80, Hospitalised: 1/6
Yung et al., 2020 (58)	Singapore	February to March 2020	0/42
Macartney et al., 2020 (19)	Australia, NSW	25 January to 9 April, 2020	All settings, all child case to child contacts 0.3% (2/649) All settings, all child case to staff member contacts 1.0% (1/103), Child close contacts 28.0% (7/25)
Stein-Zamir et al., 2020 (20)	Israel	May 2020	178 / 1312
Heavey et al., 2020 (14)	USA, Rhode Island	1 June- 31 July, 2020	n/a
Pray et al., 2020 (27)	United States, Wisconsin	July-August 2020	115/151 (76%)
Blaisdell et al., 2020 (28)	United States, Maine	June-August 2020	0
Lopez et al., 2020 (18)	USA, Utah	April-July 2020	n/a
Ehrhardt et al., 2020 (22)	Germany, Baden-Württemberg	25 May - 5 August 2020	estimation of one secondary case per roughly 25 infectious school days
Brandal et al., 2020 (26)	Norway, Oslo and Viken counties	28 August - 11 November 2020	child 2/234 (0.9%), adult 1/58 (1.7%)
Gold et al., 2021 (17)	United States, Georgia	1 December 2020 - 22 January 2021	n/a
Larosa et al., 2020 (25)	Italy	1 September -15 October 2020	38/994 (3.82%) overall 0.38% in primary schools (1/266) 6.46% in secondary schools (37/572)
Yoon et al., 2021 (24)	South Korea	27/2-16/3/2020	0

66/bmjopen-2021-023308 on 5 April 2022. Downloaded from <http://bmjopen.bmj.com/> by guest. Protected by copyright.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Yoon et al., 2020 (23)	Korea	up to 31/7/2020	2/≥ 13,100
------------------------	-------	-----------------	------------

1: The extracted the SAR based on the original definition given by the authors in each study

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURES

Figure 1. PRISMA Flowchart of study selection included in the rapid review.

For peer review only


References (imported via Endnote)

1. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward J, Hudson L, et al. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults. *JAMA Pediatrics*. 2021;175(2):143.
2. Laxminarayan R, Wahl B, Dudala SR, Gopal K, Mohan BC, Neelima S, et al. Epidemiology and transmission dynamics of COVID-19 in two Indian states. *Science*. 2020;370:691-7.
3. Jones TC, Mühlemann B, Veith T, Biele G, Zuchowski M, Hofmann J, et al. An analysis of SARS-CoV-2 viral load by patient age. *medRxiv*. 2020:2020.06.08.20125484.
4. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). *JAMA Pediatr*. 2020;174(9):902-3.
5. Götzinger F, Santiago-García B, Noguera-Julián A, Lanaspa M, Lancella L, Calò Carducci FI, et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Health*. 2020;4(9):653-61.
6. Dong QQ, Qiu LR, Cheng LM, Shu SN, Chen Y, Zhao Y, et al. Kindergartens Reopening in the Period of Regular Epidemic Prevention and Control, Beneficial or Harmful? *Curr Med Sci*. 2020;40(5):817-21.
7. Milani GP, Bottino I, Rocchi A, Marchisio P, Elli S, Agostoni C, et al. Frequency of Children vs Adults Carrying Severe Acute Respiratory Syndrome Coronavirus 2 Asymptotically. *JAMA Pediatr*. 2021;175(2):193-4.
8. Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr*. 2020;109(6):1088-95.
9. Goldstein E, Lipsitch M. Temporal rise in the proportion of younger adults and older adolescents among coronavirus disease (COVID-19) cases following the introduction of physical distancing measures, Germany, March to April 2020. *Euro Surveill*. 2020;25(17).
10. Cho EY, Choi EH, Kim JH. Interpreting Transmissibility of COVID-19 in Children. *Emerg Infect Dis*. 2020;26(12):3106-7.
11. Control ECfDPa. COVID-19 in children and the role of school settings in transmission - second update. 8 July 2021.
12. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
13. Antonakou A. The latest update on the effects of COVID-19 infection in pregnancy. *Eur J Midwifery*. 2020;4:12.
14. Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary transmission of COVID-19 from children attending school in Ireland, 2020. *Eurosurveillance*. 2020;25.
15. Danis K, Epaulard O, Bénet T, Gaymard A, Campoy S, Botelho-Nevers E, et al. Cluster of Coronavirus Disease 2019 (COVID-19) in the French Alps, February 2020. *Clinical Infectious Diseases*. 2020;71:825-32.
16. Yung CF, Kam K-Q, Nadua KD, Chong CY, Tan NWH, Li J, et al. Novel Coronavirus 2019 Transmission Risk in Educational Settings. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2021;72:1055-8.
17. Gold JA GJ, Kimball A, et al. Infection Among Elementary School Educators and Students in One School District — Georgia, December 2020–January 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:289–292. 2021.

18. Lopez A, Hill M, Antezano J, Vilven D, Rutner T, Bogdanow L, et al. Transmission Dynamics of COVID-19 Outbreaks Associated with Child Care Facilities - Salt Lake City, Utah, April-July 2020. *MMWR Morbidity and mortality weekly report*. 2020;69:1319-23.
19. Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. *The Lancet Child & Adolescent Health*. 2020;4:807-16.
20. Stein-Zamir C, Abramson N, Shoob H, Libal E, Bitan M, Cardash T, et al. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Eurosurveillance*. 2020;25.
21. Link-Gelles R, DellaGrotta AL, Molina C, Clyne A, Campagna K, Lanzieri TM, et al. Limited Secondary Transmission of SARS-CoV-2 in Child Care Programs - Rhode Island, June 1-July 31, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(34):1170-2.
22. Ehrhardt J, Ekinci A, Krehl H, Meincke M, Finci I, Klein J, et al. Transmission of SARS-CoV-2 in children aged 0 to 19 years in childcare facilities and schools after their reopening in May 2020, Baden-Wurttemberg, Germany. *Euro Surveill*. 2020;25.
23. Yoon Y, Kim KR, Park H, Kim S, Kim YJ. Stepwise School Opening and an Impact on the Epidemiology of COVID-19 in the Children. *J Korean Med Sci*. 2020;35:e414.
24. Yoon Y, Choi GJ, Kim JY, Kim KR, Park H, Chun JK, et al. Childcare Exposure to Severe Acute Respiratory Syndrome Coronavirus 2 for 4-Year-Old Presymptomatic Child, South Korea. *Emerg Infect Dis*. 2021;27:341-7.
25. Larosa E, Djuric O, Cassinadri M, Cilloni S, Bisaccia E, Vicentini M, et al. Secondary transmission of COVID-19 in preschool and school settings in northern Italy after their reopening in September 2020: a population-based study. *Euro Surveill*. 2020;25.
26. Brandal LT, Ofitserova TS, Meijerink H, Rykkvin R, Lund HM, Hungnes O, et al. Minimal transmission of SARS-CoV-2 from paediatric COVID-19 cases in primary schools, Norway, August to November 2020. *Euro Surveill*. 2021;26.
27. Pray I, Gibbons-Burgener S, Rosenberg A, Cole D, Borenstein S, Bateman A, et al. COVID-19 Outbreak at an Overnight Summer School Retreat — Wisconsin, July–August 2020. *MMWR Morbidity and Mortality Weekly Report*. 2020;69:1600-4.
28. Blaisdell L, Cohn W, Pavell J, Rubin D, Vergales J. Preventing and mitigating SARS-CoV-2 transmission - Four Overnight Camps, Maine, June-August 2020. *MMWR Morbidity and mortality weekly report*. 2020;69:1216-20.
29. Ladhani SN, Baawuah F, Beckmann J, Okike IO, Ahmad S, Garstang J, et al. SARS-CoV-2 infection and transmission in primary schools in England in June-December, 2020 (sKIDs): an active, prospective surveillance study. *Lancet Child Adolesc Health*. 2021;5(6):417-27.
30. Ismail SA, Saliba V, Lopez Bernal J, Ramsay ME, Ladhani SN. SARS-CoV-2 infection and transmission in educational settings: a prospective, cross-sectional analysis of infection clusters and outbreaks in England. *Lancet Infect Dis*. 2021;21(3):344-53.
31. Bark D, Dhillon N, St-Jean M, Kinniburgh B, McKee G, Choi A. SARS-CoV-2 transmission in kindergarten to grade 12 schools in the Vancouver Coastal Health region: a descriptive epidemiologic study. *CMAJ Open*. 2021;9(3):E810-e7.
32. Hershov RB, Wu K, Lewis NM, Milne AT, Currie D, Smith AR, et al. Low SARS-CoV-2 Transmission in Elementary Schools - Salt Lake County, Utah, December 3, 2020-January 31, 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(12):442-8.
33. Dawson P, Worrell MC, Malone S, Tinker SC, Fritz S, Maricque B, et al. Pilot Investigation of SARS-CoV-2 Secondary Transmission in Kindergarten Through Grade 12 Schools Implementing Mitigation Strategies - St. Louis County and City of Springfield, Missouri, December 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(12):449-55.

- 1
2
3 34. Volpp KG, Kraut BH, Ghosh S, Neatherlin J. Minimal SARS-CoV-2 Transmission After
4 Implementation of a Comprehensive Mitigation Strategy at a School - New Jersey, August 20-
5 November 27, 2020. *MMWR Morb Mortal Wkly Rep.* 2021;70(11):377-81.
- 6 35. Zimmerman KO, Akinboyo IC, Brookhart MA, Boutzoukas AE, McGann KA, Smith MJ, et al.
7 Incidence and Secondary Transmission of SARS-CoV-2 Infections in Schools. *Pediatrics.*
8 2021;147(4):e2020048090.
- 9 36. Ulyte A, Radtke T, Abela IA, Haile SR, Berger C, Huber M, et al. Clustering and longitudinal
10 change in SARS-CoV-2 seroprevalence in school children in the canton of Zurich, Switzerland:
11 prospective cohort study of 55 schools. *BMJ.* 2021;372:n616.
- 12 37. Willeit P, Krause R, Lamprecht B, Berghold A, Hanson B, Stelzl E, et al. Prevalence of RT-
13 qPCR-detected SARS-CoV-2 infection at schools: First results from the Austrian School-SARS-CoV-2
14 prospective cohort study. *Lancet Reg Health Eur.* 2021;5:100086.
- 15 38. Abdollahi E, Haworth-Brockman M, Keynan Y, Langley JM, Moghadas SM. Simulating the
16 effect of school closure during COVID-19 outbreaks in Ontario, Canada. *BMC medicine.*
17 2020;18:230.
- 18 39. Chin ET, Huynh BQ, Lo NC, Hastie T, Basu S. Projected geographic disparities in healthcare
19 worker absenteeism from COVID-19 school closures and the economic feasibility of child care
20 subsidies: a simulation study. *BMC medicine.* 2020;18:218.
- 21 40. Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care
22 workforce and net mortality: a modelling study. *Lancet Public Health.* 2020;5(5):e271-e8.
- 23 41. Zhang J, Litvinova M, Liang Y, Wang Y, Wang W, Zhao S, et al. Changes in contact patterns
24 shape the dynamics of the COVID-19 outbreak in China. *Science (New York, NY).* 2020;368:1481-6.
- 25 42. Koo JR, Cook AR, Park M, Sun Y, Sun H, Lim JT, et al. Interventions to mitigate early spread
26 of SARS-CoV-2 in Singapore: a modelling study. *The Lancet Infectious diseases.* 2020;20:678-88.
- 27 43. Kim S, Kim YJ, Peck KR, Jung E. School Opening Delay Effect on Transmission Dynamics of
28 Coronavirus Disease 2019 in Korea: Based on Mathematical Modeling and Simulation Study. *J*
29 *Korean Med Sci.* 2020;35:e143.
- 30 44. Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al. The effect of control
31 strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan, China: a
32 modelling study. *The Lancet Public Health.* 2020;5:e261-e70.
- 33 45. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and
34 management practices during coronavirus outbreaks including COVID-19: a rapid systematic
35 review. *Lancet Child Adolesc Health.* 2020;4(5):397-404.
- 36 46. Mendez-Brito A, El Bcheraoui C, Pozo-Martin F. Systematic review of empirical studies
37 comparing the effectiveness of non-pharmaceutical interventions against COVID-19. *J Infect.*
38 2021;83(3):281-93.
- 39 47. Sadique MZ, Adams EJ, Edmunds WJ. Estimating the costs of school closure for mitigating
40 an influenza pandemic. *BMC Public Health.* 2008;8:135.
- 41 48. Adverse consequences of school closures: UNESCO; [Available from:
42 <https://en.unesco.org/covid19/educationresponse/consequences>.
- 43 49. Yakut AD. Educators' experiences in special education institutions during the COVID-19
44 outbreak. *J Res Spec Educ Needs.* 2021.
- 45 50. Mansfield KL, Newby D, Sonesson E, Vaci N, Jindra C, Geulayov G, et al. COVID-19 partial
46 school closures and mental health problems: A cross-sectional survey of 11,000 adolescents to
47 determine those most at risk. *JCPP Adv.* 2021;1(2):e12021.
- 48 51. Merckx J, Labrecque JA, Kaufman JS. Transmission of SARS-CoV-2 by Children. *Dtsch*
49 *Arztebl Int.* 2020;117(33-34):553-60.
- 50 52. Li X, Xu W, Dozier M, He Y, Kirolos A, Theodoratou E. The role of children in transmission
51 of SARS-CoV-2: A rapid review. *J Glob Health.* 2020;10(1):011101.
- 52
53
54
55
56
57
58
59
60

- 1
2
3 53. Suk JE, Vardavas C, Nikitara K, Phalkey R, Leonardi-Bee J, Pharris A, et al. The role of
4 children in the transmission chain of SARS-CoV-2: a systematic review and update of current
5 evidence. medRxiv. 2020:2020.11.06.20227264.
6
7 54. Wiedenmann M, Goutaki M, Keiser O, Stringhini S, Tanner M, Low N. The role of children
8 and adolescents in the SARS-CoV-2 pandemic: a rapid review. Swiss Med Wkly. 2021;151:w30058.
9
10 55. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, et al. Ranking the
11 effectiveness of worldwide COVID-19 government interventions. Nat Hum Behav.
12 2020;4(12):1303-12.
13
14 56. Vardavas CI, Odani S, Nikitara K, El Banhawi H, Kyriakos CN, Taylor L, et al. Perceptions and
15 practice of personal protective behaviors to prevent COVID-19 transmission in the G7 nations.
16 Population Medicine. 2020;2(June).
17
18 57. Control ECfDPa. Overview of the implementation of COVID-19 vaccination strategies and
19 vaccine deployment plans in the EU/EEA 23 September 2021.
20
21 58. Yung CF, Kam KQ, Chong CY, Nadua KD, Li J, Tan NWH, et al. Household Transmission of
22 Severe Acute Respiratory Syndrome Coronavirus 2 from Adults to Children. J Pediatr.
23 2020;225:249-51.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Search Strategy

Database: Embase

Search Strategy:

- 1 exp coronavirus/
- 2 exp coronavirus infections/
- 3 (Coronavir* or nCov or covid or Middle East Respiratory Syndrome or MERS or Severe Acute Respiratory Syndrome or SARS).ti,ab,tw.
- 4 1 or 2 or 3
- 5 (adolescent or (pre?school adj child) or child or infant or baby or toddler or juvenile).ti,ab,tw.
- 6 (bab\$ or infant or child or boy or girl or teen\$ or school?child\$).ti,ab,tw.
- 7 5 or 6
- 8 4 and 7
- 9 8
- 10 limit 9 to human
- 11 10
- 12 limit 11 to yr="2019 -Current"
- 13 12
- 14 limit 13 to english language

Database: Ovid MEDLINE(R) ALL

Search Strategy:

- 1 exp Coronavirus/
- 2 exp Coronavirus Infections/
- 3 (Coronavir* or nCov or covid or covid-19 or Middle East Respiratory Syndrome or MERS or Severe Acute Respiratory Syndrome or SARS).ti,ab,kf.
- 4 1 or 2 or 3
- 5 (baby or babies or infant* or child* or boy* or girl* or toddler* or preschool* or pre?school* or teen* or schoolchild* or adolescen* or juvenil*).tw.
- 6 4 and 5
- 7 humans.sh.
- 8 6 and 7
- 9 limit 8 to yr="2019 -Current"
- 10 9
- 11 limit 10 to english language