

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058203
Article Type:	Original research
Date Submitted by the Author:	11-Oct-2021
Complete List of Authors:	Jalloh, Mohamed; Centers for Disease Control and Prevention, Center for Public Health Patel, Palak; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Sutton, Roberta; ICAP at Columbia University Kulkarni, Shibani; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Toure, Mame; ICAP at Columbia University Wiley, Kerrie; The University of Sydney, School of Public Health Sessay, Tom; Sierra Leone Ministry of Health and Sanitation Lahuerta, Maria; ICAP at Columbia University; Columbia University Mailman School of Public Health, Department of Epidemiology
Keywords:	Public health < INFECTIOUS DISEASES, Community child health < PAEDIATRICS, Child protection < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

TITLE PAGE

Title

Caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Authors

Mohamed F. Jalloh¹, Palak Patel^{1,2}, Roberta Sutton³, Shibani Kulkarni^{1,2}, Mame Toure⁴, Kerrie Wiley⁵, Tom Sesay,⁶ Maria Lahuerta^{3,7}, Sierra Leone Urban Immunisation Needs Assessment Group

Affiliations

1. Immunisation Systems Branch, Global Immunisation Division, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
2. Oak Ridge Institute for Science and Education's Research Participation Program, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
3. ICAP at Columbia University, Mailman School of Public Health, New York, New York, USA
4. ICAP at Columbia University, Sierra Leone Country Office, Freetown, Sierra Leone
5. School of Public Health, The University of Sydney, Sydney, NSW, Australia
6. Child Health Programme, Ministry of Health and Sanitation, Freetown, Sierra Leone
7. Department of Epidemiology, Mailman School of Public Health, New York, New York, USA

Corresponding author

Mohamed F. Jalloh, PhD, MPH

Immunisation systems Branch, Global Immunisation Division, U.S. Centres for Disease Control and Prevention, Atlanta, Georgia, USA. Email: yum8@cdc.gov. Tel: +1 (404) 401-2773.

Wordcount

Abstract: 275 words; Manuscript: 4917 words

Keywords

Vaccination, immunisation, vaccine, hesitancy, demand, caregiver, experience, qualitative, methodology, interpretative phenomenology, urban, slum, Sierra Leone

ABSTRACT

Objective: To gain in-depth understanding of the caregiver experience when navigating urban immunisation services for their children.

Design: An exploratory qualitative assessment comprising 16 in-depth interviews using an interpretative phenomenology approach.

Setting: Caregivers were purposively recruited from slums (n=8) and other urban communities (n=8) in the capital city of Sierra Leone.

Participants: Caregivers of children ages 6 to 36 months old who were fully vaccinated (n=8) or under-vaccinated (n=8).

Results: Vaccination intention was motivated by a feeling of moral duty to 'do the right thing' for the child. In one instance, wanting to do the right thing also resulted in an active refusal of vaccination when a caregiver linked vaccination to the death of a prior child. Caregivers with vaccinated children expected that their 'strong and healthy' children would be able to take care of them later in life and believed that this can be facilitated through vaccination. Trusted information exchange and social support coupled with positive clinic experiences facilitated timely vaccination. However, vaccination was constantly hindered by myriad practical constraints that were compounded by negative clinic experiences. Although childhood immunisation is free-of-charge according to national policy, some caregivers willingly gave money to health workers as a token of appreciation while others begrudgingly did so as a perceived condition to receive good quality service. Caregivers desired more convenient and positive clinic experiences and deeper community engagement.

Conclusions: Health system interventions, community engagement, and vaccination outreach need to be tailored to for urban settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is framed both around parental responsibilities to do the right thing for the child and the future benefits to the parent.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This qualitative assessment sheds light on the complexities surrounding caregivers' experiences when navigating childhood immunisation services in urban poor settings and provides insights for improving immunisation outcomes.
- Theoretical guidance from interpretative phenomenology enhanced the assessment design, data analysis, and interpretation of the results.
- Caregivers of fully vaccinated and under-vaccinated children were purposively recruited to help understand how the two outcomes may be shaped by differences in the caregiver experience.
- The sample only comprised one caregiver with a child that had experienced a serious vaccine side effect, which limits having a rich understanding of such experience among caregivers and the potential linkages to future vaccination refusal.
- Only one caregiver had a zero-dose child that had not received any vaccines in the immunisation schedule. Future sampling strategies may therefore need to be more purposively adapted to also focus on understanding the phenomenon of zero-dose children in urban poor settings.

MANUSCRIPT

1. Introduction

There have been efforts to understand urban immunisation challenges in low- and middle-income countries (LMICs), including in urban slums and informal settlements.¹ Assessment of immunisation barriers in urban areas in LMICs identified a range of practical and social issues, such as population mobility, inaccurate denominators of children due to out-of-date population estimates and poorly defined geographic catchment areas, the lack of trust in the health system among vulnerable groups, overburdened health facilities, and weak community engagement and outreach.²

In Sierra Leone, the 2014-2016 Ebola epidemic disrupted the delivery of essential health services, including immunisation services, especially in urban areas.^{3,4} Barriers that affected routine health services included the fear of contracting Ebola in health facilities, stigmatization of health workers, and shifting of resources to the epidemic.⁵ As the Ebola epidemic waned, measles outbreaks became more frequent due to the decline in vaccination.⁶ In the aftermath of the Ebola epidemic, the Government of Sierra Leone and its partners made major investments to rebuild health systems and restore public confidence in the health system.^{7,8} However, challenges in access to and the uptake of immunisation services persist, including in urban areas.⁹

The United Nations Children's Fund (UNICEF) and partners developed the Caregiver Journey Framework to guide countries in understanding the experiences, processes, and structures that shape how caregivers seek and receive health services for their children, including essential immunisation.¹⁰ In 2018, we operationalized the Caregiver Journey Framework through a qualitative approach in the Western Area Urban district (WAU) in Sierra Leone. We sought to understand the lived experiences of caregivers of vaccine-eligible children as they navigate urban immunisation services in Sierra Leone.

2. Methods

We aimed to understand how household dynamics, social factors, and formal healthcare delivery systems influence childhood immunisation uptake via the lens of primary caregivers of vaccine-eligible children in urban settings in Sierra Leone. Hence, we developed the Immunisation Caregiver Journey Interviews (ICJI) approach¹¹ based on the Caregiver Journey Framework using principles of interpretative

phenomenology,^{12 13} which focuses on elucidating the essence of common experiences to explain, interpret, and make sense of a phenomenon.¹⁴ We used a phenomenological approach to explore the lifeworld of caregivers in how they navigate childhood immunisation for their children repeatedly in low-resource, urban settings.¹¹ A semi-structured ICJI guide was used to explore the following domains: Decision-making and preparation, making the journey, experiences during vaccination visit, post-vaccination experiences, intentions to return, and perceptions of immunisation promotion activities in the community.

2.1. Setting

The WAU district in Sierra Leone comprises most of the capital city of Freetown with approximately 1.2 million inhabitants.¹⁵ The district was heavily affected by the Ebola epidemic, partly due to high population movements and crowded housing conditions.¹⁶ On average, there is less than one medical doctor per 10,000 population.¹⁷ The Government of Sierra Leone introduced the Free Health Care Initiative in 2010 to remove cost barriers for essential health services for pregnant and lactating mothers and under-five children.¹⁸ Childhood immunisation services are delivered through the Expanded Program on Immunisation using fixed sites that are supplemented by community outreach services to be conducted five times monthly.¹⁹ Each catchment community has 10 community health workers (CHWs) who support the promotion of health services on a voluntary basis.²⁰ A coverage survey in 2019 estimated 86% coverage for three doses of diphtheria-pertussis-tetanus (DPT) vaccine in slums and 92% coverage in non-slum urban areas (92%) in the WAU district. However, coverage of the second dose of measles-containing vaccine was very low in the district (33% in slums and 29% in non-slum urban areas).⁹

2.2. Sampling and data collection

The sample size for this qualitative assessment was determined based on an approach that focuses on *qualitative information power* considering the study's aim, sample specificity, application of established theory, quality of dialogue, and analysis strategy.²¹ In using interpretive phenomenology as the theoretical basis, we set out a study aim that focused on immunisation-related experiences among a well-defined sample of caregivers of vaccine eligible children. We anticipated that caregivers would provide rich narratives that offer in-depth insights into their experiences. With these considerations in mind, we purposively recruited a total 16 primary caregivers from eight communities in the WAU district, four of which were slums and four were other urban areas in the district to maximize variation in the sample. Within each community, two caregivers of children ages 6-36 months were purposively selected to capture a breadth of caregiver experiences—one whose child was fully vaccinated for age and another whose child had missed at least one scheduled vaccination visit. CHWs supported data collection teams in

1
2
3 visiting households to identify and recruit eligible caregivers in the selected communities. The data
4 collection teams were accompanied by local CHWs in visiting an initial set of households in the
5 community. Snowball sampling was then used to identify additional households with potentially eligible
6 children whereby previously visited households pointed data collectors to other households with
7 potentially eligible children. Data collectors visited such households to screen for eligibility. This process
8 continued until two caregivers of eligible children were successfully recruited and interviewed from a
9 particular community. Interviews were conducted on the same of day of recruitment after obtaining
10 informed consent from the caregiver.
11
12
13
14
15

16
17 We recruited data collectors (interviewers and notetakers) who were fluent in English and the
18 predominant local language in the WAU district (Krio). The data collectors had post-secondary
19 educational training in social sciences and were experienced in conducting qualitative data collection in
20 Sierra Leone. Two behavioural scientists trained the facilitators for a week on the assessment protocol.
21 One of the trainers was from Sierra Leone and had experience conducting social science research in Sierra
22 Leone. During the training, the English version of the guide was translated into Krio by locally hired staff
23 together with the facilitators and trainer. The data collectors were trained on how to probe on the spot to
24 obtain additional pertinent information from caregivers. Data collection occurred in August–September
25 2018. All interviews were audio-recorded with permission from participants; they were then transcribed
26 and translated into English by the local team. Interviews lasted about an hour on average and were
27 conducted in the vicinity of the homes of the caregivers. Data collection teams were trained on choosing
28 suitable interview locations to enable caregivers to speak freely. The facilitators conducted debriefing
29 sessions immediately after each interview to make note of key experiences and observations. We have
30 previously documented practical lessons learned from implementing the assessment in Sierra Leone.¹¹
31
32
33
34
35
36
37
38
39
40

41 **2.3. Data analysis**

42 We analysed the data using a combination of theoretical guidance from interpretative qualitative inquiry
43 (IPA)²² with non-theoretical aspects of qualitative content analysis for inductive coding of meaning units
44 and categorization of codes in latent constructs.²³ Two analysts (one male, one female) read all transcripts
45 and created analytic memos and then analysed the transcripts using both within-case and cross-case
46 analysis. In the within-case analysis, we developed a narrative profile for each caregiver to bring key
47 aspects of their lived experiences to the foreground—consistent with the IPA approach in qualitative
48 research. In the initial part of the cross-case analysis, each analyst coded three different transcripts (six
49 total), using an inductive approach to identify and interpret meaning units within the text. To gain
50 alternative interpretations of the coded meaning units, three of the coded transcripts were shared with a
51
52
53
54
55
56
57
58
59
60

1
2
3 third qualitative expert, who was not involved in the previous stages of the assessment, for independent
4 'blind' coding of the transcripts. Feedback from the third analyst was discussed by the two primary
5 analysts and incorporated into the coding scheme. The analysts used an iterative process to review their
6 codes, discuss their interpretations of the manifest content, and harmonize the initial set of codes that
7 were used for coding the remaining manuscripts. NVivo software (QSR International–2018, Version 12)
8 was used for the final organization and coding of the transcripts. Using a consultative process between the
9 two analysts, manifest categories of meaning units were grouped to reflect latent content that was later
10 developed into cross-cutting themes. Throughout the process, the analysts exercised reflexivity regarding
11 subjective interpretations and iteratively re-examined the transcripts to identify alternative interpretations
12 until consensus was reached.
13
14
15
16
17
18
19
20

21 **2.4. Ethical approval**

22 The assessment was approved by the Sierra Leone Ethics and Scientific Review Committee (SLESRC-
23 17052018), Columbia University Medical Centre Institutional Review Board (IRB-AAAR9031), and the
24 Centre for Global Health at the U.S. Centres for Disease Control and Prevention (CGH-2018-184). All
25 participants provided written informed consent.
26
27
28
29

30 **2.5. Patient and public involvement**

31 No patient involved.
32
33
34
35

36 **3. Results**

37 At the time of the interviews, half of the children had missed at least one scheduled vaccine dose. Four
38 themes emerged from the interviews: Caregivers felt morally obligated to vaccinate their children because
39 of the anticipated benefits (Table 1); vaccination was facilitated by trusted information exchange and
40 social support, coupled with positive clinic experiences (Table 2); vaccination was hindered by practical
41 constraints compounded by negative experiences linked to unfavourable health worker practices and
42 adverse events following immunisation (Table 3); and lastly, caregivers desired stronger community
43 engagement and better experiences at the vaccination clinic (Table 4). There were no notable differences
44 in themes between slums and other urban communities.
45
46
47
48
49
50

51 **3.1. Theme 1. Caregivers felt morally obligated to vaccinate their children and anticipated** 52 **benefits to the child and the parent (Table 1)** 53 54 55 56 57

3.1.1. Obligation to the child

Caregivers felt morally obligated to get their children vaccinated. In essence, vaccination was viewed as ‘doing the right thing’ for the child. Even caregivers who had missed scheduled vaccination felt a responsibility to try to get the child vaccinated.

“It is my duty to take my baby to the hospital for immunisation. It is my responsibility as [a] mother to ensure that my baby completes the rounds of immunisation without defaulting.” –

Caregiver whose child had missed a scheduled vaccination

3.1.2. Wanting a ‘strong and healthy baby’

Caregivers consistently expressed that immunisation has important health benefits to the child, and that missing scheduled vaccination would ‘risk the baby’s life.’ In addition, they valued having a ‘strong and healthy baby’ and felt that completing the vaccination schedule will positively impact the baby’s health.

“I think [the] vaccine is good for our children. It is important and it helps to build their immune system to keep them strong and healthy; it fights against many things in the body...” – Caregiver whose child was fully vaccinated

3.1.3. Future benefits of vaccination for the parent

The notion that vaccinated children will be able to take care of the parent later in life emerged as a dimension of vaccination benefit to the parent.

“It is very difficult for me to be absent for immunisation or not to take my baby to the hospital when sick. Sometimes people think I’m mad but I’m not. I’m trying to bring up my children in a way that they will benefit me when I’m old.” – Caregiver whose child had missed a scheduled vaccination

3.2. Theme 2. Vaccination was facilitated by trusted information exchange and social support, coupled with positive clinic experiences (Table 2)

3.2.1. Diversity of immunisation reminders

Caregivers were exposed to numerous reminders and cues to vaccinate their children, including information gleaned from the immunisation cards, health workers, community campaigns, and family members. However, the child’s immunisation card stood out as the most important reminder that caregivers and families relied on to remember the dates of the scheduled vaccination visits.

1
2
3 *"I do remind myself because they [health workers] will write at the back of the immunisation*
4 *card the date for the next visit. The immunisation card will tell you the date for the next visit. –*
5 *Caregiver whose child had missed a scheduled vaccination*
6
7

8
9 In addition, community outreach campaigns and announcements in the community were also viewed as
10 helpful reminders to vaccinate.
11

12 *"The health workers in this community and the CHWs are doing well as they do go round telling*
13 *people not to forget to take their babies to the hospital for immunisation."* – Caregiver whose
14 *child had missed a scheduled vaccination*
15
16
17

18 19 **3.2.2. Information access and trust**

20 Immunisation information sources varied, but health workers were consistently cited as trusted sources of
21 information. Nurses were more trusted than lay CHWs because nurses were viewed as more
22 knowledgeable. While waiting in line before immunisation services, caregivers appreciated the 'health
23 talk' they received from nurses who advised on health and immunisation.
24

25 *"I trust them [nurses] because they are a team of qualified nurses ... Before they come here, they*
26 *know everything about the vaccines and any implications of the vaccines. They are able to*
27 *explain more than the CHWs and other community workers in the area. The nurses will tell you*
28 *more. There are things that the community workers do not know, and they refer you to the*
29 *nurses."* – Caregiver whose child was fully vaccinated
30
31
32
33
34
35

36 37 **3.2.3. Getting fathers more involved**

38 Fathers were rarely involved in taking their children to the vaccination clinic. We uncovered only two
39 instances when fathers were actively involved. In those instances, the mothers felt supported, and their
40 children were fully vaccinated. In one situation, a father that routinely accompanied the child to the
41 vaccination clinic was celebrated by health workers and given the 'best father' award.
42

43 *"There was [a] time when he [my husband] was given the best father award [at the clinic]*
44 *because he is always with me at the hospital. That is the name I have also given to my husband.*
45 *Even when the baby is crying, I will say 'best father' take your baby."* – Caregiver whose child
46 *was fully vaccinated*
47
48
49
50
51

52 53 **3.2.4. Positive experiences with health workers**

54 Positive experiences with health workers during vaccination visits encouraged caregivers to vaccinate
55 their children. Positive experiences included a cordial relationship with the nurses and health workers.
56
57

They expressed that nurses took good care of their children, made them feel comfortable, and tried to build a good rapport. Some caregivers said they would voluntarily give small amounts of money to health workers as a token of their appreciation after vaccinating the child.

“...the nurses do encourage you and will make jokes so that you will laugh at the end of the day. There is a lot of fun [interactions], which made some of us forget about our stresses.” – Caregiver whose child was fully vaccinated

3.2.5. Post-vaccination information sharing

Upon returning home from the vaccination visit, caregivers often discussed the clinic experience with their spouses, families, and other caregivers in the community. In addition to information obtained from health workers at the clinic, caregivers also sought advice from other ‘more experienced’ caregivers in the community.

“Most times after immunisation, my baby will run a temperature, but the health workers always provide drugs to counter the fever. We have caregivers in this community with vast knowledge and since this is my first baby, I love to talk to them so that we can share our experiences which will be of advantage to me as I’m very new in the field [of parenting].” –Caregiver whose child was fully vaccinated

3.3. Theme 3. Vaccination was hindered by practical constraints compounded by negative experiences linked to unfavourable health worker practices and adverse events following immunisation (Table 3)

3.3.1. Preparing for the journey and getting to the clinic

Caregivers commonly cited the need to juggle ‘household duties’ and other activities when planning the visit as a barrier, especially in the absence of fathers’ involvement in taking the child to the clinic. In addition, mothers frequently depended on their children’s fathers for financial support to cover the expenses related to the vaccination clinic visit. Some caregivers recounted needing to travel long distances to get to the vaccination clinic, which in some instances, took up to an hour on foot.

3.3.2. Inconveniences at the clinic

Caregivers anticipated various inconveniences at the vaccination visit. The prolonged time spent waiting for the child to be vaccinated emerged as a major inconvenience, which was more pronounced when seeking immunisation services at large health facilities, such as hospitals. Anticipating the long wait, caregivers usually tried to arrive early at the vaccination site with the hopes of getting seen first. The

1
2
3 vaccination visit was reportedly prolonged due to the range of activities involved with the child health
4 visit including weighing the baby as part of growth monitoring and other health checks.

5
6 *“It is painful if you waste much time at the health facility because you have other issues to attend*
7 *to. To avoid that, that is why I always come early to the health facility.”* – Caregiver whose child
8 was fully vaccinated
9
10

11 12 13 **3.3.3. Dissatisfaction with specific practices at the clinic**

14 Caregivers complained about some of their experiences and interactions with health care workers during
15 the vaccination visit. A key complaint was that health workers shout at caregivers and sometimes used
16 vulgar language toward caregivers. In other instances, there were complaints that some health workers
17 habitually arrive late to the vaccination session, which further prolonged the time caregivers spent
18 waiting. In addition, systemically hidden costs generated substantial dissatisfaction. Caregivers said that
19 they needed to “shake hands” with health workers at different times of the visit (e.g., first time registering
20 the child to get a card, before entering the facility, and before weighing the baby). Shaking hands implied
21 giving some money during the handshake. Money was used to ‘fast-track’ the child’s vaccination.

22
23 *“Sometimes if I don’t want to spend much time at the hospital, I will shake the hand of the nurse*
24 *so that they can fast track the immunisation of my baby. I will give them something like two*
25 *thousand Leones or whatever I have with me at that moment... Health is wealth and they [health*
26 *workers] don’t need us but we do [need them]. The money we give is nothing compared to the*
27 *health of our children... At the end of the day, we will grumble on our way home as the services*
28 *are supposed to be free for our children, yet we are paying for it. The health workers are really*
29 *trying, but the idea for them to take money from us is bad. And if you don’t give them money, they*
30 *will talk to you carelessly.”* – Caregiver whose child was fully vaccinated
31
32
33
34
35
36
37
38
39
40

41 Money was also demanded as a form of ‘punishment’ to caregivers who missed their children’s scheduled
42 vaccination date.

43
44 *“If you failed to take your baby to the hospital on [the] stipulated date, you will definitely have to*
45 *pay some amount at the end of the day in the form of punishment. You must pay five thousand or*
46 *more.”* – Caregiver whose child was fully vaccinated
47
48
49

50 51 **3.3.4. Adverse events following immunisation**

52 Caregivers cited numerous instances when their children experienced vaccine side effects such as ‘fever,’
53 ‘swelling at the injection site,’ and the ‘baby becoming lethargic.’ Fever was the most common side
54 effect, and the caregivers knew to administer fever-reducing medication as instructed by health workers.
55
56
57

1
2
3 When there was swelling at the injection site, a common practice among caregivers was to massage the
4 swollen area, sometimes with an onion or a bar of soap in their effort to try to reduce the swelling.
5

6 *“Sometimes my baby’s leg becomes swollen... because some nurses are heavy-handed, and I meet*
7 *several nurses when I visit the hospital. Sometimes the leg gets swollen, and they treat him. I have*
8 *to rub the leg to avoid swelling... I use soap to rub off the swelling and I give Panadol to stop the*
9 *fever... some people say you should not allow every nurse to administer [an]injection to the child.*
10 *I should have a permanent nurse that gives injection to my child without swelling.” – Caregiver*
11 *whose child missed a scheduled vaccination*
12
13
14
15

16
17 In one rare situation, a caregiver had a prior child who experienced fever, convulsed, and died a few days
18 after getting vaccinated. Therefore, the caregiver decided to not vaccinate subsequent children.
19

20 *“After the immunisation, my baby started running temperature, I administered paracetamol as I*
21 *was told by the health workers. The baby convulsed and that was the end of that baby. I don’t*
22 *want a repeat of that in my life. I have therefore decided not to take my babies for immunisation*
23 *anymore” – Caregiver whose child missed a scheduled vaccination*
24
25
26
27

28 Besides passive acceptance of the BCG vaccine at the birth of the youngest child, this same caregiver
29 actively refused all other vaccines despite encouragement by a family member to vaccinate the child.
30

31 *“I’m not outrightly saying it was as a result of the immunisation [that my child died]; as every*
32 *death is the work of God. But from what I have gathered so far, I have personally decided not to*
33 *take my baby to the health facility to be immunized. It is not that I’m tired of going to the health*
34 *facility or because of the distance or money. I do get a lot of pressure from my aunt to take my*
35 *baby to the health facility for immunisation, but the thing is that I just don’t trust the system and*
36 *what immunisation does. – Caregiver whose child missed a scheduled vaccination*
37
38
39
40
41

42 From the perspective of this same caregiver, the vaccinated child died but the unvaccinated children
43 survived and thrived, which was cited as a reason for refusing vaccination.
44

45 *“I believe in exclusive breastfeeding, sometimes for two years and a half and sometimes [only]*
46 *two years. My baby is now two years seven months old and doing well like any of those children*
47 *that are on immunisation or have completed...Just as I was saying, sometimes my heart will tell*
48 *me to take the baby for immunisation but after thinking of the past experience, I would decide not*
49 *to. I’m now used to that... The simple fact here is that, since the other children are doing well*
50 *without immunisation, I will not take [the baby] to the health facility for immunisation and that is*
51 *all.” – Caregiver whose child missed a scheduled vaccination*
52
53
54
55
56
57

3.4. Theme 4: Caregivers desired stronger community engagement and better experiences at the vaccination clinic (Table 4)

Implementing community outreach campaigns for immunisation at regular intervals with a focus on defaulters was recommended by participants to improve vaccination outcomes. In addition, caregivers wanted health workers and community leaders to be involved in immunisation promotion along with the CHWs. They wanted the vaccination clinic experience to improve and become more conducive to caregivers, including shorter wait time at the clinic and more positive interactions with health workers. Lastly, caregivers wanted health workers to stop demanding money from them, though they may not mind giving money as a token of appreciation when they could afford it.

“In addition, you should engage the Chiefs, because in each area we have Chiefs to spread out this message. You could educate them so that they in turn can educate those in the community. Let us have Town Criers go around disseminating the messages. It would be nice for them to allocate people in the health centre who move from house to house to educate the breastfeeding mothers because some of us are stubborn to come onboard.” – Caregiver whose child was fully vaccinated

4. Discussion

Our qualitative analysis highlighted several important themes. In the backdrop of anticipated benefits to both the child and parent, vaccination intention was motivated by a feeling of moral duty to ‘do the right thing.’ Timely and trusted exchange of information together with social support and positive experiences at the vaccination clinic were important facilitators of vaccination. In contrast, vaccination was discouraged by negative interactions with health workers at the clinic, the occurrence and fear of vaccine side effects, multitude of ‘hidden’ costs, juggling vaccination with other responsibilities, and inconveniences, such as long traveling time to the clinic and long delays at the clinic. Nevertheless, caregivers were resilient in devising ways to try to get their children vaccinated. Lastly, caregivers wanted the vaccination experience to improve, and they desired stronger community engagement to help optimize vaccination outcomes. However, systemic issues, such as informal payments, overcrowding in health facilities, and the reported overburdened health workers may require interventions at the health systems level. These themes from Sierra Leone provide in-depth insights regarding the motivations, facilitators, and barriers of vaccination in an urban LMIC setting.

1
2
3 Moral values may shape vaccination attitudes.²⁴ Philosophical arguments regarding the morality of
4 vaccination have been heavily debated.²⁵⁻²⁸ Caregivers in our sample largely viewed vaccination via a
5 moral lens encompassing parental duty to do the right thing for the child. In one situation, however, we
6 found that the desire to ‘do the right thing’ can also translate into vaccination refusal in the backdrop of
7 other past refusals, observing ‘healthy unvaccinated’ children, and having distrust of the health system.
8 Quantitative research from high-income countries has shown that parents with unvaccinated children were
9 more likely to perceive their children to be at low risk of vaccine-preventable diseases and were more
10 likely to perceive low vaccine effectiveness and safety compared to parents with vaccinated children.²⁹
11 Our findings suggest that childhood immunisation communication efforts may resonate more strongly
12 with caregivers when vaccination is framed around parental responsibilities to do the right thing for the
13 child and the anticipated future benefits to parents.
14
15
16
17
18
19
20
21

22 Across the interviews, there was an apparent tension in the relationship between caregivers and health
23 workers. Caregivers often expressed their appreciation of health workers and empathized with the
24 challenging context in which they do their work. Health workers were strongly viewed as authoritative
25 sources of trusted information regarding immunisation and the child’s health, which is consistent with
26 findings from high-income countries³⁰⁻³² and LMICs.^{33 34} Our findings on the role of monetary exchange
27 in vaccination exemplify the complex relationship between caregivers and health workers in low-resource
28 urban communities in Sierra Leone. Some caregivers voluntarily gave money to health workers as a
29 ‘token of appreciation’ while others begrudgingly gave money because they viewed it as a condition for
30 receiving good quality service from health workers. Interventions at the health systems level are
31 necessary to help discourage informal payments to health workers—a practice that may perpetuate
32 vaccination inequities among poor caregivers who are unable to meet monetary expectations.
33
34
35
36
37
38
39
40

41 Our findings also illuminate the need for interventions at the household and family level. We uncovered
42 that fathers were rarely involved in taking their children to the vaccination clinic but were often engaged
43 in the decision-making processes. In the few instances when fathers were involved in taking their children
44 to the clinic, the mothers felt supported, and their children were fully vaccinated. A study in Nigeria
45 found that paternal involvement in immunisation was greater in rural settings compared to urban
46 settings.³⁵ In urban areas, the same study found that paternal involvement was greater among educated
47 fathers compared to uneducated fathers. In a separate study in Ghana, the involvement of educated fathers
48 in the vaccination decision was associated with timelier vaccination uptake compared to the involvement
49 of uneducated fathers in the decision.³⁶ More broadly, shifting from a mother-child dyad to a family triad
50 in the care of children has proven to have positive effects on paediatric health outcomes across diverse
51
52
53
54
55
56
57
58
59
60

1
2
3 contexts.³⁷ Additional assessments and interventions are needed to explore and evaluate culturally
4 appropriate ways to enhance the involvement of fathers in childhood immunisation in Sierra Leone and
5 other similar LMIC settings.
6
7

8
9 Existing evidence suggests that vaccine safety concerns, often linked to adverse events following
10 immunisation (AEFI), contribute to vaccine hesitancy.^{38 39} Serious AEFIs may be ‘triggering events’ for
11 derailing vaccine confidence and prompting active refusal among certain caregivers and their
12 communities—especially when the serious AEFI is perceived to be linked to the vaccine or the
13 vaccination process.⁴⁰ Together with prior evidence, these findings emphasize the need for robust AEFI
14 surveillance and investigations⁴¹ to identify, counsel, and follow-up with caregivers whose children
15 experience AEFI, and therefore, are potentially at risk of missing subsequent vaccination. Health workers
16 and CHWs may benefit from periodic in-service training on how to effectively communicate vaccine
17 safety and address concerns about AEFIs.⁴²
18
19
20
21
22
23
24

25 **4.1. Limitations**

26
27 There were several limitations to our assessment. First, it is possible that some nuanced meaning may
28 have been lost when translating the audio recordings from Krio to English. Second, we only identified one
29 caregiver who routinely and actively refused all vaccines for her child after experiencing a serious AEFI,
30 which may reflect the overall rarity of zero-dose unvaccinated children in Sierra Leone (approximately
31 3%).⁴³ This was the only caregiver with a child that experienced a serious AEFI in our sample, which
32 limits our ability to have a rich understanding of such experience among caregivers more broadly and the
33 potential linkages to vaccination refusal. Although such experiences of serious AEFI are rare, they may
34 have the tendency to get publicized in the community, which may negatively influence vaccination
35 decisions among other caregivers. Taken together, our results call for additional qualitative assessments to
36 get a deeper understanding of vaccination refusal within the Sierra Leonean context and other low-
37 resource LMIC settings. Sampling strategies may therefore need to be adapted accordingly to focus on
38 caregivers who actively refuse vaccination for their children. Lastly, our findings reflected gender
39 dimensions that may be based on sociocultural norms in Sierra Leonean society but may also be due to
40 sampling bias because the caregivers we conveniently recruited were mostly stay-at-home mothers.
41
42
43
44
45
46
47
48
49

50 **5. Conclusions**

51
52 As the COVID-19 pandemic disrupts childhood immunisation globally,⁴⁴ especially in LMICs, our
53 assessment provides a foundational understanding of the challenges that caregivers encounter in urban
54 settings in Sierra Leone. It also sheds light on opportunities to improve vaccination outcomes in urban
55
56
57

1
2
3 poor settings, which is a global immunisation priority. The findings show that health system
4 interventions, community engagement, and vaccination outreach may need to be tailored to for urban
5 settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is
6 framed both around parental responsibilities to do the right thing for the child and the future benefits to
7 the parent.
8
9
10

11 12 13 14 **Competing interests**

15 None declared.
16
17

18 19 **Data availability**

20 All data relevant to the study are included in the article or uploaded as supplementary information.
21
22
23

24 **Contributors**

25 MFJ, RS, MT, TS, and ML designed the assessment. MFJ, PP, SK, and KW analysed the data. MJF, PK,
26 RS, SK, MT, KW, TS, and ML contributed to the interpretation of the results. MFJ, PP and SK drafted
27 the manuscript. RS, MT, KW, TS, and ML contributed to revising the manuscript. All authors reviewed
28 and approved the final manuscript.
29
30
31
32

33 **Acknowledgements**

34 We thank the 16 caregivers who participated in the in-depth interviews.
35
36
37

38 **Funding**

39 U.S. Centers for Disease Control and Prevention (U19GGH001581).
40
41
42

43 **Disclaimer**

44 The findings and conclusions in this paper are those of the authors and do not necessarily represent the
45 official position of the US Centres for Disease Control and Prevention.
46
47
48
49
50
51
52
53
54
55
56
57

6. References

1. Nelson KN, Wallace AS, Sodha SV, et al. Assessing strategies for increasing urban routine immunisation coverage of childhood vaccines in low and middle-income countries: A systematic review of peer-reviewed literature. *Vaccine* 2016;34(46):5495-503. doi: 10.1016/j.vaccine.2016.09.038
2. JSI. Strengthening Immunisation for the Urban Poor: Experiences and Lessons Learned from Five Country Studies, 2018. Available from: <https://www.jsi.com/resource/strengthening-immunisation-for-the-urban-poor-experiences-and-lessons-learned-from-five-country-studies/> [accessed November 12 2020].
3. Brolin Ribacke KJ, Saulnier DD, Eriksson A, et al. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Frontiers in public health* 2016;4:222. doi: 10.3389/fpubh.2016.00222
4. Elston JW, Cartwright C, Ndumbi P, et al. The health impact of the 2014-15 Ebola outbreak. *Public health* 2017;143:60-70. doi: 10.1016/j.puhe.2016.10.020
5. Yerger P, Jalloh M, Coltart CEM, et al. Barriers to maternal health services during the Ebola outbreak in three West African countries: a literature review. *BMJ Glob Health* 2020;5(9) doi: 10.1136/bmjgh-2020-002974
6. Masresha BG, Luce R, Jr., Weldegebriel G, et al. The impact of a prolonged ebola outbreak on measles elimination activities in Guinea, Liberia and Sierra Leone, 2014-2015. *Pan Afr Med J* 2020;35(Suppl 1):8. doi: 10.11604/pamj.suppl.2020.35.1.19059
7. Government of Sierra Leone. National Ebola Recovery Strategy for Sierra Leone 2015 [Available from: https://ebolaresponse.un.org/sites/default/files/sierra_leone_-_national_recovery_strategy_2015-2017.pdf accessed December 5 2020].
8. U.S. Centers for Disease Control and Prevention. Global Health – Sierra Leone, 2019. Available from: <https://www.cdc.gov/globalhealth/countries/sierra-leone/default.htm> [accessed December 5 2020].
9. Feldstein LR, Sutton R, Jalloh MF, et al. Access, demand, and utilization of childhood immunisation services: A cross-sectional household survey in Western Area Urban district, Sierra Leone, 2019. *Journal of global health* 2020;10(1):010420. doi: 10.7189/jogh.10.010420
10. UNICEF. Human Centered Design 4 Health, 2018. Available from: <https://www.hcd4health.org/> [accessed November 18 2020].
11. Jalloh MF, Kinsman J, Conteh J, et al. Barriers and facilitators to reporting deaths following Ebola surveillance in Sierra Leone: implications for sustainable mortality surveillance based on an exploratory qualitative assessment. *BMJ Open* 2021;11(5):e042976. doi: 10.1136/bmjopen-2020-042976
12. Diekelmann NL. Learning-as-testing: a Heideggerian hermeneutical analysis of the lived experiences of students and teachers in nursing. *ANS Advances in nursing science* 1992;14(3):72-83. doi: 10.1097/00012272-199203000-00010
13. Dreyfus HL. Being-in-the-world: A commentary on Heidegger's Being and Time. Cambridge: The MIT Press 1991.

14. Dahlberg K. The essence of essences – the search for meaning structures in phenomenological analysis of lifeworld phenomena. *International journal of qualitative studies on health and well-being* 2006;1(1):11-19. doi: 10.1080/17482620500478405
15. Statistics Sierra Leone. 2015 Sierra Leone Population and Housing Census. 2016. Available from: <https://www.statistics.sl/index.php/census/census-2015.html> [accessed November 12, 2020]
16. Lamunu M, Olu OO, Bangura J, et al. Epidemiology of Ebola Virus Disease in the Western Area Region of Sierra Leone, 2014-2015. *Frontiers in public health* 2017;5:33. doi: 10.3389/fpubh.2017.00033 [published Online First: 2017/03/18]
17. World Health Organization. Medical doctors per 10,000 population 2021 [Available from: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/medical-doctors-\(per-10-000-population\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/medical-doctors-(per-10-000-population))] [accessed December 5 2021].
18. Witter S, Brikci N, Harris T, et al. The free healthcare initiative in Sierra Leone: Evaluating a health system reform, 2010-2015. *The International journal of health planning and management* 2018;33(2):434-48. doi: 10.1002/hpm.2484
19. Government of Sierra Leone. Comprehensive EPI Multi-Year Plan 2017-2021, 2016. Available from: <https://www.gavi.org/sites/default/files/document/2021/cMYP%20Sierra%20Leone%202017-2021.pdf> [accessed December 5 2020].
20. Government of Sierra Leone. Policy for Community Health Workers in Sierra Leone, 2012. Available from: <https://chwcentral.org/wp-content/uploads/2015/01/CHW-Policy-Sierra-Leone.pdf> [accessed December 5 2020].
21. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided by Information Power. *Qualitative health research* 2016;26(13):1753-60. doi: 10.1177/1049732315617444
22. Smith JA, Flowers P, Larkin M. Interpretative phenomenological analysis: theory, method and research. United States: Sage 2009.
23. Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse education today* 2004;24(2):105-12. doi: 10.1016/j.nedt.2003.10.001
24. Amin AB, Bednarczyk RA, Ray CE, et al. Association of moral values with vaccine hesitancy. *Nature human behaviour* 2017;1(12):873-80. doi: 10.1038/s41562-017-0256-5
25. Di Pietro ML, Poscia A, Telesman AA, et al. Vaccine hesitancy: parental, professional and public responsibility. *Annali dell'Istituto superiore di sanita* 2017;53(2):157-62. doi: 10.4415/ann_17_02_13
26. Giubilini A, Douglas T, Savulescu J. The moral obligation to be vaccinated: utilitarianism, contractualism, and collective easy rescue. *Medicine, health care, and philosophy* 2018;21(4):547-60. doi: 10.1007/s11019-018-9829-y
27. Bester JC. Measles Vaccination is Best for Children: The Argument for Relying on Herd Immunity Fails. *Journal of bioethical inquiry* 2017;14(3):375-84. doi: 10.1007/s11673-017-9799-4
28. Dawson A. What are the moral obligations of the traveller in relation to vaccination? *Travel medicine and infectious disease* 2007;5(2):90-6. doi: 10.1016/j.tmaid.2006.01.005

- 1
2
3
4 29. Omer SB, Salmon DA, Orenstein WA, et al. Vaccine refusal, mandatory immunisation, and the risks
5 of vaccine-preventable diseases. *N Engl J Med* 2009;360(19):1981-8. doi: 10.1056/NEJMsa0806477
6
7 30. Frew PM, Chung Y, Fisher AK, et al. Parental experiences with vaccine information statements:
8 Implications for timing, delivery, and parent-provider immunisation communication. *Vaccine*
9 2016;34(48):5840-44. doi: 10.1016/j.vaccine.2016.10.026
10
11 31. Drezner D, Youngster M, Klainer H, et al. Maternal vaccinations coverage and reasons for non-
12 compliance - a cross-sectional observational study. *BMC pregnancy and childbirth* 2020;20(1):541. doi:
13 10.1186/s12884-020-03243-w
14
15 32. Wiley KE, Massey PD, Cooper SC, et al. Pregnant women's intention to take up a post-partum
16 pertussis vaccine, and their willingness to take up the vaccine while pregnant: a cross sectional survey.
17 *Vaccine* 2013;31(37):3972-8. doi: 10.1016/j.vaccine.2013.06.015
18
19 33. Makwe CC, Anorlu RI. Knowledge of and attitude toward human papillomavirus infection and
20 vaccines among female nurses at a tertiary hospital in Nigeria. *International journal of women's health*
21 2011;3:313-7. doi: 10.2147/ijwh.S22792
22
23 34. Foss HS, Oldervoll A, Fretheim A, et al. Communication around HPV vaccination for adolescents in
24 low- and middle-income countries: a systematic scoping overview of systematic reviews. *Systematic*
25 *reviews* 2019;8(1):190. doi: 10.1186/s13643-019-1100-y
26
27 35. Sodeinde K, Amoran O, Abiodun O, et al. A rural/urban comparison of paternal involvement in
28 childhood immunisation in Ogun Central Senatorial District, Nigeria. *The Nigerian postgraduate medical*
29 *journal* 2020;27(4):336-42. doi: 10.4103/npmj.npmj_101_20
30
31 36. Brugha RF, Kevany JP, Swan AV. An investigation of the role of fathers in immunisation uptake. *Int*
32 *J Epidemiol* 1996;25(4):840-5. doi: 10.1093/ije/25.4.840
33
34 37. Allport BS, Johnson S, Aqil A, et al. Promoting Father Involvement for Child and Family Health.
35 *Acad Pediatr* 2018;18(7):746-53. doi: 10.1016/j.acap.2018.03.011
36
37 38. Dube E, Laberge C, Guay M, et al. Vaccine hesitancy: an overview. *Human vaccines &*
38 *immunotherapeutics* 2013;9(8):1763-73. doi: 10.4161/hv.24657
39
40 39. Larson HJ, Jarrett C, Eckersberger E, et al. Understanding vaccine hesitancy around vaccines and
41 vaccination from a global perspective: a systematic review of published literature, 2007-2012. *Vaccine*
42 2014;32(19):2150-9. doi: 10.1016/j.vaccine.2014.01.081
43
44 40. Du F, Chantler T, Francis MR, et al. The determinants of vaccine hesitancy in China: A cross-
45 sectional study following the Changchun Changsheng vaccine incident. *Vaccine* 2020;38(47):7464-71.
46 doi: 10.1016/j.vaccine.2020.09.075
47
48 41. World Health Organization. Adverse events following immunisation (AEFI), 2021. Available from:
49 https://www.who.int/immunisation/monitoring_surveillance/who-immuniz.pdf [accessed February 18
50 2021].
51
52 42. World Health Organization. Vaccine safety basics e-learning course, 2021. Available from:
53 <https://vaccine-safety-training.org/simplified-and-key-messages.html> [accessed February 19 2021].
54
55
56
57
58
59
60

1
2
3
4 43. UNICEF. Sierra Leone Multiple Indicators Cluster Survey, 2017. Available from:
5 https://mics.unicef.org/news_entries/106/SIERRA-LEONE-2017-MICS-RELEASED [accessed January
6 21 2021].
7

8
9 44. Abbas K, Procter SR, van Zandvoort K, et al. Routine childhood immunisation during the COVID-19
10 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2
11 infection. *Lancet Glob Health* 2020;8(10):e1264-e72. doi: 10.1016/s2214-109x(20)30308-9
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1. Thematic area on the motivation to get the child vaccinated–Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Theme
Responsibility to ensure full vaccination	Obligation to the child	Caregivers felt morally obligated to vaccinate their children and anticipated benefits to the child and the parent
Obligation to do the right thing		
Immunisation as a requirement		
Immunisation is important for baby's health	Anticipated benefits	
Wanting 'strong baby'		
'Health is wealth'		
Defaulting on vaccination risks baby's life		
Vaccination benefits the parent later in life		

Table 2. Thematic area on enablers of vaccination–Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Theme
Immunisation card is important	Diversity of immunisation reminders	Vaccination was facilitated by trusted information exchange and social support coupled with positive experiences
Campaign as reminder		
Reminders by health workers at vaccination visit		
Husband as reminder		
Immunisation card as reminder		
Other family member as reminder		
Nurses should lead	Information access and trust	
Nurses more trusted than CHWs		
Same information from different sources		
Immunisation promotion through radio and tv		
Immunisation promotion by health workers		
Immunisation promotion through leaders		
Immunisation promotion by NGOs	Getting fathers more involved	
Mothers take the child to the clinic		
Fathers rarely involved		
Father received an award for involvement	Positive experiences with health worker	
Cordial relationship with nurses		
Good care by nurses		
HWs encourage seeking care at HF		
Giving money to health worker as token of appreciation	Post-vaccination information sharing	
Husband asking about the visit		
Informing husband of next visit		
Telling husband about visit expenses		
Immunisation is a "learning process"		
Sharing experiences with neighbours or friends		
Other family members asking about the visit		

Table 3. Thematic area on vaccination barriers–Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Theme
Asking husband for money	Preparing for the journey and getting to the clinic	Vaccination was hindered by practical constraints compounded by negative experiences linked to unfavourable health worker practices and adverse events following immunisation
Juggling different household duties		
Competing priorities		
Time taken to get to the clinic		
Wasting time at the clinic	Inconveniences at the clinic	
Rush to arrive first and seen first		
Crowding at the clinic		
Very long wait at larger health facilities		
Spent a day and nothing happened		
Not respecting caregivers	Dissatisfaction with specific practices at the clinic	
Shouting at caregivers		
Wasting caregiver's time		
Paying for the card		
Paying for free drugs		
Paying for weighing		
Payment as punishment		
Bad care without payment		
Health workers should stop demanding money		
Health worker don't appreciate less than Le 2000		
Health workers withholding free drugs	Adverse events following immunisation	
Baby crying throughout the night		
Baby gets 'lazy' for few hours		
Fever in baby		
Swelling at injection site		
Side effects only for some vaccines		
Rub onion on swollen injection site		
Avoiding abnormalities from vaccine		
Afraid of vaccine side effects		

Table 4. Thematic area on recommendations to improve childhood immunisation services– Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Theme
Continue campaigns at repeated intervals	Improving vaccination processes and systems	Caregivers want improved vaccination processes, systems, and engagement
Do not rely on campaigns alone		
Provision of incentives for caregivers		
Concentrate on defaulters		
Compensation/incentives for nurses		
Being considerate towards health workers		
Employ more staff		
Stop demanding money		
Promote consequences of not vaccinating	Engaging communities to boost vaccine confidence	
Peer-to-peer promotion of immunisation		
Inform about importance of immunisation		
Address 'stubborn' caregivers		
Engage caregivers who do not want injections		

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

COREQ (CONsolidated criteria for REporting Qualitative research) Checklist

A checklist of items that should be included in reports of qualitative research. You must report the page number in your manuscript where you consider each of the items listed in this checklist. If you have not included this information, either revise your manuscript accordingly before submitting or note N/A.

Topic	Item No.	Guide Questions/Description	Reported on Page No.
Domain 1: Research team and reflexivity			
<i>Personal characteristics</i>			
Interviewer/facilitator	1	Which author/s conducted the interview or focus group?	
Credentials	2	What were the researcher's credentials? E.g. PhD, MD	
Occupation	3	What was their occupation at the time of the study?	
Gender	4	Was the researcher male or female?	
Experience and training	5	What experience or training did the researcher have?	
<i>Relationship with participants</i>			
Relationship established	6	Was a relationship established prior to study commencement?	
Participant knowledge of the interviewer	7	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	
Interviewer characteristics	8	What characteristics were reported about the interviewer/facilitator? e.g. Bias, assumptions, reasons and interests in the research topic	
Domain 2: Study design			
<i>Theoretical framework</i>			
Methodological orientation and Theory	9	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	
<i>Participant selection</i>			
Sampling	10	How were participants selected? e.g. purposive, convenience, consecutive, snowball	
Method of approach	11	How were participants approached? e.g. face-to-face, telephone, mail, email	
Sample size	12	How many participants were in the study?	
Non-participation	13	How many people refused to participate or dropped out? Reasons?	
<i>Setting</i>			
Setting of data collection	14	Where was the data collected? e.g. home, clinic, workplace	
Presence of non-participants	15	Was anyone else present besides the participants and researchers?	
Description of sample	16	What are the important characteristics of the sample? e.g. demographic data, date	
<i>Data collection</i>			
Interview guide	17	Were questions, prompts, guides provided by the authors? Was it pilot tested?	
Repeat interviews	18	Were repeat interviews carried out? If yes, how many?	
Audio/visual recording	19	Did the research use audio or visual recording to collect the data?	
Field notes	20	Were field notes made during and/or after the interview or focus group?	
Duration	21	What was the duration of the interviews or focus group?	
Data saturation	22	Was data saturation discussed?	
Transcripts returned	23	Were transcripts returned to participants for comment and/or	

Topic	Item No.	Guide Questions/Description	Reported on Page No.
		correction?	
Domain 3: analysis and findings			
<i>Data analysis</i>			
Number of data coders	24	How many data coders coded the data?	
Description of the coding tree	25	Did authors provide a description of the coding tree?	
Derivation of themes	26	Were themes identified in advance or derived from the data?	
Software	27	What software, if applicable, was used to manage the data?	
Participant checking	28	Did participants provide feedback on the findings?	
<i>Reporting</i>			
Quotations presented	29	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	
Data and findings consistent	30	Was there consistency between the data presented and the findings?	
Clarity of major themes	31	Were major themes clearly presented in the findings?	
Clarity of minor themes	32	Is there a description of diverse cases or discussion of minor themes?	

Developed from: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

Once you have completed this checklist, please save a copy and upload it as part of your submission. DO NOT include this checklist as part of the main manuscript document. It must be uploaded as a separate file.

BMJ Open

Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058203.R1
Article Type:	Original research
Date Submitted by the Author:	03-Mar-2022
Complete List of Authors:	Jalloh, Mohamed; Centers for Disease Control and Prevention, Center for Public Health Patel, Palak; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Sutton, Roberta; ICAP at Columbia University Kulkarni, Shibani; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Toure, Mame; ICAP at Columbia University Wiley, Kerrie; The University of Sydney, School of Public Health Sessay, Tom; Sierra Leone Ministry of Health and Sanitation Lahuerta, Maria; ICAP at Columbia University; Columbia University Mailman School of Public Health, Department of Epidemiology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Qualitative research, Public health, Health services research, Global health
Keywords:	Public health < INFECTIOUS DISEASES, Community child health < PAEDIATRICS, Child protection < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

TITLE PAGE

Title

Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Authors

Mohamed F. Jalloh¹, Palak Patel^{1,2}, Roberta Sutton³, Shibani Kulkarni^{1,2}, Mame Toure⁴, Kerrie Wiley⁵, Tom Sesay,⁶ Maria Lahuerta^{3,7}, Sierra Leone Urban Immunisation Needs Assessment Group

Affiliations

1. Immunisation Systems Branch, Global Immunisation Division, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
2. Oak Ridge Institute for Science and Education's Research Participation Program, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
3. ICAP at Columbia University, Mailman School of Public Health, New York, New York, USA
4. ICAP at Columbia University, Sierra Leone Country Office, Freetown, Sierra Leone
5. School of Public Health, The University of Sydney, Sydney, NSW, Australia
6. Child Health Programme, Ministry of Health and Sanitation, Freetown, Sierra Leone
7. Department of Epidemiology, Mailman School of Public Health, New York, New York, USA

Corresponding author

Mohamed F. Jalloh, PhD, MPH

Immunisation systems Branch, Global Immunisation Division, U.S. Centres for Disease Control and Prevention, Atlanta, Georgia, USA. Email: yum8@cdc.gov. Tel: +1 (404) 401-2773.

Wordcount

Abstract: 239 words; Manuscript: 5438 words

Keywords

Vaccination, immunisation, vaccine, hesitancy, demand, caregiver, experience, qualitative, methodology, interpretative phenomenology, urban, slum, Sierra Leone

ABSTRACT

Objective: To gain in-depth understanding of the caregiver experience when navigating urban immunisation services for their children.

Design: An exploratory qualitative assessment comprising 16 in-depth interviews using an interpretative phenomenology approach.

Setting: Caregivers were purposively recruited from slums (n=8) and other urban communities (n=8) in the capital city of Sierra Leone.

Participants: Caregivers of children ages 6 to 36 months old who were fully vaccinated (n=8) or under-vaccinated (n=8).

Results: Emotional enablers of vaccination were evident in caregivers' sense of parental obligation to their children while also anticipating reciprocal benefits in children's ability to take care of the parents later in life. Practical enablers were found in the diversity of immunization reminders, information access, information trust, getting fathers more involved, positive experiences with health workers, and post-vaccination information sharing in the community. Underlying barriers to childhood vaccination were due to practical constraints such as overcrowding and long waiting times at the clinic, feeling disrespected by health workers, expecting to give money to health workers for free services, and fear of serious vaccine side effects. To improve vaccination outcomes, caregivers desired more convenient and positive clinic experiences and deeper community engagement.

Conclusions: Health system interventions, community engagement, and vaccination outreach need to be tailored to for urban settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is framed both around parental responsibilities to do the right thing for the child and the future benefits to the parent.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This qualitative assessment describes the caregiver experience in navigating childhood immunisation services in urban settings in a low-income country to gain insights for improving vaccination outcomes.
- The sample comprised of caregivers of fully vaccinated and under-vaccinated children who were purposively recruited to help understand how the two outcomes may be shaped by differences in the caregiver experiences.
- Our sample only comprised one caregiver with a prior child that had experienced a serious vaccine side effect and whose current child had never been vaccinated due to fear of similar side effects.
- The rarity of experiencing serious side effects among children in our sample limits the ability to have a rich understanding of such experiences.

1. Introduction

There have been efforts to understand urban immunisation challenges in low- and middle-income countries (LMICs), including in urban slums and informal settlements.¹ Assessment of immunisation barriers in urban areas in LMICs identified a range of practical and social issues, such as population mobility, inaccurate denominators of children due to out-of-date population estimates and poorly defined geographic catchment areas, the lack of trust in the health system among vulnerable groups, overburdened health facilities, and weak community engagement and outreach.²

In Sierra Leone, the 2014-2016 Ebola epidemic disrupted the delivery of essential health services, including immunisation services, especially in urban areas.^{3,4} Barriers that affected routine health services included the fear of contracting Ebola in health facilities, stigmatization of health workers, and shifting of resources to the epidemic.⁵ As the Ebola epidemic waned, measles outbreaks became more frequent due to the decline in vaccination.⁶ In the aftermath of the Ebola epidemic, the Government of Sierra Leone and its partners made major investments to rebuild health systems and restore public confidence in the health system.^{7,8} However, challenges in access to and the uptake of immunisation services persist, including in urban areas.⁹

The United Nations Children's Fund (UNICEF) and partners developed the Caregiver Journey Framework to guide countries in understanding the experiences, processes, and structures that shape how caregivers seek and receive health services for their children, including essential immunisation.¹⁰ In 2018, we operationalized the Caregiver Journey Framework through a qualitative approach in the Western Area Urban district (WAU) in Sierra Leone. Implementation experiences from operationalizing the framework in the context of urban immunization has been described elsewhere.¹¹ The framework was operationalized into several domains to understand decision-making and preparation for vaccination visits, making the journey to clinics, experiences during vaccination visits, and post-vaccination experiences. Building on these domains, we aimed to describe the real-world experiences of caregivers of vaccine-eligible children as they navigate urban immunisation services in Sierra Leone to identify vaccination enablers and barriers.

2. Methods

We aimed to understand how household dynamics, social factors, and formal healthcare delivery systems influence childhood immunisation uptake via the lens of primary caregivers of vaccine-eligible children in urban settings in Sierra Leone. Hence, we developed the Immunisation Caregiver Journey Interviews

(ICJI) approach¹¹ based on the Caregiver Journey Framework using principles of interpretative phenomenology,^{12 13} which focuses on elucidating the essence of common experiences to explain, interpret, and make sense of a phenomenon.¹⁴ We used a phenomenological approach to explore the lifeworld of caregivers in how they navigate childhood immunisation for their children repeatedly in low-resource, urban settings.¹¹ A semi-structured ICJI guide was used to explore the following domains: Decision-making and preparation, making the journey, experiences during vaccination visit, post-vaccination experiences, intentions to return, and perceptions of immunisation promotion activities in the community.

2.1. Setting

The WAU district in Sierra Leone comprises most of the capital city of Freetown with approximately 1.2 million inhabitants.¹⁵ The district was heavily affected by the Ebola epidemic, partly due to high population movements and crowded housing conditions.¹⁶ On average, there is less than one medical doctor per 10,000 population.¹⁷ The Government of Sierra Leone introduced the Free Health Care Initiative in 2010 to remove cost barriers for essential health services for pregnant and lactating mothers and under-five children.¹⁸ Childhood immunisation services are delivered through the Expanded Program on Immunisation using fixed sites that are supplemented by community outreach services to be conducted five times monthly.¹⁹ Each catchment community has 10 community health workers (CHWs) who support the promotion of health services on a voluntary basis.²⁰ A coverage survey in 2019 estimated 86% coverage for three doses of diphtheria-pertussis-tetanus (DPT) vaccine in slums and 92% coverage in non-slum urban areas in the WAU district. However, coverage of the second dose of measles-containing vaccine was very low in the district (33% in slums and 29% in non-slum urban areas).⁹

2.2. Sampling and data collection

The sample size for this qualitative assessment was guided by an approach that focuses on *qualitative information power*.²¹ The concept of information power posits that researchers should determine the sample size in a qualitative assessment based on the aim (narrow versus broad), sample specificity (targeting specific group versus multiple groups), theoretical underpinning (application of theory or no theory), quality of dialogue (weak or strong), and analysis strategy (within-case only or cross-case). Sample size burden increases when the aim is broad, multiple groups are targeted in the sample, the assessment is theory-driven, the quality of the dialogue is weak, and transcripts are analysed using cross-case analysis. In our assessment, the aim was narrow, the sample targeted a specific group, we applied theory to guide the assessment, the transcripts contained rich information, and we conducted both within-case and cross-case analyses. Against these considerations, we interviewed 16 caregivers and

1
2
3 progressively reviewed debrief notes from the interviews to assess information power. In analysing the
4 transcripts, we concluded that we reached saturation with the 16 interviews and likely could have stopped
5 interviewing after the 12th interview.
6
7

8
9 We purposively recruited the caregivers from eight communities in the WAU district, four of which were
10 slums and four were other urban areas in the district to maximize variation in the sample. Within each
11 community, two caregivers of children ages 6-36 months were selected to capture a breadth of caregiver
12 experiences of caregivers with vaccine eligible children—one whose child was fully vaccinated for age
13 and another whose child had missed at least one scheduled vaccination visit. CHWs supported data
14 collection teams in visiting households to identify and recruit eligible caregivers in the selected
15 communities. Snowball sampling was used as a secondary sampling strategy when the first identified
16 caregiver declined to interview but knew of other caregivers in the community with vaccine-eligible
17 children or when CHWs were only successful in identifying just one eligible caregiver. In this form of
18 snowball sampling, a previously visited household with an eligible child would point data collectors to
19 other households with potentially eligible children (i.e., vaccine eligible children). Data collectors visited
20 such households to screen for eligibility. This process continued until two caregivers of eligible children
21 were successfully recruited and interviewed from a particular community. Interviews were conducted on
22 the same of day of recruitment after obtaining informed consent from the caregiver.
23
24
25
26
27
28
29
30
31
32

33 We recruited data collectors (interviewers and notetakers) who were fluent in English and the
34 predominant local language in the WAU district (Krio). The data collectors had post-secondary
35 educational training in social sciences and were experienced in conducting qualitative data collection in
36 Sierra Leone. Two behavioural scientists trained the facilitators for a week on the assessment protocol.
37 One of the trainers was from Sierra Leone and had experience conducting social science research in Sierra
38 Leone. During the training, the English version of the guide was translated into Krio by locally hired staff
39 together with the facilitators and trainer. The data collectors were trained on how to probe on the spot to
40 obtain additional pertinent information from caregivers. Data collection occurred in August–September
41 2018. All interviews were audio-recorded with permission from participants; they were then transcribed
42 and translated into English by the local team. Interviews lasted about an hour on average and were
43 conducted in the vicinity of the homes of the caregivers. Data collection teams were trained on choosing
44 suitable interview locations to enable caregivers to speak freely. The facilitators conducted debriefing
45 sessions immediately after each interview to make note of key experiences and observations. The
46 debriefing notes were not part of the formal analysis. However, during the field work, the debriefing notes
47 were used to progressively assess data saturation and to identify key insights emerging from the
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 interviews. We used the insights from the debrief notes to develop a preliminary report that was mostly in
4 a descriptive, narrative form. The de-identified preliminary report was shared with the Sierra Leone
5 Ministry of Health and Sanitation. We have previously documented practical lessons learned from
6 implementing the assessment in Sierra Leone.¹¹
7
8
9

10 11 **2.3. Data analysis**

12 Two analysts (one male, one female) read all transcripts and created analytic memos and then analysed
13 the transcripts using both within-case and cross-case analysis. In the within-case analysis, we developed a
14 narrative profile for each caregiver to bring key aspects of their lived experiences to the foreground. In the
15 initial part of the cross-case analysis, each analyst coded three different transcripts (six total), using an
16 inductive approach to identify and interpret meaning units within the text. To gain alternative
17 interpretations of the coded meaning units, three of the coded transcripts were shared with a third
18 qualitative expert, who was not involved in the previous stages of the assessment, for independent ‘blind’
19 coding of the transcripts. Feedback from the third analyst was discussed by the two primary analysts and
20 incorporated into the coding scheme. The analysts used an iterative process to review their codes, discuss
21 their interpretations of the manifest content, and harmonize the initial set of codes that were used for
22 coding the remaining manuscripts. NVivo software (QSR International–2018, Version 12) was used for
23 the final organization and coding of the transcripts. Manifest categories of meaning units were grouped to
24 reflect latent content that was developed into cross-cutting themes via a consultative process. Throughout
25 the process, the analysts exercised reflexivity regarding subjective interpretations and iteratively re-
26 examined the transcripts to identify alternative interpretations until consensus was reached with additional
27 inputs from the co-authors.
28
29
30
31
32
33
34
35
36
37
38
39

40 **2.4. Ethical approval**

41 The assessment was approved by the Sierra Leone Ethics and Scientific Review Committee (SLESRC-
42 17052018), Columbia University Medical Centre Institutional Review Board (IRB-AAAR9031), and the
43 Centre for Global Health at the U.S. Centres for Disease Control and Prevention (CGH-2018-184). All
44 participants provided written informed consent.
45
46
47
48

49 **2.5. Patient and public involvement**

50 No patient involved.
51
52
53
54

55 **3. Results**

1
2
3 All respondents were the biological mothers of the sampled children except for one female guardian. The
4 median age was nine months for the children included in the assessment. At the time of the interviews,
5 half of the children had missed at least one scheduled vaccine dose. Three themes emerged from the
6 interviews around vaccination enablers (Table 1), vaccination barriers (Table 2) and direct
7 recommendation to improve vaccination uptake (Table 3). There were no notable differences in themes
8 between slums and other urban communities.
9
10
11
12
13

14 **3.1. Enablers of childhood vaccination (Table 1)**

15
16 Our analysis identified emotional and practical enablers related to childhood vaccination. Emotional
17 enablers were evident in how caregivers portrayed their parental obligation to their children, wanting to
18 do the right thing for their children's health, and anticipating reciprocal benefits in children's ability to
19 take care of the parents later in life. Practical enablers were the diversity of immunization reminders,
20 information access, information trust, getting fathers more involved, positive experiences with health
21 workers, and post-vaccination information sharing at the community level.
22
23
24
25
26

27 **3.1.1. Parental responsibility**

28 A sense of parental responsibility was a major motivating factor for caregivers to seek vaccination
29 services for their children. Caregivers viewed vaccination as 'doing the right thing' for their children.
30 Even caregivers who had missed scheduled vaccination visits felt responsible for getting their children
31 caught up with their scheduled vaccine doses, which often required deprioritizing other income-
32 generating activities.
33
34
35

36 *"It is my duty to take my baby to the hospital for immunisation. It is my responsibility as [a]*
37 *mother to ensure that my baby completes the rounds of immunisation without defaulting."* –

38
39 Caregiver whose child had missed a scheduled vaccination
40
41
42

43 **3.1.2. Wanting a strong and healthy baby**

44 In addition to the affective responses regarding a sense of duty to the child, appreciating the overall health
45 benefits of vaccination to their children was another major driving force in motivating caregivers to seek
46 vaccination services. Caregivers consistently expressed that immunisation has essential health benefits to
47 the child and that missing scheduled vaccination would 'risk the baby's life.' Moreover, they valued
48 having a 'strong and healthy baby' and felt that completing the vaccination schedule would positively
49 impact the baby's health.
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

“I think [the] vaccine is good for our children. It is important and it helps to build their immune system to keep them strong and healthy; it fights against many things in the body...” – Caregiver whose child was fully vaccinated

3.1.3. Parental anticipation of reciprocal benefit

Perceptions of vaccination benefits went beyond the direct health benefits to children and extended into domains of benefit to the parent. The notion that vaccinated children will be healthier and in turn live longer and be able to take care of their parents later in life emerged as a dimension of vaccination benefit to the parent. The duality of vaccination benefit is likely grounded in the cultural context of parents expecting reciprocal care from their children when the parents can no longer care for themselves.

“It is very difficult for me to be absent for immunisation or not to take my baby to the hospital when sick. Sometimes people think I’m mad but I’m not. I’m trying to bring up my children in a way that they will benefit me when I’m old.” – Caregiver whose child had missed a scheduled vaccination

3.1.4. Diversity of immunisation reminders

Caregivers were exposed to numerous reminders and cues to vaccinate their children, including information from the immunisation cards, health workers, community campaigns, and family members. However, the child’s immunisation card stood out as the most important reminder that caregivers and families relied on to remember the dates of the scheduled vaccination visits.

“I do remind myself because they [health workers] will write at the back of the immunisation card the date for the next visit. The immunisation card will tell you the date for the next visit.” – Caregiver whose child had missed a scheduled vaccination

In addition, community outreach campaigns and announcements in the community were also viewed as helpful reminders to vaccinate.

“The health workers in this community and the CHWs are doing well as they do go round telling people not to forget to take their babies to the hospital for immunisation.” – Caregiver whose child had missed a scheduled vaccination

3.1.5. Information access and information trust

Immunisation information sources varied, but caregivers consistently cited health workers as trusted sources of information. Nurses were more trusted than lay CHWs because caregivers viewed nurses as

1
2
3 more knowledgeable. While waiting in line before immunisation services, most caregivers appreciated the
4 ‘health talk’ they received from nurses who advised on health and immunisation.
5

6 *“I trust them [nurses] because they are a team of qualified nurses ... Before they come here, they*
7 *know everything about the vaccines and any implications of the vaccines. They are able to*
8 *explain more than the CHWs and other community workers in the area. The nurses will tell you*
9 *more. There are things that the community workers do not know, and they refer you to the*
10 *nurses.”* – Caregiver whose child was fully vaccinated
11
12
13
14
15

16 **3.1.6. Getting fathers more involved**

17 Mothers desired greater involvement by the child’s fathers in supporting their children’s vaccination. In
18 most instances, fathers were rarely involved in taking their children to the vaccination clinic. We
19 uncovered only two instances when fathers actively supported their children’s vaccination visits. In those
20 instances, the mothers felt supported, and their children were fully vaccinated. In one situation, a father
21 that routinely accompanied the child to the vaccination clinic was celebrated by health workers and given
22 the ‘best father’ award.
23
24
25

26 *“There was [a] time when he [my husband] was given the best father award [at the clinic]*
27 *because he is always with me at the hospital. That is the name I have also given to my husband.*
28 *Even when the baby is crying, I will say ‘best father’ take your baby.”* – Caregiver whose child
29 was fully vaccinated
30
31
32
33
34

35 **3.1.7. Positive experiences with health workers**

36 During vaccination visits, positive experiences with health workers encouraged caregivers to vaccinate
37 their children. Positive experiences included having a cordial relationship with the health workers. In
38 particular, caregivers expressed that the nurses took good care of their children, made them feel
39 comfortable, and tried to build a good rapport. Some caregivers said they would voluntarily give small
40 amounts of money to health workers as a token of their appreciation after vaccinating the child.
41
42
43

44 *“...the nurses do encourage you and will make jokes so that you will laugh at the end of the day.*
45 *There is a lot of fun [interactions], which made some of us forget about our stresses.”* –
46 Caregiver whose child was fully vaccinated
47
48
49
50

51 **3.1.8. Post-vaccination information sharing**

52 Information exchange at the community level with trusted community members strengthened caregiver
53 confidence in childhood vaccination. Upon returning home from the vaccination visit, caregivers often
54 discussed the clinic experience with their spouses, families, and other caregivers in the community. In
55
56
57

1
2
3 addition to information obtained from health workers at the clinic, caregivers also sought advice from
4 other ‘more experienced’ caregivers in the community.

5
6 *“Most times after immunisation, my baby will run a temperature, but the health workers always*
7 *provide drugs to counter the fever. We have caregivers in this community with vast knowledge*
8 *and since this is my first baby, I love to talk to them so that we can share our experiences which*
9 *will be of advantage to me as I’m very new in the field [of parenting].” –Caregiver whose child*
10 *was fully vaccinated*
11
12
13

14 15 16 **3.2. Theme 2. Barriers related to childhood vaccination (Table 2)**

17 Practical constraints, negative experiences with health workers, and safety concerns were the underlying
18 barriers to childhood vaccination. Practical constraints included challenges faced when preparing for and
19 getting to the vaccination clinic and inconveniences encountered at the clinic, such as overcrowding and
20 long waiting times. Negative experiences among caregivers included feeling disrespected by health
21 workers while simultaneously expecting to give money to health workers for services that are supposed to
22 be free of charge. Finally, vaccine side effects led to concerns and fears about vaccine safety.
23
24
25
26
27

28 29 **3.2.1. Preparing for the journey and getting to the clinic**

30 Caregivers commonly cited the need to juggle ‘household duties’ and other income-generating activities
31 when planning the visit as a barrier, especially in the absence of fathers’ involvement in taking the child
32 to the clinic. In addition, mothers frequently depended on their children’s fathers for financial support to
33 cover the expenses related to the vaccination clinic visit. Some caregivers recounted needing to travel
34 long distances up to an hour by foot to get to the vaccination clinic, especially when they could not afford
35 public transportation.
36
37
38
39
40

41 42 **3.2.2. Inconveniences at the clinic**

43 Caregivers anticipated various inconveniences at the vaccination visit. The prolonged time spent waiting
44 for the child to be vaccinated emerged as a substantial inconvenience that was more pronounced when
45 seeking immunisation services, especially in larger facilities. Anticipating the long wait, caregivers
46 usually tried to arrive early at the vaccination site to get seen first. The range of activities involved with
47 the child health visit prolonged the visit, including weighing the baby as part of growth monitoring and
48 other health checks.
49
50
51

52 *“It is painful if you waste much time at the health facility because you have other issues to attend*
53 *to. To avoid that, that is why I always come early to the health facility.” – Caregiver whose child*
54 *was fully vaccinated*
55
56
57

3.2.3. Feeling disrespected by health workers

Caregivers often felt disrespected by health workers during the vaccination visit. A key complaint was that health workers shouted at caregivers and sometimes used vulgar language toward caregivers. In other instances, they complained that some health workers habitually arrived late to the vaccination session, which further prolonged the time caregivers spent waiting.

3.2.4. Monetary expectations

Systemically hidden costs generated substantial dissatisfaction among caregivers. Caregivers needed to “shake hands” with health workers at different times of the visit (e.g., first time registering the child to get a card, before entering the facility, and before weighing the baby). Shaking hands implied giving some money during the handshake. Caregivers used the money to ‘fast-track’ their children’s vaccination. The expectations around monetary exchange discouraged caregivers who could not afford to shake hands with health workers.

“Sometimes if I don’t want to spend much time at the hospital, I will shake the hand of the nurse so that they can fast track the immunisation of my baby. I will give them something like two thousand Leones or whatever I have with me at that moment... Health is wealth and they [health workers] don’t need us but we do [need them]. The money we give is nothing compared to the health of our children... At the end of the day, we will grumble on our way home as the services are supposed to be free for our children, yet we are paying for it. The health workers are really trying, but the idea for them to take money from us is bad. And if you don’t give them money, they will talk to you carelessly.” – Caregiver whose child was fully vaccinated

In a separate domain of monetary exchange, health workers demanded money as a form of ‘punishment’ to caregivers who missed their children’s scheduled vaccination appointments.

“If you failed to take your baby to the hospital on [the] stipulated date, you will definitely have to pay some amount at the end of the day in the form of punishment. You must pay five thousand or more.” – Caregiver whose child was fully vaccinated

3.2.5. Vaccine side effects

Caregivers cited numerous instances when their children experienced vaccine side effects such as ‘fever,’ ‘swelling at the injection site,’ and the ‘baby becoming lethargic.’ Fever was the most common side effect, and the caregivers knew to administer fever-reducing medication as instructed by health workers.

1
2
3 When there was swelling at the injection site, a common practice among caregivers was to massage the
4 swollen area, sometimes with an onion or a bar of soap to try to reduce the swelling.
5

6 *“Sometimes my baby’s leg becomes swollen... because some nurses are heavy-handed, and I meet*
7 *several nurses when I visit the hospital. Sometimes the leg gets swollen, and they treat him. I have*
8 *to rub the leg to avoid swelling... I use soap to rub off the swelling and I give Panadol to stop the*
9 *fever... some people say you should not allow every nurse to administer [an] injection to the child.*
10 *I should have a permanent nurse that gives injection to my child without swelling.” – Caregiver*
11 *whose child missed a scheduled vaccination*
12
13
14
15

16
17 In one rare situation, a caregiver had a prior child who experienced fever, convulsed, and died a few days
18 after getting vaccinated. Therefore, the caregiver decided to not vaccinate subsequent children.
19

20 *“After the immunisation, my baby started running temperature, I administered paracetamol as I*
21 *was told by the health workers. The baby convulsed and that was the end of that baby. I don’t*
22 *want a repeat of that in my life. I have therefore decided not to take my babies for immunisation*
23 *anymore” – Caregiver whose child missed a scheduled vaccination*
24
25
26
27

28 Besides passive acceptance of the BCG vaccine at the birth of the youngest child, this same caregiver
29 actively refused all other vaccines despite encouragement by a family member to vaccinate the child.
30

31 *“I’m not outrightly saying it was as a result of the immunisation [that my child died]; as every*
32 *death is the work of God. But from what I have gathered so far, I have personally decided not to*
33 *take my baby to the health facility to be immunized. It is not that I’m tired of going to the health*
34 *facility or because of the distance or money. I do get a lot of pressure from my aunt to take my*
35 *baby to the health facility for immunisation, but the thing is that I just don’t trust the system and*
36 *what immunisation does. – Caregiver whose child missed a scheduled vaccination*
37
38
39
40
41

42 From the perspective of this same caregiver, the vaccinated child died but the unvaccinated children
43 survived and thrived, which was cited as a reason for refusing vaccination.
44

45 *“I believe in exclusive breastfeeding, sometimes for two years and a half and sometimes [only]*
46 *two years. My baby is now two years seven months old and doing well like any of those children*
47 *that are on immunisation or have completed...Just as I was saying, sometimes my heart will tell*
48 *me to take the baby for immunisation but after thinking of the past experience, I would decide not*
49 *to. I’m now used to that... The simple fact here is that, since the other children are doing well*
50 *without immunisation, I will not take [the baby] to the health facility for immunisation and that is*
51 *all.” – Caregiver whose child missed a scheduled vaccination*
52
53
54
55
56
57

3.3. Recommendations to improve childhood vaccination (Table 3)

The direct recommendations provided by caregivers were categorized into (1) improving vaccination process and systems and (2) engaging communities to boost vaccine confidence. Implementing community outreach campaigns for immunisation at regular intervals with a focus on defaulters was recommended by participants to improve vaccination outcomes. In addition, caregivers wanted health workers and community leaders to be involved in immunisation promotion along with the CHWs. They wanted the vaccination clinic experience to improve and become more conducive to caregivers, including shorter wait time at the clinic and more positive interactions with health workers. Lastly, caregivers wanted health workers to stop demanding money from them, though they may not mind giving money, out of free-will, as a token of appreciation when they could afford it.

“In addition, you should engage the Chiefs, because in each area we have Chiefs to spread out this message. You could educate them so that they in turn can educate those in the community. Let us have Town Criers go around disseminating the messages. It would be nice for them to allocate people in the health centre who move from house to house to educate the breastfeeding mothers because some of us are stubborn to come onboard.” – Caregiver whose child was fully vaccinated

4. Discussion

Our qualitative analysis highlighted several important themes. In the backdrop of anticipated benefits to both the child and parent, vaccination intention was motivated by a feeling of parental responsibility to ‘do the right thing.’ Timely and trusted exchange of information together with social support and positive experiences at the vaccination clinic were important facilitators of vaccination. In contrast, vaccination was discouraged by negative interactions with health workers at the clinic, the occurrence and fear of vaccine side effects, multitude of ‘hidden’ costs, juggling vaccination with other responsibilities, and inconveniences, such as long traveling time to the clinic and long delays at the clinic. Nevertheless, caregivers were resilient in devising ways to try to get their children vaccinated such as walking on foot up to an hour to get to the clinic when they could not afford public transportation. Lastly, caregivers wanted the vaccination experience to improve, and they desired stronger community engagement to help optimize vaccination outcomes. However, systemic issues, such as informal payments, overcrowding in health facilities, and the reported overburdened health workers may require interventions at the health systems level. These themes from Sierra Leone provide in-depth insights regarding the motivations, facilitators, and barriers of vaccination in an urban LMIC setting.

1
2
3
4
5 Moral values may shape vaccination attitudes.²² Philosophical arguments regarding the morality of
6 vaccination have been heavily debated.²³⁻²⁶ Caregivers in our sample largely viewed vaccination via a
7 moral lens encompassing parental duty to do the right thing for the child. In one situation, however, we
8 found that the desire to ‘do the right thing’ may also translate into vaccination refusal in the backdrop of
9 other past refusals, observing ‘healthy unvaccinated’ children, and having distrust of the health system.
10 Quantitative research from high-income countries has shown that parents with unvaccinated children were
11 more likely to perceive their children to be at low risk of vaccine-preventable diseases and were more
12 likely to perceive low vaccine effectiveness and safety compared to parents with vaccinated children.²⁷
13 Our findings suggest that childhood immunisation communication efforts may resonate more strongly
14 with caregivers when vaccination is framed around parental responsibilities to do the right thing for the
15 child and the anticipated future benefits to parents. However, additional research is necessary to generate
16 a better understanding of the morality of childhood vaccination in the Sierra Leonean context.
17
18
19
20
21
22
23
24

25 Across the interviews, there was an apparent tension in the relationship between caregivers and health
26 workers. Caregivers often expressed their appreciation of health workers and empathized with the
27 challenging context in which they do their work. Health workers were strongly viewed as authoritative
28 sources of trusted information regarding immunisation and the child’s health, which is consistent with
29 findings from high-income countries²⁸⁻³⁰ and LMICs.^{31 32} Our findings on the role of monetary exchange
30 in vaccination exemplify the complex relationship between caregivers and health workers in low-resource
31 urban communities in Sierra Leone. Some caregivers voluntarily gave money to health workers as a
32 ‘token of appreciation’ while others begrudgingly gave money because they viewed it as a condition for
33 receiving good quality service from health workers. Interventions at the health systems level are
34 necessary to help discourage informal payments to health workers—a practice that may perpetuate
35 vaccination inequities among poor caregivers who are unable to meet monetary expectations.
36
37
38
39
40
41
42
43

44 Our findings also illuminate the need for interventions at the household and family level. Fathers were
45 rarely involved in taking their children to the vaccination clinic but were often engaged in the decision-
46 making processes. In the few instances when fathers were involved in taking their children to the clinic,
47 the mothers felt supported, and their children were fully vaccinated. A study in Nigeria found that
48 paternal involvement in immunisation was greater in rural settings compared to urban settings.³³ In urban
49 areas, the same study found that paternal involvement was greater among educated fathers compared to
50 uneducated fathers. In a separate study in Ghana, the involvement of educated fathers in the vaccination
51 decision was associated with timelier vaccination uptake compared to the involvement of uneducated
52
53
54
55
56
57
58
59
60

fathers in the decision.³⁴ More broadly, shifting from a mother-child dyad to a family triad in the care of children has proven to have positive effects on paediatric health outcomes across diverse contexts.³⁵ Additional assessments and interventions are needed to explore and evaluate culturally appropriate ways to enhance the involvement of fathers in childhood immunisation in Sierra Leone and other similar LMIC settings.

Existing evidence suggests that vaccine safety concerns, often linked to adverse events following immunisation (AEFI), contribute to vaccine hesitancy.^{36 37} Serious AEFIs may be ‘triggering events’ for derailing vaccine confidence and prompting active refusal among certain caregivers and their communities—especially when the serious AEFI is perceived to be linked to the vaccine or the vaccination process.³⁸ Together with prior evidence, these findings emphasize the need for robust AEFI surveillance and investigations³⁹ to identify, counsel, and follow-up with caregivers whose children experience AEFI, and therefore, are potentially at risk of missing subsequent vaccination. Health workers and CHWs may benefit from periodic in-service training on how to effectively communicate vaccine safety and address concerns about AEFIs.⁴⁰

4.1. Limitations

There were several limitations to our assessment. First, it is possible that some nuanced meaning may have been lost when translating the audio recordings from Krio to English—especially since the transcripts were not back-translated from English to Krio due to resource constraints. Second, we only identified one caregiver who routinely and actively refused all vaccines for her child after experiencing a serious AEFI, which may reflect the overall rarity of zero-dose unvaccinated children in Sierra Leone (approximately 3%).⁴¹ This was the only caregiver with a child that experienced a serious AEFI in our sample, which limits our ability to have a rich understanding of such experience among caregivers more broadly and the potential linkages to vaccination refusal. Although such experiences of serious AEFI are rare, they may have the tendency to get publicized in the community, which may negatively influence vaccination decisions among other caregivers. Taken together, our results call for additional qualitative assessments to get a deeper understanding of vaccination refusal within the Sierra Leonean context and other low-resource LMIC settings. Sampling strategies may therefore need to be adapted accordingly to focus on caregivers who actively refuse vaccination for their children. Lastly, our findings reflected gender dimensions that may be based on sociocultural norms in Sierra Leonean society but may also be due to sampling bias because the caregivers we conveniently recruited were mostly stay-at-home mothers.

5. Conclusions

1
2
3 As the COVID-19 pandemic disrupts childhood immunisation globally,⁴² especially in LMICs, our
4 assessment provides a foundational understanding of the challenges that caregivers encounter in urban
5 settings in Sierra Leone. It also sheds light on opportunities to improve vaccination outcomes in urban
6 poor settings, which is a global immunisation priority. The findings show that health system
7 interventions, community engagement, and vaccination outreach may need to be tailored to for urban
8 settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is
9 framed both around parental responsibilities to do the right thing for the child and the future benefits to
10 the parent.
11
12
13
14
15

16 17 **Competing interests**

18 None declared.
19
20
21

22 **Data availability**

23 All data relevant to the study are included in the article or uploaded as supplementary information.
24
25
26

27 **Contributors**

28 MFJ, RS, MT, TS, and ML designed the assessment. MFJ, PP, SK, and KW analysed the data. MJF, PP,
29 RS, SK, MT, KW, TS, and ML contributed to the interpretation of the results. MFJ, PP and SK drafted
30 the manuscript. RS, MT, KW, TS, and ML contributed to revising the manuscript. All authors reviewed
31 and approved the final manuscript.
32
33
34
35

36 **Collaborators**

37 Adewale Akinjeji, Laura Conklin, Oliver Eleeza, Lauren E Parmley, Anthony Mansaray, Dimitri
38 Prybylski, Roberta Sutton, Mame Toure, Aaron Wallace, Brent Wolff
39
40
41
42

43 **Acknowledgements**

44 We thank the 16 caregivers who participated in the in-depth interviews.
45
46
47

48 **Funding**

49 U.S. Centers for Disease Control and Prevention (U19GGH001581).
50
51
52

53 **Disclaimer**

54
55
56
57
58
59
60

The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the US Centres for Disease Control and Prevention.

6. References

1. Nelson KN, Wallace AS, Sodha SV, et al. Assessing strategies for increasing urban routine immunization coverage of childhood vaccines in low and middle-income countries: A systematic review of peer-reviewed literature. *Vaccine* 2016;34(46):5495-503. doi: 10.1016/j.vaccine.2016.09.038 [published Online First: 2016/10/25]
2. JSI. Strengthening Immunization for the Urban Poor: Experiences and Lessons Learned from Five Country Studies 2018. Available from: <https://www.jsi.com/resource/strengthening-immunization-for-the-urban-poor-experiences-and-lessons-learned-from-five-country-studies/> [accessed November 12 2020].
3. Brolin Ribacke KJ, Saulnier DD, Eriksson A, et al. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Frontiers in public health* 2016;4:222. doi: 10.3389/fpubh.2016.00222 [published Online First: 2016/10/26]
4. Elston JW, Cartwright C, Ndumbi P, et al. The health impact of the 2014-15 Ebola outbreak. *Public health* 2017;143:60-70. doi: 10.1016/j.puhe.2016.10.020 [published Online First: 2017/02/06]
5. Yerger P, Jalloh M, Coltart CEM, et al. Barriers to maternal health services during the Ebola outbreak in three West African countries: a literature review. *BMJ Glob Health* 2020;5(9) doi: 10.1136/bmjgh-2020-002974 [published Online First: 2020/09/09]
6. Masresha BG, Luce R, Jr., Weldegebriel G, et al. The impact of a prolonged ebola outbreak on measles elimination activities in Guinea, Liberia and Sierra Leone, 2014-2015. *Pan Afr Med J* 2020;35(Suppl 1):8. doi: 10.11604/pamj.supp.2020.35.1.19059 [published Online First: 2020/05/07]
7. GoSL. National Ebola Recovery Strategy for Sierra Leone 2015 [Available from: https://ebolaresponse.un.org/sites/default/files/sierra_leone_-_national_recovery_strategy_2015-2017.pdf accessed December 5 2020.
8. CDC. Global Health – Sierra Leone 2019 [Available from: <https://www.cdc.gov/globalhealth/countries/sierra-leone/default.htm> accessed December 5 2020.
9. Feldstein LR, Sutton R, Jalloh MF, et al. Access, demand, and utilization of childhood immunization services: A cross-sectional household survey in Western Area Urban district, Sierra Leone, 2019. *Journal of global health* 2020;10(1):010420. doi: 10.7189/jogh.10.010420 [published Online First: 2020/06/09]
10. UNICEF. Human Centered Design 4 Health 2018 [Available from: <https://www.hcd4health.org/> accessed November 18 2020.
11. Jalloh MF, Kinsman J, Conteh J, et al. Barriers and facilitators to reporting deaths following Ebola surveillance in Sierra Leone: implications for sustainable mortality surveillance based on an exploratory qualitative assessment. *BMJ Open* 2021;11(5):e042976. doi: 10.1136/bmjopen-2020-042976 [published Online First: 2021/05/15]
12. Diekelmann NL. Learning-as-testing: a Heideggerian hermeneutical analysis of the lived experiences of students and teachers in nursing. *ANS Advances in nursing science* 1992;14(3):72-83. doi: 10.1097/00012272-199203000-00010 [published Online First: 1992/03/01]
13. Dreyfus HL. Being-in-the-world: A commentary on Heidegger's Being and Time. Cambridge: The MIT Press 1991.
14. Dahlberg K. The essence of essences – the search for meaning structures in phenomenological analysis of lifeworld phenomena. *International journal of qualitative studies on health and well-being* 2006;1(1):11-19. doi: 10.1080/17482620500478405

15. StatisticsSierraLeone. 2015 Sierra Leone Population and Housing Census 2016 [Available from: <https://www.statistics.sl/index.php/census/census-2015.html>] accessed November 12 2020.
16. Lamunu M, Olu OO, Bangura J, et al. Epidemiology of Ebola Virus Disease in the Western Area Region of Sierra Leone, 2014-2015. *Frontiers in public health* 2017;5:33. doi: 10.3389/fpubh.2017.00033 [published Online First: 2017/03/18]
17. WHO. Medical doctors per 10,000 population 2021 [Available from: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/medical-doctors-\(per-10-000-population\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/medical-doctors-(per-10-000-population))] accessed December 5 2021.
18. Witter S, Brikci N, Harris T, et al. The free healthcare initiative in Sierra Leone: Evaluating a health system reform, 2010-2015. *The International journal of health planning and management* 2018;33(2):434-48. doi: 10.1002/hpm.2484 [published Online First: 2018/01/13]
19. Government of Sierra Leone. Comprehensive EPI Multi-Year Plan 2017-2021 2016 [Available from: <https://www.gavi.org/sites/default/files/document/2021/cMYP%20Sierra%20Leone%202017-2021.pdf>] accessed December 5 2020.
20. Government of Sierra Leone. Policy for Community Health Workers in Sierra Leone 2012 [Available from: <https://chwcentral.org/wp-content/uploads/2015/01/CHW-Policy-Sierra-Leone.pdf>] accessed December 5 2020.
21. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided by Information Power. *Qualitative health research* 2016;26(13):1753-60. doi: 10.1177/1049732315617444 [published Online First: 2015/11/29]
22. Amin AB, Bednarczyk RA, Ray CE, et al. Association of moral values with vaccine hesitancy. *Nature human behaviour* 2017;1(12):873-80. doi: 10.1038/s41562-017-0256-5 [published Online First: 2017/12/01]
23. Di Pietro ML, Poscia A, Telesman AA, et al. Vaccine hesitancy: parental, professional and public responsibility. *Annali dell'Istituto superiore di sanita* 2017;53(2):157-62. doi: 10.4415/ann_17_02_13 [published Online First: 2017/06/16]
24. Giubilini A, Douglas T, Šavulescu J. The moral obligation to be vaccinated: utilitarianism, contractualism, and collective easy rescue. *Medicine, health care, and philosophy* 2018;21(4):547-60. doi: 10.1007/s11019-018-9829-y [published Online First: 2018/02/13]
25. Bester JC. Measles Vaccination is Best for Children: The Argument for Relying on Herd Immunity Fails. *Journal of bioethical inquiry* 2017;14(3):375-84. doi: 10.1007/s11673-017-9799-4 [published Online First: 2017/08/18]
26. Dawson A. What are the moral obligations of the traveller in relation to vaccination? *Travel medicine and infectious disease* 2007;5(2):90-6. doi: 10.1016/j.tmaid.2006.01.005 [published Online First: 2007/02/15]
27. Omer SB, Salmon DA, Orenstein WA, et al. Vaccine refusal, mandatory immunization, and the risks of vaccine-preventable diseases. *N Engl J Med* 2009;360(19):1981-8. doi: 10.1056/NEJMsa0806477 [published Online First: 2009/05/08]
28. Frew PM, Chung Y, Fisher AK, et al. Parental experiences with vaccine information statements: Implications for timing, delivery, and parent-provider immunization communication. *Vaccine* 2016;34(48):5840-44. doi: 10.1016/j.vaccine.2016.10.026 [published Online First: 2016/10/30]
29. Drezner D, Youngster M, Klainer H, et al. Maternal vaccinations coverage and reasons for non-compliance - a cross-sectional observational study. *BMC pregnancy and childbirth* 2020;20(1):541. doi: 10.1186/s12884-020-03243-w [published Online First: 2020/09/18]
30. Wiley KE, Massey PD, Cooper SC, et al. Pregnant women's intention to take up a post-partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: a cross sectional survey. *Vaccine* 2013;31(37):3972-8. doi: 10.1016/j.vaccine.2013.06.015 [published Online First: 2013/06/20]
31. Makwe CC, Anorlu RI. Knowledge of and attitude toward human papillomavirus infection and vaccines among female nurses at a tertiary hospital in Nigeria. *International journal of women's health* 2011;3:313-7. doi: 10.2147/ijwh.S22792 [published Online First: 2011/10/07]

- 1
2
3 32. Foss HS, Oldervoll A, Fretheim A, et al. Communication around HPV vaccination for adolescents in
4 low- and middle-income countries: a systematic scoping overview of systematic reviews.
5 *Systematic reviews* 2019;8(1):190. doi: 10.1186/s13643-019-1100-y [published Online First:
6 2019/08/03]
7
8 33. Sodeinde K, Amoran O, Abiodun O, et al. A rural/urban comparison of paternal involvement in
9 childhood immunisation in Ogun Central Senatorial District, Nigeria. *The Nigerian postgraduate
10 medical journal* 2020;27(4):336-42. doi: 10.4103/npmj.npmj_101_20 [published Online First:
11 2020/11/07]
12
13 34. Brugha RF, Kevany JP, Swan AV. An investigation of the role of fathers in immunization uptake. *Int
14 J Epidemiol* 1996;25(4):840-5. doi: 10.1093/ije/25.4.840 [published Online First: 1996/08/01]
15
16 35. Allport BS, Johnson S, Aqil A, et al. Promoting Father Involvement for Child and Family Health.
17 *Acad Pediatr* 2018;18(7):746-53. doi: 10.1016/j.acap.2018.03.011 [published Online First:
18 2018/04/14]
19
20 36. Dube E, Laberge C, Guay M, et al. Vaccine hesitancy: an overview. *Human vaccines &
21 immunotherapeutics* 2013;9(8):1763-73. doi: 10.4161/hv.24657 [published Online First:
22 2013/04/16]
23
24 37. Larson HJ, Jarrett C, Eckersberger E, et al. Understanding vaccine hesitancy around vaccines and
25 vaccination from a global perspective: a systematic review of published literature, 2007-2012.
26 *Vaccine* 2014;32(19):2150-9. doi: 10.1016/j.vaccine.2014.01.081 [published Online First:
27 2014/03/07]
28
29 38. Du F, Chantler T, Francis MR, et al. The determinants of vaccine hesitancy in China: A cross-
30 sectional study following the Changchun Changsheng vaccine incident. *Vaccine*
31 2020;38(47):7464-71. doi: 10.1016/j.vaccine.2020.09.075 [published Online First: 2020/10/13]
32
33 39. WHO. Adverse events following immunization (AEFI) 2021 [Available from:
34 https://www.who.int/immunization/monitoring_surveillance/who-immuniz.pdf accessed February
35 18 2021.
36
37 40. WHO. Vaccine safety basics e-learning course 2021 [Available from: [https://vaccine-safety-
38 training.org/simplified-and-key-messages.html](https://vaccine-safety-training.org/simplified-and-key-messages.html) accessed February 19 2021.
39
40 41. UNICEF. Sierra Leone Multiple Indicators Cluster Survey 2017 [Available from:
41 https://mics.unicef.org/news_entries/106/SIERRA-LEONE-2017-MICS-RELEASED accessed
42 January 21 2021.
43
44 42. Abbas K, Procter SR, van Zandvoort K, et al. Routine childhood immunisation during the COVID-19
45 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2
46 infection. *Lancet Glob Health* 2020;8(10):e1264-e72. doi: 10.1016/s2214-109x(20)30308-9
47 [published Online First: 2020/07/21]
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Enablers of childhood vaccination—Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-themes	Theme
Responsibility to ensure full vaccination	Parental responsibility	Emotional enablers of childhood vaccination	Enablers of childhood vaccination
Obligation to do the right thing			
Immunisation as a requirement			
Immunisation is important for baby's health	Wanting a 'strong and healthy baby'		
Wanting 'strong baby'			
'Health is wealth'			
Defaulting on vaccination risks baby's life	Parental anticipation of reciprocal benefit		
Vaccination benefits the parent later in life			
Taking care of parents when old	Diversity of immunisation reminders	Practical enablers of childhood vaccination	
Immunisation card is important			
Campaign as reminder			
Reminders by health workers at vaccination visit			
Husband as reminder			
Immunisation card as reminder			
Other family member as reminder			
Nurses should lead			
Nurses more trusted than CHWs			
Same information from different sources			
Immunisation promotion through radio/tv	Information access and information trust		
Immunisation promotion by health workers			
Immunisation promotion through leaders			
Immunisation promotion by NGOs	Getting fathers more involved		
Mothers take the child to the clinic			
Fathers rarely involved			
Father received an award for involvement	Positive experiences with health worker		
Cordial relationship with nurses			
Good care by nurses			
HWs encourage seeking care at HF			
Giving money to health worker as token of appreciation	Post-vaccination information sharing		
Husband asking about the visit			
Informing husband of next visit			
Telling husband about visit expenses			
Immunisation is a "learning process"			
Sharing experiences with neighbours or friends			
Other family members asking about the visit			

Table 2. Barriers related to childhood vaccination–Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-themes	Themes
Asking husband for money	Preparing for the journey and getting to the clinic	Practical constraints	Barriers related to childhood vaccination
Juggling different household duties			
Competing priorities			
Time taken to get to the clinic			
Wasting time at the clinic			
Rush to arrive first and seen first			
Crowding at the clinic			
Very long wait at larger health facilities			
Spent a day and nothing happened			
Wasting caregiver's time	Inconveniences at the clinic	Negative experiences with health workers	
Not respecting caregivers			
Shouting at caregivers			
Paying for the card			
Paying for free drugs			
Paying for weighing			
Payment as punishment			
Bad care without payment			
Health workers should stop demanding money			
Health worker don't appreciate less than Le 2000	Vaccine side effects	Safety concerns	
Health workers withholding free drugs			
Baby crying throughout the night			
Baby gets 'lazy' for few hours			
Fever in baby			
Swelling at injection site			
Side effects only for some vaccines			
Rub onion on swollen injection site			
Avoiding abnormalities from vaccine			
Afraid of vaccine side effects			

Table 3. Recommendations to improve childhood vaccination– Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-theme	Theme
Continue campaigns at repeated intervals	Improving vaccination processes and systems	Caregivers want improved vaccination processes, systems, and engagement	Recommendations to improve childhood vaccination
Do not rely on campaigns alone			
Provision of incentives for caregivers			
Concentrate on defaulters			
Compensation/incentives for nurses			
Being considerate towards health workers			
Employ more staff			
Stop demanding money	Engaging communities to		
Promote consequences of not vaccinating			
Peer-to-peer promotion of immunisation			

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Inform about importance of immunisation	boost vaccine confidence		
Address ‘stubborn’ caregivers			
Engage caregivers who do not want injections			

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2021-058203 on 9 May 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

COREQ (CONsolidated criteria for REporting Qualitative research) Checklist

A checklist of items that should be included in reports of qualitative research. You must report the page number in your manuscript where you consider each of the items listed in this checklist. If you have not included this information, either revise your manuscript accordingly before submitting or note N/A.

Topic	Item No.	Guide Questions/Description	Reported on Page No.
Domain 1: Research team and reflexivity			
<i>Personal characteristics</i>			
Interviewer/facilitator	1	Which author/s conducted the interview or focus group?	
Credentials	2	What were the researcher's credentials? E.g. PhD, MD	
Occupation	3	What was their occupation at the time of the study?	
Gender	4	Was the researcher male or female?	
Experience and training	5	What experience or training did the researcher have?	
<i>Relationship with participants</i>			
Relationship established	6	Was a relationship established prior to study commencement?	
Participant knowledge of the interviewer	7	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	
Interviewer characteristics	8	What characteristics were reported about the interviewer/facilitator? e.g. Bias, assumptions, reasons and interests in the research topic	
Domain 2: Study design			
<i>Theoretical framework</i>			
Methodological orientation and Theory	9	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	
<i>Participant selection</i>			
Sampling	10	How were participants selected? e.g. purposive, convenience, consecutive, snowball	
Method of approach	11	How were participants approached? e.g. face-to-face, telephone, mail, email	
Sample size	12	How many participants were in the study?	
Non-participation	13	How many people refused to participate or dropped out? Reasons?	
<i>Setting</i>			
Setting of data collection	14	Where was the data collected? e.g. home, clinic, workplace	
Presence of non-participants	15	Was anyone else present besides the participants and researchers?	
Description of sample	16	What are the important characteristics of the sample? e.g. demographic data, date	
<i>Data collection</i>			
Interview guide	17	Were questions, prompts, guides provided by the authors? Was it pilot tested?	
Repeat interviews	18	Were repeat interviews carried out? If yes, how many?	
Audio/visual recording	19	Did the research use audio or visual recording to collect the data?	
Field notes	20	Were field notes made during and/or after the interview or focus group?	
Duration	21	What was the duration of the interviews or focus group?	
Data saturation	22	Was data saturation discussed?	
Transcripts returned	23	Were transcripts returned to participants for comment and/or	

Topic	Item No.	Guide Questions/Description	Reported on Page No.
		correction?	
Domain 3: analysis and findings			
<i>Data analysis</i>			
Number of data coders	24	How many data coders coded the data?	
Description of the coding tree	25	Did authors provide a description of the coding tree?	
Derivation of themes	26	Were themes identified in advance or derived from the data?	
Software	27	What software, if applicable, was used to manage the data?	
Participant checking	28	Did participants provide feedback on the findings?	
<i>Reporting</i>			
Quotations presented	29	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	
Data and findings consistent	30	Was there consistency between the data presented and the findings?	
Clarity of major themes	31	Were major themes clearly presented in the findings?	
Clarity of minor themes	32	Is there a description of diverse cases or discussion of minor themes?	

Developed from: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

Once you have completed this checklist, please save a copy and upload it as part of your submission. DO NOT include this checklist as part of the main manuscript document. It must be uploaded as a separate file.

BMJ Open

Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-058203.R2
Article Type:	Original research
Date Submitted by the Author:	30-Mar-2022
Complete List of Authors:	Jalloh, Mohamed; Centers for Disease Control and Prevention, Center for Public Health Patel, Palak; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Sutton, Roberta; ICAP at Columbia University Kulkarni, Shibani; Centers for Disease Control and Prevention; Oak Ridge Institute for Science and Education, Toure, Mame; ICAP at Columbia University Wiley, Kerrie; The University of Sydney, School of Public Health Sessay, Tom; Sierra Leone Ministry of Health and Sanitation Lahuerta, Maria; ICAP at Columbia University; Columbia University Mailman School of Public Health, Department of Epidemiology
Primary Subject Heading:	Public health
Secondary Subject Heading:	Qualitative research, Public health, Health services research, Global health
Keywords:	Public health < INFECTIOUS DISEASES, Community child health < PAEDIATRICS, Child protection < PAEDIATRICS

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

TITLE PAGE

Title

Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Authors

Mohamed F. Jalloh¹, Palak Patel^{1,2}, Roberta Sutton³, Shibani Kulkarni^{1,2}, Mame Toure⁴, Kerrie Wiley⁵, Tom Sesay,⁶ Maria Lahuerta^{3,7}, Sierra Leone Urban Immunisation Needs Assessment Group

Affiliations

1. Immunisation Systems Branch, Global Immunisation Division, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
2. Oak Ridge Institute for Science and Education's Research Participation Program, Centres for Disease Control and Prevention, Atlanta, Georgia, USA
3. ICAP at Columbia University, Mailman School of Public Health, New York, New York, USA
4. ICAP at Columbia University, Sierra Leone Country Office, Freetown, Sierra Leone
5. School of Public Health, The University of Sydney, Sydney, NSW, Australia
6. Child Health Programme, Ministry of Health and Sanitation, Freetown, Sierra Leone
7. Department of Epidemiology, Mailman School of Public Health, New York, New York, USA

Corresponding author

Mohamed F. Jalloh, PhD, MPH

Immunisation systems Branch, Global Immunisation Division, U.S. Centres for Disease Control and Prevention, Atlanta, Georgia, USA. Email: yum8@cdc.gov. Tel: +1 (404) 401-2773.

Wordcount

Abstract: 239 words; Manuscript: 5438 words

Keywords

Vaccination, immunisation, vaccine, hesitancy, demand, caregiver, experience, qualitative, methodology, interpretative phenomenology, urban, slum, Sierra Leone

ABSTRACT

Objective: To gain in-depth understanding of the caregiver experience when navigating urban immunisation services for their children.

Design: An exploratory qualitative assessment comprising 16 in-depth interviews using an interpretative phenomenology approach.

Setting: Caregivers were purposively recruited from slums (n=8) and other urban communities (n=8) in the capital city of Sierra Leone.

Participants: Caregivers of children ages 6 to 36 months old who were fully vaccinated (n=8) or under-vaccinated (n=8).

Results: Emotional enablers of vaccination were evident in caregivers' sense of parental obligation to their children while also anticipating reciprocal benefits in children's ability to take care of the parents later in life. Practical enablers were found in the diversity of immunization reminders, information access, information trust, getting fathers more involved, positive experiences with health workers, and post-vaccination information sharing in the community. Underlying barriers to childhood vaccination were due to practical constraints such as overcrowding and long waiting times at the clinic, feeling disrespected by health workers, expecting to give money to health workers for free services, and fear of serious vaccine side effects. To improve vaccination outcomes, caregivers desired more convenient and positive clinic experiences and deeper community engagement.

Conclusions: Health system interventions, community engagement, and vaccination outreach need to be tailored to for urban settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is framed both around parental responsibilities to do the right thing for the child and the future benefits to the parent.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This qualitative assessment describes the caregiver experience in navigating childhood immunisation services in urban settings in a low-income country to gain insights for improving vaccination outcomes.
- The sample comprised of caregivers of fully vaccinated and under-vaccinated children who were purposively recruited to help understand how the two outcomes may be shaped by differences in the caregiver experiences.
- The rarity of experiencing serious side effects among children in our sample limits the ability to have a rich understanding of such experiences on subsequent vaccination uptake.

1. Introduction

There have been efforts to understand urban immunisation challenges in low- and middle-income countries (LMICs), including in urban slums and informal settlements.¹ Assessment of immunisation barriers in urban areas in LMICs identified a range of practical and social issues, such as population mobility, inaccurate denominators of children due to out-of-date population estimates and poorly defined geographic catchment areas, the lack of trust in the health system among vulnerable groups, overburdened health facilities, and weak community engagement and outreach.²

In Sierra Leone, the 2014-2016 Ebola epidemic disrupted the delivery of essential health services, including immunisation services, especially in urban areas.^{3,4} Barriers that affected routine health services included the fear of contracting Ebola in health facilities, stigmatization of health workers, and shifting of resources to the epidemic.⁵ As the Ebola epidemic waned, measles outbreaks became more frequent due to the decline in measles vaccination.⁶ In the aftermath of the Ebola epidemic, the Government of Sierra Leone and its partners made major investments to rebuild health systems and restore public confidence in the health system.^{7,8} However, challenges in access to and the uptake of immunisation services persist, including in urban areas.⁹

The United Nations Children's Fund (UNICEF) and partners developed the Caregiver Journey Framework to guide countries in understanding the experiences, processes, and structures that shape how caregivers seek and receive health services for their children, including essential immunisation.¹⁰ In 2018, we operationalized the Caregiver Journey Framework through a qualitative approach in the Western Area Urban district (WAU) in Sierra Leone. Implementation experiences from operationalizing the framework in the context of urban immunization has been described elsewhere.¹¹ The framework was operationalized into several domains to understand decision-making and preparation for vaccination visits, making the journey to clinics, experiences during vaccination visits, and post-vaccination experiences. Building on these domains, we aimed to describe the real-world experiences of caregivers of vaccine-eligible children as they navigate urban immunisation services in Sierra Leone to identify vaccination enablers and barriers.

2. Methods

We aimed to understand how household dynamics, social factors, and formal healthcare delivery systems influence childhood immunisation uptake via the lens of primary caregivers of vaccine-eligible children in urban settings in Sierra Leone. Hence, we developed the Immunisation Caregiver Journey Interviews

(ICJI) approach¹¹ based on the Caregiver Journey Framework using principles of interpretative phenomenology,^{12 13} which focuses on elucidating the essence of common experiences to explain, interpret, and make sense of a phenomenon.¹⁴ We used a phenomenological approach to explore the lifeworld of caregivers in how they navigate childhood immunisation for their children repeatedly in low-resource, urban settings.¹¹ A semi-structured ICJI guide was used to explore the following domains: Decision-making and preparation, making the journey, experiences during vaccination visit, post-vaccination experiences, intentions to return, and perceptions of immunisation promotion activities in the community.

2.1. Setting

The WAU district in Sierra Leone comprises most of the capital city of Freetown with approximately 1.2 million inhabitants.¹⁵ The district was heavily affected by the Ebola epidemic, partly due to high population movements and crowded housing conditions.¹⁶ On average, there is less than one medical doctor per 10,000 population.¹⁷ The Government of Sierra Leone introduced the Free Health Care Initiative in 2010 to remove cost barriers for essential health services for pregnant and lactating mothers and under-five children.¹⁸ Childhood immunisation services are delivered through the Expanded Program on Immunisation using fixed sites that are supplemented by community outreach services to be conducted five times monthly.¹⁹ Each catchment community has 10 community health workers (CHWs) who support the promotion of health services on a voluntary basis.²⁰ A coverage survey in 2019 estimated 86% coverage for three doses of diphtheria-pertussis-tetanus (DPT) vaccine in slums and 92% coverage in non-slum urban areas in the WAU district. However, coverage of the second dose of measles-containing vaccine was very low in the district (33% in slums and 29% in non-slum urban areas).⁹

2.2. Sampling and data collection

The sample size for this qualitative assessment was guided by an approach that focuses on *qualitative information power*.²¹ The concept of information power posits that researchers should determine the sample size in a qualitative assessment based on the aim (narrow versus broad), sample specificity (targeting specific group versus multiple groups), theoretical underpinning (application of theory or no theory), quality of dialogue (weak or strong), and analysis strategy (within-case only or cross-case). Sample size burden increases when the aim is broad, multiple groups are targeted in the sample, the assessment is theory-driven, the quality of the dialogue is weak, and transcripts are analysed using cross-case analysis. In our assessment, the aim was narrow, the sample targeted a specific group, we applied theory to guide the assessment, the transcripts contained rich information, and we conducted both within-case and cross-case analyses. Against these considerations, we interviewed 16 caregivers and

1
2
3 progressively reviewed debrief notes from the interviews to assess information power. In analysing the
4 transcripts, we concluded that we reached saturation with the 16 interviews and likely could have stopped
5 interviewing after the 12th interview.
6
7

8
9 We purposively recruited the caregivers from eight communities in the WAU district, four of which were
10 slums and four were other urban areas in the district to maximize variation in the sample. Within each
11 community, two caregivers of children ages 6-36 months were selected to capture a breadth of
12 experiences of caregivers with vaccine eligible children—one whose child was fully vaccinated for age
13 and another whose child had missed at least one scheduled vaccination visit. CHWs supported data
14 collection teams in visiting households to identify and recruit eligible caregivers in the selected
15 communities. Snowball sampling was used as a secondary sampling strategy when the first identified
16 caregiver declined to interview but knew of other caregivers in the community with vaccine-eligible
17 children or when CHWs were only successful in identifying just one eligible caregiver. In this form of
18 snowball sampling, a previously visited household with an eligible child would point data collectors to
19 other households with potentially eligible children (i.e., vaccine eligible children). Data collectors visited
20 such households to screen for eligibility. This process continued until two caregivers of eligible children
21 were successfully recruited and interviewed from a particular community. Interviews were conducted on
22 the same of day of recruitment after obtaining informed consent from the caregiver.
23
24
25
26
27
28
29
30
31
32

33 We recruited data collectors (interviewers and notetakers) who were fluent in English and the
34 predominant local language in the WAU district (Krio). The data collectors had post-secondary
35 educational training in social sciences and were experienced in conducting qualitative data collection in
36 Sierra Leone. Two behavioural scientists trained the facilitators for a week on the assessment protocol.
37 One of the trainers was from Sierra Leone and had experience conducting social science research in Sierra
38 Leone. During the training, the English version of the guide was translated into Krio by locally hired staff
39 together with the facilitators and trainer. The data collectors were trained on how to probe on the spot to
40 obtain additional pertinent information from caregivers. Data collection occurred in August–September
41 2018. All interviews were audio-recorded with permission from participants; they were then transcribed
42 and translated into English by the local team. Interviews lasted about an hour on average and were
43 conducted in the vicinity of the homes of the caregivers. Data collection teams were trained on choosing
44 suitable interview locations to enable caregivers to speak freely. The facilitators conducted debriefing
45 sessions immediately after each interview to make note of key experiences and observations. The
46 debriefing notes were not part of the formal analysis. However, during the field work, the debriefing notes
47 were used to progressively assess data saturation and to identify key insights emerging from the
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 interviews. We used the insights from the debrief notes to develop a preliminary report that was mostly in
4 a descriptive, narrative form. The de-identified preliminary report was shared with the Sierra Leone
5 Ministry of Health and Sanitation. We have previously documented practical lessons learned from
6 implementing the assessment in Sierra Leone.¹¹
7
8
9

10 11 **2.3. Data analysis**

12 Two analysts (one male, one female) read all transcripts and created analytic memos and then analysed
13 the transcripts using both within-case and cross-case analysis. In the within-case analysis, we developed a
14 narrative profile for each caregiver to bring key aspects of their lived experiences to the foreground. In the
15 initial part of the cross-case analysis, each analyst coded three different transcripts (six total), using an
16 inductive approach to identify and interpret meaning units within the text. To gain alternative
17 interpretations of the coded meaning units, three of the coded transcripts were shared with a third
18 qualitative expert, who was not involved in the previous stages of the assessment, for independent ‘blind’
19 coding of the transcripts. Feedback from the third analyst was discussed by the two primary analysts and
20 incorporated into the coding scheme. The analysts used an iterative process to review their codes, discuss
21 their interpretations of the manifest content, and harmonize the initial set of codes that were used for
22 coding the remaining manuscripts. NVivo software (QSR International–2018, Version 12) was used for
23 the final organization and coding of the transcripts. Manifest categories of meaning units were grouped to
24 reflect latent content that was developed into cross-cutting themes via a consultative process. Throughout
25 the process, the analysts exercised reflexivity regarding subjective interpretations and iteratively re-
26 examined the transcripts to identify alternative interpretations until consensus was reached with additional
27 inputs from the co-authors.
28
29
30
31
32
33
34
35
36
37
38
39

40 **2.4. Ethical approval**

41 The assessment was approved by the Sierra Leone Ethics and Scientific Review Committee (SLESRC-
42 17052018), Columbia University Medical Centre Institutional Review Board (IRB-AAAR9031), and the
43 Centre for Global Health at the U.S. Centres for Disease Control and Prevention (CGH-2018-184). All
44 participants provided written informed consent.
45
46
47
48

49 **2.5. Patient and public involvement**

50 No patient involved.
51
52
53
54

55 **3. Results**

All respondents were the biological mothers of the sampled children except for one female guardian. The median age was nine months for the children included in the assessment. At the time of the interviews, half of the children had missed at least one scheduled vaccine dose. Three themes emerged from the interviews around vaccination enablers (Table 1), vaccination barriers (Table 2) and direct recommendation to improve vaccination uptake (Table 3). There were no notable differences in themes between slums and other urban communities.

3.1. Enablers of childhood vaccination (Table 1)

Our analysis identified emotional and practical enablers related to childhood vaccination. Emotional enablers were evident in how caregivers portrayed their parental obligation to their children, wanting to do the right thing for their children's health, and anticipating reciprocal benefits in children's ability to take care of the parents later in life. Practical enablers were the diversity of immunization reminders, information access, information trust, getting fathers more involved, positive experiences with health workers, and post-vaccination information sharing at the community level.

3.1.1. Parental responsibility

A sense of parental responsibility was a major motivating factor for caregivers to seek vaccination services for their children. Caregivers viewed vaccination as 'doing the right thing' for their children. Even caregivers who had missed scheduled vaccination visits felt responsible for getting their children caught up with their scheduled vaccine doses, which often required deprioritizing other income-generating activities.

"It is my duty to take my baby to the hospital for immunisation. It is my responsibility as [a] mother to ensure that my baby completes the rounds of immunisation without defaulting." –

Caregiver whose child had missed a scheduled vaccination

3.1.2. Wanting a strong and healthy baby

In addition to the affective responses regarding a sense of duty to the child, appreciating the overall health benefits of vaccination to their children was another major driving force in motivating caregivers to seek vaccination services. Caregivers consistently expressed that immunisation has essential health benefits to the child and that missing scheduled vaccination would 'risk the baby's life.' Moreover, they valued having a 'strong and healthy baby' and felt that completing the vaccination schedule would positively impact the baby's health.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

“I think [the] vaccine is good for our children. It is important and it helps to build their immune system to keep them strong and healthy; it fights against many things in the body...” – Caregiver whose child was fully vaccinated

3.1.3. Parental anticipation of reciprocal benefit

Perceptions of vaccination benefits went beyond the direct health benefits to children and extended into domains of benefit to the parent. The notion that vaccinated children will be healthier and in turn live longer and be able to take care of their parents later in life emerged as a dimension of vaccination benefit to the parent. This duality of vaccination benefit was grounded in the cultural context of parents expecting reciprocal care from their children when the parents can no longer care for themselves.

“It is very difficult for me to be absent for immunisation or not to take my baby to the hospital when sick. Sometimes people think I’m mad but I’m not. I’m trying to bring up my children in a way that they will benefit me when I’m old.” – Caregiver whose child had missed a scheduled vaccination

3.1.4. Diversity of immunisation reminders

Caregivers were exposed to numerous reminders and cues to vaccinate their children, including information from the immunisation cards, health workers, community campaigns, and family members. However, the child’s immunisation card stood out as the most important reminder that caregivers and families relied on to remember the dates of the scheduled vaccination visits.

“I do remind myself because they [health workers] will write at the back of the immunisation card the date for the next visit. The immunisation card will tell you the date for the next visit.” – Caregiver whose child had missed a scheduled vaccination

In addition, community outreach campaigns and announcements in the community were also viewed as helpful reminders to vaccinate.

“The health workers in this community and the CHWs are doing well as they do go round telling people not to forget to take their babies to the hospital for immunisation.” – Caregiver whose child had missed a scheduled vaccination

3.1.5. Information access and information trust

Immunisation information sources varied, but caregivers consistently cited health workers as trusted sources of information. Nurses were more trusted than lay CHWs because caregivers viewed nurses as

1
2
3 more knowledgeable. While waiting in line before immunisation services, most caregivers appreciated the
4 ‘health talk’ they received from nurses who advised on health and immunisation.
5

6 *“I trust them [nurses] because they are a team of qualified nurses ... Before they come here, they*
7 *know everything about the vaccines and any implications of the vaccines. They are able to*
8 *explain more than the CHWs and other community workers in the area. The nurses will tell you*
9 *more. There are things that the community workers do not know, and they refer you to the*
10 *nurses.”* – Caregiver whose child was fully vaccinated
11
12
13
14
15

16 **3.1.6. Getting fathers more involved**

17 Mothers desired greater involvement by the child’s fathers in supporting their children’s vaccination. In
18 most instances, fathers were rarely involved in taking their children to the vaccination clinic. We
19 uncovered only two instances when fathers actively supported their children’s vaccination visits. In those
20 instances, the mothers felt supported, and their children were fully vaccinated. In one situation, a father
21 that routinely accompanied the child to the vaccination clinic was celebrated by health workers and given
22 the ‘best father’ award.
23
24
25

26 *“There was [a] time when he [my husband] was given the best father award [at the clinic]*
27 *because he is always with me at the hospital. That is the name I have also given to my husband.*
28 *Even when the baby is crying, I will say ‘best father’ take your baby.”* – Caregiver whose child
29 was fully vaccinated
30
31
32
33
34

35 **3.1.7. Positive experiences with health workers**

36 During vaccination visits, positive experiences with health workers encouraged caregivers to vaccinate
37 their children. Positive experiences included having a cordial relationship with the health workers. In
38 particular, caregivers expressed that the nurses took good care of their children, made them feel
39 comfortable, and tried to build a good rapport. Some caregivers said they would voluntarily give small
40 amounts of money to health workers as a token of their appreciation after vaccinating the child.
41
42
43

44 *“...the nurses do encourage you and will make jokes so that you will laugh at the end of the day.*
45 *There is a lot of fun [interactions], which made some of us forget about our stresses.”* –
46 Caregiver whose child was fully vaccinated
47
48
49
50

51 **3.1.8. Post-vaccination information sharing**

52 Information exchange at the community level with trusted community members strengthened caregiver
53 confidence in childhood vaccination. Upon returning home from the vaccination visit, caregivers often
54 discussed the clinic experience with their spouses, families, and other caregivers in the community. In
55
56
57

1
2
3 addition to information obtained from health workers at the clinic, caregivers also sought advice from
4 other ‘more experienced’ caregivers in the community.
5

6 *“Most times after immunisation, my baby will run a temperature, but the health workers always*
7 *provide drugs to counter the fever. We have caregivers in this community with vast knowledge*
8 *and since this is my first baby, I love to talk to them so that we can share our experiences which*
9 *will be of advantage to me as I’m very new in the field [of parenting].” –Caregiver whose child*
10 *was fully vaccinated*
11
12
13

14 15 16 **3.2. Barriers related to childhood vaccination (Table 2)**

17 Practical constraints, negative experiences with health workers, and safety concerns were the underlying
18 barriers to childhood vaccination. Practical constraints included challenges faced when preparing for and
19 getting to the vaccination clinic and inconveniences encountered at the clinic, such as overcrowding and
20 long waiting times. Negative experiences among caregivers included feeling disrespected by health
21 workers while simultaneously expecting to give money to health workers for services that are supposed to
22 be free of charge. Finally, vaccine side effects led to concerns and fears about vaccine safety.
23
24
25
26
27

28 29 **3.2.1. Preparing for the journey and getting to the clinic**

30 Caregivers commonly cited the need to juggle ‘household duties’ and other income-generating activities
31 when planning the visit as a barrier, especially in the absence of fathers’ involvement in taking the child
32 to the clinic. In addition, mothers frequently depended on their children’s fathers for financial support to
33 cover the expenses related to the vaccination clinic visit. Some caregivers recounted needing to travel
34 long distances up to an hour by foot to get to the vaccination clinic, especially when they could not afford
35 public transportation.
36
37
38
39
40

41 42 **3.2.2. Inconveniences at the clinic**

43 Caregivers anticipated various inconveniences at the vaccination visit. The prolonged time spent waiting
44 for the child to be vaccinated emerged as a substantial inconvenience that was more pronounced when
45 seeking immunisation services, especially in larger facilities. Anticipating the long wait, caregivers
46 usually tried to arrive early at the vaccination site to get seen first. The range of activities involved with
47 the child health visit prolonged the visit, including weighing the baby as part of growth monitoring and
48 other health checks.
49
50
51

52 *“It is painful if you waste much time at the health facility because you have other issues to attend*
53 *to. To avoid that, that is why I always come early to the health facility.” – Caregiver whose child*
54 *was fully vaccinated*
55
56
57

3.2.3. Feeling disrespected by health workers

Caregivers often felt disrespected by health workers during the vaccination visit. A key complaint was that health workers shouted at caregivers and sometimes used vulgar language toward caregivers. In other instances, they complained that some health workers habitually arrived late to the vaccination session, which further prolonged the time caregivers spent waiting.

3.2.4. Monetary expectations

Systemically hidden costs generated substantial dissatisfaction among caregivers. Caregivers needed to “shake hands” with health workers at different times of the visit (e.g., first time registering the child to get a card, before entering the facility, and before weighing the baby). Shaking hands implied giving some money during the handshake. Caregivers used the money to ‘fast-track’ their children’s vaccination. The expectations around monetary exchange discouraged caregivers who could not afford to shake hands with health workers.

“Sometimes if I don’t want to spend much time at the hospital, I will shake the hand of the nurse so that they can fast track the immunisation of my baby. I will give them something like two thousand Leones or whatever I have with me at that moment... Health is wealth and they [health workers] don’t need us but we do [need them]. The money we give is nothing compared to the health of our children... At the end of the day, we will grumble on our way home as the services are supposed to be free for our children, yet we are paying for it. The health workers are really trying, but the idea for them to take money from us is bad. And if you don’t give them money, they will talk to you carelessly.” – Caregiver whose child was fully vaccinated

In a separate domain of monetary exchange, health workers demanded money as a form of ‘punishment’ to caregivers who missed their children’s scheduled vaccination appointments.

“If you failed to take your baby to the hospital on [the] stipulated date, you will definitely have to pay some amount at the end of the day in the form of punishment. You must pay five thousand or more.” – Caregiver whose child was fully vaccinated

3.2.5. Vaccine side effects

Caregivers cited numerous instances when their children experienced vaccine side effects such as ‘fever,’ ‘swelling at the injection site,’ and the ‘baby becoming lethargic.’ Fever was the most common side effect, and the caregivers knew to administer fever-reducing medication as instructed by health workers.

1
2
3 When there was swelling at the injection site, a common practice among caregivers was to massage the
4 swollen area, sometimes with an onion or a bar of soap to try to reduce the swelling.
5

6 *“Sometimes my baby’s leg becomes swollen... because some nurses are heavy-handed, and I meet*
7 *several nurses when I visit the hospital. Sometimes the leg gets swollen, and they treat him. I have*
8 *to rub the leg to avoid swelling... I use soap to rub off the swelling and I give Panadol to stop the*
9 *fever... some people say you should not allow every nurse to administer [an]injection to the child.*
10 *I should have a permanent nurse that gives injection to my child without swelling.” – Caregiver*
11 *whose child missed a scheduled vaccination*
12
13
14
15

16
17 In one rare situation, a caregiver had a prior child who experienced fever, convulsed, and died a few days
18 after getting vaccinated. Therefore, the caregiver decided to not vaccinate subsequent children.
19

20 *“After the immunisation, my baby started running temperature, I administered paracetamol as I*
21 *was told by the health workers. The baby convulsed and that was the end of that baby. I don’t*
22 *want a repeat of that in my life. I have therefore decided not to take my babies for immunisation*
23 *anymore” – Caregiver whose child missed a scheduled vaccination*
24
25
26
27

28 Besides passive acceptance of the BCG vaccine at the birth of the youngest child, this same caregiver
29 actively refused all other vaccines despite encouragement by a family member to vaccinate the child.
30

31 *“I’m not outrightly saying it was as a result of the immunisation [that my child died]; as every*
32 *death is the work of God. But from what I have gathered so far, I have personally decided not to*
33 *take my baby to the health facility to be immunized. It is not that I’m tired of going to the health*
34 *facility or because of the distance or money. I do get a lot of pressure from my aunt to take my*
35 *baby to the health facility for immunisation, but the thing is that I just don’t trust the system and*
36 *what immunisation does. – Caregiver whose child missed a scheduled vaccination*
37
38
39
40
41

42 From the perspective of this same caregiver, the vaccinated child died but the unvaccinated children
43 survived and thrived, which was cited as a reason for refusing vaccination.
44

45 *“I believe in exclusive breastfeeding, sometimes for two years and a half and sometimes [only]*
46 *two years. My baby is now two years seven months old and doing well like any of those children*
47 *that are on immunisation or have completed...Just as I was saying, sometimes my heart will tell*
48 *me to take the baby for immunisation but after thinking of the past experience, I would decide not*
49 *to. I’m now used to that... The simple fact here is that, since the other children are doing well*
50 *without immunisation, I will not take [the baby] to the health facility for immunisation and that is*
51 *all.” – Caregiver whose child missed a scheduled vaccination*
52
53
54
55
56
57

3.3. Recommendations to improve childhood vaccination (Table 3)

The direct recommendations provided by caregivers were categorized into (1) improving vaccination process and systems and (2) engaging communities to boost vaccine confidence. Implementing community outreach campaigns for immunisation at regular intervals with a focus on defaulters was recommended by participants to improve vaccination outcomes. In addition, caregivers wanted health workers and community leaders to be involved in immunisation promotion along with the CHWs. They wanted the vaccination clinic experience to improve and become more conducive to caregivers, including shorter wait time at the clinic and more positive interactions with health workers. Lastly, caregivers wanted health workers to stop demanding money from them, though they may not mind giving money, out of free-will, as a token of appreciation when they could afford it.

“In addition, you should engage the Chiefs, because in each area we have Chiefs to spread out this message. You could educate them so that they in turn can educate those in the community. Let us have Town Criers go around disseminating the messages. It would be nice for them to allocate people in the health centre who move from house to house to educate the breastfeeding mothers because some of us are stubborn to come onboard.” – Caregiver whose child was fully vaccinated

4. Discussion

Our qualitative analysis highlighted several important themes. In the backdrop of anticipated benefits to both the child and parent, vaccination intention was motivated by a feeling of parental responsibility to ‘do the right thing.’ Timely and trusted exchange of information together with social support and positive experiences at the vaccination clinic were important facilitators of vaccination. In contrast, vaccination was discouraged by negative interactions with health workers at the clinic, the occurrence and fear of vaccine side effects, multitude of ‘hidden’ costs, juggling vaccination with other responsibilities, and inconveniences, such as long traveling time to the clinic and long delays at the clinic. Nevertheless, caregivers were resilient in devising ways to try to get their children vaccinated such as walking on foot up to an hour to get to the clinic when they could not afford public transportation. Lastly, caregivers wanted the vaccination experience to improve, and they desired stronger community engagement to help optimize vaccination outcomes. However, systemic issues, such as informal payments, overcrowding in health facilities, and the reported overburdened health workers may require interventions at the health systems level. These themes from Sierra Leone provide in-depth insights regarding the motivations, facilitators, and barriers of vaccination in an urban LMIC setting.

1
2
3
4
5 Moral values may shape vaccination attitudes.²² Philosophical arguments regarding the morality of
6 vaccination have been heavily debated.²³⁻²⁶ Caregivers in our sample largely viewed vaccination via a
7 moral lens encompassing parental duty to do the right thing for the child. In one situation, however, we
8 found that the desire to ‘do the right thing’ may also translate into vaccination refusal in the backdrop of
9 other past refusals, observing ‘healthy unvaccinated’ children, and having distrust of the health system.
10 Quantitative research from high-income countries has shown that parents with unvaccinated children were
11 more likely to perceive their children to be at low risk of vaccine-preventable diseases and were more
12 likely to perceive low vaccine effectiveness and safety compared to parents with vaccinated children.²⁷
13 Our findings suggest that childhood immunisation communication efforts may resonate more strongly
14 with caregivers when vaccination is framed around parental responsibilities to do the right thing for the
15 child and the anticipated future benefits to parents. However, additional research is necessary to generate
16 a better understanding of the morality of childhood vaccination in the Sierra Leonean context.
17
18
19
20
21
22
23
24

25 Across the interviews, there was an apparent tension in the relationship between caregivers and health
26 workers. Caregivers often expressed their appreciation of health workers and empathized with the
27 challenging context in which they do their work. Health workers were strongly viewed as authoritative
28 sources of trusted information regarding immunisation and the child’s health, which is consistent with
29 findings from high-income countries²⁸⁻³⁰ and LMICs.^{31 32} Our findings on the role of monetary exchange
30 in vaccination exemplify the complex relationship between caregivers and health workers in low-resource
31 urban communities in Sierra Leone. Some caregivers voluntarily gave money to health workers as a
32 ‘token of appreciation’ while others begrudgingly gave money because they viewed it as a condition for
33 receiving good quality service from health workers. Interventions at the health systems level are
34 necessary to help discourage informal payments to health workers—a practice that may perpetuate
35 vaccination inequities among poor caregivers who are unable to meet monetary expectations.
36
37
38
39
40
41
42
43

44 Our findings also illuminate the need for interventions at the household and family level. Fathers were
45 rarely involved in taking their children to the vaccination clinic but were often engaged in the decision-
46 making processes. In the few instances when fathers were involved in taking their children to the clinic,
47 the mothers felt supported, and their children were fully vaccinated. A study in Nigeria found that
48 paternal involvement in immunisation was greater in rural settings compared to urban settings.³³ In urban
49 areas, the same study found that paternal involvement was greater among educated fathers compared to
50 uneducated fathers. In a separate study in Ghana, the involvement of educated fathers in the vaccination
51 decision was associated with timelier vaccination uptake compared to the involvement of uneducated
52
53
54
55
56
57
58
59
60

1
2
3 fathers in the decision.³⁴ More broadly, shifting from a mother-child dyad to a family triad in the care of
4 children has proven to have positive effects on paediatric health outcomes across diverse contexts.³⁵
5 Additional assessments and interventions are needed to explore and evaluate culturally appropriate ways
6 to enhance the involvement of fathers in childhood immunisation in Sierra Leone and other similar LMIC
7 settings.
8
9
10

11
12 Existing evidence suggests that vaccine safety concerns, often linked to adverse events following
13 immunisation (AEFI), contribute to vaccine hesitancy.^{36 37} Serious AEFIs may be ‘triggering events’ for
14 derailing vaccine confidence and prompting active refusal among certain caregivers and their
15 communities—especially when the serious AEFI is perceived to be linked to the vaccine or the
16 vaccination process.³⁸ Together with prior evidence, these findings emphasize the need for robust AEFI
17 surveillance and investigations³⁹ to identify, counsel, and follow-up with caregivers whose children
18 experience AEFI, and therefore, are potentially at risk of missing subsequent vaccination. Health workers
19 and CHWs may benefit from periodic in-service training on how to effectively communicate vaccine
20 safety and address concerns about AEFIs.⁴⁰
21
22
23
24
25
26
27

28 **4.1. Limitations**

29
30 There were several limitations to our assessment. First, it is possible that some nuanced meaning may
31 have been lost when translating the audio recordings from Krio to English—especially since the
32 transcripts were not back-translated from English to Krio due to resource constraints. Second, we only
33 identified one caregiver who routinely and actively refused all vaccines for her child after experiencing a
34 serious AEFI, which may reflect the overall rarity of zero-dose unvaccinated children in Sierra Leone
35 (approximately 3%).⁴¹ This was the only caregiver with a child that experienced a serious AEFI in our
36 sample, which limits our ability to have a rich understanding of such experience among caregivers more
37 broadly and the potential linkages to vaccination refusal. Although such experiences of serious AEFI are
38 rare, they may have the tendency to get publicized in the community, which may negatively influence
39 vaccination decisions among other caregivers. Taken together, our results call for additional qualitative
40 assessments to get a deeper understanding of vaccination refusal within the Sierra Leonean context and
41 other low-resource LMIC settings. Sampling strategies may therefore need to be adapted accordingly to
42 focus on caregivers who actively refuse vaccination for their children. Lastly, our findings reflected
43 gender dimensions that may be based on sociocultural norms in Sierra Leonean society but may also be
44 due to sampling bias because the caregivers we conveniently recruited were mostly stay-at-home mothers.
45
46
47
48
49
50
51
52
53
54

55 **5. Conclusions**

1
2
3 As the COVID-19 pandemic disrupts childhood immunisation globally,⁴² especially in LMICs, our
4 assessment provides a foundational understanding of the challenges that caregivers encounter in urban
5 settings in Sierra Leone. It also sheds light on opportunities to improve vaccination outcomes in urban
6 poor settings, which is a global immunisation priority. The findings show that health system
7 interventions, community engagement, and vaccination outreach may need to be tailored to for urban
8 settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is
9 framed both around parental responsibilities to do the right thing for the child and the future benefits to
10 the parent.

17 **Competing interests**

18 None declared.

22 **Data availability**

23 All data relevant to the study are included in the article or uploaded as supplementary information.

27 **Contributors**

28 MFJ, RS, MT, TS, and ML designed the assessment. MFJ, PP, SK, and KW analysed the data. MJF, PP,
29 RS, SK, MT, KW, TS, and ML contributed to the interpretation of the results. MFJ, PP and SK drafted
30 the manuscript. RS, MT, KW, TS, and ML contributed to revising the manuscript. All authors reviewed
31 and approved the final manuscript.

37 **Collaborators**

38 Adewale Akinjeji, Laura Conklin, Oliver Eleeza, Lauren E Parmley, Anthony Mansaray, Dimitri
39 Prybylski, Roberta Sutton, Mame Toure, Aaron Wallace, Brent Wolff

43 **Acknowledgements**

44 We thank the 16 caregivers who participated in the in-depth interviews.

48 **Funding**

49 U.S. Centers for Disease Control and Prevention (U19GGH001581).

53 **Disclaimer**

The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the US Centres for Disease Control and Prevention.

6. References

1. Nelson KN, Wallace AS, Sodha SV, et al. Assessing strategies for increasing urban routine immunization coverage of childhood vaccines in low and middle-income countries: A systematic review of peer-reviewed literature. *Vaccine* 2016;34(46):5495-503. doi: 10.1016/j.vaccine.2016.09.038 [published Online First: 2016/10/25]
2. John Snow International. Strengthening Immunization for the Urban Poor: Experiences and Lessons Learned from Five Country Studies, 2018. Available: <https://www.jsi.com/resource/strengthening-immunization-for-the-urban-poor-experiences-and-lessons-learned-from-five-country-studies/> [accessed November 12 2020].
3. Brolin Ribacke KJ, Saulnier DD, Eriksson A, et al. Effects of the West Africa Ebola Virus Disease on Health-Care Utilization - A Systematic Review. *Frontiers in public health* 2016;4:222. doi: 10.3389/fpubh.2016.00222 [published Online First: 2016/10/26]
4. Elston JW, Cartwright C, Ndumbi P, et al. The health impact of the 2014-15 Ebola outbreak. *Public health* 2017;143:60-70. doi: 10.1016/j.puhe.2016.10.020 [published Online First: 2017/02/06]
5. Yerger P, Jalloh M, Coltart CEM, et al. Barriers to maternal health services during the Ebola outbreak in three West African countries: a literature review. *BMJ Glob Health* 2020;5(9) doi: 10.1136/bmjgh-2020-002974 [published Online First: 2020/09/09]
6. Masresha BG, Luce R, Jr., Weldegebriel G, et al. The impact of a prolonged ebola outbreak on measles elimination activities in Guinea, Liberia and Sierra Leone, 2014-2015. *Pan Afr Med J* 2020;35(Suppl 1):8. doi: 10.11604/pamj.supp.2020.35.1.19059 [published Online First: 2020/05/07]
7. Government of Sierra Leone. National Ebola Recovery Strategy for Sierra Leone, 2015. Available: https://ebolaresponse.un.org/sites/default/files/sierra_leone_-_national_recovery_strategy_2015-2017.pdf [accessed December 5 2020].
8. U.S. Centers for Disease Control and Prevnetion. Global Health – Sierra Leone, 2019. Available: <https://www.cdc.gov/globalhealth/countries/sierra-leone/default.htm> [accessed December 5 2020].
9. Feldstein LR, Sutton R, Jalloh MF, et al. Access, demand, and utilization of childhood immunization services: A cross-sectional household survey in Western Area Urban district, Sierra Leone, 2019. *Journal of global health* 2020;10(1):010420. doi: 10.7189/jogh.10.010420 [published Online First: 2020/06/09]
10. UNICEF. Human Centered Design 4 Health, 2018. Available: <https://www.hcd4health.org/> [accessed November 18 2020].
11. Jalloh MF, Kinsman J, Conteh J, et al. Barriers and facilitators to reporting deaths following Ebola surveillance in Sierra Leone: implications for sustainable mortality surveillance based on an exploratory qualitative assessment. *BMJ Open* 2021;11(5):e042976. doi: 10.1136/bmjopen-2020-042976 [published Online First: 2021/05/15]
12. Diekelmann NL. Learning-as-testing: a Heideggerian hermeneutical analysis of the lived experiences of students and teachers in nursing. *ANS Advances in nursing science* 1992;14(3):72-83. doi: 10.1097/00012272-199203000-00010 [published Online First: 1992/03/01]
13. Dreyfus HL. Being-in-the-world: A commentary on Heidegger's Being and Time. Cambridge: The MIT Press 1991.
14. Dahlberg K. The essence of essences – the search for meaning structures in phenomenological analysis of lifeworld phenomena. *International journal of qualitative studies on health and well-being* 2006;1(1):11-19. doi: 10.1080/17482620500478405

15. Statistics Sierra Leone. 2015 Sierra Leone Population and Housing Census, 2016. Available: <https://www.statistics.sl/index.php/census/census-2015.html> [accessed November 12 2020].
16. Lamunu M, Olu OO, Bangura J, et al. Epidemiology of Ebola Virus Disease in the Western Area Region of Sierra Leone, 2014-2015. *Frontiers in public health* 2017;5:33. doi: 10.3389/fpubh.2017.00033
17. World Health Organization. Medical doctors per 10,000 population, 2021. Available: <https://www.who.int/data/gho/data/indicators/indicator-details/GHO> [accessed December 5 2021].
18. Witter S, Brikci N, Harris T, et al. The free healthcare initiative in Sierra Leone: Evaluating a health system reform, 2010-2015. *The International journal of health planning and management* 2018;33(2):434-48. doi: 10.1002/hpm.2484
19. Government of Sierra Leone. Comprehensive EPI Multi-Year Plan 2017-2021, 2016. Available: <https://www.gavi.org/sites/default/files/document/2021/cMYP%20Sierra%20Leone%202017-2021.pdf> [accessed December 5 2020].
20. Government of Sierra Leone. Policy for Community Health Workers in Sierra Leone, 2012. Available: <https://chwcentral.org/wp-content/uploads/2015/01/CHW-Policy-Sierra-Leone.pdf> [accessed December 5 2020].
21. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview Studies: Guided by Information Power. *Qualitative health research* 2016;26(13):1753-60. doi: 10.1177/1049732315617444
22. Amin AB, Bednarczyk RA, Ray CE, et al. Association of moral values with vaccine hesitancy. *Nature human behaviour* 2017;1(12):873-80. doi: 10.1038/s41562-017-0256-5
23. Di Pietro ML, Poscia A, Telesman AA, et al. Vaccine hesitancy: parental, professional and public responsibility. *Annali dell'Istituto superiore di sanita* 2017;53(2):157-62. doi: 10.4415/ann_17_02_13
24. Giubilini A, Douglas T, Savulescu J. The moral obligation to be vaccinated: utilitarianism, contractualism, and collective easy rescue. *Medicine, health care, and philosophy* 2018;21(4):547-60. doi: 10.1007/s11019-018-9829-y [published Online First: 2018/02/13]
25. Bester JC. Measles Vaccination is Best for Children: The Argument for Relying on Herd Immunity Fails. *Journal of bioethical inquiry* 2017;14(3):375-84. doi: 10.1007/s11673-017-9799-4
26. Dawson A. What are the moral obligations of the traveller in relation to vaccination? *Travel medicine and infectious disease* 2007;5(2):90-6. doi: 10.1016/j.tmaid.2006.01.005
27. Omer SB, Salmon DA, Orenstein WA, et al. Vaccine refusal, mandatory immunization, and the risks of vaccine-preventable diseases. *N Engl J Med* 2009;360(19):1981-8. doi: 10.1056/NEJMsa0806477
28. Frew PM, Chung Y, Fisher AK, et al. Parental experiences with vaccine information statements: Implications for timing, delivery, and parent-provider immunization communication. *Vaccine* 2016;34(48):5840-44. doi: 10.1016/j.vaccine.2016.10.026
29. Drezner D, Youngster M, Klainer H, et al. Maternal vaccinations coverage and reasons for non-compliance - a cross-sectional observational study. *BMC pregnancy and childbirth* 2020;20(1):541. doi: 10.1186/s12884-020-03243-w
30. Wiley KE, Massey PD, Cooper SC, et al. Pregnant women's intention to take up a post-partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: a cross sectional survey. *Vaccine* 2013;31(37):3972-8. doi: 10.1016/j.vaccine.2013.06.015
31. Makwe CC, Anorlu RI. Knowledge of and attitude toward human papillomavirus infection and vaccines among female nurses at a tertiary hospital in Nigeria. *International journal of women's health* 2011;3:313-7. doi: 10.2147/ijwh.S22792
32. Foss HS, Oldervoll A, Fretheim A, et al. Communication around HPV vaccination for adolescents in low- and middle-income countries: a systematic scoping overview of systematic reviews. *Systematic reviews* 2019;8(1):190. doi: 10.1186/s13643-019-1100-y

- 1
2
3 33. Sodeinde K, Amoran O, Abiodun O, et al. A rural/urban comparison of paternal involvement in
4 childhood immunisation in Ogun Central Senatorial District, Nigeria. *The Nigerian postgraduate*
5 *medical journal* 2020;27(4):336-42. doi: 10.4103/npmj.npmj_101_20
6
7 34. Brugha RF, Kevany JP, Swan AV. An investigation of the role of fathers in immunization uptake. *Int*
8 *J Epidemiol* 1996;25(4):840-5. doi: 10.1093/ije/25.4.840
9
10 35. Allport BS, Johnson S, Aqil A, et al. Promoting Father Involvement for Child and Family Health.
11 *Acad Pediatr* 2018;18(7):746-53. doi: 10.1016/j.acap.2018.03.011
12
13 36. Dube E, Laberge C, Guay M, et al. Vaccine hesitancy: an overview. *Human vaccines &*
14 *immunotherapeutics* 2013;9(8):1763-73. doi: 10.4161/hv.24657
15
16 37. Larson HJ, Jarrett C, Eckersberger E, et al. Understanding vaccine hesitancy around vaccines and
17 vaccination from a global perspective: a systematic review of published literature, 2007-2012.
18 *Vaccine* 2014;32(19):2150-9. doi: 10.1016/j.vaccine.2014.01.081
19
20 38. Du F, Chantler T, Francis MR, et al. The determinants of vaccine hesitancy in China: A cross-
21 sectional study following the Changchun Changsheng vaccine incident. *Vaccine*
22 2020;38(47):7464-71. doi: 10.1016/j.vaccine.2020.09.075
23
24 39. World Health Organization. Adverse events following immunization (AEFI) 2021. Available:
25 https://www.who.int/immunization/monitoring_surveillance/who-immuniz.pdf [accessed
26 February 18 2021].
27
28 40. World Health Organization. Vaccine safety basics e-learning course, 2021. Available:
29 <https://vaccine-safety-training.org/simplified-and-key-messages.html> [accessed February 19]
30 2021.
31
32 41. UNICEF. Sierra Leone Multiple Indicators Cluster Survey 2017. Available:
33 https://mics.unicef.org/news_entries/106/SIERRA-LEONE-2017-MICS-RELEASED [accessed
34 January 21 2021].
35
36 42. Abbas K, Procter SR, van Zandvoort K, et al. Routine childhood immunisation during the COVID-19
37 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2
38 infection. *Lancet Glob Health* 2020;8(10):e1264-e72. doi: 10.1016/s2214-109x(20)30308-9
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Enablers of childhood vaccination—Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-themes	Theme
Responsibility to ensure full vaccination	Parental responsibility	Emotional enablers of childhood vaccination	Enablers of childhood vaccination
Obligation to do the right thing			
Immunisation as a requirement	Wanting a ‘strong and healthy baby’		
Immunisation is important for baby’s health			
Wanting ‘strong baby’			
‘Health is wealth’	Parental anticipation of reciprocal benefit		
Defaulting on vaccination risks baby’s life			
Vaccination benefits the parent later in life	Diversity of immunisation reminders	Practical enablers of childhood vaccination	
Taking care of parents when old			
Immunisation card is important			
Campaign as reminder			
Reminders by health workers at vaccination visit			
Husband as reminder			
Immunisation card as reminder			
Other family member as reminder			
Nurses should lead			Information access and information trust
Nurses more trusted than CHWs			
Same information from different sources			
Immunisation promotion through radio/tv			
Immunisation promotion by health workers			
Immunisation promotion through leaders	Getting fathers more involved		
Immunisation promotion by NGOs			
Mothers take the child to the clinic			
Fathers rarely involved	Positive experiences with health worker		
Father received an award for involvement			
Cordial relationship with nurses			
Good care by nurses			
HWs encourage seeking care at HF	Post-vaccination information sharing		
Giving money to health worker as token of appreciation			
Husband asking about the visit			
Informing husband of next visit			
Telling husband about visit expenses			
Immunisation is a “learning process”			
Sharing experiences with neighbours or friends			
Other family members asking about the visit			

Table 2. Barriers related to childhood vaccination–Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-themes	Themes
Asking husband for money	Preparing for the journey and getting to the clinic	Practical constraints	Barriers related to childhood vaccination
Juggling different household duties			
Competing priorities			
Time taken to get to the clinic			
Wasting time at the clinic			
Rush to arrive first and seen first			
Crowding at the clinic			
Very long wait at larger health facilities			
Spent a day and nothing happened			
Wasting caregiver's time	Inconveniences at the clinic	Negative experiences with health workers	
Not respecting caregivers			
Shouting at caregivers			
Paying for the card			
Paying for free drugs			
Paying for weighing			
Payment as punishment			
Bad care without payment			
Health workers should stop demanding money			
Health worker don't appreciate less than Le 2000	Vaccine side effects	Safety concerns	
Health workers withholding free drugs			
Baby crying throughout the night			
Baby gets 'lazy' for few hours			
Fever in baby			
Swelling at injection site			
Side effects only for some vaccines			
Rub onion on swollen injection site			
Avoiding abnormalities from vaccine			
Afraid of vaccine side effects			

Table 3. Recommendations to improve childhood vaccination– Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-theme	Theme
Continue campaigns at repeated intervals	Improving vaccination processes and systems	Caregivers want improved vaccination processes, systems, and engagement	Recommendations to improve childhood vaccination
Do not rely on campaigns alone			
Provision of incentives for caregivers			
Concentrate on defaulters			
Compensation/incentives for nurses			
Being considerate towards health workers			
Employ more staff			
Stop demanding money	Engaging communities to		
Promote consequences of not vaccinating			
Peer-to-peer promotion of immunisation			

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Inform about importance of immunisation	boost vaccine confidence		
Address ‘stubborn’ caregivers			
Engage caregivers who do not want injections			

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2021-058203 on 9 May 2022. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

COREQ (CONsolidated criteria for REporting Qualitative research) Checklist

A checklist of items that should be included in reports of qualitative research. You must report the page number in your manuscript where you consider each of the items listed in this checklist. If you have not included this information, either revise your manuscript accordingly before submitting or note N/A.

Topic	Item No.	Guide Questions/Description	Reported on Page No.
Domain 1: Research team and reflexivity			
<i>Personal characteristics</i>			
Interviewer/facilitator	1	Which author/s conducted the interview or focus group?	
Credentials	2	What were the researcher's credentials? E.g. PhD, MD	
Occupation	3	What was their occupation at the time of the study?	
Gender	4	Was the researcher male or female?	
Experience and training	5	What experience or training did the researcher have?	
<i>Relationship with participants</i>			
Relationship established	6	Was a relationship established prior to study commencement?	
Participant knowledge of the interviewer	7	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	
Interviewer characteristics	8	What characteristics were reported about the interviewer/facilitator? e.g. Bias, assumptions, reasons and interests in the research topic	
Domain 2: Study design			
<i>Theoretical framework</i>			
Methodological orientation and Theory	9	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	
<i>Participant selection</i>			
Sampling	10	How were participants selected? e.g. purposive, convenience, consecutive, snowball	
Method of approach	11	How were participants approached? e.g. face-to-face, telephone, mail, email	
Sample size	12	How many participants were in the study?	
Non-participation	13	How many people refused to participate or dropped out? Reasons?	
<i>Setting</i>			
Setting of data collection	14	Where was the data collected? e.g. home, clinic, workplace	
Presence of non-participants	15	Was anyone else present besides the participants and researchers?	
Description of sample	16	What are the important characteristics of the sample? e.g. demographic data, date	
<i>Data collection</i>			
Interview guide	17	Were questions, prompts, guides provided by the authors? Was it pilot tested?	
Repeat interviews	18	Were repeat interviews carried out? If yes, how many?	
Audio/visual recording	19	Did the research use audio or visual recording to collect the data?	
Field notes	20	Were field notes made during and/or after the interview or focus group?	
Duration	21	What was the duration of the interviews or focus group?	
Data saturation	22	Was data saturation discussed?	
Transcripts returned	23	Were transcripts returned to participants for comment and/or	

Topic	Item No.	Guide Questions/Description	Reported on Page No.
		correction?	
Domain 3: analysis and findings			
<i>Data analysis</i>			
Number of data coders	24	How many data coders coded the data?	
Description of the coding tree	25	Did authors provide a description of the coding tree?	
Derivation of themes	26	Were themes identified in advance or derived from the data?	
Software	27	What software, if applicable, was used to manage the data?	
Participant checking	28	Did participants provide feedback on the findings?	
<i>Reporting</i>			
Quotations presented	29	Were participant quotations presented to illustrate the themes/findings? Was each quotation identified? e.g. participant number	
Data and findings consistent	30	Was there consistency between the data presented and the findings?	
Clarity of major themes	31	Were major themes clearly presented in the findings?	
Clarity of minor themes	32	Is there a description of diverse cases or discussion of minor themes?	

Developed from: Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International Journal for Quality in Health Care*. 2007. Volume 19, Number 6: pp. 349 – 357

Once you have completed this checklist, please save a copy and upload it as part of your submission. DO NOT include this checklist as part of the main manuscript document. It must be uploaded as a separate file.