



BMJ Open Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone

Mohamed F Jalloh ¹, Palak Patel ^{1,2}, Roberta Sutton,³ Shibani Kulkarni ^{1,2}, Mame Toure,⁴ Kerrie Wiley,⁵ Tom Sessay,⁶ Maria Lahuerta,^{3,7} Sierra Leone Urban Immunisation Needs Assessment Group

To cite: Jalloh MF, Patel P, Sutton R, *et al.* Qualitative assessment of caregiver experiences when navigating childhood immunisation in urban communities in Sierra Leone. *BMJ Open* 2022;**12**:e058203. doi:10.1136/bmjopen-2021-058203

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058203>).

Received 11 October 2021

Accepted 31 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Mohamed F Jalloh; yum8@cdc.gov

ABSTRACT

Objective To gain in-depth understanding of the caregiver experience when navigating urban immunisation services for their children.

Design An exploratory qualitative assessment comprising 16 in-depth interviews using an interpretative phenomenology approach.

Setting Caregivers were purposively recruited from slums (n=8) and other urban communities (n=8) in the capital city of Sierra Leone.

Participants Caregivers of children ages 6–36 months old who were fully vaccinated (n=8) or undervaccinated (n=8).

Results Emotional enablers of vaccination were evident in caregivers' sense of parental obligation to their children while also anticipating reciprocal benefits in children's ability to take care of their parents later in life. Practical enablers were found in the diversity of immunisation reminders, information access, information trust, getting fathers more involved, positive experiences with health workers and postvaccination information sharing in the community. Underlying barriers to childhood vaccination were due to practical constraints such as overcrowding and long waiting times at the clinic, feeling disrespected by health workers, expecting to give money to health workers for free services and fear of serious vaccine side effects. To improve vaccination outcomes, caregivers desired more convenient and positive clinic experiences and deeper community engagement.

Conclusions Health system interventions, community engagement and vaccination outreach need to be tailored for urban settings. Vaccine communication efforts may resonate more strongly with caregivers when vaccination is framed both around parental responsibilities to do the right thing for the child and the future benefits to the parent.

INTRODUCTION

There have been efforts to understand urban immunisation challenges in low-income and middle-income countries (LMICs), including in urban slums and informal settlements.¹ Assessment of immunisation barriers in urban

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This qualitative assessment describes the caregiver experience in navigating childhood immunisation services in urban settings in a low-income country to gain insights for improving vaccination outcomes.
- ⇒ The sample composed of caregivers of fully vaccinated and undervaccinated children who were purposively recruited to help understand how the two outcomes may be shaped by differences in the caregiver experiences.
- ⇒ The rarity of experiencing serious side effects among children in our sample limits the ability to have a rich understanding of such experiences on subsequent vaccination uptake.

areas in LMICs identified a range of practical and social issues, such as population mobility, inaccurate denominators of children due to out-of-date population estimates and poorly defined geographical catchment areas, the lack of trust in the health system among vulnerable groups, overburdened health facilities, and weak community engagement and outreach.²

In Sierra Leone, the 2014–2016 Ebola epidemic disrupted the delivery of essential health services, including immunisation services, especially in urban areas.^{3,4} Barriers that affected routine health services included the fear of contracting Ebola in health facilities, stigmatisation of health workers and shifting of resources to the epidemic.⁵ As the Ebola epidemic waned, measles outbreaks became more frequent due to the decline in measles vaccination.⁶ In the aftermath of the Ebola epidemic, the Government of Sierra Leone and its partners made major investments to rebuild health systems and restore public confidence in the healthcare system.^{7,8} However, challenges in access to

and the uptake of essential immunisation services persist, including in urban areas.⁹

UNICEF and partners developed the Caregiver Journey Framework to guide countries in understanding the experiences, processes and structures that shape how caregivers seek and receive health services for their children, including essential immunisation.¹⁰ In 2018, we operationalised the Caregiver Journey Framework through a qualitative approach in the Western Area Urban district (WAU) in Sierra Leone. Implementation experiences from operationalising the framework in the context of urban immunisation has been described elsewhere.¹¹ The framework was operationalised into several domains to understand decision making and preparation for vaccination visits, making the journey to clinics, experiences during vaccination visits and postvaccination experiences. Building on these domains, we aimed to understand the real-world experiences of caregivers of vaccine-eligible children as they navigate urban immunisation services in Sierra Leone to identify vaccination enablers and barriers.

METHODS

We developed the Immunisation Caregiver Journey Interviews (ICJI) approach¹¹ based on the Caregiver Journey Framework using principles of interpretative phenomenology,^{12 13} which focuses on elucidating the essence of common experiences to explain, interpret and make sense of a phenomenon.¹⁴ We used a phenomenological approach to explore the lifeworld of caregivers in how they navigate childhood immunisation for their children repeatedly in low-resource, urban settings.¹¹ A semi-structured ICJI guide was used to explore the following domains: Decision making and preparation, making the journey, experiences during vaccination visit, postvaccination experiences, intentions to return and perceptions of immunisation promotion activities in the community.

Setting

The WAU district in Sierra Leone comprises most of the capital city of Freetown with approximately 1.2 million inhabitants.¹⁵ The district was heavily affected by the Ebola epidemic, partly due to high population movements and crowded housing conditions.¹⁶ On average, there is less than one medical doctor per 10 000 population.¹⁷ The Government of Sierra Leone introduced the Free Health Care Initiative in 2010 to remove cost barriers for essential health services for pregnant and lactating mothers and under-5 children.¹⁸ Childhood immunisation services are delivered through the Expanded Programme on Immunisation using fixed sites that are supplemented by community outreach services to be conducted five times monthly.¹⁹ Each catchment community has 10 community health workers (CHWs) who support the promotion of health services on a voluntary basis.²⁰ A coverage survey in 2019 estimated 86% coverage for three doses of diphtheria–pertussis–tetanus vaccine in slums and 92% coverage in non-slum urban areas in the WAU

district. However, coverage of the second dose of measles-containing vaccine was very low in the district (33% in slums and 29% in non-slum urban areas).⁹

Sampling and data collection

The sample size for this qualitative assessment was guided by an approach that focuses on qualitative information power.²¹ The concept of information power posits that researchers should determine the sample size in a qualitative assessment based on the aim (narrow vs broad), sample specificity (targeting specific group vs multiple groups), theoretical underpinning (application of theory or no theory), quality of dialogue (weak or strong) and analysis strategy (within-case only or cross-case). Sample size burden increases when the aim is broad, multiple groups are targeted in the sample, the assessment is theory driven, the quality of the dialogue is weak and transcripts are analysed using cross-case analysis. In our assessment, the aim was narrow, the sample targeted a specific group, we applied theory to guide the assessment, the transcripts contained rich information, and we conducted both within-case and cross-case analyses. Against these considerations, we interviewed 16 caregivers and progressively reviewed debrief notes from the interviews to assess information power. In analysing the transcripts, we concluded that we reached saturation with the 16 interviews and likely could have stopped interviewing after the 12th interview.

We purposively recruited the caregivers from eight communities in the WAU district, four of which were slums and four were other urban areas in the district to maximise variation in the sample. Within each community, two caregivers of children ages 6–36 months were selected to capture a breadth of experiences of caregivers with vaccine eligible children—one whose child was fully vaccinated for age and another whose child had missed at least one scheduled vaccination visit. CHWs supported data collection teams in visiting households to identify and recruit eligible caregivers in the selected communities. Snowball sampling was used as a secondary sampling strategy when the first identified caregiver declined to interview but knew of other caregivers in the community with vaccine-eligible children or when CHWs were only successful in identifying just one eligible caregiver. In this form of snowball sampling, a previously visited household with an eligible child would point data collectors to other households with potentially eligible children (ie, vaccine eligible children). Data collectors visited such households to screen for eligibility. This process continued until two caregivers of eligible children were successfully recruited and interviewed from a particular community. Interviews were conducted on the same day of recruitment after obtaining informed consent from the caregiver.

We recruited data collectors (interviewers and note-takers) who were fluent in English and the predominant local language in the WAU district (Krio). The data collectors had postsecondary educational training in social sciences and were experienced in conducting qualitative

data collection in Sierra Leone. Two behavioural scientists trained the facilitators for a week on the assessment protocol. One of the trainers was from Sierra Leone and had experience conducting social science research in Sierra Leone. During the training, the English version of the guide was translated into Krio by locally hired staff together with the facilitators and trainer. The data collectors were trained on how to probe on the spot to obtain additional pertinent information from caregivers. Data collection occurred in August–September 2018. All interviews were audiorecorded with permission from participants; they were then transcribed and translated into English by the local team. Interviews lasted about an hour on average and were conducted in the vicinity of the homes of the caregivers. Data collection teams were trained on choosing suitable interview locations to enable caregivers to speak freely. The facilitators conducted debriefing sessions immediately after each interview to make note of key experiences and observations. The debriefing notes were not part of the formal analysis. However, during the field work, the debriefing notes were used to progressively assess data saturation and to identify key insights emerging from the interviews. We used the insights from the debrief notes to develop a preliminary report that was mostly in a descriptive, narrative form. The deidentified preliminary report was shared with the Sierra Leone Ministry of Health and Sanitation. We have previously documented practical lessons learnt from implementing the assessment in Sierra Leone.¹¹

Data analysis

Two analysts (one male, one female) read all transcripts and created analytic memos and then analysed the transcripts using both within-case and cross-case analysis. In the within-case analysis, we developed a narrative profile for each caregiver to bring key aspects of their lived experiences to the foreground. In the initial part of the cross-case analysis, each analyst coded three different transcripts (six total), using an inductive approach to identify and interpret meaning units within the text. To gain alternative interpretations of the coded meaning units, three of the coded transcripts were shared with a third qualitative expert, who was not involved in the previous stages of the assessment, for independent ‘blind’ coding of the transcripts. Feedback from the third analyst was discussed by the two primary analysts and incorporated into the coding scheme. The analysts used an iterative process to review their codes, discuss their interpretations of the manifest content and harmonise the initial set of codes that were used for coding the remaining manuscripts. NVivo software (QSR International-2018, V.12) was used for the final organisation and coding of the transcripts. Manifest categories of meaning units were grouped to reflect latent content that was developed into cross-cutting themes via a consultative process. Throughout the process, the analysts exercised reflexivity regarding subjective interpretations and iteratively re-examined the transcripts to

identify alternative interpretations until consensus was reached with additional inputs from the co-authors.

Patient and public involvement

No patient involved.

RESULTS

All respondents were the biological mothers of the sampled children except for one female guardian. The median age was 9 months for the children included in the assessment. At the time of the interviews, half of the children had missed at least one scheduled vaccine dose. Three themes emerged from the interviews around vaccination enablers (table 1), vaccination barriers (table 2) and direct recommendation to improve vaccination uptake (table 3). There were no notable differences in themes between slums and other urban communities.

Enablers of childhood vaccination

Our analysis identified emotional and practical enablers related to childhood vaccination. Emotional enablers were evident in how caregivers portrayed their parental obligation to their children, wanting to do the right thing for their children’s health and anticipating reciprocal benefits in children’s ability to take care of the parents later in life. Practical enablers were the diversity of immunisation reminders, information access, information trust, getting fathers more involved, positive experiences with health workers and postvaccination information sharing at the community level (table 1).

Parental responsibility

A sense of parental responsibility was a major motivating factor for caregivers to seek vaccination services for their children. Caregivers viewed vaccination as ‘doing the right thing’ for their children. Even caregivers who had missed scheduled vaccination visits felt responsible for getting their children caught up with their scheduled vaccine doses, which often required deprioritising other income-generating activities.

It is my duty to take my baby to the hospital for immunisation. It is my responsibility as [a] mother to ensure that my baby completes the rounds of immunisation without defaulting.—Caregiver whose child had missed a scheduled vaccination

Wanting a strong and healthy baby

In addition to the affective responses regarding a sense of duty to the child, appreciating the overall health benefits of vaccination to their children was another major driving force in motivating caregivers to seek vaccination services. Caregivers consistently expressed that immunisation has essential health benefits to the child and that missing scheduled vaccination would ‘risk the baby’s life.’ Moreover, they valued having a ‘strong and healthy baby’ and felt that completing the vaccination schedule would positively impact the baby’s health.

**Table 1** Enablers of childhood vaccination—Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Sub-themes	Theme
Responsibility to ensure full vaccination	Parental responsibility	Emotional enablers of childhood vaccination	Enablers of childhood vaccination
Obligation to do the right thing			
Immunisation as a requirement			
Immunisation is important for baby's health	Wanting a 'strong and healthy baby'		
Wanting 'strong baby'			
'Health is wealth'			
Defaulting on vaccination risks baby's life	Parental anticipation of reciprocal benefit		
Vaccination benefits the parent later in life			
Taking care of parents when old	Diversity of immunisation reminders	Practical enablers of childhood vaccination	
Immunisation card is important			
Campaign as reminder			
Reminders by health workers at vaccination visit	Information access and information trust		
Husband as reminder			
Immunisation card as reminder			
Other family member as reminder			
Nurses should lead			
Nurses more trusted than CHWs			
Same information from different sources	Getting fathers more involved		
Immunisation promotion through radio/tv			
Immunisation promotion by health workers			
Immunisation promotion through leaders			
Immunisation promotion by NGOs			
Mothers take the child to the clinic			
Fathers rarely involved	Positive experiences with health worker		
Father received an award for involvement			
Cordial relationship with nurses			
Good care by nurses	Post-vaccination information sharing		
HWs encourage seeking care at HF			
Giving money to health worker as token of appreciation			
Husband asking about the visit	Post-vaccination information sharing		
Informing husband of next visit			
Telling husband about visit expenses			
Immunisation is a 'learning process'			
Sharing experiences with neighbours or friends			
Other family members asking about the visit			

CHWs, community health workers; HF, Health facility; HWs, Health workers; NGOs, Non-governmental organisations.

I think [the] vaccine is good for our children. It is important and it helps to build their immune system to keep them strong and healthy; it fights against many things in the body...—Caregiver whose child was fully vaccinated

Parental anticipation of reciprocal benefit

Perceptions of vaccination benefits went beyond the direct health benefits to children and extended into domains of benefit to the parent. The notion that vaccinated children

will be healthier and in turn live longer and be able to take care of their parents later in life emerged as a dimension of vaccination benefit to the parent. This duality of vaccination benefit was grounded in the cultural context of parents expecting reciprocal care from their children when the parents can no longer care for themselves.

It is very difficult for me to be absent for immunisation or not to take my baby to the hospital when sick. Sometimes people think I'm mad but I'm not.

Table 2 Barriers related to childhood vaccination—Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Subthemes	Themes	
Asking husband for money	Preparing for the journey and getting to the clinic	Practical constraints	Barriers related to childhood vaccination	
Juggling different household duties				
Competing priorities				
Time taken to get to the clinic				
Wasting time at the clinic				Inconveniences at the clinic
Rush to arrive first and seen first				
Crowding at the clinic				
Very long wait at larger health facilities				
Spent a day and nothing happened				
Wasting caregiver's time				Feeling disrespected by health workers
Not respecting caregivers				
Shouting at caregivers	Monetary expectations			
Paying for the card				
Paying for free drugs				
Paying for weighing				
Payment as punishment				
Bad care without payment				
Health workers should stop demanding money				
Health worker don't appreciate less than Le 2000				
Health workers withholding free drugs	Vaccine side effects	Safety concerns		
Baby crying throughout the night				
Baby gets 'lazy' for few hours				
Fever in baby				
Swelling at injection site				
Side effects only for some vaccines				
Rub onion on swollen injection site				
Avoiding abnormalities from vaccine				
Afraid of vaccine side effects				

I'm trying to bring up my children in a way that they will benefit me when I'm old.—Caregiver whose child had missed a scheduled vaccination

Diversity of immunisation reminders

Caregivers were exposed to numerous reminders and cues to vaccinate their children, including information from the immunisation cards, health workers, community campaigns and family members. However, the child's immunisation card stood out as the most important reminder that caregivers and families relied on to remember the dates of the scheduled vaccination visits.

I do remind myself because they [health workers] will write at the back of the immunisation card the date for the next visit. The immunisation card will tell you the date for the next visit.—Caregiver whose child had missed a scheduled vaccination

In addition, community outreach campaigns and announcements in the community were also viewed as helpful reminders to vaccinate.

The health workers in this community and the CHWs are doing well as they do go around telling people not to forget to take their babies to the hospital for immunisation.—Caregiver whose child had missed a scheduled vaccination

Information access and information trust

Immunisation information sources varied, but caregivers consistently cited health workers as trusted sources of information. Nurses were more trusted than lay CHWs because caregivers viewed nurses as more knowledgeable. While waiting in line before immunisation services, most caregivers appreciated the 'health talk' they received from nurses who advised on health and immunisation.

Table 3 Recommendations to improve childhood vaccination— Western Area Urban, Sierra Leone, 2018

Meaning unit	Category	Subtheme	Theme
Continue campaigns at repeated intervals	Improving vaccination processes and systems	Caregivers want improved vaccination processes, systems and engagement	Recommendations to improve childhood vaccination
Do not rely on campaigns alone			
Provision of incentives for caregivers			
Concentrate on defaulters			
Compensation/incentives for nurses			
Being considerate towards health workers			
Employ more staff			
Stop demanding money			
Promote consequences of not vaccinating	Engaging communities to boost vaccine confidence		
Peer-to-peer promotion of immunisation			
Inform about importance of immunisation			
Address ‘stubborn’ caregivers			
Engage caregivers who do not want injections			

I trust them [nurses] because they are a team of qualified nurses ... Before they come here, they know everything about the vaccines and any implications of the vaccines. They are able to explain more than the CHWs and other community workers in the area. The nurses will tell you more. There are things that the community workers do not know, and they refer you to the nurses.—Caregiver whose child was fully vaccinated

Getting fathers more involved

Mothers desired greater involvement by the child's fathers in supporting their children's vaccination. In most instances, fathers were rarely involved in taking their children to the vaccination clinic. We uncovered only two instances when fathers actively supported their children's vaccination visits. In those instances, the mothers felt supported, and their children were fully vaccinated. In one situation, a father that routinely accompanied the child to the vaccination clinic was celebrated by health workers and given the ‘best father’ award.

There was [a] time when he [my husband] was given the best father award [at the clinic] because he is always with me at the hospital. That is the name I have also given to my husband. Even when the baby is crying, I will say ‘best father’ take your baby.—Caregiver whose child was fully vaccinated

Positive experiences with health workers

During vaccination visits, positive experiences with health workers encouraged caregivers to vaccinate their children. Positive experiences included having a cordial relationship with the health workers. In particular, caregivers expressed that the nurses took good care of their children, made them feel comfortable and tried to build a good rapport. Some caregivers said they would voluntarily give small amounts of money to health workers as a token of their appreciation after vaccinating the child.

...the nurses do encourage you and will make jokes so that you will laugh at the end of the day. There is a lot of fun [interactions], which made some of us forget about our stresses.—Caregiver whose child was fully vaccinated

Postvaccination information sharing

Information exchange at the community level with trusted community members strengthened caregiver confidence in childhood vaccination. On returning home from the vaccination visit, caregivers often discussed the clinic experience with their spouses, families and other caregivers in the community. In addition to information obtained from health workers at the clinic, caregivers also sought advice from other ‘more experienced’ caregivers in the community.

Most times after immunisation, my baby will run a temperature, but the health workers always provide

drugs to counter the fever. We have caregivers in this community with vast knowledge and since this is my first baby, I love to talk to them so that we can share our experiences which will be of advantage to me as I'm very new in the field [of parenting].—Caregiver whose child was fully vaccinated

Barriers related to childhood vaccination

Practical constraints, negative experiences with health workers and safety concerns were the underlying barriers to childhood vaccination. Practical constraints included challenges faced when preparing for and getting to the vaccination clinic and inconveniences encountered at the clinic, such as overcrowding and long waiting times. Negative experiences among caregivers included feeling disrespected by health workers while simultaneously expecting to give money to health workers for services that are supposed to be free of charge. Finally, vaccine side effects led to concerns and fears about vaccine safety (table 2).

Preparing for the journey and getting to the clinic

Caregivers commonly cited the need to juggle 'household duties' and other income-generating activities when planning the visit as a barrier, especially in the absence of fathers' involvement in taking the child to the clinic. In addition, mothers frequently depended on their children's fathers for financial support to cover the expenses related to the vaccination clinic visit. Some caregivers recounted needing to travel long distances up to an hour by foot to get to the vaccination clinic, especially when they could not afford public transportation.

Inconveniences at the clinic

Caregivers anticipated various inconveniences at the vaccination visit. The prolonged time spent waiting for the child to be vaccinated emerged as a substantial inconvenience that was more pronounced when seeking immunisation services, especially in larger facilities. Anticipating the long wait, caregivers usually tried to arrive early at the vaccination site to get seen first. The range of activities involved with the child health visit prolonged the visit, including weighing the baby as part of growth monitoring and other health checks.

It is painful if you waste much time at the health facility because you have other issues to attend to. To avoid that, that is why I always come early to the health facility.—Caregiver whose child was fully vaccinated

Feeling disrespected by health workers

Caregivers often felt disrespected by health workers during the vaccination visit. A key complaint was that health workers shouted at caregivers and sometimes used vulgar language toward caregivers. In other instances, they complained that some health workers habitually arrived late to the vaccination session, which further prolonged the time caregivers spent waiting.

Monetary expectations

Systemically hidden costs generated substantial dissatisfaction among caregivers. Caregivers needed to 'shake hands' with health workers at different times of the visit (eg, first time registering the child to get a card, before entering the facility and before weighing the baby). Shaking hands implied giving some money during the handshake. Caregivers used the money to 'fast-track' their children's vaccination. The expectations around monetary exchange discouraged caregivers who could not afford to shake hands with health workers.

Sometimes if I don't want to spend much time at the hospital, I will shake the hand of the nurse so that they can fast track the immunisation of my baby. I will give them something like two thousand Leones or whatever I have with me at that moment... Health is wealth and they [health workers] don't need us but we do [need them]. The money we give is nothing compared to the health of our children... At the end of the day, we will grumble on our way home as the services are supposed to be free for our children, yet we are paying for it. The health workers are really trying, but the idea for them to take money from us is bad. And if you don't give them money, they will talk to you carelessly.—Caregiver whose child was fully vaccinated

In a separate domain of monetary exchange, health workers demanded money as a form of 'punishment' to caregivers who missed their children's scheduled vaccination appointments.

If you failed to take your baby to the hospital on [the] stipulated date, you will definitely have to pay some amount at the end of the day in the form of punishment. You must pay five thousand or more.—Caregiver whose child was fully vaccinated

Vaccine side effects

Caregivers cited numerous instances when their children experienced vaccine side effects such as 'fever,' 'swelling at the injection site,' and the 'baby becoming lethargic.' Fever was the most common side effect, and the caregivers knew to administer fever-reducing medication as instructed by health workers. When there was swelling at the injection site, a common practice among caregivers was to massage the swollen area, sometimes with an onion or a bar of soap to try to reduce the swelling.

Sometimes my baby's leg becomes swollen... because some nurses are heavy-handed, and I meet several nurses when I visit the hospital. Sometimes the leg gets swollen, and they treat him. I have to rub the leg to avoid swelling... I use soap to rub off the swelling and I give Panadol to stop the fever... some people say you should not allow every nurse to administer [an]injection to the child. I should have a permanent nurse that gives injection to my child without

swelling.—Caregiver whose child missed a scheduled vaccination

In one rare situation, a caregiver had a prior child who experienced fever, convulsed and died a few days after getting vaccinated. Therefore, the caregiver decided to not vaccinate subsequent children.

After the immunisation, my baby started running temperature, I administered paracetamol as I was told by the health workers. The baby convulsed and that was the end of that baby. I don't want a repeat of that in my life. I have therefore decided not to take my babies for immunisation anymore.—Caregiver whose child missed a scheduled vaccination

Besides passive acceptance of the BCG vaccine at the birth of the youngest child, this same caregiver actively refused all other vaccines despite encouragement by a family member to vaccinate the child.

I'm not outrightly saying it was as a result of the immunisation [that my child died]; as every death is the work of God. But from what I have gathered so far, I have personally decided not to take my baby to the health facility to be immunized. It is not that I'm tired of going to the health facility or because of the distance or money. I do get a lot of pressure from my aunt to take my baby to the health facility for immunisation, but the thing is that I just don't trust the system and what immunisation does.—Caregiver whose child missed a scheduled vaccination

From the perspective of this same caregiver, the vaccinated child died but the unvaccinated children survived and thrived, which was cited as a reason for refusing vaccination.

I believe in exclusive breastfeeding, sometimes for two years and a half and sometimes [only] two years. My baby is now two years seven months old and doing well like any of those children that are on immunisation or have completed... Just as I was saying, sometimes my heart will tell me to take the baby for immunisation but after thinking of the past experience, I would decide not to. I'm now used to that... The simple fact here is that, since the other children are doing well without immunisation, I will not take [the baby] to the health facility for immunisation and that is all.—Caregiver whose child missed a scheduled vaccination

Recommendations to improve childhood vaccination

The direct recommendations provided by caregivers were categorised into (1) improving vaccination process and systems and (2) engaging communities to boost vaccine confidence. Implementing community outreach campaigns for immunisation at regular intervals with a focus on defaulters was recommended by participants to improve vaccination outcomes. In addition,

caregivers wanted health workers and community leaders to be involved in immunisation promotion along with the CHWs. They wanted the vaccination clinic experience to improve and become more conducive to caregivers, including shorter wait time at the clinic and more positive interactions with health workers. Lastly, caregivers wanted health workers to stop demanding money from them, though they may not mind giving money, out of free-will, as a token of appreciation when they could afford it (table 3).

In addition, you should engage the Chiefs, because in each area we have Chiefs to spread out this message. You could educate them so that they in turn can educate those in the community. Let us have Town Criers go around disseminating the messages. It would be nice for them to allocate people in the health centre who move from house to house to educate the breastfeeding mothers because some of us are stubborn to come onboard.—Caregiver whose child was fully vaccinated

DISCUSSION

Our qualitative analysis highlighted several important themes. In the backdrop of anticipated benefits to both the child and parent, vaccination intention was motivated by a feeling of parental responsibility to 'do the right thing.' Timely and trusted exchange of information together with social support and positive experiences at the vaccination clinic were important facilitators of vaccination. In contrast, vaccination was discouraged by negative interactions with health workers at the clinic, the occurrence and fear of vaccine side effects, multitude of 'hidden' costs, juggling vaccination with other responsibilities and inconveniences, such as long travelling time to the clinic and long delays at the clinic. Nevertheless, caregivers were resilient in devising ways to try to get their children vaccinated such as walking on foot up to an hour to get to the clinic when they could not afford public transportation. Lastly, caregivers wanted the vaccination experience to improve, and they desired stronger community engagement to help optimise vaccination outcomes. However, systemic issues, such as informal payments, overcrowding in health facilities and the reported overburdened health workers may require interventions at the health systems level. These themes from Sierra Leone provide in-depth insights regarding the motivations, facilitators and barriers of vaccination in an urban LMIC setting.

Moral values may shape vaccination attitudes.²² Philosophical arguments regarding the morality of vaccination have been heavily debated.^{23–26} Caregivers in our sample largely viewed vaccination via a moral lens encompassing parental duty to do the right thing for the child. In one situation, however, we found that the desire to 'do the right thing' may also translate into vaccination refusal in the backdrop of other past refusals, observing 'healthy

unvaccinated' children and having distrust of the health system. Quantitative research from high-income countries has shown that parents with unvaccinated children were more likely to perceive their children to be at low risk of vaccine-preventable diseases and were more likely to perceive low vaccine effectiveness and safety compared with parents with vaccinated children.²⁷ Our findings suggest that childhood immunisation communication efforts may resonate more strongly with caregivers when vaccination is framed around parental responsibilities to do the right thing for the child and the anticipated future benefits to parents. However, additional research is necessary to generate a better understanding of the morality of childhood vaccination in the Sierra Leonean context.

Across the interviews, there was an apparent tension in the relationship between caregivers and health workers. Caregivers often expressed their appreciation of health workers and empathised with the challenging context in which they do their work. Health workers were strongly viewed as authoritative sources of trusted information regarding immunisation and the child's health, which is consistent with findings from high-income countries^{28–30} and LMICs.^{31 32} Our findings on the role of monetary exchange in vaccination exemplify the complex relationship between caregivers and health workers in low-resource urban communities in Sierra Leone. Some caregivers voluntarily gave money to health workers as a 'token of appreciation' while others begrudgingly gave money because they viewed it as a condition for receiving good quality service from health workers. Interventions at the health systems level are necessary to help discourage informal payments to health workers—a practice that may perpetuate vaccination inequities among poor caregivers who are unable to meet monetary expectations.

Our findings also illuminate the need for interventions at the household and family level. Fathers were rarely involved in taking their children to the vaccination clinic but were often engaged in the decision-making processes. In the few instances when fathers were involved in taking their children to the clinic, the mothers felt supported and their children were fully vaccinated. A study in Nigeria found that paternal involvement in immunisation was greater in rural settings compared with urban settings.³³ In urban areas, the same study found that paternal involvement was greater among educated fathers compared with uneducated fathers. In a separate study in Ghana, the involvement of educated fathers in the vaccination decision was associated with timelier vaccination uptake compared with the involvement of uneducated fathers in the decision.³⁴ More broadly, shifting from a mother–child dyad to a family triad in the care of children has proven to have positive effects on paediatric health outcomes across diverse contexts.³⁵ Additional assessments and interventions are needed to explore and evaluate culturally appropriate ways to enhance the involvement of fathers in childhood immunisation in Sierra Leone and other similar LMIC settings.

Existing evidence suggests that vaccine safety concerns, often linked to adverse events following immunisation (AEFI), contribute to vaccine hesitancy.^{36 37} Serious AEFIs may be 'triggering events' for derailing vaccine confidence and prompting active refusal among certain caregivers and their communities—especially when the serious AEFI is perceived to be linked to the vaccine or the vaccination process.³⁸ Together with prior evidence, these findings emphasise the need for robust AEFI surveillance and investigations³⁹ to identify, counsel and follow-up with caregivers whose children experience AEFI, and therefore, are potentially at risk of missing subsequent vaccination. Health workers and CHWs may benefit from periodic in-service training on how to effectively communicate vaccine safety and address concerns about AEFIs.⁴⁰

Limitations

There were several limitations to our assessment. First, it is possible that some nuanced meaning may have been lost when translating the audio recordings from Krio to English—especially since the transcripts were not back-translated from English to Krio due to resource constraints. Second, we only identified one caregiver who routinely and actively refused all vaccines for her child, which may reflect the overall rarity of zero-dose unvaccinated children in Sierra Leone (approximately 3%).⁴¹ This was the only caregiver with a child that experienced a serious AEFI in our sample, which limits our ability to have a rich understanding of such experience among caregivers more broadly and the potential linkages to vaccination refusal. Although such experiences of serious AEFI are rare, they may have the tendency to get publicised in the community when they do occur, which may negatively influence vaccination decisions among other caregivers. Taken together, our results call for additional qualitative assessments to get a deeper understanding of vaccination refusal within the Sierra Leonean context and other low-resource LMIC settings. Sampling strategies may, therefore, need to be adapted accordingly to focus on caregivers who actively refuse vaccination for their children. Lastly, our findings reflected gender dimensions that may be based on sociocultural norms in Sierra Leonean society but may also be due to sampling bias because the caregivers we conveniently recruited were mostly stay-at-home mothers.

CONCLUSIONS

As the COVID-19 pandemic disrupts childhood immunisation globally,⁴² especially in LMICs, our assessment provides a foundational understanding of the challenges that caregivers encounter in urban settings in Sierra Leone. It also sheds light on opportunities to improve vaccination outcomes in urban poor settings, which is a global immunisation priority. The findings show that health system interventions, community engagement and vaccination outreach may need to be tailored for urban LMIC settings. Vaccine communication efforts may

resonate more strongly with caregivers when vaccination is framed both around parental responsibilities to do the right thing for the child and the future benefits to the parent.

Author affiliations

¹Global Immunization Division, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

²Oak Ridge Institute for Science and Education, Oak Ridge, Tennessee, USA

³ICAP at Columbia University, New York, New York, USA

⁴ICAP at Columbia University, Freetown, Sierra Leone

⁵School of Public Health, The University of Sydney, Sydney, New South Wales, Australia

⁶Expanded Programme on Immunization and Child Health, Sierra Leone Ministry of Health and Sanitation, Freetown, Western Area, Sierra Leone

⁷Department of Epidemiology, Columbia University Mailman School of Public Health, New York, New York, USA

Acknowledgements We thank the 16 caregivers who participated in the in-depth interviews.

Collaborators SLU Immunisation Needs Assessment Group: Adewale Akinjeji, Laura Conklin, Oliver Eleeza, Lauren E Parmley, Anthony Mansaray, Dimitri Prybylski, Aaron Wallace, Brent Woff.

Contributors MFJ, RS, MT, TS and ML designed the assessment. MFJ, PP, SK and KW analysed the data. MFJ, PP, RS, SK, MT, KW, TS and ML contributed to the interpretation of the results. MFJ, PP and SK drafted the manuscript. RS, MT, KW, TS and ML contributed to revising the manuscript. All authors reviewed and approved the final manuscript. MFJ and PP are responsible for the overall content as the guarantor.

Funding US Centers for Disease Control and Prevention (U19GGH001581).

Disclaimer The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the US Centres for Disease Control and Prevention.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The assessment was approved by the Sierra Leone Ethics and Scientific Review Committee (SLESRC-17052018), Columbia University Medical Centre Institutional Review Board (IRB-AAAR9031) and the Centre for Global Health at the US Centers for Disease Control and Prevention (CGH-2018-184). All participants provided written informed consent.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Mohamed F Jalloh <http://orcid.org/0000-0002-7206-8042>

Shibani Kulkarni <http://orcid.org/0000-0003-4999-570X>

REFERENCES

- Nelson KN, Wallace AS, Sodha SV, *et al.* Assessing strategies for increasing urban routine immunization coverage of childhood vaccines in low and middle-income countries: a systematic review of peer-reviewed literature. *Vaccine* 2016;34:5495–503.
- John Snow International. Strengthening immunization for the urban poor: experiences and lessons learned from five country studies, 2018. Available: <https://www.jsi.com/resource/strengthening-immunization-for-the-urban-poor-experiences-and-lessons-learned-from-five-country-studies/> [Accessed 12 Nov 2020].
- Brolin Ribacke KJ, Saulnier DD, Eriksson A, *et al.* Effects of the West Africa ebola virus disease on health-care utilization - a systematic review. *Front Public Health* 2016;4:222.
- Elston JWT, Cartwright C, Ndumbi P, *et al.* The health impact of the 2014–15 Ebola outbreak. *Public Health* 2017;143:60–70.
- Yerger P, Jalloh M, Coltart CEM, *et al.* Barriers to maternal health services during the Ebola outbreak in three West African countries: a literature review. *BMJ Glob Health* 2020;5:e002974.
- Masresha BG, Luce R, Weldegebriel G, *et al.* The impact of a prolonged Ebola outbreak on measles elimination activities in guinea, Liberia and Sierra Leone, 2014–2015. *Pan Afr Med J* 2020;35:8.
- Government of Sierra Leone. National Ebola recovery strategy for Sierra Leone, 2015. Available: https://ebolaresponse.un.org/sites/default/files/sierra_leone_-_national_recovery_strategy_2015-2017.pdf [Accessed 5 Dec 2020].
- U.S. Centers for Disease Control and Prevention. Global health – Sierra Leone, 2019. Available: <https://www.cdc.gov/globalhealth/countries/sierra-leone/default.htm> [Accessed 5 Dec 2020].
- Feldstein LR, Sutton R, Jalloh MF, *et al.* Access, demand, and utilization of childhood immunization services: a cross-sectional household survey in Western area urban district, Sierra Leone, 2019. *J Glob Health* 2020;10:010420.
- UNICEF. Human centered design 4 health, 2018. Available: <https://www.hcd4health.org/> [Accessed 18 Nov 2020].
- Jalloh MF, Kinsman J, Conteh J, *et al.* Barriers and facilitators to reporting deaths following Ebola surveillance in Sierra Leone: implications for sustainable mortality surveillance based on an exploratory qualitative assessment. *BMJ Open* 2021;11:e042976.
- Diekelmann NL. Learning-as-testing: a Heideggerian hermeneutical analysis of the lived experiences of students and teachers in nursing. *ANS Adv Nurs Sci* 1992;14:72–83.
- Dreyfus HL. *Being-in-the-world: a commentary on Heidegger's being and time*. Cambridge: The MIT Press, 1991.
- Dahlberg K. The essence of essences – the search for meaning structures in phenomenological analysis of lifeworld phenomena. *Int J Qual Stud Health Well-being* 2006;1:11–19.
- Statistics Sierra Leone. Sierra Leone population and housing census, 2016, 2015. Available: <https://www.statistics.sl/index.php/census/census-2015.html> [Accessed 12 Nov 2020].
- Lamunu M, Olu OO, Bangura J, *et al.* Epidemiology of Ebola virus disease in the Western area region of Sierra Leone, 2014–2015. *Front Public Health* 2017;5:33.
- World Health Organization. Medical doctors per 10,000 population, 2021. Available: <https://www.who.int/data/gho/data/indicators/indicator-details/GHO> [Accessed 5 Dec 2021].
- Witter S, Brikci N, Harris T, *et al.* The free healthcare initiative in Sierra Leone: evaluating a health system reform, 2010–2015. *Int J Health Plann Manage* 2018;33:434–48.
- Government of Sierra Leone. Comprehensive EPI multi-year plan 2017–2021, 2016. Available: <https://www.gavi.org/sites/default/files/document/2021/cMYP%20Sierra%20Leone%202017-2021.pdf> [Accessed 5 Dec 2020].
- Government of Sierra Leone. Policy for community health workers in Sierra Leone, 2012. Available: <https://chwcentral.org/wp-content/uploads/2015/01/CHW-Policy-Sierra-Leone.pdf> [Accessed 5 Dec 2020].
- Malterud K, Siersma VD, Guassora AD. Sample size in qualitative interview studies: guided by information power. *Qual Health Res* 2016;26:1753–60.
- Amin AB, Bednarczyk RA, Ray CE, *et al.* Association of moral values with vaccine hesitancy. *Nat Hum Behav* 2017;1:873–80.
- Di Pietro ML, Poscia A, Teleman AA, *et al.* Vaccine hesitancy: parental, professional and public responsibility. *Ann Ist Super Sanita* 2017;53:157–62.
- Giubilini A, Douglas T, Savulescu J. The moral obligation to be vaccinated: utilitarianism, contractualism, and collective easy rescue. *Med Health Care Philos* 2018;21:547–60.
- Bester JC. Measles vaccination is best for children: the argument for relying on herd immunity fails. *J Bioeth Inq* 2017;14:375–84.
- Dawson A. What are the moral obligations of the traveller in relation to vaccination? *Travel Med Infect Dis* 2007;5:90–6.
- Omer SB, Salmon DA, Orenstein WA, *et al.* Vaccine refusal, mandatory immunization, and the risks of vaccine-preventable diseases. *N Engl J Med* 2009;360:1981–8.
- Frew PM, Chung Y, Fisher AK, *et al.* Parental experiences with vaccine information statements: implications for timing, delivery, and parent-provider immunization communication. *Vaccine* 2016;34:5840–4.

- 29 Drezner D, Youngster M, Klainer H, *et al.* Maternal vaccinations coverage and reasons for non-compliance - a cross-sectional observational study. *BMC Pregnancy Childbirth* 2020;20:541.
- 30 Wiley KE, Massey PD, Cooper SC, *et al.* Pregnant women's intention to take up a post-partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: a cross sectional survey. *Vaccine* 2013;31:3972–8.
- 31 Makwe C, Anorlu R. Knowledge of and attitude toward human papillomavirus infection and vaccines among female nurses at a tertiary hospital in Nigeria. *Int J Womens Health* 2011;3:313–7.
- 32 Foss HS, Oldervoll A, Fretheim A, *et al.* Communication around HPV vaccination for adolescents in low- and middle-income countries: a systematic scoping overview of systematic reviews. *Syst Rev* 2019;8:190.
- 33 Sodeinde K, Amoran O, Abiodun O, *et al.* A rural/urban comparison of paternal involvement in childhood immunisation in Ogun central Senatorial district, Nigeria. *Niger Postgrad Med J* 2020;27:336–42.
- 34 Brugha RF, Kevany JP, Swan AV. An investigation of the role of fathers in immunization uptake. *Int J Epidemiol* 1996;25:840–5.
- 35 Allport BS, Johnson S, Aqil A, *et al.* Promoting father involvement for child and family health. *Acad Pediatr* 2018;18:746–53.
- 36 Dubé E, Laberge C, Guay M, *et al.* Vaccine hesitancy: an overview. *Hum Vaccin Immunother* 2013;9:1763–73.
- 37 Larson HJ, Jarrett C, Eckersberger E, *et al.* Understanding vaccine hesitancy around vaccines and vaccination from a global perspective: a systematic review of published literature, 2007-2012. *Vaccine* 2014;32:2150–9.
- 38 Du F, Chantler T, Francis MR, *et al.* The determinants of vaccine hesitancy in China: a cross-sectional study following the Changchun Changsheng vaccine incident. *Vaccine* 2020;38:7464–71.
- 39 World Health Organization. Adverse events following immunization (AEFI), 2021. Available: https://www.who.int/immunization/monitoring_surveillance/who-immuniz.pdf [Accessed 18 Feb 2021].
- 40 World Health Organization. Vaccine safety basics e-learning course, 2021. Available: <https://vaccine-safety-training.org/simplified-and-key-messages.html> [Accessed 19 Feb 2021].
- 41 UNICEF. Sierra Leone multiple indicators cluster survey, 2017. Available: https://mics.unicef.org/news_entries/106/SIERRA-LEONE-2017-MICS-RELEASED [Accessed 21 Jan 2021].
- 42 Abbas K, Procter SR, van Zandvoort K, *et al.* Routine childhood immunisation during the COVID-19 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2 infection. *Lancet Glob Health* 2020;8:e1264–72.