

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Diabetes among adults in Bangladesh: Changes in prevalence and risk factors from 2011 to 2018

| | |
|-------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2021-055044 |
| Article Type: | Original research |
| Date Submitted by the Author: | 01-Jul-2021 |
| Complete List of Authors: | Chowdhury, Muhammad Abdul Baker; University of Florida, Emergency Medicine Islam, Mirajul; University of Dhaka, Statistics Rahman, Jakia; BRAC University, Mathematics and Natural Sciences Uddin, Md. Jamal; Shahjalal University of Science and Technology, Statistics Rabiul Haque, Md; University of Dhaka, Population Sciences |
| Keywords: | Epidemiology < TROPICAL MEDICINE, DIABETES & ENDOCRINOLOGY, General diabetes < DIABETES & ENDOCRINOLOGY, PUBLIC HEALTH |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Diabetes among adults in Bangladesh: Changes in prevalence and risk factors from 2011 to 2018

Short title

Diabetes among adults in Bangladesh

Authors

Muhammad Abdul Baker Chowdhury, MPH, MPS, MS^{1*}; Mirajul Islam, MS² Jakia Rahman, BS³; Md Jamal Uddin, PhD⁴⁺; Md Rabiul Haque, PhD⁵⁺

Affiliations

1. Department of Emergency Medicine, College of Medicine, University of Florida, Gainesville, FL
2. Department of Statistics, University of Dhaka, Dhaka, Bangladesh
3. Department of Mathematics and Natural Sciences, BRAC University, Dhaka, Bangladesh
4. Department of Statistics, Shahjalal University of Science & Technology, Sylhet-3114, Bangladesh
5. Department of Population Sciences, University of Dhaka, Dhaka, Bangladesh

⁺Senior authors

* Corresponding author

* Corresponding author email: mchow023@fiu.edu

Tables and Figures: 3 Tables 2 Figures and 2 Supplemental Tables

Abstract Word Count: 264

Manuscript word count: 3440

Abstract

Objective: To investigate the change in the prevalence and risk factors of diabetes among adults in Bangladesh between 2011 and 2018.

Design: The study was conducted using nationally representative two waves of the Bangladesh Demographic and Health Survey (BDHS) in 2011 and 2017-18.

Setting: Bangladesh

Participants: Adults age 18 years and older.

Primary outcome: Diabetes mellitus.

Results The prevalence of diabetes among 18-34 years old adults in 2018 was 5.35% (95% CI: 4.62, 6.20). From 2011 to 2018, the diabetes prevalence among adults aged ≥ 35 years increased from 10.95% to 13.75% ($p < 0.001$), with the largest relative increase (90%) among obese individuals. The prevalence of diabetes remained high in 2018 for both rural (12.06%) and urban (18.95%) participants. Regression analysis identified age and BMI were the key risk factors of diabetes. Overweight and obese adults experienced significantly higher diabetes risk in the overall analysis in both survey years. Other significant risk factors of diabetes were sex, marital status, education, geographic region, wealth index, and hypertension status in both survey years.

Conclusion A high prevalence of diabetes was observed and it is increasing significantly over time. We found a substantial portion of younger adults is living with diabetes. Population-level approaches are needed to improve the identification and prevention of diabetes among adults in Bangladesh.

Keywords: Diabetes, Hypertension, trends, prevalence, risk factors, Bangladesh.

Strengths and limitations of this study

- To our knowledge this is the first study to estimate the change in prevalence and risk factors among adult population in Bangladesh using nationally representative sample including the most recent one which is released in December 2020.
- Another unique aspect of this study is the investigation of correlates of diabetes among younger adults (ages 18-34 years)
- Data of these surveys relates to anthropometric and diabetes were not self-reported rather collected by the trained and experienced health workers such as nurses, midwives, health assistance using the WHO-recommended guidelines.
- The unavailability of data of some important correlates like the types of diet, intake of fast food, intake of calories, physical exercise including the nature of work, family history of diabetes, and cholesterol level of diabetes was the major limitation of these data sets.

Introduction

Diabetes mellitus or type 2 diabetes is one of the most common chronic and preventable diseases affecting 463 million individuals worldwide in 2019.(1, 2) Furthermore, according to International Diabetes Federation (IDF), about 700 million people will have diabetes by 2045, which is a 51% increase compared to 2019.(2) This preventable disease is associated with significantly higher morbidity, mortality, and a poor quality of life. It is also associated with numerous health complications such as heart disease, stroke, renal failure, and blindness.(3-5) Moreover, diabetes causes huge financial burden on the patient, healthcare system, as well as on the country which is expected to continue to grow. Global health expenditure on diabetes is estimated to reach \$825 billion by 2030 and \$845 billion by 2045 compared to \$760 billion in 2019.(2, 6)

With more than two-thirds of people with diabetes, low- and middle-income countries have experienced a faster growth in diabetes prevalence than high-income countries.(1, 7) More specifically, the prevalence of diabetes has increased more rapidly in South East Asia than in any other regions in the world.(8) Available literature on diabetes suggests that approximately 90% to 95% of all diagnosed diabetes cases of this region are type 2 diabetes.(9, 10) Like many other countries, Bangladesh is also undergoing an epidemiological transformation, a shift from communicable to non-communicable diseases. The transition is happening due to improved socio-economic status and unplanned but rapid urban growth. From 1990 to 2014, the number of urban population increased about 2.5 times and projected to reach 112.4 million by 2050.(11) While healthy life style is less common among urban dwellers. Almost 33% of them lives in densely populated areas with poor living-environment and socio-economic conditions. They mostly have inadequate access to the basic requirements for urban life. They are subsequently

1
2
3 placing themselves at risk of contracting and suffering from both communicable and non-
4
5 communicable diseases.(12)

6
7 Bangladesh is also experiencing a nutritional transition from a conventional food habit to
8
9 a fast-food dietary practice and sedentary lifestyle which is also responsible for the emergence of
10
11 non-communicable diseases including diabetes.(13) These reasons will lead Bangladesh to
12
13 endure increased diabetes prevalence in the future. A systematic review of published studies
14
15 between 1994 and 2013 reported that the prevalence of diabetes in Bangladesh ranged between
16
17 4.5% and 35.0%.(14) The number of diabetic patients in Bangladesh is estimated to be 13.7
18
19 million by 2045.(8)

20
21
22
23
24
25
26 Several studies on diabetes conducted in Bangladesh confirmed that diabetes prevalence
27
28 among adults is rising steadily.(15, 16) However, most of those studies were confined to urban-
29
30 rural communities or some other specific groups (e.g., slum residents), which did not consider a
31
32 wide range of correlates of diabetes for the entire country. While an upward trend in the
33
34 prevalence of diabetes is evident, a very few population-based studies also reported the
35
36 prevalence of diabetes, which are outdated. Despite the rising literature of diabetes research in
37
38 Bangladesh, no study has identified trends in the prevalence of diabetes and its related risk
39
40 factors or made a comparison of its risk factors over the years.
41
42
43
44
45

46
47 In this study, we explored whether the overall prevalence of type 2 diabetes among adults
48
49 in Bangladesh changed between 2011 and 2018 and to what extent it changed by individuals'
50
51 socioeconomic and demographic characteristics. We also examined the factors that potentially
52
53 contribute to the risk of diabetes among the studied population and make comparisons among
54
55
56
57

1
2 them. It is important to recognize changes in diabetes prevalence by population subgroups to
3 ensure access to and use of available treatment required for the population living with diabetes.
4
5
6

7 8 **Methods**

9 10 *Data source*

11
12 We used two waves of BDHS's cross-sectional data, 2011 and 2017-18 (written as 2018
13 onward) to estimate the prevalence of diagnosed diabetes among the noninstitutionalized
14 Bangladeshi population aged 18 years and older. Diabetes testing and related questionnaires
15 were included only in the 2011 and 2018 surveys. The BDHS was designed to collect data to
16 monitor and evaluate population health and nutritional status of the country using two-stage
17 stratified cluster sampling from non-institutionalized households. The details of the sampling
18 procedure and sample selection are published elsewhere.(17, 18)
19
20
21
22
23
24
25
26
27
28
29
30

31 The National Institute of Population Research and Training (NIPORT) Ethics Review
32 Board approved the data collection of the Bangladesh Demographic and Health Survey (BDHS)
33 with the requirement of documented consent from all study participants. Our study was exempt
34 from the ethical review approval because we used publicly available de-identified data.
35
36
37
38
39
40
41

42 *Outcome variable*

43
44 The outcome variable for this study was the prevalence of diabetes for both survey years.
45 Diabetes Mellitus (DM)/ type-2 status was measured by fasting blood glucose (FBG) values
46 greater than or equal to 7.0 mmol/L or self-reported diabetes medication use during the
47 interview.
48
49
50
51
52
53
54
55
56
57
58
59
60

Demographic and other covariates

Demographic, household, and community-level characteristics were included to assess the prevalence and risk factors of diabetes by survey years. Individual-level characteristics were participant's (grouped into 18-24, 25-29, 30-34, 35-44, 45-54, 55-64, 65-74, and 75+ years of age), sex, marital status (currently married, not currently married), educational level (no education, primary, secondary, higher), body mass index (BMI), and hypertension status. The BMI was calculated as weight in kilograms divided by height in meters squared. We used BMI classifications for Asian population: normal weight (18.5 to 23.0), moderate risk/ overweight (23.0 to < 27.5), high risk/obese (≥ 27.5).⁽¹⁹⁾ Household and community characteristics were socio-economic status (wealth index), place of residence (urban, rural), and geographic region (division).

Statistical analysis

The full sample of each survey was used for descriptive analysis of individuals demographic and socioeconomic characteristics. Chi-square tests were performed to check the bivariate association between each characteristic and diabetes status. We also used independent proportion tests to know whether the prevalence of diabetes between the two surveys was changed significantly and calculated the relative changes between the survey periods. For the adjusted analysis, we performed multiple logistic regression models to identify the associated risk factors of diabetes in Bangladesh.

To make an appropriate comparison between the two homogeneous groups for study periods, adjusted odds ratios (AORs) were calculated for both 2011 and 2018 BDHS data of the study participants aged 35 years and older. Then we separately analyzed 18-34-year-old adults for the 2018 BDHS survey, to estimate the factors associated with diabetes among younger

1
2 adults. Moreover, we performed sensitivity analysis by splitting the datasets into rural and urban
3
4 as well as males and females. We used $P < 0.05$ at 2-sided statistical significance for all analyses.
5
6
7 Data management and statistical analyses were performed using Stata 15 (StataCorp, College
8
9 Station, TX, USA). We considered the sample weights, primary sampling units, and Strata using
10
11 the “SVY” command of Stata considering the complex nature of survey design. Comparisons by
12
13 different groups were drawn using the “svysubpop” command.
14
15

16 17 **Results**

18 19 *Sample characteristics*

20
21
22 Biomarker measurements including blood pressure and blood glucose were collected only
23
24 in 2011 and 2018 BDHS. The 2018 BDHS survey included adults ≥ 18 years representing one in
25
26 four of the households selected for the survey for biomarker data whereas the 2011 BDHS
27
28 collected biomarker measurements for adults aged ≥ 35 years only. A total of 23,541 adults were
29
30 eligible for blood glucose measurements in both surveys. After exclusion of nonresponses and
31
32 individuals with missing data, and pregnant women, 19,584 adults comprised the study
33
34 population for both survey years. Of the total included study participants 14,376 (7,556 in 2011
35
36 and 6,820 in 2018) were age 35 years or over, and the remaining 5,208 were 18-34 years
37
38 representing 2018 BDHS survey. In our study, male participants were approximately 49%, and
39
40 84% of the participants were married.
41
42
43
44
45
46

47 48 *Diabetes prevalence*

49
50
51 **Table 1** shows the socio-demographic characteristics of participants with age ≥ 35 years
52
53 in Bangladesh and their diabetes prevalence in 2011 and 2018 with relative ratios. The overall
54
55 prevalence of diabetes among adults ages ≥ 35 years increased from 10.95% in 2011 to 13.75% in
56
57

2018. The relative highest increase (38%) in diabetes prevalence was found among individuals with age 65-74 and the second-highest increase rate (36%) was found among the age group 45-54. The prevalence rate of diabetes among females increased significantly from 11.25% in 2011 to 13.81% in 2018 whereas this increment among males is not significant. The relative increase in diabetes prevalence overtime among married, currently not working individuals and rural areas were 30%, 42%, and 28%, respectively. A significant increase in diabetes prevalence was also observed among adults with no education and secondary education. The highest relative increase (54%) in diabetes prevalence was in the Dhaka region followed by 53% in the Khulna region. Diabetes prevalence among middle, richer, and the richest individuals increased significantly by 48%, 41%, and 33%, respectively. The prevalence of diabetes increased by 90% among obese individuals, this rate decreased by 9% among overweight adults. Hyperglycemia was positively and linearly associated with BMI in 2018 whereas this trend was not exactly linear in 2011 [Figure 1]. The prevalence of diabetes among males and females increased as the BMI of those individuals increased in both periods 2011 and 2018 [Figure 2].

Risk factor analysis

The adjusted results from multivariable logistic regression analysis are reported in **Table 2**. The likelihood of diabetes was the highest [AOR: 2.11, 95% CI: 1.58, 2.83] among adults ages 55-64 in 2011 whereas this rate was highest [AOR: 1.67, 95% CI: 1.21, 2.30] in the age group 65-74 in 2018 compared to adults ages 35-44. There was no significant difference in the odds of having diabetes among males and females in both periods. Although marital status was highly insignificant (p-value=0.572) in 2011, this variable was found as marginally insignificant (p-value=0.057) in 2018.

1
2
3 The findings of the study also suggest that adults with primary, secondary, and higher
4 education had 31%, 32%, and 87% higher odds of having diabetes, respectively than adults with
5 no education in 2011. However, education was not a significant factor for diabetes among adults
6 in 2018. Place of residence had no significant effect on diabetes in both periods. Compared to the
7 Dhaka division, individuals living in Barisal and Chittagong divisions had a 43% and 44%
8 higher likelihood of having diabetes, respectively in 2011. On the other hand, there exists no
9 significant difference in having diabetes among adults in Barisal, Chittagong, Dhaka, Rajshahi,
10 and Sylhet divisions in 2018. Regarding economic status, only the richest individuals had a
11 significantly higher likelihood (96%) of diabetes in 2011 compared to the poorest individuals.
12 However, both the richer and richest adults had more likelihood of diabetes [AOR: 1.84 and
13 3.09] than the poorest adults in 2018.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 Higher BMI was a significant factor in both 2011 and 2018. For example, overweight and
31 obese adults compared to normal-weight adults had 54% and 51% more likelihood of diabetes,
32 respectively in 2011, and 22% and 44% higher likelihood of diabetes in 2018. Moreover, the
33 odds of having diabetes among working adults in 2018 was lower [AOR: 0.80, 95% CI: (0.65,
34 0.99)] than non-working adults. Since there exists a strong relationship between diabetes and
35 hypertension, individuals having hypertension had 51% and 57% more likelihoods of diabetes,
36 respectively in 2011 and 2018 compared to individuals without hypertension.
37
38
39
40
41
42
43
44
45
46
47
48

49 The adjusted odds ratios of the factors with their corresponding 95% confidence intervals
50 and p-value among adults' ages 18-34 years in 2018 are presented in **Table 3**. Individuals from
51 age group 30-34 had 97% [AOR: 1.97, 95% CI: 1.31, 2.97] higher odds of having diabetes than
52 individuals from age group 18-24. Adults living in Chittagong, Khulna, and Rangpur had 49%,
53
54
55
56
57

1
2 46%, and 62% fewer odds of diabetes, respectively compared to adults living in Dhaka.
3
4 Moreover, obese individuals had a higher likelihood [AOR: 1.73, 95% CI: 1.12, 2.67] of diabetes
5
6 than individuals with normal weight. Similar to the older adults (≥ 35 years), individuals having
7
8 who were working at the time of survey had 35% lower odds of diabetes than individuals having
9
10 no job, and hypertension was also found significant ($p=0.002$) factor for diabetes.
11
12
13
14
15

16 *Subgroup analysis*

17
18 Subgroup analysis of diabetes by sex and place of residence were also performed and the
19
20 results of this analysis are presented in **Table S1** and **Table S2**. Male with higher education had
21
22 99% higher risk of diabetes compared to male with no education in 2011 whereas no significant
23
24 difference in the risk of diabetes among them was observed in 2018. However, the odds of
25
26 having diabetes between a male with primary education and no education was not significantly
27
28 different [$p=0.236$] in 2011. The opposite scenario [AOR: 1.48 [(1.06, 2.06), $p=0.022$] was
29
30 observed in 2018. Urban males had less likelihood [AOR: 0.71, 95% CI: 0.55, 0.93] of diabetes
31
32 than rural males in 2018 whereas there was no difference in the likelihood of diabetes among
33
34 them in 2011.
35
36
37
38
39
40

41
42 Marital status was found to be significant factor ($p=0.043$) of diabetes among rural adults
43
44 in 2018 only. Richer individuals had no significant difference ($p>0.05$) in having diabetes
45
46 compared to the poorest individuals except in rural areas in 2018 ($p=0.019$). Currently working
47
48 individuals had 31% (AOR: 0.69, 95% CI: 0.49, 0.99) and 27% (AOR: 0.73, 95% CI: 0.57, 0.95)
49
50 lower odds of having diabetes than non-working rural individuals, in 2011 and 2018,
51
52 respectively. However, there was no significant difference in diabetes among working and non-
53
54 working urban individuals for both periods.
55
56
57
58
59
60

Discussion

For the first time, we systematically analyses the prevalence and risk factors of diabetes among the adult population (aged 18 years) in Bangladesh using two waves of nationally representative survey data. We found some remarkable findings linked to diabetes and its risk factors. One of the important findings was the identification of upward trends in the overall prevalence of diabetes and its distribution as per individual characteristics. In 2018 the prevalence of diabetes among 18-34 years adults was 5.35%. The prevalence among adults ≥ 35 years has significantly increased from 10.95% in 2011 to 13.75% in 2018; with a relative increase of 26%. During this period, the prevalence of diabetes increased significantly not only among the overall population but also among each age groups, both sexes, married individuals, uneducated and secondary completed, rural areas, middle to richest wealth index group, obese individuals, unemployed, and hypertensive patients. The prevalence of diabetes had increased rapidly among obese, middle to rich income group, and in Dhaka and Khulna regions. During this period, the proportion of people with hyperglycemia also increased significantly among male, obese, and 45-74 age groups. This finding is consistent with the reported prevalence of diabetes among the adult populations over the years. (20-22)

We found that the likelihood of diabetes increased with an increase of age. The odds of having diabetes was almost two times for adults aged 30-34 years compared to the 18-24 years adults. Similar to the previous studies in Bangladesh(23, 24) and other developing countries, (25, 26) the odds of having diabetes increased consistently for all age groups of older adults (≥ 35 years) in both data sets. In the future, the upward trend of having diabetes is likely to be a major public health concern in Bangladesh as its population age-structure changes with lower fertility rate, steady socioeconomic growth, and increased life expectancy. This process will sharply lead

1
2
3 to an increase in the number of middle and older age population and diabetes prevalence in
4 Bangladesh. (27) The prevalence of diabetes among the working-age population may be a
5 concern because of the complex effects of diabetes on co-morbidity and economic growth in
6 Bangladesh, where about 12% of total households pay for diabetic care by selling household
7 assets or borrowing money. (28-30)
8
9
10
11
12
13
14
15

16 We observed a substantially higher relative increase in the prevalence of diabetes in rural
17 areas (28%) compared to urban areas (18%), which indicates that diabetes is no longer confined
18 to urban areas in Bangladesh and is also a matter of concern in rural areas of Bangladesh.
19
20
21 Tripathy et al.(31) found a similar trend in our neighboring country India. Our study also
22 identified the greater prevalence of having diabetes among married people compared to
23 unmarried people. This finding was significant in 2018, but was not found to be significant in
24 2011.(27) This result was broadly consistent with previous studies in which the presence of
25 diabetes was also associated with greater marital stability and satisfaction (23, 31, 32).
26
27 Moreover, male being married was also associated with a higher risk of hypertension and type-2
28 diabetes (33).
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Although higher education and socio-economic status are negatively associated with diabetes in
43 developed countries, we have found the opposite results in Bangladesh for both survey periods. Richest
44 individuals aged ≥ 35 years are 3.09 times to have diabetes, followed by richer and middle-
45 income groups compared to the poorest wealth group. These findings are in line with the
46 previous studies conducted in Asian and other developing countries (23, 34, 35). The prevalence
47 of diabetes varies also by region among individuals ≥ 35 years in Bangladesh. For example,
48 people living in Rangpur and Khulna regions have experienced a significantly lower risk of
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 having diabetes than those from Dhaka and other regions. The relatively lower socioeconomic
4
5 status of people in these two divisions compared to Dhaka is plausibly linked with their lower
6
7 odds of living with diabetes (23, 35). The rapid increase in the prevalence of diabetes in all
8
9 regions, particularly in Dhaka and Khulna regions, in which it has increased 54% and 53%
10
11 respectively between 2011 and 2018, is plausibly associated with the rapid growth of
12
13 urbanization and its consequences on healthy lifestyles (36).
14
15

16
17
18
19 High BMI and abdominal obesity are independent risk factors of diabetes reported in
20
21 most of the previous studies. (23, 28, 30) Our study findings also pointed out that adults ages ≥ 35
22
23 years with higher BMI have a greater likelihood of having diabetes compared to normal weight
24
25 adults in both 2011 and 2018. For instance, the odds of having diabetes among overweight and
26
27 obese individuals were 1.22 and 1.44 times compared to the normal weight individuals in 2018.
28
29
30

31
32
33 Another unique aspect of this study is the investigation of correlates of diabetes among
34
35 younger adults (ages 18-34 years). This study reports the prevalence of diabetes among younger
36
37 adults for the first time in Bangladesh. We observed 5.35% younger adults had diabetes in
38
39 Bangladesh in 2018. Although the prevalence of diabetes among < 35 years adults is low; in
40
41 Bangladesh, the population age structure is predominantly young, this is an alarming rate with
42
43 the total number of young adults with diabetes. While the annual cost of caring for older people
44
45 with diabetes may be higher than caring for younger people with diabetes, since they have longer
46
47 to live with the disease, the lifetime health burden and cost may also be greater for young people.
48
49 Moreover, due to change in diet habits younger population are at greater risk of obesity than in
50
51 older adults resulting them into higher risk group of diabetes. Similar to the older adults, the
52
53
54
55
56
57
58
59
60

1
2
3 increasing age, higher BMI and, having hypertension were found as significant risk factors of
4
5 diabetes among younger adults.

6
7 One of the major strengths of our study is the use of nationally representative cross-
8
9 sectional survey data of the last two waves, including the most recent one which is released in
10
11 December 2020. Data of these surveys relates to anthropometric and diabetes were not self-
12
13 reported rather collected by the trained and experienced health workers such as nurses,
14
15 midwives, health assistance using the WHO-recommended guidelines. Compared to other
16
17 context-specific cross-sectional survey data, the measurement error and bias are likely to be less
18
19 for this data sets due to the use of globally standard and valid measures in Bangladesh. To our
20
21 knowledge, this study for the first time estimated the national diabetes prevalence and its risk
22
23 factors among general adults of Bangladesh. Another important strength is that we compared
24
25 changes in estimates of diabetes predictors between 2011 and 2018 surveys along with
26
27 subgroups analyses: such as 18-34 years, by sex, and by place of residence. Despite having some
28
29 strength, our study is not beyond the limitations. The unavailability of data of some important
30
31 correlates like the types of diet, intake of fast food, intake of calories, physical exercise including
32
33 the nature of work, family history of diabetes, and cholesterol level of diabetes was the major
34
35 limitation of these data sets. Moreover, as a cross-sectional survey, blood sugar level was
36
37 measured for one day only and thus we do not have follow-up and or longitudinal data on
38
39 diabetes and its correlates.
40
41
42
43
44
45
46
47
48

49 The study shows that among Bangladeshi adults, there is a high prevalence of diabetes
50
51 and it is escalating over time. The study also reports a significant portion of younger adults with
52
53 diabetes. Age and overweight/obesity are the two most important risk factors for diabetes for all
54
55 adults, irrespective of sex, residence, educational attainment and wealth index. There is evidence
56
57
58
59

1
2 of an increase in the magnitude of diabetes over time and in the younger population; indeed, as
3 age increases the chances of developing Diabetes mellitus significantly. These findings, together
4 with an increase in the prevalence of type 2 diabetes among Bangladeshi general adults,
5 underscore the need for primary and secondary prevention efforts tailored to age-specific
6 populations.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

List of Abbreviations

| | |
|------|--|
| AOR | Adjusted Odds Ratio |
| BMI | Body mass index |
| CI | Confidence interval |
| CVD | Cardiovascular diseases |
| WHO | World Health Organization |
| BDHS | Bangladesh Demographic and Health Survey |

Availability of data and material: All data presented here in the manuscript is freely available at dhsprogram.com.

Prior Publication: This data has not been published previously and is not under consideration elsewhere.

Author Contribution: M.A.B. Chowdhury conceptualized the study, designed the analytic approach, managed and performed the analysis, interpreted the results, and drafted the manuscript. M. Islam: helped with the analysis, interpreted the results, and drafted the manuscript. J. Rahman, M.J. Uddin reviewed, edited and updated the manuscript and M.R.Haque reviewed, edited, and supervised the study.

Disclosures/Conflict of Interest: There are no potential conflicts (financial, professional, or personal) to disclose by any of the authors.

Funding: None

Acknowledgement: We thank DHS program for providing the data.

Ethical approval and consent

All BDHS surveys received ethical approval from ICF Macro Institutional Review Board, Maryland, USA and National Research Ethics Committee of Bangladesh Medical Research Council (BMRC), Dhaka, Bangladesh. Informed consent was obtained from each participant of the survey before enrolling in the survey by using the Introduction and Consent form of the survey. It was also explained that the information will be kept strictly confidential and will not be shared with anyone except members of the survey team.

Patient and Public Involvement

The BDHS questionnaires were based on the MEASURE DHS model questionnaires. These model questionnaires were adapted for use in Bangladesh during a series of meetings with a technical working group (TWG) that consisted of representatives from NIPORT, Mitra and Associates, International Centre for Diarrheal Diseases and Control, Bangladesh (icDDR, b), USAID/Bangladesh, and MEASURE DHS. Patients were not directly involved in the study however; the TWG involved representatives from the government, non-government, ministry of health and family welfare representatives and donor organizations were involved the study design and questionnaire development. The results will be used by the health researchers, policy makers of the country.

Description of tables and figures

FIGURE 1: Trends in diabetes class by age group and sex

FIGURE 2: Trends in diabetes class by BMI groups and sex

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

TABLE 2. Adjusted odds ratios for factors predicting diabetes among adults 35 years and older BDHS 2011-2018

TABLE 3. Adjusted odds Ratios for factors predicting diabetes among adults 18-34 years BDHS 2017-18.

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|-----------------------------------|---------------------------------|----------------------------------|----------------------------------|--------------------------------|-------|
| All adults age 35 years and older | 100% | 10.95 (0.0048) | 13.75 (0.0056) | <0.001 | 1.26 |
| Age group | | | | | |
| 35-44 | 35.99 | 8.82 (0.0065) | 11.21 (0.0078) | 0.081 | 1.27 |
| 45-54 | 27.43 | 10.86 (0.0079) | 14.82 (0.01) | 0.008 | 1.36 |
| 55-64 | 18.65 | 14.69 (0.012) | 15.78 (0.0115) | 0.6286 | 1.07 |
| 65-74 | 11.26 | 11.60 (0.0128) | 15.97 (0.016) | 0.1123 | 1.38 |
| 75+ | 6.68 | 12.11 (0.0174) | 13.43 (0.0189) | 0.793 | 1.11 |
| Sex | | | | | |
| Male | 48.98 | 10.65 (0.0061) | 13.69 (0.0072) | 0.091 | 1.29 |
| Female | 51.02 | 11.25 (0.006) | 13.81 (0.0072) | 0.003 | 1.23 |
| Marital status | | | | | |
| Not married | 15.88 | 12.13 (0.0112) | 12.54 (0.0111) | 0.953 | 1.03 |
| Married | 84.12 | 10.73 (0.0049) | 13.98 (0.006) | <0.001 | 1.3 |
| Educational level | | | | | |
| No education | 44.8 | 8.39 (0.0058) | 10.64 (0.0072) | 0.021 | 1.27 |
| Primary | 29.07 | 11.10 (0.0078) | 13.59 (0.009) | 0.141 | 1.22 |
| Secondary | 18.15 | 13.03 (0.0108) | 17.8 (0.0121) | 0.015 | 1.37 |
| Higher | 7.98 | 21.79 (0.018) | 20.27 (0.0185) | 0.209 | 0.93 |
| Place of residence | | | | | |
| Urban | 23.91 | 16.08 (0.0106) | 18.95 (0.0106) | 0.129 | 1.18 |
| Rural | 76.09 | 9.39 (0.0051) | 12.06 (0.0065) | 0.003 | 1.28 |
| Geographic region | | | | | |
| Barisal | 5.79 | 12.54 (0.0117) | 12.09 (0.0161) | 0.975 | 0.96 |
| Chittagong | 16.45 | 14.28 (0.0125) | 17.33 (0.0169) | 0.087 | 1.21 |
| Dhaka | 38.99 | 11.26 (0.0107) | 17.32 (0.0118) | 0.002 | 1.54 |
| Khulna | 13.45 | 7.30 (0.007) | 11.14 (0.0114) | 0.002 | 1.53 |
| Rajshahi | 13.15 | 10.73 (0.01) | 11.32 (0.0134) | 0.664 | 1.05 |
| Rangpur | 9.26 | 8.59 (0.0111) | 7.93 (0.0097) | 0.667 | 0.92 |
| Sylhet | 2.91 | 11.85 (0.011) | 12.71 (0.0169) | 0.998 | 1.07 |
| Wealth index | | | | | |
| Poorest | 19.91 | 7.28 (0.0088) | 7.26 (0.0091) | 0.69 | 1.00 |
| Poorer | 19.74 | 7.35 (0.0084) | 7.70 (0.0085) | 0.891 | 1.05 |
| Middle | 20.19 | 7.56 (0.0075) | 11.17 (0.01) | 0.005 | 1.48 |
| Richer | 19.70 | 11.33 (0.0098) | 16.01 (0.0128) | <0.001 | 1.41 |
| Richest | 20.46 | 20.49 (0.0122) | 27.18 (0.0134) | 0.006 | 1.33 |
| Body Mass Index | | | | | |
| Underweight | 19.87 | 7.27 (0.0081) | 7.79 (0.0087) | 0.437 | 1.07 |
| Normal weight | 36.10 | 9.59 (0.0071) | 11.21 (0.0074) | 0.343 | 1.17 |
| Overweight | 20.14 | 18.57 (0.0149) | 16.95 (0.0095) | 0.045 | 0.91 |
| Obese | 23.88 | 11.90 (0.0078) | 22.56 (0.0156) | <0.001 | 1.9 |
| Currently working | | | | | |

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|--------------|---------------------------------|----------------------------------|----------------------------------|--------------------------------|-------|
| No | 43.63 | 12.11 (0.0064) | 17.16 (0.0089) | <0.001 | 1.42 |
| Yes | 56.37 | 9.70 (0.0061) | 12.00 (0.006) | 0.10 | 1.24 |
| Hypertension | | | | | |
| No | 67.45 | 9.06 (0.0048) | 10.55 (0.0063) | 0.522 | 1.16 |
| Yes | 32.55 | 16.33 (0.0105) | 18.57 (0.009) | 0.178 | 1.14 |

For peer review only

TABLE 2. Adjusted odds ratios for factors predicting diabetes among adults 35 years and older BDHS 2011-2018.

| Variables | 2011 BDHS | | 2017-18 BDHS | |
|---------------------------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | |
| 35-44 | Ref | | Ref | |
| 45-54 | 1.36 (1.07, 1.72) | 0.012 | 1.41 (1.13, 1.76) | 0.002 |
| 55-64 | 2.11 (1.58, 2.83) | <0.001 | 1.58 (1.24, 2.03) | <0.001 |
| 65-74 | 1.60 (1.13, 2.27) | 0.008 | 1.67 (1.21, 2.30) | 0.002 |
| 75+ | 1.77 (1.14, 2.74) | 0.011 | 1.32 (0.86, 2.01) | 0.202 |
| Sex | | | | |
| Male | Ref | | Ref | |
| Female | 0.78 (0.58, 1.05) | 0.104 | 0.92 (0.75, 1.13) | 0.435 |
| Marital status | | | | |
| Not married | Ref | | | |
| Married | 0.93 (0.72, 1.20) | 0.572 | 1.28 (0.99, 1.64) | 0.057 |
| Educational level | | | | |
| No education | Ref | | Ref | |
| Primary | 1.31 (1.05, 1.64) | 0.017 | 1.23 (0.99, 1.53) | 0.059 |
| Secondary | 1.32 (1.01, 1.73) | 0.045 | 1.23 (0.96, 1.57) | 0.108 |
| Higher | 1.87 (1.35, 2.60) | <0.001 | 1.15 (0.83, 1.60) | 0.397 |
| Place of residence | | | | |
| Urban | 1.08 (0.87, 1.33) | 0.7 | 0.96 (0.79, 1.16) | 0.678 |
| Rural | Ref | | Ref | |
| Geographic region | | | | |
| Barisal | 1.43 (1.04, 1.96) | 0.027 | 0.75 (0.54, 1.06) | 0.103 |
| Chittagong | 1.44 (1.09, 1.89) | 0.010 | 0.88 (0.67, 1.16) | 0.369 |
| Dhaka | Ref | | Ref | |
| Khulna | 0.66 (0.50, 0.89) | 0.007 | 0.60 (0.46, 0.79) | <0.001 |
| Rajshahi | 1.17 (0.87, 1.57) | 0.309 | 0.74 (0.55, 1.00) | 0.054 |
| Rangpur | 1.00 (0.69, 1.45) | 0.986 | 0.56 (0.41, 0.76) | <0.001 |
| Sylhet | 1.22 (0.91, 1.63) | 0.183 | 0.75 (0.53, 1.06) | 0.101 |
| Wealth index | | | | |
| Poorest | Ref | | Ref | |
| Poorer | 0.89 (0.62, 1.27) | 0.514 | 0.99 (0.70, 1.40) | 0.957 |
| Middle | 0.86 (0.62, 1.19) | 0.365 | 1.33 (0.94, 1.88) | 0.103 |
| Richer | 1.18 (0.84, 1.65) | 0.345 | 1.84 (1.29, 2.61) | 0.001 |
| Richest | 1.96 (1.40, 2.76) | <0.001 | 3.09 (2.18, 4.38) | <0.001 |
| Body Mass Index | | | | |
| Underweight | 0.82 (0.61, 1.10) | 0.177 | 0.77 (0.58, 1.01) | 0.055 |
| Normal weight | Ref | | Ref | |
| Overweight | 1.54 (1.20, 1.97) | 0.001 | 1.22 (1.00, 1.50) | 0.052 |
| Obese | 1.51 (1.16, 1.97) | 0.003 | 1.44 (1.13, 1.84) | 0.003 |
| Currently working | | | | |
| No | Ref | | Ref | |
| Yes | 0.77 (0.59, 1.01) | 0.056 | 0.80 (0.65, 0.99) | 0.039 |
| Hypertension | | | | |
| No | Ref | | Ref | |
| Yes | 1.51 (1.26, 1.81) | <0.001 | 1.57 (1.32, 1.87) | <0.001 |

TABLE 3. Adjusted odds Ratios for factors predicting diabetes among adults 18-34 years
BDHS 2017-18.

| Variables | OR (95% CI) | p-value |
|--------------------|-------------------|---------|
| Age groups | | |
| 18-24 | Ref | |
| 25-19 | 1.36 (0.93, 2.00) | 0.113 |
| 30-34 | 1.97 (1.31, 2.97) | 0.001 |
| Sex | | |
| Male | Ref | |
| Female | 0.88 (0.64, 1.22) | 0.445 |
| Marital status | | |
| Not married | | |
| Married | 0.88 (0.61, 1.26) | 0.473 |
| Educational level | | |
| No education | Ref | |
| Primary | 1.48 (0.81, 2.69) | 0.198 |
| Secondary | 1.27 (0.67, 2.41) | 0.464 |
| Higher | 1.17 (0.59, 2.34) | 0.654 |
| Place of residence | | |
| Urban | 1.26 (0.9, 1.75) | 0.179 |
| Rural | | |
| Geographic region | | |
| Barisal | 0.95 (0.58, 1.55) | 0.840 |
| Chittagong | 0.51 (0.32, 0.81) | 0.004 |
| Dhaka | Ref | |
| Khulna | 0.54 (0.35, 0.85) | 0.007 |
| Rajshahi | 0.60 (0.35, 1.01) | 0.053 |
| Rangpur | 0.38 (0.21, 0.68) | 0.001 |
| Sylhet | 0.97 (0.57, 1.64) | 0.896 |
| Wealth index | | |
| Poorest | Ref | |
| Poorer | 1.02 (0.60, 1.72) | 0.933 |
| Middle | 1.04 (0.63, 1.70) | 0.902 |
| Richer | 1.53 (0.92, 2.53) | 0.122 |
| Richest | 1.58 (0.94, 2.69) | 0.088 |
| Body Mass Index | | |
| Underweight | 1.07 (0.67, 1.70) | 0.778 |
| Normal weight | Ref | |
| Overweight | 1.31 (0.93, 1.84) | 0.128 |
| Obese | 1.73 (1.12, 2.67) | 0.013 |
| Currently working | | |
| No | Ref | |
| Yes | 0.65 (0.48, 0.89) | 0.007 |
| Hypertension | | |
| No | Ref | |
| Yes | 1.73 (1.23, 2.42) | 0.002 |

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|---------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.56 (1.1, 2.21) | 0.013 | 1.41 (0.98, 2.03) | 0.065 | 1.18 (0.87, 1.62) | 0.288 | 1.41 (1.06, 1.86) | 0.018 |
| 55-64 | 2.36 (1.6, 3.47) | <0.001 | 1.58 (1.09, 2.29) | 0.015 | 1.84 (1.15, 2.94) | 0.012 | 1.53 (1.09, 2.14) | 0.013 |
| 65-74 | 1.79 (1.14, 2.8) | 0.011 | 1.80 (1.18, 2.75) | 0.007 | 1.35 (0.76, 2.39) | 0.304 | 1.44 (0.9, 2.33) | 0.132 |
| 75+ | 1.52 (0.85, 2.72) | 0.153 | 1.41 (0.79, 2.51) | 0.24 | 1.92 (0.98, 3.76) | 0.059 | 1.11 (0.62, 1.99) | 0.727 |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.66 (0.35, 1.26) | 0.205 | 1.83 (0.96, 3.47) | 0.064 | 1.00 (0.75, 1.34) | 0.992 | 1.17 (0.89, 1.56) | 0.263 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.22 (0.88, 1.68) | 0.236 | 1.48 (1.06, 2.06) | 0.022 | 1.37 (1, 1.86) | 0.047 | 1.12 (0.84, 1.48) | 0.434 |
| Secondary | 1.14 (0.80, 1.63) | 0.47 | 1.15 (0.78, 1.69) | 0.482 | 1.53 (1.05, 2.23) | 0.029 | 1.37 (0.97, 1.92) | 0.071 |
| Higher | 1.99 (1.31, 3.01) | 0.001 | 1.19 (0.76, 1.86) | 0.449 | 1.85 (1.1, 3.11) | 0.02 | 1.98 (0.59, 1.64) | 0.941 |
| Place of residence | | | | | | | | |
| Urban | 1.03 (0.76, 1.4) | 0.838 | 0.71 (0.55, 0.93) | 0.012 | 1.1 (0.83, 1.45) | 0.497 | 1.24 (0.96, 1.61) | 0.097 |
| Rural | Ref | | Ref | | Ref | | Ref | |
| Geographic region | | | | | | | | |
| Barisal | 1.42 (0.91, 2.22) | 0.121 | 0.83 (0.53, 1.30) | 0.415 | 1.44 (0.97, 2.14) | 0.072 | 0.72 (0.46, 1.12) | 0.148 |
| Chittagong | 1.55 (1.05, 2.27) | 0.027 | 0.88 (0.62, 1.26) | 0.484 | 1.36 (0.97, 1.91) | 0.071 | 0.88 (0.61, 1.26) | 0.48 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.75 (0.50, 1.12) | 0.159 | 0.56 (0.39, 0.82) | 0.003 | 0.61 (0.4, 0.92) | 0.02 | 0.65 (0.46, 0.93) | 0.019 |
| Rajshahi | 1.14 (0.73, 1.77) | 0.569 | 0.78 (0.52, 1.2) | 0.259 | 1.21 (0.82, 1.77) | 0.338 | 0.71 (0.46, 1.09) | 0.116 |
| Rangpur | 1.04 (0.64, 1.68) | 0.885 | 0.60 (0.39, 0.92) | 0.019 | 0.99 (0.63, 1.57) | 0.975 | 0.53 (0.34, 0.82) | 0.004 |
| Sylhet | 1.41 (0.92, 2.15) | 0.112 | 0.66 (0.41, 1.07) | 0.091 | 1.11 (0.77, 1.6) | 0.58 | 0.82 (0.54, 1.22) | 0.325 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |
| Poorer | 0.81 (0.51, 1.30) | 0.389 | 1.20 (0.72, 2.00) | 0.485 | 0.96 (0.58, 1.59) | 0.88 | 0.87 (0.58, 1.32) | 0.522 |
| Middle | 0.75 (0.46, 1.21) | 0.239 | 1.90 (1.16, 3.12) | 0.011 | 0.99 (0.63, 1.57) | 0.981 | 1.02 (0.66, 1.57) | 0.945 |

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|-------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Richer | 0.95 (0.59, 1.51) | 0.815 | 2.58 (1.53, 4.34) | <0.001 | 1.43 (0.9, 2.25) | 0.129 | 1.44 (0.94, 2.21) | 0.093 |
| Richest | 1.54 (0.94, 2.53) | 0.088 | 5.47 (3.25, 9.19) | <0.001 | 2.38 (1.49, 3.79) | <0.001 | 2.03 (1.33, 3.09) | 0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.86 (0.6, 1.22) | 0.382 | 0.76 (0.51, 1.13) | 0.177 | 0.69 (0.44, 1.08) | 0.101 | 0.78 (0.52, 1.18) | 0.238 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 1.54 (1.13, 2.12) | 0.007 | 1.25 (0.94, 1.66) | 0.122 | 1.63 (1.09, 2.42) | 0.016 | 1.15 (0.86, 1.52) | 0.343 |
| Obese | 1.71 (1.1, 2.68) | 0.018 | 1.82 (1.22, 2.69) | 0.003 | 1.31 (0.86, 1.98) | 0.207 | 1.29 (0.95, 1.76) | 0.105 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.71 (0.48, 1.06) | 0.09 | 0.80 (0.54, 1.17) | 0.249 | 0.84 (0.55, 1.26) | 0.394 | 0.79 (0.61, 1.03) | 0.076 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.17 (0.88, 1.56) | 0.269 | 1.42 (1.1, 1.83) | 0.007 | 1.77 (1.4, 2.24) | <0.001 | 1.70 (1.33, 2.18) | <0.001 |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.58 (1.07, 2.36) | 0.023 | 1.19 (0.85, 1.65) | 0.312 | 1.22 (1.66, 0.9) | 0.194 | 1.59 (1.18, 2.13) | 0.002 |
| 55-64 | 3.57 (2.16, 5.9) | <0.001 | 1.50 (0.98, 2.29) | 0.059 | 1.60 (2.3, 1.11) | 0.011 | 1.67 (1.24, 2.27) | 0.001 |
| 65-74 | 2.25 (1.24, 4.1) | 0.008 | 1.50 (0.84, 2.66) | 0.168 | 1.32 (2.04, 0.86) | 0.200 | 1.75 (1.19, 2.57) | 0.004 |
| 75+ | 2.54 (1.17, 5.5) | 0.018 | 1.89 (1.01, 3.51) | 0.045 | 1.45 (2.45, 0.85) | 0.171 | 1.14 (0.65, 2.01) | 0.645 |
| Sex | | | | | | | | |
| Female | 0.82 (0.49, 1.38) | 0.461 | 1.29 (0.9, 1.85) | 0.173 | 0.76 (1.11, 0.52) | 0.15 | 0.88 (0.63, 1.03) | 0.087 |
| Male | Ref | | Ref | | Ref | | Ref | |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.8 (0.52, 1.24) | 0.322 | 1.07 (0.7, 1.62) | 0.759 | 1.01 (1.38, 0.74) | 0.962 | 1.39 (1.01, 1.92) | 0.043 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.44 (0.96, 2.14) | 0.076 | 1 (0.69, 1.45) | 0.992 | 1.26 (1.64, 0.97) | 0.086 | 1.30 (1.00, 1.69) | 0.048 |
| Secondary | 1.46 (0.94, 2.27) | 0.089 | 1.35 (0.89, 2.03) | 0.155 | 1.22 (1.72, 0.86) | 0.272 | 1.15 (0.83, 1.59) | 0.394 |
| Higher | 1.9 (1.13, 3.19) | 0.015 | 1.19 (0.74, 1.93) | 0.465 | 1.85 (2.94, 1.16) | 0.009 | 1.15 (0.74, 1.81) | 0.532 |
| Geographic region | | | | | | | | |
| Barisal | 0.93 (0.58, 1.49) | 0.76 | 0.99 (0.54, 1.83) | 0.982 | 1.63 (2.45, 1.09) | 0.018 | 0.70 (0.46, 1.06) | 0.088 |
| Chittagong | 1.04 (0.69, 1.58) | 0.845 | 0.58 (0.4, 0.84) | 0.004 | 1.66 (2.41, 1.15) | 0.007 | 1.04 (0.72, 1.5) | 0.841 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.82 (0.52, 1.28) | 0.381 | 0.70 (0.5, 1.00) | 0.05 | 0.61 (0.91, 0.4) | 0.017 | 0.57 (0.4, 0.81) | 0.002 |
| Rajshahi | 0.94 (0.60, 1.48) | 0.802 | 0.74 (0.51, 1.06) | 0.095 | 1.28 (1.9, 0.86) | 0.228 | 0.75 (0.51, 1.12) | 0.165 |
| Rangpur | 0.77 (0.46, 1.29) | 0.318 | 0.5 (0.33, 0.78) | 0.002 | 1.07 (1.71, 0.67) | 0.775 | 0.58 (0.4, 0.84) | 0.005 |
| Sylhet | 1.43 (0.89, 2.31) | 0.143 | 0.71 (0.47, 1.09) | 0.115 | 1.22 (1.79, 0.83) | 0.318 | 0.77 (0.5, 1.18) | 0.226 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Poorer | 1.23 (0.49, 3.07) | 0.653 | 0.87 (0.36, 2.08) | 0.754 | 0.87 (1.27, 0.6) | 0.468 | 0.98 (0.67, 1.42) | 0.896 |
| Middle | 1.05 (0.49, 2.24) | 0.900 | 0.67 (0.3, 1.51) | 0.335 | 0.84 (1.2, 0.58) | 0.327 | 1.42 (0.97, 2.08) | 0.07 |
| Richer | 0.92 (0.49, 1.72) | 0.787 | 1.74 (0.81, 3.76) | 0.155 | 1.27 (1.85, 0.87) | 0.218 | 1.54 (1.08, 2.47) | 0.019 |
| Richest | 1.67 (0.89, 3.11) | 0.108 | 2.54 (1.21, 5.34) | 0.014 | 2.05 (3.05, 1.38) | <0.001 | 3.05 (2.03, 4.60) | <0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.96 (0.55, 1.68) | 0.884 | 1.24 (0.73, 2.1) | 0.422 | 0.77 (1.07, 0.55) | 0.119 | 0.58 (0.49, 0.94) | 0.02 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 2.27 (1.47, 3.5) | <0.001 | 0.99 (0.7, 1.41) | 0.971 | 1.32 (1.8, 0.97) | 0.077 | 1.66 (1.06, 1.73) | 0.015 |
| Obese | 2.86 (1.88, 4.34) | <0.001 | 1.5 (1.04, 2.16) | 0.029 | 1.1 (1.55, 0.78) | 0.581 | 1.66 (0.98, 1.89) | 0.068 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.91 (0.6, 1.38) | 0.664 | 1.04 (0.73, 1.48) | 0.828 | 0.69 (0.49, 0.99) | 0.044 | 0.73 (0.57, 0.95) | 0.02 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.52 (1.15, 2.02) | 0.004 | 1.58 (1.18, 2.11) | 0.002 | 1.50 (1.91, 1.18) | 0.001 | 1.66 (1.29, 1.99) | <0.001 |

References

1. Ogurtsova K, da Rocha Fernandes JD, Huang Y, Linnenkamp U, Guariguata L, Cho NH, et al. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes research and clinical practice*. 2017;128:40-50.
2. Federation ID. IDF Diabetes Atlas Ninth. Dunia: IDF. 2019.
3. Kumar SS, Houlden RL. Ethnocultural diversity and the diabetes epidemic in Canada: a call to action. *Can J Diab*. 2005;29(2):84-5.
4. Oldroyd J, Banerjee M, Heald A, Cruickshank K. Diabetes and ethnic minorities. *Postgraduate medical journal*. 2005;81(958):486-90.
5. World Health O. World health statistics 2013: a wealth of information on global public health. World Health Organization; 2013.
6. Williams R, Karuranga S, Malanda B, Saeedi P, Basit A, Besançon S, et al. Global and regional estimates and projections of diabetes-related health expenditure: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract*. 2020;162:108072.
7. Dagenais GR, Gerstein HC, Zhang X, McQueen M, Lear S, Lopez-Jaramillo P, et al. Variations in diabetes prevalence in low-, middle-, and high-income countries: results from the prospective urban and rural epidemiological study. *Diabetes care*. 2016;39(5):780-7.
8. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes research and clinical practice*. 2018;138:271-81.
9. Niti S, Amrit V, Gupta B, Jasdeep S. Prevalence and risk factors of diabetes mellitus among adults residing in field practice area of a teaching Hospital in Punjab. *Health line J*. 2015;6(1):57-62.
10. Hussain A, Vaaler S, Sayeed MA, Mahtab H, Ali SMK, Khan AKA. Type 2 diabetes and impaired fasting blood glucose in rural Bangladesh: a population-based study. *The European Journal of Public Health*. 2007;17(3):291-6.
11. Desa UN. World urbanization prospects, the 2011 revision. Population Division, Department of Economic and Social Affairs, United Nations Secretariat. 2014.
12. Goli S, Arokiasamy P, Chattopadhyay A. Living and health conditions of selected cities in India: Setting priorities for the National Urban Health Mission. *Cities*. 2011;28(5):461-9.
13. Amuna P, Zotor FB. Epidemiological and nutrition transition in developing countries: impact on human health and development. *Proc Nutr Soc*. 2008;67(1):82-90.
14. Biswas T, Islam A, Rawal LB, Islam SM. Increasing prevalence of diabetes in Bangladesh: a scoping review. *Public Health*. 2016;138:4-11.
15. Rahim MA, Hussain A, Khan AKA, Sayeed MA, Ali SMK, Vaaler S. Rising prevalence of type 2 diabetes in rural Bangladesh: a population based study. *Diabetes research and clinical practice*. 2007;77(2):300-5.
16. Bhowmik B, Afsana F, My Diep L, Binte Munir S, Wright E, Mahmood S, et al. Increasing prevalence of type 2 diabetes in a rural Bangladeshi population: a population based study for 10 years. *Diabetes & metabolism journal*. 2013;37(1):46-53.
17. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2011. 2011.
18. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2014. 2014.
19. Who EC. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet (London, England)*. 2004;363(9403):157.
20. Menke A, Casagrande S, Geiss L, Cowie CC. Prevalence of and Trends in Diabetes Among Adults in the United States, 1988-2012. *JAMA*. 2015;314(10):1021-9.

21. Bullard KM, Saydah SH, Imperatore G, Cowie CC, Gregg EW, Geiss LS, et al. Secular changes in U.S. Prediabetes prevalence defined by hemoglobin A1c and fasting plasma glucose: National Health and Nutrition Examination Surveys, 1999-2010. *Diabetes Care*. 2013;36(8):2286-93.
22. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. *The Australasian medical journal*. 2014;7(1):45-8.
23. Chowdhury MAB, Uddin MJ, Khan HMR, Haque MR. Type 2 diabetes and its correlates among adults in Bangladesh: a population based study. *BMC Public Health*. 2015;15(1):1070.
24. Swasey KK, Gupta RD, Nayeem J, Kibria GMA. Determinants of diabetes in Bangladesh using two approaches: an analysis of the Demographic and Health Survey 2011. *Journal of Biosocial Science*. 2019;52(4):585-95.
25. Thapa S, Kayastha P, Mishra DK. Assessment of Risk of Type 2 Diabetes Among Adults of Banepa Municipality, Nepal: Community Based Cross-Sectional Study. *International Journal of Travel Medicine and Global Health*. 2020;8(1):31-6.
26. Hills AP, Arena R, Khunti K, Yajnik CS, Jayawardena R, Henry CJ, et al. Epidemiology and determinants of type 2 diabetes in south Asia. *The Lancet Diabetes & Endocrinology*. 2018;6(12):966-78.
27. BBS. Sample Vital Registration System, 2019. Dhaka: Bangladesh Bureau of Statistics (BBS), Ministry of Planning, Government of Bangladesh; 2019.
28. Afroz A, Hird TR, Zomer E, Owen A, Chen L, Ademi Z, et al. The impact of diabetes on the productivity and economy of Bangladesh. *BMJ Global Health*. 2020;5(6):e002420.
29. Rahman MM, Gilmour S, Saito E, Sultana P, Shibuya K. Self-reported illness and household strategies for coping with health-care payments in Bangladesh. *Bulletin of the World Health Organization*. 2013;91(6):449-58.
30. Akter S, Rahman MM, Abe SK, Sultana P. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bulletin of the World Health Organization*. 2014;92(3):204-13A.
31. Tripathy JP, Thakur JS, Jeet G, Chawla S, Jain S, Pal A, et al. Prevalence and risk factors of diabetes in a large community-based study in North India: results from a STEPS survey in Punjab, India. *Diabetology & Metabolic Syndrome*. 2017;9(1):8.
32. Trief PM, Himes CL, Orendorff R, Weinstock RS. The Marital Relationship and Psychosocial Adaptation and Glycemic Control of Individuals With Diabetes. *Diabetes Care*. 2001;24(8):1384-9.
33. Ramezankhani A, Azizi F, Hadaegh F. Associations of marital status with diabetes, hypertension, cardiovascular disease and all-cause mortality: A long term follow-up study. *PloS one*. 2019;14(4):e0215593-e.
34. Chande AT, Rowell J, Rishishwar L, Conley AB, Norris ET, Valderrama-Aguirre A, et al. Influence of genetic ancestry and socioeconomic status on type 2 diabetes in the diverse Colombian populations of Chocó and Antioquia. *Scientific Reports*. 2017;7(1):17127.
35. Xu Z, Yu D, Yin X, Zheng F, Li H. Socioeconomic status is associated with global diabetes prevalence. *Oncotarget*. 2017;8(27):44434-9.
36. Hossain MBH, Khan MH, Haque MA, Roy S, Hasan MS. Changing Patterns of Urbanization in Bangladesh: An Analysis of Census Data. *Population Monograph-2*. Dhaka, Bangladesh: Department of Population Sciences, University of Dhaka, and Bangladesh Bureau of Statistics; 2015. p. 1-85.




Figure 1: Trends in diabetes class by age group and sex




Figure 2: Trends in diabetes class by BMI groups and sex

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

| Section/Topic | Item # | Recommendation | Reported on page # |
|------------------------------|--------|--|--------------------|
| Title and abstract | 1 | (a) Indicate the study's design with a commonly used term in the title or the abstract | 2 |
| | | (b) Provide in the abstract an informative and balanced summary of what was done and what was found | 2 |
| Introduction | | | |
| Background/rationale | 2 | Explain the scientific background and rationale for the investigation being reported | 4-6 |
| Objectives | 3 | State specific objectives, including any prespecified hypotheses | 5 - 6 |
| Methods | | | |
| Study design | 4 | Present key elements of study design early in the paper | 6 |
| Setting | 5 | Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection | 6 |
| Participants | 6 | (a) Give the eligibility criteria, and the sources and methods of selection of participants | 6-7 |
| Variables | 7 | Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable | 7 |
| Data sources/ measurement | 8* | For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group | Not applicable |
| Bias | 9 | Describe any efforts to address potential sources of bias | Not applicable |
| Study size | 10 | Explain how the study size was arrived at | 6 |
| Quantitative variables | 11 | Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why | 8 |
| Statistical methods | 12 | (a) Describe all statistical methods, including those used to control for confounding | 8 |
| | | (b) Describe any methods used to examine subgroups and interactions | 8 |
| | | (c) Explain how missing data were addressed | 6-8 |
| | | (d) If applicable, describe analytical methods taking account of sampling strategy | 8 |

| | | | |
|--------------------------|-----|--|----------------|
| | | (e) Describe any sensitivity analyses | 8 |
| Results | | | |
| Participants | 13* | (a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed | 6- 9 |
| | | (b) Give reasons for non-participation at each stage | Not applicable |
| | | (c) Consider use of a flow diagram | Not applicable |
| Descriptive data | 14* | (a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders | 9 |
| | | (b) Indicate number of participants with missing data for each variable of interest | Not applicable |
| Outcome data | 15* | Report numbers of outcome events or summary measures | 9-10 |
| Main results | 16 | (a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included | 12-13 |
| | | (b) Report category boundaries when continuous variables were categorized | 7 |
| | | (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period | 11-12 |
| Other analyses | 17 | Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses | 12 |
| Discussion | | | |
| Key results | 18 | Summarise key results with reference to study objectives | 13 |
| Limitations | 19 | Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias | 14-16 |
| Interpretation | 20 | Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence | 3, 14-16 |
| Generalisability | 21 | Discuss the generalisability (external validity) of the study results | 14-16 |
| Other information | | | |
| Funding | 22 | Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based | Not applicable |

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

BMJ Open

Diabetes among adults in Bangladesh: changes in prevalence and risk factors from 2011 to 2018

| | |
|---------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2021-055044.R1 |
| Article Type: | Original research |
| Date Submitted by the Author: | 02-Mar-2022 |
| Complete List of Authors: | Chowdhury, Muhammad Abdul Baker; University of Florida, Emergency Medicine Islam, Mirajul; University of Dhaka, Statistics Rahman, Jakia; BRAC University, Mathematics and Natural Sciences Uddin, Md. Jamal; Shahjalal University of Science and Technology, Statistics Rabiul Haque, Md; University of Dhaka, Population Sciences |
| Primary Subject Heading: | Diabetes and endocrinology |
| Secondary Subject Heading: | Diabetes and endocrinology, Epidemiology, Global health |
| Keywords: | Epidemiology < TROPICAL MEDICINE, DIABETES & ENDOCRINOLOGY, General diabetes < DIABETES & ENDOCRINOLOGY, PUBLIC HEALTH |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Diabetes among adults in Bangladesh: changes in prevalence and risk factors from 2011 to 2018

Short title

Diabetes among adults in Bangladesh

Authors

Muhammad Abdul Baker Chowdhury, MPH, MPS, MS^{1*}; Mirajul Islam, MS^{2,3} Jakia Rahman, BS⁴; Md Jamal Uddin, PhD⁵⁺; Md Rabiul Haque, PhD⁶⁺

Affiliations

1. Department of Neurosurgery, College of Medicine, University of Florida, Gainesville, FL
2. Statistics Department, University of Florida, Gainesville FL
3. Department of Statistics, University of Dhaka, Dhaka, Bangladesh
4. Department of Mathematics and Natural Sciences, BRAC University, Dhaka, Bangladesh
5. Department of Statistics, Shahjalal University of Science & Technology, Sylhet-3114, Bangladesh
6. Department of Population Sciences, University of Dhaka, Dhaka, Bangladesh

⁺Senior authors

^{*} Corresponding author

^{*} Corresponding author email: mchow023@fiu.edu

Tables: 2 Tables and 2 Supplemental Tables

Abstract Word Count: 194

Manuscript word count: 3139

Abstract

Objective/ Research Question: To investigate the change in the prevalence and risk factors of diabetes among adults in Bangladesh between 2011 and 2018.

Design: The study was conducted using nationally representative two waves of cross-sectional data extracted from 2011 and 2017-18 Bangladesh Demographic and Health Survey.

Setting: Bangladesh

Participants: Adults age 35 years and older.

Primary outcome: Diabetes mellitus.

Results: From 2011 to 2018, the diabetes prevalence among adults aged ≥ 35 years increased from 10.95% to 13.75% ($p < 0.001$), with the largest relative increase (90%) among obese individuals. Multivariable logistic regression analysis identified age and BMI were the key risk factors of diabetes. Overweight and obese adults experienced significantly higher diabetes risk in the overall analysis in both survey years. Other significant risk factors of diabetes were marital status, education, geographic region, wealth index, and hypertension status in both survey years.

Conclusion A high prevalence of diabetes was observed and it is increasing significantly over time. Population-level approaches are needed to improve the identification and prevention of diabetes among adults in Bangladesh.

Keywords: *Diabetes, Hypertension, trends, prevalence, risk factors, Bangladesh.*

Strengths and limitations of this study

- We estimated the change in prevalence and risk factors of diabetes among Bangladeshi adults between 2011 and 2018.
- Data were obtained from nationally representative two cross-sectional surveys including the most recent one.
- Our study uniquely investigates the correlates of diabetes among adults
- Anthropometric and diabetes data collected using the WHO-recommended guidelines.
- Unavailability of information of some important correlates was the major data limitation.

Introduction

Diabetes mellitus or type 2 diabetes is one of the most common chronic and preventable diseases affecting 463 million individuals worldwide in 2019.^{1,2} By 2045, the International Diabetes Federation (IDF) predicts 700 million people will have diabetes worldwide, a 51% increase from 2019.² Morbidity, mortality, and poor quality of life are associated with this preventable disease and it is also linked to heart disease, stroke, renal failure, and blindness.³⁻⁵ Moreover, diabetes causes huge financial burden on the patient and healthcare system of the country which is expected to continue to grow. Global health expenditure on diabetes is estimated to reach \$825 billion by 2030 and \$845 billion by 2045 compared to \$760 billion in 2019.^{2,6}

With more than two-thirds of people with diabetes, low- and middle-income countries have experienced a faster growth in diabetes prevalence than high-income countries.^{1,7-10} Bangladesh, like many other countries, is transitioning from communicable to non-communicable diseases due to improved socio-economic status and unplanned but rapid urbanization.¹¹ Bangladesh is also going through a nutritional transition from traditional eating habits to a fast-food diet and sedentary lifestyle, which is contributing to the rise of non-communicable diseases like diabetes.¹² These reasons may lead Bangladesh to endure increased diabetes prevalence in the future. A systematic review of published studies between 1994 and 2013 found that diabetes prevalence in Bangladesh ranged from 4.5 to 35.0 percent.¹³ Furthermore, the number of diabetic patients in Bangladesh is estimated to be 13.7 million by 2045.¹⁴

Several studies on diabetes conducted in Bangladesh confirmed that diabetes prevalence among adults is rising steadily.^{15,16} In Bangladesh, people being living in urban areas, being member of higher income households, higher age groups, having higher education and hypertension experienced greater prevalence of diabetes.^{11,17-24} Systematic review study on

1
2 prevalence of diabetes and pre-diabetes in Bangladesh mentioned that the prevalence of diabetes
3 was significantly higher in urban areas compared with rural areas, while there was no significant
4 gender difference. ¹⁹Another cross-sectional study found that longer duration of diabetes, use of
5 insulin, and presence of diabetes complications were significantly related to the average annual
6 cost per patient in Bangladesh. ¹⁸People with lower socioeconomic status are less aware as well
7 as spend less for care of diabetes. However, most of those studies were confined to urban-rural
8 communities or some other specific groups (e.g., slum residents), which did not consider a wide
9 range of correlates of diabetes for the entire country. While an upward trend in the prevalence of
10 diabetes is evident, a very few population-based studies also reported the prevalence of diabetes,
11 which are outdated. Despite the rising literature of diabetes research in Bangladesh, no study has
12 identified trends in the prevalence of diabetes and its related risk factors or made a comparison of
13 its risk factors over the years.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 In this study, we explored whether the overall prevalence of type 2 diabetes among adults
33 in Bangladesh changed between 2011 and 2018 and to what extent it changed by individuals'
34 socioeconomic and demographic characteristics. We also examined the factors that potentially
35 contribute to the risk of diabetes among the studied population and make comparisons among
36 them. It is important to recognize changes in diabetes prevalence by population subgroups to
37 ensure access to and use of available treatment required for the population living with diabetes.
38
39
40
41
42
43
44
45
46
47
48

49 **Methods**

50 *Data source*

51 We used two waves of Bangladesh Demographic and Health Survey (BDHS) cross-
52 sectional data from 2011 and 2017-18 (written as 2018 onward) to estimate the prevalence of
53 diagnosed diabetes among the noninstitutionalized Bangladeshi population aged 35 years and
54
55
56
57
58
59
60

1
2
3 older. Diabetes testing and related questionnaires were included only in the 2011 and 2018
4 surveys. The BDHS was designed to collect data to monitor and evaluate population health and
5 nutritional status of the country using two-stage stratified cluster sampling from non-
6 institutionalized households. The details of the sampling procedure and sample selection are
7 published elsewhere.^{25 26}

8
9
10
11
12
13
14 The National Institute of Population Research and Training (NIPORT) Ethics Review
15 Board approved the data collection of the BDHS with the requirement of documented consent
16 from all study participants. Our study was exempt from the ethical review approval because we
17 used freely available de-identified data.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Biomarker measurements including blood pressure and blood glucose were collected only
in 2011 and 2018 BDHS. A total of 23,541 adults were eligible for blood glucose measurements
in both surveys. After exclusion of nonresponses and individuals with missing data, and pregnant
women, 19,584 adults comprised the study population for both survey years. Of the total
included study participants 14,376 (7,556 in 2011 and 6,820 in 2018) were age 35 years or over.

Outcome variable

The outcome variable for this study was the prevalence of diabetes for both survey years.
Diabetes status was measured by fasting blood glucose (FBG) values greater than or equal to 7.0
mmol/L or self-reported use of blood glucose lowering medication during the interview.²⁷

Demographic and other covariates

Demographic, household, and community-level characteristics were included to assess
the prevalence and risk factors of diabetes by survey years. Individual-level characteristics were
participant's (grouped into 35-44, 45-54, 55-64, 65-74, and 75+ years of age), sex, marital status
(currently married, not currently married), educational level (no education, primary, secondary,

1
2
3 higher), body mass index (BMI), and hypertension status. The BMI was calculated as weight in
4 kilograms divided by height in meters squared. We used BMI classifications for Asian
5 population: underweight (<18.5) normal weight (18.5 to 23.0), moderate risk/ overweight (23.0
6 to < 27.5), high risk/obese (≥ 27.5).²⁸ Household and community characteristics were socio-
7 economic status (wealth index), place of residence (urban, rural), and geographic region
8 (division).
9
10
11
12
13
14
15

16 17 18 *Statistical analysis*

19
20
21 The full sample of each survey was used for descriptive analysis of individuals
22 demographic and socioeconomic characteristics. Chi-square tests were performed to check the
23 bivariate association between each characteristic and diabetes status. We also used independent
24 proportion tests to know whether the prevalence of diabetes between the two surveys was
25 changed significantly and calculated the relative changes between the survey periods. For the
26 adjusted analysis in each survey years, we performed multivariable logistic regression models
27 using the to identify the associated risk factors of diabetes in Bangladesh by entering the
28 variables that were significantly associated with the outcome in the univariate logistic regression
29 analysis. Before entering the models, tests for multi-collinearity between explanatory variables
30 were performed. To select the best model, we checked the values of -2LogLikelihood ratio test,
31 AIC, and the area under the receiver operating characteristic (ROC) curve. The best model had
32 lower values of -2Log Likelihood ratio test and lower AIC value.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 To make an appropriate comparison between the two homogeneous groups for study
51 periods, adjusted odds ratios (AORs) were calculated for both 2011 and 2018 BDHS data of the
52 study participants aged 35 years and older. Moreover, we performed sensitivity analysis by
53
54
55
56
57

1
2
3 splitting the datasets into rural and urban as well as males and females. We used $P < 0.05$ at 2-
4
5 sided statistical significance for all analyses. Data management and statistical analyses were
6
7 performed using Stata 15 (StataCorp, College Station, TX, USA). We considered the sample
8
9 weights, primary sampling units, and Strata using the “SVY” command of Stata considering the
10
11 complex nature of survey design. Comparisons by different groups were drawn using the
12
13 “svysubpop” command.
14
15

16 17 **Results**

18 19 *Diabetes prevalence*

20
21 **Table 1** shows the socio-demographic characteristics of participants with age ≥ 35 years
22
23 in Bangladesh and their diabetes prevalence in 2011 and 2018 with relative ratios. The overall
24
25 prevalence of diabetes among adults ages ≥ 35 years increased from 10.95% in 2011 to 13.75% in
26
27 2018. The relative highest increase (38%) in diabetes prevalence was found among individuals
28
29 with age 65-74 and the second-highest increase rate (36%) was found among the age group 45-
30
31 54. The prevalence rate of diabetes among females increased significantly from 11.25% in 2011
32
33 to 13.81% in 2018 whereas this increment among males is not significant. The relative increase
34
35 in diabetes prevalence overtime among married, currently not working individuals and rural
36
37 areas were 30%, 42%, and 28%, respectively. A significant increase in diabetes prevalence was
38
39 also observed among adults with no education and secondary education. The highest relative
40
41 increase (54%) in diabetes prevalence was in the Dhaka region followed by 53% in the Khulna
42
43 region. Diabetes prevalence among middle, richer, and the richest individuals increased
44
45 significantly by 48%, 41%, and 33%, respectively. The prevalence of diabetes increased by 90%
46
47 among obese individuals, this rate decreased by 9% among overweight adults.
48
49
50
51
52
53
54
55
56
57
58
59
60

Risk factor analysis

The adjusted results from multivariable logistic regression analysis are reported in **Table 2**. The likelihood of diabetes was the highest [AOR: 2.11, 95% CI: 1.58, 2.83] among adults ages 55-64 in 2011 whereas this rate was highest [AOR: 1.67, 95% CI: 1.21, 2.30] in the age group 65-74 in 2018 compared to adults ages 35-44. There was no significant difference in the odds of having diabetes among males and females in both periods. Although marital status was highly insignificant (p-value=0.572) in 2011, this variable was found as marginally insignificant (p-value=0.057) in 2018.

The findings of the study also suggest that adults with primary, secondary, and higher education had 31%, 32%, and 87% higher odds of having diabetes, respectively than adults with no education in 2011. However, education was not a significant factor for diabetes among adults in 2018. Place of residence had no significant effect on diabetes in both periods. Compared to the Dhaka division, individuals living in Barisal and Chittagong divisions had a 43% and 44% higher likelihood of having diabetes, respectively in 2011. On the other hand, there exists no significant difference in having diabetes among adults in Barisal, Chittagong, Dhaka, Rajshahi, and Sylhet divisions in 2018. Regarding economic status, only the richest individuals had a significantly higher likelihood (96%) of diabetes in 2011 compared to the poorest individuals. However, both the richer and richest adults had more likelihood of diabetes [AOR: 1.84 and 3.09] than the poorest adults in 2018.

Higher BMI was a significant factor in both 2011 and 2018. For example, overweight and obese adults compared to normal-weight adults had 54% and 51% more likelihood of diabetes, respectively in 2011, and 22% and 44% higher likelihood of diabetes in 2018. Moreover, the

1
2 odds of having diabetes among working adults in 2018 was lower [AOR: 0.80, 95% CI: (0.65,
3 0.99)] than non-working adults. Since there exists a strong relationship between diabetes and
4 hypertension, individuals having hypertension had 51% and 57% more likelihoods of diabetes,
5 respectively in 2011 and 2018 compared to individuals without hypertension.
6
7
8
9
10

11 12 13 14 *Subgroup analysis*

15
16 Subgroup analysis of diabetes by sex and place of residence were also performed and the
17 results of this analysis are presented in **Table S1** and **Table S2**. Male with higher education had
18 99% higher risk of diabetes compared to male with no education in 2011 whereas no significant
19 difference in the risk of diabetes among them was observed in 2018. However, the odds of
20 having diabetes between a male with primary education and no education was not significantly
21 different [p=0.236] in 2011. The opposite scenario [AOR: 1.48 [(1.06, 2.06), p=0.022] was
22 observed in 2018. Urban males had less likelihood [AOR: 0.71, 95% CI: 0.55, 0.93] of diabetes
23 than rural males in 2018 whereas there was no difference in the likelihood of diabetes among
24 them in 2011.
25
26
27
28
29
30
31
32
33
34
35

36
37 Marital status was found to be significant factor (p=0.043) of diabetes among rural adults
38 in 2018 only. Richer individuals had no significant difference (p>0.05) in having diabetes
39 compared to the poorest individuals except in rural areas in 2018 (p=0.019). Currently working
40 individuals had 31% (AOR: 0.69, 95% CI: 0.49, 0.99) and 27% (AOR: 0.73, 95% CI: 0.57, 0.95)
41 lower odds of having diabetes than non-working rural individuals, in 2011 and 2018,
42 respectively. However, there was no significant difference in diabetes among working and non-
43 working urban individuals for both periods.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

We systematically analyze the prevalence and risk factors of diabetes among the adult population (aged ≥ 35 years) in Bangladesh using two waves of nationally representative survey data (2011 and 2018). The purpose of this study was to compare the prevalence of diabetes among people aged >35 years in 2018 with 2011 in Bangladesh. To fulfill the purpose, we analyzed two nationally representative survey data and found several remarkable findings linked to diabetes and its risk factors. One of the important findings was the identification of upward trends in the overall prevalence of diabetes and its distribution as per individual characteristics. The prevalence of diabetes among adults ≥ 35 years has significantly increased from 2011 to 2018; with a relative increase of 26%. During this period, the prevalence increased significantly not only among the overall population but also among different age groups, both sexes, married individuals, uneducated and secondary completed, rural areas, middle to richest wealth index group, obese individuals, unemployed, and hypertensive patients. This finding is consistent with the reported prevalence of diabetes among the adult populations over the years.²⁹⁻³¹

We found that the likelihood of diabetes increased with an increase of age. The odds of having diabetes was higher for older age individuals compared to the younger adults. Similar to the previous studies in Bangladesh^{32 33} and other developing countries,^{34 35} the odds of having diabetes increased consistently for all age groups of older adults (≥ 35 years) in both data sets. In the future, the upward trend of having diabetes is likely to be a major public health concern in Bangladesh owing to changes in population age-structure with lower fertility rate, steady socioeconomic growth, and increased life expectancy. This process will sharply lead to an increase in the number of middle and older age population and diabetes prevalence in Bangladesh.³⁶ The prevalence of diabetes among the working-age population may be a concern because of the complex effects of diabetes on co-morbidity and economic growth in Bangladesh,

1
2
3 where about 12% of total households pay for diabetic care by selling household assets or
4
5 borrowing money.³⁷⁻³⁹
6

7 We found a significantly higher relative increase in diabetes prevalence in rural areas
8
9 compared to urban areas, indicating that diabetes is no longer confined to urban areas in
10
11 Bangladesh because, in recent years, rural residents have adopted an unhealthy lifestyle,
12
13 consuming fast food with more carbs. Moreover, they are also less aware of the disease and seek
14
15 medical attention at a later stage. Hira et al found a similar results in a study conducted in
16
17 Bangladesh⁴⁰ and Tripathy et al.⁴¹ found a similar trend in our neighboring country India.
18
19
20

21 Our study also identified the greater prevalence of having diabetes among married people
22
23 compared to unmarried people. This finding was significant in 2018, but was not found to be
24
25 significant in 2011.³⁶ This result was broadly consistent with previous studies in which the
26
27 presence of diabetes was also associated with greater marital stability and satisfaction^{32 41 42}.
28
29 Furthermore, this could be a proxy for age because married people are older than unmarried
30
31 people, and married women who have children sometimes develop diabetes at a younger age⁴³
32
33 Moreover, male being married was also associated with a higher risk of hypertension and type-2
34
35 diabetes.⁴⁴
36
37
38

39 Although higher education and socio-economic status are negatively associated with
40
41 diabetes in developed countries, we have found the opposite results in Bangladesh for both
42
43 survey periods. Richest individuals aged ≥ 35 years are three times to have diabetes, followed by
44
45 richer and middle-income groups compared to the poorest wealth group. These findings are in
46
47 line with the previous studies conducted in Asian and other developing countries.^{32 45 46} The
48
49 greater likelihoods of diabetes among people with no education and secondary education are likely to be
50
51 associated with their less awareness about lifestyle. The prevalence of diabetes varies also by region
52
53 among individuals in Bangladesh. For example, people living in Rangpur and Khulna regions
54
55
56
57
58
59
60

1
2
3 have experienced a significantly lower risk of having diabetes than those from Dhaka and other
4 regions. The relatively lower socioeconomic status of people in these two divisions compared to
5 Dhaka is plausibly linked with their lower odds of living with diabetes.^{32 46} The rapid increase in
6 the prevalence of diabetes in all regions, particularly in Dhaka and Khulna regions, in which it
7 has increased 54% and 53% respectively between 2011 and 2018, is plausibly associated with
8 the rapid growth of urbanization and its consequences on healthy lifestyles.⁴⁷ High BMI and
9 hypertension are important factors of diabetes reported in most of the previous studies.^{11 32 37 39}
10
11 Our study findings also pointed out that with higher BMI have a greater likelihood of having
12 diabetes compared to normal weight adults in both 2011 and 2018.⁴⁸⁻⁵²
13
14
15
16
17
18
19
20
21
22

23 One of the major strengths of our study is the use of nationally representative cross-
24 sectional survey data of the two waves, including the most recent one which is released in
25 December 2020. Data of these surveys relates to anthropometric and diabetes were not self-
26 reported rather collected by the trained and experienced health workers such as nurses,
27 midwives, health assistance using the WHO-recommended guidelines. To our knowledge, this
28 study for the first time estimated the national diabetes prevalence and its risk factors among
29 adults of Bangladesh. Another important strength is that we compared changes in estimates of
30 diabetes predictors between 2011 and 2018 surveys along with subgroups analyses, by sex, and
31 by place of residence. Despite having some strength, our study is not beyond the limitations. Due
32 to lack of diabetes data for younger adults (<35 years) in previous surveys, we could not
33 compare their current diabetes prevalence. Moreover, the unavailability of data of some
34 important correlates like the types of diet, intake of fast food, intake of calories, physical exercise
35 including the nature of work, family history of diabetes, and cholesterol level of diabetes was the
36 major limitation of these data sets. As a cross-sectional survey, blood sugar level was measured
37 for one day only and thus we do not have follow-up and or longitudinal data on diabetes and its
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 correlates. Due to the nature of survey data, this study identified the risk factors of diabetes only
4
5 rather than causality.
6

7 The study shows that among Bangladeshi adults, there is a high prevalence of diabetes
8
9 and it is escalating over time. The study also reports a significant portion of younger adults with
10
11 diabetes. Age and overweight/obesity are the two most important risk factors for diabetes for all
12
13 adults, irrespective of sex, residence, educational attainment and wealth index. There is evidence
14
15 of an increase in the magnitude of diabetes over time and in the younger population; indeed, as
16
17 age increases the chances of developing Diabetes mellitus significantly. These findings, together
18
19 with an increase in the prevalence of type 2 diabetes among Bangladeshi adults, underscore the
20
21 need for primary (awareness campaign about the adversity of fast food habit and impotence of
22
23 physical mobility/work/exercise) and secondary (incorporate the issue text on the causes and
24
25 consequences of diabetes and NCDs in school level curriculum) prevention efforts tailored to
26
27 age-specific populations.
28
29
30
31
32

33 **List of Abbreviations**

| | |
|---------|--|
| 34 AOR | Adjusted Odds Ratio |
| 35 BMI | Body mass index |
| 36 CI | Confidence interval |
| 37 CVD | Cardiovascular diseases |
| 38 WHO | World Health Organization |
| 39 BDHS | Bangladesh Demographic and Health Survey |

40
41
42
43
44
45

46 **Availability of data and material:** All data presented here in the manuscript is freely available at
47
48 dhsprogram.com.
49

50
51 **Prior Publication:** This data has not been published previously and is not under consideration
52
53 elsewhere.
54

55 **Author Contribution:** M.A.B. Chowdhury conceptualized the study, designed the analytic
56
57 approach, managed and performed the analysis, interpreted the results, and drafted the
58

1
2
3 manuscript. M. Islam: helped with the analysis, interpreted the results, and drafted the
4 manuscript. J. Rahman, M.J. Uddin reviewed, edited and updated the manuscript and M.R.Haque
5 reviewed, edited, and supervised the study.
6
7
8

9
10 **Disclosures/Conflict of Interest:** There are no potential conflicts (financial, professional, or
11 personal) to disclose by any of the authors.
12
13

14
15 **Funding:** None

16
17 **Acknowledgement:** We thank DHS program for providing the data.
18
19

20 **Ethical approval and consent**

21 All BDHS surveys received ethical approval from ICF Macro Institutional Review Board,
22 Maryland, USA and National Research Ethics Committee of Bangladesh Medical Research
23 Council (BMRC), Dhaka, Bangladesh. Informed consent was obtained from each participant of
24 the survey before enrolling in the survey by using the Introduction and Consent form of the
25 survey. It was also explained that the information will be kept strictly confidential and will not
26 be shared with anyone except members of the survey team.
27
28
29
30
31
32

33 **Patient and Public Involvement**

34 The BDHS questionnaires were based on the MEASURE DHS model questionnaires. These
35 model questionnaires were adapted for use in Bangladesh during a series of meetings with a
36 technical working group (TWG) that consisted of representatives from NIPORT, Mitra and
37 Associates, International Centre for Diarrheal Diseases and Control, Bangladesh (icddr, b),
38 USAID/Bangladesh, and MEASURE DHS. Patients were not directly involved in the study
39 however; the TWG involved representatives from the government, non-government, ministry of
40 health and family welfare representatives and donor organizations were involved the study
41 design and questionnaire development. The results will be used by the health researchers, policy
42 makers of the country.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Description of tables

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

TABLE 2. Adjusted odds ratios for factors predicting diabetes among adults 35 years and older BDHS 2011-2018

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|-----------------------------------|---------------------------------|----------------------------------|----------------------------------|--------------------------------|-------|
| All adults age 35 years and older | 100% | 10.95 (0.0048) | 13.75 (0.0056) | <0.001 | 1.26 |
| Age group | | | | | |
| 35-44 | 35.99 | 8.82 (0.0065) | 11.21 (0.0078) | 0.081 | 1.27 |
| 45-54 | 27.43 | 10.86 (0.0079) | 14.82 (0.01) | 0.008 | 1.36 |
| 55-64 | 18.65 | 14.69 (0.012) | 15.78 (0.0115) | 0.6286 | 1.07 |
| 65-74 | 11.26 | 11.60 (0.0128) | 15.97 (0.016) | 0.1123 | 1.38 |
| 75+ | 6.68 | 12.11 (0.0174) | 13.43 (0.0189) | 0.793 | 1.11 |
| Sex | | | | | |
| Male | 48.98 | 10.65 (0.0061) | 13.69 (0.0072) | 0.091 | 1.29 |
| Female | 51.02 | 11.25 (0.006) | 13.81 (0.0072) | 0.003 | 1.23 |
| Marital status | | | | | |
| Not married | 15.88 | 12.13 (0.0112) | 12.54 (0.0111) | 0.953 | 1.03 |
| Married | 84.12 | 10.73 (0.0049) | 13.98 (0.006) | <0.001 | 1.3 |
| Educational level | | | | | |
| No education | 44.8 | 8.39 (0.0058) | 10.64 (0.0072) | 0.021 | 1.27 |
| Primary | 29.07 | 11.10 (0.0078) | 13.59 (0.009) | 0.141 | 1.22 |
| Secondary | 18.15 | 13.03 (0.0108) | 17.8 (0.0121) | 0.015 | 1.37 |
| Higher | 7.98 | 21.79 (0.018) | 20.27 (0.0185) | 0.209 | 0.93 |
| Place of residence | | | | | |
| Urban | 23.91 | 16.08 (0.0106) | 18.95 (0.0106) | 0.129 | 1.18 |
| Rural | 76.09 | 9.39 (0.0051) | 12.06 (0.0065) | 0.003 | 1.28 |
| Geographic region | | | | | |
| Barisal | 5.79 | 12.54 (0.0117) | 12.09 (0.0161) | 0.975 | 0.96 |
| Chittagong | 16.45 | 14.28 (0.0125) | 17.33 (0.0169) | 0.087 | 1.21 |
| Dhaka | 38.99 | 11.26 (0.0107) | 17.32 (0.0118) | 0.002 | 1.54 |
| Khulna | 13.45 | 7.30 (0.007) | 11.14 (0.0114) | 0.002 | 1.53 |
| Rajshahi | 13.15 | 10.73 (0.01) | 11.32 (0.0134) | 0.664 | 1.05 |
| Rangpur | 9.26 | 8.59 (0.0111) | 7.93 (0.0097) | 0.667 | 0.92 |
| Sylhet | 2.91 | 11.85 (0.011) | 12.71 (0.0169) | 0.998 | 1.07 |
| Wealth index | | | | | |
| Poorest | 19.91 | 7.28 (0.0088) | 7.26 (0.0091) | 0.69 | 1.00 |
| Poorer | 19.74 | 7.35 (0.0084) | 7.70 (0.0085) | 0.891 | 1.05 |
| Middle | 20.19 | 7.56 (0.0075) | 11.17 (0.01) | 0.005 | 1.48 |
| Richer | 19.70 | 11.33 (0.0098) | 16.01 (0.0128) | <0.001 | 1.41 |
| Richest | 20.46 | 20.49 (0.0122) | 27.18 (0.0134) | 0.006 | 1.33 |
| Body Mass Index | | | | | |
| Underweight | 19.87 | 7.27 (0.0081) | 7.79 (0.0087) | 0.437 | 1.07 |
| Normal weight | 36.10 | 9.59 (0.0071) | 11.21 (0.0074) | 0.343 | 1.17 |
| Overweight | 20.14 | 18.57 (0.0149) | 16.95 (0.0095) | 0.045 | 0.91 |
| Obese | 23.88 | 11.90 (0.0078) | 22.56 (0.0156) | <0.001 | 1.9 |
| Currently working | | | | | |

TABLE 1. Sociodemographic characteristics of adults age 35 years and older and diabetes rate in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|--------------|---------------------------------|----------------------------------|----------------------------------|--------------------------------|-------|
| No | 43.63 | 12.11 (0.0064) | 17.16 (0.0089) | <0.001 | 1.42 |
| Yes | 56.37 | 9.70 (0.0061) | 12.00 (0.006) | 0.10 | 1.24 |
| Hypertension | | | | | |
| No | 67.45 | 9.06 (0.0048) | 10.55 (0.0063) | 0.522 | 1.16 |
| Yes | 32.55 | 16.33 (0.0105) | 18.57 (0.009) | 0.178 | 1.14 |

For peer review only

TABLE 2. Adjusted odds ratios for factors predicting diabetes among adults 35 years and older BDHS 2011-2018.

| Variables | 2011 BDHS | | 2017-18 BDHS | |
|---------------------------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | |
| 35-44 | Ref | | Ref | |
| 45-54 | 1.36 (1.07, 1.72) | 0.012 | 1.41 (1.13, 1.76) | 0.002 |
| 55-64 | 2.11 (1.58, 2.83) | <0.001 | 1.58 (1.24, 2.03) | <0.001 |
| 65-74 | 1.60 (1.13, 2.27) | 0.008 | 1.67 (1.21, 2.30) | 0.002 |
| 75+ | 1.77 (1.14, 2.74) | 0.011 | 1.32 (0.86, 2.01) | 0.202 |
| Sex | | | | |
| Male | Ref | | Ref | |
| Female | 0.78 (0.58, 1.05) | 0.104 | 0.92 (0.75, 1.13) | 0.435 |
| Marital status | | | | |
| Not married | Ref | | | |
| Married | 0.93 (0.72, 1.20) | 0.572 | 1.28 (0.99, 1.64) | 0.057 |
| Educational level | | | | |
| No education | Ref | | Ref | |
| Primary | 1.31 (1.05, 1.64) | 0.017 | 1.23 (0.99, 1.53) | 0.059 |
| Secondary | 1.32 (1.01, 1.73) | 0.045 | 1.23 (0.96, 1.57) | 0.108 |
| Higher | 1.87 (1.35, 2.60) | <0.001 | 1.15 (0.83, 1.60) | 0.397 |
| Place of residence | | | | |
| Urban | 1.08 (0.87, 1.33) | 0.7 | 0.96 (0.79, 1.16) | 0.678 |
| Rural | Ref | | Ref | |
| Geographic region | | | | |
| Barisal | 1.43 (1.04, 1.96) | 0.027 | 0.75 (0.54, 1.06) | 0.103 |
| Chittagong | 1.44 (1.09, 1.89) | 0.010 | 0.88 (0.67, 1.16) | 0.369 |
| Dhaka | Ref | | Ref | |
| Khulna | 0.66 (0.50, 0.89) | 0.007 | 0.60 (0.46, 0.79) | <0.001 |
| Rajshahi | 1.17 (0.87, 1.57) | 0.309 | 0.74 (0.55, 1.00) | 0.054 |
| Rangpur | 1.00 (0.69, 1.45) | 0.986 | 0.56 (0.41, 0.76) | <0.001 |
| Sylhet | 1.22 (0.91, 1.63) | 0.183 | 0.75 (0.53, 1.06) | 0.101 |
| Wealth index | | | | |
| Poorest | Ref | | Ref | |
| Poorer | 0.89 (0.62, 1.27) | 0.514 | 0.99 (0.70, 1.40) | 0.957 |
| Middle | 0.86 (0.62, 1.19) | 0.365 | 1.33 (0.94, 1.88) | 0.103 |
| Richer | 1.18 (0.84, 1.65) | 0.345 | 1.84 (1.29, 2.61) | 0.001 |
| Richest | 1.96 (1.40, 2.76) | <0.001 | 3.09 (2.18, 4.38) | <0.001 |
| Body Mass Index | | | | |
| Underweight | 0.82 (0.61, 1.10) | 0.177 | 0.77 (0.58, 1.01) | 0.055 |
| Normal weight | Ref | | Ref | |
| Overweight | 1.54 (1.20, 1.97) | 0.001 | 1.22 (1.00, 1.50) | 0.052 |
| Obese | 1.51 (1.16, 1.97) | 0.003 | 1.44 (1.13, 1.84) | 0.003 |
| Currently working | | | | |
| No | Ref | | Ref | |
| Yes | 0.77 (0.59, 1.01) | 0.056 | 0.80 (0.65, 0.99) | 0.039 |
| Hypertension | | | | |
| No | Ref | | Ref | |
| Yes | 1.51 (1.26, 1.81) | <0.001 | 1.57 (1.32, 1.87) | <0.001 |

References

1. Ogurtsova K, da Rocha Fernandes JD, Huang Y, et al. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes research and clinical practice* 2017;128:40-50.
2. Federation ID. IDF Diabetes Atlas Ninth. *Dunia: IDF* 2019
3. Kumar SS, Houlden RL. Ethnocultural diversity and the diabetes epidemic in Canada: a call to action. *Can J Diab* 2005;29(2):84-5.
4. Oldroyd J, Banerjee M, Heald A, et al. Diabetes and ethnic minorities. *Postgraduate medical journal* 2005;81(958):486-90.
5. World Health O. World health statistics 2013: a wealth of information on global public health: World Health Organization, 2013.
6. Williams R, Karuranga S, Malanda B, et al. Global and regional estimates and projections of diabetes-related health expenditure: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract* 2020;162:108072. doi: 10.1016/j.diabres.2020.108072 [published Online First: 2020/02/18]
7. Dagenais GR, Gerstein HC, Zhang X, et al. Variations in diabetes prevalence in low-, middle-, and high-income countries: results from the prospective urban and rural epidemiological study. *Diabetes care* 2016;39(5):780-87.
8. Mkuu R, Barry A, Yonga G, et al. Prevalence and factors associated with overweight and obesity in Kenya. *Preventive Medicine Reports* 2021;22:101340. doi: <https://doi.org/10.1016/j.pmedr.2021.101340>
9. Mkuu RS, Gilreath TD, Barry AE, et al. Identifying individuals with multiple non-communicable disease risk factors in Kenya: a latent class analysis. *Public Health* 2021;198:180-86. doi: <https://doi.org/10.1016/j.puhe.2021.07.031>
10. Chowdhury MAB, Epnere K, Haque MA, et al. Urban rural differences in prevalence and risk factors of self-reported hypertension among Kenyan women: a population-based study. *Journal of Human Hypertension* 2021;35(10):912-20. doi: 10.1038/s41371-020-00435-x
11. Chowdhury MAB, Islam M, Rahman J, et al. Changes in prevalence and risk factors of hypertension among adults in Bangladesh: An analysis of two waves of nationally representative surveys. *PLOS ONE* 2021;16(12):e0259507. doi: 10.1371/journal.pone.0259507
12. Amuna P, Zotor FB. Epidemiological and nutrition transition in developing countries: impact on human health and development. *Proc Nutr Soc* 2008;67(1):82-90. doi: 10.1017/s0029665108006058 [published Online First: 2008/02/01]
13. Biswas T, Islam A, Rawal LB, et al. Increasing prevalence of diabetes in Bangladesh: a scoping review. *Public Health* 2016;138:4-11. doi: 10.1016/j.puhe.2016.03.025 [published Online First: 2016/05/14]
14. Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes research and clinical practice* 2018;138:271-81.
15. Rahim MA, Hussain A, Khan AKA, et al. Rising prevalence of type 2 diabetes in rural Bangladesh: a population based study. *Diabetes research and clinical practice* 2007;77(2):300-05.
16. Bhowmik B, Afsana F, My Diep L, et al. Increasing prevalence of type 2 diabetes in a rural Bangladeshi population: a population based study for 10 years. *Diabetes & metabolism journal* 2013;37(1):46-53.
17. Alam DS, Talukder SH, Chowdhury MAH, et al. Overweight and abdominal obesity as determinants of undiagnosed diabetes and pre-diabetes in Bangladesh. *BMC Obesity* 2016;3(1):19. doi: 10.1186/s40608-016-0099-z

18. Afroz A, Alam K, Ali L, et al. Type 2 diabetes mellitus in Bangladesh: a prevalence based cost-of-illness study. *BMC Health Services Research* 2019;19(1):601. doi: 10.1186/s12913-019-4440-3
19. Akhtar S, Nasir JA, Sarwar A, et al. Prevalence of diabetes and pre-diabetes in Bangladesh: a systematic review and meta-analysis. *BMJ open* 2020;10(9):e036086. doi: 10.1136/bmjopen-2019-036086
20. Haque M, Islam S, Kamal ZM, et al. Ongoing efforts to improve the management of patients with diabetes in Bangladesh and the implications. *Hospital Practice* 2021;49(4):266-72. doi: 10.1080/21548331.2021.1906083
21. Islam SMS, Niessen LW, Seissler J, et al. Diabetes knowledge and glycemic control among patients with type 2 diabetes in Bangladesh. *SpringerPlus* 2015;4(1):284. doi: 10.1186/s40064-015-1103-7
22. Rahman MS, Akter S, Abe SK, et al. Awareness, treatment, and control of diabetes in Bangladesh: a nationwide population-based study. *PloS one* 2015;10(2):e0118365-e65. doi: 10.1371/journal.pone.0118365
23. Shariful Islam SM, Lechner A, Ferrari U, et al. Healthcare use and expenditure for diabetes in Bangladesh. *BMJ Global Health* 2017;2(1):e000033. doi: 10.1136/bmjgh-2016-000033
24. Talukder A, Hossain MZ. Prevalence of Diabetes Mellitus and Its Associated Factors in Bangladesh: Application of Two-level Logistic Regression Model. *Scientific Reports* 2020;10(1):10237. doi: 10.1038/s41598-020-66084-9
25. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2011, 2011.
26. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2014, 2014.
27. World Health O. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia: report of a WHO/IDF consultation. 2006
28. Who EC. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet (London, England)* 2004;363(9403):157.
29. Menke A, Casagrande S, Geiss L, et al. Prevalence of and Trends in Diabetes Among Adults in the United States, 1988-2012. *JAMA* 2015;314(10):1021-29. doi: 10.1001/jama.2015.10029
30. Bullard KM, Saydah SH, Imperatore G, et al. Secular changes in U.S. Prediabetes prevalence defined by hemoglobin A1c and fasting plasma glucose: National Health and Nutrition Examination Surveys, 1999-2010. *Diabetes Care* 2013;36(8):2286-93. doi: 10.2337/dc12-2563 [published Online First: 2013/04/23]
31. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. *The Australasian medical journal* 2014;7(1):45-48. doi: 10.4066/AMJ.2013.1979
32. Chowdhury MAB, Uddin MJ, Khan HMR, et al. Type 2 diabetes and its correlates among adults in Bangladesh: a population based study. *BMC Public Health* 2015;15(1):1070. doi: 10.1186/s12889-015-2413-y
33. Swasey KK, Gupta RD, Nayeem J, et al. Determinants of diabetes in Bangladesh using two approaches: an analysis of the Demographic and Health Survey 2011. *Journal of Biosocial Science* 2019;52(4):585-95. doi: 10.1017/S002193201900066X [published Online First: 10/21]
34. Thapa S, Kayastha P, Mishra DK. Assessment of Risk of Type 2 Diabetes Among Adults of Banepa Municipality, Nepal: Community Based Cross-Sectional Study. *International Journal of Travel Medicine and Global Health* 2020;8(1):31-36. doi: 10.34172/ijtmgh.2020.05
35. Hills AP, Arena R, Khunti K, et al. Epidemiology and determinants of type 2 diabetes in south Asia. *The Lancet Diabetes & Endocrinology* 2018;6(12):966-78. doi: [https://doi.org/10.1016/S2213-8587\(18\)30204-3](https://doi.org/10.1016/S2213-8587(18)30204-3)

- 1
- 2
- 3
- 4 36. BBS. Sample Vital Registration System, 2019. Dhaka: Bangladesh Bureau of Statistics (BBS), Ministry
- 5 of Planning, Government of Bangladesh 2019.
- 6 37. Afroz A, Hird TR, Zomer E, et al. The impact of diabetes on the productivity and economy of
- 7 Bangladesh. *BMJ Global Health* 2020;5(6):e002420. doi: 10.1136/bmjgh-2020-002420
- 8 38. Rahman MM, Gilmour S, Saito E, et al. Self-reported illness and household strategies for coping with
- 9 health-care payments in Bangladesh. *Bulletin of the World Health Organization* 2013;91(6):449-
- 10 58. doi: 10.2471/BLT.12.115428
- 11 39. Akter S, Rahman MM, Abe SK, et al. Prevalence of diabetes and prediabetes and their risk factors
- 12 among Bangladeshi adults: a nationwide survey. *Bulletin of the World Health Organization*
- 13 2014;92(3):204-13A. doi: 10.2471/BLT.13.128371 [published Online First: 01/10]
- 14 40. Hira R, Miah MAW, Akash DH. Prevalence of type 2 diabetes mellitus in rural adults (> _31years) in
- 15 Bangladesh. *Faridpur Medical College Journal* 2018;13(1):20-23.
- 16 41. Tripathy JP, Thakur JS, Jeet G, et al. Prevalence and risk factors of diabetes in a large community-
- 17 based study in North India: results from a STEPS survey in Punjab, India. *Diabetology &*
- 18 *Metabolic Syndrome* 2017;9(1):8. doi: 10.1186/s13098-017-0207-3
- 19 42. Trief PM, Himes CL, Orendorff R, et al. The Marital Relationship and Psychosocial Adaptation and
- 20 Glycemic Control of Individuals With Diabetes. *Diabetes Care* 2001;24(8):1384-89. doi:
- 21 10.2337/diacare.24.8.1384
- 22 43. Cdcgov. Gestational Diabetes and Pregnancy | CDC: @CDCgov; 2020 [updated 2020-07-
- 23 14T06:42:06Z. Available from: <https://www.cdc.gov/pregnancy/diabetes-gestational.html>.
- 24 44. Ramezankhani A, Azizi F, Hadaegh F. Associations of marital status with diabetes, hypertension,
- 25 cardiovascular disease and all-cause mortality: A long term follow-up study. *PloS one*
- 26 2019;14(4):e0215593-e93. doi: 10.1371/journal.pone.0215593
- 27 45. Chande AT, Rowell J, Rishishwar L, et al. Influence of genetic ancestry and socioeconomic status on
- 28 type 2 diabetes in the diverse Colombian populations of Chocó and Antioquia. *Scientific Reports*
- 29 2017;7(1):17127. doi: 10.1038/s41598-017-17380-4
- 30 46. Xu Z, Yu D, Yin X, et al. Socioeconomic status is associated with global diabetes prevalence.
- 31 *Oncotarget* 2017;8(27):44434-39. doi: 10.18632/oncotarget.17902
- 32 47. Hossain MBH, Khan MH, Haque MA, et al. Changing Patterns of Urbanization in Bangladesh: An
- 33 Analysis of Census Data. Population Monograph-2. Dhaka, Bangladesh: Department of
- 34 Population Sciences, University of Dhaka, and Bangladesh Bureau of Statistics, 2015:1-85.
- 35 48. Rahman MA, Halder HR, Yadav UN, et al. Prevalence of and factors associated with hypertension
- 36 according to JNC 7 and ACC/AHA 2017 guidelines in Bangladesh. *Scientific Reports*
- 37 2021;11(1):15420. doi: 10.1038/s41598-021-94947-2
- 38 49. Mkuu RS, Epnere K, Chowdhury MAB. Prevalence and Predictors of Overweight and Obesity Among
- 39 Kenyan Women. *Prev Chronic Dis* 2018;15:E44. doi: 10.5888/pcd15.170401
- 40 50. Chowdhury MAB, Adnan MM, Hassan MZ. Trends, prevalence and risk factors of overweight and
- 41 obesity among women of reproductive age in Bangladesh: a pooled analysis of five national
- 42 cross-sectional surveys. *BMJ open* 2018;8(7):e018468. doi: 10.1136/bmjopen-2017-018468
- 43 51. Mohammad M, Chowdhury MAB, Islam MN, et al. Factors associated with body mass index among
- 44 university students in Bangladesh. *Journal of Public Health* 2019 doi: 10.1007/s10389-019-
- 45 01124-x
- 46 52. Mohammad M, Chowdhury MAB, Islam MN, et al. Factors associated with body mass index among
- 47 university students in Bangladesh. *Journal of Public Health* 2021;29(2):345-51. doi:
- 48 10.1007/s10389-019-01124-x
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|---------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.56 (1.1, 2.21) | 0.013 | 1.41 (0.98, 2.03) | 0.065 | 1.18 (0.87, 1.62) | 0.288 | 1.41 (1.06, 1.86) | 0.018 |
| 55-64 | 2.36 (1.6, 3.47) | <0.001 | 1.58 (1.09, 2.29) | 0.015 | 1.84 (1.15, 2.94) | 0.012 | 1.53 (1.09, 2.14) | 0.013 |
| 65-74 | 1.79 (1.14, 2.8) | 0.011 | 1.80 (1.18, 2.75) | 0.007 | 1.35 (0.76, 2.39) | 0.304 | 1.44 (0.9, 2.33) | 0.132 |
| 75+ | 1.52 (0.85, 2.72) | 0.153 | 1.41 (0.79, 2.51) | 0.24 | 1.92 (0.98, 3.76) | 0.059 | 1.11 (0.62, 1.99) | 0.727 |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.66 (0.35, 1.26) | 0.205 | 1.83 (0.96, 3.47) | 0.064 | 1.00 (0.75, 1.34) | 0.992 | 1.17 (0.89, 1.56) | 0.263 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.22 (0.88, 1.68) | 0.236 | 1.48 (1.06, 2.06) | 0.022 | 1.37 (1, 1.86) | 0.047 | 1.12 (0.84, 1.48) | 0.434 |
| Secondary | 1.14 (0.80, 1.63) | 0.47 | 1.15 (0.78, 1.69) | 0.482 | 1.53 (1.05, 2.23) | 0.029 | 1.37 (0.97, 1.92) | 0.071 |
| Higher | 1.99 (1.31, 3.01) | 0.001 | 1.19 (0.76, 1.86) | 0.449 | 1.85 (1.1, 3.11) | 0.02 | 1.98 (0.59, 1.64) | 0.941 |
| Place of residence | | | | | | | | |
| Urban | 1.03 (0.76, 1.4) | 0.838 | 0.71 (0.55, 0.93) | 0.012 | 1.1 (0.83, 1.45) | 0.497 | 1.24 (0.96, 1.61) | 0.097 |
| Rural | Ref | | Ref | | Ref | | Ref | |
| Geographic region | | | | | | | | |
| Barisal | 1.42 (0.91, 2.22) | 0.121 | 0.83 (0.53, 1.30) | 0.415 | 1.44 (0.97, 2.14) | 0.072 | 0.72 (0.46, 1.12) | 0.148 |
| Chittagong | 1.55 (1.05, 2.27) | 0.027 | 0.88 (0.62, 1.26) | 0.484 | 1.36 (0.97, 1.91) | 0.071 | 0.88 (0.61, 1.26) | 0.48 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.75 (0.50, 1.12) | 0.159 | 0.56 (0.39, 0.82) | 0.003 | 0.61 (0.4, 0.92) | 0.02 | 0.65 (0.46, 0.93) | 0.019 |
| Rajshahi | 1.14 (0.73, 1.77) | 0.569 | 0.78 (0.52, 1.2) | 0.259 | 1.21 (0.82, 1.77) | 0.338 | 0.71 (0.46, 1.09) | 0.116 |
| Rangpur | 1.04 (0.64, 1.68) | 0.885 | 0.60 (0.39, 0.92) | 0.019 | 0.99 (0.63, 1.57) | 0.975 | 0.53 (0.34, 0.82) | 0.004 |
| Sylhet | 1.41 (0.92, 2.15) | 0.112 | 0.66 (0.41, 1.07) | 0.091 | 1.11 (0.77, 1.6) | 0.58 | 0.82 (0.54, 1.22) | 0.325 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |
| Poorer | 0.81 (0.51, 1.30) | 0.389 | 1.20 (0.72, 2.00) | 0.485 | 0.96 (0.58, 1.59) | 0.88 | 0.87 (0.58, 1.32) | 0.522 |
| Middle | 0.75 (0.46, 1.21) | 0.239 | 1.90 (1.16, 3.12) | 0.011 | 0.99 (0.63, 1.57) | 0.981 | 1.02 (0.66, 1.57) | 0.945 |

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Richer | 0.95 (0.59, 1.51) | 0.815 | 2.58 (1.53, 4.34) | <0.001 | 1.43 (0.9, 2.25) | 0.129 | 1.44 (0.94, 2.21) | 0.093 |
| Richest | 1.54 (0.94, 2.53) | 0.088 | 5.47 (3.25, 9.19) | <0.001 | 2.38 (1.49, 3.79) | <0.001 | 2.03 (1.33, 3.09) | 0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.86 (0.6, 1.22) | 0.382 | 0.76 (0.51, 1.13) | 0.177 | 0.69 (0.44, 1.08) | 0.101 | 0.78 (0.52, 1.18) | 0.238 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 1.54 (1.13, 2.12) | 0.007 | 1.25 (0.94, 1.66) | 0.122 | 1.63 (1.09, 2.42) | 0.016 | 1.15 (0.86, 1.52) | 0.343 |
| Obese | 1.71 (1.1, 2.68) | 0.018 | 1.82 (1.22, 2.69) | 0.003 | 1.31 (0.86, 1.98) | 0.207 | 1.29 (0.95, 1.76) | 0.105 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.71 (0.48, 1.06) | 0.09 | 0.80 (0.54, 1.17) | 0.249 | 0.84 (0.55, 1.26) | 0.394 | 0.79 (0.61, 1.03) | 0.076 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.17 (0.88, 1.56) | 0.269 | 1.42 (1.1, 1.83) | 0.007 | 1.77 (1.4, 2.24) | <0.001 | 1.70 (1.33, 2.18) | <0.001 |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.58 (1.07, 2.36) | 0.023 | 1.19 (0.85, 1.65) | 0.312 | 1.22 (1.66, 0.9) | 0.194 | 1.59 (1.18, 2.13) | 0.002 |
| 55-64 | 3.57 (2.16, 5.9) | <0.001 | 1.50 (0.98, 2.29) | 0.059 | 1.60 (2.3, 1.11) | 0.011 | 1.67 (1.24, 2.27) | 0.001 |
| 65-74 | 2.25 (1.24, 4.1) | 0.008 | 1.50 (0.84, 2.66) | 0.168 | 1.32 (2.04, 0.86) | 0.200 | 1.75 (1.19, 2.57) | 0.004 |
| 75+ | 2.54 (1.17, 5.5) | 0.018 | 1.89 (1.01, 3.51) | 0.045 | 1.45 (2.45, 0.85) | 0.171 | 1.14 (0.65, 2.01) | 0.645 |
| Sex | | | | | | | | |
| Female | 0.82 (0.49, 1.38) | 0.461 | 1.29 (0.9, 1.85) | 0.173 | 0.76 (1.11, 0.52) | 0.15 | 0.88 (0.63, 1.03) | 0.087 |
| Male | Ref | | Ref | | Ref | | Ref | |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.8 (0.52, 1.24) | 0.322 | 1.07 (0.7, 1.62) | 0.759 | 1.01 (1.38, 0.74) | 0.962 | 1.39 (1.01, 1.92) | 0.043 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.44 (0.96, 2.14) | 0.076 | 1 (0.69, 1.45) | 0.992 | 1.26 (1.64, 0.97) | 0.086 | 1.30 (1.00, 1.69) | 0.048 |
| Secondary | 1.46 (0.94, 2.27) | 0.089 | 1.35 (0.89, 2.03) | 0.155 | 1.22 (1.72, 0.86) | 0.272 | 1.15 (0.83, 1.59) | 0.394 |
| Higher | 1.9 (1.13, 3.19) | 0.015 | 1.19 (0.74, 1.93) | 0.465 | 1.85 (2.94, 1.16) | 0.009 | 1.15 (0.74, 1.81) | 0.532 |
| Geographic region | | | | | | | | |
| Barisal | 0.93 (0.58, 1.49) | 0.76 | 0.99 (0.54, 1.83) | 0.982 | 1.63 (2.45, 1.09) | 0.018 | 0.70 (0.46, 1.06) | 0.088 |
| Chittagong | 1.04 (0.69, 1.58) | 0.845 | 0.58 (0.4, 0.84) | 0.004 | 1.66 (2.41, 1.15) | 0.007 | 1.04 (0.72, 1.5) | 0.841 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.82 (0.52, 1.28) | 0.381 | 0.70 (0.5, 1.00) | 0.05 | 0.61 (0.91, 0.4) | 0.017 | 0.57 (0.4, 0.81) | 0.002 |
| Rajshahi | 0.94 (0.60, 1.48) | 0.802 | 0.74 (0.51, 1.06) | 0.095 | 1.28 (1.9, 0.86) | 0.228 | 0.75 (0.51, 1.12) | 0.165 |
| Rangpur | 0.77 (0.46, 1.29) | 0.318 | 0.5 (0.33, 0.78) | 0.002 | 1.07 (1.71, 0.67) | 0.775 | 0.58 (0.4, 0.84) | 0.005 |
| Sylhet | 1.43 (0.89, 2.31) | 0.143 | 0.71 (0.47, 1.09) | 0.115 | 1.22 (1.79, 0.83) | 0.318 | 0.77 (0.5, 1.18) | 0.226 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Poorer | 1.23 (0.49, 3.07) | 0.653 | 0.87 (0.36, 2.08) | 0.754 | 0.87 (1.27, 0.6) | 0.468 | 0.98 (0.67, 1.42) | 0.896 |
| Middle | 1.05 (0.49, 2.24) | 0.900 | 0.67 (0.3, 1.51) | 0.335 | 0.84 (1.2, 0.58) | 0.327 | 1.42 (0.97, 2.08) | 0.07 |
| Richer | 0.92 (0.49, 1.72) | 0.787 | 1.74 (0.81, 3.76) | 0.155 | 1.27 (1.85, 0.87) | 0.218 | 1.54 (1.08, 2.47) | 0.019 |
| Richest | 1.67 (0.89, 3.11) | 0.108 | 2.54 (1.21, 5.34) | 0.014 | 2.05 (3.05, 1.38) | <0.001 | 3.05 (2.03, 4.60) | <0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.96 (0.55, 1.68) | 0.884 | 1.24 (0.73, 2.1) | 0.422 | 0.77 (1.07, 0.55) | 0.119 | 0.58 (0.49, 0.94) | 0.02 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 2.27 (1.47, 3.5) | <0.001 | 0.99 (0.7, 1.41) | 0.971 | 1.32 (1.8, 0.97) | 0.077 | 1.16 (1.06, 1.73) | 0.015 |
| Obese | 2.86 (1.88, 4.34) | <0.001 | 1.5 (1.04, 2.16) | 0.029 | 1.1 (1.55, 0.78) | 0.581 | 1.36 (0.98, 1.89) | 0.068 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.91 (0.6, 1.38) | 0.664 | 1.04 (0.73, 1.48) | 0.828 | 0.69 (0.49, 0.99) | 0.044 | 0.73 (0.57, 0.95) | 0.02 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.52 (1.15, 2.02) | 0.004 | 1.58 (1.18, 2.11) | 0.002 | 1.50 (1.91, 1.18) | 0.001 | 1.16 (1.29, 1.99) | <0.001 |

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

| Section/Topic | Item # | Recommendation | Reported on page # |
|------------------------------|--------|--|--------------------|
| Title and abstract | 1 | (a) Indicate the study's design with a commonly used term in the title or the abstract | 2 |
| | | (b) Provide in the abstract an informative and balanced summary of what was done and what was found | 2 |
| Introduction | | | |
| Background/rationale | 2 | Explain the scientific background and rationale for the investigation being reported | 4-6 |
| Objectives | 3 | State specific objectives, including any prespecified hypotheses | 5 - 6 |
| Methods | | | |
| Study design | 4 | Present key elements of study design early in the paper | 6 |
| Setting | 5 | Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection | 6 |
| Participants | 6 | (a) Give the eligibility criteria, and the sources and methods of selection of participants | 6-7 |
| Variables | 7 | Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable | 7 |
| Data sources/ measurement | 8* | For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group | Not applicable |
| Bias | 9 | Describe any efforts to address potential sources of bias | Not applicable |
| Study size | 10 | Explain how the study size was arrived at | 6 |
| Quantitative variables | 11 | Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why | 8 |
| Statistical methods | 12 | (a) Describe all statistical methods, including those used to control for confounding | 8 |
| | | (b) Describe any methods used to examine subgroups and interactions | 8 |
| | | (c) Explain how missing data were addressed | 6-8 |
| | | (d) If applicable, describe analytical methods taking account of sampling strategy | 8 |

| | | | |
|--------------------------|-----|--|----------------|
| | | (e) Describe any sensitivity analyses | 8 |
| Results | | | |
| Participants | 13* | (a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed | 6- 9 |
| | | (b) Give reasons for non-participation at each stage | Not applicable |
| | | (c) Consider use of a flow diagram | Not applicable |
| Descriptive data | 14* | (a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders | 9 |
| | | (b) Indicate number of participants with missing data for each variable of interest | Not applicable |
| Outcome data | 15* | Report numbers of outcome events or summary measures | 9-10 |
| Main results | 16 | (a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included | 12-13 |
| | | (b) Report category boundaries when continuous variables were categorized | 7 |
| | | (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period | 11-12 |
| Other analyses | 17 | Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses | 12 |
| Discussion | | | |
| Key results | 18 | Summarise key results with reference to study objectives | 13 |
| Limitations | 19 | Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias | 14-16 |
| Interpretation | 20 | Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence | 3, 14-16 |
| Generalisability | 21 | Discuss the generalisability (external validity) of the study results | 14-16 |
| Other information | | | |
| Funding | 22 | Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based | Not applicable |

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

BMJ Open

Diabetes among adults in Bangladesh: Changes in prevalence and risk factors between two cross sectional surveys

| | |
|---------------------------------|---|
| Journal: | <i>BMJ Open</i> |
| Manuscript ID | bmjopen-2021-055044.R2 |
| Article Type: | Original research |
| Date Submitted by the Author: | 13-Jun-2022 |
| Complete List of Authors: | Chowdhury, Muhammad Abdul Baker; University of Florida, Emergency Medicine Islam, Mirajul; University of Dhaka, Statistics Rahman, Jakia; BRAC University, Mathematics and Natural Sciences Uddin, Md. Jamal; Shahjalal University of Science and Technology, Statistics Rabiul Haque, Md; University of Dhaka, Population Sciences |
| Primary Subject Heading: | Diabetes and endocrinology |
| Secondary Subject Heading: | Diabetes and endocrinology, Epidemiology, Global health |
| Keywords: | Epidemiology < TROPICAL MEDICINE, DIABETES & ENDOCRINOLOGY, General diabetes < DIABETES & ENDOCRINOLOGY, PUBLIC HEALTH |
| | |

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Title

Diabetes among adults in Bangladesh: Changes in prevalence and risk factors between two cross sectional surveys

Short title

Diabetes among adults in Bangladesh

Authors

Muhammad Abdul Baker Chowdhury, MPH, MPS, MS^{1*}; Mirajul Islam, MS^{2,3} Jakia Rahman, BS⁴; Md Jamal Uddin, PhD^{5,7+}; Md Rabiul Haque, PhD⁶⁺

Affiliations

1. Department of Neurosurgery, College of Medicine, University of Florida, Gainesville, FL
2. Statistics Department, University of Florida, Gainesville FL
3. Department of Statistics, University of Dhaka, Dhaka, Bangladesh
4. Department of Mathematics and Natural Sciences, BRAC University, Dhaka, Bangladesh
5. Department of Statistics, Shahjalal University of Science & Technology, Sylhet-3114, Bangladesh
6. Department of Population Sciences, University of Dhaka, Dhaka, Bangladesh
7. Department of General Educational and Development, Daffodil International University, Dhaka, Bangladesh

⁺Senior authors

^{*} Corresponding author

^{*} Corresponding author email: mchow023@fiu.edu

Tables: 2 Tables and 2 Supplemental Tables

Abstract Word Count: 194

Manuscript word count: 3139

Abstract

Objective/ Research Question: To investigate the change in the prevalence and risk factors of diabetes among adults in Bangladesh between 2011 and 2018.

Design: The study used two waves of nationally representative cross-sectional data extracted from the Bangladesh Demographic and Health Surveys in 2011 and 2017-18.

Setting: Bangladesh

Participants: 14,376 adults aged ≥ 35 years.

Primary outcome: Diabetes mellitus (Type-II diabetes).

Results: From 2011 to 2018, the diabetes prevalence among adults aged ≥ 35 years increased from 10.95% (880) to 13.75% (922) ($p < 0.001$), with the largest relative increase (90%) among obese individuals. Multivariable logistic regression analysis identified age and body mass index (BMI) were the key risk factors for diabetes. Adults who were overweight or obese were 1.54 times (AOR: 1.54, 95% CI: 1.20, 1.97) more likely to develop diabetes than normal-weight individuals in 2011, and 1.22 times (AOR: 1.22, 95% CI: 1.00, 1.50) and 1.44 times (AOR: 1.44, 95% CI: 1.13, 1.84) more prone to develop diabetes in 2018. Other significant risk factors for diabetes were marital status, education, geographic region, wealth index, and hypertension status in both survey years.

Conclusion A high prevalence of diabetes was observed and it has been steadily increasing over time. To enhance diabetes detection and prevention among adults in Bangladesh, population-level interventions focusing on health education, including a healthy diet and lifestyle, are required.

Keywords: *Diabetes, Hypertension, trends, prevalence, risk factors, Bangladesh.*

Strengths and limitations of this study

- We estimated the change in prevalence and risk factors of diabetes among Bangladeshi adults between 2011 and 2018.
- Data were obtained from nationally representative two cross-sectional surveys including the most recent one.
- Our study uniquely investigates the risk factors of diabetes among adults
- Anthropometric and diabetes data were collected using the WHO-recommended guidelines.
- Unavailability of information on some important risk factors was the major data limitation.

Introduction

Diabetes mellitus or type 2 diabetes is one of the most common chronic and preventable diseases affecting 463 million individuals worldwide in 2019.^{1,2} By 2045, the International Diabetes Federation (IDF) predicts 700 million people will have diabetes worldwide, a 51% increase from 2019.² This preventable disease is linked to heart disease, stroke, renal failure, and blindness, as well as morbidity, mortality, and poor quality of life.³⁻⁵ Moreover, diabetes causes a huge financial burden on the patient and the healthcare system of the country which is expected to continue to grow. Global health expenditure on diabetes is estimated to reach \$825 billion by 2030 and \$845 billion by 2045 compared to \$760 billion in 2019.^{2,6}

Low- and middle-income countries have seen a faster increase in diabetes prevalence than high-income countries, with more than two-thirds of the population suffering from the disease.⁷⁻¹⁰ Bangladesh, like many other countries, is transitioning from communicable to non-communicable diseases due to improved socio-economic status and unplanned but rapid urbanization.¹¹ Bangladesh is also going through a nutritional transition from traditional eating habits to a fast-food diet and sedentary lifestyle, which is contributing to the rise of non-communicable diseases like diabetes.¹² These reasons may lead Bangladesh to endure increased diabetes prevalence in the future. A systematic review of published studies between 1994 and 2013 found that diabetes prevalence in Bangladesh ranged from 4.5 to 35.0 percent.¹³ Furthermore, the number of diabetic patients in Bangladesh is estimated to be 13.7 million by 2045.¹⁴

Several studies on diabetes conducted in Bangladesh confirmed that diabetes prevalence among adults is rising steadily.^{15,16} People in Bangladesh who live in urban areas, belong to higher-income households, are older, have more education, and have hypertension get a higher prevalence of diabetes.¹⁷⁻²⁴ Systematic review study on the prevalence of diabetes and pre-diabetes in Bangladesh mentioned that the prevalence of diabetes was significantly higher in urban areas compared with rural areas, while there was no significant gender difference.¹⁹ Another cross-sectional study found that longer duration of

1
2 diabetes, use of insulin, and presence of diabetes complications were significantly related to the average
3 annual cost per patient in Bangladesh.¹⁸ People with lower socioeconomic status are less aware as well as
4 spend less on the care of diabetes. However, most of those studies were confined to urban-rural
5 communities or some other specific groups (e.g., slum residents), which did not consider a wide range of
6 correlates of diabetes for the entire country. While an upward trend in the prevalence of diabetes is
7 evident, very few population-based studies also reported the prevalence of diabetes, which are outdated.
8
9 Despite the rising literature on diabetes research in Bangladesh, no study has identified trends in the
10 prevalence of diabetes and its related risk factors or made a comparison of its risk factors over the years.
11
12

13
14
15 In this study, we explored whether the overall prevalence of type 2 diabetes among adults in
16 Bangladesh changed between 2011 and 2018 and to what extent it changed by socioeconomic and
17 demographic characteristics of individuals. We also examined the factors that potentially contribute to the
18 risk of diabetes among the studied population and make comparisons among them. It is important to
19 recognize changes in diabetes prevalence by population subgroups to ensure access to and use of
20 available treatment required for the population living with diabetes.
21
22

23 **Methods**

24 *Data source*

25
26 We used two waves of cross-sectional data from the Bangladesh Demographic and Health Survey
27 (BDHS) from 2011 and 2017-18 to estimate the prevalence of diagnosed diabetes among the
28 noninstitutionalized Bangladeshi population aged 35 and older. Diabetes testing and related
29 questionnaires were included only in the 2011 and 2018 surveys. The BDHS was designed to collect data
30 to monitor and evaluate the population health and nutritional status of the country using two-stage
31 stratified cluster sampling from non-institutionalized households. The details of the sampling procedure
32 and sample selection are published elsewhere.^{25 26}
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The National Institute of Population Research and Training (NIPORT) Ethics Review Board
4 approved the data collection of the BDHS with the requirement of documented consent from all study
5 participants. Our study was exempt from the ethical review approval because we used freely available de-
6 identified data.
7
8
9

10
11 Biomarker measurements including blood pressure and blood glucose were collected only in 2011
12 and 2018 BDHS. A total of 23,541 adults were eligible for blood glucose measurements in both surveys.
13
14 After exclusion of nonresponses and individuals with missing data and pregnant women, 19,584 adults
15 comprised the study population for both survey years. Of the total included study participants 14,376
16
17 (7,556 in 2011 and 6,820 in 2018) were aged 35 years or over.
18
19
20

21 22 *Outcome variable*

23
24
25 The outcome variable for this study was the prevalence of diabetes for both survey years.
26
27 Diabetes status was measured by fasting blood glucose (FBG) values greater than or equal to 7.0 mmol/L
28
29 or self-reported use of blood glucose-lowering medication during the interview.²⁷
30
31

32 *Demographic and other covariates*

33
34
35 Demographic, household and community-level characteristics were included to assess the
36
37 prevalence and risk factors of diabetes by survey years. Individual-level characteristics were participant's
38
39 age (grouped into 35-44, 45-54, 55-64, 65-74, and 75+ years of age), sex, marital status (currently
40
41 married, not currently married), educational level (no education, primary, secondary, higher), body mass
42
43 index (BMI), and hypertension status. The BMI was calculated as weight in kilograms divided by height
44
45 in meters squared. We used BMI classifications for Asian population: underweight (<18.5) normal weight
46
47 (18.5 to 23.0), moderate risk/ overweight (23.0 to < 27.5), high risk/obese (≥ 27.5).²⁸ Household and
48
49 community characteristics were socio-economic status (wealth index), place of residence (urban, rural),
50
51 and geographic region (division).
52
53
54
55
56
57
58
59
60

Statistical analysis

The full sample of each survey was used for descriptive analysis of individuals' demographic and socioeconomic characteristics. Chi-square tests were performed to check the bivariate association between each characteristic and diabetes status. We also used independent proportion tests to know whether the prevalence of diabetes between the two surveys was changed significantly and calculated the relative changes between the survey periods. For the adjusted analysis in each survey year, we performed multivariable logistic regression models to identify the associated risk factors of diabetes in Bangladesh by entering the variables that were significantly associated with the outcome in the univariate logistic regression analysis. Before entering the models, tests for multi-collinearity between explanatory variables were performed. To select the best model, we checked the values of -2Log Likelihood ratio test, AIC, and the area under the receiver operating characteristic (ROC) curve. The best model had lower values of -2Log Likelihood ratio test and lower AIC value.

To make an appropriate comparison between the two homogeneous groups for study periods, adjusted odds ratios (AORs) were calculated for both 2011 and 2018 BDHS data of the study participants aged 35 years and older. Moreover, we performed sensitivity analysis by splitting the datasets into rural and urban as well as males and females. We used $P < 0.05$ at 2-sided statistical significance for all analyses. Data management and statistical analyses were performed using Stata 15 (StataCorp, College Station, TX, USA). We considered the sample weights, primary sampling units, and Strata using the “SVY” command of Stata considering the complex nature of the survey design. Comparisons by different groups were drawn using the “svysubpop” command.

Results

Diabetes prevalence

Table 1 shows the socio-demographic characteristics of participants with age ≥ 35 years in Bangladesh. The highest percentage of individuals came from the age group 35-44 in 2011 (35.77%) and

2018 (36.03%). In both years' male and female percentages are approximately 50%, and a similar decreasing trend in education level was observed. Urban individuals were higher than rural in both years. We observe that the lowest percentages of individuals came from overweight (12.97%) in 2011 but from obese (13.36%) in 2018.

The diabetes prevalence in 2011 and 2018 with relative ratios are presented in Table 2. The overall prevalence of diabetes among adults ages ≥ 35 years increased from 10.95% in 2011 to 13.75% in 2018. The relative highest increase (38%) in diabetes prevalence was found among individuals with age 65-74 and the second-highest increase rate (36%) was found among the age group 45-54. The prevalence rate of diabetes among females increased significantly from 11.25% in 2011 to 13.81% in 2018 whereas this increment among males is not significant. The relative increase in diabetes prevalence over time among married, currently not working individuals and rural areas were 30%, 42%, and 28%, respectively. A significant increase in diabetes prevalence was also observed among adults with no education and secondary education. The highest relative increase (54%) in diabetes prevalence was in the Dhaka region followed by 53% in the Khulna region. Diabetes prevalence among middle, richer, and the richest individuals increased significantly by 48%, 41%, and 33%, respectively. The prevalence of diabetes increased by 90% among obese individuals, this rate decreased by 9% among overweight adults.

Risk factor analysis

The adjusted results from multivariable logistic regression analysis are reported in **Table 3**. The likelihood of diabetes was the highest [AOR: 2.11, 95% CI: 1.58, 2.83] among adults ages 55-64 in 2011 whereas this rate was highest [AOR: 1.67, 95% CI: 1.21, 2.30] in the age group 65-74 in 2018 compared to adults ages 35-44. There was no significant difference in the odds of having diabetes among males and females in both periods. Although marital status was highly insignificant (p-value=0.572) in 2011, this variable was found as marginally insignificant (p-value=0.057) in 2018.

1
2
3 The findings of the study also suggest that adults with primary, secondary, and higher education
4 had 31%, 32%, and 87% higher odds of having diabetes, respectively than adults with no education in
5 2011. However, education was not a significant factor for diabetes among adults in 2018. Place of
6 residence had no significant effect on diabetes in both periods. Compared to the Dhaka division,
7 individuals living in Barisal and Chittagong divisions had a 43% and 44% higher likelihood of having
8 diabetes, respectively in 2011. On the other hand, there exists no significant difference in having diabetes
9 among adults in Barisal, Chittagong, Dhaka, Rajshahi, and Sylhet divisions in 2018. Regarding economic
10 status, only the richest individuals had a significantly higher likelihood (96%) of diabetes in 2011
11 compared to the poorest individuals. However, both the richer and richest adults had more likelihood of
12 diabetes [AOR: 1.84 and 3.09] than the poorest adults in 2018.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Higher BMI was a significant factor in both 2011 and 2018. For example, overweight and obese
adults compared to normal-weight adults had 54% and 51% more likelihood of diabetes, respectively in
2011, and 22% and 44% higher likelihood of diabetes in 2018. Moreover, the odds of having diabetes
among working adults in 2018 was lower [AOR: 0.80, 95% CI: (0.65, 0.99)] than in non-working adults.
Since there exists a strong relationship between diabetes and hypertension, individuals having
hypertension had 51% and 57% more likelihood of diabetes, respectively in 2011 and 2018 compared to
individuals without hypertension.

Subgroup analysis

Subgroup analysis of diabetes by sex and place of residence were also performed and the results
of this analysis are presented in **Table S1** and **Table S2**. Male with higher education had 99% higher risk
of diabetes compared to male with no education in 2011 whereas no significant difference in the risk of
diabetes among them was observed in 2018. However, the odds of having diabetes between a male with
primary education and no education was not significantly different [p=0.236] in 2011. The opposite
scenario [AOR: 1.48 [(1.06, 2.06), p=0.022] was observed in 2018. Urban males had less likelihood

1
2 [AOR: 0.71, 95% CI: 0.55, 0.93] of diabetes than rural males in 2018 whereas there was no difference in
3
4 the likelihood of diabetes among them in 2011.
5
6

7 Marital status was found to be a significant factor ($p=0.043$) of diabetes among rural adults in
8
9 2018 only. Richer individuals had no significant difference ($p>0.05$) in having diabetes compared to the
10
11 poorest individuals except in rural areas in 2018 ($p=0.019$). Currently working individuals had 31%
12
13 (AOR: 0.69, 95% CI: 0.49, 0.99) and 27% (AOR: 0.73, 95% CI: 0.57, 0.95) lower odds of having
14
15 diabetes than non-working rural individuals, in 2011 and 2018, respectively. However, there was no
16
17 significant difference in diabetes among working and non-working urban individuals for both periods.
18
19

20 21 Discussion

22
23 We systematically analyze the prevalence and risk factors of diabetes among the adult population
24
25 (aged ≥ 35 years) in Bangladesh using two waves of nationally representative survey data (2011 and
26
27 2018). The purpose of this study was to compare the prevalence of diabetes among people aged >35 years
28
29 in 2018 with 2011 in Bangladesh. We found several remarkable findings linked to diabetes and its risk
30
31 factors. One of the important findings was the identification of upward trends in the overall prevalence of
32
33 diabetes and its distribution as per individual characteristics. The prevalence of diabetes increased by 26%
34
35 between 2011 and 2018.. During this period, the prevalence increased significantly not only among the
36
37 overall population but also among different age groups, both sexes, married individuals, uneducated and
38
39 secondary completed, rural areas, middle to richest wealth index group, obese individuals, unemployed,
40
41 and hypertensive patients. This finding is consistent with the reported prevalence of diabetes among the
42
43 adult populations over the years.²⁹⁻³¹
44
45
46

47 We found that the likelihood of diabetes increased with an increase of age. The odds of having
48
49 diabetes was higher for older age individuals compared to younger adults. Similar to the previous studies
50
51 in Bangladesh^{32 33} and other developing countries,^{34 35} the odds of having diabetes increased consistently
52
53 for all age groups of older adults (≥ 35 years) in both data sets. In the future, the upward trend of having
54
55

1
2
3 diabetes is likely to be a major public health concern in Bangladesh owing to changes in population age
4 structure with lower fertility rate, steady socioeconomic growth, and increased life expectancy. This
5 process will sharply lead to an increase in the number of middle and older age population and diabetes
6 prevalence in Bangladesh.³⁶ The prevalence of diabetes among the working-age population may be a
7 concern because of the complex effects of diabetes on co-morbidity and economic growth in Bangladesh,
8 where about 12% of total households pay for diabetic care by selling household assets or borrowing
9 money.³⁷⁻³⁹

10
11 We found a significantly higher relative increase in diabetes prevalence in rural areas compared
12 to urban areas, indicating that diabetes is no longer confined to urban areas in Bangladesh because, in
13 recent years, rural residents have adopted an unhealthy lifestyle, consuming fast food with more carbs.
14 Moreover, they are also less aware of the disease and seek medical attention at a later stage. Hira et al⁴⁰
15 found similar results in a study conducted in Bangladesh and Tripathy et al.⁴¹ found a similar trend in our
16 neighboring country India.

17
18 Our study also identified the greater prevalence of having diabetes among married people
19 compared to unmarried people. This finding was significant in 2018 but was not found to be significant in
20 2011.³⁶ This result was broadly consistent with previous studies in which the presence of diabetes was
21 also associated with greater marital stability and satisfaction.^{32 41 42} Furthermore, this could be a proxy for
22 age because married people are older than unmarried people, and married women who have children
23 sometimes develop diabetes at a younger age.⁴³ Moreover, male being married was also associated with a
24 higher risk of hypertension and type-2 diabetes.⁴⁴

25
26 Although higher education and socio-economic status are negatively associated with diabetes in
27 developed countries, we have found the opposite results in Bangladesh for both survey periods. The
28 richest 35-year-olds are three times more likely to have diabetes than the poorest wealth group. These
29 findings are in line with the previous studies conducted in Asian and other developing countries.^{32 45 46}
30 The greater likelihoods of diabetes among people with no education and secondary education are likely to

1
2
3 be associated with their less awareness about lifestyle, and may not consider it as a threat to their
4 health.⁴⁷⁻⁵⁰ The prevalence of diabetes varies also by region among individuals in Bangladesh. For
5 example, people living in Rangpur and Khulna regions have experienced a significantly lower risk of
6 having diabetes than those from Dhaka and other regions. People in these two divisions have a lower
7 socioeconomic status than those in Dhaka, which may be linked to a lower risk of diabetes.^{32 46} The rapid
8 increase in the prevalence of diabetes in all regions, particularly in Dhaka and Khulna regions, in which it
9 has increased by 54% and 53% respectively between 2011 and 2018, is plausibly associated with the
10 rapid growth of urbanization and its consequences on healthy lifestyles.^{9 51} High BMI and hypertension
11 are important factors of diabetes reported in most of the previous studies.^{8 11 32 37 39 52 53} Our study findings
12 also pointed out that with higher BMI have a greater likelihood of having diabetes compared to normal-
13 weight adults in both 2011 and 2018.⁵⁴⁻⁵⁷

24
25
26 One of the major strengths of our study is the use of nationally representative cross-sectional
27 survey data of the two waves, including the most recent one which is released in December 2020. Data of
28 these surveys related to anthropometrics and diabetes were not self-reported but rather collected by
29 trained and experienced health workers such as nurses, midwives, and health assistants using the WHO-
30 recommended guidelines. To our knowledge, this study for the first time estimated the national diabetes
31 prevalence and its risk factors among adults in Bangladesh. Another important strength is that we
32 compared changes in estimates of diabetes predictors between 2011 and 2018 surveys along with
33 subgroups analyses, by sex, and by place of residence. Despite having some strengths, our study is not
34 beyond the limitations. Due to the lack of diabetes data for younger adults (<35 years) in previous
35 surveys, we could not compare their current diabetes prevalence. Moreover, the unavailability of data on
36 some important correlates like the types of diet, intake of fast food, intake of calories, physical exercise
37 including the nature of work, family history of diabetes, and cholesterol level of diabetes was the major
38 limitation of these data sets. As a cross-sectional survey, blood sugar level was measured for one day only

1
2 and thus we do not have follow-up and or longitudinal data on diabetes and its correlates. Due to the
3
4 nature of survey data, this study identified the risk factors of diabetes only rather than causality.
5
6

7
8 The study shows that among Bangladeshi adults, there is a high prevalence of diabetes and it is
9
10 escalating over time. The study also reports a significant portion of younger adults with diabetes. Age and
11
12 overweight/obesity are the two most important risk factors for diabetes for all adults, irrespective of sex,
13
14 residence, educational attainment, and wealth index. There is evidence of an increase in the magnitude of
15
16 diabetes over time and in the younger population; indeed, age increases the chances of developing
17
18 diabetes mellitus significantly. These findings, together with an increase in the prevalence of type 2
19
20 diabetes among Bangladeshi adults, underscore the need for primary (awareness campaign about the
21
22 adversity of fast food habit and impotence of physical mobility/work/exercise) and secondary (incorporate
23
24 the issue text on the causes and consequences of diabetes and NCDs in school level curriculum)
25
26 prevention efforts tailored to age-specific populations.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

List of Abbreviations

| | |
|------|--|
| AOR | Adjusted Odds Ratio |
| BMI | Body mass index |
| CI | Confidence interval |
| CVD | Cardiovascular diseases |
| WHO | World Health Organization |
| BDHS | Bangladesh Demographic and Health Survey |

Availability of data and material: All data presented here in the manuscript is freely available at dhsprogram.com.

Prior Publication: This data has not been published previously and is not under consideration elsewhere.

Author Contribution: M.A.B. Chowdhury conceptualized the study, designed the analytic approach, managed and performed the analysis, interpreted the results, and drafted the manuscript. M. Islam: helped with the analysis, interpreted the results, and drafted the manuscript. J. Rahman, M.J. Uddin reviewed, edited, and updated the manuscript, and M.R.Haque reviewed, edited, and supervised the study.

Disclosures/Conflict of Interest: There are no potential conflicts (financial, professional, or personal) to disclose by any of the authors.

Funding: None

Acknowledgment: We thank DHS program for providing the data.

Ethical approval and consent

All BDHS surveys received ethical approval from ICF Macro Institutional Review Board, Maryland, USA, and the National Research Ethics Committee of Bangladesh Medical Research Council (BMRC),

1
2 Dhaka, Bangladesh. Informed consent was obtained from each participant of the survey before enrolling
3 in the survey by using the Introduction and Consent form of the survey. It was also explained that the
4 information will be kept strictly confidential and will not be shared with anyone except members of the
5 survey team.
6
7
8

9 **Patient and Public Involvement**

10
11 The BDHS questionnaires were based on the MEASURE DHS model questionnaires. These model
12 questionnaires were adapted for use in Bangladesh during a series of meetings with a technical working
13 group (TWG) that consisted of representatives from NIPORT, Mitra and Associates, International Centre
14 for Diarrheal Diseases and Control, Bangladesh (icddr, b), USAID/Bangladesh, and MEASURE DHS.
15 Patients were not directly involved in the study however; the TWG involved representatives from the
16 government, non-government, ministry of health and family welfare representatives and donor
17 organizations were involved in the study design and questionnaire development. The results will be used
18 by the health researchers and policymakers of the country.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE 1. Socio-demographic characteristics of the study sample by survey year, Bangladesh Demographic and Health Surveys.

| Variables | 2011, n (%) | 2017-18, n (%) |
|---------------------------|--------------|----------------|
| Age group | | |
| 35-44 | 2703 (35.77) | 2457 (36.03) |
| 45-54 | 2236 (29.59) | 1721 (25.23) |
| 55-64 | 1292 (17.1) | 1394 (20.44) |
| 65-74 | 809 (10.71) | 809 (11.86) |
| 75+ | 516 (6.83) | 439 (6.44) |
| Sex | | |
| Female | 3823 (50.6) | 3510 (51.47) |
| Male | 3733 (49.4) | 3310 (48.53) |
| Marital status | | |
| Not married | 1214 (16.07) | 1140 (16.72) |
| Married | 6342 (83.93) | 5680 (83.28) |
| Wealth index | | |
| Poorest | 1346 (17.81) | 1380 (20.23) |
| Poorer | 1351 (17.88) | 1328 (19.47) |
| Middle | 1463 (19.36) | 1353 (19.84) |
| Richer | 1584 (20.96) | 1278 (18.74) |
| Richest | 1812 (23.98) | 1481 (21.72) |
| Educational level | | |
| No education | 3424 (45.31) | 2663 (39.07) |
| Primary | 2082 (27.55) | 2175 (31.91) |
| Secondary | 1403 (18.57) | 1314 (19.28) |
| Higher | 647 (8.56) | 664 (9.74) |
| Place of residence | | |
| Urban | 5071 (67.11) | 4514 (66.19) |
| Rural | 2485 (32.89) | 2306 (33.81) |

TABLE 1. Socio-demographic characteristics of the study sample by survey year, Bangladesh Demographic and Health Surveys.

| Variables | 2011, n (%) | 2017-18, n (%) |
|-------------------|--------------|----------------|
| Geographic region | | |
| Dhaka | 1316 (17.42) | 1608 (23.58) |
| Barisal | 866 (11.46) | 748 (10.97) |
| Chittagong | 1116 (14.77) | 865 (12.68) |
| Khulna | 1205 (15.95) | 1034 (15.16) |
| Rajshahi | 1067 (14.12) | 890 (13.05) |
| Rangpur | 1066 (14.11) | 924 (13.55) |
| Sylhet | 920 (12.18) | 751 (11.01) |
| Body Mass Index | | |
| Underweight | 1594 (21.1) | 1188 (17.42) |
| Normal weight | 2424 (32.08) | 2711 (39.75) |
| Overweight | 980 (12.97) | 2010 (29.47) |
| Obese | 2558 (33.85) | 911 (13.36) |
| Currently working | | |
| No | 3901 (51.64) | 2367 (34.71) |
| Yes | 3653 (48.36) | 4453 (65.29) |
| Hypertension | | |
| No | 5534 (73.24) | 4036 (59.18) |
| Yes | 2022 (26.76) | 2784 (40.82) |

TABLE 2. Prevalence of diabetes among adults age 35 years and older in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|--------------------------------------|------------------------------|----------------------------------|----------------------------------|-----------------------------|-------|
| All adults age 35 years and older | 100% | 10.95 (0.0048) | 13.75 (0.0056) | <0.001 | 1.26 |
| Age group | | | | | |
| 35-44 | 35.99 | 8.82 (0.0065) | 11.21 (0.0078) | 0.081 | 1.27 |
| 45-54 | 27.43 | 10.86 (0.0079) | 14.82 (0.01) | 0.008 | 1.36 |
| 55-64 | 18.65 | 14.69 (0.012) | 15.78 (0.0115) | 0.6286 | 1.07 |
| 65-74 | 11.26 | 11.60 (0.0128) | 15.97 (0.016) | 0.1123 | 1.38 |
| 75+ | 6.68 | 12.11 (0.0174) | 13.43 (0.0189) | 0.793 | 1.11 |
| Sex | | | | | |
| Male | 48.98 | 10.65 (0.0061) | 13.69 (0.0072) | 0.091 | 1.29 |
| Female | 51.02 | 11.25 (0.006) | 13.81 (0.0072) | 0.003 | 1.23 |
| Marital status | | | | | |
| Not married | 15.88 | 12.13 (0.0112) | 12.54 (0.0111) | 0.953 | 1.03 |
| Married | 84.12 | 10.73 (0.0049) | 13.98 (0.006) | <0.001 | 1.3 |
| Educational level | | | | | |
| No education | 44.8 | 8.39 (0.0058) | 10.64 (0.0072) | 0.021 | 1.27 |
| Primary | 29.07 | 11.10 (0.0078) | 13.59 (0.009) | 0.141 | 1.22 |
| Secondary | 18.15 | 13.03 (0.0108) | 17.8 (0.0121) | 0.015 | 1.37 |
| Higher | 7.98 | 21.79 (0.018) | 20.27 (0.0185) | 0.209 | 0.93 |
| Place of residence | | | | | |
| Urban | 23.91 | 16.08 (0.0106) | 18.95 (0.0106) | 0.129 | 1.18 |
| Rural | 76.09 | 9.39 (0.0051) | 12.06 (0.0065) | 0.003 | 1.28 |
| Geographic region | | | | | |
| Barisal | 5.79 | 12.54 (0.0117) | 12.09 (0.0161) | 0.975 | 0.96 |
| Chittagong | 16.45 | 14.28 (0.0125) | 17.33 (0.0169) | 0.087 | 1.21 |
| Dhaka | 38.99 | 11.26 (0.0107) | 17.32 (0.0118) | 0.002 | 1.54 |

TABLE 2. Prevalence of diabetes among adults age 35 years and older in Bangladesh, 2011-2018

| Variables | Distribution 2011-2018, % | Diabetes 2011 BDHS, % (SE) | Diabetes 2018 BDHS, % (SE) | p-value 2011 vs. 2018 | Ratio |
|-------------------|------------------------------|----------------------------------|----------------------------------|-----------------------------|-------|
| Khulna | 13.45 | 7.30 (0.007) | 11.14 (0.0114) | 0.002 | 1.53 |
| Rajshahi | 13.15 | 10.73 (0.01) | 11.32 (0.0134) | 0.664 | 1.05 |
| Rangpur | 9.26 | 8.59 (0.0111) | 7.93 (0.0097) | 0.667 | 0.92 |
| Sylhet | 2.91 | 11.85 (0.011) | 12.71 (0.0169) | 0.998 | 1.07 |
| Wealth index | | | | | |
| Poorest | 19.91 | 7.28 (0.0088) | 7.26 (0.0091) | 0.69 | 1.00 |
| Poorer | 19.74 | 7.35 (0.0084) | 7.70 (0.0085) | 0.891 | 1.05 |
| Middle | 20.19 | 7.56 (0.0075) | 11.17 (0.01) | 0.005 | 1.48 |
| Richer | 19.70 | 11.33 (0.0098) | 16.01 (0.0128) | <0.001 | 1.41 |
| Richest | 20.46 | 20.49 (0.0122) | 27.18 (0.0134) | 0.006 | 1.33 |
| Body Mass Index | | | | | |
| Underweight | 19.87 | 7.27 (0.0081) | 7.79 (0.0087) | 0.437 | 1.07 |
| Normal weight | 36.10 | 9.59 (0.0071) | 11.21 (0.0074) | 0.343 | 1.17 |
| Overweight | 20.14 | 18.57 (0.0149) | 16.95 (0.0095) | 0.045 | 0.91 |
| Obese | 23.88 | 11.90 (0.0078) | 22.56 (0.0156) | <0.001 | 1.9 |
| Currently working | | | | | |
| No | 43.63 | 12.11 (0.0064) | 17.16 (0.0089) | <0.001 | 1.42 |
| Yes | 56.37 | 9.70 (0.0061) | 12.00 (0.006) | 0.10 | 1.24 |
| Hypertension | | | | | |
| No | 67.45 | 9.06 (0.0048) | 10.55 (0.0063) | 0.522 | 1.16 |
| Yes | 32.55 | 16.33 (0.0105) | 18.57 (0.009) | 0.178 | 1.14 |

TABLE 3. Adjusted odds ratios for factors predicting diabetes among adults 35 years and older BDHS 2011-2018.

| Variables | 2011 BDHS | | 2017-18 BDHS | |
|---------------------------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | |
| 35-44 | Ref | | Ref | |
| 45-54 | 1.36 (1.07, 1.72) | 0.012 | 1.41 (1.13, 1.76) | 0.002 |
| 55-64 | 2.11 (1.58, 2.83) | <0.001 | 1.58 (1.24, 2.03) | <0.001 |
| 65-74 | 1.60 (1.13, 2.27) | 0.008 | 1.67 (1.21, 2.30) | 0.002 |
| 75+ | 1.77 (1.14, 2.74) | 0.011 | 1.32 (0.86, 2.01) | 0.202 |
| Sex | | | | |
| Male | Ref | | Ref | |
| Female | 0.78 (0.58, 1.05) | 0.104 | 0.92 (0.75, 1.13) | 0.435 |
| Marital status | | | | |
| Not married | Ref | | | |
| Married | 0.93 (0.72, 1.20) | 0.572 | 1.28 (0.99, 1.64) | 0.057 |
| Educational level | | | | |
| No education | Ref | | Ref | |
| Primary | 1.31 (1.05, 1.64) | 0.017 | 1.23 (0.99, 1.53) | 0.059 |
| Secondary | 1.32 (1.01, 1.73) | 0.045 | 1.23 (0.96, 1.57) | 0.108 |
| Higher | 1.87 (1.35, 2.60) | <0.001 | 1.15 (0.83, 1.60) | 0.397 |
| Place of residence | | | | |
| Urban | 1.08 (0.87, 1.33) | 0.7 | 0.96 (0.79, 1.16) | 0.678 |
| Rural | Ref | | Ref | |
| Geographic region | | | | |
| Barisal | 1.43 (1.04, 1.96) | 0.027 | 0.75 (0.54, 1.06) | 0.103 |
| Chittagong | 1.44 (1.09, 1.89) | 0.010 | 0.88 (0.67, 1.16) | 0.369 |
| Dhaka | Ref | | Ref | |
| Khulna | 0.66 (0.50, 0.89) | 0.007 | 0.60 (0.46, 0.79) | <0.001 |

| | | | | | |
|----|-------------------|-------------------|--------|-------------------|--------|
| 1 | | | | | |
| 2 | | | | | |
| 3 | Rajshahi | 1.17 (0.87, 1.57) | 0.309 | 0.74 (0.55, 1.00) | 0.054 |
| 4 | Rangpur | 1.00 (0.69, 1.45) | 0.986 | 0.56 (0.41, 0.76) | <0.001 |
| 5 | | | | | |
| 6 | Sylhet | 1.22 (0.91, 1.63) | 0.183 | 0.75 (0.53, 1.06) | 0.101 |
| 7 | | | | | |
| 8 | Wealth index | | | | |
| 9 | | | | | |
| 10 | Poorest | Ref | | Ref | |
| 11 | | | | | |
| 12 | Poorer | 0.89 (0.62, 1.27) | 0.514 | 0.99 (0.70, 1.40) | 0.957 |
| 13 | | | | | |
| 14 | Middle | 0.86 (0.62, 1.19) | 0.365 | 1.33 (0.94, 1.88) | 0.103 |
| 15 | | | | | |
| 16 | Richer | 1.18 (0.84, 1.65) | 0.345 | 1.84 (1.29, 2.61) | 0.001 |
| 17 | | | | | |
| 18 | Richest | 1.96 (1.40, 2.76) | <0.001 | 3.09 (2.18, 4.38) | <0.001 |
| 19 | | | | | |
| 20 | Body Mass Index | | | | |
| 21 | | | | | |
| 22 | Underweight | 0.82 (0.61, 1.10) | 0.177 | 0.77 (0.58, 1.01) | 0.055 |
| 23 | | | | | |
| 24 | Normal weight | Ref | | Ref | |
| 25 | | | | | |
| 26 | Overweight | 1.54 (1.20, 1.97) | 0.001 | 1.22 (1.00, 1.50) | 0.052 |
| 27 | | | | | |
| 28 | Obese | 1.51 (1.16, 1.97) | 0.003 | 1.44 (1.13, 1.84) | 0.003 |
| 29 | | | | | |
| 30 | Currently working | | | | |
| 31 | | | | | |
| 32 | No | Ref | | Ref | |
| 33 | | | | | |
| 34 | Yes | 0.77 (0.59, 1.01) | 0.056 | 0.80 (0.65, 0.99) | 0.039 |
| 35 | | | | | |
| 36 | Hypertension | | | | |
| 37 | | | | | |
| 38 | No | Ref | | Ref | |
| 39 | | | | | |
| 40 | Yes | 1.51 (1.26, 1.81) | <0.001 | 1.57 (1.32, 1.87) | <0.001 |

References

1. Ogurtsova K, da Rocha Fernandes JD, Huang Y, et al. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes research and clinical practice* 2017;128:40-50.
2. Federation ID. IDF Diabetes Atlas Ninth. *Dunia: IDF* 2019
3. Kumar SS, Houlden RL. Ethnocultural diversity and the diabetes epidemic in Canada: a call to action. *Can J Diab* 2005;29(2):84-5.
4. Oldroyd J, Banerjee M, Heald A, et al. Diabetes and ethnic minorities. *Postgraduate medical journal* 2005;81(958):486-90.
5. World Health O. World health statistics 2013: a wealth of information on global public health: World Health Organization, 2013.
6. Williams R, Karuranga S, Malanda B, et al. Global and regional estimates and projections of diabetes-related health expenditure: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract* 2020;162:108072. doi: 10.1016/j.diabres.2020.108072 [published Online First: 2020/02/18]
7. Dagenais GR, Gerstein HC, Zhang X, et al. Variations in diabetes prevalence in low-, middle-, and high-income countries: results from the prospective urban and rural epidemiological study. *Diabetes care* 2016;39(5):780-87.
8. Mkuu R, Barry A, Yonga G, et al. Prevalence and factors associated with overweight and obesity in Kenya. *Preventive Medicine Reports* 2021;22:101340. doi: <https://doi.org/10.1016/j.pmedr.2021.101340>
9. Mkuu RS, Gilreath TD, Barry AE, et al. Identifying individuals with multiple non-communicable disease risk factors in Kenya: a latent class analysis. *Public Health* 2021;198:180-86. doi: <https://doi.org/10.1016/j.puhe.2021.07.031>
10. Chowdhury MAB, Epnere K, Haque MA, et al. Urban rural differences in prevalence and risk factors of self-reported hypertension among Kenyan women: a population-based study. *Journal of Human Hypertension* 2021;35(10):912-20. doi: 10.1038/s41371-020-00435-x
11. Chowdhury MAB, Islam M, Rahman J, et al. Changes in prevalence and risk factors of hypertension among adults in Bangladesh: An analysis of two waves of nationally representative surveys. *PLOS ONE* 2021;16(12):e0259507. doi: 10.1371/journal.pone.0259507
12. Amuna P, Zotor FB. Epidemiological and nutrition transition in developing countries: impact on human health and development. *Proc Nutr Soc* 2008;67(1):82-90. doi: 10.1017/s0029665108006058 [published Online First: 2008/02/01]
13. Biswas T, Islam A, Rawal LB, et al. Increasing prevalence of diabetes in Bangladesh: a scoping review. *Public Health* 2016;138:4-11. doi: 10.1016/j.puhe.2016.03.025 [published Online First: 2016/05/14]
14. Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes research and clinical practice* 2018;138:271-81.
15. Rahim MA, Hussain A, Khan AKA, et al. Rising prevalence of type 2 diabetes in rural Bangladesh: a population based study. *Diabetes research and clinical practice* 2007;77(2):300-05.
16. Bhowmik B, Afsana F, My Diep L, et al. Increasing prevalence of type 2 diabetes in a rural Bangladeshi population: a population based study for 10 years. *Diabetes & metabolism journal* 2013;37(1):46-53.

17. Alam DS, Talukder SH, Chowdhury MAH, et al. Overweight and abdominal obesity as determinants of undiagnosed diabetes and pre-diabetes in Bangladesh. *BMC Obesity* 2016;3(1):19. doi: 10.1186/s40608-016-0099-z
18. Afroz A, Alam K, Ali L, et al. Type 2 diabetes mellitus in Bangladesh: a prevalence based cost-of-illness study. *BMC Health Services Research* 2019;19(1):601. doi: 10.1186/s12913-019-4440-3
19. Akhtar S, Nasir JA, Sarwar A, et al. Prevalence of diabetes and pre-diabetes in Bangladesh: a systematic review and meta-analysis. *BMJ open* 2020;10(9):e036086. doi: 10.1136/bmjopen-2019-036086
20. Haque M, Islam S, Kamal ZM, et al. Ongoing efforts to improve the management of patients with diabetes in Bangladesh and the implications. *Hospital Practice* 2021;49(4):266-72. doi: 10.1080/21548331.2021.1906083
21. Islam SMS, Niessen LW, Seissler J, et al. Diabetes knowledge and glycemic control among patients with type 2 diabetes in Bangladesh. *SpringerPlus* 2015;4(1):284. doi: 10.1186/s40064-015-1103-7
22. Rahman MS, Akter S, Abe SK, et al. Awareness, treatment, and control of diabetes in Bangladesh: a nationwide population-based study. *PloS one* 2015;10(2):e0118365-e65. doi: 10.1371/journal.pone.0118365
23. Shariful Islam SM, Lechner A, Ferrari U, et al. Healthcare use and expenditure for diabetes in Bangladesh. *BMJ Global Health* 2017;2(1):e000033. doi: 10.1136/bmjgh-2016-000033
24. Talukder A, Hossain MZ. Prevalence of Diabetes Mellitus and Its Associated Factors in Bangladesh: Application of Two-level Logistic Regression Model. *Scientific Reports* 2020;10(1):10237. doi: 10.1038/s41598-020-66084-9
25. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2011, 2011.
26. (NIPORT) NIOPRaT, Associates Ma, International I. Bangladesh Demographic and Health Survey 2014, 2014.
27. World Health O. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia: report of a WHO/IDF consultation. 2006
28. Who EC. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet (London, England)* 2004;363(9403):157.
29. Menke A, Casagrande S, Geiss L, et al. Prevalence of and Trends in Diabetes Among Adults in the United States, 1988-2012. *JAMA* 2015;314(10):1021-29. doi: 10.1001/jama.2015.10029
30. Bullard KM, Saydah SH, Imperatore G, et al. Secular changes in U.S. Prediabetes prevalence defined by hemoglobin A1c and fasting plasma glucose: National Health and Nutrition Examination Surveys, 1999-2010. *Diabetes Care* 2013;36(8):2286-93. doi: 10.2337/dc12-2563 [published Online First: 2013/04/23]
31. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. *The Australasian medical journal* 2014;7(1):45-48. doi: 10.4066/AMJ.2013.1979
32. Chowdhury MAB, Uddin MJ, Khan HMR, et al. Type 2 diabetes and its correlates among adults in Bangladesh: a population based study. *BMC Public Health* 2015;15(1):1070. doi: 10.1186/s12889-015-2413-y
33. Swasey KK, Gupta RD, Nayeem J, et al. Determinants of diabetes in Bangladesh using two approaches: an analysis of the Demographic and Health Survey 2011. *Journal of Biosocial Science* 2019;52(4):585-95. doi: 10.1017/S002193201900066X [published Online First: 10/21]
34. Thapa S, Kayastha P, Mishra DK. Assessment of Risk of Type 2 Diabetes Among Adults of Banepa Municipality, Nepal: Community Based Cross-Sectional Study. *International Journal of Travel Medicine and Global Health* 2020;8(1):31-36. doi: 10.34172/ijtmgh.2020.05

- 1
- 2
- 3
- 4 35. Hills AP, Arena R, Khunti K, et al. Epidemiology and determinants of type 2 diabetes in south Asia. *The Lancet Diabetes & Endocrinology* 2018;6(12):966-78. doi: [https://doi.org/10.1016/S2213-8587\(18\)30204-3](https://doi.org/10.1016/S2213-8587(18)30204-3)
- 5
- 6
- 7 36. BBS. Sample Vital Registration System, 2019. Dhaka: Bangladesh Bureau of Statistics (BBS), Ministry of Planning, Government of Bangladesh 2019.
- 8
- 9 37. Afroz A, Hird TR, Zomer E, et al. The impact of diabetes on the productivity and economy of Bangladesh. *BMJ Global Health* 2020;5(6):e002420. doi: 10.1136/bmjgh-2020-002420
- 10
- 11 38. Rahman MM, Gilmour S, Saito E, et al. Self-reported illness and household strategies for coping with health-care payments in Bangladesh. *Bulletin of the World Health Organization* 2013;91(6):449-58. doi: 10.2471/BLT.12.115428
- 12
- 13
- 14 39. Akter S, Rahman MM, Abe SK, et al. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bulletin of the World Health Organization* 2014;92(3):204-13A. doi: 10.2471/BLT.13.128371 [published Online First: 01/10]
- 15
- 16
- 17 40. Hira R, Miah MAW, Akash DH. Prevalence of type 2 diabetes mellitus in rural adults (>_31years) in Bangladesh. *Faridpur Medical College Journal* 2018;13(1):20-23.
- 18
- 19 41. Tripathy JP, Thakur JS, Jeet G, et al. Prevalence and risk factors of diabetes in a large community-based study in North India: results from a STEPS survey in Punjab, India. *Diabetology & Metabolic Syndrome* 2017;9(1):8. doi: 10.1186/s13098-017-0207-3
- 20
- 21 42. Trief PM, Himes CL, Orendorff R, et al. The Marital Relationship and Psychosocial Adaptation and Glycemic Control of Individuals With Diabetes. *Diabetes Care* 2001;24(8):1384-89. doi: 10.2337/diacare.24.8.1384
- 22
- 23 43. Cdcgov. Gestational Diabetes and Pregnancy | CDC: @CDCgov; 2020 [updated 2020-07-14T06:42:06Z. Available from: <https://www.cdc.gov/pregnancy/diabetes-gestational.html>.
- 24
- 25
- 26 44. Ramezankhani A, Azizi F, Hadaegh F. Associations of marital status with diabetes, hypertension, cardiovascular disease and all-cause mortality: A long term follow-up study. *PLoS one* 2019;14(4):e0215593-e93. doi: 10.1371/journal.pone.0215593
- 27
- 28 45. Chande AT, Rowell J, Rishishwar L, et al. Influence of genetic ancestry and socioeconomic status on type 2 diabetes in the diverse Colombian populations of Chocó and Antioquia. *Scientific Reports* 2017;7(1):17127. doi: 10.1038/s41598-017-17380-4
- 29
- 30 46. Xu Z, Yu D, Yin X, et al. Socioeconomic status is associated with global diabetes prevalence. *Oncotarget* 2017;8(27):44434-39. doi: 10.18632/oncotarget.17902
- 31
- 32 47. Nyanzi R, Wamala R, Atuhaire LK. Diabetes and quality of life: a Ugandan perspective. *J Diabetes Res* 2014;2014:402012. doi: 10.1155/2014/402012 [published Online First: 2014/04/12]
- 33
- 34 48. Saydah S, Lochner K. Socioeconomic status and risk of diabetes-related mortality in the U.S. *Public health reports (Washington, DC : 1974)* 2010;125(3):377-88. doi: 10.1177/003335491012500306
- 35
- 36 49. Hasan MM, Tasnim F, Tariqujjaman M, et al. Socioeconomic Inequalities of Undiagnosed Diabetes in a Resource-Poor Setting: Insights from the Cross-Sectional Bangladesh Demographic and Health Survey 2011. *International Journal of Environmental Research and Public Health* 2019;16(1) doi: 10.3390/ijerph16010115
- 37
- 38 50. Rahman MS, Akter S, Abe SK, et al. Awareness, Treatment, and Control of Diabetes in Bangladesh: A Nationwide Population-Based Study. *PLOS ONE* 2015;10(2):e0118365. doi: 10.1371/journal.pone.0118365
- 39
- 40 51. Hossain MBH, Khan MH, Haque MA, et al. Changing Patterns of Urbanization in Bangladesh: An Analysis of Census Data. Population Monograph-2. Dhaka, Bangladesh: Department of Population Sciences, University of Dhaka, and Bangladesh Bureau of Statistics, 2015:1-85.
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 52. Chowdhury MAB, Uddin MJ, Haque MR, et al. Hypertension among adults in Bangladesh: evidence
4 from a national cross-sectional survey. *BMC Cardiovasc Disord* 2016;16(1):22. doi:
5 10.1186/s12872-016-0197-3
6
7 53. Kibria GMA, Crispen R, Chowdhury MAB, et al. Disparities in absolute cardiovascular risk, metabolic
8 syndrome, hypertension, and other risk factors by income within racial/ethnic groups among
9 middle-aged and older US people. *Journal of Human Hypertension* 2021 doi: 10.1038/s41371-
10 021-00513-8
11
12 54. Rahman MA, Halder HR, Yadav UN, et al. Prevalence of and factors associated with hypertension
13 according to JNC 7 and ACC/AHA 2017 guidelines in Bangladesh. *Scientific Reports*
14 2021;11(1):15420. doi: 10.1038/s41598-021-94947-2
15
16 55. Mkuu RS, Epnere K, Chowdhury MAB. Prevalence and Predictors of Overweight and Obesity Among
17 Kenyan Women. *Prev Chronic Dis* 2018;15:E44. doi: 10.5888/pcd15.170401
18
19 56. Chowdhury MAB, Adnan MM, Hassan MZ. Trends, prevalence and risk factors of overweight and
20 obesity among women of reproductive age in Bangladesh: a pooled analysis of five national
21 cross-sectional surveys. *BMJ open* 2018;8(7):e018468. doi: 10.1136/bmjopen-2017-018468
22
23 57. Mohammad M, Chowdhury MAB, Islam MN, et al. Factors associated with body mass index among
24 university students in Bangladesh. *Journal of Public Health* 2019 doi: 10.1007/s10389-019-
25 01124-x
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|---------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.56 (1.1, 2.21) | 0.013 | 1.41 (0.98, 2.03) | 0.065 | 1.18 (0.87, 1.62) | 0.288 | 1.41 (1.06, 1.86) | 0.018 |
| 55-64 | 2.36 (1.6, 3.47) | <0.001 | 1.58 (1.09, 2.29) | 0.015 | 1.84 (1.15, 2.94) | 0.012 | 1.53 (1.09, 2.14) | 0.013 |
| 65-74 | 1.79 (1.14, 2.8) | 0.011 | 1.80 (1.18, 2.75) | 0.007 | 1.35 (0.76, 2.39) | 0.304 | 1.44 (0.9, 2.33) | 0.132 |
| 75+ | 1.52 (0.85, 2.72) | 0.153 | 1.41 (0.79, 2.51) | 0.24 | 1.92 (0.98, 3.76) | 0.059 | 1.11 (0.62, 1.99) | 0.727 |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.66 (0.35, 1.26) | 0.205 | 1.83 (0.96, 3.47) | 0.064 | 1.00 (0.75, 1.34) | 0.992 | 1.17 (0.89, 1.56) | 0.263 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.22 (0.88, 1.68) | 0.236 | 1.48 (1.06, 2.06) | 0.022 | 1.37 (1, 1.86) | 0.047 | 1.12 (0.84, 1.48) | 0.434 |
| Secondary | 1.14 (0.80, 1.63) | 0.47 | 1.15 (0.78, 1.69) | 0.482 | 1.53 (1.05, 2.23) | 0.029 | 1.37 (0.97, 1.92) | 0.071 |
| Higher | 1.99 (1.31, 3.01) | 0.001 | 1.19 (0.76, 1.86) | 0.449 | 1.85 (1.1, 3.11) | 0.02 | 1.98 (0.59, 1.64) | 0.941 |
| Place of residence | | | | | | | | |
| Urban | 1.03 (0.76, 1.4) | 0.838 | 0.71 (0.55, 0.93) | 0.012 | 1.1 (0.83, 1.45) | 0.497 | 1.24 (0.96, 1.61) | 0.097 |
| Rural | Ref | | Ref | | Ref | | Ref | |
| Geographic region | | | | | | | | |
| Barisal | 1.42 (0.91, 2.22) | 0.121 | 0.83 (0.53, 1.30) | 0.415 | 1.44 (0.97, 2.14) | 0.072 | 0.72 (0.46, 1.12) | 0.148 |
| Chittagong | 1.55 (1.05, 2.27) | 0.027 | 0.88 (0.62, 1.26) | 0.484 | 1.36 (0.97, 1.91) | 0.071 | 0.88 (0.61, 1.26) | 0.48 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.75 (0.50, 1.12) | 0.159 | 0.56 (0.39, 0.82) | 0.003 | 0.61 (0.4, 0.92) | 0.02 | 0.65 (0.46, 0.93) | 0.019 |
| Rajshahi | 1.14 (0.73, 1.77) | 0.569 | 0.78 (0.52, 1.2) | 0.259 | 1.21 (0.82, 1.77) | 0.338 | 0.71 (0.46, 1.09) | 0.116 |
| Rangpur | 1.04 (0.64, 1.68) | 0.885 | 0.60 (0.39, 0.92) | 0.019 | 0.99 (0.63, 1.57) | 0.975 | 0.53 (0.34, 0.82) | 0.004 |
| Sylhet | 1.41 (0.92, 2.15) | 0.112 | 0.66 (0.41, 1.07) | 0.091 | 1.11 (0.77, 1.6) | 0.58 | 0.82 (0.54, 1.22) | 0.325 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |
| Poorer | 0.81 (0.51, 1.30) | 0.389 | 1.20 (0.72, 2.00) | 0.485 | 0.96 (0.58, 1.59) | 0.88 | 0.87 (0.58, 1.32) | 0.522 |
| Middle | 0.75 (0.46, 1.21) | 0.239 | 1.90 (1.16, 3.12) | 0.011 | 0.99 (0.63, 1.57) | 0.981 | 1.02 (0.66, 1.57) | 0.945 |

TABLE S1. Adjusted odds ratios predicting diabetes and adults age 35 and older by gender and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Male | | 2018 BDHS-Male | | 2011 BDHS-Female | | 2018 BDHS-Female | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Richer | 0.95 (0.59, 1.51) | 0.815 | 2.58 (1.53, 4.34) | <0.001 | 1.43 (0.9, 2.25) | 0.129 | 1.44 (0.94, 2.21) | 0.093 |
| Richest | 1.54 (0.94, 2.53) | 0.088 | 5.47 (3.25, 9.19) | <0.001 | 2.38 (1.49, 3.79) | <0.001 | 2.03 (1.33, 3.09) | 0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.86 (0.6, 1.22) | 0.382 | 0.76 (0.51, 1.13) | 0.177 | 0.69 (0.44, 1.08) | 0.101 | 0.78 (0.52, 1.18) | 0.238 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 1.54 (1.13, 2.12) | 0.007 | 1.25 (0.94, 1.66) | 0.122 | 1.63 (1.09, 2.42) | 0.016 | 1.15 (0.86, 1.52) | 0.343 |
| Obese | 1.71 (1.1, 2.68) | 0.018 | 1.82 (1.22, 2.69) | 0.003 | 1.31 (0.86, 1.98) | 0.207 | 1.29 (0.95, 1.76) | 0.105 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.71 (0.48, 1.06) | 0.09 | 0.80 (0.54, 1.17) | 0.249 | 0.84 (0.55, 1.26) | 0.394 | 0.79 (0.61, 1.03) | 0.076 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.17 (0.88, 1.56) | 0.269 | 1.42 (1.1, 1.83) | 0.007 | 1.77 (1.4, 2.24) | <0.001 | 1.70 (1.33, 2.18) | <0.001 |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Age groups | | | | | | | | |
| 35-44 | Ref | | Ref | | Ref | | Ref | |
| 45-54 | 1.58 (1.07, 2.36) | 0.023 | 1.19 (0.85, 1.65) | 0.312 | 1.22 (1.66, 0.9) | 0.194 | 1.59 (1.18, 2.13) | 0.002 |
| 55-64 | 3.57 (2.16, 5.9) | <0.001 | 1.50 (0.98, 2.29) | 0.059 | 1.60 (2.3, 1.11) | 0.011 | 1.67 (1.24, 2.27) | 0.001 |
| 65-74 | 2.25 (1.24, 4.1) | 0.008 | 1.50 (0.84, 2.66) | 0.168 | 1.32 (2.04, 0.86) | 0.200 | 1.75 (1.19, 2.57) | 0.004 |
| 75+ | 2.54 (1.17, 5.5) | 0.018 | 1.89 (1.01, 3.51) | 0.045 | 1.45 (2.45, 0.85) | 0.171 | 1.14 (0.65, 2.01) | 0.645 |
| Sex | | | | | | | | |
| Female | 0.82 (0.49, 1.38) | 0.461 | 1.29 (0.9, 1.85) | 0.173 | 0.76 (1.11, 0.52) | 0.15 | 0.88 (0.63, 1.03) | 0.087 |
| Male | Ref | | Ref | | Ref | | Ref | |
| Marital status | | | | | | | | |
| Not married | Ref | | | | Ref | | | |
| Married | 0.8 (0.52, 1.24) | 0.322 | 1.07 (0.7, 1.62) | 0.759 | 1.01 (1.38, 0.74) | 0.962 | 1.39 (1.01, 1.92) | 0.043 |
| Educational level | | | | | | | | |
| No education | Ref | | Ref | | Ref | | Ref | |
| Primary | 1.44 (0.96, 2.14) | 0.076 | 1 (0.69, 1.45) | 0.992 | 1.26 (1.64, 0.97) | 0.086 | 1.30 (1.00, 1.69) | 0.048 |
| Secondary | 1.46 (0.94, 2.27) | 0.089 | 1.35 (0.89, 2.03) | 0.155 | 1.22 (1.72, 0.86) | 0.272 | 1.15 (0.83, 1.59) | 0.394 |
| Higher | 1.9 (1.13, 3.19) | 0.015 | 1.19 (0.74, 1.93) | 0.465 | 1.85 (2.94, 1.16) | 0.009 | 1.15 (0.74, 1.81) | 0.532 |
| Geographic region | | | | | | | | |
| Barisal | 0.93 (0.58, 1.49) | 0.76 | 0.99 (0.54, 1.83) | 0.982 | 1.63 (2.45, 1.09) | 0.018 | 0.70 (0.46, 1.06) | 0.088 |
| Chittagong | 1.04 (0.69, 1.58) | 0.845 | 0.58 (0.4, 0.84) | 0.004 | 1.66 (2.41, 1.15) | 0.007 | 1.04 (0.72, 1.5) | 0.841 |
| Dhaka | Ref | | Ref | | Ref | | Ref | |
| Khulna | 0.82 (0.52, 1.28) | 0.381 | 0.70 (0.5, 1.00) | 0.05 | 0.61 (0.91, 0.4) | 0.017 | 0.57 (0.4, 0.81) | 0.002 |
| Rajshahi | 0.94 (0.60, 1.48) | 0.802 | 0.74 (0.51, 1.06) | 0.095 | 1.28 (1.9, 0.86) | 0.228 | 0.75 (0.51, 1.12) | 0.165 |
| Rangpur | 0.77 (0.46, 1.29) | 0.318 | 0.5 (0.33, 0.78) | 0.002 | 1.07 (1.71, 0.67) | 0.775 | 0.58 (0.4, 0.84) | 0.005 |
| Sylhet | 1.43 (0.89, 2.31) | 0.143 | 0.71 (0.47, 1.09) | 0.115 | 1.22 (1.79, 0.83) | 0.318 | 0.77 (0.5, 1.18) | 0.226 |
| Wealth index | | | | | | | | |
| Poorest | Ref | | Ref | | Ref | | Ref | |

TABLE S2. Adjusted odds ratios predicting diabetes and adults age 35 and older by rural-urban and survey year in Bangladesh, BDHS 2011-2018.

| Variables | 2011 BDHS-Urban | | 2018 BDHS-Urban | | 2011 BDHS-Rural | | 2018 BDHS-Rural | |
|--------------------------|-------------------|---------|-------------------|---------|-------------------|---------|-------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| Poorer | 1.23 (0.49, 3.07) | 0.653 | 0.87 (0.36, 2.08) | 0.754 | 0.87 (1.27, 0.6) | 0.468 | 0.98 (0.67, 1.42) | 0.896 |
| Middle | 1.05 (0.49, 2.24) | 0.900 | 0.67 (0.3, 1.51) | 0.335 | 0.84 (1.2, 0.58) | 0.327 | 1.42 (0.97, 2.08) | 0.07 |
| Richer | 0.92 (0.49, 1.72) | 0.787 | 1.74 (0.81, 3.76) | 0.155 | 1.27 (1.85, 0.87) | 0.218 | 1.54 (1.08, 2.47) | 0.019 |
| Richest | 1.67 (0.89, 3.11) | 0.108 | 2.54 (1.21, 5.34) | 0.014 | 2.05 (3.05, 1.38) | <0.001 | 3.05 (2.03, 4.60) | <0.001 |
| Body Mass Index | | | | | | | | |
| Underweight | 0.96 (0.55, 1.68) | 0.884 | 1.24 (0.73, 2.1) | 0.422 | 0.77 (1.07, 0.55) | 0.119 | 0.58 (0.49, 0.94) | 0.02 |
| Normal weight | Ref | | Ref | | Ref | | Ref | |
| Overweight | 2.27 (1.47, 3.5) | <0.001 | 0.99 (0.7, 1.41) | 0.971 | 1.32 (1.8, 0.97) | 0.077 | 1.16 (1.06, 1.73) | 0.015 |
| Obese | 2.86 (1.88, 4.34) | <0.001 | 1.5 (1.04, 2.16) | 0.029 | 1.1 (1.55, 0.78) | 0.581 | 1.36 (0.98, 1.89) | 0.068 |
| Currently working | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 0.91 (0.6, 1.38) | 0.664 | 1.04 (0.73, 1.48) | 0.828 | 0.69 (0.49, 0.99) | 0.044 | 0.73 (0.57, 0.95) | 0.02 |
| Hypertension | | | | | | | | |
| No | Ref | | Ref | | Ref | | Ref | |
| Yes | 1.52 (1.15, 2.02) | 0.004 | 1.58 (1.18, 2.11) | 0.002 | 1.50 (1.91, 1.18) | 0.001 | 1.16 (1.29, 1.99) | <0.001 |

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

| Section/Topic | Item # | Recommendation | Reported on page # |
|------------------------------|--------|--|--------------------|
| Title and abstract | 1 | (a) Indicate the study's design with a commonly used term in the title or the abstract | 2 |
| | | (b) Provide in the abstract an informative and balanced summary of what was done and what was found | 2 |
| Introduction | | | |
| Background/rationale | 2 | Explain the scientific background and rationale for the investigation being reported | 4-6 |
| Objectives | 3 | State specific objectives, including any prespecified hypotheses | 5 - 6 |
| Methods | | | |
| Study design | 4 | Present key elements of study design early in the paper | 6 |
| Setting | 5 | Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection | 6 |
| Participants | 6 | (a) Give the eligibility criteria, and the sources and methods of selection of participants | 6-7 |
| Variables | 7 | Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable | 7 |
| Data sources/ measurement | 8* | For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group | Not applicable |
| Bias | 9 | Describe any efforts to address potential sources of bias | Not applicable |
| Study size | 10 | Explain how the study size was arrived at | 6 |
| Quantitative variables | 11 | Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why | 8 |
| Statistical methods | 12 | (a) Describe all statistical methods, including those used to control for confounding | 8 |
| | | (b) Describe any methods used to examine subgroups and interactions | 8 |
| | | (c) Explain how missing data were addressed | 6-8 |
| | | (d) If applicable, describe analytical methods taking account of sampling strategy | 8 |

| | | | |
|--------------------------|-----|--|----------------|
| | | (e) Describe any sensitivity analyses | 8 |
| Results | | | |
| Participants | 13* | (a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed | 6- 9 |
| | | (b) Give reasons for non-participation at each stage | Not applicable |
| | | (c) Consider use of a flow diagram | Not applicable |
| Descriptive data | 14* | (a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders | 9 |
| | | (b) Indicate number of participants with missing data for each variable of interest | Not applicable |
| Outcome data | 15* | Report numbers of outcome events or summary measures | 9-10 |
| Main results | 16 | (a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included | 12-13 |
| | | (b) Report category boundaries when continuous variables were categorized | 7 |
| | | (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period | 11-12 |
| Other analyses | 17 | Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses | 12 |
| Discussion | | | |
| Key results | 18 | Summarise key results with reference to study objectives | 13 |
| Limitations | 19 | Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias | 14-16 |
| Interpretation | 20 | Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence | 3, 14-16 |
| Generalisability | 21 | Discuss the generalisability (external validity) of the study results | 14-16 |
| Other information | | | |
| Funding | 22 | Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based | Not applicable |

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>