Parenteral Pethidine for labour pain relief and substance use disorder: 20-year follow-up cohort study in offspring

Robert Rodrigues Pereira,^{1,2,3} Humphrey Kanhai,⁴ Frits Rosendaal,⁵ Paula van Dommelen,⁶ Dick Swaab,⁷ Erik Rodrigues Pereira,⁸ Ben van de Wetering⁹

To cite: Rodrigues Pereira R, Kanhai H, Rosendaal F, *et al*. Parenteral Pethidine for labour pain relief and substance use disorder: 20-year follow-up cohort study in offspring. *BMJ Open* 2012;2:e000719. doi:10.1136/bmjopen-2011-000719

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2011-000719>).

Received 5 December 2011
Accepted 1 May 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

ABSTRACT

Objective: To determine whether use of intrapartum Pethidine pain analgesia increases the risk for substance use disorder in adult offspring.

Design: Analysis of data from a cohort study.

Setting: Academic hospital in Leiden, the Netherlands.

Participants: 133 cases and 164 control individuals, aged 18–20 years at follow-up.

Main outcome measure: Incidence of substance use disorder or use of alcohol and tobacco.

Results: The lifetime use of addictive substances in children exposed to intrapartum Pethidine analgesia was 45% of 133 children versus 48% of 164 not-exposed subjects (adjusted OR=0.79, 95% CI 0.48 to 1.29). Recent use of alcohol, tobacco and hard drugs showed no statistical difference either.

Conclusion: Pethidine for labour pain medication appears not to be associated with substance misuse or smoking in later life.

INTRODUCTION

Analgesia during labour is common worldwide. Methods used include barbiturates, nitrous oxide, opioids, epidural analgesia, transcutaneous electric nerve stimulation, psychoprophylaxis and hypnosis. In the Netherlands, opiates had been used in 7%–15% of all deliveries between 2000 and 2007, which is in 15 000–30 000 mothers a year.¹ In the 1970s, intramuscular Pethidine® was the most common drug for pain relief during labour. In recent years, a rise in the use of epidural or spinal analgesia and in PCA (patient-controlled analgesia) with very short acting opiates intravenously is observed. However, Pethidine® is the most commonly used opioid worldwide because it is cheap and easy to administer. Concerns have been raised about its effectiveness and potential maternal, fetal and neonatal side effects.²

ARTICLE SUMMARY

Article focus

- Research question: does perinatal Pethidine for labour pain relief lead to substance use disorder in later life?
- Is it possible to confirm the published data on an OR of around 5?
- We hypothesise that there is no relationship.

Key messages

- We cannot confirm earlier data on the risk of substance use disorder after perinatal analgesia.
- We did not find a risk for smoking and drinking alcohol either.
- Perinatal Pethidine analgesia appears not to be associated with substance misuse in later life.

Strengths and limitations of this study

- This is a well-designed large cohort study on long-term effects of perinatal Pethidine analgesia.
- The main limitation is the relatively low response and the possible selective response.

After parenteral therapy in the mother, the opiates can be detected in cord blood with a plasma level of 50% of that of the mother. After birth, the child is often sleepy and slightly respiratory depressed for a few hours.^{3 4}

In 1987, the issue of development of substance use disorders (SUDs) and behavioural problems in the offspring after perinatal analgesic medication was addressed.⁵ The authors reported ORs of 4.7 compared with individuals who did not receive perinatal analgesic drugs. These results were derived from case–control studies in patients with SUD.^{6–8}

The SUD was attributed to the ‘imprinting hypothesis’ and was first published in the BMJ in 1990 by Jacobson *et al.*⁷ The brain, when exposed to a noxe during a window in time before or during birth, could be

For numbered affiliations see end of article.

Correspondence to

Dr Robert Rodrigues Pereira;
pereirar@maasstadziekenhuis.nl

affected permanently by changing neurotransmitter receptors, synaptogenesis, myelination, proliferation, apoptosis, migration of neuronal cells or by stunting of dendrite growth.^{9–12} An epidemiological and clinical study showed that cannabis exposure before birth is associated with impulsive and psychiatric disorders in later life.¹³ Conflicting results have been published about the association of autism spectrum disorders, developmental delay and learning disorders in offspring after peripartum exposure to analgesics.^{14–17}

No long-term follow-up study in children born after opioid use of the mother during pregnancy or after opioid labour analgesia has been published. A recent Cochrane Review has been published looking at the effectiveness and side effects of intrapartum parenteral opioids.¹⁸ Short-term follow-up studies did not show sequelae in children in their development up to 5 years.^{19–22}

We investigated the association between Pethidine® use and the risk on smoking, drinking alcohol or drug abuse in offspring 20 years after birth.

METHODS

Sample and study design

Power calculation

Considering the prevalence of drug abuse, smoking and drinking alcohol, we needed 160 participants in each group. In total, 85 individuals in each group were sufficient to detect an OR of 4.7 or more between the groups with 80% power and a type I error of 0.05,⁷ assuming a prevalence of major drug abuse in the control group of 5%.¹ Because of the much more frequent use of tobacco and alcohol, this sample size was sufficient to detect ORs around two.

Data about lifetime prevalence and recent substance use are available for Europe²³ and for the Netherlands²⁴ (table 1).

Identification of the cohort variables

After ethical approval by the Medical Ethical Committee of the Leiden University Medical Center, the birth files from the Academic Hospital Leiden from 1986 to 1987 were used to compose two groups of participants: one cohort with and the other without labour analgesia by Pethidine®. Included were only healthy babies without

congenital anomalies, born at term after an uncomplicated delivery and who had not been admitted in the paediatric ward. Information was available on maternal characteristics and obstetric history including medication as well as the postpartum condition of the newborn.

Data collection

After finding the recent addresses, we were able to send validated questionnaires about lifetime and recent (last month) use of cigarettes, alcohol and drugs. The first is the National Drugs Questionnaire as a part of the Permanent National Life Style Inquiry that is used from 1997 (with computer-assisted personal interviewing or in case of drug questions with computer-assisted self-interviewing method) that meet with the European Statistics Code of Practice. The second questionnaire is a validated questionnaire for young adults about life events, schooling and behaviour.²⁵

Analysis

The primary analysis was a comparison of the prevalence of substance abuse at adult age between the Pethidine®-exposed and not-exposed groups. Subsequently, the results were compared with the national drug monitor study that was done in the same period. A multivariate logistic regression model was used to calculate ORs and 95% CIs for the outcome measures with age, sex, religion and parental education as potential confounders. The analyses were performed with SPSS V.11.5 for Windows (SPSS Inc).

Non-response

Non-responders received a short questionnaire about yes or no lifetime or recent use of alcohol, smoking and drugs. The individuals who responded to the short questionnaire and those who only partially filled in the questionnaire were analysed separately.

RESULTS

From a total of 715 deliveries, 91% of the addresses were found. Of these 651 individuals, 347 (53%) returned the questionnaires. One hundred and thirty-three participants with and 164 without Pethidine® analgesia could be fully evaluated (n=297, 46%). After a second call, 26 individuals with and 24 without Pethidine® analgesia completed the questionnaire and 92 (53 with and 39 without Pethidine®) completed the short questionnaire.

All together, 439 children (67% of 651 children) were analysed. Both the index and the control group showed the same distribution of age and sex. There were no differences in the distribution of parental education or religion (table 2). These were also similar to national data. The peripartum data of both groups showed no differences in birth weight, sex and Apgar scores. All newborns were healthy and none was admitted in the paediatric ward.

No differences were found in lifetime or recent use of drugs, alcohol or tobacco use between both groups. The

Table 1 Lifetime prevalence and recent tobacco, alcohol and drug use in the Netherlands²⁴

	Lifetime prevalence (%)	Recent use, last month (%)
Smoking	72	37
Alcohol	90	79
Cannabis	37	17
XTC	6.2	2.2
Amfetamine	5	1.9
Cocaine	4.3	1.7

Table 2 Distribution of parental education or religion stratified by Pethidine® and no Pethidine® groups

	Total group (n=297)	% Pethidine® (n=133)	% No Pethidine® (n=164)
Religion			
Christian	41.4	39.8	42.7
Other	4.4	6.8	2.4
No	54.2	53.4	54.9
Education			
Low	38.4	32.3	43.3
Intermediate	27.9	32.3	24.4
High	33.7	35.3	32.3
Sex			
Male	39.4	42.1	37.2
Female	60.6	57.9	62.8

Low, elementary school and vocational education; intermediate, high school; high, university.

OR for ever drug abuse was 0.79 (95% CI 0.48 to 1.29) and for recent drug abuse 1.08 (95% CI 0.49 to 2.37). We also gathered data about alcohol use, cannabis and smoking during and after secondary school. These data did not show differences between the groups either, the adjusted ORs are presented in table 3. In total, 92 children responded to the short questionnaire and 50 children partially filled out the standard questionnaire. No difference was found in lifetime prevalence of smoking, alcohol and drug abuse between the Pethidine® and the non-Pethidine® group independent of the dose that ranged from 75 to 150 mg. The ORs were all close to unity, varying between 0.58 and 1.42.

DISCUSSION

This 20-year follow-up study showed that labour analgesia by Pethidine® did not significantly increase the risk of SUD in offspring. We found no association for

hard drugs, tranquilisers, hallucinogenics, anabolics or soft drugs (including tobacco and alcohol use). This is in line with the results of the study from the Mayo clinics on the risk of learning disability after exposure to peripartum analgesia.¹⁴

The strength of this study is that it is a large, long-term follow-up study of a healthy cohort of neonates.

We evaluated putative confounders, such as socioeconomic status, sex, religion and parental education, but did not find differences between the index and the control group. Uncontrolled confounding is always possible, but in this instance, one would expect these to lead to spurious positive associations and not to the absence of an association. Further investigation of potential confounders could be valuable. Given the result of the previous studies reporting a fivefold increased risk, our sample size with a minimal detectable RR of 2 was conservative. However, this implies that we could not detect small risk increases. Nevertheless, the most likely interpretation of our findings, which exclude a risk increase of more than 29% of lifetime drug abuse in children exposed to Pethidine, is the absence of any association.

The main limitation of the study is the low response and the possibility of selected response. It is not inconceivable that drug users responded less than others. It is of some concern that response was lower in the group exposed during labour (54%) than in those who were not (42%). Still, if selective response was only associated to drug use in adult life, one would expect an attenuated effect but not the absence of any effect. Bias would have resulted from a differential response related to opioid use during delivery, irrespective of drug use later in life. This seems implausible, as most young adults will not be aware of the analgesic medication of their mothers during birth. Reassuring for the absence of major bias is the fact that the non-respondents to the first questionnaire, who responded to the second short questionnaire

Table 3 Lifetime or recent use of alcohol, tobacco and drugs stratified by Pethidine® and no Pethidine® groups

	Total group (n=297)	% Pethidine® (n=133)	% No Pethidine® (n=164)	Adjusted OR (95% CI)*
Smoking cigarettes ever	70.4	68.4	72.0	0.84 (0.50 to 1.41)
Smoking cigarettes recent	41.4	40.6	41.5	0.98 (0.61 to 1.58)
Alcohol ever	98.7	98.5	98.8	0.81 (0.10 to 6.23)
Alcohol recent	86.5	86.5	86.6	0.91 (0.45 to 1.81)
Smoking cannabis ever	46.5	45.1	47.6	0.76 (0.46 to 1.24)
Smoking cannabis recent	8.8	9.8	7.9	1.14 (0.50 to 2.63)
XTC ever	10.1	12.0	8.5	1.42 (0.64 to 3.17)
XTC recent	1.0	2.3	0	—
Amphetamine ever	5.4	4.5	6.1	0.66 (0.22 to 1.99)
Amphetamine recent	1.0	0.8	1.2	0.58 (0.05 to 6.61)
Cocaine ever	6.4	6.0	6.7	0.85 (0.32 to 2.29)
Cocaine recent	2.7	2.3	3.0	0.58 (0.12 to 2.68)
Drug abuse† ever	46.8	45.9	47.6	0.79 (0.48 to 1.29)
Drug abuse† recent	10.1	10.5	9.8	1.08 (0.49 to 2.37)

*Adjusted for age, religion and parental education.

†This includes cannabis, XTC, Amphetamine and cocaine.

showed the same negative association for SUD, drinking and smoking. The non-responders showed an unequal distribution in sex compared with the responders: 41% vs 59% male/female ratio.

In this large follow-up study, we cannot confirm the results of previous studies that parenteral Pethidine® for labour pain relief is associated with SUD in later life. Because of the above-mentioned limitations, further research is needed to assess possible associations between other forms of intrapartum analgesia and SUD.

Author affiliations

¹Department of Paediatrics, Maasstadziekenhuis, Rotterdam, the Netherlands

²Bouman Mental Health Care, Rotterdam, the Netherlands

³Department of Prevention and Care Netherlands Institute of Applied Sciences TNO, Leiden, the Netherlands

⁴Department of Obstetrics, Leiden University Medical Center, Leiden, the Netherlands

⁵Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands

⁶Department of statistics, Netherlands Institute of Applied Sciences TNO, Leiden, the Netherlands

⁷Netherlands Institute for Neuroscience, an Institute of the KNAW, Amsterdam, the Netherlands

⁸Central Office for Motor Vehicle Driving Testing (CBR), Amsterdam, the Netherlands

⁹Bouman Mental Health Care, Rotterdam, the Netherlands

Contributors All authors contributed to the following criteria: (1) substantial contribution to conception and design, acquisition of data or analysis and interpretation of data; (2) drafting the article or revising it critically for important intellectual content and (3) final approval of the version to be published.

Funding This research received no specific grant from any funding agency in public, commercial or not-for-profit sectors.

Competing interests None.

Patient consent Obtained.

Ethics approval Ethics approval was provided by Medical Ethical Committee of the Leiden University Medical Center, the Netherlands.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data are uploaded to the Dryad data repository and available for any scientist.

REFERENCES

1. The Netherlands Perinatal Registry. http://www.perinatreg.nl/home_english
2. Bricker L, Lavender T. Parenteral opioids for labour pain relief: a systematic review. *Am J Obstet Gynecol* 2002;186(5 Suppl Nature): S94–109.
3. Thomas H, Asskali F, Vettermann J. Addition of fentanyl to bupivacaine—peridural analgesia in cesarian section. *Anaesthesist* 1996;45:635–42.
4. Nicolle E, Devillier P, Durand C, et al. Therapeutic monitoring of nalbuphine: transplacental transfer and estimated pharmacokinetics in the neonate. *Eur J Clin Pharmacol* 1996;49:485–9.
5. Jacobson B, Eklund G, Hamberger L, et al. Perinatal origin of adult self-destructing behaviour. *Acta Psychiatr Scand* 1987;76:364–71.
6. Jacobson B, Nyberg K, Eklund G, et al. Obstetric pain medication and eventual adult amphetamine addiction in offspring. *Acta Obstet Gynecol Scand* 1988;67:677–82.
7. Jacobson B, Nyberg K, Grönbladh L, et al. Opiate addiction in adult offspring through possible imprinting after obstetric treatment. *BMJ* 1990;30:1067–70.
8. Nyberg K, Buka S, Lipsitt LP. Perinatal medication as a potential risk factor for adult abuse in a North American cohort. *Epidemiology* 2000;11:715–16.
9. Tchernitchin AN, Tchernitchin NN, Mena MA, et al. Imprinting: perinatal exposures cause the development of diseases during the adult age. *Acta Biol Hung* 1999;50:425–40.
10. Fernandez-Ruiz J, Berrendero F, Hernández ML, et al. The endogenous cannabinoid system and brain development. *Trends Neurosci* 2000;23:14–20.
11. Swaab DF, Boer K, Mirmian M. Functional teratogenic effects of chemicals on the developing brain. In: Levene MI, Chervenak FA, eds. *Fetal and Neonatal Neurology and Neurosurgery*. Toronto: Churchill Livingstone, 2001:263–77.
12. Ricalde A, Hammer RP. Perinatal opiate treatment delays growth of cortical dendrites. *Neurosci Lett* 1990;115:137–43.
13. Spano MS, Ellgren M, Wang X, et al. Prenatal cannabis exposure increases heroin seeking with allostatic changes in limbic enkefalin systems in adulthood. *Biol Psychiatry* 2007;61:554–63.
14. Flick RP, Lee K, Hofer RE, et al. Neuraxial labor analgesia for vaginal delivery and its effect on childhood learning disabilities. *Anaesth Analg* 2011;112:1424–31.
15. Hattori R, Desimaru M, Nagayami I, et al. Autistic and developmental disorders after general anesthetic delivery. *Lancet* 1991;337:1357–8.
16. Sun LS. Editorial: labor analgesia and the developing human brain. *Anaesth Analg* 2011;112:1265–7.
17. Wong CA, Scavone BM, Peaceman AM, et al. The risk of caesarean delivery with neuraxial analgesia given early versus late in labor. *N Engl J Med* 2005;352:655–65.
18. Ullman R, Smith LA, Burns E, et al. Parenteral opioids for maternal pain management in labour. *Cochrane Database Syst Rev* 2010;(9): CD007396.
19. Kaye K, Elkind L, Goldberg D, et al. Birth outcomes for infants of drug abusing mothers. *N Y State J Med* 1989;89:256–61.
20. Doberczak TM, Kandall SR, Friedmann P. Relationship between maternal methadone dosage, maternal-neonatal methadone levels, and neonatal withdrawal. *Obstet Gynecol* 1993;81:936–40.
21. Hagopian GS, Wolfe HM, Sokol RJ, et al. Neonatal outcome following methadone exposure in utero. *J Matern Fetal Med* 1996;5:348–54.
22. Thaithumyanon P, Limpongsanurak S, Praisuwanna P, et al. Perinatal effects of amphetamine and heroin use during pregnancy on the mother and infant. *J Med Assoc Thai* 2007;88:1506–13.
23. *Jaarverslag over de stand van de drugsproblematiek in de Europese Unie*. Lisbon: E.W.D.D, 2000. ISBN 92-9168-100-8.
24. *Trimbos-instituut, Jeugd en riskant gedrag, kerngegevens 1999*, Utrecht: Cf international equivalent, 2000. <http://www.espad.org/espado-reports>
25. Stam H, Grootenhuys MA, Last BF. The course of life of survivors of childhood cancer. *Psychooncology* 2005;14:227–38.