Social inequality and infant health in the UK: systematic review and meta-analyses

Alison L Weightman,¹ Helen E Morgan,¹ Michael A Shepherd,² Hilary Kitcher,¹ Chris Roberts,³ Frank D Dunstan⁴

To cite: Weightman AL, Morgan HE, Shepherd MA, *et al*. Social inequality and infant health in the UK: systematic review and meta-analyses. *BMJ Open* 2012;**2**:e000964. doi:10.1136/bmjopen-2012-000964

► Prepublication history and additional appendices for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-000964>).

Received 2 February 2012
Accepted 10 April 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

¹Support Unit for Research Evidence (SURE), Information Services, Cardiff University, Cardiff, UK

²Freelance public health researcher, Somerset, UK

³Health, Social Services and Children Analytical Team, Welsh Government, Cardiff, UK

⁴Cochrane Institute of Primary Care and Public Health, School of Medicine, Cardiff University, Cardiff, UK

Correspondence to

Dr Alison Lesley Weightman; weightmanal@cardiff.ac.uk

ABSTRACT

Objectives: To determine the association between area and individual measures of social disadvantage and infant health in the UK.

Design: Systematic review and meta-analyses.

Data sources: 26 databases and websites, reference lists, experts in the field and hand-searching.

Study selection: 36 prospective and retrospective observational studies with socioeconomic data and health outcomes for infants in the UK, published from 1994 to May 2011.

Data extraction and synthesis: 2 independent reviewers assessed the methodological quality of the studies and abstracted data. Where possible, study outcomes were reported as ORs for the highest versus the lowest deprivation quintile.

Results: In relation to the highest versus lowest area deprivation quintiles, the odds of adverse birth outcomes were 1.81 (95% CI 1.71 to 1.92) for low birth weight, 1.67 (95% CI 1.42 to 1.96) for premature birth and 1.54 (95% CI 1.39 to 1.72) for stillbirth. For infant mortality rates, the ORs were 1.72 (95% CI 1.37 to 2.15) overall, 1.61 (95% CI 1.08 to 2.39) for neonatal and 2.31 (95% CI 2.03 to 2.64) for post-neonatal mortality. For lowest versus highest social class, the odds were 1.79 (95% CI 1.43 to 2.24) for low birth weight, 1.52 (95% CI 1.44 to 1.61) for overall infant mortality, 1.42 (95% CI 1.33 to 1.51) for neonatal and 1.69 (95% CI 1.53 to 1.87) for post-neonatal mortality. There are similar patterns for other infant health outcomes with the possible exception of failure to thrive, where there is no clear association.

Conclusions: This review quantifies the influence of social disadvantage on infant outcomes in the UK. The magnitude of effect is similar across a range of area and individual deprivation measures and birth and mortality outcomes. Further research should explore the factors that are more proximal to mothers and infants, to help throw light on the most appropriate times to provide support and the form(s) that this support should take.

ARTICLE SUMMARY

Article focus

- To determine the association between area and individual measures of social disadvantage and infant health in the UK.

Key messages

- It is possible to be reasonably confident of the magnitude of effect of social disadvantage on health outcomes for UK infants, in particular the effects of area or individual deprivation on birth outcomes and infant mortality.
- The magnitude of effect, of a range of measures of social disadvantage on infant health outcomes, is broadly similar and has remained so over the past 30 years.
- Future research should concentrate on the more proximal determinants of infant health outcomes (such as individual maternal and infant factors) and the impact of any interventions employed to mitigate these determinants.

Strengths and limitations of this study

- A rigorous systematic review and accompanying meta-analyses provide confidence in the findings.
- Specific relevance to the UK.
- A risk of publication bias (missing studies) cannot be ruled out though, given the large size of many included studies, the likely effect is minimal.
- The review concentrates on health outcomes only.

INTRODUCTION

Socioeconomic status, as indicated by level of income, education, wealth, occupation and access to resources, is well established as associated with an individual's health and well-being. Furthermore, the correlation is evident throughout the life course. Such social variables are key elements of the aetiology of ill health and disease.^{1–3} The same social factors also determine the health and life chances of the unborn and newborn child and impact on early child development. These interactions have been theorised in the social–ecological model of health, which acknowledges the

cumulative impact of individual, familial, community and societal forces on people's health.^{4 5}

The UK government and devolved administrations are committed to tackling the social, economic and environmental factors that influence health through evidence-based approaches to the underlying social determinants of ill health and to interventions to address these inequalities.^{6–9} This approach is in line with the conclusions of WHO sponsored Commission on the Social Determinants of Health.³

Birth and early development play a vital role in creating and maintaining socioeconomic health inequalities through adulthood.^{6 10} From a pathway approach, early development from conception to 5 years of age is widely accepted as establishing the foundation for learning, behaviour and health throughout the life cycle.¹¹ WHO Commission on the Social Determinants of Health in particular stressed the critical role of child health in addressing inequities in health. Investments in child health and development are seen as forces to equalise health status through the life course.³ These points were reiterated in the recent Marmot Review, which particularly stressed the importance of giving every child the best possible start in life.⁶

There is an implicit recognition of the importance of social determinants to the well-being of children in national policy. For example, a stated aim of 'Healthy Lives, Brighter Futures' is to get the right services, advice and support to all parents with more intensive support for the most vulnerable.¹²

The UK performs very poorly in comparison with similar countries on mortality among the under 5s.¹³ A recent index of child well-being in the European Union¹⁴ suggested that the UK ranked 24/29 for both child health from birth and overall child well-being. Furthermore, the UK had a stillbirth rate of 3.5 per 1000 births in 2009. Within high-income nations, only France and Austria had higher levels.¹⁵ Such poor outcomes in comparison to our nearest neighbours suggest that significant action is required to improve health and well-being in childhood.

The objectives of this review were to assess the current evidence for the effects of social disadvantage on birth and infant outcomes for children born in the UK to provide a firm basis for practitioners and policy makers on which to measure the effect of interventions to address these inequalities. This systematic review builds on a previous review of prospective studies carried out for the Welsh government to examine the social determinants of child health in the UK and to develop social indicators from these determinants.¹⁶

This analysis provides a current summary of relevant and well-conducted epidemiological research into the links between social determinants and infant health in the UK.

METHODS

Inclusion/exclusion criteria

Research studies were sought that reported socioeconomic data and health outcomes for infants (0–12

months) and were carried out in the UK and published between January 1994 and May 2011. The most recent update search was completed on 19 May 2011. There were no language exclusions. A 'best evidence' approach was adopted by using data from longitudinal and record linkage studies. Prospective cohort, case–control and retrospective cohort studies with a sample size of 200 or more were included, as well as record linkage analyses of routinely collected data. Case studies and cross-sectional surveys (with data from a single time point) were excluded.

The outcome criteria for the studies were preterm birth, birth weight, mortality, diagnosed illness, attendance at primary or secondary care in relation to ill health, infection, injury or disability, growth and development. The social determinants explored were specific area-based and individual measures of social disadvantage which included area deprivation scores^{17–20} and individual measures of deprivation (occupational social class, household income/poverty, parental educational status). Lone parenthood was also included as a social determinant, albeit not a direct measure of social disadvantage.

To ensure that the review focused on specific infant health outcomes for the general current population, papers on the following topics were excluded: congenital malformations, in-care or adopted infants, infants with pre-existing medical conditions, specific subgroups of mothers (eg, previous major pregnancy complication) and data from pre-1970. Access to treatment or screening services, immunisation uptake, child abuse or domestic violence studies were excluded unless reported with a social determinant and child health outcome.

Search sources and search strategy

The following databases, websites and other sources were searched: ASSIA (Cambridge Scientific Abstracts); British Nursing Index (OVID); ChildData; CINAHL (Ebsco); Community Wise; Conference Proceedings Citation Index—Science and Social Science & Humanities (Web of Science); Embase (OVID); EPPI Centre DoPHER; HMIC (OVID); Joseph Rowntree Foundation; Local Government Data Unit Wales; Medline (OVID); Medline in Process (OVID); NHS Plans and National Service Frameworks for Wales and England in relation to children; National Institute for Health and Clinical Excellence (NICE); Office for National Statistics; PsycINFO (OVID); ReFER (archive up to September 2007); Science Citation Index (Web of Science); Open SIGLE (replaced by OpenGrey), Social Care Online; Social Science Citation Index (Web of Science); Social Services Abstracts; Sociological Abstracts (Cambridge Scientific Abstracts); WHO Health Evidence Network.

The search terms were developed and tested by qualified librarians (HK and ALW) in one database, Medline, to a high recall of relevant studies (sensitive) without too many irrelevant studies (specific).

The following search terms were used in Medline (where * is the truncation symbol; ab=abstract;

cp=country of publication, pt=publication type; ti=title; /=subject heading):

Infant health

Exp infants/OR (Babies or baby or birth* or infant* or neonatal or newborn or perinatal or postnatal or neonatal or perinatal).ab,ti.

AND

Social determinants

Exp socioeconomic factors/OR exp social class/OR (benefit recipient* or deprivation or economic inactivity or educational achievement or educational attainment or employment status or financial hardship or home own* or house own* or housing tenure or inequalit* or inequit* or job opportunit* or lone parent or low income or low pay or marital separation) OR (neglect or overcrowd* or poor environment or poor housing or poverty or property own* or prosperity or single parent* or social adversity or social capital or social class or social disadvantage or social disparit* or social exclusion or social inclusion or social gradient or social housing or social integration or social interaction) OR (social isolation or social mobility or social network* or social position or social relationship* or social security or social status or social stigma or social trend* or social welfare or sociodemographic or socioeconomic or socio-demographic or socio-economic or sole parent or standard of living or unemployment*).ab,ti.

AND

Study designs

exp Meta-Analysis/ OR review.pt OR (census* or cohort* or survey* or evaluat* or longitudinal* or questionnaire* or meta-analys* or metaanaly* or meta analys* or registry or registries or systematic* review* or systematic overview).ti,ab.

AND

Exp Great Britain/ OR (UK OR United Kingdom OR England OR Wales OR Scotland OR Britain OR British OR English OR Welsh OR Ireland OR Irish OR Scottish).ti,ab,cp.

Searches were adapted to other databases to replicate, as closely as possible, the Medline search. In addition, and to minimise the potential for publication bias, a range of supplementary ('snowballing') techniques was used to increase the sensitivity of the search and to ensure coverage of grey literature and unpublished studies. These included reference list follow-up, citation tracking of relevant studies (to find newer studies), contact with subject experts and organisations, and table of contents scanning for the journals that appeared most frequently in the list of relevant studies: *Journal of Epidemiology and Community Health*, *Archives of Disease in Childhood* (including the *Fetal and Neonatal edition*) and *British Medical Journal*. Finally, all systematic reviews on relevant topics were unpicked for primary studies

meeting inclusion/exclusion criteria. This included the publication on which this review builds¹⁶ and a more recent systematic search carried out by members of the same team for the Welsh government to support the development of a Child Health Monitor.²¹

Duplicate references were excluded. All titles and, where necessary, abstracts were then screened to eliminate duplicates and obviously irrelevant citations. The full text of all potentially relevant papers was retrieved and screened independently for eligibility by two reviewers using a standardised eligibility form, with adjudication from a third reviewer in cases of disagreement (HK, HEM, MAS and ALW).

Following screening of 5173 citations found in the literature search, 88 papers potentially met the inclusion criteria and were examined in full text. Thirty-six studies met all inclusion and critical appraisal criteria and were included in the final review. Details are given in the flow diagram of figure 1

Classification of social determinants

Many studies, particularly those using routine data, use area-based deprivation data such as the Townsend²⁰ and Carstairs¹⁷ scores, the UK Indices of Multiple Deprivation (IMD)¹⁹ and the Jarman Index.¹⁸ These are derived from routine data sources, including census data. Most researchers divide geographical units into classes, such as quintiles, based on these deprivation scores. Different geographical areas are used, ranging from enumeration districts, with a population of around 500, to Electoral Wards, with about 3000 people. Many recent studies use the 2001 UK Census geographical unit of Lower Super Output Area containing, on average, a population of 1500.

Some studies used social class, usually based on the Registrar General's classification, instead of an area measure. This has the advantage of being an individual measure and is recorded by the Office for National Statistics on a randomly selected 10% of births. Most studies which used this measure compared class I, consisting of professionals, with class V, made up of unskilled manual workers.

Critical appraisal and data extraction

Following pre-tests to confirm inter-rater reliability and iron out any queries, data were extracted independently by two reviewers (shared by FDD, HEM, MAS, ALW). Each study was assessed for quality against a critical appraisal checklist based on the Newcastle–Ottawa Quality Assessment Scale (NOS).²² Any disagreements were resolved by a third reviewer. There is, as yet, no internationally established quality assessment tool for observational epidemiological studies,²³ and the NOS has not been validated.²⁴ Thus, the tool was used to help identify potential methodological weaknesses rather than to provide a definitive quality score for each study. Methodological quality indicators (study design, data sources, response rate and any adjustments to control for baseline variables) were summarised in the Evidence Table (online appendix 1).

Figure 1 Flow diagram.

A paper was excluded if it did not provide usable data to allow a quantitative comparison of the risk for a health outcome in relation to a measure of individual or area social inequality with an indication of statistical significance (CI or p value). If a research publication reported a non-significant result without supporting data, but the statistical rigour suggested that this was a reliable result, a non-significant result was recorded. In the latter case, it was clearly not possible to include these papers in any meta-analysis.

The key components of the data extraction and critical appraisal outcomes are presented in the Evidence Table (online appendix 1). Details of the eligibility, critical appraisal and data extraction methods are available from the authors.

Study outcomes were analysed by a statistician (FDD) and reported as described by the authors of each paper. Where possible, outcomes were reported as an 'effect size' observed with subjects from highest versus lowest deprivation exposures in the Evidence Table (online appendix 1 and 2) to allow comparability between studies. In some cases, the authors did not report the data in the most suitable way, but it was possible to calculate effect sizes and CIs from the data shown. Where possible, an OR comparing the highest and lowest deprivation levels was calculated, as this was the most commonly reported measure. As most outcomes measured were relatively rare, differences between relative risks and ORs will be modest. Some studies adjusted the ORs to allow for confounders, and where possible, the most adjusted data have been reported.

Meta-analyses were carried out, where possible, when a group of papers were sufficiently homogeneous in respect of both outcome and deprivation measures. Because adjustments varied between studies, these anal-

yses were carried out on unadjusted results. Random effects models were used to allow for heterogeneity. The number of studies on any one outcome was too small for funnel plots to be of any value so the risk of publication bias cannot be excluded. However, given the huge size of many of the studies included, CIs are quite narrow and missing small studies would likely have little effect.

RESULTS

Thirty-six studies met the inclusion criteria, of which 13 were prospective cohort (longitudinal) or case-control studies and 23 were based on data collected retrospectively from medical records or routinely collected data (eg, record linkage between a health outcomes register and census data). Findings are summarised in table 1 and detailed in the text. Summary data from each included study are given in online appendix 1.

Birth weight, prematurity and cerebral palsy

Birth weight, prematurity and cerebral palsy are very closely linked outcomes,⁴¹ which are considered together.

Low birth weight

Nine studies (Paranjothy, 2010, unpublished)^{25–31 42} considered the outcome of low birth weight or very low birth weight, defined as <2500 g or <1500 g, respectively; none considered birth weight as a continuous outcome.

These are all based on large routinely collected data sets but they differ in the measures of deprivation used and in the adjustments made in analyses. In general, the studies did not control for gestational age to distinguish premature from small-for-dates babies.

Table 1 Social determinants related to specific infant health outcomes: summary results from meta-analyses

Health outcome	Social determinant	Prospective and case-control studies	Retrospective studies	Summary magnitude of effect*, OR: most vs least deprived quintiles
Birth				
Very low birth weight	Area deprivation		Bundred <i>et al</i> ²⁵ Dibben <i>et al</i> ²⁶ Paranjothy (2010, unpublished) Spencer <i>et al</i> ²⁷	1.61 (1.31 to 1.97)
Low birth weight	Area deprivation	Collingwood Bakeo and Clarke ²⁸	Bundred <i>et al</i> ²⁵ Dibben <i>et al</i> ²⁶ Gray <i>et al</i> ²⁹ Paranjothy (2010, unpublished) Pattenden <i>et al</i> ³⁰ Spencer <i>et al</i> ²⁷	1.81 (1.71 to 1.92)
	Lower social class		Fairley and Leyland ³¹ Maher and Macfarlane ³² Pattenden <i>et al</i> ³⁰ Spencer <i>et al</i> ²⁷	1.79 (1.43 to 2.24)
Preterm birth (preterm/very preterm combined)	Area deprivation	Smith <i>et al</i> ³³ Yuan <i>et al</i> ³⁵	Gray <i>et al</i> ³⁴ Paranjothy (2010, unpublished)	1.67 (1.42 to 1.96)
Stillbirth	Area deprivation		Gray <i>et al</i> ²⁹ Paranjothy (2010, unpublished)	1.54 (1.39 to 1.72)
Infant health				
Infant mortality overall	Area deprivation		Dummer and Parker ³⁶ Gray <i>et al</i> ²⁹ Oakley <i>et al</i> ³⁷	1.72 (1.37 to 2.15)
	Lower social class		Dummer and Parker ³⁶ Whitehead and Drever ³⁸ Oakley <i>et al</i> ³⁷	1.52 (1.44 to 1.61)
Neonatal mortality	Area deprivation		Dummer and Parker ³⁶ Gray <i>et al</i> ²⁹ Smith <i>et al</i> ³⁹	1.61 (1.08 to 2.39)
	Lower social class		Dummer and Parker ³⁶ Oakley <i>et al</i> ³⁷ Whitehead and Drever ³⁸	1.42 (1.33 to 1.51)
Postneonatal mortality	Area deprivation		Dummer and Parker ³⁶ Gray <i>et al</i> ²⁹ Oakley <i>et al</i> ³⁷	2.31 (2.03 to 2.64)
	Lower social class		Dummer and Parker ³⁶ Whitehead and Drever ³⁸ Oakley <i>et al</i> ³⁷ Petrou <i>et al</i> ⁴⁰	1.69 (1.53 to 1.87)

*To avoid double counting, meta-analyses were carried out using only those studies without overlapping data.

Different studies used different time periods, and it is possible that the effect of deprivation has changed over recent decades. However, two studies which subdivided the data into different decades^{28–31} suggest that the effects have not changed markedly since 1980. Some studies adjusted for confounders, typically maternal age, gender of child and whether the registration was joint or single parent (mother) only. ORs, adjusted for these potential confounders, were generally considerably closer to the null value of 1 than unadjusted ones, reflecting the importance of these factors.

In seven studies that compared area measures (Paranjothy, 2010, unpublished),^{25–30} the unadjusted ORs, comparing the most and least deprived, varied

from 1.67 to 2.01. Combining the estimates in a random effects meta-analysis, the resulting OR was 1.81 (95% CI 1.71 to 1.92) (figure 2). The study by Wright *et al*⁴² was not included in the meta-analysis because the definitions of area deprivation are unclear and it is not possible to calculate ORs with any degree of certainty.

One further study⁴³ looked at the specific issue of small for gestational age babies in the West Midlands (birth weight <10th customised centile). For UK IMD quintile 5 versus IMD 1–3 (personal communication from author), the RR of small for gestational age was 1.2 (95% CI 1.1 to 1.3). Since both outcome and comparator measures vary from the other studies, this result was not included in the meta-analysis.

Figure 2 Low birth weight versus area deprivation.

Five studies^{26 27 30–32} measured deprivation at an individual level using the Registrar General's social class categories. All but Maher and Macfarlane³² compared social class V with social class I; Maher and Macfarlane compared manual and non-manual workers, and this is probably a less extreme comparison, leading to a smaller OR. For social class, as opposed to area deprivation, Dibben *et al*²⁶ only reported a heavily adjusted OR of 1.20 and thus was not included in the meta-analysis. The unadjusted estimates of the OR varied from 1.45 to 2.17. Excluding Dibben *et al*,²⁶ a random effects model pooling these gave an overall estimate of 1.79 (95% CI 1.43 to 2.24), very similar to the estimate for the area-based measures (figure 3).

Several authors, including Fairley and Leyland³¹ and Paranjothy (2010, unpublished), noted that the effects of deprivation appeared to be greater in older mothers. This may merit further investigation.

Four of the studies (Paranjothy, 2010, unpublished)^{25–27} also examined the incidence of very low birth weight against area-based measures. There was greater heterogeneity in this case, with estimates of the

OR varying between 1.29 and 2.54. The meta-analysis gave a combined OR of 1.61 (95% CI 1.31 to 1.97) (figure 4). Again there was evidence from Paranjothy that this OR increased with maternal age.

In all the above cases, there was considerable heterogeneity; the reasons for this are not clear. All studies used large samples of routine data and, within each analysis, used broadly comparable measures of deprivation.

Overall, it is clear that deprivation is strongly associated with low and very low birth weight. The effect may vary with maternal factors such as age, and this needs further investigation.

Two studies^{26 30} found a statistically significant relationship between lone parenthood and low birth weight with estimated ORs of 1.16 (95% CI 1.09 to 1.25) and 1.46 (95% CI 1.41 to 1.52), respectively, though Pattenden *et al*³⁰ used single registration, not identical to lone parenthood, and the result of Dibben *et al*²⁶ was heavily adjusted.

Prematurity

Six papers considered preterm birth as an outcome (Paranjothy, 2010, unpublished).^{33–35 42 44} Smith *et al*⁴⁴ used 549 618 births from 1994 to 2003 from the Trent region and defined very preterm birth as 22–32 weeks and extremely preterm birth as 22–28 weeks. Combining their deciles into quintiles, the unadjusted OR for very preterm, comparing the most and least deprived quintiles, was 1.72 (95% CI 1.59 to 1.86). For extremely preterm birth, this was 1.74 (95% CI 1.54 to 1.98). Their analysis included all babies alive at the start of labour. A later paper³³ reported on a different but overlapping 10-year period from the same region. The ORs were a little higher. Paranjothy (2010, unpublished) included stillbirths in an analysis of 408 445 births in Wales between 1994 and 2005 and found an OR of 1.44 (95% CI 1.32 to 1.56) for very preterm births, comparing the most and least deprived quintiles based on the Townsend score, and the corresponding OR for extremely preterm births was 1.38 (95% CI 1.20 to 1.57). Excluding stillbirths had little effect. Gray *et al*³⁴

Figure 3 Low birth weight versus social class.

Figure 4 Very low birth weight versus area deprivation.

considered all preterm births, defined as being <37 weeks, the OR was 1.47 (95% CI 1.42 to 1.52). Since the majority of preterm births are after 32 weeks, this broadly similar result suggests that the association with deprivation is similar at different stages of prematurity. The study by Wright *et al*⁴² in Newcastle found that the RR for most deprived versus most affluent areas was 2.95 (95% CI 1.61 to 5.38). The definitions of area deprivation are unclear and thus it is not possible to calculate ORs with any degree of certainty. Thus, this paper was not included in the meta-analysis.

Yuan *et al*³⁵ used a case-control study to compare the prevalence of social problems in those whose pregnancy ended at term (≥ 37 weeks) compared with preterm (<35 weeks), the OR was 2.42 (95% CI 1.60 to 3.66). The definition of social problems is likely to identify more seriously deprived mothers than those studies using social class, and this may account for the more extreme result.

The varying definitions of prematurity in these studies mean it is thus arguable that a meta-analysis is not appropriate in this case. However, to give an indication of effect sizes, combining four studies, with useable data and no overlaps (Paranjthy, 2010, unpublished),^{33–35} all of which related to area deprivation, gave an OR of 1.67 (95% CI 1.42 to 1.96) (figure 5). The conclusions on prematurity were therefore similar to those for low birth weight, which is strongly related to prematurity. Stillbirth is discussed further under infant mortality.

Cerebral palsy

Two studies considered the relationship between cerebral palsy and area deprivation.^{45–46} Sundrum *et al*⁴⁵ used births in West Sussex and found an OR of 1.65 (95% CI 1.14 to 2.39) comparing extreme quintiles based on the Townsend score for enumeration districts. Adjusting for birth weight and gestation age gave a revised estimate of 1.55 (95% CI 1.06 to 2.25). Dolk *et al*⁴⁶ examined over 1.5 million births in five regions of the UK. Cases of cerebral palsy were divided into those acquired postnatally and those present at birth. For the former, the OR, comparing extreme quintiles based on the Carstairs score at electoral ward level, was 1.86 (95% CI 1.19 to

2.88). For non-acquired cases, the association with deprivation had the corresponding OR of 1.16 (95% CI 1.00 to 1.35). When this was adjusted for birth weight, it was reduced to 1.05 (95% CI 0.86 to 1.29), a non-significant result. This suggests that the effect of deprivation acts largely through low birth weight. There was evidence of considerable heterogeneity between regions, and this requires further investigation to seek possible explanations.

Infant mortality

Infant mortality is traditionally classified by the age of the child at death. Neonatal mortality refers to deaths up to 27 days after the birth, with those up to day 6 classed as early neonatal; the remaining neonatal deaths are termed late neonatal. Those after 27 days but up to a year are called postneonatal, while infant deaths are those between birth and a year. A stillborn baby is a baby born after the 24th week of pregnancy who does not show any signs of life. Perinatal mortality combines stillbirths and early neonatal deaths. An analysis of infant mortality needs to be separated into these different time periods as any possible effects of deprivation may operate differently in the neonatal period from the postneonatal period, for example.

For stillbirths, there were four studies (Paranjthy, 2010, unpublished)^{29–47–48} all of which postdate the change in definition resulting from the Still-Birth (Definition) Act of 1992; when births from 24-week gestation were included, the data by Guildea *et al*⁴⁷ were subsumed by that of Paranjthy. Published data from Guildea *et al*⁴⁷ are, however, included in the Evidence Table (online appendix 1). Gray *et al*,²⁹ using data from Scotland from 1994 to 2003, found an OR of 1.56 (95% CI 1.38 to 1.77) comparing extreme quintiles based on Carstairs scores calculated for postcode sectors. Paranjthy (2010, unpublished) found a very similar OR of 1.54 (95% CI 1.32 to 1.80) comparing extreme quintiles of the Townsend score calculated for lower super output areas. Combining these two studies (Paranjthy, 2010, unpublished),²⁹ a pooled estimate of the OR is 1.54 (95% CI 1.39 to 1.72) (figure 6).

Figure 5 Preterm birth versus area deprivation.

Figure 6 Stillbirth versus area deprivation.

Smeeton *et al*⁴⁸ used a case–control study in a Health Authority in London to study stillbirths, using the Jarman score to measure deprivation. The presumption is that the results were not statistically significant as no useable data were shown.

Bambang *et al*⁴⁹ was most concerned with specific causes of perinatal death in one English region; the association with deprivation varied relatively little with cause, and combining the causes gave an OR of 1.91 (95% CI 1.66 to 2.20) comparing extreme quintiles based on Townsend scores.

Four studies^{36 40 47 48} examined early neonatal death as a separate outcome; all but Smeeton also considered late neonatal mortality. Two of these studies^{40 47} found that the risk of late neonatal mortality was higher in more deprived areas, with very different ORs. Generally, results varied considerably and details are shown in online appendix 1. There was more evidence for an association of deprivation with late neonatal than early neonatal deaths.

Whitehead and Drever³⁸ gave results for perinatal mortality, which includes stillbirths and early neonatal deaths within 7 days of birth. Using data on over 14 million births from a 22-year period from England and Wales, they found an OR of 1.35 (95% CI 1.30 to 1.40).

Whitehead and Drever³⁸ also compared overall neonatal mortality between classes IV and V and classes I and II. The rate was significantly higher in classes IV and V, with an OR of 1.41 with approximate 95% CI 1.32 to 1.51. As about 75% of neonatal deaths are early neonatal deaths, it is likely that an association existed with early neonatal death. Gray *et al*²⁹ studied over 500 000 births in Scotland between 1994 and 2003; the OR for comparing quintiles based on Carstairs–Morris scores was 1.37 (95% CI 1.16 to 1.62). Oakley *et al*³⁷ took all 1 276 198 singleton births in England and Wales in 2005 and 2006 and found an OR of 2.00 (95% CI 1.79 to 2.24) comparing quintiles based on Carstairs scores. For a 10% sample, those who are routine or manual workers were compared with the category of Higher and Professional workers; the OR was 1.54 (95% CI 1.20 to 1.98). Smith *et al*³⁹ took all live births in England between 1997 and 2007, more than 6 million in total. Comparing the most extreme quintiles, based on the Index of Multiple Deprivation, the OR was 2.19 (95% CI 2.09 to 2.29). Dummer and Parker³⁶ found an OR of 1.28 (95% CI 0.82 to 2.01) comparing extreme deprivation quintiles in Cumbria. A meta-analysis was performed but Oakley *et al*³⁷ was omitted, as its study population was largely subsumed by that of Smith *et al*³⁹. The pooled OR, comparing the extreme quintiles from these three studies,^{29 36 39} was 1.61 (95% CI 1.08 to 2.39) (figure 7).

A meta-analysis comparing the extremes of social class from three studies looking at neonatal mortality^{36–38} gave a pooled OR of 1.42 (95% CI 1.33 to 1.51) (figure 8).

In summary, there does seem to be evidence for an effect of both area deprivation and social class on

Figure 7 Neonatal mortality versus area deprivation.

neonatal mortality, although there is considerable heterogeneity between studies.

Six studies^{29 36–38 40 47} assessed the effect of deprivation on postneonatal mortality. All found that the unadjusted rates were significantly higher in the most deprived areas compared with the least deprived, with ORs of 2.26 (95% CI 1.29 to 3.93) in Dummer and Parker,³⁶ 2.20 (95% CI 1.77 to 2.72) in Guilda *et al*,⁴⁷ 2.55 (95% CI 2.01 to 3.23) in Gray *et al*,²⁹ and 2.21 (95% CI 1.87 to 2.61) in Oakley *et al*.³⁷ The data from Guilda *et al*⁴⁷ were not available in a form that could be included in the meta-analysis. A pooled OR was 2.31 (95% CI 2.03 to 2.64) for the other three studies^{29 36 37} (figure 9). If raw data from Paranjothy (2010, unpublished) are included, which represent (and subsume) the Guilda *et al*⁴⁷ population, the pooled OR is very similar, 2.28 (95% CI 2.04 to 2.56).

Comparing social class, Dummer and Parker³⁶ found an OR of 2.24 (95% CI 0.81 to 6.23) and Petrou *et al*⁴⁰ recorded 1.70 (95% CI 1.14 to 2.54) comparing classes V and I. Whitehead and Drever³⁸ found an OR of 1.69 (95% CI 1.52 to 1.88) for comparing classes IV and V with I and II. Oakley *et al*³⁷ used the newer NS_SEC

Figure 8 Neonatal mortality versus social class.

Figure 9 Postneonatal mortality versus area deprivation.

classification¹⁹ and compared Higher and Professional with Routine and manual, finding an OR of 2.63 (95% CI 1.41 to 1.91). The pooled OR was 1.69 (95% CI 1.53 to 1.87) (figure 10). Petrou *et al*⁴⁰ and Oakley *et al*³⁷ adjusted for a number of factors, leading to reductions in the estimated ORs as shown in online appendix 1.

Many of these studies also gave results for overall infant mortality. Combining their results, the pooled OR was 1.72 (95% CI 1.37 to 2.15) for comparing extreme quintiles of deprivation^{29 36 37} (figure 11) and 1.52 (95% CI 1.44 to 1.61) for comparing the extreme social classes^{36–38 40} (figure 12).

Neonatal admissions

Two studies examined associations between neonatal admissions and deprivation. Manning *et al*,⁵⁰ in a study on the Wirral in England, showed a strong linear association between deprivation, based on quartiles of Townsend scores, and the rate of neonatal admissions. The OR comparing the most and least deprived quartiles was 1.91 (95% CI 1.71 to 2.11). Jenkins *et al*⁵¹ examined intensive care neonatal admissions in Northern Ireland, dividing wards into quintiles based on the Northern

Figure 10 Postneonatal mortality versus social class.

Figure 11 Overall infant mortality versus area deprivation.

Ireland Multiple Deprivation index.⁵² There was no clear pattern of the rate of admissions by quintile for the four least deprived quintiles, but the admission rate in the most deprived quintile was significantly higher than in other quintiles with an OR of 1.22 (95% CI 1.11 to 1.33) relative to the least deprived quintile. In both studies, one of the most common causes for admission was prematurity and so the strong association of prematurity with deprivation is clearly a factor.

Sudden infant death syndrome

Sudden infant death syndrome (SIDS) is a common cause of death in infancy, especially in the postneonatal period when, in spite of reducing incidence, from one recent study it accounted for about 30% of deaths in 2004–2008 (Paranjothy, 2010, unpublished). Guilda *et al*⁴⁷ found a strong association with deprivation with an OR of 3.07 (95% CI 1.97 to 4.56) comparing extreme quintiles based on the Townsend score. Sanderson *et al*⁵³ compared areas of poverty with those not of poverty and found an OR of 2.33 (95% CI 1.06 to 5.11); the definition of an area of poverty was not clear and so this is hard to compare with Guilda's result. Blair *et al*⁵⁴ analysed

Figure 12 Overall infant mortality versus social class.

a population-based case-control study and found an OR of 18.49 (95% CI 3.62 to 94.48) for overcrowding, one aspect of deprivation. The definition used here was quite extreme, based on at least two people per room in a house, and a less extreme one may give a smaller OR. Brooke *et al*⁵⁵ also used a case-control study. Deprivation was assessed using Carstairs-Morris scores,¹⁷ and the most deprived group had a significantly higher risk of SIDS than the remainder, even after adjustment for other factors, though the definition of the group was not clear. For social class, there was a significantly higher risk in classes IV and V than I and II in a univariate analysis; the OR remained >1 but was no longer significant in a multivariate analysis. Smith *et al*³⁹ looked at SIDS in the neonatal period, though the great majority occur later. Comparing the extreme quintiles, the OR was 2.20 (95% CI 1.66 to 2.91). Fleming *et al*⁶⁶ used a case-control study to examine risk factors. Comparing social classes V and I, the OR was 13.74 (95% CI 4.80 to 39.34). Using receipt of income support as a measure of individual deprivation, the OR was 4.80 (95% CI 3.68 to 6.25). Finally, Blair *et al*⁶⁷ carried out a case-control study which explored individual deprivation measures. The OR for SIDS in infants from social classes IV, V or never employed versus social classes I-III was 3.64 (95% CI 1.84 to 7.21). This became non-significant in a multivariate analysis. There is clearly great heterogeneity here, perhaps arising from different measures of deprivation, and no meta-analysis has been performed.

Overall, there appears to be some evidence for an association in keeping with the findings of an earlier worldwide systematic review, which found a link between deprivation and SIDS in 51 of 52 case-control and cohort studies.⁵⁸

Failure to thrive

Two analyses of the same Avon Longitudinal Study of Parents and Children cohort of children were published.^{59 60} Their definition of failure to thrive was based on the rate of growth, conditional on birth weight to allow for regression to the mean effects. They considered birth to 8 weeks and 8 weeks to 9 months separately and also jointly, leading to a large number of comparisons. No overall association with social class was found.

Wright *et al*⁶¹ also used standardised rates of growth conditional on birth weight, through their thrive index, on a small birth cohort of 923 children in Gateshead, analysed by Townsend quintile. They regressed the thrive index on quintile of deprivation and reported a significant result, but the relationship appeared to be U-shaped and so interpretation is complex. An earlier study by the same group⁴² on a different population did find a statistically significant result. The OR for failure to thrive at 12 months for most deprived versus intermediate area deprivation levels was 2.15 (95% CI 1.46 to 3.14). However, there was a wide variation in results at earlier ages, and those in more affluent areas had lower weight gain than in intermediate areas.

Wheeze

A single study⁶² suggested a potential link between both rented housing and lower maternal education (up to and beyond GCSE stage) and wheeze in infants with ORs of 1.45 (95% CI 1.28 to 1.66) and 1.12 (95% CI 1.00 to 1.27), respectively.

Diarrhoea

A single study⁶² found a significant association between diarrhoea incidence and lower maternal education (OR 1.32, 95% CI 1.19 to 1.45).

DISCUSSION

This review quantifies the evidence of inequalities in the health of infants in the UK. The socioeconomic circumstances in which an infant is conceived and born have a major effect on their early life chances and may have life course impacts.

Principal findings

The analysis demonstrates that a large number of health outcomes for UK infants are closely linked to measures of social disadvantage. There is a strong indication that the effect of social disadvantage can be measured via a range of social determinants (individual and area based) and health outcomes (eg, low birth weight, infant mortality) and found to be similar. These results are not surprising, given that prematurity, low birth weight and infant morbidity and mortality are extremely closely linked.^{39 63 64} As many as two-thirds of neonatal deaths in England and Wales in 2006 were accounted for by prematurity.⁶⁵ Furthermore, some studies show that the magnitude of effect of social disadvantage on low birth weight is durable and has not markedly changed since the early 1980s.^{28 31} The only infant health outcome which may not follow this pattern is failure to thrive, where there was no clear link based on evidence from the three cohorts identified.^{42 60 61}

For most outcomes, the association with area deprivation was quite similar to that with social class. Social class is an individual characteristic and might be expected to have a stronger association, but it is different from the factors that are combined to give an area deprivation score. Area scores are often criticised as they are ecological measures, but the results here are quite reassuring about its use in large studies. It might be expected that when a small area was used as the basis of a deprivation score, the resulting measure would better reflect the deprivation of individuals living there than if a larger area was used. However, there was no obvious variation in its effect with the size of the areas, though these were mostly quite small. Since few studies reported an analysis in which both individual and area measures were incorporated into a single analysis, it is difficult to add to the debate about the relative importance of area and individual characteristics on health outcomes.⁶⁶

Many of the meta-analyses showed heterogeneity so that results varied more than might be expected by

chance. Random effects models allow for this to some extent but identifying, and if possible modelling, the causes of heterogeneity is better practice. In most cases, there was insufficient detail available to do this. In the case of preterm births, the different definitions of the outcome could lead to heterogeneity. In other cases, there was variation in the comparisons of deprivation levels, as noted in low birth weight and social class. The effects of deprivation could also vary between areas in which studies were carried out and that might be another cause of heterogeneity.

Based on the results of this review, it can be strongly argued that no more epidemiological research needs to be carried out in the UK to address this general effect of area and individual measures of social deprivation on birth and infant mortality outcomes. Further research should seek to explore less researched areas such as any specific area effects and, most importantly, to explore the effect of interventions on those factors that are more proximal to maternal and infant health. This may help to throw light on the most appropriate times to provide support and the form(s) that such support should take.

Such studies could usefully build on recent research examining behavioural change interventions regarding the known intermediate determinants of infant health, for example, maternal nutrition and overweight,^{67–69} and smoking.^{29 67} The impact of teenage pregnancy, which is high in the UK relative to other countries, is also strongly associated with social disadvantage.⁷⁰ A recent review exploring rates of stillbirth found that maternal overweight and smoking appear to be linked to around 8000 and 2800 stillbirths annually, respectively, in the five high-income countries of Australia, Canada, the USA, the UK and the Netherlands.⁶⁷

Another potential determinant is maternal stress. There is now clear evidence of the association of maternally reported stress,^{71–73} domestic violence⁷⁴ and unintended pregnancies⁷³ with preterm birth or low birth weight. Although the evidence is yet to be definitive, there are a number of well-established hypothetical but evidence-based biological pathways for stressors leading to preterm birth, essentially related to hormonal, immune and vascular dysfunction.⁷⁵

Antenatal interventions carried out to date within high-income countries for socially disadvantaged women are few in number and of varying quality. In a Cochrane review, of support during pregnancy for women at risk of low birthweight babies, Hodnett *et al*⁷⁶ suggested that the women involved in the trials exhibit such a degree of deprivation that the support programmes evaluated are just not powerful enough to overcome the disadvantage experienced.

Nevertheless, there are promising approaches. Further exploration of WHO Baby Friendly Hospital Initiative (BFI) in properly evaluated and controlled settings appears to be warranted.⁷⁷ In a recent systematic review of 36 studies, the authors concluded that, while there is insufficient evidence to recommend any particular

programme at the moment, some targeted antenatal care programmes are promising and warrant rigorous evaluation.⁷⁸ Harden *et al*⁷⁹ concluded that there is promise for appropriately designed early childhood interventions and youth development programmes to reduce unintended teenage pregnancy, especially if such interventions take into account the known views and emotions correlated with teenage pregnancy, such as dislike of school, low expectations of life and poor material circumstances.

Given the pervasive impact of socioeconomic conditions on almost all aspects of child health and development, unless justified to the contrary, all studies of population health, including routinely collected population data, should include robust measures of socioeconomic status and maternal and infant stress levels. Intervention outcomes should also be analysed for differential effects as well as taking into account other influences, such as a tendency to low birth weight across generations, which may be independent of economic conditions.⁸⁰

Comparison with other studies

The findings of the review in terms of social disadvantage and low birth weight are remarkably similar to the findings of studies within the Republic of Ireland⁸¹ and to a recent global review of reviews carried out by the Canadian Institute of Health Economics⁸² which also concluded that the causes of low birth weight are multifactorial and closely inter-related. In addition, these results are very much in keeping with a recent review of worldwide studies published to 2007⁸³ which concluded that socioeconomic differences in low birth weight and preterm birth remain pervasive at both individual and neighbourhood levels with 93/106 studies reporting a significant association.

Conclusions and implications

It is anticipated that the results of this review will be of interest and value to policy makers and all involved directly with the care of infants and young children, as well as funders of research in this area, with two key implications emerging. First, given the clear association between child health outcomes and social disadvantage at individual and area level, governments must continue to focus on tackling social determinants, which in turn will require a cross cutting approach that includes those working in health, education, child poverty and other related policy portfolios.

Second, the absence of research on interventions through which the effects of disadvantage might be mediated is a serious shortcoming of appropriately designed interventional research.⁸² Further research is urgently required to evaluate approaches to intervention, including individual behavioural studies and studies of more upstream approaches that seek to alter the material and environmental conditions before and immediately after birth. It is imperative that policy is designed and rolled-out in a manner which allows the best possible chance of a robust assessment of outcomes and costs.^{84 85}

Acknowledgements The authors acknowledge, with thanks, contributions by Samia Addis (Cardiff Institute of Society and Health, Cardiff University) and Fiona Morgan (Support Unit for Research Evidence, Cardiff University) to early drafts of the review. We also acknowledge valuable contributions from researchers active in the field who reviewed the initial list of included studies and suggested other studies that should be screened: Sohinee Bhattacharya, Manning Donal, Ron Gray, Lucy Smith and Charlotte Wright. Lucy Smith read through and commented on the draft of the paper.

Contributors HK, HEM and ALW carried out the literature searching and selected relevant papers. FDD, HEM, MAS and ALW completed critical appraisal, data extraction and summary of the included studies. FDD carried out statistical analyses including meta-analyses of studies where feasible. The author from the Welsh Government (CR) contributed to the conception of the study, drafted and revised sections on the policy context. He had no role in the searching, analysis or interpretation of data. Each author's contribution to the paper meets the three ICMJE guidelines for authorship. All were responsible for drafting sections of the article, revising it critically for important intellectual content and approving the final version for publication.

Funding Funding from the Welsh Government was received in partial support of preparation and publication costs. No grant number is available. All authors completed the unified competing interest form. FDD does consultancy for Jacob, Medinger and Finnegan and MAS does consultancy for the Welsh Government. There are no other relationships or activities that could appear to have influenced the submitted work. The researchers operated independently from the funders and views expressed in the paper are those of the authors, based on the evidence reviewed, and are not necessarily those of the Welsh government.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All original data extraction and critical appraisal forms are available from the corresponding author. Dryad doi is 10.5061/dryad.35db6.

REFERENCES

- Marmot M, Wilkinson RG. *The Social Determinants of Health*. Oxford: Oxford University Press, 1999.
- Phelan JC, Link BG. Controlling disease and creating disparities: a fundamental cause perspective. *J Gerontol B Psychol Sci Soc Sci* 2005;60:S27–33.
- World Health Organisation. *Closing the Gap in a Generation. Health Equity through Action on the Social Determinants of Health*. Geneva: WHO, 2008. http://whqlibdoc.who.int/publications/2008/9789241563703_eng.pdf (accessed 16 Jan 2012).
- McLeroy KR, Bibeau D, Steckler A, et al. An ecological perspective on health promotion programs. *Health Educ Q* 1988;15:351–77.
- Dahlgren G, Whitehead M. *Policies and Strategies to Promote Social Equality in Health*. Stockholm: Institute of Future Studies, 1991.
- Department of Health. *Fair Society, Healthy Lives. The Marmot review. Strategic Review of Health Inequalities in England Post-2010*. London: Department of Health, 2010. <http://www.instituteofhealthequity.org/projects/fair-society-healthy-lives-the-marmot-review> (accessed 16 Jan 2012).
- Lansley A. *Healthy Lives, Healthy People: Update And Way Forward*. Cm 8134. London: The Stationery Office Limited, 2011.
- Scottish Government. *Equally Well Review 2010. Report by the Ministerial Task Force on Implementing Equally Well, the Early Years Framework and Achieving Our Potential*. Edinburgh: Scottish Government, 2010. <http://www.scotland.gov.uk/Resource/Doc/315880/0100454.pdf> (accessed 16 Jan 2012).
- Welsh Assembly Government. *Cardiff: Welsh Assembly government, 2011. Fairer Health Outcomes For All Reducing Inequities Health Strategic Action Plan Moving Agenda Forward Cardiff Welsh Assembly Government*. 2011. <http://wales.gov.uk/docs/phhs/publications/110329working2en.pdf> (accessed 16 Jan 2012).
- Keating DP, Hertzman C, eds. *Development Health and the Wealth of Nations. Social, Biological and Educational Dynamics*. New York: Guilford Press, 1999.
- Li J, Mattes E, Stanley F, et al. Social determinants of child health and wellbeing (editorial). *Health Soc Rev* 2009;18:3–11.
- Department of Health. *Healthy Lives, Brighter Futures. The Strategy for Children and Young People's Health*. London: Department of Health, 2009. http://www.dh.gov.uk/prod_consum_dh/groups/
- dh_digitalassets/documents/digitalasset/dh_094397.pdf (accessed 16 Jan 2012).
- Rajaratnam JK, Marcus JR, Flaxman AD, et al. Neonatal, postneonatal, childhood, and under-5 mortality for 187 countries, 1970–2010: a systematic analysis of progress towards millennium development goal 4. *Lancet* 2010;375:1988–2008.
- Bradshaw J, Richardson D. An index of child well-being in Europe. *Child Indicators Res* 2009;2:319–51.
- Cousens S, Blencowe H, Stanton C, et al. National, regional, and worldwide estimates of stillbirth rates in 2009 with trends since 1995: a systematic analysis. *Lancet* 2011;377:1319–30.
- Weightman AL, Addis SG, Morgan HE, et al. *Social Determinants For Child Health: A Systematic Review*. Cardiff: Welsh Assembly Government, 2008. <http://new.wales.gov.uk/dphhp/publication/research/health-gain-targets/child/childhealth-exec-e.pdf?lang=en> (accessed 16 Jan 2012).
- Carstairs V, Morris R. *Deprivation and Health in Scotland*. Aberdeen: Aberdeen University Press, 1991.
- Jarman B. Underprivileged areas: validation and distribution of scores. *BMJ* 1984;289:1587–92.
- Office for National Statistics. *The National Statistics Socio-economic Classification (NS-SEC based on the SOC2010)*. <http://www.ons.gov.uk/ons/guide-method/classifications/current-standard-classifications/soc2010/soc2010-volume-3-ns-sec-rebased-on-soc2010-user-manual/index.html> (accessed 16 Jan 2012).
- Townsend P, Phillimore P, Beattie A. *Health and Deprivation. Inequality and the North*. Beckenham, Kent: Croom Helm, 1988.
- Welsh Assembly Government. *2011 Children And Young People's Wellbeing Monitor For Wales*. Cardiff: Welsh Assembly Government, 2011. <http://wales.gov.uk/docs/caecd/research/110328cypmonitoren.pdf> (accessed 16 Jan 2012).
- Wells GA, Shea B, O'Connell D, et al. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomized Studies in Meta-analyses*. Canada: Department of Epidemiology and Community Medicine, University of Ottawa. http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm (accessed 16 Jan 2012).
- Sanderson S, Tatt ID, Higgins JP. Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography. *Int J Epidemiol* 2007;36:666–76.
- Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomised studies in meta-analysis. *Eur J Epidemiol* 2010;25:603–5.
- Bundred P, Manning D, Brewster B, et al. Social trends in singleton births and birth weight in Wirral residents, 1990–2001. *Arch Dis Child* 2003;88:F421–4.
- Dibben C, Sigala M, Macfarlane A. Area deprivation, individual factors and low birth weight in England: is there evidence of an "area effect"? *J Epidemiol Community Health* 2006;60:1053–9.
- Spencer N, Bambang S, Logan S, et al. Socioeconomic status and birth weight: comparison of an area-based measure with the registrar general's social class. *J Epidemiol Community Health* 1999;53:495–8.
- Collingwood Bakeo A, Clarke L. Risk factors for low birthweight based on birth registration and census information, England and Wales, 1981–2000. *Health Stat Q* 2006;30:15–21.
- Gray R, Bonellie SR, Chalmers J, et al. Contribution of smoking during pregnancy to inequalities in stillbirth and infant death in Scotland 1994–2003: retrospective population based study using hospital maternity records. *BMJ* 2009;339:b3754.
- Pattenden S, Dolk H, Vrijheid M. Inequalities in low birth weight: parental social class, area deprivation, and "lone mother" status. *J Epidemiol Community Health* 1999;53:355–8.
- Fairley L, Leyland AH. Social class inequalities in perinatal outcomes: Scotland 1980–2000. *J Epidemiol Community Health* 2006;60:31–6.
- Maher J, Macfarlane A. Trends in live births and birthweight by social class, marital status and mother's age, 1976–2000. *Health Stat Q* 2004;23:34–42.
- Smith LK, Draper ES, Manktelow BN, et al. Socioeconomic inequalities in survival and provision of neonatal care: population based study of very preterm infants. *BMJ* 2009;339:b4702.
- Gray R, Bonellie SR, Chalmers J, et al. Social inequalities in preterm birth in Scotland 1980–2003: findings from an area-based measure of deprivation. *Br J Obstet Gynaecol* 2008;115:82–90.
- Yuan W, Duffner AM, Chen L, et al. Analysis of preterm deliveries below 35 weeks' gestation in a tertiary referral hospital in the UK. A case-control survey. *BMC Res Notes* 2010;3:119.
- Dummer TJB, Parker L. Changing socioeconomic inequality in infant mortality in Cumbria. *Arch Dis Child* 2005;90:157–62.
- Oakley L, Maconochie N, Doyle P, et al. Multivariate analysis of infant death in England and Wales in 2005–06, with focus on

- socio-economic status and deprivation. *Health Stat Q* 2009;42:22–39.
38. Whitehead M, Drever F. Narrowing social inequalities in health? Analysis of trends in mortality among babies of lone mothers. *BMJ* 1999;318:908–12.
39. Smith LK, Manktelow BN, Draper ES, *et al.* Nature of socioeconomic inequalities in neonatal mortality: population based study. *BMJ* 2010;341:c6654.
40. Petrou S, Kupek E, Hockley C, *et al.* Social class inequalities in childhood mortality and morbidity in an English population. *Paediatr Perinat Epidemiol* 2006;20:14–23.
41. Platt MJ, Cans C, Johnson A, *et al.* Trends in cerebral palsy among infants of very low birthweight (<1500g) or born prematurely (<32 weeks) in 16 European centres: a database study. *Lancet* 2007;369:43–50.
42. Wright CM, Waterston A, Aynsley-Green A. Effect of deprivation on weight gain in infancy. *Acta Paediatr* 1994;83:357–9.
43. Gardosi J, Francis A. Social deprivation, smoking and maternal characteristics as risk factors for small-for-gestational age birthweight. *Arch Dis Child Fetal Neonatal Ed* 2010;95(Suppl X):90–111.
44. Smith LK, Draper ES, Manktelow BN, *et al.* Socioeconomic inequalities in very preterm birth rates. *Arch Dis Child Fetal Neonatal Ed* 2007;92:F11–14.
45. Sundrum R, Logan S, Wallace A, *et al.* Cerebral palsy and socioeconomic status: a retrospective cohort study. *Arch Dis Child* 2005;90:15–18.
46. Dolk H, Pattenden S, Bonellie S, *et al.* Socio-economic inequalities in cerebral palsy prevalence in the United Kingdom: a register-based study. *Paediatr Perinat Epidemiol* 2010;24:149–55.
47. Guillea ZES, Fone DL, Dunstan FD, *et al.* Social deprivation and the causes of stillbirth and infant mortality. *Arch Dis Child* 2001;84:307–10.
48. Smeeton NC, Rona RJ, Dobson P, *et al.* Assessing the determinants of stillbirths and early neonatal deaths using routinely collected data in an inner city area. *BMC Med* 2004;2:27.
49. Bambang S, Spencer NJ, Logan S, *et al.* Cause-specific perinatal death rates, birth weight and deprivation in the West Midlands, 1991–93. *Child Care Health Dev* 2000;26:73–82.
50. Manning D, Brewster B, Bundred P. Social deprivation and admission for neonatal care. *Arch Dis Child Fetal Neonatal Ed* 2005;90:F337–8.
51. Jenkins J, McCall E, Gardner E, *et al.* Socioeconomic inequalities in neonatal intensive care admission rates. *Arch Dis Child Fetal Neonatal Ed* 2009;94:F423–8.
52. Office for National Statistics. *Indices of Deprivation Across the UK*. 2011. <http://www.neighbourhood.statistics.gov.uk/dissemination/Info.do?page=analysisandguidance/analysisarticles/indices-of-deprivation.htm> (accessed 16 Jan 2012).
53. Sanderson CA, Cowden B, Hall DMB, *et al.* Is postnatal depression a risk factor for sudden infant death? *Br J Gen Pract* 2002;52:636–40.
54. Blair PS, Fleming PJ, Smith IJ, *et al.* Babies sleeping with parents: case control study of factors influencing the risk of the sudden infant death syndrome. *BMJ* 1999;319:1457–62.
55. Brooke AG, Gibson A, Tappin D, *et al.* Case-control study of sudden infant death syndrome in Scotland. 1992–5. *BMJ* 1997;314:1516–20.
56. Fleming PJ, Blair PS, Ward PM, *et al.* CESDI SUDI Research Group. Sudden infant death syndrome and social deprivation: assessing epidemiological factors after post-matching for deprivation. *Paediatr Perinat Epidemiol* 2003;17:272–80.
57. Blair PS, Sidebotham P, Evason-Coombe C, *et al.* Hazardous cosleeping environments and risk factors amenable to change: case-control study of SIDS in south west England. *BMJ* 2009;339:b3666.
58. Spencer N, Logan S. Sudden unexpected death in infancy and socioeconomic status: a systematic review. *J Epidemiol Community Health* 2004;58:366–73.
59. Blair PS, Drewett RF, Emmett PM, *et al.* Family, socioeconomic and prenatal factors associated with failure to thrive in the avon longitudinal study of parents and children (ALSPAC). *Int J Epidemiol* 2004;33:839–47.
60. Emond A, Drewett R, Blair P, *et al.* Postnatal factors associated with failure to thrive in term infants in the avon longitudinal study of parents and children. *Arch Dis Child* 2007;92:115–19.
61. Wright CM, Parkinson KN, Drewett RF. The influence of maternal socioeconomic and emotional factors on infant weight gain and weight faltering (failure to thrive): data from a prospective birth cohort. *Arch Dis Child* 2006;91:312–17.
62. Baker D, Taylor H, Henderson J. Inequality in infant morbidity: causes and consequences in England in the 1990s. ALSPAC study team. Avon longitudinal study of pregnancy and childhood. *J Epidemiol Community Health* 1998;52:451–8.
63. The Poverty Site. *Low Birthweight Babies*. 2008. United Kingdom. 2010. <http://www.poverty.org.uk/20/index.shtml> (accessed 16 Jan 2012).
64. Stevens-Simon C, Orleans M. Low birthweight prevention programs: the enigma of failure. *Birth* 1999;26:184–91.
65. Office for National Statistics. *Gestation-Specific Infant Mortality By Social And Biological Factors, 2006. Statistical Bulletin*. Newport: Office for National Statistics, 2009. <http://www.ons.gov.uk/ons/rel/child-health/gestation-specific-infant-mortality-in-england-and-wales/2006/stb-gestation-specific-infant-mortality-2006.html> (accessed 16 Jan 2012).
66. Macintyre S, Ellaway A, Cummins S. Place effects on health: how can we conceptualise, operationalise and measure them? *Soc Sci Med* 2002;55:125–39.
67. Flenady V, Koopmans L, Middleton P, *et al.* Major risk factors for stillbirth in high-income countries: a systematic review and meta-analysis. *Lancet* 2011;377:1331–40.
68. Haggarty P, Campbell DM, Duthie S, *et al.* Diet and deprivation in pregnancy. *Br J Nutr* 2009;102:1487–97.
69. McDonald SD, Han Z, Mulla S. Overweight and obesity in mothers and risk of preterm birth and low birth weight infants: systematic review and metaanalysis. *BMJ* 2010;341:c3428. <http://www.bmj.com/content/341/bmj.c3428.full> (accessed 16 Jan 2012).
70. Department for Education and Skills. *Teenage Pregnancy: Accelerating The Strategy to 2010*. London: Department for Education and Skills, 2006. <http://media.education.gov.uk/assets/files/pdf/t/teenage%20pregnancy%20%20%20accelerating%20the%20strategy%20to%202010.pdf> (accessed 16 Jan 2012).
71. Savitz DA, Pastore LM. Causes of prematurity. In: McCormick MC, Siegel JE, eds. *Prenatal Care: Effectiveness and Implementation*. Cambridge: Cambridge University Press, 1999.
72. Hobel CJ, Goldstein A, Barrett ES. Psychosocial stress and pregnancy outcomes. *Clin Obstet Gynecol* 2008;51:333–48.
73. Messer LC, Dole N, Kaufman JS, *et al.* Pregnancy intendedness, maternal psychosocial factors and preterm birth. *Matern Child Health J* 2005;9:403–12.
74. Shah PS, Shah J. Maternal exposure to domestic violence and pregnancy and birth outcomes: a systematic review and meta-analysis. *J Womens Health* 2010;19:2017–31.
75. Kramer MR, Hogue CR. What causes racial disparities in very preterm birth? A biosocial perspective. *Epidemiol Rev* 2009;31:84–98.
76. Hodnett ED, Fredericks S, Weston J. Support during pregnancy for women at increased risk of low birthweight babies. *Cochrane Database Syst Rev* 2010;(6):CD000198. http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD000198/pdf_fs.html (accessed 16 Jan 2012).
77. Spijby H, McCormick F, Wallace L, *et al.* A systematic review of education and evidence-based practice interventions with health professionals and breast feeding counsellors on duration of breast feeding. *Midwifery* 2009;25:50–61.
78. Hollowell J, Oakley L, Kurinczuk JJ, *et al.* The effectiveness of antenatal care programmes to reduce infant mortality and preterm birth in socially disadvantaged and vulnerable women in high-income countries: a systematic review. *BMC Pregnancy Childbirth* 2011;11:13.
79. Harden A, Brunton G, Fletcher A, *et al.* Teenage pregnancy and social disadvantage: systematic review integrating controlled trials and qualitative studies. *BMJ* 2009;339:b4254.
80. Collins J, Rankin K, David R. Low birth weight across generations: the effect of economic environment. *Matern Child Health J* 2011;15:438–45.
81. McAvooy H, Sturley J, Burke S, *et al.* *Unequal at Birth—Inequalities in the Occurrence Of Low Birthweight Babies in Ireland*. Ireland: Institute of Public Health in Ireland, 2006. http://www.publichealth.ie/files/file/Unequal_at_Birth.pdf (accessed 16 Jan 2012).
82. Ohlsson A, Shah P. *Determinants And Prevention Of Low Birth Weight: A Synopsis Of The Evidence*. Alberta, Canada: Institute of Health Economics, 2008. <http://www.ihe.ca/documents/IHE%20Report%20LowBirthWeight%20final.pdf> (accessed 16 Jan 2012).
83. Blumenshine P, Egarter S, Barclay CJ, *et al.* Socioeconomic disparities in adverse birth outcomes: a systematic review. *Am J Prev Med* 2010;39:263–72.
84. Macintyre S. Good intentions and received wisdom are not good enough: the need for controlled trials in public health. *J Epidemiol Community Health* 2011;65:564–7.
85. House of Commons Health Committee. *Health Inequalities. Third Report of Session 2008–09. Volume 1. Report, Together with Formal Minutes HC 286-I*. London: The Stationary Office Limited, 2009. <http://www.publications.parliament.uk/pa/cm200809/cmselect/cmhealth/286/286.pdf> (accessed 16 Jan 2012).