Correction

Allemani C, Rachet B, Weir HK, *et al.* Colorectal cancer survival in the USA and Europe: a CONCORD high-resolution study. *BMJ Open* 2013;3:e003055. Table 2 of this article was published incorrectly. The correct table is below.

Table 2 Advanced stage, resection with curative intent, 30-day postoperative mortality and proportion of patients with information on stage: colorectal cancer, Europe and the USA, 1996–1998

EUROPE	Registry	All cases			Resected with curative intent*							
		N	Advanced stage†		N	%	Deaths within 30 days		Staged			
			N	%			N	%	Colon		Rectum	
		N			N		N		N	%	N	%
<i>European registries‡</i>		8762	2535	29	6584	75	248	4	3895	95	2374	95
	Northern Europe	523	134	26	385	74	16	4	192	84	142	90
	Western Europe§	2497	609	24	2092	84	24	6	1299	93	646	92
	Southern Europe¶	3818	1131	30	2912	76	152	5	1748	97	1081	97
	Eastern Europe	1924	661	34	1195	62	56	5	656	98	505	97
<i>US registries</i>		3337	676	20	2832	85	124	4	2039	97	677	93
	California	495	112	23	415	84	15	4	294	96	102	93
	Colorado	548	113	21	468	85	18	4	335	95	109	93
	Illinois	505	112	22	422	84	21	5	320	97	85	93
	Louisiana	511	105	21	431	84	26	6	315	100	111	97
	New York	492	80	16	411	84	22	5	287	95	102	94
	Rhode Island	418	78	19	369	88	9	2	268	99	93	94
	South Carolina	368	76	21	316	86	13	4	220	96	75	87
<i>Total</i>		12 099										

†All metastatic cases, plus unresected cases for which no stage data were available.

*Curative intent: surgery not specified as palliative or tumour entirely resected.

‡Northern Europe: Finland; Western Europe: France (Côte d'Or) and the Netherlands (North East Netherlands); Southern Europe: Italy (Genova, Ragusa and Varese), Slovenia and Spain (Granada, Navarra and Tarragona); Eastern Europe: Estonia, Poland (Cracow and Kielce) and Slovakia.

§Data for North East Netherlands (1936) are not included in the proportion of deaths within 30 days of surgery for Western Europe because the date of surgery was not available.

¶Data for Ragusa (424) are not included in the percentages of Dukes' stage for Southern Europe.

BMJ Open 2013;3:e003055corr1. doi:10.1136/bmjopen-2013-003055corr1