BMJ Open Multi-institution case-control and cohort study of risk factors for the development and mortality of *Clostridium difficile* infections in Japan

Masahiko Takahashi,¹ Nobuaki Mori,² Seiji Bito²

To cite: Takahashi M, Mori N, Bito S. Multi-institution case-control and cohort study of risk factors for the development and mortality of *Clostridium difficile* infections in Japan. *BMJ Open* 2014;**4**:e005665. doi:10.1136/bmjopen-2014-005665

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2014-005665>).

Received 9 May 2014

Revised 5 August 2014

Accepted 8 August 2014

CrossMark

¹Department of Gastroenterology, National Hospital Organization Tokyo Medical Center, Tokyo, Japan

²Department of General Internal Medicine, National Hospital Organization Tokyo Medical Center, Tokyo, Japan

Correspondence to

Dr Masahiko Takahashi; matakaha@ntmc.hosp.go.jp

ABSTRACT

Objective: To examine risk factors for *Clostridium difficile* infection (CDI) morbidity and mortality in Japan.

Design: Multimethod investigation including a case-control study and cohort study.

Setting: 47 participating facilities of the National Hospital Organization (NHO).

Participants: 1026 patients with CDI and 878 patients in the control group over the age of 18 years admitted to the subject NHO facilities from November 2010 to October 2011.

Main outcome measures: In a case-control study, we identify risk factors for CDI development. Next, in a cohort study, we identify risk factors for all-cause mortality within 30 days following CDI onset.

Results: A total of 1026 cases of CDI meeting the definitions of this investigation were identified, encompassing 878 patients at 42 of the 47 subject facilities. In the case-control study, we identified, compared with no antibiotics use, use of first-generation and second-generation cephem antibiotics (OR 1.44; 95% CI 1.10 to 1.87), use of third-generation and fourth-generation cephem antibiotics (OR 1.86; 95% CI 1.48 to 2.33) and use of carbapenem antibiotics (OR 1.87; 95% CI 1.44 to 2.42) the risk factors for CDI development. However, use of penicillin was not identified as a risk factor. In the cohort study, sufficient data for analysis was available for 924 CDI cases; 102 of them (11.0%) resulted in death within 30 days of CDI onset. Compared with no anti-CDI drug use, use of vancomycin was associated with reduced risk of mortality (OR 0.43; 95% CI 0.25 to 0.75) whereas metronidazole was not.

Conclusions: The findings mirror those of previous studies from Europe and North America, identifying the administration of broad-spectrum antibiotics as a risk factor for CDI development. The use of vancomycin is associated with a decreased risk of mortality.

INTRODUCTION

Clostridium difficile is the main causative pathogen of antibiotic-associated colitis. Since 2000, outbreaks of BI/NAP1/027

Strengths and limitations of this study

- This study is the first large-scale nationwide multicentre *Clostridium difficile* infection (CDI) investigation in Japan.
- Most of the epidemiological data of CDI have been limited to North America and Europe. Our data plays a role of completion of the missing data in Asia.
- Use of β -lactam antibiotics, except penicillin, was the risk factor for CDI development in the first Japanese large-scale investigation. Appropriate antibiotic use is necessary in order to control the incidence of CDI.
- Vancomycin administration for CDI was associated with decreased risk of mortality. Although the cost-effective treatment of CDI may necessitate the appropriate use of the less-expensive metronidazole, vancomycin should be administered in cases expected to become severe or life-threatening.
- The limitation of this study is the low number of registered CDI cases from quite a few participants and the existence of many confounding factors.

strain *C. difficile* infections (CDI) have been reported in North American and European hospitals and elder care facilities. The numbers of patients with CDI as well as severe and intractable cases have increased simultaneously. Consequently, epidemiological surveillance systems have been set up in several countries. However, very few countries have implemented such national-level measures.

CDI epidemiological studies in Japan to date have been based on scattered data from individual medical facilities. Consequently, the phenomenon of CDI in Japan is not sufficiently understood.^{1–9} Previous studies report that antibiotic administration is the largest risk factor for CDI development. Other risk factors include advanced age and proton pump inhibitor use.^{10 11} CDI

mortality rates differ depending on the presence or absence of an outbreak as well as the relevant definitions of epidemiological surveillance. Furthermore, it is especially difficult to objectively determine precise CDI-related mortality rates because of factors such as underlying patient conditions.¹²

This report documents a case-control study of CDI in Japan based on data from the National Hospital Organization (NHO), which is Japan's largest group of hospitals and includes facilities located nationwide. In addition, a cohort investigation of mortality among CDI cases was conducted.

MATERIALS AND METHODS

Research design

This multicenter study is a collaborative effort of the 47 facilities that met our facility standards from among the 143 NHO facilities in Japan. The study was planned as a part of the NHO's 'National Hospital Organization Multi-Center Clinical Research for Evidence-Based Medicine' project. This study was conducted with the approval of the Central Ethics Committee of the NHO. The CDI group in this study included, in principal, all newly diagnosed CDI cases among patients hospitalised from 1 November 2010 to 31 October 2011; cases were registered continuously.

In the case-control study of CDI development, CDI cases newly diagnosed during the investigation period were registered in the CDI group; meanwhile, age-matched, sex-matched and underlying disease-matched patients in the same facilities were registered to the control group. In addition, a prospective cohort study of CDI group patients who died within 30 days of CDI development was conducted. This investigation is a multimethod study using standard case-control and cohort study designs.

Definition of CDI

CDI was defined as the presence of any gastrointestinal symptoms accompanied by a clinical suspicion of CDI as well as a positive result for *C. difficile* toxins from rapid stool testing or *C. difficile* isolation from stool cultures or both. Final determinations were carried out by the attending physician or the facility's infection control team.

Enzyme immunoassay testing kits for *C. difficile* toxins A and B were used as the rapid testing method (Immunocard CD toxin A&B, Meridian Bioscience Inc, Cincinnati, Ohio, USA; *C. Diff* Quik Chek, Alere Medical Co., Ltd, Tokyo, Japan; Tox A/B Quik Chek, Nissui Pharmaceutical Co., Ltd, Tokyo, Japan; X/pect Toxin A/B, Kanto Chemical Co., Ltd, Tokyo, Japan). Cycloserine-cefoxitin mannitol agar (Nissuiure-to CCMA baichi EX, Nissui Pharmaceutical Co., Ltd, Tokyo, Japan), cycloserine-cefoxitin fructose agar (CCFA baichi, Becton, Dickinson and Company Co., Ltd, Tokyo, Japan; Poamedhia CCFA kairyoubaiichi, Eiken Chemical Co., Ltd, Tokyo, Japan), and brucella HK agar

(RS) (brucella HK agar (RS), Kyokuto Pharmaceutical Industrial Co., Ltd, Tokyo, Japan) were used in the *C. difficile* isolation cultures.

Case-control study of CDI development

No additional information besides age, sex and date of diagnosis was gathered when new patients were registered in the CDI group. After the end of the study registration period, additional patient clinical data were gathered, including clinical department, underlying diseases, dates of hospital admittance and discharge and medical treatments administered for ≥ 3 days between admittance and CDI development. Recorded medical treatments included disruption of feeding, parenteral nutrition, enteral feeding, surgery with general anaesthetics, cancer drugs, antibiotics (excluding external-use antibiotics) and proton pump inhibitors (oral or intravenous). We also collected data regarding the use of intravenous antibiotics including penicillins, first-generation and second-generation cepheems, third-generation and fourth-generation cepheems, carbapenems, fluoroquinolones, clindamycin/lincomycin, anti-methicillin-resistant *Staphylococcus aureus* (MRSA) drugs and antifungal drugs, and others. Finally, we collected data regarding the use of oral antibiotics including cepheems, fluoroquinolones and others.

The control group was divided into three subgroups according to age: ≤ 74 , 75–84 and ≥ 85 years. Control patients were selected from among patients at the same facilities who did not contract CDI and were matched to the patients with CDI with respect to age, sex, underlying disease and hospital stays of ≥ 5 days within the same month as a counterpart's CDI diagnosis. The control group cases were selected regardless of gastrointestinal symptoms such as diarrhoea. We strove to ensure that the CDI and control groups were as matched as possible. The same data were collected from both groups. The control patients were registered and relevant patient data were gathered after the end of the CDI group study registration period.

Cohort study on mortality among patients with CDI

The prospective cohort study of registered CDI group patients from the case-control study examined all-cause mortality within 30 days as the primary outcome. If the registered patients were discharged within 30 days, clinical outcomes were not investigated after discharge in this study. The following data were collected: whether the underlying disease was infectious and whether comorbidities were related to malignant tumours (ie, gastrointestinal, respiratory, blood/lymph, gynaecological, urological, or other tumours including cancers of the ear, nose and throat), diabetes, renal failure, heart failure, respiratory failure or cirrhosis. In the cohort study, the definitions of severe complications were gastrointestinal perforations, toxic megacolon and CDI-related surgeries. Severe complications and the all-cause in-hospital mortality of patients within 30 days of CDI development were recorded. We

also considered patient nutritional status including whether the patient was subjected to parenteral nutrition or enteral feeding as well as serum albumin levels measured within 30 days prior to CDI development (ie, ≥ 3.5 , 2.7–3.4 or ≤ 2.6 g/dL). In addition, we examined CDI treatment factors including whether antibiotic use was halted, probiotic use and the type of anti-CDI drugs used (ie, vancomycin and metronidazole). All patient data for the cohort investigation were collected after the end of the registration period.

Data management and statistical analysis

All input data were verified by a designated study data manager. Data from each facility were entered directly into a web-based case report form and subsequently encrypted for security. The data management centre was responsible for confirming any missing data and directly inquiring of the relevant facilities as necessary.

During the case–control phase of the study, CDI development was treated as the outcome and ORs were calculated from bivariate analysis comparing the use of different types of antibiotics as outcome causes. For each type of antibiotic, those used for ≥ 3 days were designated ‘used’ while all others were designated ‘unused.’ A dummy variable regression was subsequently performed. Statistical significance in the bivariate analysis was tested by the χ^2 test. Logistic regression analysis was performed using the individual patient characteristics and other assumed confounding variables as independent variables. The 95% CIs for each variable were used to determine the relationships between the various predictive variables and outcomes.

The clinical outcome of mortality within 30 days was set as the dependent variable and the relationships among the underlying diseases, nutritional status, probiotic use and types of anti-CDI drugs used were subjected to bivariate and multivariate analyses. Like the case–control phase, bivariate analysis was conducted using the χ^2 test and the multivariate analysis was conducted using logistic regression. The significance level for all analyses was set at $p < 0.05$. We used IBM SPSS Statistics V.20 for statistical analysis.

Ethics committee approval and informed consent

This study was conducted with the approval of the Central Ethics Committee of the NHO. In principle, individual patients who met the inclusion criteria were not given direct explanations of the study and no direct consent was sought. Information about the study was made public through postings on facility notice boards and webpages. Patients and their representative agents had the right to refuse study participation.

RESULTS

Participating facilities

Among the 47 facilities, a total of 1026 CDI cases were registered at 42 facilities throughout Japan, from

Hokkaido in the north to Okinawa in the south. No CDI cases were recorded at the remaining 5 participating facilities, more than 280 patient beds (table 1).

Patient grouping

A total of 1026 CDI cases that met the study definitions were recorded at the various institutions. We were unable to collect clinical records regarding medical treatments for 1 case; therefore, this case was excluded from the case–control study, and the remaining 1025 cases were analysed. A total of 962 patients (93.9%) developed CDI within 48 h after hospital admittance. The control group comprised 878 patients who were selected from 41 of the 42 facilities. In the cohort study, we analysed the data from 924 of the 1025 CDI group patients, excluding 101 patients with no available recent serum albumin level data (ie, within 30 days prior to CDI development (figure 1).

Case–control study of CDI development

The mean ages of the CDI and control groups were 75.8 and 75.4 years, respectively. The majority of the participants were of advanced age: 64% and 62.5% of the CDI and control group patients were aged ≥ 75 years, respectively. No significant differences were identified between the CDI and control groups in the univariate analysis of age distribution, sex differences or underlying disease (table 2). Among the medical treatments administered before CDI development, the following were significantly more prevalent in the CDI group than the control group: disruption of feeding (48.6% vs 30.4%), parenteral nutrition (24.7% vs 10.3%) and enteral feeding (24.8% vs 9.1%). Antibiotics were used prior to CDI development in 85.8% of cases. The use of all types of intravenous antibiotics was significantly more prevalent in the CDI group. No significant differences were identified between the two groups with respect to oral antibiotic use. Meanwhile, in the univariate analysis, proton pump inhibitor use was significantly more prevalent in the CDI group than the control group (40.3% vs 31.2%).

We used logistic regression analysis to determine the risk factors for CDI development. The following medical treatments prior to CDI development were identified as significant risk factors in comparison to the control group: disruption of feeding (OR 1.31; 95% CI 1.05 to 1.64), parenteral nutrition (OR 1.63; 95% CI 1.21 to 2.20) and enteral feeding (OR 2.16; 95% CI 1.60 to 2.92). The following intravenous antibiotics were also identified as statistically significant risk factors for CDI development: first-generation and second-generation cepheems (OR 1.44; 95% CI 1.10 to 1.87), third-generation and fourth-generation cepheems (OR 1.86; 95% CI 1.48 to 2.33) and carbapenems (OR 1.87; 95% CI 1.44 to 2.42). However, penicillin (OR 1.04; 95% CI 0.82 to 1.33), fluoroquinolones (OR 1.16; 95% CI 0.74 to 1.83),

Table 1 Number of registered cases of CDI and characteristics of hospitals included in the surveillance of CDI in the NHO (from November 2010 through October 2011)

Region	Number of patient beds	Number of patient days	Number of patients registered		30-day all-cause mortality in CDI group	Laboratory tests used	
			CDI group	Control group		EIA for toxins A and B	Culture
Hokkaido, Tohoku	698	208 388	55	55	3 (5%)	+	+
	500	150 603	42	32	1 (2%)	+	+
	310	82 687	28	19	2 (7%)		+
	310	72 144	17	12	2 (12%)	+	+
	220	76 539	1	1	0 (0%)	+	+
Kanto, Koshinetsu	780	238 420	124	121	15 (12%)	+	+
	455	151 622	36	36	3 (8%)	+	
	560	158 921	35	30	4 (11%)	+	+
	243	60 155	34	34	6 (18%)	+	+
	350	109 025	22	22	4 (18%)	+	+
	500	159 432	15	14	1 (7%)	+	
	510	166 668	4	4	0 (0%)	+	
	380	109 482	3	2	0 (0%)	+	+
	455	132 483	3	1	0 (0%)	+	
	429	104 802	0	0	– (–)	+	
Tokai, Hokuriku, Kinki	430	195 209	42	26	10 (24%)	+	+
	280	56 475	0	0	– (–)	+	
	316	103 677	24	22	1 (4%)	+	
	220	47 354	23	23	1 (4%)	+	+
	600	191 041	20	20	3 (15%)	+	
	494	70 455	15	15	6 (40%)	+	+
	520	145 299	13	9	1 (8%)	+	
	500	142 409	6	6	1 (17%)	+	
	180	55 721	3	3	1 (33%)	+	
	346	118 014	2	2	0 (0%)	+	
Chugoku, Shikoku	370	94 722	0	0	– (–)	+	
	388	99 728	54	49	5 (9%)	+	+
	700	211 595	49	48	4 (8%)	+	+
	506	119 356	33	8	1 (3%)	+	+
	400	122 846	30	30	5 (17%)	+	
	401	108 303	26	0	2 (8%)	+	+
	250	80 558	21	21	0 (0%)	+	
	424	128 868	12	10	0 (0%)	+	
	365	125 645	10	10	3 (30%)	+	+
	300	87 061	0	0	– (–)		+
Kyushu, Okinawa	459	66 454	0	0	– (–)	+	
	424	137 827	46	22	5 (11%)	+	
	702	239 448	38	37	1 (3%)	+	
	190	54 038	33	31	9 (27%)	+	
	550	189 417	27	26	3 (11%)	+	
	285	58 185	25	25	3 (12%)	+	
	500	140 371	24	23	2 (8%)	+	
	300	90 457	14	14	4 (29%)	+	
	320	103 315	6	5	1 (17%)	+	+
	280	79 580	4	4	2 (50%)	+	
Total	366	112 906	4	4	0 (0%)	+	
	368	89 195	3	2	2 (67%)	+	
Total	19 486	5 592 077	1026	878	117 (11%)	45	20

CDI, *Clostridium difficile* infection; EIA, enzyme immunoassay; NHO, National Hospital Organization.

clindamycin/lincomycin (OR 1.35; 95% CI 0.81 to 2.26) and proton pump inhibitor use (OR 1.17; 95% CI 0.95 to 1.44) were not identified as risk factors.

Cohort study on mortality among patients with CDI

The cohort study examined mortality among the 924 patients from the 1025 CDI group patients in the

Figure 1 Study populations for the analysis of patients with *Clostridium difficile* infection (CDI) and controls.

case-control study for whom serum albumin level data before CDI development were available.

Among the 924 patients, 102 (11.0%) died within 30 days of developing CDI. Among those cases, the cause of death was attributed to CDI in 11 cases (1.2%). Of 11 patients, a patient had gastrointestinal perforation, another patient had CDI-related surgery and the others were not reported as severe complications. Toxic megacolon was reported in two patients, however, they had not died within 30 days of CDI development. The mean age of the 102 patients who died during the study was 80.1 ± 8.3 years. Patients ≥ 75 years old were especially prevalent in this subgroup, accounting for 77.5% (79/102) of the cases.

Among the 714 cases in which CDI was treated directly, recurrence within 30 days was observed in 34 cases (4.8%).

The univariate analysis indicated that comorbidities of heart and respiratory failure were significantly more prevalent among patients with CDI. In addition, lower serum albumin levels were significantly associated with mortality. Among CDI treatments, mortality was significantly lower among cases in which probiotics were administered.

A logistic regression analysis of the 102 cases in which the patients died within 30 days of CDI development was performed to identify the factors associated with the risk of mortality. Compared to patients ≤ 74 years old, the OR of mortality among patients aged 75–84 years was 2.08 (95% CI 1.19 to 3.62). Among underlying diseases, heart failure (OR 2.12; 95% CI 1.26 to 3.55) and

respiratory failure (OR 1.98; 95% CI 1.19 to 3.32) were identified as risk factors for mortality within 30 days of CDI development. Regarding nutritional status, neither parenteral nutrition nor enteral nutrition was identified as a risk factor for mortality. However, low serum albumin level (ie, ≤ 2.6 g/dL) was identified as a significant risk factor for mortality (OR 3.50; 95% CI 1.33 to 9.22). Among CDI treatments, probiotic use (OR 0.66; 95% CI 0.42 to 1.04) was not identified as a risk factor for mortality. However, compared to cases in which no anti-CDI drugs were administered, vancomycin administration yielded an OR of 0.43 (95% CI 0.25 to 0.75), indicating a significantly lowered risk of mortality in the CDI group. Meanwhile, no such lowered mortality was observed in cases treated with metronidazole (OR 0.85; 95% CI 0.48 to 1.51; table 3).

DISCUSSION

This is the first large-scale clinical study of CDI in Japan. This study examined 1026 cases of CDI recorded over 1 year at the nationwide facilities of Japan's largest hospital group. The findings of this investigation are similar to those reported in previous studies conducted in Europe, North America and Australia with respect to the identification of several risk factors for CDI development including age, severity of the underlying condition, artificial feeding and mortality.^{11 13 14} Antibiotic use is a known risk factor for CDI development.¹⁵ The present case-control study confirms that intravenous cepheems and carbapenems are important risk factors. Some

Table 2 Univariate and multivariate analyses of CDI development-related risk factors

Characteristics	CDI group Per cent	Control group Per cent	Univariate analysis p Value	Multivariate analysis	
				OR (95% CI)	p Value
All	(1025)	(878)	–	–	–
Age (years)					
≤74	36.0 (369)	37.5 (329)	0.67	Ref.	–
75–84	37.0 (379)	37.2 (327)		1.02 (0.81 to 1.28)	0.88
≥85	27.0 (277)	25.3 (222)		1.09 (0.84 to 1.41)	0.52
Sex					
Women	43.0 (441)	42.6 (374)	0.85	1.11 (0.91 to 1.36)	0.28
Underlying disease					
Respiratory infections	15.8 (162)	17.5 (154)	0.14	–	–
Other infectious conditions	16.9 (173)	14.2 (125)		–	–
Gastrointestinal conditions	8.1 (83)	9.0 (79)		–	–
Malignant tumours	22.6 (232)	24.3 (213)		–	–
Cardiovascular conditions	7.7 (79)	9.8 (86)		–	–
Other conditions	28.9 (296)	25.2 (221)		–	–
Medical treatment prior to CDI development					
Disruption of feeding	48.6 (498)	30.4 (267)	<0.001	1.31 (1.05 to 1.64)	<0.05
Parenteral nutrition	24.7 (253)	10.3 (90)	<0.001	1.63 (1.21 to 2.20)	<0.01
Enteral feeding	24.8 (254)	9.1 (80)	<0.001	2.16 (1.60 to 2.92)	<0.001
Surgery with general anaesthetic	18.2 (187)	15.6 (137)	0.14	0.89 (0.67 to 1.18)	0.41
Cancer drugs	11.3 (116)	14.2 (125)	0.06	0.86 (0.62 to 1.18)	0.35
Antibiotics use	85.8 (879)	66.5 (584)	<0.001	–	–
Intravenous					
Penicillins	27.6 (283)	21.0 (184)	<0.01	1.04 (0.82 to 1.33)	0.75
First-generation/second-generation cepheims	22.7 (233)	15.6 (137)	<0.001	1.44 (1.10 to 1.87)	<0.01
Third-generation/fourth-generation cepheims	35.2 (361)	19.9 (175)	<0.001	1.86 (1.48 to 2.33)	<0.001
Carbapenems	31.8 (326)	15.0 (132)	<0.001	1.87 (1.44 to 2.42)	<0.001
Fluoroquinolones	7.5 (77)	4.0 (35)	<0.01	1.16 (0.74 to 1.83)	0.52
Clindamycin/lincomycin	6.5 (67)	2.8 (25)	<0.001	1.35 (0.81 to 2.26)	0.25
MRSA drugs	10.7 (110)	4.3 (38)	<0.001	1.10 (0.71 to 1.72)	0.66
Antifungal drugs	6.9 (71)	3.2 (28)	<0.001	1.01 (0.60 to 1.70)	0.96
Others (aminoglycosides, monobactam, etc)	8.5 (87)	5.9 (52)	<0.05	1.19 (0.80 to 1.77)	0.39
Oral					
Cepheims	5.6 (57)	4.4 (39)	0.29	1.49 (0.95 to 2.32)	0.08
Fluoroquinolones	14.5 (149)	11.5 (101)	0.06	1.11 (0.82 to 1.51)	0.49
Others (macrolides, penicillins, etc)	14.0 (144)	13.9 (122)	0.95	0.84 (0.63 to 1.13)	0.26
Proton pump inhibitors	40.3 (413)	31.2 (274)	<0.001	1.17 (0.95 to 1.44)	0.14

CDI, *Clostridium difficile* infection; MRSA, methicillin-resistant *Staphylococcus aureus*.

studies report a low risk of CDI development owing to intravenous penicillin administration.^{16 17} Concordantly, penicillin use was not identified as a risk factor in the present study. The proton pump inhibitor use was discussed as a risk factor for CDI development in the previous studies.^{18–20} In the present logistic regression analysis, it was not identified as a risk factor.

In this study, 11.0% of patients with CDI died within 30 days. In comparison, higher 30-day mortality rates have been reported in previous outbreaks: 24.8% in the ribotype 027 strain outbreak in Canada and 36.7% in an examination of a single intensive care unit in the USA.^{21 22} However, reports of non-outbreak conditions indicate mortality rates of 13%, similar to the findings of the present study.²³ Some reports state that the CDI-associated mortality rate has increased 2.5-fold, possibly indicating that CDI cases are more severe and

contribute more significantly to mortality than previously thought.^{12 23} The mortality rate of patients with CDI is reported to increase with age.²⁴ Concordantly, the present study also found a significantly elevated risk of death in patients ≥75 years old.

The findings of this study indicate that the mortality risk of patients with CDI was not reduced as a result of metronidazole treatment but was reduced with vancomycin treatment, corroborating the existing recommendation.²⁵ It is worth noting that metronidazole is less expensive than vancomycin, making it economically advantageous. A patient's condition must be carefully evaluated when selecting anti-CDI drugs. In particular, for patients in the present study who had conditions associated with a greater mortality risk, including advanced age (ie, ≥75 years), heart or respiratory failure or malnutrition as determined by low serum albumin levels, the

Table 3 Univariate and multivariate analyses of all-cause mortality in patients with *Clostridium difficile* infection (CDI)

Characteristics	All-cause mortality rate Per cent	Univariate analysis p Value	Multivariate analysis	
			OR (95% CI)	p Value
All	11.0 (102/924)	–	–	–
Age (years)				
≤74	7.1 (23/326)	<0.05	Ref.	<0.05
75–84	13.3 (47/353)		2.08 (1.19 to 3.62)	
≥85	13.1 (32/245)		1.86 (0.98 to 3.55)	
Sex				
Men	12.2 (64/524)	0.21	Ref.	0.29
Women	9.5 (38/400)		0.78 (0.49 to 1.24)	
Underlying disease				
Non-infectious	10.3 (64/619)	0.37	Ref.	0.97
Infectious	12.5 (38/305)		0.99 (0.60 to 1.62)	
Comorbidities				
Malignant tumours				
Not present	10.6 (67/630)	0.57	Ref.	0.09
Present	11.9 (35/294)		1.54 (0.94 to 2.53)	
Diabetes				
Not present	11.6 (89/765)	0.27	Ref.	0.29
Present	8.2 (13/159)		0.71 (0.37 to 1.35)	
Renal failure				
Not present	10.7 (84/784)	0.46	Ref.	0.73
Present	12.9 (18/140)		0.90 (0.49 to 1.65)	
Heart failure				
Not present	9.3 (70/756)	<0.01	Ref.	<0.01
Present	19.0 (32/168)		2.12 (1.26 to 3.55)	
Respiratory failure				
Not present	9.2 (69/754)	<0.001	Ref.	<0.01
Present	19.4 (33/170)		1.98 (1.19 to 3.32)	
Cirrhosis				
Not present	11.2 (100/895)	0.76	Ref.	0.53
Present	6.9 (2/29)		0.61 (0.13 to 2.83)	
Indicators of nutritional status				
Parenteral nutrition or enteral feeding				
Not present	9.4 (53/563)	0.05	Ref.	0.53
Present	13.6 (49/361)		1.16 (0.73 to 1.84)	
Serum albumin (g/dL)				
≥3.5	4.0 (5/124)	<0.001	Ref.	0.39
2.7–3.4	7.2 (27/376)		1.55 (0.57 to 4.21)	
≤2.6	16.5 (70/424)		3.50 (1.33 to 9.22)	
CDI treatments				
Cessation of antibiotics				
Not present	12.5 (65/519)	0.11	Ref.	0.26
Present	9.1 (37/405)		0.77 (0.48 to 1.22)	
Probiotics (for intestine treatment)				
Not present	13.8 (52/378)	<0.05	Ref.	0.08
Present	9.2 (50/546)		0.66 (0.42 to 1.04)	
Anti-CDI drugs				
Not present	15.2 (32/210)	<0.05	Ref.	<0.01
Vancomycin alone	7.4 (32/433)		0.43 (0.25 to 0.75)	
Metronidazole alone	13.5 (32/237)		0.85 (0.48 to 1.51)	
Vancomycin and metronidazole	13.6 (6/44)		0.75 (0.27 to 2.08)	

use of vancomycin was expected to reduce the mortality. The recurrence rate was low (4.8%) in this study compared to the previous studies.^{11 26} We did not investigate those patients who developed CDI after 30 days nor did we investigate those patients who were discharged, even if

within 30 days of CDI development. Therefore, the recurrence rate might be underestimated.

Regardless, this study also has several methodological limitations. The most salient being the low number of registered CDI cases from quite a few participants. In

the definition of CDI, the times of diarrhoea were not investigated. Another limitation of the case-control study phase is the existence of many confounding factors. In particular, probiotic use, which was recently discussed to be correlated with CDI prevention, was not included in the predictive model of this study.^{10 11 27}

When interpreting the findings of this study, it is necessary to consider the influence of confounding factors that were not included in the analytical models. Regarding antibiotic use, the present analyses included independent explanatory variables for each antibiotic. However, actual antibiotic use is more complicated. Therefore, it is difficult to clearly determine the roles of individual antibiotics as risk factors for CDI development. Concerning matching process, we tried to adopt 1 to 1 pair sampling matched with sex, age group and main diagnosis. Some hospitals could not find appropriate control samples well matched with case samples. So the total number of the control group was less than that of the case sample. In addition, although data for the control group were analysed during the entire study period until hospital discharge, only data from the period prior to CDI development were analysed in the CDI group. Therefore, the risks might be underestimated, because the control group had a longer period of exposure risk than the CDI group did. Confounding factors that were not included in the present analyses also represent a limitation of the cohort study phase. Furthermore, issues of data quality among the facilities affect all aspects of this study. More than 40 different facilities participated in this study. While some facilities registered nearly all of their patients with CDI, other facilities registered smaller proportions of patients. Only *C. difficile* culture but not toxin test was used for the laboratory test in two facilities. Finally, there might have been differences with regard to individual researchers' understanding of the outcome definitions.

In order to ensure appropriate antibiotic use and control the incidence of CDI, it is important to create institutional measures such as infection control teams. The cost-effective treatment of CDI may necessitate the appropriate use of less-expensive metronidazole. However, in cases expected to become severe or life-threatening, the more expensive drug vancomycin should be administered. CDI is one of many issues concerning medicine and medical treatment costs. Accordingly, further and more proactive research into CDI epidemiology is needed.

Acknowledgements The authors would like to express their sincere gratitude to Dr Haru Kato and the Department of Bacteriology II, National Institute of Infectious Diseases, Tokyo, Japan for their expert advice regarding CDI and the provision of CDI training to the participating facilities. They also wish to thank the participating institutions in the CD-NHO Study Group for their collaboration with data and sample collection: Hisaji Oshima (NHO Tokyo Medical Center); Hiroshi Miki (NHO Sendai Medical Center); Keisei Shimoe (NHO Fukuyama Medical Center); Harumi Tominaga (NHO Kure Medical Center); Toyomitsu Sawai and Eisuke Sasaki (NHO Ureshino Medical Center); Shie Nishijima and Naoko Maeda (NHO Shizuoka Medical Center); Masaru Amishima (NHO Hokkaido Medical Center); Miki Odawara (NHO Kyushu

Medical Center); Mitsuhiro Kamimura (NHO National Disaster Medical Center); Hideaki Nagai (NHO Tokyo National Hospital); Kiyoshi Furuta (NHO Matsumoto Medical Center, Matsumoto Hospital); Tohru Yamanaka (NHO Kumamoto Minami Hospital); Ikuko Mizouchi (NHO Minimi-Okayama Medical Center); Yutaka Sato (NHO Kanmon Medical Center); Keita Ato and Hiroki Saito (NHO Asahikawa Medical Center); Yoshio Haga (NHO Kumamoto Medical Center); Isao Murakami (NHO Higashihiroshima Medical Center); Takeshi Yamaryo (NHO Nagasaki Kawatana Medical Center); Hiroyuki Akiyama and Yukino Yoshikura (NHO Minami Wakayama Medical Center); Akiko Muratake (NHO Beppu Medical Center); Masato Hasegawa (NHO Higashi-Ohmi General Medical Center); Isamu Kamimaki (NHO Saitama National Hospital); Tomoaki Kosyoubu (NHO Yonago Medical Center); Takao Odagaki (NHO Kyoto Medical Center); Nozomu Iwashiro (NHO Hakodate National Hospital); Hiroyasu Ishida (NHO Mito Medical Center); Hiroshi Komatsu (NHO Maizuru Medical Center); Kaoru Nakama (NHO Oita Medical Center); Yoshiko Yamamoto (NHO Osaka Minami Medical Center); Yoshihito Iwahara (NHO Kochi National Hospital); Fumiko Okino (NHO Yamaguchi-Ube Medical Center); Daisuke Higuchi (NHO Okinawa National Hospital); Kazuhiro Satonaka (NHO Hyogo-Chuo National Hospital); Takayoshi Soga and Haruko Ideguchi (NHO Yokohama Medical Center); Mayuko Watanabe (NHO Kagoshima Medical Center); Kozaburo Hiramatsu (NHO Nagasaki National Hospital); Mitsugu Saito (NHO Awara National Hospital); Morio Sawamura (NHO Nishigunma National Hospital); Satoru Kaneda (NHO Chiba Medical Center); Kenji Okada (NHO Fukuoka National Hospital); Katsuhiro Suzuki (NHO Kinki-Chuo Chest Medical Center); Tetsuko Chiba and Keiji Chida (NHO Iwate National Hospital); Akihiko Tamura (NHO Tochigi Medical Center); Shunji Matsuda (NHO Ehime Medical Center); Takaya Maruyama (NHO Mie National Hospital); Shigeaki Kimura (NHO Tokushima National Hospital); Shin Oguri (NHO Minami Kyoto National Hospital).

Contributors MT conceived the idea for the study, designed the study, developed the protocol, was responsible for study management and data collection, interpreted the findings and drafted the paper. NM contributed to the data analysis and interpretation of the findings and drafted the paper. SB designed the study, developed the protocol, performed the data analysis and interpreted the findings, and drafted the paper. All authors have read and approved the final manuscript.

Funding This study was supported by a grant from the National Hospital Organization (multicenter clinical studies for evidence-based medicine).

Competing interests None.

Ethics approval The Central Ethics Committee of the NHO.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Honda H, Yamazaki A, Sato Y, *et al.* Incidence and mortality associated with *Clostridium difficile* infection at a Japanese tertiary care center. *Anaerobe* 2014;25:5–10.
2. Kato H, Ito Y, van den Berg RJ, *et al.* First isolation of *Clostridium difficile* 027 in Japan. *Euro Surveill* 2007;12:E070111.3.
3. Kato H, Kato N, Watanabe K, *et al.* Analysis of *Clostridium difficile* isolates from nosocomial outbreaks at three hospitals in diverse areas of Japan. *J Clin Microbiol* 2001;39:1391–5.
4. Tagashira Y, Kato H, Senoh M, *et al.* Two cases of fulminant colitis due to binary toxin-positive *Clostridium difficile* that are not PCR ribotype 027 or type 078. *J Med Microbiol* 2013;62:1486–9.
5. Sawabe E, Kato H, Osawa K, *et al.* Molecular analysis of *Clostridium difficile* at a university teaching hospital in Japan: a shift in the predominant type over a five-year period. *Eur J Clin Microbiol Infect Dis* 2007;26:695–703.
6. Iwashima Y, Nakamura A, Kato H, *et al.* A retrospective study of the epidemiology of *Clostridium difficile* infection at a University Hospital

- in Japan: genotypic features of the isolates and clinical characteristics of the patients. *J Infect Chemother* 2010;16:329–33.
7. Kato H, Ito Y, Akahane T, *et al.* Typing of *Clostridium difficile* isolates endemic in Japan by sequencing of *slpA* and its application to direct typing. *J Med Microbiol* 2010;59:556–62.
 8. Collins DA, Hawkey PM, Riley TV. Epidemiology of *Clostridium difficile* infection in Asia. *Antimicrob Resist Infect Control* 2013;2:21.
 9. Nakamura I, Yamaguchi T, Tsukimori A, *et al.* Fulminant colitis from *Clostridium difficile* infection, the epidemic strain ribotype 027, in Japan. *J Infect Chemother* 2014;20:380–3.
 10. Rupnik M, Wilcox MH, Gerding DN. *Clostridium difficile* infection: new developments in epidemiology and pathogenesis. *Nat Rev Microbiol* 2009;7:526–36.
 11. Cohen SH, Gerding DN, Johnson S, *et al.* Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the society for healthcare epidemiology of America (SHEA) and the infectious diseases society of America (IDSA). *Infect Control Hosp Epidemiol* 2010;31:431–55.
 12. Mitchell BG, Gardner A. Mortality and *Clostridium difficile* infection: a review. *Antimicrob Resist Infect Control* 2012;1:20.
 13. Freeman J, Bauer MP, Baines SD, *et al.* The changing epidemiology of *Clostridium difficile* infections. *Clin Microbiol Rev* 2010;23:529–49.
 14. Bliss DZ, Johnson S, Savik K, *et al.* Acquisition of *Clostridium difficile* and *Clostridium difficile*-associated diarrhea in hospitalized patients receiving tube feeding. *Ann Intern Med* 1998;129:1012–19.
 15. Owens RC Jr, Donskey CJ, Gaynes RP, *et al.* Antimicrobial-associated risk factors for *Clostridium difficile* infection. *Clin Infect Dis* 2008;46(Suppl 1):S19–31.
 16. Brown KA, Khanafer N, Daneman N, *et al.* Meta-analysis of antibiotics and the risk of community-associated *Clostridium difficile* infection. *Antimicrob Agents Chemother* 2013;57:2326–32.
 17. Deshpande A, Pasupuleti V, Thota P, *et al.* Community-associated *Clostridium difficile* infection and antibiotics: a meta-analysis. *J Antimicrob Chemother* 2013;68:1951–61.
 18. Dial S, Delaney JA, Barkun AN, *et al.* Use of gastric acid-suppressive agents and the risk of community-acquired *Clostridium difficile*-associated disease. *JAMA* 2005;294:2989–95.
 19. Kwok CS, Arthur AK, Anibueze CI, *et al.* Risk of *Clostridium difficile* infection with acid suppressing drugs and antibiotics: meta-analysis. *Am J Gastroenterol* 2012;107:1011–19.
 20. Tleyjeh IM, Bin Abdulhak AA, Riaz M, *et al.* Association between proton pump inhibitor therapy and *Clostridium difficile* infection: a contemporary systematic review and meta-analysis. *PLoS ONE* 2012;7:e50836.
 21. Loo VG, Poirier L, Miller MA, *et al.* A predominantly clonal multi-institutional outbreak of *Clostridium difficile*-associated diarrhea with high morbidity and mortality. *N Engl J Med* 2005;353:2442–9.
 22. Kenneally C, Rosini JM, Skrupky LP, *et al.* Analysis of 30-day mortality for *Clostridium difficile*-associated disease in the ICU setting. *Chest* 2007;132:418–24.
 23. Hensgens MP, Goorhuis A, Dekkers OM, *et al.* All-cause and disease-specific mortality in hospitalized patients with *Clostridium difficile* infection: a multicenter cohort study. *Clin Infect Dis* 2013;56:1108–16.
 24. Bloomfield MG, Sherwin JC, Gkrania-Klotsas E. Risk factors for mortality in *Clostridium difficile* infection in the general hospital population: a systematic review. *J Hosp Infect* 2012;82:1–12.
 25. Zar FA, Bakkanagari SR, Moorthi KM, *et al.* A comparison of vancomycin and metronidazole for the treatment of *Clostridium difficile*-associated diarrhea, stratified by disease severity. *Clin Infect Dis* 2007;45:302–7.
 26. Otete EH, Ahankari AS, Jones H, *et al.* Parameters for the mathematical modelling of *Clostridium difficile* acquisition and transmission: a systematic review. *PLoS ONE* 2013;8:e84224.
 27. Johnston BC, Ma SS, Goldenberg JZ, *et al.* Probiotics for the prevention of *Clostridium difficile*-associated diarrhea: a systematic review and meta-analysis. *Ann Intern Med* 2012;157:878–88.