

# BMJ Open

## The pathway to diagnosis of type 1 diabetes in children: a questionnaire study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-006470
Article Type:	Research
Date Submitted by the Author:	26-Aug-2014
Complete List of Authors:	Usher-Smith, Juliet; The Primary Care Unit, Institute of Public Health Thompson, Matthew; Oxford University, Department of Primary Care Health Sciences Zhu, Hannah; Cambridge University Hospitals NHS Foundation Trust, Sharp, Stephen; University of Cambridge, Walter, Fiona; University of Cambridge, Dept of Public Health and Primary Care
<b>Primary Subject Heading</b>:	Diabetes and endocrinology
Secondary Subject Heading:	Diagnostics, General practice / Family practice
Keywords:	Paediatric endocrinology < DIABETES & ENDOCRINOLOGY, PRIMARY CARE, Paediatric A&E and ambulatory care < PAEDIATRICS

SCHOLARONE™  
Manuscripts

The pathway to diagnosis of type 1 diabetes in children: a questionnaire study

Juliet A Usher-Smith<sup>1</sup>, Matthew J Thompson<sup>2</sup>, Hannah Zhu<sup>3</sup>, Stephen J Sharp<sup>4</sup>, Fiona M Walter<sup>5</sup>

<sup>1</sup> Clinical Lecturer in General Practice, The Primary Care Unit, University of Cambridge CB1 8RN

<sup>2</sup> Professor, Department of Family Medicine, University of Washington, Seattle, WA 98195 USA

<sup>3</sup> Foundation Year 2 Doctor, Cambridge University Hospitals NHS Foundation Trust, CB2 0QQ

<sup>4</sup> Senior Statistician, MRC Epidemiology Unit, University of Cambridge, Institute of Metabolic Science, Cambridge, CB2 0QQ

<sup>5</sup> GP & Clinician Scientist, The Primary Care Unit, University of Cambridge CB1 8RN

Correspondence to: J Usher-Smith [jau20@medschl.cam.ac.uk](mailto:jau20@medschl.cam.ac.uk)

ABSTRACT

**Objective** To explore the pathway to diagnosis of type 1 diabetes (T1D) in children

**Design** Questionnaire completed by parents

**Participants** Parents of children aged 1 month to 16 years diagnosed with T1D within the previous three months

**Setting** Children and parents from 11 hospitals within the East of England.

**Results** 88/164 (54%) of invited families returned the questionnaire. Children had mean±SD age of 9.41±4.5 years. 35 (39.8%) presented with DKA at diagnosis. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%) and all children presented with at least one of those symptoms. The time from symptom onset to diagnosis ranged from 2 to 315 days (median 25 days). Most of this was the appraisal interval from symptom onset until perceiving the need to seek medical advice. Access to healthcare was good but one in five children presenting to primary care were not diagnosed at first encounter, most commonly due to waiting for fasting blood tests or alternative diagnoses. Children diagnosed at first consultation had a shorter duration of symptoms (p=0.022) and children whose parents suspected the diagnosis were 1.3 times more likely (RR 1.3, 95% CI 1.02-1.67) to be diagnosed at first consultation.

**Conclusions** Children present with the known symptoms of T1D but there is considerable scope to improve the diagnostic pathway. Future interventions targeted at primary care physicians may help reduce delays in a small number of children but targeting parents in the appraisal interval is likely to have greater effect.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study uses a questionnaire developed from a previous interview study to explore the diagnostic pathway of children with newly diagnosed T1D
- It uses the Model of Pathway to Treatment as a framework to allow analysis of the factors acting at different stages in the pathway
- The inclusion of a calendar with key events in the questionnaires and use of free text responses for internal validation and checking of prompted responses reduced bias but the data was necessarily collected retrospectively and so subject to recall and framing bias

## INTRODUCTION

Type 1 diabetes (T1D) is one of the commonest endocrine diseases in children, with an estimated 65,000 children world-wide under 15 years developing the disease each year and the incidence increasing at a rate of 3% per year<sup>1,2</sup>. Despite this, in a typical primary care practice a child with new onset T1D will be seen only about once every two years<sup>3</sup> and the symptoms are often non-specific in the early stages. Distinguishing the rare child with T1D from the large number with similar symptoms and minor undifferentiated illness is therefore challenging for both primary care physicians and families. The mean duration of symptoms prior to diagnosis is over two weeks; a significant number of children experience delay in diagnosis or misdiagnosis<sup>4</sup> with only one in five diagnosed at first encounter<sup>5-9</sup>; and worldwide up to 80% present in diabetic ketoacidosis (DKA)<sup>10</sup> which has both immediate life-threatening complications and is associated with poorer long term diabetic control<sup>11-13</sup>.

Whilst several studies have highlighted these difficulties in making the diagnosis and the features associated with diabetic ketoacidosis at diagnosis<sup>4,9,14</sup>, few have explored the period between symptom onset and diagnosis. Our recent qualitative interview study of parents and General Practitioners (GPs) of children newly diagnosed with T1D suggested that the longest component in the diagnostic pathway is the time between onset of symptoms and the decision to seek medical help (known as the appraisal interval)<sup>15</sup>. The early symptoms are subtle, and even with some knowledge of T1D it took many parents several weeks of a complex decision making process and often a physical trigger, such as weight loss or vomiting, to decide to consult a healthcare professional. Once the decision to seek help had been made almost all children were seen immediately and diagnoses were mostly prompt and managed appropriately. Parents continued to play a key role during the diagnostic interval however, with many having already made or suspected the diagnosis themselves, and several feeling that their GP did not take their concerns seriously.

This study builds on this earlier work by using a questionnaire developed from the interview findings to further explore the pathway to diagnosis of T1D in children. By using a structured questionnaire to survey a larger number of families we aimed to quantify the symptoms and their time course prior to diagnosis, the triggers and barriers to seeking help, the influence of parental prior knowledge of diabetes, and the role of healthcare services.

## METHODS

### Design

A questionnaire about the pathway from first symptom(s) to diagnosis was completed by the parent(s)/guardian(s)/step-parents (hereafter referred to as parents) of children aged 1 month to 16 years diagnosed with T1D within the previous three months.

### Recruitment

Children and parents were identified and recruited via the paediatric diabetes specialist nurses and research nurses at 11 hospitals within the East of England Diabetes Children and Young People's Network. Parents of all children aged 1 month to 16 years diagnosed with T1D diagnosed within the previous 3 months at participating hospitals were eligible for inclusion unless their clinical team felt that this was not appropriate. Parents who failed to respond within one month were sent a reminder letter with a further copy of the questionnaire. Recruitment began at each site between February 2013 and April 2013, and continued across all sites until January 2014.

The clinical or research teams at all sites collected data on the age and gender of each child diagnosed during the study period and whether they had DKA at diagnosis. Each hospital used a

slightly different definition of DKA but all included either pH < 7.3, bicarbonate < 15 mmol/L or clinical acidosis.

**The questionnaire**

The questionnaire was developed from the findings of our previous qualitative study of parents and children recently diagnosed with T1D<sup>15</sup>. It was first reviewed by an expert panel comprising paediatric diabetes consultants, a paediatric diabetes research nurse and primary care researchers, and then piloted with parents of four children recently diagnosed with T1D. In addition to their specific feedback, parents were asked to talk aloud whilst completing the questionnaire and then interviewed after completion to ensure face validity. Based on feedback from the parents, the questionnaire was revised.

The final questionnaire included 5 sections. The first included questions about the child’s age, gender, postcode, ethnic background, family history of diabetes, any medically trained family members, the number of children in the household and whether the parents knew about the symptoms of diabetes prior to their child being diagnosed. The second section asked about the symptoms the children had experienced with yes/no responses for 14 symptoms and space to add the date they noticed the symptoms, what they thought the symptoms were due to at the time and how much it concerned them. The third section focused on help-seeking and asked where parents had looked for information, who they spoke to and then details on when and how they had sought medical advice. It also asked them to describe their main concern at their first appointment and whether they had considered diabetes. Parents were also asked in this section about factors contributing to their decision to seek medical advice sooner or later. The fourth section asked about the diagnosis, including whether it was made at their first appointment with a healthcare professional and, if not, how many subsequent consultations they had, and the investigations that were done before diagnosis. The final section then asked parents if they felt there was anything that prolonged them finding out their child had diabetes and had further space for free text comments.

**Analysis**

Data from the questionnaires were entered into a database and then double checked by a second researcher. Socioeconomic status was computed using postcode and the English indices of deprivation 2010 available online<sup>16</sup>. The presence of DKA at diagnosis was obtained from hospital records rather than self-report. Walter et al’s Model of Pathways to Treatment<sup>17,18</sup> provided a theoretic model of the intervals that occur prior to a diagnosis. This model divides the pathway to diagnosis into two intervals prior to presentation to healthcare about a symptom (the appraisal interval from the onset of symptoms to perceiving a reason to discuss symptoms with a healthcare professional, and the help-seeking interval from that decision until presentation to a healthcare professional), and then the diagnostic interval from first presentation to a healthcare professional until diagnosis. The help-seeking interval was further sub-divided into the behavioural interval (the time between perceiving the reason to discuss the symptoms with a healthcare professional to making the decision to seek help) and the scheduling interval (the time between making the decision to seek help and the first consultation)<sup>19</sup>. Intervals were calculated from responses to the questionnaire. Where dates were incomplete we applied midpoint rules to estimate the actual date<sup>20</sup>. In cases where the responses in free text differed from the dates entered as numbers, the free text was assumed to be correct, and where there was uncertainty the researchers met to agree consensus.

Characteristics (age, gender, presence of DKA) were compared between children whose parents had and had not returned a questionnaire using a t-test for age and chi-squared test for gender and presence of DKA. All further analyses used only data from returned questionnaires. The frequency of the 14 symptoms was compared between those with and without DKA using a chi-squared test.

Cox regression was used to estimate the association between various factors and the hazard of diagnosis; if a factor was associated with an increased hazard (i.e. hazard ratio greater than 1), this implied that that factor was associated with a shorter time to diagnosis, and vice versa. Time to diagnosis was from the date of the earliest symptom to the date of diagnosis, and the factors assessed were age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the parents suspected T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis. A similar approach was used to assess factors associated with the length of the appraisal and help-seeking intervals (with the end of the interval being defined as the “event” in the Cox model), but only the first six variables in the list above were considered, as the others do not relate to those time intervals. The Schoenfeld residuals test was used to assess the proportional hazards (PH) assumption for each covariate in each model. Whether parents suspected the diagnosis of T1D did not meet the PH assumption for the total diagnostic interval and so the Cox regression model was stratified by that variable. Logistic regression was used to estimate the association between the same factors and presence of DKA at diagnosis. All analyses were performed using STATA version 12.

Free text responses were grouped into similar categories and coded. Where individual free text responses contained several comments, these were each coded individually.

## RESULTS

A total of 172 children were diagnosed with T1D in the 11 hospitals during the study period. Of those, 8 families were not invited to take part in the study: 5 lived outside the hospital catchment area; 1 emigrated the week after diagnosis; and the clinical team felt it was not appropriate to include 2. From the remaining 164 families invited to take part in the study, 88 (54%) completed and returned the questionnaire. There were no significant differences in the proportion presenting in DKA ( $p=0.27$ ), mean age ( $p=0.77$ ) or gender ( $p=0.77$ ) between children of responders and non-responders.

Table 1 shows the characteristics of the 88 children and families included in the study. The mean age was  $9.41 \pm 4.5$  years, 49 (55.7%) were male and 35 (39.8%) presented with DKA at diagnosis. The majority (90.9%) were white and as a group they were generally from less deprived areas.

One child was excluded from subsequent analysis as the parents were intermittently testing the child’s blood glucose prior to diagnosis in the absence of symptoms as they had an older child with T1D. 87 children are therefore included in the analysis that follows.

## Symptoms

Table 2 shows the frequency and duration of the 14 symptoms that were specifically asked about in the questionnaire. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%). Most symptoms were present for a median of between 13 and 17 days. Faster breathing and vomiting both had much shorter median (IQR) durations of 0.5 (0-7.5) and 2.5 (1.5-5.5) days respectively and weight loss, vomiting and faster breathing were significantly more frequent in those children who presented in DKA. All the children had at least one of the 4 main symptoms (polydipsia, polyuria or nocturia, weight loss or tiredness), 97.7% had 2 or more, 79.3% 3 or more and over half (50.6%) had all 4 symptoms.

A very small number of parents mentioned symptoms other than those listed in the questionnaire, these included constipation (9), headaches (3), thrush (3), blurred vision (2), dry skin (2) and different smelling urine (1).



**Diagnostic intervals**

Table 3 shows the mean  $\pm$  SD and median (IQR) for the diagnostic intervals. Additional details on the diagnostic intervals for different subgroups are shown in Appendix Table 1. The total diagnostic interval ranged from 2 to 315 days with a median (IQR) of 25 days (14-50). In unadjusted Cox regression analysis (data not shown) the time to diagnosis was significantly shorter for children diagnosed at first appointment compared to a subsequent appointment ( $p=0.046$ ) and for those seen in secondary care rather than primary care ( $p=0.01$ ). No evidence of associations with time to diagnosis was found for age, gender, family history of T1D, deprivation, prior knowledge of symptoms or DKA at diagnosis. Adjusting for age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis (Figure 1a), the association between whether the diagnosis of T1D was made at the first or subsequent appointments and total diagnostic interval remained statistically significant ( $p=0.022$ ).

**The appraisal interval**

The appraisal interval was the longest of all the intervals in the pathway for all but 3 of the families with a mean  $\pm$  SD of  $39.1 \pm 49$  days and median (IQR) 20 (8-42) days. During this period nearly two thirds (64%) of parents discussed the symptoms with family members, 40% with friends and 41% looked on the internet. Only 16% spoke to the child's nursery, school or playgroup and very few (6%) looked for information in books. Over half of parents (49, 56%) reported being aware of some symptoms of T1D in children prior to their child's diagnosis: 40 (45%) were aware of increased thirst, 24 (27%) of polyuria, 17 (19%) of weight loss and 13 (15%) tiredness.

Cox-regression analysis (Figure 1b) showed no significant associations between parent/child characteristics and the appraisal interval.

Analysis of the free text showed that most parents found explanations for their child's symptoms (Table 4). For example, polydipsia was attributed most commonly to hot weather (19/58, 33%) or infection (13/58, 22%), polyuria and nocturia were frequently explained by drinking more (29/47, 62% and 26/40, 65%) and tiredness was thought to be school related (12/44, 27%) or secondary to infection (5/44, 12%) or nocturia (4/44, 10%).

The majority of parents (61/87, 70%) additionally reported that they had suspected diabetes before their first consultation with a healthcare professional. When asked what had made them suspect diabetes, the most common reason given was that they knew the symptoms (22/59, 37%), especially thirst (12/59, 20%). Others cited information from the internet (12/59, 20%) or having a family history of diabetes (11/59, 19%).

**The help-seeking interval**

24 (28%) children were seen on the same day their parents first thought about seeking medical advice and 64 (74%) within 5 days. Most of this time was the behavioural interval (mean  $\pm$  SD  $2.1 \pm 3.7$  days, median (IQR) 0 (0-3) days) rather than the scheduling interval (mean  $\pm$  SD  $1.1 \pm 2.6$  days, median (IQR) 0 (0-1) days).

Cox-regression analysis (Figure 1c) showed no significant associations between parent/child characteristics and the help-seeking interval.

The most common reasons that parents cited for seeking medical advice sooner rather than later (Table 5) were that the symptoms were not getting better or were getting worse, wanting reassurance or concern something serious was wrong. This was also reflected in the free text responses where 22% of parents noted that worsening or persistent symptoms was the reason they decided to seek help. In general, fewer parents reported factors that led to them seeking medical advice later. Of those that did, the most common reason for waiting was hope that the symptoms would go away (51.6%) but 29.8% felt difficulty getting an appointment contributed and 27.6% and 25.2% were worried about wasting the GPs time or that the GP would not take them seriously respectively.

### ***The diagnostic interval***

The diagnostic interval was the shortest of the intervals with a mean  $\pm$  SD of 5 days  $\pm$  34.8 and median 0 (IQR) (0-0) days. 69 (78%) of children were diagnosed at first consultation. Cox regression was not possible given the high number of children with a diagnostic interval of zero. However, children whose parents suspected the diagnosis were more likely (unadjusted RR 1.30, 1.02-1.67) to be diagnosed at first consultation ( $p=0.046$ ) than those in whom there was no suspicion. All children (10) who were seen first in secondary care were diagnosed at first consultation compared to 76.6% (59/77) of those seen first in primary care, but this difference was not statistically significant ( $p=0.114$ ). None of the variables considered were significantly associated with risk of DKA (Figure 2).

Further details from the questionnaires were available from 14 of the 18 children who were not diagnosed at first encounter with primary care. Of these, 6 had fasting glucose blood tests arranged by the GP and 4 were given alternative diagnoses (urine infection, viral infection, tonsillitis, puberty) and diagnosed at a second appointment. Two children were diagnosed with psychological problems: In one case the child's mother had seen the GP alone to discuss her child's 'obsessive drinking' and was advised to see the school counsellor, and in the second the GP apparently felt the symptoms were psychological and the child was diagnosed in the emergency department four consultations later. One other family had already done a finger prick glucose test at home which was high but the GP did not trust the result and asked the child to come back later in the day with a urine sample. In the final case, the child's mother had spoken to a health visitor and suggested diabetes but was told 'no, not unless the child is lifeless'. The mother took the child to the GP 12 days later and the diagnosis was made at that consultation.

## **DISCUSSION**

### **Principal findings**

This study shows that all children with new onset T1D present with one, and 98% present with two, of the four main symptoms of diabetes (polydipsia, polyuria, weight loss and tiredness). Moreover, over half have had symptoms for over three weeks before diagnosis. Most of that time is the appraisal interval during which parents found alternative explanations for the symptoms, discussed the symptoms with family and friends and looked on the internet for information. Once they made the decision to seek advice, access to healthcare was generally not difficult with 28% consulting with a healthcare professional on the same day. However, when asked about factors contributing to their decision to seek help, nearly a third of parents felt that difficulty getting an appointment contributed to them waiting to seek help and over a quarter felt that worry about wasting the doctor's time influenced their decision. This suggests that even if access is not difficult, it is perceived as such.

Once parents had sought help, one in five children were then not diagnosed at their first consultation with a healthcare professional, mainly due to being given an alternative diagnosis,

most commonly infection, or waiting for further investigations. Diagnosis at first consultation was associated with a shorter total diagnostic interval and children were more likely to be diagnosed at first consultation when their parents suspected the diagnosis of T1D. The association between diagnosis at first consultation and total diagnostic interval may simply reflect the additional time between consultations, or it may be due to biological differences causing some children to develop symptoms more slowly which are then more difficult for both parents and primary care physicians to recognise.

**Strengths and weaknesses**

By using a questionnaire developed from a previous interview study<sup>15</sup> and the Model of Pathway to Treatment<sup>17,18</sup> as a framework for analysis, this study provides in-depth insights into the diagnostic pathway of children with newly diagnosed T1D and allows factors acting at different stages in the pathway to be explored.

The main weakness is that the data was necessarily collected retrospectively and so subject to recall and framing bias. Parents have multiple contacts with different healthcare professionals in the period immediately following diagnosis and so their responses to the questionnaire reflect a post-hoc rationalisation of events framed by those subsequent encounters and increased knowledge since the diagnosis. The inclusion of a calendar with key events in the questionnaires minimised the error in recall of dates, and the free text responses allowed internal validation and checking of prompted responses. Despite these efforts, we still only have the parents' perspective on the pathway and were not able to confirm the number of healthcare contacts, diagnostic tests or the parental reports of missed opportunities for diagnosis.

**Comparison with existing literature**

The median duration of symptoms prior to diagnosis was 13-17 days for the nine most frequent symptoms, with a mean of 30-50 days. This is longer than previous studies relying on retrospective review of medical records<sup>21-24</sup> but similar to studies which have used a checklist to identify subtle symptoms<sup>25</sup> or asked parents soon after diagnosis<sup>14,15</sup>. The wide range (a few days to over six months) has been described previously<sup>15,22,24</sup> and highlights the heterogeneous nature of the disease.

The frequency of individual symptoms we report is also similar to previous studies<sup>14,15,21,23,26</sup>. Additionally we showed that all the children had at least one of 4 symptoms (polydipsia, polyuria, weight loss and fatigue) and over half (50.6%) had all four. Consistent with the known course of the disease and previous studies, vomiting<sup>5,23,25</sup>, weight loss<sup>14,26,27</sup>, and dyspnoea<sup>23</sup> were more common in those children who presented in DKA.

This is the first quantitative study to compare the time periods during the pathway to diagnosis of T1D in children. The finding that most of the total diagnostic interval was the appraisal interval is consistent with a previous qualitative study<sup>15</sup> and the free text analysis confirms that during that time the parents find alternative explanations for the symptoms initially and make use of a social network of extended family, friends and work colleagues, or the internet<sup>15,28,29</sup>. That children were more likely to be diagnosed at their first encounter with a healthcare professional when their parents suspected diabetes prior to that consultation may also reflect the findings of previous qualitative work in which a number of parents prompted the GP to consider T1D and pushed for investigations<sup>15</sup>. However, whilst parental suspicion of T1D has also been shown to be associated with a reduced risk of DKA in a parental survey<sup>14</sup>, in that study the incidence of DKA at presentation was no different whether or not the parents discussed their concerns with the healthcare professional, suggesting other factors may be contributing. The absence of an effect of parental



prior knowledge of diabetes either on the total diagnostic interval or the risk of DKA further highlights the complexities around the role of knowledge on help-seeking behaviour.

The finding that parents worry about wasting the doctor's time has also been shown in previous qualitative studies in children<sup>30,31</sup> and in studies of help-seeking behaviour for adults with symptoms of cancer in the UK<sup>32,33</sup> and so it may reflect a particular British trait rather than be specific to T1D or children.

### **Implications for clinicians and policymakers**

Clinicians should remain alert to the possibility of T1D in all children presenting with one or more symptoms of polyuria, polydipsia, weight loss and tiredness – as almost all children have at least two of these. Interventions targeted at increasing public awareness, such as the 4 T's campaign launched by Diabetes UK to raise awareness of the four most common symptoms of T1D (Toilet, Thirsty, Tired and Thinner)<sup>34</sup>, should continue to focus on these established symptoms.

As most of the time between symptom onset and diagnosis is the appraisal interval, the greatest benefit is likely to be seen from interventions directed towards parents and their social network, probably via the internet. Despite ongoing government pressure for better access to primary care, improving access is unlikely to have much impact on the pathway. Instead efforts should be made to address the perception that access is difficult and the general concern in the UK about wasting healthcare professional time, particularly for children with acute or sub-acute health concerns.

Additionally, although the diagnostic interval itself was generally short, one in five children presenting to primary care were not diagnosed at first consultation. Similar numbers have been reported in a recent survey in the UK which found that 24% were not diagnosed at first contact with a healthcare professional<sup>14</sup>, and studies in the USA, Canada and Poland noted between 14 and 35% of children had more than one consultation before diagnosis<sup>7,8,35–37</sup>. As in those studies, the most common reasons for not being diagnosed at first encounter was either being given an alternative diagnosis, most commonly infection, or waiting for further investigations. In this study 33% of those not diagnosed at first consultation were waiting for fasting glucose tests and in other studies the number waiting for further investigations is as high as 46%<sup>7,14</sup>. This suggests that healthcare professionals may have considered a diagnosis of T1D but either lack ready access to rapid tests to confirm or exclude the diagnosis, or are reluctant to use existing tests in children<sup>15</sup>. Access to point of care urine and finger-prick testing and the use of those tests should be routine management for all children presenting with one or more of the four main symptoms of diabetes. The increased use of point of care testing in Emergency departments may also explain why all children seen in secondary care were diagnosed at their first consultation. Whilst educational interventions aimed at primary care physicians may help a small number of children not currently diagnosed at first encounter, finding ways to overcome barriers to point-of-care tests in primary care may be more effective and this approach may also improve the diagnosis of other serious illnesses in children and adults.

### **Unanswered questions and future research**

Whilst this study contributes to our understanding of the pathway to diagnosis and the stages at which this may be improved, the findings are unable to explain the large variability in the overall duration of the pathway to diagnosis and why some children develop DKA within a few weeks whilst others can be symptomatic for up to six months before requiring treatment. Further studies are, therefore, needed into the natural course and biology of the disease to better understand these variations. The findings also highlight the need for continuing research into the presentation of serious but rare conditions in primary care and the best ways to improve diagnosis of these conditions.

**ACKNOWLEDGEMENTS**

We thank all the parents who kindly gave up their time and shared their personal stories with us to either help develop the questionnaire or take part in the study. We also thank the Paediatric diabetes clinical teams at the 11 hospitals, particularly Jon Hassler-Hurst (Ipswich), Holly Roper (Norfolk and Norwich), Elissa Harwood (Cambridgeshire community services), Nazia Bhatti and Claire Pesterfield (Addenbrooke’s hospital NHS foundation trust), Julia Harding and Jacky Plumb (West Suffolk), Philippa Corbishley and Laura Ardrey (QE2 and Lister), Suzie Williams (SEPT), Mandy Stevenson (Harlow), Claire Gibbs and Allyson Davison (James Paget) and Meeta Patel (Luton and Dunstable). We are also grateful to the Diabetes Research Network and the East of England Children and Young People’s Diabetes Network and network manager Kate Wilson for support, James Brimicombe for advice on data management, Dr Carlo Acerini for comments on the questionnaire and final manuscript and support through his role as Chair of the East of England Children and Young People’s Diabetes Network, Professor Jon Emery for comments on the questionnaire, and Professor David Dunger for his support and encouragement throughout the study and comments on the final manuscript.

**Contributors**

JUS, MT, FMW and SJS were involved in the design of the study and all authors were involved in analysis of the data. JUS and FMW developed the questionnaire. JUS wrote the first draft of the manuscript and all authors reviewed and edited the manuscript.

**Funding**

The study was funded by the Royal College of General Practitioners Scientific Foundation Board (SFB-2011-15). JUS was supported by a National Institute of Health Research (NIHR) Academic Clinical Fellowship and subsequently Clinical Lectureship, and FMW by an NIHR Clinician Scientist award. SJS was supported by the Medical Research Council [www.mrc.ac.uk](http://www.mrc.ac.uk) [Unit Programme number MC\_UU\_12015/1]. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

All researchers were independent of the funding body (although FMW subsequently became a member of the Royal College of General Practitioners Scientific Foundation Board), and the study sponsors and funder had no role in study design; data collection, analysis and interpretation of data; in the writing of the report; or decision to submit the article for publication.

**Ethics approval**

The study obtained ethical approval from the East of England Hertfordshire REC (reference number 12/EE/0390).

**Data sharing**

The questionnaire is available from the corresponding author on request.

**Competing Interests**

All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare that (1) JUS, MT, HZ, SJS and FMW have no support from or relationships with companies that might have an interest in the submitted work in the previous 3 years; (2) their spouses, partners, or children have no financial relationships that may be relevant to the submitted work; and (3) JUS, MT, HZ, SJS and FMW have no non-financial interests that may be relevant to the submitted work.

The corresponding author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in BMJ editions and any other BMJ PGL products and sublicences to exploit all subsidiary rights, as set out in our licence (<http://resources.bmj.com/bmj/authors/checklists-forms/licence-for-publication>).

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

The corresponding author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

## REFERENCES

- 1 DIAMOND Project Group. Incidence and trends of childhood Type 1 diabetes worldwide 1990–1999. *Diabet Med* 2006;**23**:857–66.
- 2 EURODIAB ACE Study. Variation and trends in incidence of childhood diabetes in Europe. *Lancet* 2000;**355**:873–6.
- 3 Ali K, Harnden A, Edge J. Type 1 diabetes in children. *Br Med J* 2011;**342**:d294.
- 4 Usher-Smith JA, Thompson MJ, Sharp SJ, *et al*. Factors associated with the presence of diabetic ketoacidosis at diagnosis of diabetes in children and young adults: a systematic review. *Br Med J* 2011;**343**:d4092.
- 5 Blanc N, Lucidarme N, Tubiana-Rufi N. Factors associated to ketoacidosis at diagnosis of type 1 diabetes in children. *Arch Pediatr* 2003;**10**:320–5.
- 6 Hamilton D V, Mundia SS, Lister JC-1687347. Mode of presentation of juvenile diabetes. *Br Med J* 1976;**2**:211–2.
- 7 Mallare JT, Cordice CC, Ryan BA, *et al*. Identifying risk factors for the development of diabetic ketoacidosis in new onset type 1 diabetes mellitus. *Clin Pediatr (Phila)* 2003;**42**:591–7.
- 8 Pawłowicz M, Birkholz D, Niedźwiecki M, *et al*. Difficulties or mistakes in diagnosing type 1 diabetes mellitus in children? The consequences of delayed diagnosis. *Endokrynol Diabetol i Chor Przemiany Mater Wieku Rozw* 2008;**14**:7–12.
- 9 Soliman AT, ElZalabany MM, Bappal B, *et al*. Permanent neonatal diabetes mellitus: epidemiology, mode of presentation, pathogenesis and growth. *Indian J Pediatr* 1999;**66**:363–73.
- 10 Usher-Smith JA, Thompson M, Ercole A, *et al*. Variation between countries in the frequency of diabetic ketoacidosis at first presentation of type 1 diabetes in children: a systematic review. *Diabetologia* 2012;**55**:2878–94.

11     Rewers A, Brown A, Rewers M. Diabetic Ketoacidosis at Diagnosis Predicts Poorer Glycemic Control in the Initial Course of Type 1 Diabetes. *Abstr Present Pediatr Soc Meet Toronto, Canada*

12     Bowden SA, Duck MM, Hoffman RP. Young children (< 5 yr) and adolescents (> 12 yr) with type 1 diabetes mellitus have low rate of partial remission: diabetic ketoacidosis is an important risk factor. *Pediatr Diabetes* 2008;**9**:197–201.

13     Abdul-Rasoul M, Habib H, Al-Khouly M. “The honeymoon phase” in children with type 1 diabetes mellitus: frequency, duration, and influential factors. *Pediatr Diabetes* 2006;**7**:101–7.

14     Lokulo-Sodipe K, Moon RJ, Edge J a, *et al.* Identifying targets to reduce the incidence of diabetic ketoacidosis at diagnosis of type 1 diabetes in the UK. *Arch Dis Child* 2014;**99**:438–42.

15     Usher-Smith JA, Thompson MJ, Walter FM. “Looking for the needle in the haystack”: a qualitative study of the pathway to diagnosis of type 1 diabetes in children. *BMJ Open* 2013;**3**:e004068.

16     English Indices of Deprivation 2010. <https://www.gov.uk/government/publications/english-indices-of-deprivation-2010> (accessed 7 Feb2014).

17     Walter F, Webster A, Scott S, *et al.* The Andersen Model of Total Patient Delay: a systematic review of its application in cancer diagnosis. *J Health Serv Res Policy* 2012;**17**:110–8.

18     Scott SE, Walter FM, Webster a, *et al.* The model of pathways to treatment: conceptualization and integration with existing theory. *Br J Health Psychol* 2013;**18**:45–65.

19     Andersen BL, Cacioppo JT. Delay in seeking a cancer diagnosis: delay stages and psychophysiological comparison processes. *Br J Soc Psychol* 1995;**34**:33–52.

20     Allgar VL, Neal RD. Delays in the diagnosis of six cancers: analysis of data from the National Survey of NHS Patients: Cancer. *Br J Cancer* 2005;**92**:1959–70.

21     Neu A, Willasch A, Eehalt S, *et al.* Ketoacidosis at onset of type 1 diabetes mellitus in children - Frequency and clinical presentation. *Pediatr Diabetes* 2003;**4**:77–81.

22     Roche EF, Menon A, Gill D, *et al.* Clinical presentation of type 1 diabetes. *Pediatr Diabetes* 2005;**6**:75–8.

23     Xin Y, Yang M, Chen XJ, *et al.* Clinical features at the onset of childhood type 1 diabetes mellitus in Shenyang, China. *J Paediatr Child Health* 2010;**46**:171–5.

24     Neu A, Eehalt S, Willasch A, *et al.* Varying clinical presentations at onset of type 1 diabetes mellitus in children--epidemiological evidence for different subtypes of the disease? *Pediatr Diabetes* 2001;**2**:157–53.

- 25 Ting WH, Huang CY, Lo FS, *et al.* Clinical and laboratory characteristics of type 1 diabetes in children and adolescents: experience from a medical center. *Acta Paediatr Taiwanica* 2007;**48**:119–24.
- 26 Kapellen TM, Galler A, Nietzschmann U, *et al.* Prevalence of diabetic ketoacidosis in newly diagnosed children and adolescents with type 1 diabetes mellitus. Experience of a center for pediatric diabetology in Germany. [Article in German]. *Monatsschr Kinderheilkd* 2001;**149**:679–82.
- 27 Lévy-Marchal C, Patterson CC, Green A. Geographical variation of presentation at diagnosis of type I diabetes in children: The EURODIAB study. *Diabetologia* 2001;**44**:B75–80.
- 28 Neill SJ. Acute childhood illness at home: the parents' perspective. *J Adv Nurs* 2000;**31**:821–32.
- 29 Kai J. What worries parents when their preschool children are acutely ill, and why: a qualitative study. *Br Med J* 1996;**313**:983–6.
- 30 Dixon-Woods M, Findlay M, Young B, *et al.* Parents' accounts of obtaining a diagnosis of childhood cancer. *Lancet* 2001;**357**:670–4.
- 31 Francis N a, Crocker JC, Gamper A, *et al.* Missed opportunities for earlier treatment? A qualitative interview study with parents of children admitted to hospital with serious respiratory tract infections. *Arch Dis Child* 2011;**96**:154–9.
- 32 Forbes LJJ, Atkins L, Thurnham A, *et al.* Breast cancer awareness and barriers to symptomatic presentation among women from different ethnic groups in East London. *Br J Cancer* 2011;**105**:1474–9.
- 33 Forbes LJJ, Simon AE, Warburton F, *et al.* Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer* 2013;**108**:292–300.
- 34 Diabetes UK. 4 T's campaign. [https://www.diabetes.org.uk/Get\\_involved/Campaigning/Our-campaigns/4-Ts-campaign/](https://www.diabetes.org.uk/Get_involved/Campaigning/Our-campaigns/4-Ts-campaign/) (accessed 19 Aug2014).
- 35 Bui H, To T, Stein R, *et al.* Is diabetic ketoacidosis at disease onset a result of missed diagnosis? *J Pediatr* 2010;**156**:472–7.
- 36 Quinn M, Fleischman A, Rosner B, *et al.* Characteristics at diagnosis of type 1 diabetes in children younger than 6 years. *J Pediatr* 2006;**148**:366–71.
- 37 Pawlowicz M, Birkholz D, Niedzwiecki M, *et al.* Difficulties or mistakes in diagnosing type 1 diabetes in children?--demographic factors influencing delayed diagnosis. *Pediatr Diabetes* 2009;**10**:542–9.



TABLES

Table 1. Child and family characteristics for those included in the study

Child and family characteristics	Number	Percentage (%)
<b>Gender</b>		
Male	49	55.7
Female	39	44.3
<b>Age</b>		
0-5	26	29.5
6-10	20	22.7
11-16	42	47.7
Mean ± SD	9.41 ± 4.5	
<b>Ethnicity</b>		
White	80	90.9
Asian	2	2.3
Black	3	3.4
Mixed	3	3.4
<b>Family history</b>		
First degree relative(s) with T1D	8	9.1
First degree relative(s) with T2D	8	9.1
Second or third degree relative(s) with T1D	13	14.8
Second or third degree relative(s) with T2D	24	27.3
<b>Indices of deprivation</b>		
Least deprived tertile	43	48.9
Middle tertile	34	38.6
Most deprived tertile	9	10.2
Missing	2	2.3
<b>Medically trained family member</b>	9	10.2
<b>DKA at diagnosis</b>		
Yes	35	39.8
No	53	60.2

Table 2. Frequency of symptoms amongst all children and those with and without DKA and duration of individual symptoms

\* p &lt; 0.05

	Frequency of symptoms						Duration of symptoms		
	All (n=87)		DKA (n=35)		No-DKA (n=52)		Mean ± SD	Median (IQR)	n
	n	%	n	%	n	%			
Polydipsia	85	97.7	33	94.3	52	100	31.9 ± 48	16 (8,36)	77
Polyuria	73	83.9	27	77.1	46	88.5	29.8 ± 53	14 (5,26)	65
Tiredness	66	75.9	28	80.0	38	73.1	34.5 ± 49.2	17 (10,39)	53
Nocturia	64	73.6	28	80.0	36	69.2	31.3 ± 52.1	15.5 (7,28.5)	56
Weight loss	56	64.4	28	80.0*	28	53.8*	50.1 ± 82.7	13.5 (7,44)	42
Changes in behaviour/mood	48	55.2	17	48.6	31	59.6	34.3 ± 40.8	15 (8,42)	34
Change in appetite	45	51.7	18	51.4	27	51.9	30.7 ± 48	14.5 (7,39)	38
Abdominal pain	37	42.5	17	48.6	20	38.5	41.4 ± 64.1	17 (7,38)	25
Nocturnal enuresis	33	37.9	14	40.0	19	36.5	28.4 ± 49.2	15 (5.5,21.5)	28
Different smelling breath	31	35.6	14	40.0	17	32.7	17.5 ± 28.7	6.5 (3,17)	22
Vomiting	17	19.5	15	42.9*	2	3.8*	7.3 ± 12.6	2.5 (1.5,5.5)	8
Faster breathing	15	17.2	12	34.3*	3	5.8*	3.8 ± 5.8	0.5 (0,7.5)	8
Urinary incontinence	14	16.1	4	11.4	10	19.2	36.6 ± 77.2	10 (3,21)	10
Fever	12	13.8	6	17.1	6	11.5	25 ± 35.8	8 (2,55)	7

Table 3. Duration of diagnostic intervals

	Mean ± SD (days)	Median (IQR) (days)	<i>n</i>
Appraisal Interval	41 ± 51.7	20 (9,40)	75
Help-seeking Interval	3 ± 4.6	1 (0,4.5)	83
Diagnostic Interval	5 ± 34.8	0 (0,0)	83
Total Diagnostic Interval	48 ± 60.4	25 (14,50)	74

Table 4. Parents’ explanations for the ten most common symptoms

Symptom	<i>n</i>	Number with explanation for symptom <i>n</i> (%)	Most common explanations <i>n</i> (%)
Polydipsia	85	58 (68.2)	Hot weather 19 (32.8) Infection 13 (22.4) Activity/Travel 10 (17.2)
Polyuria	73	47 (64.4)	Drinking more 29 (61.7) Urine infection 6 (12.7) Diabetes 4 (8.5)
Tiredness	66	44 (66.7)	School related 12 (27.3) Infection 5 (11.9) Nocturia 4 (9.5)
Nocturia	64	40 (62.5)	Drinking more 26 (65.0) Diabetes 4 (10.0) Urine infection 3 (7.5)
Weight loss	56	33 (58.9)	Growth related 15 (45.5) Decreased appetite 4 (12.1) Increased activity 3 (9.1)
Changes in behaviour/mood	48	31 (64.6)	Tiredness 10 (32.3) Age related/puberty 7 (22.6) Infection/illness 6 (19.4)
Change in appetite	45	28 (62.2)	Growth related 14 (50.0) Infection 5 (17.9) Holiday related 2 (7.1)
Abdominal pain	37	19 (51.4)	Infection 4 (21.1) School related 3 (15.8) Period pains 3 (15.8)
Nocturnal enuresis	33	23 (69.7)	Drinking more 13 (56.5) Tired 4 (17.4) School related 3 (13.0)
Different smelling breath	31	14 (45.2)	Poor dental hygiene 4 (28.6) Infection 3 (21.4) Diabetes 3 (21.4)

Table 5. Factors influencing parents' decisions to seek medical advice sooner or later

	Not at all <i>n</i> (%)	A little <i>n</i> (%)	Quite a lot <i>n</i> (%)	Very much <i>n</i> (%)	Did not answer <i>n</i> (%)
<b>Factors influencing seeking medical advice <i>sooner</i></b>					
Concern something serious	9 (10.3)	16 (18.4)	18 (20.7)	42 (48.3)	2 (2.3)
Symptoms getting worse	7 (8.0)	19 (21.8)	14 (16.1)	46 (52.9)	1 (1.1)
Symptoms not getting better	4 (4.6)	12 (13.8)	22 (25.3)	45 (51.7)	4 (4.6)
Wanting reassurance	8 (9.2)	15 (17.2)	16 (18.4)	46 (52.9)	2 (2.3)
Comments from family	30 (34.5)	28 (32.2)	11 (12.6)	13 (14.9)	5 (5.7)
Comments from school	63 (72.4)	10 (11.5)	4 (4.6)	4 (4.6)	6 (6.9)
Comments from friends	49 (56.3)	20 (23.0)	7 (8.0)	5 (5.7)	6 (6.9)
Written information	50 (57.5)	8 (9.2)	10 (11.5)	15 (17.2)	4 (4.6)
<b>Factors influencing seeking medical advice <i>later</i></b>					
Difficulty getting appointment	60 (69.0)	8 (9.2)	7 (8.0)	11 (12.6)	1 (1.1)
Waiting for a particular doctor or nurse	68 (78.2)	7 (8.0)	4 (4.6)	6 (6.9)	2 (2.3)
Concern about having to wait at the surgery	72 (82.8)	6 (6.9)	4 (4.6)	3 (3.4)	2 (2.3)
Worry about wasting the doctor or nurse's time	61 (70.1)	10 (11.5)	8 (9.2)	6 (6.9)	2 (2.3)
Worry the doctor would not take them seriously	62 (71.3)	12 (13.8)	3 (3.4)	7 (8.0)	3 (3.4)
Symptoms weren't very serious	55 (63.2)	20 (23.0)	9 (10.3)	0 (0)	3 (3.4)
Hope the symptoms would go away	42 (48.3)	21 (24.1)	9 (10.3)	15 (17.2)	0 (0)
Fear of serious diagnosis	58 (66.7)	16 (18.4)	5 (5.7)	7 (8.0)	1 (1.1)

Figure 1a. Associations between parent/child characteristics and the total diagnostic interval. IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure. Cox model stratified by whether parents suspected the diagnosis or not.






Figure 1b. Associations between parent/child characteristics and the appraisal interval.  
IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure




Figure 1c. Associations between parent/child characteristics and the help-seeking interval.

IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure





Figure 2. Associations between parent/child characteristics and presence/absence of DKA.

IMD – index of multiple deprivation. Odds ratios adjusted for all variables in the figure



	Appraisal interval			Help-seeking interval			Diagnostic interval			Total diagnostic interval		
	Mean ± SD	Median (IQR)	<i>n</i>	Mean ± SD	Median (IQR)	<i>n</i>	Mean ± SD	Median (IQR)	<i>n</i>	Mean ± SD	Median (IQR)	<i>n</i>
All	41 ± 51.7	20 (9,40)	75	3 ± 4.6	1 (0,4.5)	83	5 ± 34.8	0 (0,0)	83	48 ± 60.4	25 (14,50)	74
Age												
0-5 years	47.8 ± 53.8	35 (11,67)	23	2.4 ± 2.9	1 (0,3)	25	1.1 ± 3.2	0 (0,0)	24	47.7 ± 51.9	36 (15,64)	21
6-10 years	29.2 ± 40.7	15.5 (11,31)	18	2.5 ± 3.7	1 (0,3)	18	16.1 ± 69.0	0 (0,0)	20	47.3 ± 75.6	21 (15,36)	19
11-16 years	38.5 ± 49.9	17 (7,38)	34	4.1 ± 5.6	3 (0,5)	39	2.2 ± 7.4	0(0,0)	38	45.5 ± 51.9	22 (11,50)	34
Gender												
Male	43.8 ± 56.6	20 (10,53)	43	2.9 ± 4.7	1 (0,4)	47	1.8 ± 6.5	0 (0,0)	45	46.9 ± 56.9	22 (13,59)	41
Female	32.8 ± 36.2	19.5 (7,38)	32	3.7 ± 4.5	1 (1,7)	35	9.4 ± 50.8	0 (0,0)	37	46.2 ± 60.2	29 (15,45)	33
Family history of T1D												
No FH	37.7 ± 51.8	19 (8,37)	41	4 ± 5.5	2 (1,5)	46	1.8 ± 6.4	0 (0,0)	46	44.6 ± 52.6	23 (15,50)	41
1st degree relative	42.5 ± 54.3	29 (11,37)	13	1.5 ± 2.5	0 (0,3)	14	0.8 ± 2.2	0 (0,0)	13	32.8 ± 47.8	15.5 (9,38.5)	12
2nd or 3rd degree relative	39.7 ± 41.7	32 (12,56)	21	2.8 ± 2.8	1.5 (1,5)	22	14.7 ± 64.3	0 (0,0)	23	58.3 ± 72.3	40 (17,64)	21
Prior knowledge of symptoms												
Yes	38.4 ± 44.7	20 (11,42)	46	3.7 ± 5.3	1 (0,6)	47	1.1 ± 3.7	0(0,0)	48	42.3 ± 46.5	23 (14,49)	47
No	42.7 ± 57.3	22 (7,53)	27	2.8 ± 3.5	2 (0,3)	33	11.6 ± 54.8	0 (0,0)	32	57.5 ± 76.3	28 (15,64)	25
Deprivation												
Low	33.4 ± 18.9	31 (27,38)	9	3.8 ± 4.7	2 (0,7)	9	2.3 ± 7.3	0 (0,0)	42	36.8 ± 36.8	20 (11,58)	37
Middle	46.9 ± 62.1	19 (11,38)	29	2.2 ± 2.8	1 (0,3)	30	11.6 ± 57.3	0 (0,0)	29	58.1 ± 79.1	27 (15, 49)	27
High	30.3 ± 34.5	14 (6.5, 47)	36	3.9 ± 5.5	2 (0,6)	41	0 ± 0	0 (0,0)	9	37.2 ± 17.8	36 (28,40)	9
Parents considered diabetes												
Yes	40.2 ± 45.8	23.5 (12,50)	54	3.8 ± 5	2 (1,6.5)	60	1 ± 3.4	0 (0,0)	60	45.3 ± 47.3	28.5 (16,50)	54
No	36.3 ± 57.6	14 (7,37)	21	1.7 ± 2.6	0.5 (0,3)	22	17 ± 65.8	0 (0,2)	22	50.0 ± 81.7	19 (7,50)	20
Diagnosis at first appointment												
Yes	35.2 ± 45.4	20 (11,38)	61	3.0 ± 3.7	1.5 (0,5)	66	1.1 ± 2.4	0 (0,0)	32	38.4 ± 44.8	22.5 (14,41)	62
No	56.1 ± 61.3	25.5 (7,92)	14	4.1 ± 7.2	1(0,5)	16	7.9 ± 44	0 (0,0)	50	89.0 ± 94.1	61.6 (15, 145)	12
First contact with healthcare												
Primary care	42.5 ± 51.0	23.5 (11,53)	66	3.5 ± 4.7	2 (1,5)	73	5.9 ± 36.4	0 (0,0)	73	51.0 ± 60.4	28 (16,59)	65
Secondary care	14.4 ± 16.4	8 (4,16)	9	0.7 ± 1.3	0 (0,1)	9	0 ± 0	0 (0,0)	9	15.1 ± 16.4	11 (4,17)	9
DKA												
Yes	42.5 ± 59.4	21 (7,36.5)	28	2.9 ± 5.6	1 (0,3)	34	0.1 ± 0.7	0 (0,0)	68	43 ± 59.6	21 (13,36)	26
No	37.1 ± 42.2	20 (11,50)	47	3.4 ± 3.7	2 (1,5.5)	48	30.4 ± 80.9	6.5 (2,12)	14	48.5 ± 57.7	35.5 (14.5, 61.5)	48

Appendix Table 1. Time intervals along the pathway to diagnosis.

For peer review only



# BMJ Open

## The pathway to diagnosis of type 1 diabetes in children: a questionnaire study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-006470.R1
Article Type:	Research
Date Submitted by the Author:	17-Nov-2014
Complete List of Authors:	Usher-Smith, Juliet; The Primary Care Unit, Institute of Public Health Thompson, Matthew; Oxford University, Department of Primary Care Health Sciences Zhu, Hannah; Cambridge University Hospitals NHS Foundation Trust, Sharp, Stephen; University of Cambridge, Walter, Fiona; University of Cambridge, Dept of Public Health and Primary Care
<b>Primary Subject Heading</b>:	Diabetes and endocrinology
Secondary Subject Heading:	Diagnostics, General practice / Family practice
Keywords:	Paediatric endocrinology < DIABETES & ENDOCRINOLOGY, PRIMARY CARE, Paediatric A&E and ambulatory care < PAEDIATRICS

SCHOLARONE™  
Manuscripts

The pathway to diagnosis of type 1 diabetes in children: a questionnaire study

Juliet A Usher-Smith<sup>1</sup>, Matthew J Thompson<sup>2</sup>, Hannah Zhu<sup>3</sup>, Stephen J Sharp<sup>4</sup>, Fiona M Walter<sup>5</sup>

<sup>1</sup> Clinical Lecturer in General Practice, The Primary Care Unit, University of Cambridge CB1 8RN

<sup>2</sup> Professor, Department of Family Medicine, University of Washington, Seattle, WA 98195 USA

<sup>3</sup> Foundation Year 2 Doctor, Cambridge University Hospitals NHS Foundation Trust, CB2 0QQ

<sup>4</sup> Senior Statistician, MRC Epidemiology Unit, University of Cambridge, Institute of Metabolic Science, Cambridge, CB2 0QQ

<sup>5</sup> GP & Clinician Scientist, The Primary Care Unit, University of Cambridge CB1 8RN

Correspondence to: J Usher-Smith [jau20@medschl.cam.ac.uk](mailto:jau20@medschl.cam.ac.uk)

ABSTRACT

**Objective** To explore the pathway to diagnosis of type 1 diabetes (T1D) in children

**Design** Questionnaire completed by parents

**Participants** Parents of children aged 1 month to 16 years diagnosed with T1D within the previous three months

**Setting** Children and parents from 11 hospitals within the East of England.

**Results** 88/164 (54%) of invited families returned the questionnaire. Children had mean±SD age of 9.41±4.5 years. 35 (39.8%) presented with DKA at diagnosis. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%) and all children presented with at least one of those symptoms. The time from symptom onset to diagnosis ranged from 2 to 315 days (median 25 days). Most of this was the appraisal interval from symptom onset until perceiving the need to seek medical advice. Access to healthcare was good but one in five children presenting to primary care were not diagnosed at first encounter, most commonly due to waiting for fasting blood tests or alternative diagnoses. Children diagnosed at first consultation had a shorter duration of symptoms (p=0.022) and children whose parents suspected the diagnosis were 1.3 times more likely (RR 1.3, 95% CI 1.02-1.67) to be diagnosed at first consultation.

**Conclusions** Children present with the known symptoms of T1D but there is considerable scope to improve the diagnostic pathway. Future interventions targeted at parents need to address the tendency of parents to find alternative explanations for symptoms and the perceived barriers to access, in addition to symptom awareness.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study uses a questionnaire developed from a previous interview study to explore the diagnostic pathway of children with newly diagnosed T1D
- It uses the Model of Pathway to Treatment as a framework to allow analysis of the factors acting at different stages in the pathway
- The inclusion of a calendar with key events in the questionnaires and use of free text responses for internal validation and checking of prompted responses reduced bias but the data was necessarily collected retrospectively and so subject to recall and framing bias

## INTRODUCTION

Approximately 65,000 children are diagnosed with type 1 diabetes (T1D) each year and the incidence is continuing to increase at a rate of approximately 3% per year<sup>1,2</sup>. The most common symptoms are well described and include polyuria, polydipsia, weight loss and tiredness. At the early stages of the disease, however, these symptoms are often non-specific and distinguishing the children with T1D from the large number with similar symptoms and minor undifferentiated illness can therefore be difficult. This is reflected in studies which have shown that the mean duration of symptoms prior to diagnosis is over two weeks with a significant number of children experiencing delay in diagnosis or misdiagnosis<sup>3</sup> and only one in five diagnosed at first encounter<sup>4-8</sup>. Up to 80% of children additionally present in diabetic ketoacidosis (DKA)<sup>9</sup> which has both immediate life-threatening complications and is associated with poorer long term diabetic control<sup>10-12</sup>.

Whilst several studies have highlighted these difficulties in making the diagnosis and the features associated with diabetic ketoacidosis at diagnosis<sup>3-8,13</sup>, few have explored the period between symptom onset and diagnosis. Our recent qualitative interview study of parents and General Practitioners (GPs) of children newly diagnosed with T1D suggested that the longest component in the diagnostic pathway is the time between onset of symptoms and the decision to seek medical help (known as the appraisal interval)<sup>14</sup>. The early symptoms are subtle, and even with some knowledge of T1D it took many parents several weeks of a complex decision making process and often a physical trigger, such as weight loss or vomiting, to decide to consult a healthcare professional. Once the decision to seek help had been made almost all children were seen immediately and diagnoses were mostly prompt and managed appropriately. Parents continued to play a key role during the diagnostic interval however, with many having already made or suspected the diagnosis themselves, and several feeling that their GP did not take their concerns seriously.

This study builds on this earlier work by using a questionnaire developed from the interview findings to further explore the pathway to diagnosis of T1D in children. By using a structured questionnaire to survey a larger number of families we aimed to quantify the symptoms and their time course prior to diagnosis, the triggers and barriers to seeking help, the influence of parental prior knowledge of diabetes, and the role of healthcare services.

## METHODS

### Design

A questionnaire about the pathway from first symptom(s) to diagnosis was completed by the parent(s)/guardian(s)/step-parents (hereafter referred to as parents) of children aged 1 month to 16 years diagnosed with T1D within the previous three months.

### Recruitment

Children and parents were identified and recruited via the paediatric diabetes specialist nurses and research nurses at 11 hospitals within the East of England Diabetes Children and Young People's Network. Parents of all children aged 1 month to 16 years diagnosed with T1D diagnosed within the previous 3 months at participating hospitals were eligible for inclusion unless their clinical team felt that this was not appropriate. Parents who failed to respond within one month were sent a reminder letter with a further copy of the questionnaire. Recruitment began at each site between February 2013 and April 2013, and continued across all sites until January 2014.

The clinical or research teams at all sites collected data on the age and gender of each child diagnosed during the study period and whether they had DKA at diagnosis. Each hospital used a slightly different definition of DKA but all included either pH < 7.3, bicarbonate < 15 mmol/L (see Appendix Table 1).

**The questionnaire**

The questionnaire was developed from the findings of our previous qualitative study of parents and children recently diagnosed with T1D<sup>14</sup>. It was first reviewed by an expert panel comprising paediatric diabetes consultants, a paediatric diabetes research nurse and primary care researchers, and then piloted with parents of four children recently diagnosed with T1D. In addition to their specific feedback, parents were asked to talk aloud whilst completing the questionnaire and then interviewed after completion to ensure face validity. Based on feedback from the parents, the questionnaire was revised.

The final questionnaire included 5 sections (see Supplementary file). The first included questions about the child's age, gender, postcode, ethnic background, family history of diabetes, any medically trained family members and the number of children in the household. Parents were also asked if they knew what the symptoms of diabetes in children are before their child was diagnosed, and if so, to give details of those symptoms they were aware of. The second section asked about the symptoms the children had experienced with yes/no responses for 14 symptoms and space to add the date they noticed the symptoms, what they thought the symptoms were due to at the time and how much it concerned them. The third section focused on help-seeking and asked where parents had looked for information, who they spoke to and then details on when and how they had sought medical advice. It also asked them to describe their main concern at their first appointment and whether they had considered diabetes. Parents were also asked in this section about factors contributing to their decision to seek medical advice sooner or later. The fourth section asked about the diagnosis, including whether it was made at their first appointment with a healthcare professional and, if not, how many subsequent consultations they had, and the investigations that were done before diagnosis. The final section then asked parents if they felt there was anything that prolonged them finding out their child had diabetes and had further space for free text comments.

**Analysis**

Data from the questionnaires were entered into a database and then double checked by a second researcher. Socioeconomic status was computed using postcode and the English indices of deprivation 2010 available online<sup>15</sup>. The presence of DKA at diagnosis was obtained from hospital records rather than self-report. Walter et al's Model of Pathways to Treatment<sup>16,17</sup> provided a theoretic model of the intervals that occur prior to a diagnosis. This model divides the pathway to diagnosis into two intervals prior to presentation to healthcare about a symptom (the appraisal interval from the onset of symptoms to perceiving a reason to discuss symptoms with a healthcare professional, and the help-seeking interval from that decision until presentation to a healthcare professional), and then the diagnostic interval from first presentation to a healthcare professional until diagnosis. The help-seeking interval was further sub-divided into the behavioural interval (the time between perceiving the reason to discuss the symptoms with a healthcare professional to making the decision to seek help) and the scheduling interval (the time between making the decision to seek help and the first consultation)<sup>18</sup>. Intervals were calculated from responses to the questionnaire. Where dates were incomplete we applied midpoint rules to estimate the actual date<sup>19</sup>. In cases where the responses in free text differed from the dates entered as numbers, the free text was assumed to be correct, and where there was uncertainty the researchers met to agree consensus.

Characteristics (age, gender, presence of DKA) were compared between children whose parents had and had not returned a questionnaire using a t-test for age and chi-squared test for gender and presence of DKA. All further analyses used only data from returned questionnaires. The frequency of the 14 symptoms was compared between those with and without DKA using a chi-squared test. Cox regression was used to estimate the association between various factors and the hazard of

diagnosis; if a factor was associated with an increased hazard (i.e. hazard ratio greater than 1), this implied that that factor was associated with a shorter time to diagnosis, and vice versa. Time to diagnosis was from the date of the earliest symptom to the date of diagnosis, and the factors assessed were age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the parents suspected T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis. A similar approach was used to assess factors associated with the length of the appraisal and help-seeking intervals (with the end of the interval being defined as the “event” in the Cox model), but only the first six variables in the list above were considered, as the others do not relate to those time intervals. The Schoenfeld residuals test was used to assess the proportional hazards (PH) assumption for each covariate in each model. Whether parents suspected the diagnosis of T1D did not meet the PH assumption for the total diagnostic interval and so the Cox regression model was stratified by that variable. Logistic regression was used to estimate the association between the same factors and presence of DKA at diagnosis. All analyses were performed using STATA version 12.

Free text responses were grouped into similar categories and coded. Where individual free text responses contained several comments, these were each coded individually.

## RESULTS

A total of 172 children were diagnosed with T1D in the 11 hospitals during the study period. Of those, 8 families were not invited to take part in the study: 5 lived outside the hospital catchment area; 1 emigrated the week after diagnosis; and the clinical team felt it was not appropriate to include 2. From the remaining 164 families invited to take part in the study, 88 (54%) completed and returned the questionnaire. There were no significant differences in the proportion presenting in DKA ( $p=0.27$ ), mean age ( $p=0.77$ ) or gender ( $p=0.77$ ) between children of responders and non-responders.

One child was excluded from the analysis as they had no symptoms and the diagnosis was made on a random blood glucose test that the parents were doing at home on an intermittent basis as they had an older child with T1D. Children whose parents checked blood glucose at home after noticing symptoms remain in the analysis. 87 children are therefore included in the analysis that follows.

Table 1 shows the characteristics of the 87 children and families included in the study. The mean age was  $9.34 \pm 4.5$  years, 49 (56.3%) were male and 35 (40.2%) presented with DKA at diagnosis. The majority (90.8%) were white and as a group they were generally from less deprived areas of England, with 49.4% from the least deprived tertile of English Indices of Deprivation and only 10.3% from the most deprived.

## Symptoms

Table 2 shows the frequency and duration of the 14 symptoms that were specifically asked about in the questionnaire. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%). Most symptoms were present for a median of between 13 and 17 days. Faster breathing and vomiting both had much shorter median (IQR) durations of 0.5 (0-7.5) and 2.5 (1.5-5.5) days respectively than the other symptoms. Weight loss, vomiting and faster breathing were significantly more frequent in those children who presented in DKA ( $p = 0.014$ ,  $<0.0005$  and  $0.001$  respectively). All the children had at least one of the 4 main symptoms (polydipsia, polyuria or nocturia, weight loss or tiredness), 97.7% had 2 or more, 79.3% 3 or more and over half (50.6%) had all 4 symptoms.



A very small number of parents mentioned symptoms other than those listed in the questionnaire, these included constipation (9), headaches (3), thrush (3), blurred vision (2), dry skin (2) and different smelling urine (1).

**Diagnostic intervals**

Table 3 shows the mean  $\pm$  SD and median (IQR) for the diagnostic intervals. Additional details on the diagnostic intervals for different subgroups are shown in Appendix Table 2. The total diagnostic interval ranged from 2 to 315 days with a median (IQR) of 25 days (14-50). In unadjusted Cox regression analysis (data not shown) the time to diagnosis was significantly shorter for children diagnosed at first appointment compared to a subsequent appointment ( $p=0.046$ ) and for those seen in secondary care rather than primary care ( $p=0.01$ ). No evidence of associations with time to diagnosis was found for age, gender, family history of T1D, deprivation, prior knowledge of symptoms or DKA at diagnosis. In multivariable cox regression including age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis (Figure 1a), the association between whether the diagnosis of T1D was made at the first or subsequent appointments and total diagnostic interval remained statistically significant ( $p=0.022$ ).

***The appraisal interval***

The appraisal interval was the longest of all the intervals in the pathway for all but 3 of the families with a mean  $\pm$  SD of  $41.0 \pm 51.7$  days and median (IQR) 20 (9-40) days. During this period nearly two thirds (64%) of parents discussed the symptoms with family members, 40% with friends and 41% looked on the internet. Only 16% spoke to the child’s nursery, school or playgroup and very few (6%) looked for information in books. Over half of parents (49, 56%) reported being aware of some symptoms of T1D in children prior to their child’s diagnosis: 40 (45%) were aware of increased thirst, 24 (27%) of polyuria, 17 (19%) of weight loss and 13 (15%) tiredness.

Cox-regression analysis (Figure 1b) showed no significant associations between parent/child characteristics and the appraisal interval.

Analysis of the free text showed that most parents found explanations for their child’s symptoms (Table 4). For example, polydipsia was attributed most commonly to hot weather (19/58, 33%) or infection (13/58, 22%), polyuria and nocturia were frequently explained by drinking more (29/47, 62% and 26/40, 65%) and tiredness was thought to be school related (12/44, 27%) or secondary to infection (5/44, 12%) or nocturia (4/44, 10%).

The majority of parents (61/87, 70%) additionally reported that they had suspected diabetes before their first consultation with a healthcare professional. When asked what had made them suspect diabetes, the most common reason given was that they knew the symptoms (22/59, 37%), especially thirst (12/59, 20%). Others cited information from the internet (12/59, 20%) or having a family history of diabetes (11/59, 19%).

***The help-seeking interval***

24 (28%) children were seen on the same day their parents first thought about seeking medical advice and 64 (74%) within 5 days. Most of this time was the behavioural interval (mean  $\pm$  SD  $2.1 \pm 3.7$  days, median (IQR) 0 (0-3) days) rather than the scheduling interval (mean  $\pm$  SD  $1.1 \pm 2.6$  days, median (IQR) 0 (0-1) days).

Cox-regression analysis (Figure 1c) showed no significant associations between parent/child characteristics and the help-seeking interval.

The most common reasons that parents cited for seeking medical advice sooner rather than later (Table 5) were that the symptoms were not getting better or were getting worse, wanting reassurance or concern something serious was wrong. This was also reflected in the free text responses where 22% of parents noted that worsening or persistent symptoms was the reason they decided to seek help. In general, fewer parents reported factors that led to them seeking medical advice later. Of those that did, the most common reason for waiting was hope that the symptoms would go away (51.6%) but 29.8% felt difficulty getting an appointment contributed and 27.6% and 25.2% were worried about wasting the GPs time or that the GP would not take them seriously respectively.

### ***The diagnostic interval***

The diagnostic interval was the shortest of the intervals with a mean  $\pm$  SD of 5 days  $\pm$  34.8 and median 0 (IQR) (0-0) days. 69 (78%) of children were diagnosed at first consultation. Cox regression was not possible given the high number of children with a diagnostic interval of zero. However, children whose parents suspected the diagnosis ( $n=61$ , 70.1%) were more likely (unadjusted RR 1.30, 1.02-1.67,  $p=0.046$ ) to be diagnosed at first consultation ( $n=52$ , 85.2%) than those in whom there was no suspicion ( $n=26$ , 29.9% with 17 (65.4%) diagnosed at first consultation). All children (10) who were seen first in secondary care were diagnosed at first consultation compared to 76.6% (59/77) of those seen first in primary care, but this difference was not statistically significant ( $p=0.114$ ). None of the variables considered were significantly associated with risk of DKA (Figure 2).

Further details from the questionnaires were available from 14 of the 18 children who were not diagnosed at first encounter with primary care. Of these, 6 had fasting glucose blood tests arranged by the GP and 4 were given alternative diagnoses (urine infection, viral infection, tonsillitis, puberty) and diagnosed at a second appointment. Two children were diagnosed with psychological problems: In one case the child's mother had seen the GP alone to discuss her child's 'obsessive drinking' and was advised to see the school counsellor, and in the second the GP apparently felt the symptoms were psychological and the child was diagnosed in the emergency department four consultations later. One other family had already done a finger prick glucose test at home which was high but the GP did not trust the result and asked the child to come back later in the day with a urine sample. In the final case, the child's mother had spoken to a health visitor and suggested diabetes but was told 'no, not unless the child is lifeless'. The mother took the child to the GP 12 days later and the diagnosis was made at that consultation.

## **DISCUSSION**

### **Principal findings**

This study shows that all children with new onset T1D present with one, and 98% present with two, of the four main symptoms of diabetes (polydipsia, polyuria, weight loss and tiredness). Moreover, over half have had symptoms for over three weeks before diagnosis. Most of that time is the appraisal interval during which parents found alternative explanations for the symptoms, discussed the symptoms with family and friends and looked on the internet for information. Once they made the decision to seek advice, access to healthcare was generally not difficult with 28% consulting with a healthcare professional on the same day. However, when asked about factors contributing to their decision to seek help, nearly a third of parents felt that difficulty getting an appointment contributed to them waiting to seek help and over a quarter felt that worry about wasting the

doctor’s time influenced their decision. This suggests that even if access is not difficult, it is perceived as such.

Once parents had sought help, one in five children were then not diagnosed at their first consultation with a healthcare professional, mainly due to being given an alternative diagnosis, most commonly infection, or waiting for further investigations. Diagnosis at first consultation was associated with a shorter total diagnostic interval and children were more likely to be diagnosed at first consultation when their parents suspected the diagnosis of T1D. The association between diagnosis at first consultation and total diagnostic interval may simply reflect the additional time between consultations, or it may be due to biological differences causing some children to develop symptoms more slowly which are then more difficult for both parents and primary care physicians to recognise.

**Strengths and weaknesses**

By using a questionnaire developed from a previous interview study<sup>14</sup> and the Model of Pathway to Treatment<sup>16,17</sup> as a framework for analysis, this study provides in-depth insights into the diagnostic pathway of children with newly diagnosed T1D and allows factors acting at different stages in the pathway to be explored.

The main weakness is that the data was necessarily collected retrospectively and so subject to recall and framing bias. Parents have multiple contacts with different healthcare professionals in the period immediately following diagnosis and so their responses to the questionnaire reflect a post-hoc rationalisation of events framed by those subsequent encounters and increased knowledge since the diagnosis. The inclusion of a calendar with key events in the questionnaires minimised the error in recall of dates, and the free text responses allowed internal validation and checking of prompted responses. Despite these efforts, we still only have the parents’ perspective on the pathway and were not able to confirm the number of healthcare contacts, diagnostic tests or the parental reports of missed opportunities for diagnosis. We were, however, able to confirm the diagnosis of DKA from clinical records and, although there was variation in the definition of DKA used across the 11 sites, all included a biochemical measurement of either pH or bicarbonate.

Our results are also based on the views of 88 parents. Although not a large number, they were recruited from 11 sites across a large region of the UK and the response rate was over 50% with no significant differences in gender, age or DKA status between the children whose parents responded and those who did not. The fact that they were a predominantly white group from less deprived areas of England limits the generalizability of the results outside the East of England but the main findings are likely to be relevant across the UK and other countries with similar primary care healthcare provision. The questionnaire also did not include questions specifically for the children to complete and so we are unable to comment on the views of the children during this time.

**Comparison with existing literature**

The median duration of symptoms prior to diagnosis was 13-17 days for the nine most frequent symptoms, with a mean of 30-50 days. This is longer than previous studies relying on retrospective review of medical records<sup>20-23</sup> but similar to studies which have used a checklist to identify subtle symptoms<sup>24</sup> or asked parents soon after diagnosis<sup>13,14</sup>. The wide range (a few days to over six months) has been described previously<sup>14,21,23</sup> and highlights the heterogeneous nature of the disease.

The frequency of individual symptoms we report is also similar to previous studies<sup>13,14,20,22,25</sup>. Additionally we showed that all the children had at least one of 4 symptoms (polydipsia, polyuria, weight loss and fatigue) and over half (50.6%) had all four. Consistent with the known course of

the disease and previous studies, vomiting<sup>4,22,24</sup>, weight loss<sup>13,25,26</sup>, and dyspnoea<sup>22</sup> were more common in those children who presented in DKA.

This is the first quantitative study to compare the time periods during the pathway to diagnosis of T1D in children. The finding that most of the total diagnostic interval was the appraisal interval is consistent with a previous qualitative study<sup>14</sup> and the free text analysis confirms that during that time the parents find alternative explanations for the symptoms initially and make use of a social network of extended family, friends and work colleagues, or the internet<sup>14,27,28</sup>. That children were more likely to be diagnosed at their first encounter with a healthcare professional when their parents suspected diabetes prior to that consultation may also reflect the findings of previous qualitative work in which a number of parents prompted the GP to consider T1D and pushed for investigations<sup>14</sup>. However, whilst parental suspicion of T1D has also been shown to be associated with a reduced risk of DKA in a parental survey<sup>13</sup>, in that study the incidence of DKA at presentation was no different whether or not the parents discussed their concerns with the healthcare professional, suggesting other factors may be contributing. The absence of an effect of parental prior knowledge of diabetes either on the total diagnostic interval or the risk of DKA further highlights the complexities around the role of knowledge on help-seeking behaviour.

The finding that parents worry about wasting the doctor's time has also been shown in previous qualitative studies in children<sup>29,30</sup> and in studies of help-seeking behaviour for adults with symptoms of cancer in the UK<sup>31,32</sup> and so it may reflect a particular British trait rather than be specific to T1D or children.

### **Implications for clinicians and policymakers**

Clinicians should remain alert to the possibility of T1D in all children presenting with one or more symptoms of polyuria, polydipsia, weight loss and tiredness – as almost all children have at least two of these. Interventions targeted at increasing public awareness, such as the 4 T's campaign launched by Diabetes UK to raise awareness of the four most common symptoms of T1D (Toilet, Thirsty, Tired and Thinner)<sup>33</sup>, should continue to focus on these established symptoms.

As most of the time between symptom onset and diagnosis is the appraisal interval, the greatest benefit is likely to be seen from interventions directed towards parents and their social network, probably via the internet. Despite ongoing government pressure for better access to primary care, improving access is unlikely to have much impact on the pathway. Instead efforts should be made to address the perception that access is difficult and the general concern in the UK about wasting healthcare professional time, particularly for children with acute or sub-acute health concerns.

Additionally, although the diagnostic interval itself was generally short, one in five children presenting to primary care were not diagnosed at first consultation. Similar numbers have been reported in a recent survey in the UK which found that 24% were not diagnosed at first contact with a healthcare professional<sup>13</sup>, and studies in the USA, Canada and Poland noted between 14 and 35% of children had more than one consultation before diagnosis<sup>6,7,34–36</sup>. As in those studies, the most common reasons for not being diagnosed at first encounter was either being given an alternative diagnosis, most commonly infection, or waiting for further investigations. In this study 33% of those not diagnosed at first consultation were waiting for fasting glucose tests and in other studies the number waiting for further investigations is as high as 46%<sup>6 13</sup>. This suggests that healthcare professionals may have considered a diagnosis of T1D but either lack ready access to rapid tests to confirm or exclude the diagnosis, or are reluctant to use existing tests in children<sup>14</sup>. Access to point of care urine and finger-prick testing and the use of those tests should be routine management for all children presenting with one or more of the four main symptoms of diabetes. The increased use of



point of care testing in Emergency departments may also explain why all children seen in secondary care were diagnosed at their first consultation. Whilst educational interventions aimed at primary care physicians may help a small number of children not currently diagnosed at first encounter, finding ways to overcome barriers to point-of-care tests in primary care may be more effective and this approach may also improve the diagnosis of other serious illnesses in children and adults.

**Unanswered questions and future research**

Whilst this study contributes to our understanding of the pathway to diagnosis and the stages at which this may be improved, the findings are unable to explain the large variability in the overall duration of the pathway to diagnosis and why some children develop DKA within a few weeks whilst others can be symptomatic for up to six months before requiring treatment. Further studies are, therefore, needed into the natural course and biology of the disease to better understand these variations. The findings also highlight the need for continuing research into the presentation of serious but rare conditions in primary care and the best ways to improve diagnosis of these conditions.

**ACKNOWLEDGEMENTS**

We thank all the parents who kindly gave up their time and shared their personal stories with us to either help develop the questionnaire or take part in the study. We also thank the Paediatric diabetes clinical teams at the 11 hospitals, particularly Jon Hassler-Hurst (Ipswich), Holly Roper (Norfolk and Norwich), Elissa Harwood (Cambridgeshire community services), Nazia Bhatti and Claire Pesterfield (Addenbrooke’s hospital NHS foundation trust), Julia Harding and Jacky Plumb (West Suffolk), Philippa Corbishley and Laura Ardrey (QE2 and Lister), Suzie Williams (SEPT), Mandy Stevenson (Harlow), Claire Gibbs and Allyson Davison (James Paget) and Meeta Patel (Luton and Dunstable). We are also grateful to the Diabetes Research Network and the East of England Children and Young People’s Diabetes Network and network manager Kate Wilson for support, James Brimicombe for advice on data management, Dr Carlo Acerini for comments on the questionnaire and final manuscript and support through his role as Chair of the East of England Children and Young People’s Diabetes Network, Professor Jon Emery for comments on the questionnaire, and Professor David Dunger for his support and encouragement throughout the study and comments on the final manuscript.

**Contributors**

JUS, MT, FMW and SJS were involved in the design of the study and all authors were involved in analysis of the data. JUS and FMW developed the questionnaire. JUS wrote the first draft of the manuscript and all authors reviewed and edited the manuscript.

**Funding**

The study was funded by the Royal College of General Practitioners Scientific Foundation Board (SFB-2011-15). JUS was supported by a National Institute of Health Research (NIHR) Academic Clinical Fellowship and subsequently Clinical Lectureship, and FMW by an NIHR Clinician Scientist award. SJS was supported by the Medical Research Council [www.mrc.ac.uk](http://www.mrc.ac.uk) [Unit Programme number MC\_UU\_12015/1]. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

All researchers were independent of the funding body (although FMW subsequently became a member of the Royal College of General Practitioners Scientific Foundation Board), and the study sponsors and funder had no role in study design; data collection, analysis and interpretation of data; in the writing of the report; or decision to submit the article for publication.

### Ethics approval

The study obtained ethical approval from the East of England Hertfordshire REC (reference number 12/EE/0390).

### Data sharing

The questionnaire is available from the corresponding author on request.

### Competing Interests

All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare that (1) JUS, MT, HZ, SJS and FMW have no support from or relationships with companies that might have an interest in the submitted work in the previous 3 years; (2) their spouses, partners, or children have no financial relationships that may be relevant to the submitted work; and (3) JUS, MT, HZ, SJS and FMW have no non-financial interests that may be relevant to the submitted work.

The corresponding author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in BMJ editions and any other BMJ PGL products and sublicences to exploit all subsidiary rights, as set out in our licence (<http://resources.bmj.com/bmj/authors/checklists-forms/licence-for-publication>).

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

The corresponding author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

### REFERENCES

- 1 Lawrence JM, Imperatore G, Dabelea D, *et al.* Trends in incidence of type 1 diabetes among non-Hispanic white youth in the u.s., 2002-2009. *Diabetes* 2014;**63**:3938–45.
- 2 Fazeli Farsani S, Souverein PC, van der Vorst MMJ, *et al.* Increasing trends in the incidence and prevalence rates of type 1 diabetes among children and adolescents in the Netherlands. *Pediatr Diabetes* Published Online First: 7 November 2014. doi:10.1111/pedi.12232
- 3 Usher-Smith JA, Thompson MJ, Sharp SJ, *et al.* Factors associated with the presence of diabetic ketoacidosis at diagnosis of diabetes in children and young adults: a systematic review. *Br Med J* 2011;**343**:d4092.
- 4 Blanc N, Lucidarme N, Tubiana-Rufi N. Factors associated to ketoacidosis at diagnosis of type 1 diabetes in children. *Arch Pediatr* 2003;**10**:320–5.
- 5 Hamilton D V, Mundia SS, Lister JC-1687347. Mode of presentation of juvenile diabetes. *Br Med J* 1976;**2**:211–2.
- 6 Mallare JT, Cordice CC, Ryan BA, *et al.* Identifying risk factors for the development of diabetic ketoacidosis in new onset type 1 diabetes mellitus. *Clin Pediatr (Phila)* 2003;**42**:591–7.

7 Pawłowicz M, Birkholz D, Niedźwiecki M, *et al.* Difficulties or mistakes in diagnosing type 1 diabetes mellitus in children? The consequences of delayed diagnosis. *Endokrynol Diabetol i Chor Przemiany Mater Wieku Rozw* 2008;**14**:7–12.

8 Soliman AT, ElZalabany MM, Bappal B, *et al.* Permanent neonatal diabetes mellitus: epidemiology, mode of presentation, pathogenesis and growth. *Indian J Pediatr* 1999;**66**:363–73.

9 Usher-Smith JA, Thompson M, Ercole A, *et al.* Variation between countries in the frequency of diabetic ketoacidosis at first presentation of type 1 diabetes in children: a systematic review. *Diabetologia* 2012;**55**:2878–94.

10 Abdul-Rasoul M, Habib H, Al-Khouly M. “The honeymoon phase” in children with type 1 diabetes mellitus: frequency, duration, and influential factors. *Pediatr Diabetes* 2006;**7**:101–7.

11 Bowden SA, Duck MM, Hoffman RP. Young children (< 5 yr) and adolescents (> 12 yr) with type 1 diabetes mellitus have low rate of partial remission: diabetic ketoacidosis is an important risk factor. *Pediatr Diabetes* 2008;**9**:197–201.

12 Fredheim S, Johannesen J, Johansen A, *et al.* Diabetic ketoacidosis at the onset of type 1 diabetes is associated with future HbA1c levels. *Diabetologia* 2013;**56**:995–1003.

13 Lokulo-Sodipe K, Moon RJ, Edge J a, *et al.* Identifying targets to reduce the incidence of diabetic ketoacidosis at diagnosis of type 1 diabetes in the UK. *Arch Dis Child* 2014;**99**:438–42.

14 Usher-Smith JA, Thompson MJ, Walter FM. “Looking for the needle in the haystack”: a qualitative study of the pathway to diagnosis of type 1 diabetes in children. *BMJ Open* 2013;**3**:e004068.

15 English Indices of Deprivation 2010. <https://www.gov.uk/government/publications/english-indices-of-deprivation-2010> (accessed 7 Feb2014).

16 Walter F, Webster A, Scott S, *et al.* The Andersen Model of Total Patient Delay: a systematic review of its application in cancer diagnosis. *J Health Serv Res Policy* 2012;**17**:110–8.

17 Scott SE, Walter FM, Webster a, *et al.* The model of pathways to treatment: conceptualization and integration with existing theory. *Br J Health Psychol* 2013;**18**:45–65.

18 Andersen BL, Cacioppo JT. Delay in seeking a cancer diagnosis: delay stages and psychophysiological comparison processes. *Br J Soc Psychol* 1995;**34**:33–52.

19 Allgar VL, Neal RD. Delays in the diagnosis of six cancers: analysis of data from the National Survey of NHS Patients: Cancer. *Br J Cancer* 2005;**92**:1959–70.

20 Neu A, Willasch A, Ehehalt S, *et al.* Ketoacidosis at onset of type 1 diabetes mellitus in children - Frequency and clinical presentation. *Pediatr Diabetes* 2003;**4**:77–81.



- 21 Roche EF, Menon A, Gill D, *et al.* Clinical presentation of type 1 diabetes. *Pediatr Diabetes* 2005;**6**:75–8.
- 22 Xin Y, Yang M, Chen XJ, *et al.* Clinical features at the onset of childhood type 1 diabetes mellitus in Shenyang, China. *J Paediatr Child Health* 2010;**46**:171–5.
- 23 Neu A, Ehehalt S, Willasch A, *et al.* Varying clinical presentations at onset of type 1 diabetes mellitus in children--epidemiological evidence for different subtypes of the disease? *Pediatr Diabetes* 2001;**2**:157–53.
- 24 Ting WH, Huang CY, Lo FS, *et al.* Clinical and laboratory characteristics of type 1 diabetes in children and adolescents: experience from a medical center. *Acta Paediatr Taiwanica* 2007;**48**:119–24.
- 25 Kapellen TM, Galler A, Nietzschmann U, *et al.* Prevalence of diabetic ketoacidosis in newly diagnosed children and adolescents with type 1 diabetes mellitus. Experience of a center for pediatric diabetology in Germany. [Article in German]. *Monatsschr Kinderheilkd* 2001;**149**:679–82.
- 26 Lévy-Marchal C, Patterson CC, Green A. Geographical variation of presentation at diagnosis of type I diabetes in children: The EURODIAB study. *Diabetologia* 2001;**44**:B75–80.
- 27 Neill SJ. Acute childhood illness at home: the parents' perspective. *J Adv Nurs* 2000;**31**:821–32.
- 28 Kai J. What worries parents when their preschool children are acutely ill, and why: a qualitative study. *Br Med J* 1996;**313**:983–6.
- 29 Dixon-Woods M, Findlay M, Young B, *et al.* Parents' accounts of obtaining a diagnosis of childhood cancer. *Lancet* 2001;**357**:670–4.
- 30 Francis N a, Crocker JC, Gamper A, *et al.* Missed opportunities for earlier treatment? A qualitative interview study with parents of children admitted to hospital with serious respiratory tract infections. *Arch Dis Child* 2011;**96**:154–9.
- 31 Forbes LJL, Atkins L, Thurnham A, *et al.* Breast cancer awareness and barriers to symptomatic presentation among women from different ethnic groups in East London. *Br J Cancer* 2011;**105**:1474–9.
- 32 Forbes LJL, Simon AE, Warburton F, *et al.* Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer* 2013;**108**:292–300.
- 33 Diabetes UK. 4 T's campaign. [https://www.diabetes.org.uk/Get\\_involved/Campaigning/Our-campaigns/4-Ts-campaign/](https://www.diabetes.org.uk/Get_involved/Campaigning/Our-campaigns/4-Ts-campaign/) (accessed 19 Aug2014).
- 34 Bui H, To T, Stein R, *et al.* Is diabetic ketoacidosis at disease onset a result of missed diagnosis? *J Pediatr* 2010;**156**:472–7.

35 Quinn M, Fleischman A, Rosner B, *et al.* Characteristics at diagnosis of type 1 diabetes in children younger than 6 years. *J Pediatr* 2006;**148**:366–71.

36 Pawlowicz M, Birkholz D, Niedzwiecki M, *et al.* Difficulties or mistakes in diagnosing type 1 diabetes in children?--demographic factors influencing delayed diagnosis. *Pediatr Diabetes* 2009;**10**:542–9.

For peer review only

## TABLES

Table 1. Child and family characteristics for those included in the study

Child and family characteristics	Number	Percentage (%)
<b>Gender</b>		
Male	49	56.3
Female	38	43.7
<b>Age</b>		
0-5	26	29.9
6-10	20	23.0
11-16	41	47.1
Mean $\pm$ SD	9.34 $\pm$ 4.5	
<b>Ethnicity</b>		
White	79	90.8
Asian	2	2.3
Black	3	3.4
Mixed	3	3.4
<b>Family history</b>		
First degree relative(s) with T1D	7	8.0
First degree relative(s) with T2D	8	9.2
Second or third degree relative(s) with T1D	13	14.9
Second or third degree relative(s) with T2D	24	27.6
<b>Indices of deprivation</b>		
Least deprived tertile	43	49.4
Middle tertile	33	37.9
Most deprived tertile	9	10.3
Missing	2	2.3
<b>Medically trained family member</b>	9	10.3
<b>DKA at diagnosis</b>		
Yes	35	40.2
No	52	60.0

Table 2. Frequency of symptoms amongst all children and those with and without DKA and duration of individual symptoms  
\* p < 0.05

	Frequency of symptoms						Duration of symptoms		
	All (n=87)		DKA (n=35)		No-DKA (n=52)		Mean ± SD	Median (IQR)	n
	n	%	n	%	n	%			
Polydipsia	85	97.7	33	94.3	52	100	31.9 ± 48	16 (8,36)	77
Polyuria	73	83.9	27	77.1	46	88.5	29.8 ± 53	14 (5,26)	65
Tiredness	66	75.9	28	80.0	38	73.1	34.5 ± 49.2	17 (10,39)	53
Nocturia	64	73.6	28	80.0	36	69.2	31.3 ± 52.1	15.5 (7,28.5)	56
Weight loss	56	64.4	28	80.0*	28	53.8*	50.1 ± 82.7	13.5 (7,44)	42
Changes in behaviour/mood	48	55.2	17	48.6	31	59.6	34.3 ± 40.8	15 (8,42)	34
Change in appetite	45	51.7	18	51.4	27	51.9	30.7 ± 48	14.5 (7,39)	38
Abdominal pain	37	42.5	17	48.6	20	38.5	41.4 ± 64.1	17 (7,38)	25
Nocturnal enuresis	33	37.9	14	40.0	19	36.5	28.4 ± 49.2	15 (5.5,21.5)	28
Different smelling breath	31	35.6	14	40.0	17	32.7	17.5 ± 28.7	6.5 (3,17)	22
Vomiting	17	19.5	15	42.9*	2	3.8*	7.3 ± 12.6	2.5 (1.5,5.5)	8
Faster breathing	15	17.2	12	34.3*	3	5.8*	3.8 ± 5.8	0.5 (0,7.5)	8
Urinary incontinence	14	16.1	4	11.4	10	19.2	36.6 ± 77.2	10 (3,21)	10
Fever	12	13.8	6	17.1	6	11.5	25 ± 35.8	8 (2,55)	7

Table 3. Duration of diagnostic intervals

	Mean $\pm$ SD (days)	Median (IQR) (days)	<i>n</i>
Appraisal Interval	41 $\pm$ 51.7	20 (9,40)	75
Help-seeking Interval	3 $\pm$ 4.6	1 (0,4.5)	83
Diagnostic Interval	5 $\pm$ 34.8	0 (0,0)	83
Total Diagnostic Interval	48 $\pm$ 60.4	25 (14,50)	74

Table 4. Parents' explanations for the ten most common symptoms

Symptom	<i>n</i>	Number with explanation for symptom <i>n</i> (%)	Most common explanations <i>n</i> (%)
Polydipsia	85	58 (68.2)	Hot weather 19 (32.8) Infection 13 (22.4) Activity/Travel 10 (17.2)
Polyuria	73	47 (64.4)	Drinking more 29 (61.7) Urine infection 6 (12.7) Diabetes 4 (8.5)
Tiredness	66	44 (66.7)	School related 12 (27.3) Infection 5 (11.9) Nocturia 4 (9.5)
Nocturia	64	40 (62.5)	Drinking more 26 (65.0) Diabetes 4 (10.0) Urine infection 3 (7.5)
Weight loss	56	33 (58.9)	Growth related 15 (45.5) Decreased appetite 4 (12.1) Increased activity 3 (9.1)
Changes in behaviour/mood	48	31 (64.6)	Tiredness 10 (32.3) Age related/puberty 7 (22.6) Infection/illness 6 (19.4)
Change in appetite	45	28 (62.2)	Growth related 14 (50.0) Infection 5 (17.9) Holiday related 2 (7.1)
Abdominal pain	37	19 (51.4)	Infection 4 (21.1) School related 3 (15.8) Period pains 3 (15.8)
Nocturnal enuresis	33	23 (69.7)	Drinking more 13 (56.5) Tired 4 (17.4) School related 3 (13.0)
Different smelling breath	31	14 (45.2)	Poor dental hygiene 4 (28.6) Infection 3 (21.4) Diabetes 3 (21.4)

Table 5. Factors influencing parents’ decisions to seek medical advice sooner or later

	Not at all <i>n</i> (%)	A little <i>n</i> (%)	Quite a lot <i>n</i> (%)	Very much <i>n</i> (%)	Did not answer <i>n</i> (%)
<b>Factors influencing seeking medical advice <i>sooner</i></b>					
Concern something serious	9 (10.3)	16 (18.4)	18 (20.7)	42 (48.3)	2 (2.3)
Symptoms getting worse	7 (8.0)	19 (21.8)	14 (16.1)	46 (52.9)	1 (1.1)
Symptoms not getting better	4 (4.6)	12 (13.8)	22 (25.3)	45 (51.7)	4 (4.6)
Wanting reassurance	8 (9.2)	15 (17.2)	16 (18.4)	46 (52.9)	2 (2.3)
Comments from family	30 (34.5)	28 (32.2)	11 (12.6)	13 (14.9)	5 (5.7)
Comments from school	63 (72.4)	10 (11.5)	4 (4.6)	4 (4.6)	6 (6.9)
Comments from friends	49 (56.3)	20 (23.0)	7 (8.0)	5 (5.7)	6 (6.9)
Written information	50 (57.5)	8 (9.2)	10 (11.5)	15 (17.2)	4 (4.6)
<b>Factors influencing seeking medical advice <i>later</i></b>					
Difficulty getting appointment	60 (69.0)	8 (9.2)	7 (8.0)	11 (12.6)	1 (1.1)
Waiting for a particular doctor or nurse	68 (78.2)	7 (8.0)	4 (4.6)	6 (6.9)	2 (2.3)
Concern about having to wait at the surgery	72 (82.8)	6 (6.9)	4 (4.6)	3 (3.4)	2 (2.3)
Worry about wasting the doctor or nurse’s time	61 (70.1)	10 (11.5)	8 (9.2)	6 (6.9)	2 (2.3)
Worry the doctor would not take them seriously	62 (71.3)	12 (13.8)	3 (3.4)	7 (8.0)	3 (3.4)
Symptoms weren't very serious	55 (63.2)	20 (23.0)	9 (10.3)	0 (0)	3 (3.4)
Hope the symptoms would go away	42 (48.3)	21 (24.1)	9 (10.3)	15 (17.2)	0 (0)
Fear of serious diagnosis	58 (66.7)	16 (18.4)	5 (5.7)	7 (8.0)	1 (1.1)

## The pathway to diagnosis of type 1 diabetes in children: a questionnaire study

Juliet A Usher-Smith<sup>1</sup>, Matthew J Thompson<sup>2</sup>, Hannah Zhu<sup>3</sup>, Stephen J Sharp<sup>4</sup>, Fiona M Walter<sup>5</sup>

<sup>1</sup> Clinical Lecturer in General Practice, The Primary Care Unit, University of Cambridge CB1 8RN

<sup>2</sup> Professor, Department of Family Medicine, University of Washington, Seattle, WA 98195 USA

<sup>3</sup> Foundation Year 2 Doctor, Cambridge University Hospitals NHS Foundation Trust, CB2 0QQ

<sup>4</sup> Senior Statistician, MRC Epidemiology Unit, University of Cambridge, Institute of Metabolic Science, Cambridge, CB2 0QQ

<sup>5</sup> GP & Clinician Scientist, The Primary Care Unit, University of Cambridge CB1 8RN

Correspondence to: J Usher-Smith jau20@medschl.cam.ac.uk

### ABSTRACT

**Objective** To explore the pathway to diagnosis of type 1 diabetes (T1D) in children

**Design** Questionnaire completed by parents

**Participants** Parents of children aged 1 month to 16 years diagnosed with T1D within the previous three months

**Setting** Children and parents from 11 hospitals within the East of England.

**Results** 88/164 (54%) of invited families returned the questionnaire. Children had mean±SD age of 9.41±4.5 years. 35 (39.8%) presented with DKA at diagnosis. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%) and all children presented with at least one of those symptoms. The time from symptom onset to diagnosis ranged from 2 to 315 days (median 25 days). Most of this was the appraisal interval from symptom onset until perceiving the need to seek medical advice. Access to healthcare was good but one in five children presenting to primary care were not diagnosed at first encounter, most commonly due to waiting for fasting blood tests or alternative diagnoses. Children diagnosed at first consultation had a shorter duration of symptoms ( $p=0.022$ ) and children whose parents suspected the diagnosis were 1.3 times more likely (RR 1.3, 95% CI 1.02-1.67) to be diagnosed at first consultation.

**Conclusions** Children present with the known symptoms of T1D but there is considerable scope to improve the diagnostic pathway. **Future interventions targeted at parents need to address the tendency of parents to find alternative explanations for symptoms and the perceived barriers to access, in addition to symptom awareness.**

### STRENGTHS AND LIMITATIONS OF THIS STUDY

- This study uses a questionnaire developed from a previous interview study to explore the diagnostic pathway of children with newly diagnosed T1D
- It uses the Model of Pathway to Treatment as a framework to allow analysis of the factors acting at different stages in the pathway
- The inclusion of a calendar with key events in the questionnaires and use of free text responses for internal validation and checking of prompted responses reduced bias but the data was necessarily collected retrospectively and so subject to recall and framing bias



**INTRODUCTION**

Approximately 65,000 children are diagnosed with type 1 diabetes (T1D) each year and the incidence is continuing to increase at a rate of approximately 3% per year<sup>1,2</sup>. The most common symptoms are well described and include polyuria, polydipsia, weight loss and tiredness. At the early stages of the disease, however, these symptoms are often non-specific and distinguishing the children with T1D from the large number with similar symptoms and minor undifferentiated illness can therefore be difficult. This is reflected in studies which have shown that the mean duration of symptoms prior to diagnosis is over two weeks with a significant number of children experiencing delay in diagnosis or misdiagnosis<sup>3</sup> and only one in five diagnosed at first encounter<sup>4-8</sup>. Up to 80% of children additionally present in diabetic ketoacidosis (DKA)<sup>9</sup> which has both immediate life-threatening complications and is associated with poorer long term diabetic control<sup>10-12</sup>.

Whilst several studies have highlighted these difficulties in making the diagnosis and the features associated with diabetic ketoacidosis at diagnosis<sup>3-8,13</sup>, few have explored the period between symptom onset and diagnosis. Our recent qualitative interview study of parents and General Practitioners (GPs) of children newly diagnosed with T1D suggested that the longest component in the diagnostic pathway is the time between onset of symptoms and the decision to seek medical help (known as the appraisal interval)<sup>14</sup>. The early symptoms are subtle, and even with some knowledge of T1D it took many parents several weeks of a complex decision making process and often a physical trigger, such as weight loss or vomiting, to decide to consult a healthcare professional. Once the decision to seek help had been made almost all children were seen immediately and diagnoses were mostly prompt and managed appropriately. Parents continued to play a key role during the diagnostic interval however, with many having already made or suspected the diagnosis themselves, and several feeling that their GP did not take their concerns seriously.

This study builds on this earlier work by using a questionnaire developed from the interview findings to further explore the pathway to diagnosis of T1D in children. By using a structured questionnaire to survey a larger number of families we aimed to quantify the symptoms and their time course prior to diagnosis, the triggers and barriers to seeking help, the influence of parental prior knowledge of diabetes, and the role of healthcare services.

**METHODS**

**Design**

A questionnaire about the pathway from first symptom(s) to diagnosis was completed by the parent(s)/guardian(s)/step-parents (hereafter referred to as parents) of children aged 1 month to 16 years diagnosed with T1D within the previous three months.

**Recruitment**

Children and parents were identified and recruited via the paediatric diabetes specialist nurses and research nurses at 11 hospitals within the East of England Diabetes Children and Young People's Network. Parents of all children aged 1 month to 16 years diagnosed with T1D diagnosed within the previous 3 months at participating hospitals were eligible for inclusion unless their clinical team felt that this was not appropriate. Parents who failed to respond within one month were sent a reminder letter with a further copy of the questionnaire. Recruitment began at each site between February 2013 and April 2013, and continued across all sites until January 2014.

The clinical or research teams at all sites collected data on the age and gender of each child diagnosed during the study period and whether they had DKA at diagnosis. Each hospital used a slightly different definition of DKA but all included either pH < 7.3, bicarbonate < 15 mmol/L (see Appendix Table 1).

### The questionnaire

The questionnaire was developed from the findings of our previous qualitative study of parents and children recently diagnosed with T1D<sup>14</sup>. It was first reviewed by an expert panel comprising paediatric diabetes consultants, a paediatric diabetes research nurse and primary care researchers, and then piloted with parents of four children recently diagnosed with T1D. In addition to their specific feedback, parents were asked to talk aloud whilst completing the questionnaire and then interviewed after completion to ensure face validity. Based on feedback from the parents, the questionnaire was revised.

The final questionnaire included 5 sections (see Supplementary file). The first included questions about the child's age, gender, postcode, ethnic background, family history of diabetes, any medically trained family members and the number of children in the household. Parents were also asked if they knew what the symptoms of diabetes in children are before their child was diagnosed, and if so, to give details of those symptoms they were aware of. The second section asked about the symptoms the children had experienced with yes/no responses for 14 symptoms and space to add the date they noticed the symptoms, what they thought the symptoms were due to at the time and how much it concerned them. The third section focused on help-seeking and asked where parents had looked for information, who they spoke to and then details on when and how they had sought medical advice. It also asked them to describe their main concern at their first appointment and whether they had considered diabetes. Parents were also asked in this section about factors contributing to their decision to seek medical advice sooner or later. The fourth section asked about the diagnosis, including whether it was made at their first appointment with a healthcare professional and, if not, how many subsequent consultations they had, and the investigations that were done before diagnosis. The final section then asked parents if they felt there was anything that prolonged them finding out their child had diabetes and had further space for free text comments.

### Analysis

Data from the questionnaires were entered into a database and then double checked by a second researcher. Socioeconomic status was computed using postcode and the English indices of deprivation 2010 available online<sup>15</sup>. The presence of DKA at diagnosis was obtained from hospital records rather than self-report. Walter et al's Model of Pathways to Treatment<sup>16,17</sup> provided a theoretic model of the intervals that occur prior to a diagnosis. This model divides the pathway to diagnosis into two intervals prior to presentation to healthcare about a symptom (the appraisal interval from the onset of symptoms to perceiving a reason to discuss symptoms with a healthcare professional, and the help-seeking interval from that decision until presentation to a healthcare professional), and then the diagnostic interval from first presentation to a healthcare professional until diagnosis. The help-seeking interval was further sub-divided into the behavioural interval (the time between perceiving the reason to discuss the symptoms with a healthcare professional to making the decision to seek help) and the scheduling interval (the time between making the decision to seek help and the first consultation)<sup>18</sup>. Intervals were calculated from responses to the questionnaire. Where dates were incomplete we applied midpoint rules to estimate the actual date<sup>19</sup>. In cases where the responses in free text differed from the dates entered as numbers, the free text was assumed to be correct, and where there was uncertainty the researchers met to agree consensus.

Characteristics (age, gender, presence of DKA) were compared between children whose parents had and had not returned a questionnaire using a t-test for age and chi-squared test for gender and presence of DKA. All further analyses used only data from returned questionnaires. The frequency of the 14 symptoms was compared between those with and without DKA using a chi-squared test. Cox regression was used to estimate the association between various factors and the hazard of

diagnosis; if a factor was associated with an increased hazard (i.e. hazard ratio greater than 1), this implied that that factor was associated with a shorter time to diagnosis, and vice versa. Time to diagnosis was from the date of the earliest symptom to the date of diagnosis, and the factors assessed were age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the parents suspected T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis. A similar approach was used to assess factors associated with the length of the appraisal and help-seeking intervals (with the end of the interval being defined as the “event” in the Cox model), but only the first six variables in the list above were considered, as the others do not relate to those time intervals. The Schoenfeld residuals test was used to assess the proportional hazards (PH) assumption for each covariate in each model. Whether parents suspected the diagnosis of T1D did not meet the PH assumption for the total diagnostic interval and so the Cox regression model was stratified by that variable. Logistic regression was used to estimate the association between the same factors and presence of DKA at diagnosis. All analyses were performed using STATA version 12.

Free text responses were grouped into similar categories and coded. Where individual free text responses contained several comments, these were each coded individually.

RESULTS

A total of 172 children were diagnosed with T1D in the 11 hospitals during the study period. Of those, 8 families were not invited to take part in the study: 5 lived outside the hospital catchment area; 1 emigrated the week after diagnosis; and the clinical team felt it was not appropriate to include 2. From the remaining 164 families invited to take part in the study, 88 (54%) completed and returned the questionnaire. There were no significant differences in the proportion presenting in DKA (p=0.27), mean age (p=0.77) or gender (p=0.77) between children of responders and non-responders.

One child was excluded from the analysis as they had no symptoms and the diagnosis was made on a random blood glucose test that the parents were doing at home on an intermittent basis as they had an older child with T1D. Children whose parents checked blood glucose at home after noticing symptoms remain in the analysis. 87 children are therefore included in the analysis that follows.

Table 1 shows the characteristics of the 87 children and families included in the study. The mean age was 9.34 ± 4.5 years, 49 (56.3%) were male and 35 (40.2%) presented with DKA at diagnosis. The majority (90.8%) were white and as a group they were generally from less deprived areas of England, with 49.4% from the least deprived tertile of English Indices of Deprivation and only 10.3% from the most deprived.

Symptoms

Table 2 shows the frequency and duration of the 14 symptoms that were specifically asked about in the questionnaire. The most common symptoms were polydipsia (97.7%), polyuria (83.9%), tiredness (75.9%), nocturia (73.6%) and weight loss (64.4%). Most symptoms were present for a median of between 13 and 17 days. Faster breathing and vomiting both had much shorter median (IQR) durations of 0.5 (0-7.5) and 2.5 (1.5-5.5) days respectively than the other symptoms. Weight loss, vomiting and faster breathing were significantly more frequent in those children who presented in DKA (p = 0.014, <0.0005 and 0.001 respectively). All the children had at least one of the 4 main symptoms (polydipsia, polyuria or nocturia, weight loss or tiredness), 97.7% had 2 or more, 79.3% 3 or more and over half (50.6%) had all 4 symptoms.

A very small number of parents mentioned symptoms other than those listed in the questionnaire, these included constipation (9), headaches (3), thrush (3), blurred vision (2), dry skin (2) and different smelling urine (1).

### Diagnostic intervals

Table 3 shows the mean  $\pm$  SD and median (IQR) for the diagnostic intervals. Additional details on the diagnostic intervals for different subgroups are shown in Appendix Table 2. The total diagnostic interval ranged from 2 to 315 days with a median (IQR) of 25 days (14-50). In unadjusted Cox regression analysis (data not shown) the time to diagnosis was significantly shorter for children diagnosed at first appointment compared to a subsequent appointment ( $p=0.046$ ) and for those seen in secondary care rather than primary care ( $p=0.01$ ). No evidence of associations with time to diagnosis was found for age, gender, family history of T1D, deprivation, prior knowledge of symptoms or DKA at diagnosis. In multivariable cox regression including age, gender, family history of T1D, index of multiple deprivation, prior knowledge of symptoms of T1D, whether the diagnosis was made at the first consultation, whether the first consultation was with primary or secondary care and whether the child had DKA at diagnosis (Figure 1a), the association between whether the diagnosis of T1D was made at the first or subsequent appointments and total diagnostic interval remained statistically significant ( $p=0.022$ ).

### The appraisal interval

The appraisal interval was the longest of all the intervals in the pathway for all but 3 of the families with a mean  $\pm$  SD of  $41.0 \pm 51.7$  days and median (IQR) 20 (9-40) days. During this period nearly two thirds (64%) of parents discussed the symptoms with family members, 40% with friends and 41% looked on the internet. Only 16% spoke to the child's nursery, school or playgroup and very few (6%) looked for information in books. Over half of parents (49, 56%) reported being aware of some symptoms of T1D in children prior to their child's diagnosis: 40 (45%) were aware of increased thirst, 24 (27%) of polyuria, 17 (19%) of weight loss and 13 (15%) tiredness.

Cox-regression analysis (Figure 1b) showed no significant associations between parent/child characteristics and the appraisal interval.

Analysis of the free text showed that most parents found explanations for their child's symptoms (Table 4). For example, polydipsia was attributed most commonly to hot weather (19/58, 33%) or infection (13/58, 22%), polyuria and nocturia were frequently explained by drinking more (29/47, 62% and 26/40, 65%) and tiredness was thought to be school related (12/44, 27%) or secondary to infection (5/44, 12%) or nocturia (4/44, 10%).

The majority of parents (61/87, 70%) additionally reported that they had suspected diabetes before their first consultation with a healthcare professional. When asked what had made them suspect diabetes, the most common reason given was that they knew the symptoms (22/59, 37%), especially thirst (12/59, 20%). Others cited information from the internet (12/59, 20%) or having a family history of diabetes (11/59, 19%).

### The help-seeking interval

24 (28%) children were seen on the same day their parents first thought about seeking medical advice and 64 (74%) within 5 days. Most of this time was the behavioural interval (mean  $\pm$  SD 2.1  $\pm$  3.7 days, median (IQR) 0 (0-3) days) rather than the scheduling interval (mean  $\pm$  SD 1.1  $\pm$  2.6 days, median (IQR) 0 (0-1) days).



Cox-regression analysis (Figure 1c) showed no significant associations between parent/child characteristics and the help-seeking interval.

The most common reasons that parents cited for seeking medical advice sooner rather than later (Table 5) were that the symptoms were not getting better or were getting worse, wanting reassurance or concern something serious was wrong. This was also reflected in the free text responses where 22% of parents noted that worsening or persistent symptoms was the reason they decided to seek help. In general, fewer parents reported factors that led to them seeking medical advice later. Of those that did, the most common reason for waiting was hope that the symptoms would go away (51.6%) but 29.8% felt difficulty getting an appointment contributed and 27.6% and 25.2% were worried about wasting the GPs time or that the GP would not take them seriously respectively.

**The diagnostic interval**

The diagnostic interval was the shortest of the intervals with a mean  $\pm$  SD of 5 days  $\pm$  34.8 and median 0 (IQR) (0-0) days. 69 (78%) of children were diagnosed at first consultation. Cox regression was not possible given the high number of children with a diagnostic interval of zero. However, children whose parents suspected the diagnosis (n=61, 70.1%) were more likely (unadjusted RR 1.30, 1.02-1.67, p=0.046) to be diagnosed at first consultation (n=52, 85.2%) than those in whom there was no suspicion (n=26, 29.9% with 17 (65.4%) diagnosed at first consultation). All children (10) who were seen first in secondary care were diagnosed at first consultation compared to 76.6% (59/77) of those seen first in primary care, but this difference was not statistically significant (p=0.114). None of the variables considered were significantly associated with risk of DKA (Figure 2).

Further details from the questionnaires were available from 14 of the 18 children who were not diagnosed at first encounter with primary care. Of these, 6 had fasting glucose blood tests arranged by the GP and 4 were given alternative diagnoses (urine infection, viral infection, tonsillitis, puberty) and diagnosed at a second appointment. Two children were diagnosed with psychological problems: In one case the child’s mother had seen the GP alone to discuss her child’s ‘obsessive drinking’ and was advised to see the school counsellor, and in the second the GP apparently felt the symptoms were psychological and the child was diagnosed in the emergency department four consultations later. One other family had already done a finger prick glucose test at home which was high but the GP did not trust the result and asked the child to come back later in the day with a urine sample. In the final case, the child’s mother had spoken to a health visitor and suggested diabetes but was told ‘no, not unless the child is lifeless’. The mother took the child to the GP 12 days later and the diagnosis was made at that consultation.

**DISCUSSION**

**Principal findings**

This study shows that all children with new onset T1D present with one, and 98% present with two, of the four main symptoms of diabetes (polydipsia, polyuria, weight loss and tiredness). Moreover, over half have had symptoms for over three weeks before diagnosis. Most of that time is the appraisal interval during which parents found alternative explanations for the symptoms, discussed the symptoms with family and friends and looked on the internet for information. Once they made the decision to seek advice, access to healthcare was generally not difficult with 28% consulting with a healthcare professional on the same day. However, when asked about factors contributing to their decision to seek help, nearly a third of parents felt that difficulty getting an appointment contributed to them waiting to seek help and over a quarter felt that worry about wasting the

doctor's time influenced their decision. This suggests that even if access is not difficult, it is perceived as such.

Once parents had sought help, one in five children were then not diagnosed at their first consultation with a healthcare professional, mainly due to being given an alternative diagnosis, most commonly infection, or waiting for further investigations. Diagnosis at first consultation was associated with a shorter total diagnostic interval and children were more likely to be diagnosed at first consultation when their parents suspected the diagnosis of T1D. The association between diagnosis at first consultation and total diagnostic interval may simply reflect the additional time between consultations, or it may be due to biological differences causing some children to develop symptoms more slowly which are then more difficult for both parents and primary care physicians to recognise.

### Strengths and weaknesses

By using a questionnaire developed from a previous interview study<sup>14</sup> and the Model of Pathway to Treatment<sup>16,17</sup> as a framework for analysis, this study provides in-depth insights into the diagnostic pathway of children with newly diagnosed T1D and allows factors acting at different stages in the pathway to be explored.

The main weakness is that the data was necessarily collected retrospectively and so subject to recall and framing bias. Parents have multiple contacts with different healthcare professionals in the period immediately following diagnosis and so their responses to the questionnaire reflect a post-hoc rationalisation of events framed by those subsequent encounters and increased knowledge since the diagnosis. The inclusion of a calendar with key events in the questionnaires minimised the error in recall of dates, and the free text responses allowed internal validation and checking of prompted responses. Despite these efforts, we still only have the parents' perspective on the pathway and were not able to confirm the number of healthcare contacts, diagnostic tests or the parental reports of missed opportunities for diagnosis. **We were, however, able to confirm the diagnosis of DKA from clinical records and, although there was variation in the definition of DKA used across the 11 sites, all included a biochemical measurement of either pH or bicarbonate.**

**Our results are also based on the views of 88 parents. Although not a large number, they were recruited from 11 sites across a large region of the UK and the response rate was over 50% with no significant differences in gender, age or DKA status between the children whose parents responded and those who did not. The fact that they were a predominantly white group from less deprived areas of England limits the generalizability of the results outside the East of England but the main findings are likely to be relevant across the UK and other countries with similar primary care healthcare provision. The questionnaire also did not include questions specifically for the children to complete and so we are unable to comment on the views of the children during this time.**

### Comparison with existing literature

The median duration of symptoms prior to diagnosis was 13-17 days for the nine most frequent symptoms, with a mean of 30-50 days. This is longer than previous studies relying on retrospective review of medical records<sup>20-23</sup> but similar to studies which have used a checklist to identify subtle symptoms<sup>24</sup> or asked parents soon after diagnosis<sup>13,14</sup>. The wide range (a few days to over six months) has been described previously<sup>14,21,23</sup> and highlights the heterogeneous nature of the disease.

The frequency of individual symptoms we report is also similar to previous studies<sup>13,14,20,22,25</sup>. Additionally we showed that all the children had at least one of 4 symptoms (polydipsia, polyuria, weight loss and fatigue) and over half (50.6%) had all four. Consistent with the known course of

the disease and previous studies, vomiting<sup>4,22,24</sup>, weight loss<sup>13,25,26</sup>, and dyspnoea<sup>22</sup> were more common in those children who presented in DKA.

This is the first quantitative study to compare the time periods during the pathway to diagnosis of T1D in children. The finding that most of the total diagnostic interval was the appraisal interval is consistent with a previous qualitative study<sup>14</sup> and the free text analysis confirms that during that time the parents find alternative explanations for the symptoms initially and make use of a social network of extended family, friends and work colleagues, or the internet<sup>14,27,28</sup>. That children were more likely to be diagnosed at their first encounter with a healthcare professional when their parents suspected diabetes prior to that consultation may also reflect the findings of previous qualitative work in which a number of parents prompted the GP to consider T1D and pushed for investigations<sup>14</sup>. However, whilst parental suspicion of T1D has also been shown to be associated with a reduced risk of DKA in a parental survey<sup>13</sup>, in that study the incidence of DKA at presentation was no different whether or not the parents discussed their concerns with the healthcare professional, suggesting other factors may be contributing. The absence of an effect of parental prior knowledge of diabetes either on the total diagnostic interval or the risk of DKA further highlights the complexities around the role of knowledge on help-seeking behaviour.

The finding that parents worry about wasting the doctor's time has also been shown in previous qualitative studies in children<sup>29,30</sup> and in studies of help-seeking behaviour for adults with symptoms of cancer in the UK<sup>31,32</sup> and so it may reflect a particular British trait rather than be specific to T1D or children.

**Implications for clinicians and policymakers**

Clinicians should remain alert to the possibility of T1D in all children presenting with one or more symptoms of polyuria, polydipsia, weight loss and tiredness – as almost all children have at least two of these. Interventions targeted at increasing public awareness, such as the 4 T's campaign launched by Diabetes UK to raise awareness of the four most common symptoms of T1D (Toilet, Thirsty, Tired and Thinner)<sup>33</sup>, should continue to focus on these established symptoms.

As most of the time between symptom onset and diagnosis is the appraisal interval, the greatest benefit is likely to be seen from interventions directed towards parents and their social network, probably via the internet. Despite ongoing government pressure for better access to primary care, improving access is unlikely to have much impact on the pathway. Instead efforts should be made to address the perception that access is difficult and the general concern in the UK about wasting healthcare professional time, particularly for children with acute or sub-acute health concerns.

Additionally, although the diagnostic interval itself was generally short, one in five children presenting to primary care were not diagnosed at first consultation. Similar numbers have been reported in a recent survey in the UK which found that 24% were not diagnosed at first contact with a healthcare professional<sup>13</sup>, and studies in the USA, Canada and Poland noted between 14 and 35% of children had more than one consultation before diagnosis<sup>6,7,34–36</sup>. As in those studies, the most common reasons for not being diagnosed at first encounter was either being given an alternative diagnosis, most commonly infection, or waiting for further investigations. In this study 33% of those not diagnosed at first consultation were waiting for fasting glucose tests and in other studies the number waiting for further investigations is as high as 46%<sup>6 13</sup>. This suggests that healthcare professionals may have considered a diagnosis of T1D but either lack ready access to rapid tests to confirm or exclude the diagnosis, or are reluctant to use existing tests in children<sup>14</sup>. Access to point of care urine and finger-prick testing and the use of those tests should be routine management for all children presenting with one or more of the four main symptoms of diabetes. The increased use of



point of care testing in Emergency departments may also explain why all children seen in secondary care were diagnosed at their first consultation. Whilst educational interventions aimed at primary care physicians may help a small number of children not currently diagnosed at first encounter, finding ways to overcome barriers to point-of-care tests in primary care may be more effective and this approach may also improve the diagnosis of other serious illnesses in children and adults.

### Unanswered questions and future research

Whilst this study contributes to our understanding of the pathway to diagnosis and the stages at which this may be improved, the findings are unable to explain the large variability in the overall duration of the pathway to diagnosis and why some children develop DKA within a few weeks whilst others can be symptomatic for up to six months before requiring treatment. Further studies are, therefore, needed into the natural course and biology of the disease to better understand these variations. The findings also highlight the need for continuing research into the presentation of serious but rare conditions in primary care and the best ways to improve diagnosis of these conditions.

### ACKNOWLEDGEMENTS

We thank all the parents who kindly gave up their time and shared their personal stories with us to either help develop the questionnaire or take part in the study. We also thank the Paediatric diabetes clinical teams at the 11 hospitals, particularly Jon Hassler-Hurst (Ipswich), Holly Roper (Norfolk and Norwich), Elissa Harwood (Cambridgeshire community services), Nazia Bhatti and Claire Pesterfield (Addenbrooke's hospital NHS foundation trust), Julia Harding and Jacky Plumb (West Suffolk), Philippa Corbishley and Laura Ardrey (QE2 and Lister), Suzie Williams (SEPT), Mandy Stevenson (Harlow), Claire Gibbs and Allyson Davison (James Paget) and Meeta Patel (Luton and Dunstable). We are also grateful to the Diabetes Research Network and the East of England Children and Young People's Diabetes Network and network manager Kate Wilson for support, James Brimicombe for advice on data management, Dr Carlo Acerini for comments on the questionnaire and final manuscript and support through his role as Chair of the East of England Children and Young People's Diabetes Network, Professor Jon Emery for comments on the questionnaire, and Professor David Dunger for his support and encouragement throughout the study and comments on the final manuscript.

### Contributors

JUS, MT, FMW and SJS were involved in the design of the study and all authors were involved in analysis of the data. JUS and FMW developed the questionnaire. JUS wrote the first draft of the manuscript and all authors reviewed and edited the manuscript.

### Funding

The study was funded by the Royal College of General Practitioners Scientific Foundation Board (SFB-2011-15). JUS was supported by a National Institute of Health Research (NIHR) Academic Clinical Fellowship and subsequently Clinical Lectureship, and FMW by an NIHR Clinician Scientist award. SJS was supported by the Medical Research Council [www.mrc.ac.uk](http://www.mrc.ac.uk) [Unit Programme number MC\_UU\_12015/1]. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

All researchers were independent of the funding body (although FMW subsequently became a member of the Royal College of General Practitioners Scientific Foundation Board), and the study sponsors and funder had no role in study design; data collection, analysis and interpretation of data; in the writing of the report; or decision to submit the article for publication.

**Ethics approval**

The study obtained ethical approval from the East of England Hertfordshire REC (reference number 12/EE/0390).

**Data sharing**

The questionnaire is available from the corresponding author on request.

**Competing Interests**

All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare that (1) JUS, MT, HZ, SJS and FMW have no support from or relationships with companies that might have an interest in the submitted work in the previous 3 years; (2) their spouses, partners, or children have no financial relationships that may be relevant to the submitted work; and (3) JUS, MT, HZ, SJS and FMW have no non-financial interests that may be relevant to the submitted work.

The corresponding author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in BMJ editions and any other BMJ PGL products and sublicences to exploit all subsidiary rights, as set out in our licence (<http://resources.bmj.com/bmj/authors/checklists-forms/licence-for-publication>).

All authors had full access to all of the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis

The corresponding author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

**REFERENCES**

- 1 Lawrence JM, Imperatore G, Dabelea D, *et al.* Trends in incidence of type 1 diabetes among non-Hispanic white youth in the u.s., 2002-2009. *Diabetes* 2014;**63**:3938–45.
- 2 Fazeli Farsani S, Souverein PC, van der Vorst MMJ, *et al.* Increasing trends in the incidence and prevalence rates of type 1 diabetes among children and adolescents in the Netherlands. *Pediatr Diabetes* Published Online First: 7 November 2014. doi:10.1111/pedi.12232
- 3 Usher-Smith JA, Thompson MJ, Sharp SJ, *et al.* Factors associated with the presence of diabetic ketoacidosis at diagnosis of diabetes in children and young adults: a systematic review. *Br Med J* 2011;**343**:d4092.
- 4 Blanc N, Lucidarme N, Tubiana-Rufi N. Factors associated to ketoacidosis at diagnosis of type 1 diabetes in children. *Arch Pediatr* 2003;**10**:320–5.
- 5 Hamilton D V, Mundia SS, Lister JC-1687347. Mode of presentation of juvenile diabetes. *Br Med J* 1976;**2**:211–2.
- 6 Mallare JT, Cordice CC, Ryan BA, *et al.* Identifying risk factors for the development of diabetic ketoacidosis in new onset type 1 diabetes mellitus. *Clin Pediatr (Phila)* 2003;**42**:591–7.

- 1  
2  
3 7 Pawłowicz M, Birkholz D, Niedźwiecki M, *et al.* Difficulties or mistakes in diagnosing type  
4 1 diabetes mellitus in children? The consequences of delayed diagnosis. *Endokrynol Diabetol*  
5 *i Chor Przemiany Mater Wieku Rozw* 2008;**14**:7–12.  
6  
7 8 Soliman AT, ElZalabany MM, Bappal B, *et al.* Permanent neonatal diabetes mellitus:  
9 epidemiology, mode of presentation, pathogenesis and growth. *Indian J Pediatr*  
10 1999;**66**:363–73.  
11  
12 9 Usher-Smith JA, Thompson M, Ercole A, *et al.* Variation between countries in the frequency  
13 of diabetic ketoacidosis at first presentation of type 1 diabetes in children: a systematic  
14 review. *Diabetologia* 2012;**55**:2878–94.  
15  
16 10 Abdul-Rasoul M, Habib H, Al-Khouly M. “The honeymoon phase” in children with type 1  
17 diabetes mellitus: frequency, duration, and influential factors. *Pediatr Diabetes* 2006;**7**:101–  
18 7.  
19  
20 11 Bowden SA, Duck MM, Hoffman RP. Young children (< 5 yr) and adolescents (> 12 yr)  
21 with type 1 diabetes mellitus have low rate of partial remission: diabetic ketoacidosis is an  
22 important risk factor. *Pediatr Diabetes* 2008;**9**:197–201.  
23  
24 12 Fredheim S, Johannesen J, Johansen A, *et al.* Diabetic ketoacidosis at the onset of type 1  
25 diabetes is associated with future HbA1c levels. *Diabetologia* 2013;**56**:995–1003.  
26  
27 13 Lokulo-Sodipe K, Moon RJ, Edge J a, *et al.* Identifying targets to reduce the incidence of  
28 diabetic ketoacidosis at diagnosis of type 1 diabetes in the UK. *Arch Dis Child* 2014;**99**:438–  
29 42.  
30  
31 14 Usher-Smith JA, Thompson MJ, Walter FM. “Looking for the needle in the haystack”: a  
32 qualitative study of the pathway to diagnosis of type 1 diabetes in children. *BMJ Open*  
33 2013;**3**:e004068.  
34  
35 15 English Indices of Deprivation 2010. [https://www.gov.uk/government/publications/english-](https://www.gov.uk/government/publications/english-indices-of-deprivation-2010)  
36 [indices-of-deprivation-2010](https://www.gov.uk/government/publications/english-indices-of-deprivation-2010) (accessed 7 Feb2014).  
37  
38 16 Walter F, Webster A, Scott S, *et al.* The Andersen Model of Total Patient Delay: a systematic  
39 review of its application in cancer diagnosis. *J Health Serv Res Policy* 2012;**17**:110–8.  
40  
41 17 Scott SE, Walter FM, Webster a, *et al.* The model of pathways to treatment:  
42 conceptualization and integration with existing theory. *Br J Health Psychol* 2013;**18**:45–65.  
43  
44 18 Andersen BL, Cacioppo JT. Delay in seeking a cancer diagnosis: delay stages and  
45 psychophysiological comparison processes. *Br J Soc Psychol* 1995;**34**:33–52.  
46  
47 19 Allgar VL, Neal RD. Delays in the diagnosis of six cancers: analysis of data from the  
48 National Survey of NHS Patients: Cancer. *Br J Cancer* 2005;**92**:1959–70.  
49  
50 20 Neu A, Willasch A, Ehehalt S, *et al.* Ketoacidosis at onset of type 1 diabetes mellitus in  
51 children - Frequency and clinical presentation. *Pediatr Diabetes* 2003;**4**:77–81.  
52  
53  
54  
55  
56  
57  
58  
59  
60

21 Roche EF, Menon A, Gill D, *et al.* Clinical presentation of type 1 diabetes. *Pediatr Diabetes* 2005;**6**:75–8.

22 Xin Y, Yang M, Chen XJ, *et al.* Clinical features at the onset of childhood type 1 diabetes mellitus in Shenyang, China. *J Paediatr Child Health* 2010;**46**:171–5.

23 Neu A, Ehehalt S, Willasch A, *et al.* Varying clinical presentations at onset of type 1 diabetes mellitus in children--epidemiological evidence for different subtypes of the disease? *Pediatr Diabetes* 2001;**2**:157–53.

24 Ting WH, Huang CY, Lo FS, *et al.* Clinical and laboratory characteristics of type 1 diabetes in children and adolescents: experience from a medical center. *Acta Paediatr Taiwanica* 2007;**48**:119–24.

25 Kapellen TM, Galler A, Nietzschmann U, *et al.* Prevalence of diabetic ketoacidosis in newly diagnosed children and adolescents with type 1 diabetes mellitus. Experience of a center for pediatric diabetology in Germany. [Article in German]. *Monatsschr Kinderheilkd* 2001;**149**:679–82.

26 Lévy-Marchal C, Patterson CC, Green A. Geographical variation of presentation at diagnosis of type I diabetes in children: The EURODIAB study. *Diabetologia* 2001;**44**:B75–80.

27 Neill SJ. Acute childhood illness at home: the parents' perspective. *J Adv Nurs* 2000;**31**:821–32.

28 Kai J. What worries parents when their preschool children are acutely ill, and why: a qualitative study. *Br Med J* 1996;**313**:983–6.

29 Dixon-Woods M, Findlay M, Young B, *et al.* Parents' accounts of obtaining a diagnosis of childhood cancer. *Lancet* 2001;**357**:670–4.

30 Francis N a, Crocker JC, Gamper A, *et al.* Missed opportunities for earlier treatment? A qualitative interview study with parents of children admitted to hospital with serious respiratory tract infections. *Arch Dis Child* 2011;**96**:154–9.

31 Forbes LJL, Atkins L, Thurnham A, *et al.* Breast cancer awareness and barriers to symptomatic presentation among women from different ethnic groups in East London. *Br J Cancer* 2011;**105**:1474–9.

32 Forbes LJL, Simon AE, Warburton F, *et al.* Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer* 2013;**108**:292–300.

33 Diabetes UK. 4 T's campaign. [https://www.diabetes.org.uk/Get\\_involved/Campaigning/Our-campaigns/4-Ts-campaign/](https://www.diabetes.org.uk/Get_involved/Campaigning/Our-campaigns/4-Ts-campaign/) (accessed 19 Aug2014).

34 Bui H, To T, Stein R, *et al.* Is diabetic ketoacidosis at disease onset a result of missed diagnosis? *J Pediatr* 2010;**156**:472–7.

- 1  
2  
3 35 Quinn M, Fleischman A, Rosner B, *et al.* Characteristics at diagnosis of type 1 diabetes in  
4 children younger than 6 years. *J Pediatr* 2006;**148**:366–71.  
5  
6 36 Pawlowicz M, Birkholz D, Niedzwiecki M, *et al.* Difficulties or mistakes in diagnosing type  
7 1 diabetes in children?--demographic factors influencing delayed diagnosis. *Pediatr Diabetes*  
8 2009;**10**:542–9.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

TABLES

Table 1. Child and family characteristics for those included in the study

Child and family characteristics	Number	Percentage (%)
<b>Gender</b>		
Male	49	56.3
Female	38	43.7
<b>Age</b>		
0-5	26	29.9
6-10	20	23.0
11-16	41	47.1
Mean ± SD	9.34 ± 4.5	
<b>Ethnicity</b>		
White	79	90.8
Asian	2	2.3
Black	3	3.4
Mixed	3	3.4
<b>Family history</b>		
First degree relative(s) with T1D	7	8.0
First degree relative(s) with T2D	8	9.2
Second or third degree relative(s) with T1D	13	14.9
Second or third degree relative(s) with T2D	24	27.6
<b>Indices of deprivation</b>		
Least deprived tertile	43	49.4
Middle tertile	33	37.9
Most deprived tertile	9	10.3
Missing	2	2.3
<b>Medically trained family member</b>	9	10.3
<b>DKA at diagnosis</b>		
Yes	35	40.2
No	52	60.0



Table 2. Frequency of symptoms amongst all children and those with and without DKA and duration of individual symptoms

\*  $p < 0.05$ 

	Frequency of symptoms						Duration of symptoms		
	All ( <i>n</i> =87)		DKA ( <i>n</i> =35)		No-DKA ( <i>n</i> =52)		Mean ± SD	Median (IQR)	<i>n</i>
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%			
Polydipsia	85	97.7	33	94.3	52	100	31.9 ± 48	16 (8,36)	77
Polyuria	73	83.9	27	77.1	46	88.5	29.8 ± 53	14 (5,26)	65
Tiredness	66	75.9	28	80.0	38	73.1	34.5 ± 49.2	17 (10,39)	53
Nocturia	64	73.6	28	80.0	36	69.2	31.3 ± 52.1	15.5 (7,28.5)	56
Weight loss	56	64.4	28	80.0*	28	53.8*	50.1 ± 82.7	13.5 (7,44)	42
Changes in behaviour/mood	48	55.2	17	48.6	31	59.6	34.3 ± 40.8	15 (8,42)	34
Change in appetite	45	51.7	18	51.4	27	51.9	30.7 ± 48	14.5 (7,39)	38
Abdominal pain	37	42.5	17	48.6	20	38.5	41.4 ± 64.1	17 (7,38)	25
Nocturnal enuresis	33	37.9	14	40.0	19	36.5	28.4 ± 49.2	15 (5.5,21.5)	28
Different smelling breath	31	35.6	14	40.0	17	32.7	17.5 ± 28.7	6.5 (3,17)	22
Vomiting	17	19.5	15	42.9*	2	3.8*	7.3 ± 12.6	2.5 (1.5,5.5)	8
Faster breathing	15	17.2	12	34.3*	3	5.8*	3.8 ± 5.8	0.5 (0,7.5)	8
Urinary incontinence	14	16.1	4	11.4	10	19.2	36.6 ± 77.2	10 (3,21)	10
Fever	12	13.8	6	17.1	6	11.5	25 ± 35.8	8 (2,55)	7

Table 3. Duration of diagnostic intervals

	Mean ± SD (days)	Median (IQR) (days)	<i>n</i>
Appraisal Interval	41 ± 51.7	20 (9,40)	75
Help-seeking Interval	3 ± 4.6	1 (0,4.5)	83
Diagnostic Interval	5 ± 34.8	0 (0,0)	83
Total Diagnostic Interval	48 ± 60.4	25 (14,50)	74

Table 4. Parents’ explanations for the ten most common symptoms

Symptom	<i>n</i>	Number with explanation for symptom <i>n</i> (%)	Most common explanations <i>n</i> (%)	
Polydipsia	85	58 (68.2)	Hot weather	19 (32.8)
			Infection	13 (22.4)
			Activity/Travel	10 (17.2)
Polyuria	73	47 (64.4)	Drinking more	29 (61.7)
			Urine infection	6 (12.7)
Tiredness	66	44 (66.7)	Diabetes	4 (8.5)
			School related	12 (27.3)
			Infection	5 (11.9)
Nocturia	64	40 (62.5)	Nocturia	4 (9.5)
			Drinking more	26 (65.0)
			Diabetes	4 (10.0)
Weight loss	56	33 (58.9)	Urine infection	3 (7.5)
			Growth related	15 (45.5)
			Decreased appetite	4 (12.1)
Changes in behaviour/mood	48	31 (64.6)	Increased activity	3 (9.1)
			Tiredness	10 (32.3)
			Age related/puberty	7 (22.6)
Change in appetite	45	28 (62.2)	Infection/illness	6 (19.4)
			Growth related	14 (50.0)
			Infection	5 (17.9)
Abdominal pain	37	19 (51.4)	Holiday related	2 (7.1)
			Infection	4 (21.1)
			School related	3 (15.8)
Nocturnal enuresis	33	23 (69.7)	Period pains	3 (15.8)
			Drinking more	13 (56.5)
			Tired	4 (17.4)
Different smelling breath	31	14 (45.2)	School related	3 (13.0)
			Poor dental hygiene	4 (28.6)
			Infection	3 (21.4)
			Diabetes	3 (21.4)

Table 5. Factors influencing parents' decisions to seek medical advice sooner or later

	Not at all <i>n</i> (%)	A little <i>n</i> (%)	Quite a lot <i>n</i> (%)	Very much <i>n</i> (%)	Did not answer <i>n</i> (%)
<b>Factors influencing seeking medical advice <i>sooner</i></b>					
Concern something serious	9 (10.3)	16 (18.4)	18 (20.7)	42 (48.3)	2 (2.3)
Symptoms getting worse	7 (8.0)	19 (21.8)	14 (16.1)	46 (52.9)	1 (1.1)
Symptoms not getting better	4 (4.6)	12 (13.8)	22 (25.3)	45 (51.7)	4 (4.6)
Wanting reassurance	8 (9.2)	15 (17.2)	16 (18.4)	46 (52.9)	2 (2.3)
Comments from family	30 (34.5)	28 (32.2)	11 (12.6)	13 (14.9)	5 (5.7)
Comments from school	63 (72.4)	10 (11.5)	4 (4.6)	4 (4.6)	6 (6.9)
Comments from friends	49 (56.3)	20 (23.0)	7 (8.0)	5 (5.7)	6 (6.9)
Written information	50 (57.5)	8 (9.2)	10 (11.5)	15 (17.2)	4 (4.6)
<b>Factors influencing seeking medical advice <i>later</i></b>					
Difficulty getting appointment	60 (69.0)	8 (9.2)	7 (8.0)	11 (12.6)	1 (1.1)
Waiting for a particular doctor or nurse	68 (78.2)	7 (8.0)	4 (4.6)	6 (6.9)	2 (2.3)
Concern about having to wait at the surgery	72 (82.8)	6 (6.9)	4 (4.6)	3 (3.4)	2 (2.3)
Worry about wasting the doctor or nurse's time	61 (70.1)	10 (11.5)	8 (9.2)	6 (6.9)	2 (2.3)
Worry the doctor would not take them seriously	62 (71.3)	12 (13.8)	3 (3.4)	7 (8.0)	3 (3.4)
Symptoms weren't very serious	55 (63.2)	20 (23.0)	9 (10.3)	0 (0)	3 (3.4)
Hope the symptoms would go away	42 (48.3)	21 (24.1)	9 (10.3)	15 (17.2)	0 (0)
Fear of serious diagnosis	58 (66.7)	16 (18.4)	5 (5.7)	7 (8.0)	1 (1.1)

Figure 1a. Associations between parent/child characteristics and the total diagnostic interval. IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure. Cox model stratified by whether parents suspected the diagnosis or not.




Figure 1a  
125x90mm (300 x 300 DPI)

Figure 1b. Associations between parent/child characteristics and the appraisal interval.  
IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure




Figure 1b  
122x86mm (300 x 300 DPI)

Figure 1c. Associations between parent/child characteristics and the help-seeking interval.  
IMD – Index of multiple deprivation. Hazard ratios adjusted for all variables in the figure




Figure 1c  
122x85mm (300 x 300 DPI)



Figure 2. Associations between parent/child characteristics and presence/absence of DKA.  
IMD – index of multiple deprivation. Odds ratios adjusted for all variables in the figure




Figure 2  
124x89mm (300 x 300 DPI)

Appendix Table 1. Definitions of DKA used across the 11 hospital sites

Hospital	Glucose > 11mmol/L	Ketones > 3mmol/L	pH < 7.3	pH < 7.3 AND bicarb < 15mmol/L	pH < 7.3 OR bicarb < 15mmol/L	Ketones in urine or blood	Glucose and ketones in urine	Symptoms	Treated according to DKA protocol
1	•		•			•			
2	•	•			•		•		
3	•	•			•				
4	•			•				•	
5					•				
6	•				•				
7	•	•			•				
8	•			•					•
9	•			•					
10	•	•			•				
11	•				•				

	Appraisal interval			Help-seeking interval			Diagnostic interval			Total diagnostic interval		
	Mean ± SD	Median (IQR)	n	Mean ± SD	Median (IQR)	n	Mean ± SD	Median (IQR)	n	Mean ± SD	Median (IQR)	n
<b>All</b>	41 ± 51.7	20 (9,40)	75	3 ± 4.6	1 (0,4.5)	83	5 ± 34.8	0 (0,0)	83	48 ± 60.4	25 (14,50)	74
<b>Age</b>												
0-5 years	47.8 ± 53.8	35 (11,67)	23	2.4 ± 2.9	1 (0,3)	25	1.1 ± 3.2	0 (0,0)	24	47.7 ± 51.9	36 (15,64)	21
6-10 years	29.2 ± 40.7	15.5 (11,31)	18	2.5 ± 3.7	1 (0,3)	18	16.1 ± 69.0	0 (0,0)	20	47.3 ± 75.6	21 (15,36)	19
11-16 years	38.5 ± 49.9	17 (7,38)	34	4.1 ± 5.6	3 (0,5)	39	2.2 ± 7.4	0(0,0)	38	45.5 ± 51.9	22 (11,50)	34
<b>Gender</b>												
Male	43.8 ± 56.6	20 (10,53)	43	2.9 ± 4.7	1 (0,4)	47	1.8 ± 6.5	0 (0,0)	45	46.9 ± 56.9	22 (13,59)	41
Female	32.8 ± 36.2	19.5 (7,38)	32	3.7 ± 4.5	1 (1,7)	35	9.4 ± 50.8	0 (0,0)	37	46.2 ± 60.2	29 (15,45)	33
<b>Family history of T1D</b>												
No FH	37.7 ± 51.8	19 (8,37)	41	4 ± 5.5	2 (1,5)	46	1.8 ± 6.4	0 (0,0)	46	44.6 ± 52.6	23 (15,50)	41
1st degree relative	42.5 ± 54.3	29 (11,37)	13	1.5 ± 2.5	0 (0,3)	14	0.8 ± 2.2	0 (0,0)	13	32.8 ± 47.8	15.5 (9,38.5)	12
2nd or 3rd degree relative	39.7 ± 41.7	32 (12,56)	21	2.8 ± 2.8	1.5 (1,5)	22	14.7 ± 64.3	0 (0,0)	23	58.3 ± 72.3	40 (17,64)	21
<b>Prior knowledge of symptoms</b>												
Yes	38.4 ± 44.7	20 (11,42)	46	3.7 ± 5.3	1 (0,6)	47	1.1 ± 3.7	0(0,0)	48	42.3 ± 46.5	23 (14,49)	47
No	42.7 ± 57.3	22 (7,53)	27	2.8 ± 3.5	2 (0,3)	33	11.6 ± 54.8	0 (0,0)	32	57.5 ± 76.3	28 (15,64)	25
<b>Deprivation</b>												
Low	33.4 ± 18.9	31 (27,38)	9	3.8 ± 4.7	2 (0,7)	9	2.3 ± 7.3	0 (0,0)	42	36.8 ± 36.8	20 (11,58)	37
Middle	46.9 ± 62.1	19 (11,38)	29	2.2 ± 2.8	1 (0,3)	30	11.6 ± 57.3	0 (0,0)	29	58.1 ± 79.1	27 (15, 49)	27
High	30.3 ± 34.5	14 (6.5, 47)	36	3.9 ± 5.5	2 (0,6)	41	0 ± 0	0 (0,0)	9	37.2 ± 17.8	36 (28,40)	9
<b>Parents considered diabetes</b>												
Yes	40.2 ± 45.8	23.5 (12,50)	54	3.8 ± 5	2 (1,6.5)	60	1 ± 3.4	0 (0,0)	60	45.3 ± 47.3	28.5 (16,50)	54
No	36.3 ± 57.6	14 (7,37)	21	1.7 ± 2.6	0.5 (0,3)	22	17 ± 65.8	0 (0,2)	22	50.0 ± 81.7	19 (7,50)	20
<b>Diagnosis at first appointment</b>												
Yes	35.2 ± 45.4	20 (11,38)	61	3.0 ± 3.7	1.5 (0,5)	66	1.1 ± 2.4	0 (0,0)	32	38.4 ± 44.8	22.5 (14,41)	62
No	56.1 ± 61.3	25.5 (7,92)	14	4.1 ± 7.2	1(0,5)	16	7.9 ± 44	0 (0,0)	50	89.0 ± 94.1	61.6 (15, 145)	12
<b>First contact with healthcare</b>												
Primary care	42.5 ± 51.0	23.5 (11,53)	66	3.5 ± 4.7	2 (1,5)	73	5.9 ± 36.4	0 (0,0)	73	51.0 ± 60.4	28 (16,59)	65
Secondary care	14.4 ± 16.4	8 (4,16)	9	0.7 ± 1.3	0 (0,1)	9	0 ± 0	0 (0,0)	9	15.1 ± 16.4	11 (4,17)	9
<b>DKA</b>												
Yes	42.5 ± 59.4	21 (7,36.5)	28	2.9 ± 5.6	1 (0,3)	34	0.1 ± 0.7	0 (0,0)	68	43 ± 59.6	21 (13,36)	26
No	37.1 ± 42.2	20 (11,50)	47	3.4 ± 3.7	2 (1,5.5)	48	30.4 ± 80.9	6.5 (2,12)	14	48.5 ± 57.7	35.5 (14.5, 61.5)	48

Appendix Table 2. Time intervals along the pathway to diagnosis.

For peer review only

# Pathway to diagnosis of type 1 diabetes in children questionnaire

We are interested in your experience of the time before your child was diagnosed with diabetes. We know that recognising the symptoms of diabetes is difficult and we really want to know what you noticed and what made you first ask for medical advice from a doctor or nurse. The symptoms are different for every child so don't worry if your child didn't have all the symptoms mentioned. We are hoping to also find out how long children have symptoms for before they are diagnosed so please try to add dates wherever possible and be as accurate as you can. We have included a calendar on the next page with school holidays and bank holidays on it so please use this and your own diaries to help you remember.

## Section 1 - Information about your child and family

In this section we are interested in details about your child and family. This allows us to make sure that we have included children of different ages and from different places to make the results as useful as possible.

Gender of your child: Male ☐ Female ☐ Date of birth of your child: DD/MM/YY  
 Ethnic background? White ☐ Asian ☐ Black ☐ Chinese ☐ Mixed ☐ Other \_\_\_\_\_

Does your child have any other medical problems? Yes ☐ No ☐ (If yes, please give details)  
 \_\_\_\_\_  
 \_\_\_\_\_

Does your child take any medication other than for diabetes? Yes ☐ No ☐ (If yes, please give details)  
 \_\_\_\_\_  
 \_\_\_\_\_

Does anyone else in the family have diabetes?

	Type 1	Type 2
Child's parent (s)	<input type="checkbox"/>	<input type="checkbox"/>
Child's brother or sister?	<input type="checkbox"/>	<input type="checkbox"/>
Other (please specify)		
_____	<input type="checkbox"/>	<input type="checkbox"/>
_____	<input type="checkbox"/>	<input type="checkbox"/>

What are the child's parents' current occupations?

Child's mother \_\_\_\_\_  
 Child's father \_\_\_\_\_

What is your postcode?

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Is anyone in the family medical / healthcare trained? e.g. a doctor, a nurse or a paramedic

Yes ☐ No ☐ (If yes, please give details)  
 \_\_\_\_\_  
 \_\_\_\_\_

Before your child was diagnosed did you know what the symptoms of diabetes in children are?

Yes ☐ No ☐ (If yes, please give details of those symptoms you knew of)  
 \_\_\_\_\_  
 \_\_\_\_\_

How many other children live in the same house as your child?

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4+ ☐



Calendar

This calendar shows the school holidays and bank holidays. We have included it to help you remember when you noticed symptoms and other dates relating to your child’s diagnosis. You may find it helpful to add important family dates such as birthdays and holidays.

### August 2012

Mon	Tue	Wed	Thu	Fri	Sat	Sun
30	31	1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31	1	2
3	4	5	6	7	8	9

27<sup>th</sup> - Bank holiday  
27<sup>th</sup> July – 12<sup>th</sup> August - Olympics

### September 2012

Mon	Tue	Wed	Thu	Fri	Sat	Sun
27	28	29	30	31	1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
1	2	3	4	5	6	7

3<sup>rd</sup> – school term starts

### October 2012

Mon	Tue	Wed	Thu	Fri	Sat	Sun
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31	1	2	3	4
5	6	7	8	9	10	11

29<sup>th</sup> Oct – 2<sup>nd</sup> Nov - half term  
31<sup>st</sup> - Halloween

### November 2012

Mon	Tue	Wed	Thu	Fri	Sat	Sun
29	30	31	1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	1	2
3	4	5	6	7	8	9

29<sup>th</sup> Oct – 2<sup>nd</sup> Nov - half term  
5<sup>th</sup> – Guy Fawkes day

### December 2012

Mon	Tue	Wed	Thu	Fri	Sat	Sun
26	27	28	29	30	1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31	1	2	3	4	5	6

21<sup>st</sup> - school term ends  
25<sup>th</sup> – Christmas Day

### January 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
31	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31	1	2	3
4	5	6	7	8	9	10

8<sup>th</sup> – school term starts

### February 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
28	29	30	31	1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	1	2	3
4	5	6	7	8	9	10

11<sup>th</sup> – 15<sup>th</sup> Half term

### March 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
25	26	27	28	1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31
1	2	3	4	5	6	7

28<sup>th</sup> - school term ends  
31<sup>st</sup> - Easter Sunday

### April 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	1	2	3	4	5
6	7	8	9	10	11	12

1<sup>st</sup> – Easter Monday  
15<sup>th</sup> – school term starts

### May 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
29	30	1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31	1	2
3	4	5	6	7	8	9

27<sup>th</sup> – 31<sup>st</sup> - half term  
6<sup>th</sup> and 27<sup>th</sup> - Bank holidays  
11<sup>th</sup> – FA cup final

### June 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
27	28	29	30	31	1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
1	2	3	4	5	6	7

### July 2013

Mon	Tue	Wed	Thu	Fri	Sat	Sun
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31	1	2	3	4
5	6	7	8	9	10	11

23<sup>rd</sup> - school term ends



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Section 2 – Information about the symptoms you noticed before your child was diagnosed**

In this section we are interested in all the symptoms your child had in the weeks or months leading up to when they were diagnosed with diabetes.

For each symptom please tick yes or no to indicate whether you noticed that symptom. If you did notice it, please add the date you first noticed it and what you thought the symptom was due to at the time.

Symptom	Did you notice this symptom?	If yes, when did you notice this symptom?	What did you think the symptom was due to at the time?
Drinking more than usual	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Weeing (passing urine) more than usual	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Changes in appetite	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Going to the loo at night more than usual	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Being more tired than usual	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Wetting the bed at night	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Losing weight	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Vomiting	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Having accidents when passing urine	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Tummy pain	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Fever	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Constipation	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Different smelling breath	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Skin infections	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Faster breathing	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
Other changes in behaviour / mood (please give details)	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
_____			
_____			
Other (please specify)	YES <input type="checkbox"/> NO <input type="checkbox"/>	DD/MM/YY	
_____			
_____			



Still thinking about the symptoms that you noticed at the time before your child was diagnosed with diabetes, when you first noticed each symptom, how much did each of them concern you?

Symptom	Not applicable, my child did not have this symptom	Not at all	A little	Quite a lot	Very much
Drinking more than usual	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Weeing (passing urine) more than usual	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Changes in appetite	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Going to the loo at night more than usual	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Being more tired than usual	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Wetting the bed at night	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Losing weight	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Vomiting	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Having accidents when passing urine	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Tummy pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Fever	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Constipation	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Different smelling breath	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Skin infections	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Faster breathing	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Other changes in behaviour/mood	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Other (please specify)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Section 3 – Information about what made you decide to seek medical advice

In this section we are interested in what made you decide to seek medical advice and where you went for that advice.

Did you look for information about the symptoms your child had in any of the places below?

Books	<input type="checkbox"/>	Magazines	<input type="checkbox"/>
The internet	<input type="checkbox"/>	Other _____	<input type="checkbox"/>

If you have ticked any of the boxes above, please give details below of where you looked and what information you read \_\_\_\_\_

Did you discuss the symptoms your child had with any of the following groups of people? (Please tick all that apply)

Family members	<input type="checkbox"/>	School / nursery / play group	<input type="checkbox"/>
Friends	<input type="checkbox"/>	Other _____	<input type="checkbox"/>

If you have ticked any of the boxes above, please give details below of who you spoke to and what advice they gave \_\_\_\_\_

When did you first think about seeking medical advice about the symptoms? DD/MM/YY

When did you decide to seek medical advice about the symptoms? DD/MM/YY

What was it that made you decide to seek medical advice then?

Where did you go **first** for that medical advice?

GP	<input type="checkbox"/>	Out of hours GP	<input type="checkbox"/>	Emergency department	<input type="checkbox"/>
NHS Direct	<input type="checkbox"/>	Health visitor	<input type="checkbox"/>	Other _____	<input type="checkbox"/>
Pharmacy	<input type="checkbox"/>	Minor illness centre	<input type="checkbox"/>		

Where did you **first** see a doctor or nurse?

GP	<input type="checkbox"/>	Out of hours GP	<input type="checkbox"/>	Emergency department	<input type="checkbox"/>
Minor illness centre	<input type="checkbox"/>	Other _____	<input type="checkbox"/>		

When was that first appointment with a doctor or nurse? DD/MM/YY

What was the main concern that you mentioned at that first appointment?

Had you considered diabetes at that stage? Yes ☐ No ☐ If yes, please explain what had made you think it might be diabetes



Thinking about your decision to seek medical advice, how much do you think each of the following made you seek medical advice sooner?

Not at all      A little      Quite a lot      Very much

Concern there was something serious wrong

☐
☐
☐
☐

The symptoms were getting worse

☐
☐
☐
☐

The symptoms were not getting any better

☐
☐
☐
☐

Wanting reassurance from a doctor

☐
☐
☐
☐

Comments from other family members

☐
☐
☐
☐

Comments from school

☐
☐
☐
☐

Comments from friends

☐
☐
☐
☐

Written information from books, magazines, posters or the internet

☐
☐
☐
☐

And how much do you think each of the following made you wait and seek medical advice later?

Not at all      A little      Quite a lot      Very much

Difficulty getting an appointment with a doctor or nurse

☐
☐
☐
☐

Waiting to get an appointment with a particular doctor or nurse

☐
☐
☐
☐

Concern about having to wait at the surgery to see a doctor or nurse

☐
☐
☐
☐

Worry about wasting the time of the doctor or nurse

☐
☐
☐
☐

Fear of getting a serious diagnosis

☐
☐
☐
☐

Worry that the doctor would not take you seriously

☐
☐
☐
☐

The symptoms weren't very serious

☐
☐
☐
☐

Hope that the symptoms would go away

☐
☐
☐
☐



Section 4 – Information about the diagnosis

In this section we are interested who made the diagnosis, how the diagnosis was made and how your child was at the time.

Was the diagnosis of diabetes made or suggested at that first appointment? Yes ☐ No ☐

If not, how many more times did you see a health professional before you were told your child might have diabetes?

1 ☐ 2 ☐ 3 ☐ 4 ☐ 5+ ☐

When were you told your child had diabetes? DD/MM/YY

Who told you your child might have diabetes?

GP	<input type="checkbox"/>	Out of hours GP	<input type="checkbox"/>	Emergency department	<input type="checkbox"/>
Hospital doctor	<input type="checkbox"/>	Health visitor	<input type="checkbox"/>	Other	_____

Which of the following tests did your child have before the diagnosis?

Urine dipstick	<input type="checkbox"/>	Finger prick blood test	<input type="checkbox"/>	Fasting blood test	<input type="checkbox"/>
----------------	--------------------------	-------------------------	--------------------------	--------------------	--------------------------

Did your child need to have fluids through a tube (a drip)? Yes ☐ No ☐

How long did your child stay in hospital after the diagnosis was made? \_\_\_\_\_ nights

Did your child have diabetic ketoacidosis? Yes ☐ No ☐ I don't know ☐

Section 5 – Other information

In your opinion, do you feel there was anything that prolonged you finding out that your child has diabetes? (Please continue over the page if you need more space)

Do you have any other comments about the symptoms your child had or how the diagnosis was made? (Please continue over the page if you need more space)

Thank you very much for completing this questionnaire.

Please now put it in the pre-paid envelope and return it to:  
For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>  
Dr Juliet Usher-Smith, University of Cambridge, Dept. Public Health & Primary Care, Strangeways  
Research Laboratory, 2 Worts Causeway, Cambridge CB1 8RN.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60