

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

Using Structural Equation Modeling to Explore Associated Factors of Intimate Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019465
Article Type:	Research
Date Submitted by the Author:	05-Sep-2017
Complete List of Authors:	Hou, Fengsu; Sun Yat-sen Global Health Institute, School of Public Health, Sun Yat-sen University; Department of Psychiatry, University of Rochester Medical Center Cerulli, Catherine; Injury Control Research Center for Suicide Prevention, University of Rochester Medical Center; Department of Psychiatry, University of Rochester Medical Center Wittink, Marsha; Department of Psychiatry, University of Rochester Medical Center Caine, Eric; Injury Control Research Center for Suicide Prevention; Department of Psychiatry, University of Rochester Medical Center Qiu, Peiyuan; Department of Epidemiology and Health Statistics, West China School of Public Health, Sichuan University
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Sociology, Legal and forensic medicine
Keywords:	Intimate partner violence, Violence against women, Rural Chinese women, Social support

SCHOLARONE™
Manuscripts

1
2
3 **1 Using Structural Equation Modeling to Explore Associated Factors of Intimate**
4 **2 Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study**

5
6
7
8
9 **4 Fengsu Hou** Ph.D. ^{1,2}, **Catherine Cerulli** J.D., Ph.D. ^{2,3}, **Marsha N. Wittink** M.D., M.B.E. ²,
10
11 **Eric D. Caine** M.D. ^{2,3}, **Peiyuan Qiu** Ph.D. ^{4, #}

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

6
7 **Institutions:**

8 1. Sun Yat-sen Global Health Institute, School of Public Health, Sun Yat-sen University
9 Address: No. 74 Zhongshan 2nd Road, Guangzhou, Guangdong, China, 510080.

10 2. Department of Psychiatry, University of Rochester Medical Center
11 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

12 3. Injury Control Research Center for Suicide Prevention, University of Rochester
13 Medical Center.
14 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

15 4. Department of Epidemiology and Health Statistics, West China School of Public
16 Health, Sichuan University
17 Address: No.17, 3 section South Renmin Road, Chengdu, Sichuan, China, 610041

18 # Corresponding author

19
20 **Corresponding author:** Peiyuan Qiu

21 Address: Department of Epidemiology and Health Statistics, West China School of Public
22 Health, Sichuan University, No.17, 3 section South Renmin Road, Chengdu, Sichuan,
23 China, 610041.

24 Telephone number: +86- 15882349145.

25 Fax: +86 028-85501604.

26

27 **Word count:** 4172 words

28

1 **Abstract**

2 **Objectives**

3 To estimate the prevalence of IPV, and explore associated factors for IPV based on a
4 rural sample of women.

5 **Setting**

6 Rural area of Guangyuan city, Sichuan, China.

7 **Participants**

8 We recruited women in total aged 16 years and above, had lived locally for at least 2
9 years and reported being married or in a relationship in the preceding year. In total,
10 there were 1501 women eligible for this study, and the response rate was 100%.

11 **Methods**

12 Participants completed demographic and social economic measures, the Revised
13 Conflict Tactics Scale, and the Duke Social Support Index. We applied chi-square test and
14 ANOVA for data description, and confirmatory factor analysis for the exploration of
15 associated factors of IPV.

16 **Results**

17 The total prevalence of IPV in the past 12 months was 29.05%, and the prevalence of
18 physical, psychological, and sexual violence were 7.66%, 26.58%, and 3.20%
19 respectively. The total prevalence was the highest among women aged 16 to 29,
20 received junior high school education, and believing family financial status was very
21 poor and same as before. Comparing with IPV victims, non-victims showed higher level

1 of social support. Confirmatory factor analysis showed that when social support
2 increased 1 unit, the risk for physical, psychological and sexual violence decreased 0.12,
3 0.35 and 0.12 units respectively; when economic status increased 1 unit, the risk for
4 physical, psychological and sexual violence indirectly decreased 0.047, 0.014 and 0.047
5 units respectively, but the total effects were not significant; when education increased 1
6 unit, the risk for psychological violence decreased 0.056.

7 **Conclusion**

8 We suggest future work is needed to create, disseminate and test potential IPV
9 interventions focusing on building social support network and increasing individuals'
10 perceived social support level to ameliorate the morbidity and mortality associated
11 with IPV.

13 **Keywords**

14 Intimate partner violence; Violence against women; Rural Chinese women; Social
15 support;

17 **Strength and limitations**

18 1. Strengths of this study are that it is the first population-based study in this rural
19 region to estimate the prevalence and associated factors of IPV among women who
20 were married or in a relationship in proceeding 12 months. In particular, we
21 considered three forms of IPV including physical, psychological and sexual violence.

- 1
2
3
4 2. We applied confirmatory factor analysis to explore associated factors of IPV, which
5
6 2 could reveal the underlying structure of observed factors and estimate the strength
7
8
9 3 of relationships between latent and observed factors.
10
11 4 3. Limitations include that we might miss residents who had day-jobs and were not at
12
13 5 home; we did not investigate women's financial contributions to their families which
14
15 6 could be an important factor associated with IPV. And at last, cultural barriers could
16
17 7 prevent women from reporting victimized experiences.
18
19 8 4. Due to the vast territory and many nationalities of China, it must be cautious to
20
21 9 generalize results.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Introduction

2 Intimate partner violence (IPV), and violence against women in particular, is a prevalent
3 and global public health concern¹. The World Health Organization (WHO) reports four
4 types of IPV including physical violence, psychological violence, sexual violence, and
5 controlling behaviors; and victims may experience different types of violence at a same
6 time ². A large WHO multi-country study reported lifetime rates of physical and sexual
7 violence by an intimate partner ranged from 13% to 61% and 6% to 59% respectively^{1 3}.

8 Although much is known about IPV worldwide, very little is known about it in China.
9 Since 2000, Hong Kong researchers have published myriad articles focusing on child
10 abuse⁴, incest ⁵and the emergence of non-governmental entities to address violence
11 against women ⁶. While these often stigmatizing topics are addressed in the literature,
12 the research on violence against women in China remains nascent. A 2004 study
13 examined data from the 1999-2000 Chinese Health and Family Life Survey and reported
14 among adults aged 20 to 64 years, 34% of women were hit during their current
15 relationship and 19% of respondents reported male-to-female IPV⁷. Other studies were
16 conducted in urban medical settings. For example, Xiao and colleagues surveyed 600
17 women in an urban outpatient gynecology clinic and found 43% reported lifetime
18 violence (including physical violence, sexual violence, or both), placing them squarely in
19 the mid-range when compared to international communities ⁸. A recent multi-country
20 study, including the USA, India, Nigeria, South Africa and China, reported that the IPV
21 rate in the past year was 10.2% among ever-partnered women in Shanghai ⁹. Pan Teng

1 and colleagues recruited 1,368 women in Guangzhou and reported the prevalence of IPV
2 over the past year was 39.2% for local women and 41.2% for migrant women ¹⁰.

3 Compared to studies in urban settings, few studies discuss IPV against women
4 living in rural China. A study based on 3,998 married women, rural Chinese women
5 reported prevalence for IPV in past-year was 65.0%, and the prevalence for physical,
6 psychological and sexual violence was 29.8%, 58.3% and 16.9% respectively¹¹.
7 However, when compared to a community-based survey that sampled patients from
8 urban, rural and industrial areas in Hunan, the lifetime reported rate of IPV was 12.4%,
9 3.3% and 14.5% respectively ¹². It is unclear whether such discrepancies accurately
10 reflect true differences or variations in sampling, measures, reporting, and other social
11 factors that can impede valid epidemiological assessments.

12 In addition to variable prevalence rates, there is a dearth of information about
13 common IPV risk factors in rural Chinese settings. Though little is known about the risk
14 factors specifically associated with IPV in rural China, there have been many studies
15 across cultures, not specific to female victimization, that provide some clues. Commonly
16 reported IPV risk factors include young age, low education, low socio-economic
17 status/income, substance use, male dominance in the family, infidelity, high proportion
18 of neighborhood poverty, acceptance of violence and divorce regulations by the
19 government^{23 13-16}. The purpose of this paper is to explore these risk factors among a
20 sample of rural Chinese women living in Hunan province. We use the social ecological
21 model to contextualize commonly studied risk factors within a framework (see Figure 1)

1 and to generate hypotheses regarding IPV risk factors ¹⁷.

2 *Insert Figure 1 here.*

3 In light of China's extraordinary economic transformation and the major social
4 changes that are sweeping across rural regions of the country, we see a need for further
5 studies of IPV. Family structures and roles are changing rapidly, with unmatched levels of
6 internal workforce migration from west to east and inland rural to urban coastal regions.
7 This "floating population" predominately includes men ¹⁸; consequently, the term
8 "left-behind" implies too much homogeneity among the women, elders, and children
9 who do not migrate for work, and it denotes populations that have emerged during the
10 past two decades. Among the "left-behind" population, most of them are rural women
11 who devote themselves to family responsibilities. However, we have little understanding
12 about how these dramatic changes have affected these women of common social and
13 interpersonal problems such as IPV.

14 Our study focuses on data collected from the rural areas of Guangyuan City in Sichuan
15 province to estimate the prevalence of IPV and to explore associated factors. Of noted, a
16 "city" in China is most often comprised of multiple regions including counties, towns,
17 villages, and rural areas; and Guangyuan region, especially its rural towns and villages in
18 mountainous areas, has been stricken by the Wenchuan earthquake in 2008. We
19 hypothesize lower prevalence rates in Guangyuan region than in rural areas of northeast
20 China, where IPV against women is acceptable in the regional culture¹¹. We also
21 hypothesize social ecological factors from all levels (personal and interpersonal level)

1 will have a direct influence on IPV.

2

3 **Method**

4 This study is part of a larger epidemiological study conducted in rural areas of
5 Guangyuan City in Sichuan province in July 2012. The larger study assessed the
6 prevalence of distress and diagnosed psychopathology among rural women, and
7 explored how women understand their conditions^{19 20}. Guangyuan City is located in the
8 north of Sichuan, a southwestern province in China, with approximately 820,000 people
9 in urban areas and 1.66 million in rural areas²¹. This regional area is economically
10 underdeveloped with the per capital net income of 4035.5 Yuan (equivalent to 585.3
11 USD) for rural household in 2010, one of the lowest in the province ²².

12 **Participants**

13 The sampling strategy is discussed in greater detail elsewhere²⁰. We recruited a
14 socio-economically diverse sample and used multi-stage sampling to randomly select
15 towns and villages for this study. We included all women, aged 16 years and older (16
16 years is the age of consent in China), who had lived locally for at least 2 years and
17 reported being married or in a relationship in the preceding year. Local hospitals
18 provided a list of eligible women based on the Chinese household registrations system
19 (the *hukou*), which excluded women if: 1) they had diagnosed mental or cognitive
20 problems, such as schizophrenia, autism, dementia and mental retardation, which would

1
2
3
4 1 impede their abilities to comprehend and answer questions; 2) they were unable to
5
6 2 communicate due to being deaf or mute.

7
8
9 3 Considering rural Chinese women usually have low education or could be illiterate,
10
11 4 we provided verbal informed consent without an information sheet. Participants joined
12
13 5 the study only when they provided oral consent after being informed of the study
14
15 6 purpose during a private, face-to-face interview in their homes. We provided
16
17 7 participants toiletry items (such as toothpaste and soap) worth 5 Yuan (about 0.8 USD)
18
19 8 to compensate them for their time. Additionally, interviewers provided helpful IPV
20
21 9 resources with participants who endorsed IPV.

22 23 24 25 26 10 Procedure

27
28
29 11 We conducted the field survey in July 2012. Local government and Guangyuan Mental
30
31 12 Health Center staff assisted us with recruitment. They coordinated with village leaders
32
33 13 and village doctors, and held public information sessions about this study before the
34
35 14 survey began. During the field survey, village leaders, doctors and reputable seniors led
36
37 15 interviewers door-to-door to conduct the interviews. As some villages have low
38
39 16 population density, local residents helped interviewers by transporting them
40
41 17 door-to-door on motorcycles. When an eligible participant was not at home, or
42
43 18 unavailable, the interviewer would return twice more. Interviewers conducted the
44
45 19 surveys on their personal computers during the face-to-face interviews. Interviewers
46
47 20 deleted the data their computers after putting the data on the research leader's flash
48
49 21 drive.
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Measures

2 *Demographic information questionnaire*

3 We designed the demographic questionnaire to collect socio-demographic information
4 from participants. Items included age and education.

5 *Family economic status*

6 The family economic status questionnaire asks about family annual income, family
7 property information, perceived family economic status, family economic status
8 compared with others in the village and compared with previous years.

9 Information provided about family annual income depended on participants' recall,
10 and was usually an estimation; therefore, we also collected information on property to
11 evaluate participants' economic status. We asked whether a participant's family owned
12 a TV, personal computer, modern kitchen range, mobile phone, DVD/VCD, refrigerator,
13 sofa, modern living furniture (Western style composite furniture), two wheeled
14 motorized vehicles, three wheeled or above motorized vehicles, air conditioner, washing
15 machine, and bank account with over 10,000 Yuan (about 1,571 USD). We assigned a
16 one for each of the items the participant's family owned, and a zero for those they did
17 not; the score ranged from 0 to 13.

18 *Social support*

19 We applied the 23-item Duke Social Support Index (DSSI) to evaluate participants' social
20 support²³. The Chinese version has already been used in research, and studies in
21 Chinese rural samples reported internal consistency over 0.79^{24 25}. DSSI assesses social

1 interaction, perceived support and instrumental social support. Every answer has an
2 assigned score, and these are added up to determine the total score (possible total
3 scores ranged from 11 to 45); higher scores indicate higher social support levels. Our
4 Cronbach's α was 0.835.

5 *IPV experience*

6 We applied the Short Form of the Revised Conflict Tactics Scale (CTS2S) to investigate
7 participants' IPV experiences, which measures negotiation, psychological aggression,
8 physical assault, injury, and sexual coercion, has acceptable validity and sensitivity²⁶. In
9 our previous study, the CTS2S showed good internal reliability and structural validity in
10 rural China²⁷.

11 The CTS2S contains statements about participants' experiences during the past 12
12 months and examines the frequencies of those events. We administered six victimization
13 questions from the survey, across three IPV domains (physical violence, psychological
14 violence, and sexual violence). For this study, participants were considered positive for
15 IPV if they endorsed any of the six questions. Our Cronbach's α was 0.845.

16 *Quality control*

17 The larger epidemiological study described interview training and quality control
18 measures in detail²⁰. Briefly, we recruited interviewers from West China School of
19 Public Health of Sichuan University; faculty members from Sichuan University and
20 University of Rochester Medical Center conducted training sessions related to methods,
21 interviewing skills, qualitative methods, and safety regarding IPV identification. We

1
2
3
4 1 deployed three research teams, each had eight interviewers and was led by experienced
5
6 2 senior researchers. Questions were routinely checked for missing items post-interview
7
8
9 3 to reduce missing data issues.
10

11 4 12 13 14 5 **Analysis**

15
16 6 Given that several risk factors can influence IPV both separately and collectively and are
17
18 7 likely to be highly collinear³, we used confirmatory factor analysis (CFA). CFA is a type
19
20 8 of structural equation modeling (SEM), which allows researchers to determine the
21
22 9 underlying structure of a set of observed factors based on *a priori* hypotheses, and to
23
24 10 estimate the strength of relationships between latent and observed factors^{28 29}.

25
26
27
28
29 11 We hypothesized that there were six latent factors including objective economic
30
31 12 status, subjective economic status, social support, physical violence, psychological
32
33 13 violence, and sexual violence) and two observed factors (including age and education).
34
35 14 Factors' labels and assignments are shown in Appendix 1.
36
37
38

39 15 We have a priori hypotheses based on the social ecological model according to an
40
41 16 report on violence and health³⁰, more specifically they are: 1) age, education, objective
42
43 17 economic status, subjective economic status, and social support have direct effects on
44
45 18 physical violence, psychological violence and sexual violence; 2) age and education may
46
47 19 have indirect effects on physical violence, psychological violence and sexual violence
48
49 20 through objective economic status, subjective economic status and social support; 3)
50
51 21 objective economic status may have indirect effects on physical violence, psychological
52
53
54
55
56
57
58
59
60

1 violence and sexual violence through subjective economic status and social support.

2 We ran the analysis with Mplus 7.3. and applied mean and variance-adjusted WLS
3 estimator (WLSMV) as the default estimation method in order to handle categorical
4 continuous variable in Mplus ³¹. The default model estimators in WLSMV include
5 chi-square value (χ^2), degrees of freedom (df), χ^2/df , root-mean-squared error of
6 approximation (RMSEA) and its 95% confidence interval (90%CI), comparative fit index
7 (CFI), Tucker-Lewis Index (TLI), and weighted root-mean-square residual (WRMSR).
8 Criteria to assess the model include: the lower chi-square value and degrees of freedom,
9 the better the model; CFI and TLI values should be 0.90 and over; the RMSEA value
10 should be 0.06 and below, the lower limit of 90%CI should be 0 or close to 0, and its
11 upper limit of 90%CI should be 0.08 and below; the value of χ^2/df should be under 5.0 ³¹
12 ³².

13 With respect to age, and consistent with our previous research^{19 20}, we categorized
14 participants into six age groups based on frequencies. For educational attainment, we
15 divided participants into five groups. We divided participants into six groups based on
16 family annual income and four groups based on perceived family economic status:
17 affluent, basic needs met, poor, and very poor. We divided participants into three groups
18 based on their family economic status compared with others: wealthier, same, and
19 poorer, and three groups based on their family economic status compared with previous
20 years: better, same, and worse. With respect to perceived health status, we divided
21 participants into four groups: very good, good, average and bad. Finally, we divided

1 participants into five groups based perceived activity status: “normal,” “cannot do heavy
2 farm work, but can do light farm work,” “cannot do any farm work, but can do house
3 work,” “cannot do house work but can take care of myself,” and “cannot take care of
4 myself.” We applied chi-square test and ANOVA for data description, and the statistical
5 significance level was 0.05.

7 **Results**

8 Demographic information

9 We recruited 1,501 women who were eligible for this study, and all of them completed
10 surveys with a response rate of 100%. However, we found there were 17 participants,
11 1.13% of total sample, who did not answer to either perceived family financial status,
12 family economic status compared with others or family economic status compared with
13 previous years. Since the 17 participants provided all other demographic information,
14 social support, and IPV experiences, we only exclude them in confirmatory factor
15 analysis.

16 Participants' age ranged from 16 to 87 years old, with a mean (SD) of 46.44 (13.11)
17 years. Overall, participants were not well educated: 33.11% had never been educated,
18 41.51% had received primary school education, and only 7.99% had received high
19 school education and above. Most participants' annual family income status was under
20 40,000 Yuan: 17.65% were under 9,999 Yuan; 24.72% were between 10,000 and 19,999
21 Yuan; 24.25% were between 20,000 and 29,999 Yuan; and 14.19% were between
22 30,000 and 39,999 Yuan. Meanwhile, 52.50% felt their family economic status was basic

1 enough, and 31.25% felt it was poor. However, if participants compared their family
2 economic status to others in the village, 57.16% felt it was the same comparing with that
3 36.38% felt it was poorer. Most participants, 74.55%, considered their family economic
4 status better than in previous years, compared with 9.13% who considered it worse.
5 With respect to the score of family properties, normality test showed it was negatively
6 skewed distribution with the coefficients of kurtosis and skewness as -0.631 and -0.311;
7 hence, we calculated its range, median and quartile, which were 0 to 13, 8 and 5
8 respectively. The mean score for DSSI was 37.33 ± 5.13 ; and the mean scores for social
9 interaction, perceived social support and instrumental social support were 7.81 ± 1.70 ,
10 18.89 ± 2.74 , and 10.62 ± 2.38 respectively. Details are shown in Table 1.

11 *Insert Table 1 here*

12 IPV experience

13 Participants reported an IPV prevalence rate of 29.05% (436/1501). With respect to
14 physical, psychological and sexual violence, shown in Table 2, the prevalence was 7.66%
15 (115/1501), 26.58% (399/1501) and 3.20% (48/1501) respectively.

16 The overall IPV prevalence was the highest for women aged 16-29 years old
17 (37.35%), followed by 31.20% for women aged 40-49 years old; it was the lowest for
18 women 70 years old and above. Prevalence rates also differed by education level. The
19 prevalence was the highest (35.33%) for women with junior high school education,
20 followed by 31.82% for women with college education and above, and it was the lowest
21 (24.55%) for women with no education. IPV prevalence rates varied by perceived

1 economic status in this sample. Prevalence was the highest (40.43%) for women whose
2 perceived family economic status was very poor, followed by 32.20% for women whose
3 perceived family economic status was poor, and it was the lowest (26.14%) for women
4 whose perceived family economic status was basic enough; prevalence was also highest
5 (35.53%) for women whose perceived family economic status was the same as previous
6 years, followed by 31.39% for women whose perceived family economic status was
7 worse than previous years, and it was the lowest (27.61%) for women whose perceived
8 family economic status was better than previous years. The study also found IPV victims
9 had lower total social support levels; the total score of DSSI was 36.28 ± 5.86 for victims,
10 and 37.77 ± 4.74 for non-victims. Victims also had lower perceived and instrumental
11 social support. Details are shown in Table 1 and Table 2.

12 *Insert Table 2 here*

13 Confirmatory factor analysis

14 *Model testing*

15 After seven iterations, we had the best fitting model. In Model 7, the χ^2 was 129.23, the
16 df was 50, the χ^2/df was 2.58, the RMSEA was 0.032 and its 95% confident interval
17 (95%CI) was from 0.026 to 0.039, the CFI was 0.991, the TLI was 0.987 and the WRMSR
18 was 1.116. According to the modification indexes, there was no error covariance that
19 could be set as free parameters based on knowledge, hence we chose Model 7 as the
20 final model. Table 3 showed the factor loadings and coefficients of the final model.

21 *Insert Table 3 here*

1
2
3
4 1 *Direct and indirect effects of the final model*

5
6 2 The results supported the a priori hypotheses that: 1) social support had direct effects
7
8 3 on physical, psychological and sexual violence; 2) objective economic status had indirect
9
10 4 effects on physical, psychological and sexual violence through social support; 3)
11
12 5 education had indirect effect on psychological violence through social support and
13
14 6 objective economic status. Table 3 and Figure 2 show the effects.

15
16
17
18 7 As displayed in Table 4, we found that: 1) as education increased 1 unit, the risk for
19
20 8 psychological violence decreased 0.056; 2) as economic status increased 1 unit, the risk
21
22 9 for physical, psychological and sexual violence indirectly decreased 0.047, 0.014 and
23
24 10 0.047 units respectively, but the total effects were not significant; 3) as social support
25
26 11 increased 1 unit, the risk for physical, psychological and sexual violence decreased 0.12,
27
28 12 0.35 and 0.12 units respectively.

29
30
31
32
33
34 13 *Insert Table 4 and Figure 2 here.*

35
36
37 14

38
39 15 **Discussion**

40
41 16 IPV is well-recognized as an important global health challenge, but it is not yet well
42
43 17 understood in China, particularly given the wide variability in reported prevalence. The
44
45 18 variability in earlier work may be attributed to the deep cultural roots that may impede
46
47 19 accurate reporting, methods in the relatively scant research that has already been done,
48
49 20 and the geographic variation among different studies.

50
51
52
53
54 21 The prevalence rates from studies focusing on rural Chinese women experiencing IPV

1 were not consistent. A study focusing on married women under 37 years old in central
2 China reported that the total lifetime IPV prevalence was 7.3%, the prevalence of minor
3 and severe physical violence was 6.4% and 5.8%, and the prevalence of psychological and
4 sexual violence were 3% and 1% respectively^{33 34}; another study in north China,
5 reported that the prevalence of physical, psychological and sexual violence among
6 women were 12.4%, 20.6% and 11.2%³⁵. Our results were close to Gao and Tamara's
7 study in Ning Xia, the north west of China, which reported the prevalence of physical,
8 psychological and sexual violence was 4.4%, 23.9% and 1.1%³⁶.

9 There may be two possible explanations for the variability. First, studies were
10 conducted in different areas of China, where differences in local cultural context and
11 economic development may impact attitudes towards violence against women. Thus, in
12 some cases, victims may perceive their experiences of violence as normal and/or
13 private, resulting in response bias. Traditional Chinese practices, influenced by
14 Confucian doctrine, emphasize the inferior social status of women. A famous Confucian
15 doctrine states that there are three kinds of obedience for women: "*San Cong*" (obeying
16 your father before you are married, obeying your husband during marriage, and obeying
17 your sons after your husband dies), "*Si De*" (four kinds of virtues including fidelity,
18 tidiness, propriety in speech and commitment to needle work)^{34 37}. During our CTS2S
19 assessments, several women cited these Confucian credos, when asked about hitting or
20 fighting, to rationalize violent behaviors as their punishment for failing to obey their
21 husbands or partners. Hence, we can expect that the deeper traditional culture roots, the

1 more underestimated IPV prevalence will be. Second, different studies applied varied
2 research tools such as the Abuse Assessment Screen, the Conflict Tactics Scale, the
3 Revised Conflict Tactics Scale, the Composite Abuse Scale, and self-designed
4 questionnaires or items. The difference in sensitivities and specificities of these tools
5 could account for variability in rates. Despite lower prevalence rates than expected, our
6 findings here revealed important risk and protective factors for IPV among rural Chinese
7 women.

8 This study also confirmed findings from previous studies suggesting that social
9 support is an important protective factor for IPV against women. Several considerations
10 may explain the association. Several considerations may explain the association. First,
11 social support is often a source of empowerment for women ³⁸. For example, an
12 ethnographic study revealed that attending social activities could increase women's
13 influence and prestige, and, in turn, decrease the risk for IPV ³⁹. Similarly, another
14 anthropological study suggested that the more social support women have, the greater
15 their social resources, and the more they pay attention to their rights ⁴⁰. Women with
16 higher social support might decide to end the violent relationships and decrease their
17 risks ⁴⁰. Second, perpetrators' controlling behaviors usually limit victims' interactions
18 with other people, isolate victims, lower their social support level, and eventually lead to
19 an increase of IPV risk and a vicious cycle ⁴¹. Third, social support will buffer the
20 negative and traumatic experience victims have been through. The Buffer Theory
21 suggests that social support can buffer adverse life events and the negative impacts;

1 individuals with high social support levels thus could cope with adverse events well and
2 maintain physical and mental wellness ⁴². In a 2002 study, Ann Coker and colleagues
3 reported that, among American female IPV victims aged 18 to 65 years old, victims with
4 high social support had greater perceptions of their mental health, better physical
5 health, lower prevalence of depression, anxiety, suicidal ideation and post-traumatic
6 stress disorder (PTSD) ⁴³.

7 Unlike social support, the association between sociodemographic factors and IPV are
8 not consistent with previous studies. We did not find a significant relationship between
9 age and IPV, but other studies have shown that younger age is a risk factor for both male
10 perpetrators' violent behaviors and female victims' violent experiences ⁴⁴⁻⁴⁸. This study
11 found that the higher education level rural women had, the lower their risk for
12 psychological violence, which was consistent with other studies ^{13 49 50}. Though
13 education level had insignificant effects on physical and sexual violence in this study, we
14 attributed this to the fact that the prevalence of physical and sexual violence was lower
15 than psychological violence in this sample. The sample also had a low proportion of
16 women with a relatively high education level; only 120 women received high school
17 education and above.

18 This study found an indirect relationship between objective economic status and IPV.
19 This relationship remains unsettled in current literature. Some studies reported that low
20 family economic status was a critical risk factor for male to female violence ⁴⁵. Faced
21 with the stress of poverty, males are more likely to use violent behaviors as a solution to

1 release pressure ³⁸. However, a study in South Africa reported that extremely poor family
2 economic status protects women from IPV ⁵¹. Other studies have reported that,
3 compared with objective family economic status, the contribution women made to
4 family income was a more important factor—women who made little contribution or
5 were totally dependent on their partners faced increasing risk of IPV ^{34 38 50 52}.

6 We recognize several limitations. As a cross-sectional field study, interviewers were
7 only able to recruit residents at home during the survey days; thus, some who had
8 day-jobs may have been missed. As China has a vast territory and many nationalities,
9 rural women in different areas face various living environments, cultural backgrounds
10 and customs in which attitudes towards IPV may vary; hence we must be cautious to
11 generalize our results. We didn't investigate the relationship between IPV and women's
12 financial contributions to their families. And lastly, it was possible that cultural barriers,
13 those Confucian credos, prevented women from sharing their stories. Another
14 traditional Chinese expression worth mentioning is "*Jia chou bu ke wai yang*," which
15 means one should not reveal family disgrace to outsiders ⁵³⁻⁵⁶. It is possible that the
16 belief in this notion of family disgrace could lead some participants to intentionally
17 under-report male-to-female violence.

18

19 **Conclusion**

20 IPV is a common phenomenon in rural China. The overall IPV prevalence in the last 12
21 months in Guangyuan rural area was 29.05%, and the prevalence of physical,

1 psychological and sexual violence were 7.66%, 26.58% and 3.20% respectively. We
2 found that social support, education and family objective economic status were
3 important to IPV. In 2016, China implemented its first law against violence, which
4 emphasizes global responsibility to stop violence: government departments, judiciary
5 authorities, non-governmental organizations, enterprises and institutions, and the
6 citizens. We suggest future work is needed to create, disseminate and test potential IPV
7 interventions focusing on building social support network and increasing individuals'
8 perceived social support level to ameliorate the morbidity and mortality associated with
9 IPV.

11 **Funding**

12 This study was funded by NIH Grants D43 TW009101 and D43 TW009101-01S1 (E.D.
13 Caine, PI).

15 **Conflict of interest**

16 The authors declare that they have not conflict of interest.

18 **Ethical Statement**

19 The Ethics Committee of Sichuan University reviewed and approved the protocol,
20 including the verbal informed consent process (NO.2011004-1). The University of
21 Rochester Research Subjects Review Board reviewed the approval from Sichuan
22 University and approved analyses of de-identified data.

24 **Contribution**

1 FH designed the survey instruments, implemented the field survey, monitored data
2 collection, cleaned the data, developed the plan for analysis, analyzed the data, drafted
3 and revised the paper. CC designed the survey instruments, trained interviewers,
4 assisted with the analysis plan, revised the paper, and supervised FH. MNW designed the
5 survey instruments, trained interviews, and revised the paper. EDC initiated the project,
6 revised the paper, and supervised FH. PQ initiated the project, designed the survey
7 instruments, monitored data collection, cleaned the data, and revised the paper. All
8 authors had full access to all the data in the study and take responsibility for the
9 integrity of the data and the accuracy of the data analysis. All authors read and approved
10 the final manuscript.

11

12 Acknowledgments

13 We acknowledge the important input of Wang Fang, Wang Ying, Feng Fei, and Hua Junjie
14 et al. from Sichuan University, Zhang Yinghui and Ma Mei from Guangyuan Mental Health
15 Center, Chen Shulin from Zhejiang University, and Jennifer Thompson Stone from
16 University of Rochester.

18 Data sharing statement

19 The data set is available from the corresponding author at qiupeiyuan@scu.edu.cn.

21 Reference

- 22 1. Krantz G, Garcia-Moreno C. Violence against women. *Journal of Epidemiology and*
23 *Community Health* 2005;59(10):818-21. doi: 10.1136/jech.2004.022756
- 24 2. World Health Organization. Understanding and addressing violence against women.
25 2012.

- 1
2
3 1 http://apps.who.int/iris/bitstream/10665/77432/1/WHO_RHR_12.36_eng.pdf
4 2 (accessed September 18th, 2015).
5
6 3 3. World Health Organization. Violence against women: Intimate partner and sexual
7 4 violence against women 2013 [Available from:
8 5 <http://www.who.int/mediacentre/factsheets/fs239/en/> accessed January 17
9 6 2014.
10 7
11 4. Tang CS-k. Corporal punishment and physical maltreatment against children: a
12 8 community study on Chinese parents in Hong Kong. *Child Abuse & Neglect*
13 9 2006;30(8):893-907.
14 10
15 5. Wang X, Ho PSY. Violence and desire in Beijing a young Chinese woman's strategies of
16 11 resistance in father–daughter incest and dating relationships. *Violence Against*
17 12 *Women* 2007;13(12):1319-38.
18 13
19 6. Tam SY, Tang CS-k. Comparing wife abuse perceptions between Chinese police officers
20 14 and social workers. *J Fam Viol* 2005;20(1):29-38.
21 15
22 7. Parish WL, Wang T, Laumann EO, et al. Intimate partner violence in China: national
23 16 prevalence, risk factors and associated health problems. *International Family*
24 17 *Planning Perspectives* 2004:174-81.
25 18
26 8. Xu X, Zhu F, O'Campo P, et al. Prevalence of and risk factors for intimate partner
27 19 violence in China. *American journal of public health* 2005;95(1):78-85.
28 20
29 9. Decker MR, Peitzmeier S, Olumide A, et al. Prevalence and health impact of intimate
30 21 partner violence and non-partner sexual violence among female adolescents
31 22 aged 15–19 years in vulnerable urban environments: a multi-country study. *J*
32 23 *Adolescent Health* 2014;55(6, Supplement):S58-S67. doi:
33 24 <http://dx.doi.org/10.1016/j.jadohealth.2014.08.022>
34 25
35 10. Pan T, Hall BJ, Ling L. The association between social resources and depression
36 26 among female migrants affected by domestic violence. *European journal of*
37 27 *psychotraumatology* 2014;5:10.3402/ejpt.v5.26528. doi: 10.3402/ejpt.v5.26528
38 28
39 11. Guo S, Zhao F, Wu J. Study on prevalence of domestic violence and its related factors
40 29 in rural area of China. *Chinese Journal of Public Health* 2007;1:002.
41 30
42 12. Cao Y, Zhang Y, Guo G, et al. A comparison study on domestic violence by geographic
43 31 settings in Hunan. *Chinese Mental Health Journal* 2007;21(12):849-52.
44 32
45 13. Cao Y, Zhang Y, Yang S, et al. A comparartive study on the patterns of domestic
46 33 violence *Chinese Journal of Clinical Psychology* 2008;16(1):28-30.
47 34
48 14. Tsai AC. Intimate partner violence and population mental health: why poverty and
49 35 gender inequities matter. *PLoS medicine* 2013;10(5):e1001440.
50 36
51 15. Walton-Moss BJ, Manganello J, Frye V, et al. Risk factors for intimate partner violence
52 37 and associated injury among urban women. *Journal of Community Health*
53 38 2005;30(5):377-89.
54 39
55 16. Xu X, Campbell JC, Zhu F. Intimate partner violence against Chinese women-the past,
56 40 present, and future. *Trauma, Violence, & Abuse* 2001;2(4):296-315.
57 41
58 17. Krugg E, Dahlberg L, Mercy J, et al. Violence—a global public health approach. World
59 42 report on violence and health: Geneva, Switz: World Health Organization, 2002.

- 1
2
3 18. De Brauw A, Li Q, Liu C, et al. Feminization of agriculture in China? Myths
4 surrounding women's participation in farming. *The China Quarterly*
5 2008;194:327-48.
6
7 19. Hou F, Cerulli C, Wittink M, et al. Depression, social support and associated factors
8 among women living in rural China: a cross-sectional study. *BMC women's health*
9 2015;15(1):28.
10
11 20. Qiu P, Caine E, Hou F, et al. The prevalence of distress and depression among women
12 in rural Sichuan province *PLoS one* 2016;11(8):e0161097. doi: 10.1371/
13 journal.pone.0161097
14
15 21. Guangyuan Statistics Intranet. Statistical Communiqué on the 2012 Guangyuan
16 economic and social development. 2012
17
18 22. Statistic Bureau of Sichuan Province China. Sichuan Province Statistics Yearbook
19 2011
20 <http://www.sc.stats.gov.cn/tjcbw/tjnj/2011/chinese/dir/chinesemenu.htm>
21 [1](#) [Available from:
22 <http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&sub=1>
23 [lse](#) accessed 12.31.2012.
24
25 23. Koenig HG, Westlund RE, George LK, et al. Abbreviating the Duke Social Support
26 Index for use in chronically ill elderly individuals. *Psychosomatics*
27 1993;34(1):61-9. doi: 10.1016/s0033-3182(93)71928-3 [published Online First:
28 1993/01/01]
29
30 24. Zhang J, Conwell Y, Wiczorek WF, et al. Studying Chinese suicide with proxy-based
31 data: reliability and validity of the methodology and instruments in China. *The*
32 *Journal of nervous and mental disease* 2003;191(7):450.
33
34 25. Jia C, Zhang J. Psychometric characteristics of the Duke Social Support Index in a
35 young rural Chinese population. *Death studies* 2012;36(9):858-69.
36
37 26. Straus MA, Douglas EM. A short form of the Revised Conflict Tactics Scales, and
38 typologies for severity and mutuality. *Violence and Victims* 2004;19(5):507-20.
39
40 27. Zhang N, Gao Y, Wu R, et al. Validity and reliability of the short form of the revised
41 Conflict Tactics Scales in women suffering from domestic violence in rural areas.
42 *Chinese Mental Health Journal* 2014;28(5):381-84.
43
44 28. Kääriäinen M, Kanste O, Elo S, et al. Testing and verifying nursing theory by
45 confirmatory factor analysis. *Journal of Advanced Nursing* 2011;67(5):1163-72.
46
47 29. Meyers LS, Gamst G, Guarino AJ. Applied multivariate research: design and
48 interpretation: SAGE Publications, Incorporated 2006:539-68.
49
50 30. Krug EG, Mercy JA, Dahlberg LL, et al. The world report on violence and health. *The*
51 *lancet* 2002;360(9339):1083-88.
52
53 31. Muthén LK, Muthén BO. Mplus user's guide: The comprehensive modeling program
54 for applied researchers 2012.
55
56 32. Byrne BM. Structural equation modeling with Mplus: Basic concepts, applications,
57 and programming. New York: Routledge 2012.
58
59 33. Li Z, Chen Y, Wang X. A survey of domestic violence in rural China. *Chinese Journal of*
60

- 1
2
3 1 *Behavioral Medicine and Brain Science* 2003;12(2):228-30.
- 4 2 34. Tang CS-K, Lai BP-Y. A review of empirical literature on the prevalence and risk
5 3 markers of male-on-female intimate partner violence in contemporary China,
6 4 1987–2006. *Aggression and Violent Behavior* 2008;13(1):10-28.
- 7 5 35. Xue L, Pang S, Wu L, et al. Study on the influential factors of domestic violence against
8 6 married women in rural area of Tangshan. *Modern Preventive Medicine*
9 7 2007;34(17):3343-44.
- 10 8 36. Gao Y, Jacka T. Prevalence of intimate partner violence and its impact on women's
11 9 mental health in rural Western China: a study of a county in Ningxia. *Journal of*
12 10 *Peking University Health Sciences* 2012;44(3):379-86.
- 13 11 37. Taylor RL, Choy HYF. The illustrated encyclopedia of Confucianism: N-Z: The Rosen
14 12 Publishing Group 2005:496.
- 15 13 38. Jewkes R. Intimate partner violence: causes and prevention. *The Lancet*
16 14 2002;359(9315):1423-29.
- 17 15 39. Levinson D. Family violence in cross-cultural perspective. Thousand Oaks, CA, US:
18 16 Sage Publications, Inc 1989.
- 19 17 40. Mager AK. Gender and the making of a South African Bantustan: a social history of
20 18 the Ciskei, 1945-1959. 1999
- 21 19 41. Heise L, Ellsberg M, Gottemoeller M. Ending violence against women: population
22 20 reports Baltimore: Johns Hopkins University 1999.
- 23 21 42. Callaghan P, Morrissey J. Social support and health: a review. *Journal of Advanced*
24 22 *Nursing* 1993(18):203-10.
- 25 23 43. Coker AL, Smith PH, Thompson MP, et al. Social support protects against the negative
26 24 effects of partner violence on mental health. *Journal of Women's Health &*
27 25 *Gender-Based Medicine* 2002;11(5):465-76.
- 28 26 44. Harwell TS, Spence MR. Population surveillance for physical violence among adult
29 27 men and women, Montana 1998. *American Journal of Preventive Medicine*
30 28 2000;19(4):321-24.
- 31 29 45. World Health Organization. Preventing intimate partner and sexual violence against
32 30 women: taking action and generating evidence: World Health Organization 2010.
- 33 31 46. Romans S, Forte T, Cohen MM, et al. Who is most at risk for intimate partner violence?
34 32 A Canadian population-based study. *Journal of Interpersonal Violence*
35 33 2007;22(12):1495-514.
- 36 34 47. Vest JR, Catlin TK, Chen JJ, et al. Multistate analysis of factors associated with intimate
37 35 partner violence. *American Journal of Preventive Medicine* 2002;22(3):156-64.
- 38 36 48. Beydoun HA, Beydoun MA, Kaufman JS, et al. Intimate partner violence against adult
39 37 women and its association with major depressive disorder, depressive symptoms
40 38 and postpartum depression: a systematic review and meta-analysis. *Social*
41 39 *Science & Medicine* 2012;75(6):959-75. doi: 10.1016/j.socscimed.2012.04.025
- 42 40 49. Zhao F, Guo S, Wang L, et al. Investigation on the patterns and knowledge regarding
43 41 domestic violence among married women in rural areas of China. *Chinese Journal*
44 42 *of Epidemiology* 2006;27(8):664-68.

- 1 50. Chan KL. Study on child abuse and spouse battering: report on findings of household
 2 survey : a consultancy study commissioned by the Social Welfare Department of
 3 the Hong Kong SAR. Hong Kong: Department of Social Work and Social
 4 Admission, the University of Hong Kong. Hong Kong, 2005.
- 5 51. Jewkes R, Levin J, Penn-Kekana L. Risk factors for domestic violence: findings from a
 6 South African cross-sectional study. *Social Science & Medicine*
 7 2002;55(9):1603-17.
- 8 52. Guo S, Wu J, Qu C, et al. Physical and sexual abuse of women before, during, and after
 9 pregnancy. *International Journal of Gynecology & Obstetrics* 2004;84(3):281-86.
- 10 53. Gao G. An initial analysis of the effects of face and concern for other in Chinese
 11 interpersonal communication. *International Journal of Intercultural Relations*
 12 1998;22(4):467-82.
- 13 54. Pattie Yuk Yee LF. Hybridity in a guidance curriculum in Hong Kong. *International*
 14 *Journal for the Advancement of Counselling* 2006;28(4):331-42.
- 15 55. Liu L, Xudong Z, Miller JK. Use of metaphors in Chinese family therapy: a qualitative
 16 study. *Journal of Family Therapy* 2012
- 17 56. Sze YT, Hou J, Lan J, et al. Brief report: Profiling family therapy users of a therapy
 18 center in Beijing. *The American Journal of Family Therapy* 2011;39(4):299-306.

19
 20 Table 1 Demographic information of participants

Demographic Characters	n	IPV victim		Prevalence (%)	χ^2	p	
		No	Yes				
Age	16-29	204	127	77	37.75	19.873	0.001
	30-39	225	159	66	29.33		
	40-49	468	320	146	31.20		
	50-59	364	276	88	24.18		
	60-69	188	138	50	26.60		
Education	70-	52	45	7	13.46	11.378	0.023
	Never educated	497	375	122	24.55		
	Primary school	623	442	181	29.10		
	Junior high school	283	183	100	35.33		
	High school	98	50	26	26.53		
Family annual income	College and above	22	15	7	31.82	3.237	0.663
	0-9999 Yuan	265	194	71	26.79		
	10000-19999 Yuan	371	263	108	29.11		
	20000-29999 Yuan	346	248	98	28.32		
	30000-39999 Yuan	213	152	61	28.64		
Perceived family	40000-49999 Yuan	116	83	33	28.45	9.091	0.028
	≥50000 Yuan	190	125	65	34.21		
	Affluent	182	124	58	31.87		
	Basic enough	788	582	206	26.14		

financial status*	Difficult	469	318	151	32.20		
	Very difficult	47	28	19	40.43		
Comparing with others*	Wealthier	81	62	19	23.46	1.363	0.506
	Same	858	606	252	29.37		
	Poorer	546	384	162	29.67		
Comparing with previous years*	Better	1119	810	309	27.61	6.095	0.047
	Same	228	147	81	35.53		
	Worse	137	94	43	31.39		

	Mean	IPV victim		F	p
		No	Yes		
Social support					
Total score	37.33±5.13	37.77±4.74	36.28±5.86	26.289	0.000
Social interaction	7.81±1.70	7.82±1.70	7.81±1.68	0.002	0.962
Perceived social support	18.89±2.74	19.15±2.58	18.24±3.00	35.154	0.000
Instrumental social support	10.62±2.38	10.78±2.18	10.23±2.77	16.778	0.000
Family properties	7.54±2.97	7.50±2.93	7.63±3.05	0.604	0.437

1 Note: "*" indicates missing data.

2 "#" indicates using Fisher exact test.

3

1 Table 2 IPV experiences of participants

IPV experiences	No (%)	Yes (%)
Ever experienced in the past 12 months	1065 (70.95%)	436 (29.05%)
Physical violence	1386 (92.34%)	115 (7.66%)
Psychological violence	1102 (73.42%)	399 (26.58%)
Sexual violence	1453 (96.80%)	48 (3.20%)

2

For peer review only

1 Table 3 Factor loadings and path coefficients of the final model

Model	Dependent variables	Independent variables	Statistics results		
			Standardized coefficient	<i>p</i>	
Measurement model	OE	A3	0.61	0.00	
		A4	0.79	0.00	
		C1	0.84	0.00	
	PHV	C2	0.98	0.00	
		C3	0.45	0.00	
		C4	0.71	0.00	
	PSV	C5	0.76	0.00	
		C6	0.65	0.00	
		D1	0.55	0.00	
Structure model	SS	D2	0.42	0.00	
		D3	0.40	0.00	
		PHV	SS	-0.12	0.005
	PHV	OE	0.060	0.14	
		PSV	SS	-0.35	0.00
		SV	SS	-0.12	0.021
	SV	OE	0.062	0.11	
		OE	A1	-0.18	0.00
		OE	A2	0.30	0.00
SS		A1	-0.068	0.18	
SS		A2	0.13	0.012	
OE		OE	0.41	0.00	
Factor related	PSV	PHV	0.91	0.00	
		SV	PHV	0.30	0.00
	SV	PSV	0.46	0.00	
	C5	C1	0.31	0.00	
	D3	D2	0.41	0.00	

2 Note: OE=Objective economic status; SS=Social support; PHV=Physical violence; PSV=

3 Psychological violence; SV=Sexual violence; A1=Age; A2=Education; B1=Family annual income;

4 B2=Family properties; C1= Being pushed, shoved or slapped; C2=Being punched, kicked or

5 beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4= Being threatened to destroy

6 belongings; C5= Sex against will with physical force; C6= Sex against will without physical force;

7 D1=Social interaction; D2=Perceived social support; D3=Instrumental social support.

8

1 Table 4 Direct and indirect effects of the final model

Factors	Physical violence			Psychological violence			Sexual violence		
	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect
Age	—	0.005	0.005	—	0.030	0.030	—	0.005	0.005
Education	—	-0.011	-0.011	—	-0.056*	-0.056*	—	-0.011	-0.011
Objective economic status	0.060	-0.047*	0.013	0.11	-0.14*	-0.033	0.062	-0.047*	0.015
Social support	-0.12*	—	-0.12*	-0.35*	—	-0.35*	-0.12*	—	-0.12*

2 Note: "*" indicates $p < 0.05$.


3

For peer review only

Societal level	Community level	Interpersonal level	Personal level
Divorce regulations by government	Acceptance of traditional gender roles	Education disparity	Young age
Lack of legislation on IPV within marriage	High proportion of poverty	Male dominance in the family	Low socio-economic status/ income
Protective marriage law	High proportion of unemployment	Economic stress	Low education
Traditional gender norms and social norms supportive of violence	High proportion of female literacy	Low female contribution to household income	Mental disorder
	Low proportion of women with high level of autonomy	Number of children	Substance use
	Social isolation of nuclear families	Infidelity	Acceptance of violence and exposure to prior abuse
		Sexual jealousy	Child maltreatment
		Marital dissatisfaction	

2 Figure 1 The socio-ecological model of risk factors of IPV

1



2

3 Figure 2 The final confirmatory factor analysis model

4 Note: OE=Objective economic status; SS=Social support; SE= Subjective economic status;
 5 PHV=Physical violence; PSV= Psychological violence; SV=Sexual violence; A1=Age; A2=Education;
 6 A3=Family annual income; A3=Family properties; C1= Being pushed, shoved or slapped;
 7 C2=Being punched, kicked or beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4=
 8 Being threatened to destroy belongings; C5= Sex against will with physical force; C6= Sex against
 9 will without physical force; D1=Social interaction; D2=Perceived social support; D3=Instrumental
 10 social support.

11
12
13
14

Appendix 1 Latent and observation factors

Latent factors	Observed factors
	Age, A1
	Education, A2
Objective economic status, OE	Family annual income, A3
	Family properties, A4
Subjective economic status, SE	Perceived family financial status, B1
	Comparing with others, B2
	Comparing with previous years, B3
Physical violence, PHV	Being pushed, shoved or slapped, C1
	Being punched, kicked or beat-me-up, C2
Psychological violence, PSV	Being Insulted, swore, shouted, yelled at, C3
	Being threatened to destroy belongings, C4
Sexual violence, SV	Sex against will with physical force, C5
	Sex against will without physical force, C6
Social support, SS	Social interaction, D1
	Perceived social support, D2
	Instrumental social support, D3

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page number
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2,3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5
Objectives	3	State specific objectives, including any prespecified hypotheses	7,8
Methods			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	8,9
		<i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	
Variables	7	(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	-
		Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	10-12
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	10-12
Bias	9	Describe any efforts to address potential sources of bias	21
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	13,14
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	12-14
		(b) Describe any methods used to examine subgroups and interactions	No applicable
		(c) Explain how missing data were addressed	14
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	8
		(e) Describe any sensitivity analyses	No applicable

Continued on next page

Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	14
		(b) Give reasons for non-participation at each stage	21
		(c) Consider use of a flow diagram	-
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	14, 15
		(b) Indicate number of participants with missing data for each variable of interest	14
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	-
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	-
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	-
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	15,16
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	No applicable
		(b) Report category boundaries when continuous variables were categorized	14, 15
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	No applicable
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	No applicable
Discussion			
Key results	18	Summarise key results with reference to study objectives	17-21
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	21
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	17-21
Generalisability	21	Discuss the generalisability (external validity) of the study results	21, 22
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	22

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Using Confirmatory Factor Analysis to Explore Associated Factors of Intimate Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019465.R1
Article Type:	Research
Date Submitted by the Author:	26-Oct-2017
Complete List of Authors:	Hou, Fengsu; Sun Yat-sen University School of Public Health, Sun Yat-sen Global Health Institute; University of Rochester Medical Center, Department of Psychiatry Cerulli, Catherine; University of Rochester Medical Center, Injury Control Research Center for Suicide Prevention; University of Rochester Medical Center, Department of Psychiatry Wittink, Marsha; University of Rochester Medical Center, Department of Psychiatry Caine, Eric; University of Rochester Medical Center, Injury Control Research Center for Suicide Prevention ; University of Rochester Medical Center, Department of Psychiatry Qiu, Peiyuan; Sichuan University West China School of Public Health, West China School of Public Health; University of Rochester Medical Center, Department of Psychiatry
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Sociology, Legal and forensic medicine
Keywords:	Intimate partner violence, Violence against women, Rural Chinese women, Social support

SCHOLARONE™
Manuscripts

1
2
3 1 **Using Confirmatory Factor Analysis to Explore Associated Factors of Intimate**
4 **Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study**

5
6
7 3
8
9 4 **Fengsu Hou** Ph.D. ^{1,2}, **Catherine Cerulli** J.D., Ph.D. ^{2,3}, **Marsha N. Wittink** M.D., M.B.E. ²,
10
11 **Eric D. Caine** M.D. ^{2,3}, **Peiyuan Qiu** Ph.D. ^{4, #}
12

13 6

14
15 7 **Institutions:**

16
17 8 1. Sun Yat-sen Global Health Institute, School of Public Health, Sun Yat-sen University
18
19 9 Address: No. 74 Zhongshan 2nd Road, Guangzhou, Guangdong, China, 510080.

20
21 10 2. Department of Psychiatry, University of Rochester Medical Center

22
23 11 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

24
25 12 3. Injury Control Research Center for Suicide Prevention, University of Rochester
26
27 13 Medical Center.

28
29 14 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

30
31 15 4. Department of Epidemiology and Health Statistics, West China School of Public
32
33 16 Health, Sichuan University

34
35 17 Address: No.17, 3 section South Renmin Road, Chengdu, Sichuan, China, 610041

36
37 18 # Corresponding author

38
39 19

40
41 20 **Corresponding author:** Peiyuan Qiu

42
43 21 Address: Department of Epidemiology and Health Statistics, West China School of Public
44
45 22 Health, Sichuan University, No.17, 3 section South Renmin Road, Chengdu, Sichuan,
46
47 23 China, 610041.

48
49 24 Telephone number: +86- 15882349145.

50
51 25 Fax: +86 028-85501604.

52
53 26

54
55 27 **Word count:** 4495 words

56
57 28

1 **Abstract**

2 **Objectives**

3 To estimate the prevalence of intimate partner violence (IPV) against women and to
4 explore associated factors based on a sample of rural Chinese women.

5 **Design**

6 Cross-sectional study

7 **Setting**

8 Rural areas of Guangyuan city, Sichuan, China.

9 **Participants**

10 We recruited women in total aged 16 years and above, had lived locally for at least 2
11 years and reported being married or in a relationship in the past 12 months. In total,
12 there were 1501 women eligible for this study, and the response rate was 100%.

13 **Methods**

14 Participants completed demographic and social economic measures, the Short Form of
15 the Revised Conflict Tactics Scale, and the Duke Social Support Index. We applied
16 chi-square test, ANOVA and confirmatory factor analysis for analysis.

17 **Results**

18 The overall prevalence of IPV in the past 12 months was 29.05%, and the prevalence of
19 physical, psychological, and sexual violence was 7.66%, 26.58%, and 3.20%
20 respectively. The overall prevalence was the highest among women aged 16 to 29,
21 received junior high school education, and believing family financial status was very

1 poor and same as before. Non-victims showed higher level of social support.
2 Confirmatory factor analysis showed that the total effects of social support on physical,
3 psychological and sexual violence was -0.12, -0.35 and -0.12 respectively; the indirect
4 effects of objective economic status on physical, psychological and sexual violence was
5 -0.047, -0.014 and -0.047 respectively, but the total effects were not significant; the
6 indirect effect of education on psychological violence was -0.056.

7 **Conclusion**

8 The prevalence of IPV in Guangyuan is close to that in northwest of China. Social
9 support is an important protective factor. Future work is needed to create, disseminate
10 and test potential IPV interventions focusing on building social support network cross
11 societal sectors and increasing individuals' perceived social support level to ameliorate
12 the morbidity and mortality associated with IPV.

14 **Keywords**

15 Intimate partner violence; Violence against women; Rural Chinese women; Social
16 support;

18 **Strength and limitations**

19 1. Strengths of this study are that it is the first population-based study in this rural
20 region to estimate the prevalence and associated factors of IPV among women who
21 were married or in a relationship in the past 12 months. In particular, we considered

- 1 three forms of IPV including physical, psychological and sexual violence.
- 2 2. We applied confirmatory factor analysis to explore associated factors of IPV, which
- 3 could reveal the underlying structure of observed factors and estimate the strength
- 4 of relationships between latent and observed factors.
- 5 3. Limitations include that we might miss residents who had day-jobs and were not at
- 6 home; we did not investigate factors including child maltreatment, marital
- 7 dissatisfaction and education disparity between couples, and women's financial
- 8 contributions to their families which could be important factors associated with IPV.
- 9 And at last, cultural barriers could prevent women from reporting victimized
- 10 experiences.
- 11 4. Due to the vast territory and many nationalities of China, it must be cautious to
- 12 generalize results.

1 Introduction

2 Intimate partner violence (IPV), and violence against women in particular, is a prevalent
3 and global public health concern¹. The World Health Organization (WHO) reports four
4 types of IPV including physical violence, psychological violence, sexual violence, and
5 controlling behaviors; and victims may experience different types of violence at a same
6 time ². A large WHO multi-country study reported lifetime rates of physical and sexual
7 violence by an intimate partner ranged from 13% to 61% and 6% to 59% respectively^{1 3}.

8 Although much is known about IPV worldwide, very little is known about it in China.
9 Since 2000, Hong Kong researchers have published myriad articles focusing on child
10 abuse⁴, incest ⁵and the emergence of non-governmental entities to address violence
11 against women ⁶. While these often stigmatizing topics are addressed in the literature,
12 the research on violence against women in China remains nascent. A 2004 study
13 examined data from the 1999-2000 Chinese Health and Family Life Survey and reported
14 among adults aged 20 to 64 years, 34% of women were hit during their current
15 relationship and 19% of respondents reported male-to-female IPV⁷. Other studies were
16 conducted in urban medical settings. For example, Xiao and colleagues surveyed 600
17 women in an urban outpatient gynecology clinic and found 43% reported lifetime
18 violence (including physical violence, sexual violence, or both), placing them squarely in
19 the mid-range when compared to international communities ⁸. A recent multi-country
20 study, including the USA, India, Nigeria, South Africa and China, reported that the IPV
21 rate in the past year was 10.2% among ever-partnered women in Shanghai ⁹. Pan Teng

1 and colleagues recruited 1,368 women in Guangzhou and reported the prevalence of IPV
2 over the past year was 39.2% for local women and 41.2% for migrant women ¹⁰.

3 Compared to studies in urban settings, few studies discuss IPV against women
4 living in rural China. A study based on 3,998 married women, rural Chinese women
5 reported prevalence for IPV in past-year was 65.0%, and the prevalence for physical,
6 psychological and sexual violence was 29.8%, 58.3% and 16.9% respectively¹¹.
7 However, when compared to a community-based survey that sampled patients from
8 urban, rural and industrial areas in Hunan, the lifetime reported rate of IPV was 12.4%,
9 3.3% and 14.5% respectively ¹². It is unclear whether such discrepancies accurately
10 reflect true differences or variations in sampling, measures, reporting, and other social
11 factors that can impede valid epidemiological assessments.

12 In addition to variable prevalence rates, there is a dearth of information about
13 common IPV risk factors in rural Chinese settings. Though little is known about the risk
14 factors specifically associated with IPV in rural China, there have been many studies
15 across cultures, not specific to female victimization, that provide some clues. Commonly
16 reported IPV risk factors include young age, low education, low socio-economic
17 status/income, substance use, male dominance in the family, infidelity, high proportion
18 of neighborhood poverty, acceptance of violence and divorce regulations by the
19 government^{23 13-16}. The purpose of this paper is to explore these risk factors among a
20 sample of rural Chinese women living in Hunan province. We use the social ecological
21 model to contextualize commonly studied risk factors within a framework (see Figure 1)

1 and to generate hypotheses regarding IPV risk factors ¹⁷.

2 *Insert Figure 1 here.*

3 In light of China's extraordinary economic transformation and the major social
4 changes that are sweeping across rural regions of the country, we see a need for further
5 studies of IPV. Family structures and roles are changing rapidly, with unmatched levels of
6 internal workforce migration from west to east and inland rural to urban coastal regions.
7 This "floating population" predominately includes men ¹⁸; consequently, the term
8 "left-behind" implies too much homogeneity among the women, elders, and children
9 who do not migrate for work, and it denotes populations that have emerged during the
10 past two decades. Among the "left-behind" population, most of them are rural women
11 who devote themselves to family responsibilities. However, we have little understanding
12 about how these dramatic changes have affected these women of common social and
13 interpersonal problems such as IPV.

14 Based on data collected from the rural areas of Guangyuan City in Sichuan province,
15 we aim to estimate the prevalence of IPV, and hypothesize the prevalence would be
16 lower in Guangyuan region than in rural areas of north China, where IPV against women
17 is acceptable in the regional culture¹¹. We also aim to explore associated factors for IPV,
18 and hypothesize factors of personal and interpersonal levels of social ecological model
19 will have a direct influence on IPV.

20 21 **Method**

1 This study is part of a larger epidemiological study conducted in rural areas of
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 This study is part of a larger epidemiological study conducted in rural areas of
2 Guangyuan City in Sichuan province in July 2012. The larger study assessed the
3 prevalence of distress and diagnosed psychopathology among rural women, and
4 explored how women understand their conditions^{19 20}. Guangyuan City is located in the
5 north of Sichuan, a southwestern province in China, with approximately 820,000 people
6 in urban areas and 1.66 million in rural areas²¹. Of noted, a “city” in China is most often
7 comprised of multiple regions including counties, towns, villages, and rural areas; and
8 Guangyuan region, especially its rural towns and villages in mountainous areas, has
9 been stricken by the Wenchuan earthquake in 2008. This regional area is economically
10 underdeveloped with the per capital net income of 4035.5 Yuan (equivalent to 585.3
11 USD) for rural household in 2010, one of the lowest in the province ²².

12 Participants

13 The sampling strategy is discussed in greater detail elsewhere²⁰. We recruited a
14 socio-economically diverse sample and used multi-stage sampling to randomly select
15 towns and villages for this study. We included all women, aged 16 years and older (16
16 years is the age of consent in China), who had lived locally for at least 2 years and
17 reported being married or in a relationship in the past 12 months. Local hospitals
18 provided a list of eligible women based on the Chinese household registrations system
19 (the *hukou*), which excluded women if: 1) they had diagnosed mental or cognitive
20 problems, such as schizophrenia, autism, dementia and mental retardation, which would

1
2
3
4 1 impede their abilities to comprehend and answer questions; 2) they were unable to

5
6 2 communicate due to being deaf or mute.

7
8 3 Considering rural Chinese women usually have low education or could be illiterate,

9
10 4 we provided verbal informed consent without an information sheet. Participants joined

11
12 5 the study only when they provided oral consent after being informed of the study

13
14 6 purpose during a private, face-to-face interview in their homes; and the interviewer

15
16 7 would only start the interview when a spouse, any family members, neighbors or friends

17
18 8 of the participant were not at present. We provided participants toiletry items (such as

19
20 9 toothpaste and soap) worth 5 Yuan (about 0.8 USD) to compensate them for their time.

21
22 10 Additionally, during the interview, we required interviewers to orally provide advice

23
24 11 and suggestions on coping strategies to participants who endorsed IPV, including

25
26 12 turning to family members and friends for help, searching help from local village

27
28 13 committee, women's federation, and civil affairs department, calling police for

29
30 14 immediate intervention, and seeing a doctor; we also required interviewers to explain

31
32 15 these strategies with phrases that participants could understand. However, we did not

33
34 16 deliver advice and suggestions through any physical materials, because we were

35
36 17 concerned that the materials may indicate the disclosure of IPV which would irritate

37
38 18 perpetrators.

39
40 19 Procedure

41
42 20 We conducted the field survey in July 2012. Local government and Guangyuan Mental

43
44 21 Health Center staff assisted us with recruitment. They coordinated with village leaders

1 and village doctors, and held public information sessions about this study before the
2 survey began. During the field survey, village leaders, doctors and reputable seniors led
3 interviewers door-to-door to conduct the interviews. As some villages have low
4 population density, local residents helped interviewers by transporting them
5 door-to-door on motorcycles. When an eligible participant was not at home, or
6 unavailable, the interviewer would return twice more. Interviewers conducted the
7 surveys on their personal computers during the face-to-face interviews. Interviewers
8 deleted the data their computers after putting the data on the research leader's flash
9 drive.

10 Measures

11 *Demographic information questionnaire*

12 We designed the demographic questionnaire to collect socio-demographic information
13 from participants. Items included age and education.

14 *Family economic status*

15 The family economic status questionnaire asks about family annual income, family
16 property information, perceived family economic status, family economic status
17 compared with others in the village and compared with previous years.

18 Information provided about family annual income depended on participants' recall,
19 and was usually an estimation; therefore, we also collected information on property to
20 evaluate participants' economic status. We asked whether a participant's family owned
21 a TV, personal computer, modern kitchen range, mobile phone, DVD/VCD, refrigerator,

1 sofa, modern living furniture (Western style composite furniture), two wheeled
2 motorized vehicles, three wheeled or above motorized vehicles, air conditioner, washing
3 machine, and bank account with over 10,000 Yuan (about 1,571 USD). We assigned a
4 one for each of the items the participant's family owned, and a zero for those they did
5 not; the score ranged from 0 to 13.

6 *Social support*

7 We applied the 23-item Duke Social Support Index (DSSI) to evaluate participants' social
8 support²³. The Chinese version has already been used in research, and studies in
9 Chinese rural samples reported internal consistency over 0.79^{24 25}. DSSI assesses social
10 interaction, perceived support and instrumental social support. Every answer has an
11 assigned score, and these are added up to determine the total score (possible total
12 scores ranged from 11 to 45); higher scores indicate higher social support levels. Our
13 Cronbach's α was 0.835.

14 *IPV experience*

15 We applied the Short Form of the Revised Conflict Tactics Scale (CTS2S) to investigate
16 participants' IPV experiences, which measures negotiation, psychological aggression,
17 physical assault, injury, and sexual coercion, has acceptable validity and sensitivity²⁶. In
18 our previous study, the CTS2S showed good internal reliability and structural validity in
19 rural China²⁷.

20 The CTS2S contains statements about participants' experiences during the past year
21 and examines the frequencies of those events. For example, one of the statements is "my

1 partner pushed, shoved, or slapped me.” Participants’ answers will be categorized into
2 eight categories: once in the past year, twice in the past year, 3-5 times in the past year,
3 6-10 times in the past year, 11-20 in the past year, more than 20 times in the past year,
4 not in the past year but it did happen before, and this has never happened. We
5 administered six victimization questions from the survey, across three IPV domains
6 (physical violence, psychological violence, and sexual violence). For this study, we
7 defined the phrase “in the past year” in the CTS2S as in the past 12 months before the
8 survey, and participants were considered positive for IPV if they endorsed any of the six
9 questions. Our Cronbach’s α was 0.845.

10 Quality control

11 The larger epidemiological study described interview training and quality control
12 measures in detail ²⁰. Briefly, we recruited interviewers, who could speak and
13 understand local dialect in Guangyuan, from West China School of Public Health of
14 Sichuan University; faculty members from Sichuan University and University of
15 Rochester Medical Center conducted training sessions related to methods, interviewing
16 skills, qualitative methods, and safety regarding IPV identification. Considering rural
17 women in Guangyuan might not speak or understand phrases/expression in “*putonghua*
18 (mandarin)”, we required interviewers to explain the study consent, the purpose of the
19 study and the questionnaire including statements and phrases in the CTS2S in local
20 dialect. We deployed three research teams, each had eight interviewers and was led by

1 experienced senior researchers. Questions were routinely checked for missing items
2 post-interview to reduce missing data issues.

3

4 **Analysis**

5 Given that several risk factors can influence IPV both separately and collectively and are
6 likely to be highly collinear³, we used confirmatory factor analysis (CFA). CFA is a type
7 of structural equation modeling (SEM), which allows researchers to determine the
8 underlying structure of a set of observed factors based on *a priori* hypotheses, and to
9 estimate the strength of relationships between latent and observed factors^{28 29}.

10 We hypothesized that there were six latent factors including objective economic
11 status, subjective economic status, social support, physical violence, psychological
12 violence, and sexual violence) and two observed factors (including age and education).
13 Factors' labels and assignments are shown in Supplementary Appendix 1.

14 We have a priori hypotheses based on the social ecological model according to an
15 report on violence and health³⁰, more specifically they are: 1) age, education, objective
16 economic status, subjective economic status, and social support have direct effects on
17 physical violence, psychological violence and sexual violence; 2) age and education may
18 have indirect effects on physical violence, psychological violence and sexual violence
19 through objective economic status, subjective economic status and social support; 3)
20 objective economic status may have indirect effects on physical violence, psychological
21 violence and sexual violence through subjective economic status and social support.

1 We ran the analysis with Mplus 7.3. and applied mean and variance-adjusted WLS
2 estimator (WLSMV) as the default estimation method in order to handle categorical
3 continuous variable in Mplus³¹. The default model estimators in WLSMV include
4 chi-square value (χ^2), degrees of freedom (df), χ^2 /df, root-mean-squared error of
5 approximation (RMSEA) and its 95% confidence interval (90%CI), comparative fit index
6 (CFI), Tucker-Lewis Index (TLI), and weighted root-mean-square residual (WRMSR).
7 Criteria to assess the model include: the lower chi-square value and degrees of freedom,
8 the better the model; CFI and TLI values should be 0.90 and over; the RMSEA value
9 should be 0.06 and below, the lower limit of 90%CI should be 0 or close to 0, and its
10 upper limit of 90%CI should be 0.08 and below; the value of χ^2 /df should be under 5.0³¹
11³².

12 With respect to age, and consistent with our previous research^{19 20}, we categorized
13 participants into six age groups based on frequencies. For educational attainment, we
14 divided participants into five groups. We divided participants into six groups based on
15 family annual income and four groups based on perceived family economic status:
16 affluent, basic needs met, poor, and very poor. We divided participants into three groups
17 based on their family economic status compared with others: wealthier, same, and
18 poorer, and three groups based on their family economic status compared with previous
19 years: better, same, and worse. With respect to perceived health status, we divided
20 participants into four groups: very good, good, average and bad. Finally, we divided
21 participants into five groups based perceived activity status: "normal," "cannot do heavy

1 farm work, but can do light farm work," "cannot do any farm work, but can do house
2 work," "cannot do house work but can take care of myself," and "cannot take care of
3 myself." We applied chi-square test and ANOVA for data description, and the statistical
4 significance level was 0.05.

6 **Results**

7 Demographic information

8 We recruited 1,501 women who were eligible for this study, and all of them completed
9 surveys with a response rate of 100%. However, we found there were 17 participants,
10 1.13% of total sample, who did not answer to either perceived family financial status,
11 family economic status compared with others or family economic status compared with
12 previous years. Since the 17 participants provided all other demographic information,
13 social support, and IPV experiences, we only exclude them in confirmatory factor
14 analysis.

15 Participants' age ranged from 16 to 87 years old, with a mean (SD) of 46.44 (13.11)
16 years. Overall, participants were not well educated: 33.11% had never been educated,
17 41.51% had received primary school education, and only 7.99% had received high
18 school education and above. Most participants' annual family income status was under
19 40,000 Yuan: 17.65% were under 9,999 Yuan; 24.72% were between 10,000 and 19,999
20 Yuan; 24.25% were between 20,000 and 29,999 Yuan; and 14.19% were between
21 30,000 and 39,999 Yuan. Meanwhile, 52.50% felt their family economic status was basic
22 enough, and 31.25% felt it was poor. However, if participants compared their family

1 economic status to others in the village, 57.16% felt it was the same comparing with that
2 36.38% felt it was poorer. Most participants, 74.55%, considered their family economic
3 status better than in previous years, compared with 9.13% who considered it worse.
4 With respect to the score of family properties, normality test showed it was negatively
5 skewed distribution with the coefficients of kurtosis and skewness as -0.631 and -0.311;
6 hence, we calculated its range, median and quartile, which were 0 to 13, 8 and 5
7 respectively. The mean score for DSSI was 37.33 ± 5.13 ; and the mean scores for social
8 interaction, perceived social support and instrumental social support were 7.81 ± 1.70 ,
9 18.89 ± 2.74 , and 10.62 ± 2.38 respectively. Details are shown in Table 1.1 and Table 1.2.

10 *Insert Table 1.1 and Table 1.2 here*

11 IPV experience

12 Participants reported an IPV prevalence rate of 29.05% (436/1501). With respect to
13 physical, psychological and sexual violence, shown in Table 2, the prevalence was 7.66%
14 (115/1501), 26.58% (399/1501) and 3.20% (48/1501) respectively.

15 The overall IPV prevalence was the highest for women aged 16-29 years old
16 (37.35%), followed by 31.20% for women aged 40-49 years old; it was the lowest for
17 women 70 years old and above. Prevalence rates also differed by education level. The
18 prevalence was the highest (35.33%) for women with junior high school education,
19 followed by 31.82% for women with college education and above, and it was the lowest
20 (24.55%) for women with no education. IPV prevalence rates varied by perceived
21 economic status in this sample. Prevalence was the highest (40.43%) for women whose

1 perceived family economic status was very poor, followed by 32.20% for women whose
2 perceived family economic status was poor, and it was the lowest (26.14%) for women
3 whose perceived family economic status was basic enough; prevalence was also highest
4 (35.53%) for women whose perceived family economic status was the same as previous
5 years, followed by 31.39% for women whose perceived family economic status was
6 worse than previous years, and it was the lowest (27.61%) for women whose perceived
7 family economic status was better than previous years. The study also found IPV victims
8 had lower total social support levels; the total score of DSSI was 36.28 ± 5.86 for victims,
9 and 37.77 ± 4.74 for non-victims. Victims also had lower perceived and instrumental
10 social support. Details are shown in Table 1 and Table 2.

11 *Insert Table 2 here*

12 Confirmatory factor analysis

13 *Model testing*

14 After seven iterations, we had the best fitting model. In Model 7, the χ^2 was 129.23, the
15 df was 50, the χ^2/df was 2.58, the RMSEA was 0.032 and its 95% confident interval
16 (95%CI) was from 0.026 to 0.039, the CFI was 0.991, the TLI was 0.987 and the WRMSR
17 was 1.116. According to the modification indexes, there was no error covariance that
18 could be set as free parameters based on knowledge, hence we chose Model 7 as the
19 final model. Table 3 showed the factor loadings and coefficients of the final model.

20 *Insert Table 3 here*

21 *Direct and indirect effects of the final model*

1 The results supported the a priori hypotheses that: 1) social support had direct effects
2 on physical, psychological and sexual violence; 2) objective economic status had indirect
3 effects on physical, psychological and sexual violence through social support; 3)
4 education had indirect effect on psychological violence through social support and
5 objective economic status. Table 3 and Figure 2 show the effects.

6 As displayed in Table 4, we found that: 1) as education increased 1 unit, the risk for
7 psychological violence decreased 0.056; 2) as economic status increased 1 unit, the risk
8 for physical, psychological and sexual violence indirectly decreased 0.047, 0.014 and
9 0.047 units respectively, but the total effects were not significant; 3) as social support
10 increased 1 unit, the risk for physical, psychological and sexual violence decreased 0.12,
11 0.35 and 0.12 units respectively.

12 *Insert Table 4 and Figure 2 here.*

14 **Discussion**

15 IPV is well-recognized as an important global health challenge, but it is not yet well
16 understood in China, particularly given the wide variability in reported prevalence. The
17 variability in earlier work may be attributed to the deep cultural roots that may impede
18 accurate reporting, methods in the relatively scant research that has already been done,
19 and the geographic variation among different studies.

20 The prevalence rates from studies focusing on rural Chinese women experiencing IPV
21 were not consistent. A study focusing on married women under 37 years old in central

1 China reported that the total lifetime IPV prevalence was 7.3%, the prevalence of minor
2 and severe physical violence was 6.4% and 5.8%, and the prevalence of psychological and
3 sexual violence were 3% and 1% respectively^{33 34}; another study in north China,
4 reported that the prevalence of physical, psychological and sexual violence among
5 women were 12.4%, 20.6% and 11.2%³⁵. Comparing with these studies, our results did
6 not fully support the hypothesis that the prevalence was lower in Guangyuan than that
7 in north China. The prevalence of psychological violence was higher in Guangyuan.
8 However our results were close to Gao and Tamara 's study in Ning Xia, the northwest of
9 China, which reported the prevalence of physical, psychological and sexual violence was
10 4.4%, 23.9% and 1.1%³⁶.

11 There may be two possible explanations for the variability. First, studies were
12 conducted in different areas of China, where differences in local cultural context and
13 economic development may impact attitudes towards violence against women. Thus, in
14 some cases, victims may perceive their experiences of violence as normal and/or
15 private, resulting in response bias. Traditional Chinese practices, influenced by
16 Confucian doctrine, emphasize the inferior social status of women. A famous Confucian
17 doctrine states that there are three kinds of obedience for women: "*San Cong*" (obeying
18 your father before you are married, obeying your husband during marriage, and obeying
19 your sons after your husband dies), "*Si De*" (four kinds of virtues including fidelity,
20 tidiness, propriety in speech and commitment to needle work)^{34 37}. During our CTS2S
21 assessments, several women cited these Confucian credos, when asked about hitting or

1 fighting, to rationalize violent behaviors as their punishment for failing to obey their
2 husbands or partners. Hence, we can expect that the deeper traditional culture roots, the
3 more underestimated IPV prevalence will be. Second, different studies applied varied
4 research tools such as the Abuse Assessment Screen, the Conflict Tactics Scale, the
5 Revised Conflict Tactics Scale, the Composite Abuse Scale, and self-designed
6 questionnaires or items. The difference in sensitivities and specificities of these tools
7 could account for variability in rates. Despite lower prevalence rates than expected, our
8 findings here revealed important risk and protective factors for IPV among rural Chinese
9 women.

10 This study also confirmed findings from previous studies suggesting that social
11 support is an important protective factor for IPV against women. Several considerations
12 may explain the association. First, social support is often a source of empowerment for
13 women ³⁸. For example, an ethnographic study revealed that attending social activities
14 could increase women's influence and prestige, and, in turn, decrease the risk for IPV ³⁹.
15 Similarly, another anthropological study suggested that the more social support women
16 have, the greater their social resources, and the more they pay attention to their rights
17 ⁴⁰. Women with higher social support might decide to end the violent relationships and
18 decrease their risks ⁴⁰. Second, perpetrators' controlling behaviors usually limit victims'
19 interactions with other people, isolate victims, lower their social support level, and
20 eventually lead to an increase of IPV risk and a vicious cycle ⁴¹. Third, social support will
21 buffer the negative and traumatic experience victims have been through. The Buffer

1 Theory suggests that social support can buffer adverse life events and the negative
2 impacts; individuals with high social support levels thus could cope with adverse events
3 well and maintain physical and mental wellness⁴². In a 2002 study, Ann Coker and
4 colleagues reported that, among American female IPV victims aged 18 to 65 years old,
5 victims with high social support had greater perceptions of their mental health, better
6 physical health, lower prevalence of depression, anxiety, suicidal ideation and
7 post-traumatic stress disorder (PTSD)⁴³.

8 Unlike social support, the association between sociodemographic factors and IPV are
9 not consistent with previous studies. We did not find a significant relationship between
10 age and IPV, but other studies have shown that younger age is a risk factor for both male
11 perpetrators' violent behaviors and female victims' violent experiences⁴⁴⁻⁴⁸. This study
12 found that the higher education level rural women had, the lower their risk for
13 psychological violence, which was consistent with other studies^{13 49 50}. Though
14 education level had insignificant effects on physical and sexual violence in this study, we
15 attributed this to the fact that the prevalence of physical and sexual violence was lower
16 than psychological violence in this sample. The sample also had a low proportion of
17 women with a relatively high education level; only 120 women received high school
18 education and above.

19 This study found an indirect relationship between objective economic status and IPV.
20 This relationship remains unsettled in current literature. Some studies reported that low
21 family economic status was a critical risk factor for male to female violence⁴⁵. Faced

1 with the stress of poverty, males are more likely to use violent behaviors as a solution to
2 release pressure ³⁸. However, a study in South Africa reported that extremely poor family
3 economic status protects women from IPV ⁵¹. Other studies have reported that,
4 compared with objective family economic status, the contribution women made to
5 family income was a more important factor—women who made little contribution or
6 were totally dependent on their partners faced increasing risk of IPV ^{34 38 50 52}.

7 We recognize several limitations. As a cross-sectional field study, interviewers were
8 only able to recruit residents at home during the survey days; thus, some who had
9 day-jobs may have been missed. As China has a vast territory and many nationalities,
10 rural women in different areas face various living environments, cultural backgrounds
11 and customs in which attitudes towards IPV may vary; hence we must be cautious to
12 generalize our results. We didn't investigate the relationship between IPV and other
13 important factors including childhood maltreatment, marital satisfaction and education
14 disparity between couples, and women's financial contributions to their families.
15 Therefore, we encourage future research to investigate these factors and their
16 relationships with IPV, to fully understand IPV against women in China, so as to develop
17 and implement effective interventions. And lastly, it was possible that cultural barriers,
18 those Confusion credos, prevented women from sharing their stories. Another
19 traditional Chinese expression worth mentioning is "*Jia chou bu ke wai yang*," which
20 means one should not reveal family disgrace to outsiders ⁵³⁻⁵⁶. It is possible that the
21 belief in this notion of family disgrace could lead some participants to intentionally

1 under-report male-to-female violence.

2

3 **Conclusion**

4 Findings from this study indicate that the overall IPV prevalence in Guangyuan rural
5 areas is close to that in northwest of China. And we found personal and interpersonal
6 factors, especially social support, were important to IPV. In 2016, China has
7 implemented its first law against violence, which emphasizes global responsibility of
8 different societal sectors to stop violence, including government departments, judiciary
9 authorities, non-governmental organizations, enterprises and institutions, and the
10 citizens. We suggest future work is needed to create, disseminate and test potential IPV
11 interventions focusing on building social support network cross different societal
12 sectors and increasing individuals' perceived social support level to ameliorate the
13 morbidity and mortality associated with IPV.

14 **Funding**

15 This study was funded by NIH Grants D43 TW009101 and D43 TW009101-01S1 (E.D.
16 Caine, PI).

17 **Conflict of interest**

18 The authors declare that they have not conflict of interest.

19 **Ethical Statement**

20 The Ethics Committee of Sichuan University reviewed and approved the protocol,

1 including the verbal informed consent process (NO.2011004-1). The University of
2 Rochester Research Subjects Review Board reviewed the approval from Sichuan
3 University and approved analyses of de-identified data.

4 5 **Contribution**

6 FH designed the survey instruments, implemented the field survey, monitored data
7 collection, cleaned the data, developed the plan for analysis, analyzed the data, drafted
8 and revised the paper. CC designed the survey instruments, trained interviewers,
9 assisted with the analysis plan, revised the paper, and supervised FH. MNW designed the
10 survey instruments, trained interviews, and revised the paper. EDC initiated the project,
11 revised the paper, and supervised FH. PQ initiated the project, designed the survey
12 instruments, monitored data collection, cleaned the data, and revised the paper. All
13 authors had full access to all the data in the study and take responsibility for the
14 integrity of the data and the accuracy of the data analysis. All authors read and approved
15 the final manuscript.

16 17 **Acknowledgments**

18 We acknowledge the important input of Wang Fang, Wang Ying, Feng Fei, and Hua Junjie
19 et al. from Sichuan University, Zhang Yinghui and Ma Mei from Guangyuan Mental Health
20 Center, Chen Shulin from Zhejiang University, and Jennifer Thompson Stone from
21 University of Rochester.

22 23 **Data sharing statement**

24 The data set is available from the corresponding author at qiupeiyuan@scu.edu.cn.

1

2 **Reference**

- 3 1. Krantz G, Garcia-Moreno C. Violence against women. *Journal of Epidemiology*
4 *and Community Health* 2005;59(10):818-21. doi:
5 10.1136/jech.2004.022756
- 6 2. World Health Organization. Understanding and addressing violence against
7 women: World Health Organization; 2012 [updated September 18th, 2015].
8 Available from:
9 [http://apps.who.int/iris/bitstream/10665/77432/1/WHO_RHR_12.36_e](http://apps.who.int/iris/bitstream/10665/77432/1/WHO_RHR_12.36_eng.pdf)
10 [ng.pdf](http://apps.who.int/iris/bitstream/10665/77432/1/WHO_RHR_12.36_eng.pdf).
- 11 3. World Health Organization. Violence against women: Intimate partner and
12 sexual violence against women 2013 [Available from:
13 <http://www.who.int/mediacentre/factsheets/fs239/en/> accessed
14 January 17 2014.
- 15 4. Tang CS-k. Corporal punishment and physical maltreatment against children: a
16 community study on Chinese parents in Hong Kong. *Child Abuse & Neglect*
17 2006;30(8):893-907.
- 18 5. Wang X, Ho PSY. Violence and desire in Beijing a young Chinese woman's
19 strategies of resistance in father–daughter incest and dating relationships.
20 *Violence Against Women* 2007;13(12):1319-38.
- 21 6. Tam SY, Tang CS-k. Comparing wife abuse perceptions between Chinese police
22 officers and social workers. *J Fam Viol* 2005;20(1):29-38.
- 23 7. Parish WL, Wang T, Laumann EO, et al. Intimate partner violence in China:
24 national prevalence, risk factors and associated health problems.
25 *International Family Planning Perspectives* 2004:174-81.
- 26 8. Xu X, Zhu F, O'Campo P, et al. Prevalence of and risk factors for intimate partner
27 violence in China. *American journal of public health* 2005;95(1):78-85.
- 28 9. Decker MR, Peitzmeier S, Olumide A, et al. Prevalence and health impact of
29 intimate partner violence and non-partner sexual violence among female
30 adolescents aged 15–19 years in vulnerable urban environments: a
31 multi-country study. *J Adolescent Health* 2014;55(6, Supplement):S58-S67.
32 doi: <http://dx.doi.org/10.1016/j.jadohealth.2014.08.022>
- 33 10. Pan T, Hall BJ, Ling L. The association between social resources and
34 depression among female migrants affected by domestic violence.
35 *European journal of psychotraumatology* 2014;5:10.3402/ejpt.v5.26528.
36 doi: 10.3402/ejpt.v5.26528
- 37 11. Guo S, Zhao F, Wu J. Study on prevalence of domestic violence and its related
38 factors in rural area of China. *Chinese Journal of Public Health* 2007;1:002.
- 39 12. Cao Y, Zhang Y, Guo G, et al. A comparison study on domestic violence by
40 geographic settings in Hunan. *Chinese Mental Health Journal*

- 1 2007;21(12):849-52.
- 2 13. Cao Y, Zhang Y, Yang S, et al. A comparative study on the patterns of domestic
3 violence *Chinese Journal of Clinical Psychology* 2008;16(1):28-30.
- 4 14. Tsai AC. Intimate partner violence and population mental health: why poverty
5 and gender inequities matter. *PLoS medicine* 2013;10(5):e1001440.
- 6 15. Walton-Moss BJ, Manganello J, Frye V, et al. Risk factors for intimate partner
7 violence and associated injury among urban women. *Journal of*
8 *Community Health* 2005;30(5):377-89.
- 9 16. Xu X, Campbell JC, Zhu F. Intimate partner violence against Chinese
10 women—the past, present, and future. *Trauma, Violence, & Abuse*
11 2001;2(4):296-315.
- 12 17. Krugg E, Dahlberg L, Mercy J, et al. Violence—a global public health approach.
13 World report on violence and health: Geneva, Switz: World Health
14 Organization, 2002.
- 15 18. De Brauw A, Li Q, Liu C, et al. Feminization of agriculture in China? Myths
16 surrounding women's participation in farming. *The China Quarterly*
17 2008;194:327-48.
- 18 19. Hou F, Cerulli C, Wittink M, et al. Depression, social support and associated
19 factors among women living in rural China: a cross-sectional study. *BMC*
20 *women's health* 2015;15(1):28.
- 21 20. Qiu P, Caine E, Hou F, et al. The prevalence of distress and depression among
22 women in rural Sichuan province *PloS one* 2016;11(8):e0161097. doi:
23 10.1371/journal.pone.0161097
- 24 21. Guangyuan Statistics Intranet. Statistical Communique on the 2012
25 Guangyuan economic and social development. 2012
- 26 22. Statistic Bureau of Sichuan Province China. Sichuan province statistics
27 yearbook 2011
28 [http://www.sc.stats.gov.cn/tjcbw/tjnj/2011/chinese/dir/chinesemenu.h](http://www.sc.stats.gov.cn/tjcbw/tjnj/2011/chinese/dir/chinesemenu.htm2011)
29 [tm2011](http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&sub=1,false) [Available from:
30 [http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&s](http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&sub=1,false)
31 [ub=1,false](http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&sub=1,false) accessed 12.31.2012.
- 32 23. Koenig HG, Westlund RE, George LK, et al. Abbreviating the Duke Social
33 Support Index for use in chronically ill elderly individuals. *Psychosomatics*
34 1993;34(1):61-9. doi: 10.1016/s0033-3182(93)71928-3 [published
35 Online First: 1993/01/01]
- 36 24. Zhang J, Conwell Y, Wiczorek WF, et al. Studying Chinese suicide with
37 proxy-based data: Reliability and validity of the methodology and
38 instruments in China. *The Journal of nervous and mental disease*
39 2003;191(7):450.
- 40 25. Jia C, Zhang J. Psychometric characteristics of the Duke Social Support Index
41 in a young rural Chinese population. *Death studies* 2012;36(9):858-69.
- 42 26. Straus MA, Douglas EM. A short form of the Revised Conflict Tactics Scales,

- 1 and typologies for severity and mutuality. *Violence and Victims*
2 2004;19(5):507-20.
- 3 27. Zhang N, Gao Y, Wu R, et al. Validity and reliability of the short form of the
4 revised Conflict Tactics Scales in women suffering from domestic violence
5 in rural areas. *Chinese Mental Health Journal* 2014;28(5):381-84.
- 6 28. Kääriäinen M, Kanste O, Elo S, et al. Testing and verifying nursing theory by
7 confirmatory factor analysis. *Journal of Advanced Nursing*
8 2011;67(5):1163-72.
- 9 29. Meyers LS, Gamst G, Guarino AJ. Applied multivariate research: design and
10 interpretation: SAGE Publications, Incorporated 2006:539-68.
- 11 30. Krug EG, Mercy JA, Dahlberg LL, et al. The world report on violence and
12 health. *The Lancet* 2002;360(9339):1083-88.
- 13 31. Muthén LK, Muthén BO. Mplus user's guide: the comprehensive modeling
14 program for applied researchers 2012.
- 15 32. Byrne BM. Structural equation modeling with Mplus: basic concepts,
16 applications, and programming. New York: Routledge 2012.
- 17 33. Li Z, Chen Y, Wang X. A survey of domestic violence in rural China. *Chinese*
18 *Journal of Behavioral Medicine and Brain Science* 2003;12(2):228-30.
- 19 34. Tang CS-K, Lai BP-Y. A review of empirical literature on the prevalence and
20 risk markers of male-on-female intimate partner violence in
21 contemporary China, 1987–2006. *Aggression and Violent Behavior*
22 2008;13(1):10-28.
- 23 35. Xue L, Pang S, Wu L, et al. Study on the influential factors of domestic violence
24 against married women in rural area of Tangshan. *Modern Preventive*
25 *Medicine* 2007;34(17):3343-44.
- 26 36. Gao Y, Jacka T. Prevalence of intimate partner violence and its impact on
27 women's mental health in rural Western China: a study of a county in
28 Ningxia. *Journal of Peking University Health Sciences* 2012;44(3):379-86.
- 29 37. Taylor RL, Choy HYF. The illustrated encyclopedia of Confucianism: N-Z: The
30 Rosen Publishing Group 2005:496.
- 31 38. Jewkes R. Intimate partner violence: causes and prevention. *The Lancet*
32 2002;359(9315):1423-29.
- 33 39. Levinson D. Family violence in cross-cultural perspective. Thousand Oaks, CA,
34 US: Sage Publications, Inc 1989.
- 35 40. Mager AK. Gender and the making of a South African Bantustan: a social
36 history of the Ciskei, 1945-1959. 1999
- 37 41. Heise L, Ellsberg M, Gottemoeller M. Ending violence against women:
38 population reports Baltimore: Johns Hopkins University 1999.
- 39 42. Callaghan P, Morrissey J. Social support and health: a review. *Journal of*
40 *Advanced Nursing* 1993(18):203-10.
- 41 43. Coker AL, Smith PH, Thompson MP, et al. Social support protects against the
42 negative effects of partner violence on mental health. *Journal of Women's*

- 1
2
3 1 *Health & Gender-Based Medicine* 2002;11(5):465-76.
- 4 2 44. Harwell TS, Spence MR. Population surveillance for physical violence among
5 3 adult men and women, Montana 1998. *American Journal of Preventive*
6 4 *Medicine* 2000;19(4):321-24.
- 7 5 45. World Health Organization. Preventing intimate partner and sexual violence
8 6 against women: taking action and generating evidence: World Health
9 7 Organization 2010.
- 10 8 46. Romans S, Forte T, Cohen MM, et al. Who is most at risk for intimate partner
11 9 violence? A Canadian population-based study. *Journal of Interpersonal*
12 10 *Violence* 2007;22(12):1495-514.
- 13 11 47. Vest JR, Catlin TK, Chen JJ, et al. Multistate analysis of factors associated with
14 12 intimate partner violence. *American Journal of Preventive Medicine*
15 13 2002;22(3):156-64.
- 16 14 48. Beydoun HA, Beydoun MA, Kaufman JS, et al. Intimate partner violence
17 15 against adult women and its association with major depressive disorder,
18 16 depressive symptoms and postpartum depression: a systematic review
19 17 and meta-analysis. *Social Science & Medicine* 2012;75(6):959-75. doi:
20 18 10.1016/j.socscimed.2012.04.025
- 21 19 49. Zhao F, Guo S, Wang L, et al. Investigation on the patterns and knowledge
22 20 regarding domestic violence among married women in rural areas of
23 21 China. *Chinese Journal of Epidemiology* 2006;27(8):664-68.
- 24 22 50. Chan KL. Study on child abuse and spouse battering: report on findings of
25 23 household survey : a consultancy study commissioned by the Social
26 24 Welfare Department of the Hong Kong SAR. Hong Kong: Department of
27 25 Social Work and Social Admission, the University of Hong Kong. Hong
28 26 Kong, 2005.
- 29 27 51. Jewkes R, Levin J, Penn-Kekana L. Risk factors for domestic violence: findings
30 28 from a South African cross-sectional study. *Social Science & Medicine*
31 29 2002;55(9):1603-17.
- 32 30 52. Guo S, Wu J, Qu C, et al. Physical and sexual abuse of women before, during,
33 31 and after pregnancy. *International Journal of Gynecology & Obstetrics*
34 32 2004;84(3):281-86.
- 35 33 53. Gao G. An initial analysis of the effects of face and concern for other in
36 34 Chinese interpersonal communication. *International Journal of*
37 35 *Intercultural Relations* 1998;22(4):467-82.
- 38 36 54. Pattie Yuk Yee LF. Hybridity in a guidance curriculum in Hong Kong.
39 37 *International Journal for the Advancement of Counselling*
40 38 2006;28(4):331-42.
- 41 39 55. Liu L, Xudong Z, Miller JK. Use of metaphors in Chinese family therapy: a
42 40 qualitative study. *Journal of Family Therapy* 2012
- 43 41 56. Sze YT, Hou J, Lan J, et al. Brief report: Profiling family therapy users of a
44 42 therapy center in Beijing. *The American Journal of Family Therapy*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 2011;39(4):299-306.
2
3

For peer review only

1 Table 1.1 Demographic information of participants

Demographic Characters	n	IPV victim		Prevalence (%)	χ^2	p
		No	Yes			
Age	16-29	204	127	77	19.873	0.001
	30-39	225	159	66		
	40-49	468	320	146		
	50-59	364	276	88		
	60-69	188	138	50		
	70-	52	45	7	13.46	
Education	Never educated	497	375	122	11.378	0.023
	Primary school	623	442	181		
	Junior high school	283	183	100		
	High school	98	50	26		
	College and above	22	15	7		
Family annual income	0-9999 Yuan	265	194	71	3.237	0.663
	10000-19999 Yuan	371	263	108		
	20000-29999 Yuan	346	248	98		
	30000-39999 Yuan	213	152	61		
	40000-49999 Yuan	116	83	33		
	≥50000 Yuan	190	125	65	34.21	
Perceived family financial status*	Affluent	182	124	58	9.091	0.028
	Basic enough	788	582	206		
	Difficult	469	318	151		
	Very difficult	47	28	19		
Comparing with others*	Wealthier	81	62	19	1.363	0.506
	Same	858	606	252		
	Poorer	546	384	162		
Comparing with previous years*	Better	1119	810	309	6.095	0.047
	Same	228	147	81		
	Worse	137	94	43		

2 Note: "*" indicates missing data; "#" indicates using Fisher exact test; "IPV" is the abbreviation
3 of intimate partner violence.

4

1 Table 1.2 The social support level of participants

Social support	Mean	IPV victim		F	<i>p</i>
		No	Yes		
Total score	37.33±5.13	37.77±4.74	36.28±5.86	26.289	0.000
Social interaction	7.81±1.70	7.82±1.70	7.81±1.68	0.002	0.962
Perceived social support	18.89±2.74	19.15±2.58	18.24±3.00	35.154	0.000
Instrumental social support	10.62±2.38	10.78±2.18	10.23±2.77	16.778	0.000
Family properties	7.54±2.97	7.50±2.93	7.63±3.05	0.604	0.437

2 Note: "IPV" is the abbreviation of intimate partner violence.

3

For peer review only

1 Table 2 IPV experiences of participants

IPV experiences	No (%)	Yes (%)
Ever experienced in the past 12 months	1065 (70.95%)	436 (29.05%)
Physical violence	1386 (92.34%)	115 (7.66%)
Psychological violence	1102 (73.42%)	399 (26.58%)
Sexual violence	1453 (96.80%)	48 (3.20%)

2 Note: "IPV" is the abbreviation of intimate partner violence.

3

For peer review only

1 Table 3 Factor loadings and path coefficients of the final model

Model	Dependent variables	Independent variables	Statistics results	
			Standardized coefficient	<i>p</i>
Measurement model	OE	A3	0.61	0.00
		A4	0.79	0.00
	PHV	C1	0.84	0.00
		C2	0.98	0.00
		C3	0.45	0.00
	PSV	C4	0.71	0.00
		C5	0.76	0.00
	SV	C6	0.65	0.00
		D1	0.55	0.00
		D2	0.42	0.00
Structure model	PHV	D3	0.40	0.00
		SS	-0.12	0.005
	PSV	OE	0.060	0.14
		SS	-0.35	0.00
	SV	OE	0.11	0.060
		SS	-0.12	0.021
	OE	OE	0.062	0.11
		A1	-0.18	0.00
		A2	0.30	0.00
		SS	A1	-0.068
A2			0.13	0.012
OE		0.41	0.00	
Factor related	PSV	PHV	0.91	0.00
		PHV	0.30	0.00
	PSV	0.46	0.00	
	C5	C1	0.31	0.00
	D3	D2	0.41	0.00

2 Note: OE=Objective economic status; SS=Social support; PHV=Physical violence; PSV=

3 Psychological violence; SV=Sexual violence; A1=Age; A2=Education; B1=Family annual income;

4 B2=Family properties; C1= Being pushed, shoved or slapped; C2=Being punched, kicked or

5 beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4= Being threatened to destroy

6 belongings; C5= Sex against will with physical force; C6= Sex against will without physical force;

7 D1=Social interaction; D2=Perceived social support; D3=Instrumental social support.

8

1 Table 4 Direct and indirect effects of the final model

Factors	Physical violence			Psychological violence			Sexual violence		
	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect
Age	—	0.005	0.005	—	0.030	0.030	—	0.005	0.005
Education	—	-0.011	-0.011	—	-0.056*	-0.056*	—	-0.011	-0.011
Objective economic status	0.060	-0.047*	0.013	0.11	-0.014*	-0.033	0.062	-0.047*	0.015
Social support	-0.12*	—	-0.12*	-0.35*	—	-0.35*	-0.12*	—	-0.12*

2 Note: "*" indicates $p < 0.05$.

3

For peer review only

1 Figure 1 The socio-ecological model of risk factors of IPV

2

3 Figure 2 The final confirmatory factor analysis model

4 Note: OE=Objective economic status; SS=Social support; SE= Subjective economic status;

5 PHV=Physical violence; PSV= Psychological violence; SV=Sexual violence; A1=Age; A2=Education;

6 A3=Family annual income; A3=Family properties; C1= Being pushed, shoved or slapped;

7 C2=Being punched, kicked or beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4=

8 Being threatened to destroy belongings; C5= Sex against will with physical force; C6= Sex against

9 will without physical force; D1=Social interaction; D2=Perceived social support; D3=Instrumental

10 social support

For peer review only

Societal level	Community level	Interpersonal level	Personal level
Divorce regulations by government	Acceptance of traditional gender roles	Education disparity	Young age
Lack of legislation on IPV within marriage	High proportion of poverty	Male dominance in the family	Low socio-economic status/ income
Protective marriage law	High proportion of unemployment	Economic stress	Low education
Traditional gender norms and social norms supportive of violence	High proportion of female literacy	Low female contribution to household income	Mental disorder
	Low proportion of women with high level of autonomy	Number of children	Substance use
	Social isolation of nuclear families	Infidelity	Acceptance of violence and exposure to prior abuse
		Sexual jealousy	Child maltreatment
		Marital dissatisfaction	

Figure 1 The socio-ecological model of risk factors of IPV

303x121mm (300 x 300 DPI)




Figure 2 The final confirmatory factor analysis model

499x240mm (300 x 300 DPI)

Appendix 1 Latent and observation factors

Latent factors	Observed factors
	Age, A1
	Education, A2
Objective economic status, OE	Family annual income, A3
	Family properties, A4
Subjective economic status, SE	Perceived family financial status, B1
	Comparing with others, B2
	Comparing with previous years, B3
Physical violence, PHV	Being pushed, shoved or slapped, C1
	Being punched, kicked or beat-me-up, C2
Psychological violence, PSV	Being Insulted, swore, shouted, yelled at, C3
	Being threatened to destroy belongings, C4
Sexual violence, SV	Sex against will with physical force, C5
	Sex against will without physical force, C6
Social support, SS	Social interaction, D1
	Perceived social support, D2
	Instrumental social support, D3

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page number
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2,3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5
Objectives	3	State specific objectives, including any prespecified hypotheses	7,8
Methods			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	8,9
		<i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	
Variables	7	(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	-
		Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	10-12
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	10-12
Bias	9	Describe any efforts to address potential sources of bias	21
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	13,14
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	12-14
		(b) Describe any methods used to examine subgroups and interactions	No applicable
		(c) Explain how missing data were addressed	14
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	8
		(e) Describe any sensitivity analyses	No applicable

Continued on next page

Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	14
		(b) Give reasons for non-participation at each stage	21
		(c) Consider use of a flow diagram	-
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	14, 15
		(b) Indicate number of participants with missing data for each variable of interest	14
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	-
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	-
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	-
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	15,16
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	No applicable
		(b) Report category boundaries when continuous variables were categorized	14, 15
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	No applicable
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	No applicable
Discussion			
Key results	18	Summarise key results with reference to study objectives	17-21
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	21
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	17-21
Generalisability	21	Discuss the generalisability (external validity) of the study results	21, 22
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	22

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Using Confirmatory Factor Analysis to Explore Associated Factors of Intimate Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019465.R2
Article Type:	Research
Date Submitted by the Author:	27-Nov-2017
Complete List of Authors:	Hou, Fengsu; Sun Yat-sen University School of Public Health, Sun Yat-sen Global Health Institute; University of Rochester Medical Center, Department of Psychiatry Cerulli, Catherine; University of Rochester Medical Center, Injury Control Research Center for Suicide Prevention; University of Rochester Medical Center, Department of Psychiatry Wittink, Marsha; University of Rochester Medical Center, Department of Psychiatry Caine, Eric; University of Rochester Medical Center, Injury Control Research Center for Suicide Prevention ; University of Rochester Medical Center, Department of Psychiatry Qiu, Peiyuan; Sichuan University West China School of Public Health, West China School of Public Health; University of Rochester Medical Center, Department of Psychiatry
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology, Sociology, Legal and forensic medicine
Keywords:	Intimate partner violence, Violence against women, Rural Chinese women, Social support

SCHOLARONE™
Manuscripts

1
2
3 **1 Using Confirmatory Factor Analysis to Explore Associated Factors of Intimate**
4 **2 Partner Violence in a Sample of Chinese Rural Women: A Cross-sectional Study**

5
6
7
8
9 **4 Fengsu Hou** Ph.D. ^{1,2}, **Catherine Cerulli** J.D., Ph.D. ^{2,3}, **Marsha N. Wittink** M.D., M.B.E. ²,
10
11 **Eric D. Caine** M.D. ^{2,3}, **Peiyuan Qiu** Ph.D. ^{4, #}

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

6
7 **Institutions:**

8 1. Sun Yat-sen Global Health Institute, School of Public Health, Sun Yat-sen University
9 Address: No. 74 Zhongshan 2nd Road, Guangzhou, Guangdong, China, 510080.

10 2. Department of Psychiatry, University of Rochester Medical Center
11 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

12 3. Injury Control Research Center for Suicide Prevention, University of Rochester
13 Medical Center.
14 Address: 601 Elmwood Ave, Box PSYCH, Rochester, NY, USA 14642

15 4. Department of Epidemiology and Health Statistics, West China School of Public
16 Health, Sichuan University
17 Address: No.17, 3 section South Renmin Road, Chengdu, Sichuan, China, 610041

18 # Corresponding author

19
20 **Corresponding author:** Peiyuan Qiu

21 Address: Department of Epidemiology and Health Statistics, West China School of Public
22 Health, Sichuan University, No.17, 3 section South Renmin Road, Chengdu, Sichuan,
23 China, 610041.

24 Telephone number: +86- 15882349145.

25 Fax: +86 028-85501604.

26

27 **Word count:** 4551 words

28

1 **Abstract**

2 **Objectives**

3 To estimate the prevalence of intimate partner violence (IPV) among a sample of rural
4 Chinese women and to explore associated factors.

5 **Design**

6 Cross-sectional study

7 **Setting**

8 Rural areas of Guangyuan city, Sichuan, China.

9 **Participants**

10 We recruited 1501 women, ages 16 years and older, who lived locally for at least 2 years
11 and reported being married or in a relationship during the past 12 months. They were
12 among a larger sample 1898 potential participants from our larger parent study on the
13 prevalence of depressive-distressed symptoms.

14 **Methods**

15 Participants completed demographic and social economic measures, the Short Form of
16 the Revised Conflict Tactics Scale, and the Duke Social Support Index. We applied
17 chi-square test, ANOVA and confirmatory factor analysis for analysis.

18 **Results**

19 The overall prevalence of IPV in the past 12 months was 29.05%; the prevalence of
20 physical, psychological, and sexual violence was 7.66%, 26.58%, and 3.20%,
21 respectively. The overall prevalence was highest among women ages 16 to 29 years,

1 and was more common among those without a high school diploma and who saw their
2 family's financial status as very poor or stagnant. Women who were not victims of IPV
3 had higher levels of social support.

4 Confirmatory factor analysis showed that the total effects of social support on physical,
5 psychological and sexual violence was -0.12, -0.35 and -0.12, respectively. The indirect
6 effects of objective economic status on physical, psychological and sexual violence was
7 -0.047, -0.014 and -0.047, respectively, but the total effect was not significant. The
8 indirect effect of education on psychological violence was -0.056.

9 **Conclusion**

10 IPV is common in rural Guangyuan. Our data are comparable to findings from
11 Northwest of China. Social support is an important protective factor. Future work is
12 needed to develop, test, and later disseminate potential IPV interventions, with a focus
13 on building actual and perceived supportive social networks.

14 **Keywords**

15 Intimate partner violence; Violence against women; Rural Chinese women; Social
16 support;

17 **Strength and limitations**

18
19
20 1. Strengths of this study are that it is the first population-based study in this rural
21 region to estimate the prevalence and associated factors of IPV including physical,

- 1
2
3
4 1 psychological and sexual violence among women who were married or in a
5
6 2 relationship during the past 12 months.
7
8
9 3 2. We applied confirmatory factor analysis to explore associated factors of IPV, which
10
11 4 could reveal the underlying structure of observed factors and estimate the strength
12
13 5 of relationships between latent and observed factors.
14
15
16 6 3. We might miss residents who had day-jobs and were not at home; we did not
17
18 7 investigate factors including child maltreatment, marital dissatisfaction, and
19
20 8 education disparity between couples, and women's financial contributions to their
21
22 9 families, which could be important factors associated with IPV.
23
24
25
26 10 4. Cultural barriers could prevent women from reporting victimized experiences.
27
28
29 11 5. Due to China's vast territory and diverse nationalities of China, we must be cautious
30
31 12 generalizing these results.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Introduction

2 Intimate partner violence (IPV), and violence against women in particular, is a prevalent
3 and global public health concern¹. The World Health Organization (WHO) reports four
4 types of IPV including physical violence, psychological violence, sexual violence, and
5 controlling behaviors; and victims may experience different types of violence at a same
6 time ². A large WHO multi-country study reported lifetime rates of physical and sexual
7 violence by an intimate partner ranged from 13% to 61% and 6% to 59% respectively^{1 3}.

8 Although much is known about IPV worldwide, very little is known about its
9 occurrence in China. Since 2000, Hong Kong researchers have published myriad articles
10 focusing on child abuse⁴, incest, ⁵and the emergence of non-governmental entities to
11 address violence against women ⁶. In contrast, research on violence against women in
12 Mainland China remains nascent. A 2004 study examined data from the 1999-2000
13 Chinese Health and Family Life Survey and reported among adults, ages 20 to 64 years,
14 34% of women were hit during their current relationship and 19% of respondents
15 reported male-to-female IPV⁷. Other studies were conducted in urban medical settings.
16 For example, Xiao and colleagues surveyed 600 women in an urban outpatient
17 gynecology clinic and found 43% reported lifetime violence (including physical violence,
18 sexual violence, or both), placing them squarely in the mid-range when compared to
19 international communities ⁸. A recent multi-country study, including the USA, India,
20 Nigeria, South Africa and China, reported that the IPV rate during the past year was
21 10.2% among ever-partnered women in Shanghai ⁹. Pan Teng and colleagues recruited

1 1,368 women in Guangzhou and reported the prevalence of IPV over the past year was
2 39.2% for local women and 41.2% for migrant women ¹⁰.

3 Compared to studies in urban settings, few studies discuss IPV against women
4 living in rural China. A study based on 3,998 married women, rural Chinese women
5 reported prevalence for IPV in past-year was 65.0%, and the prevalence for physical,
6 psychological and sexual violence was 29.8%, 58.3% and 16.9% respectively¹¹.
7 However, when compared to a community-based survey that sampled patients from
8 urban, rural and industrial areas in Hunan, the lifetime reported rate of IPV was 12.4%,
9 3.3% and 14.5% respectively ¹². It is unclear whether such discrepancies accurately
10 reflect true differences or variations in sampling, measures, reporting, and other social
11 factors that can impede valid epidemiological assessments.

12 In addition to variable prevalence rates, there is a dearth of information about
13 common IPV risk factors in rural Chinese settings. Though little is known about the risk
14 factors specifically associated with IPV in rural China, there have been many studies
15 across cultures, not specific to female victimization, that provide some clues. Commonly
16 reported IPV risk factors include young age, low education, low socio-economic
17 status/income, substance use, male dominance in the family, infidelity, high proportion
18 of neighborhood poverty, acceptance of violence and divorce regulations by the
19 government^{23 13-16}. The purpose of this paper is to explore these risk factors among a
20 sample of rural Chinese women living in Sichuan province. We use the social ecological
21 model to contextualize commonly studied risk factors within a framework (see Figure 1)

1 and to generate hypotheses regarding IPV risk factors ¹⁷.

2 *Insert Figure 1 here.*

3 In light of China's extraordinary economic transformation and the major social
4 changes that are sweeping across rural regions of the country, we see a need for further
5 studies of IPV. Family structures and roles are changing rapidly, with unmatched levels of
6 internal workforce migration from west to east and inland rural to urban coastal regions.
7 This "floating population" predominately includes men ¹⁸; consequently, the term
8 "left-behind" implies too much homogeneity among the women, elders, and children
9 who do not migrate for work, and it denotes populations that have emerged during the
10 past two decades. Among the "left-behind" population, most of them are rural women
11 who devote themselves to family responsibilities. However, we have little understanding
12 about how these dramatic changes have affected these women of common social and
13 interpersonal problems such as IPV.

14 Based on data collected from the rural areas of Guangyuan City in northern Sichuan
15 Province, we estimated the prevalence of IPV. We also tested whether the prevalence
16 would be lower in Guangyuan region than in rural areas of northern China, where IPV
17 against women has been described as an acceptable aspect of regional culture¹¹. We also
18 examined associated factors, hypothesizing that measures sensitive to personal and
19 interpersonal levels of the social ecological model will have a direct influence on IPV.

20

21 **Method**

1 This study is part of a larger epidemiological study conducted in rural areas of
2
3
4 1 This study is part of a larger epidemiological study conducted in rural areas of
5
6 2 Guangyuan City in Sichuan province in July 2012. The larger study assessed the
7
8 3 prevalence of distress and diagnosed psychopathology among rural women, and
9
10 4 explored how women understand their conditions^{19 20}. Guangyuan City is located in the
11
12 5 north of Sichuan, a southwestern province in China, with approximately 820,000 people
13
14 6 in urban areas and 1.66 million in rural areas²¹. Of noted, a “city” in China is most often
15
16 7 comprised of multiple regions including counties, towns, villages, and rural areas. Of
17
18 8 note, the Guangyuan region of Sichuan Province, especially its rural towns and villages
19
20 9 in mountainous areas, was stricken by the 2008 Wenchuan earthquake. This regional
21
22 10 area is economically underdeveloped with a 2010 net per capita net household income
23
24 11 of 4035.5 yuan (USD\$585.3), one of the lowest in the province and the country ²².
25
26
27
28
29
30

31 Participants

32
33
34 13 The sampling strategy is discussed in greater detail elsewhere²⁰. We recruited a
35
36 14 socio-economically diverse sample and used multi-stage sampling to randomly select
37
38 15 towns and villages for this study. We included all women, ages 16 years and older (16
39
40 16 years is the age of consent in China), who had lived locally for at least 2 years and
41
42 17 reported being married or in a relationship during the past 12 months. Local hospitals
43
44 18 provided a list of eligible women based on the Chinese household registrations system
45
46 19 (the *hukou*), which excluded women if: 1) they had diagnosed mental or cognitive
47
48 20 problems, such as schizophrenia, autism, dementia and mental retardation, which would
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 1 impede their abilities to comprehend and answer questions; 2) they were unable to
5
6 2 communicate due to being deaf or mute.

7
8 3 This study was reviewed and approved by the Ethics Committee of Sichuan
9
10 4 University and the University of Rochester Research Subjects Review Board. To
11
12 5 ascertain informed consent among these women, with many having low educational
13
14 6 levels and a substantial proportion with functional illiteracy, we conducted a verbal
15
16 7 consent with language attuned to participants needs and without a written information
17
18 8 sheet. This process did not begin until the potential participants was alone. We insisted
19
20 9 that spouse, family members, neighbors, or friends of the participant were not present
21
22 10 before beginning face-to-face interviews. We provided participants toiletry items (such
23
24 11 as toothpaste and soap) worth 5 yuan (about USD\$0.80) to compensate them for their
25
26 12 time. If the participant reported any affirmative response regarding IPV, interviewers
27
28 13 provided oral advice and suggestions on coping strategies to participants who endorsed
29
30 14 IPV, including turning to family members and friends for help, searching help from the
31
32 15 local village committee, women's federation, or civil affairs department, calling police
33
34 16 for immediate intervention, and seeing a doctor. We also trained interviewers to explain
35
36 17 these strategies with simple phrases that participants could understand. However, we
37
38 18 did not provide any physical brochures or other materials, as these materials may have
39
40 19 indicated the disclosure of IPV, which potentially could provoke perpetrators.

41
42
43
44
45
46
47
48
49
50
51
52 20 Procedure
53
54
55
56
57
58
59
60

1 We conducted the field survey in July 2012. Local government and Guangyuan Mental
2 Health Center staff assisted us with recruitment. They coordinated with village leaders
3 and village doctors, and held public information sessions about this study before the
4 survey began. During the field survey, village leaders, doctors and reputable seniors led
5 interviewers door-to-door to conduct the interviews. As some villages have low
6 population density, local residents helped interviewers by transporting them
7 door-to-door on motorcycles. When an eligible participant was not at home, or
8 unavailable, the interviewer would return twice more. Interviewers conducted the
9 surveys on their personal computers during the face-to-face interviews. Interviewers
10 deleted the data their computers after putting the data on the research leader's flash
11 drive.

12 Measures

13 *Demographic information questionnaire*

14 We designed the demographic questionnaire to collect socio-demographic information
15 from participants. Items included age and education.

16 *Family economic status*

17 The family economic status questionnaire asks about family annual income, family
18 property information, perceived family economic status, family economic status
19 compared with others in the village and compared with previous years.

20 Information provided about family annual income depended on participants' recall,
21 and was usually an estimation; therefore, we also collected information on property to

1 evaluate participants' economic status. We asked whether a participant's family owned
2 a TV, personal computer, modern kitchen range, mobile phone, DVD/VCD, refrigerator,
3 sofa, modern living furniture (Western style composite furniture), two wheeled
4 motorized vehicles, three wheeled or above motorized vehicles, air conditioner, washing
5 machine, and bank account with over 10,000 Yuan (about 1,571 USD). We assigned a
6 one for each of the items the participant's family owned, and a zero for those they did
7 not; the score ranged from 0 to 13.

8 *Social support*

9 We applied the 23-item Duke Social Support Index (DSSI) to evaluate participants' social
10 support²³. The Chinese version has already been used in research, and studies in
11 Chinese rural samples reported internal consistency over 0.79^{24 25}. DSSI assesses social
12 interaction, perceived support and instrumental social support. Every answer has an
13 assigned score, and these are added up to determine the total score (possible total
14 scores ranged from 11 to 45); higher scores indicate higher social support levels. Our
15 Cronbach's α was 0.835.

16 *IPV experience*

17 We applied the Short Form of the Revised Conflict Tactics Scale (CTS2S) to investigate
18 participants' IPV experiences, which measures negotiation, psychological aggression,
19 physical assault, injury, and sexual coercion, has acceptable validity and sensitivity²⁶. In
20 our previous study, the CTS2S showed good internal reliability and structural validity in
21 rural China²⁷.

1 The CTS2S contains statements about participants' experiences during the past year
2 and examines the frequencies of those events. For example, one of the statements is "my
3 partner pushed, shoved, or slapped me." Participants' answers were categorized into
4 eight categories: once in the past year, twice in the past year, 3-5 times in the past year,
5 6-10 times in the past year, 11-20 in the past year, more than 20 times in the past year,
6 not in the past year but it did happen before, and this has never happened. We
7 administered six victimization questions across three IPV domains (physical violence,
8 psychological violence, and sexual violence). For this study, we defined the phrase "in
9 the past year" in the CTS2S as occurring during the 12 months preceding the survey, and
10 participants were considered positive for IPV if they endorsed any of the six questions.
11 Our Cronbach's α was 0.845.

12 Quality control

13 The larger epidemiological study described interview training and quality control
14 measures in detail ²⁰. Briefly, we recruited interviewers who could speak and
15 understand the local dialect in Guangyuan from the West China School of Public Health
16 of Sichuan University. Faculty from Sichuan University and University of Rochester
17 Medical Center conducted training sessions related to methods, interviewing skills,
18 qualitative methods, and safety regarding IPV identification. Considering that many
19 women in Guangyuan might not speak or understand phrases/expression in *putonghua*
20 (mandarin)", we required interviewers to explain the study consent, the purpose of the
21 study and the questionnaire including statements and phrases in the CTS2S in local

1 dialect. We deployed three research teams, each had eight interviewers and was led by
2 experienced senior researchers. Questions were routinely checked for missing items
3 post-interview to reduce missing data issues.

4 **Analysis**

5
6 Given that several risk factors can influence IPV both separately and collectively and are
7 likely to be highly collinear³, we used confirmatory factor analysis (CFA). CFA is a type
8 of structural equation modeling (SEM), which allows researchers to determine the
9 underlying structure of a set of observed factors based on *a priori* hypotheses, and to
10 estimate the strength of relationships between latent and observed factors^{28 29}.

11 We hypothesized that there were six latent factors including objective economic
12 status, subjective economic status, social support, physical violence, psychological
13 violence, and sexual violence) and two observed factors (including age and education).
14 Factors' labels and assignments are shown in Supplementary Appendix 1.

15 We had *a priori* hypotheses based on the social ecological model according to an
16 report on violence and health³⁰, more specifically they were: 1) age, education, objective
17 economic status, subjective economic status, and social support would have direct
18 effects on physical violence, psychological violence and sexual violence; 2) age and
19 education would have indirect effects on physical violence, psychological violence and
20 sexual violence through objective economic status, subjective economic status and social
21 support; 3) objective economic status would have indirect effects on physical violence,

1 psychological violence and sexual violence through subjective economic status and
2 social support.

3 We ran the analysis with Mplus 7.3. and applied mean and variance-adjusted WLS
4 estimator (WLSMV) as the default estimation method in order to handle categorical
5 continuous variable in Mplus³¹. The default model estimators in WLSMV included
6 chi-square value (χ^2), degrees of freedom (df), χ^2 /df, root-mean-squared error of
7 approximation (RMSEA) and its 95% confidence interval (90%CI), comparative fit index
8 (CFI), Tucker-Lewis Index (TLI), and weighted root-mean-square residual (WRMSR).
9 Criteria to assess the model included: the lower chi-square value and degrees of
10 freedom, the better the model; CFI and TLI values should be 0.90 and over; the RMSEA
11 value should be 0.06 and below, the lower limit of 90%CI should be 0 or close to 0, and
12 its upper limit of 90%CI should be 0.08 and below; the value of χ^2 /df should be under
13 5.0^{31 32}.

14 With respect to age, and consistent with our previous research^{19 20}, we categorized
15 participants into six age groups based on frequencies. For educational attainment, we
16 divided participants into five groups based on the hierarchy of Chinese education
17 system. We divided participants into six groups based on family annual income and four
18 groups based on perceived family economic status: affluent, basic needs met, poor, and
19 very poor. We divided participants into three groups based on their family economic
20 status compared with others: wealthier, same, and poorer, and three groups based on
21 their family economic status compared with previous years: better, same, and worse.

1 With respect to perceived health status, we divided participants into four groups: very
2 good, good, average and bad. Finally, we divided participants into five groups based
3 perceived activity status: "normal," "cannot do heavy farm work, but can do light farm
4 work," "cannot do any farm work, but can do house work," "cannot do house work but
5 can take care of myself," and "cannot take care of myself." We applied chi-square test and
6 ANOVA for data description, and the statistical significance level was 0.05.

7 8 **Results**

9 Demographic information

10 For this aspect of our larger study, we recruited 1,501 of a potential 1898 women who
11 were eligible for this study; all who consented completed their surveys. However, we
12 found there were 17 participants (1.13% of the total) did not answer items related to
13 perceived family financial status, family economic status compared with others, or
14 family economic status compared with previous years. Since the 17 participants
15 provided all other demographic information, social support, and IPV experiences, we
16 only excluded them in confirmatory factor analysis.

17 Participants' age ranged from 16 to 87 years old, with a mean (SD) of 46.44 (13.11)
18 years. Overall, participants were not well educated: 33.11% had never been educated,
19 41.51% had received primary school education, and only 7.99% had received high
20 school education and above. Most participants' annual family income status was under
21 40,000 Yuan: 17.65% were under 9,999 Yuan; 24.72% were between 10,000 and 19,999
22 Yuan; 24.25% were between 20,000 and 29,999 Yuan; and 14.19% were between

1 30,000 and 39,999 Yuan. Meanwhile, 52.50% felt their family economic status was basic
2 enough, and 31.25% felt it was poor. However, if participants compared their family
3 economic status to others in the village, 57.16% felt it was the same comparing with that
4 36.38% felt it was poorer. Most participants, 74.55%, considered their family economic
5 status better than in previous years, compared with 9.13% who considered it worse.
6 With respect to the score of family properties, normality test showed it was negatively
7 skewed distribution with the coefficients of kurtosis and skewness as -0.631 and -0.311;
8 hence, we calculated its range, median and quartile, which were 0 to 13, 8 and 5
9 respectively. The mean score for DSSI was 37.33 ± 5.13 ; and the mean scores for social
10 interaction, perceived social support and instrumental social support were 7.81 ± 1.70 ,
11 18.89 ± 2.74 , and 10.62 ± 2.38 respectively. Details are shown in Table 1.1 and Table 1.2.

12 *Insert Table 1.1 and Table 1.2 here*

13 IPV experience

14 Participants reported an IPV prevalence rate of 29.05% (436/1501). With respect to
15 physical, psychological and sexual violence, shown in Table 2, the prevalence was 7.66%
16 (115/1501), 26.58% (399/1501) and 3.20% (48/1501) respectively.

17 The overall IPV prevalence was the highest for women aged 16-29 years old
18 (37.35%), followed by 31.20% for women aged 40-49 years old; it was the lowest for
19 women 70 years old and above. Prevalence rates also differed by education level. The
20 prevalence was the highest (35.33%) for women with junior high school education,
21 followed by 31.82% for women with college education and above, and it was the lowest

1 (24.55%) for women with no education. IPV prevalence rates varied by perceived
2 economic status in this sample. Prevalence was the highest (40.43%) for women whose
3 perceived family economic status was very poor, followed by 32.20% for women whose
4 perceived family economic status was poor, and it was the lowest (26.14%) for women
5 whose perceived family economic status was basic enough; prevalence was also highest
6 (35.53%) for women whose perceived family economic status was the same as previous
7 years, followed by 31.39% for women whose perceived family economic status was
8 worse than previous years, and it was the lowest (27.61%) for women whose perceived
9 family economic status was better than previous years. The study also found IPV victims
10 had lower total social support levels; the total score of DSSI was 36.28 ± 5.86 for victims,
11 and 37.77 ± 4.74 for non-victims. Victims also had lower perceived and instrumental
12 social support. Details are shown in Table 1 and Table 2.

13 *Insert Table 2 here*

14 Confirmatory factor analysis

15 *Model testing*

16 After seven iterations, we had the best fitting model. In Model 7, the χ^2 was 129.23, the
17 df was 50, the χ^2/df was 2.58, the RMSEA was 0.032 and its 95% confident interval
18 (95%CI) was from 0.026 to 0.039, the CFI was 0.991, the TLI was 0.987 and the WRMSR
19 was 1.116. According to the modification indexes, there was no error covariance that
20 could be set as free parameters based on knowledge, hence we chose Model 7 as the
21 final model. Table 3 showed the factor loadings and coefficients of the final model.

1
2
3
4 1 *Insert Table 3 here*

5
6 2 *Direct and indirect effects of the final model*

7
8 3 The results supported our *a priori* hypotheses that: 1) social support had direct effects
9
10 4 on physical, psychological and sexual violence; 2) objective economic status had indirect
11
12 5 effects on physical, psychological and sexual violence through social support; 3)
13
14 6 education had indirect effect on psychological violence through social support and
15
16 7 objective economic status. Table 3 and Figure 2 show the effects.

17
18
19 8 As displayed in Table 4, we found that: 1) as education increased 1 unit, the risk for
20
21 9 psychological violence decreased 0.056; 2) as economic status increased 1 unit, the risk
22
23 10 for physical, psychological and sexual violence indirectly decreased 0.047, 0.014 and
24
25 11 0.047 units respectively, but the total effects were not significant; 3) as social support
26
27 12 increased 1 unit, the risk for physical, psychological and sexual violence decreased 0.12,
28
29 13 0.35 and 0.12 units respectively.

30
31
32 14 *Insert Table 4 and Figure 2 here.*

33
34
35
36
37
38
39
40
41
42 15

43 16 **Discussion**

44 17 IPV is well-recognized as an important global health challenge, but it has not been
45
46 18 well-studied or well-understood in China, particularly given the wide variability in
47
48 19 reported prevalence. This variability may reflect: deeply ingrained cultural practices that
49
50 20 may impede accurate reporting, inadequately developed research or survey methods
51
52 21 and limited samples involved in past studies, and substantial geographic and social
53
54
55
56
57
58
59
60

1 variations among those samples that have been involved in the past.

2 The prevalence rates from studies focusing on rural Chinese women experiencing IPV
3 have not been consistent. A study focusing on married women under 37 years old in
4 central China reported that the total lifetime IPV prevalence was 7.3%, with a prevalence
5 of minor and severe physical violence as 6.4% and 5.8%, respectively, and the
6 prevalence of psychological and sexual violence were 3% and 1%, respectively^{33 34}. A
7 study from north China reported that the respective prevalence of physical,
8 psychological and sexual violence among women were 12.4%, 20.6%, and 11.2%³⁵. Our
9 results did not fully support the expect finding that the prevalence of IPV was lower in
10 Guangyuan as compared to results from northern China. The prevalence of psychological
11 violence was higher in Guangyuan. Our results were close to those from Gao and Tamara
12 's study in Ning Xia, the northwest of China, which reported the prevalence of physical,
13 psychological and sexual violence was 4.4%, 23.9% and 1.1%³⁶.

14 There may be two possible explanations for the variability. First, studies were
15 conducted in different areas of China, where differences in local cultural context and
16 economic development may impact attitudes towards violence against women. In some
17 cases, victims may perceive their experiences of violence as normative or private,
18 resulting in response bias. Traditional Chinese practices, influenced by Confucian
19 doctrine, emphasize the inferior social status of women. A famous Confucian doctrine
20 states that there are three kinds of obedience for women, "*San Cong*" (obeying your
21 father before you are married, obeying your husband during marriage, and obeying your

1
2
3
4 1 sons after your husband dies), and four kinds of virtue, “*Si De*” (fidelity, tidiness,
5
6 2 propriety in speech and commitment to needle work) ³⁴⁻³⁷. During our CTS2S
7
8 3 assessments, several women cited these Confucian credos when asked about hitting or
9
10 4 fighting, to rationalize violent behaviors as their punishment for failing to obey their
11
12 5 husbands or partners. Hence, we can expect that the deeper traditional culture roots, the
13
14 6 more underestimated IPV prevalence will be. Second, different studies applied varied
15
16 7 research tools such as the Abuse Assessment Screen, the Conflict Tactics Scale, the
17
18 8 Revised Conflict Tactics Scale, the Composite Abuse Scale, and self-designed
19
20 9 questionnaires or items. The difference in sensitivities and specificities of these tools
21
22 10 could account for variability in rates. Despite lower prevalence rates than expected, our
23
24 11 findings here revealed important risk and protective factors for IPV among rural Chinese
25
26 12 women.

27
28
29
30
31
32
33
34 13 This study also confirmed findings from previous studies suggesting that social
35
36 14 support is an important protective factor for IPV against women. Several considerations
37
38 15 may explain the association. First, social support is often a source of empowerment for
39
40 16 women ³⁸. For example, an ethnographic study revealed that attending social activities
41
42 17 could increase women’s influence and prestige, and, in turn, decrease the risk for IPV ³⁹.
43
44 18 Similarly, another anthropological study suggested that the more social support women
45
46 19 have, the greater their social resources, and the more they pay attention to their rights
47
48 20 ⁴⁰. Women with higher social support might decide to end the violent relationships and
49
50 21 decrease their risks ⁴⁰. Second, perpetrators’ controlling behaviors usually limit victims’
51
52
53
54
55
56
57
58
59
60

1 interactions with other people, isolate victims, lower their social support level, and
2 eventually lead to an increase of IPV risk and a vicious cycle ⁴¹. Third, social support will
3 buffer the negative and traumatic experience victims have been through. The Buffer
4 Theory suggests that social support can buffer adverse life events and the negative
5 impacts; individuals with high social support levels thus could cope with adverse events
6 well and maintain physical and mental wellness ⁴². In a 2002 study, Ann Coker and
7 colleagues reported that, among American female IPV victims aged 18 to 65 years old,
8 victims with high social support had greater perceptions of their mental health, better
9 physical health, lower prevalence of depression, anxiety, suicidal ideation and
10 post-traumatic stress disorder (PTSD) ⁴³.

11 Unlike social support, the association between sociodemographic factors and IPV are
12 not consistent with previous studies. We did not find a significant relationship between
13 age and IPV, but other studies have shown that younger age is a risk factor for both male
14 perpetrators' violent behaviors and female victims' violent experiences ⁴⁴⁻⁴⁸. This study
15 found that the higher education level rural women had, the lower their risk for
16 psychological violence, which was consistent with other studies ^{13 49 50}. Though
17 education level had insignificant effects on physical and sexual violence in this study, we
18 attributed this to the fact that the prevalence of physical and sexual violence was lower
19 than psychological violence in this sample. The sample also had a low proportion of
20 women with a relatively high education level; only 120 women received high school
21 education and above.

1 This study found an indirect relationship between objective economic status and IPV.
2 This relationship remains unsettled in current literature. Some studies reported that low
3 family economic status was a critical risk factor for male to female violence ⁴⁵. Faced
4 with the stress of poverty, males may be more likely to use violent behaviors as a
5 solution to release pressure ³⁸. However, a study in South Africa reported that extremely
6 poor family economic status protects women from IPV ⁵¹. Other studies have reported
7 that, compared with objective family economic status, the contribution women made to
8 family income was a more important factor—women who made little contribution or
9 were totally dependent on their partners faced increasing risk of IPV ^{34 38 50 52}.

10 We recognize several limitations. As a cross-sectional field study, interviewers were
11 only able to recruit residents at home during the survey days; thus, some who had
12 day-jobs may have been missed. As China has a vast territory and many nationalities,
13 rural women in different areas face various living environments, cultural backgrounds
14 and customs in which attitudes towards IPV may vary; hence we must be cautious to
15 generalize our results. We did not investigate the relationship between IPV and other
16 important factors, including childhood maltreatment, marital satisfaction, education
17 disparity between couples, and women's financial contributions to their families.

18 We encourage future investigators to investigate these factors and their relationships
19 with IPV, to fully understand IPV against women in China to develop and implement
20 effective interventions. It will be important to examine cultural barriers, such as
21 Confucian precepts, to explore how these affect normative assumptions and openness to

1 speaking with others about their experiences. Another traditional Chinese expression is
2
3
4
5
6 "Jia chou bu ke wai yang," which means one should not reveal family disgrace to
7
8
9 outsiders⁵³⁻⁵⁶. It is possible that the belief in this notion of family disgrace could lead
10
11 some participants to under-report victimization and male-to-female violence.
12
13
14
15

16 **Conclusion**

17
18
19 Findings from this study indicated that the overall IPV prevalence in Guangyuan rural
20
21 areas was close to that in northwest of China. And we found personal and interpersonal
22
23 factors, especially social support, were linked to the occurrence of IPV.
24
25

26 In 2016, China implemented its first law against violence, which emphasizes global
27
28 responsibility of different societal sectors to stop violence, including government
29
30 departments, judiciary authorities, non-governmental organizations, enterprises and
31
32 institutions, and the citizens. Future work is needed to develop, test, and then
33
34 disseminate IPV prevention and intervention programs. Our data suggested that
35
36 reinforcing social support networks offers the potential to enhance real and perceived
37
38 protection, which in turn may reduce the morbidity and mortality associated with IPV.
39
40
41
42
43
44
45

46 **Funding**

47
48
49 This study was funded by NIH Grants D43 TW009101 and D43 TW009101-01S1 (E.D.
50
51 Caine, PI).
52

53 **Conflict of interest**

1 The authors declare that they have not conflict of interest.

2

3 **Ethical Statement**

4 The Ethics Committee of Sichuan University reviewed and approved the protocol,
5 including the verbal informed consent process (NO.2011004-1). The University of
6 Rochester Research Subjects Review Board reviewed the approval from Sichuan
7 University and approved analyses of de-identified data.

8

9 **Contribution**

10 FH designed the survey instruments, implemented the field survey, monitored data
11 collection, cleaned the data, developed the plan for analysis, analyzed the data, drafted
12 and revised the paper. CC designed the survey instruments, trained interviewers,
13 assisted with the analysis plan, revised the paper, and supervised FH. MNW designed the
14 survey instruments, trained interviews, and revised the paper. EDC initiated the project,
15 revised the paper, and supervised FH. PQ initiated the project, designed the survey
16 instruments, monitored data collection, cleaned the data, and revised the paper. All
17 authors had full access to all the data in the study and take responsibility for the
18 integrity of the data and the accuracy of the data analysis. All authors read and approved
19 the final manuscript.

20

21 **Acknowledgments**

22 We acknowledge the important input of Wang Fang, Wang Ying, Feng Fei, and Hua Junjie
23 et al. from Sichuan University, Zhang Yinghui and Ma Mei from Guangyuan Mental Health
24 Center, Chen Shulin from Zhejiang University, and Jennifer Thompson Stone from
25 University of Rochester.

1

2 **Data sharing statement**3 The data set is available from the corresponding author at qiupeiyuan@scu.edu.cn.

4

5 **Reference**

- 6 1. Krantz G, Garcia-Moreno C. Violence against women. *Journal of Epidemiology and*
7 *Community Health* 2005;59(10):818-21. doi: 10.1136/jech.2004.022756
- 8 2. World Health Organization. Understanding and addressing violence against women.
9 2012.
10 http://apps.who.int/iris/bitstream/10665/77432/1/WHO_RHR_12.36_eng.pdf
11 (accessed September 18th, 2015).
- 12 3. World Health Organization. Violence against women: Intimate partner and sexual
13 violence against women 2013 [Available from:
14 <http://www.who.int/mediacentre/factsheets/fs239/en/> accessed January 17
15 2014.
- 16 4. Tang CS-k. Corporal punishment and physical maltreatment against children: a
17 community study on Chinese parents in Hong Kong. *Child Abuse & Neglect*
18 2006;30(8):893-907.
- 19 5. Wang X, Ho PSY. Violence and desire in Beijing a young Chinese woman's strategies of
20 resistance in father-daughter incest and dating relationships. *Violence Against*
21 *Women* 2007;13(12):1319-38.
- 22 6. Tam SY, Tang CS-k. Comparing wife abuse perceptions between Chinese police officers
23 and social workers. *J Fam Viol* 2005;20(1):29-38.
- 24 7. Parish WL, Wang T, Laumann EO, et al. Intimate partner violence in China: national
25 prevalence, risk factors and associated health problems. *International Family*
26 *Planning Perspectives* 2004:174-81.
- 27 8. Xu X, Zhu F, O'Campo P, et al. Prevalence of and risk factors for intimate partner
28 violence in China. *American journal of public health* 2005;95(1):78-85.
- 29 9. Decker MR, Peitzmeier S, Olumide A, et al. Prevalence and health impact of intimate
30 partner violence and non-partner sexual violence among female adolescents
31 aged 15-19 years in vulnerable urban environments: a multi-country study. *J*
32 *Adolescent Health* 2014;55(6, Supplement):S58-S67. doi:
33 <http://dx.doi.org/10.1016/j.jadohealth.2014.08.022>
- 34 10. Pan T, Hall BJ, Ling L. The association between social resources and depression
35 among female migrants affected by domestic violence. *European journal of*
36 *psychotraumatology* 2014;5:10.3402/ejpt.v5.26528. doi: 10.3402/ejpt.v5.26528
- 37 11. Guo S, Zhao F, Wu J. Study on prevalence of domestic violence and its related factors

- 1 in rural area of China. *Chinese Journal of Public Health* 2007;1:002.
- 2 12. Cao Y, Zhang Y, Guo G, et al. A comparison study on domestic violence by geographic
3 settings in Hunan. *Chinese Mental Health Journal* 2007;21(12):849-52.
- 4 13. Cao Y, Zhang Y, Yang S, et al. A comparartive study on the patterns of domestic
5 violence *Chinese Journal of Clinical Psychology* 2008;16(1):28-30.
- 6 14. Tsai AC. Intimate partner violence and population mental health: why poverty and
7 gender inequities matter. *PLoS medicine* 2013;10(5):e1001440.
- 8 15. Walton-Moss BJ, Manganello J, Frye V, et al. Risk factors for intimate partner violence
9 and associated injury among urban women. *Journal of Community Health*
10 2005;30(5):377-89.
- 11 16. Xu X, Campbell JC, Zhu F. Intimate partner violence against Chinese women-the past,
12 present, and future. *Trauma, Violence, & Abuse* 2001;2(4):296-315.
- 13 17. Krugg E, Dahlberg L, Mercy J, et al. Violence—a global public health approach. World
14 report on violence and health: Geneva, Switz: World Health Organization, 2002.
- 15 18. De Brauw A, Li Q, Liu C, et al. Feminization of agriculture in China? Myths
16 surrounding women's participation in farming. *The China Quarterly*
17 2008;194:327-48.
- 18 19. Hou F, Cerulli C, Wittink M, et al. Depression, social support and associated factors
19 among women living in rural China: a cross-sectional study. *BMC women's health*
20 2015;15(1):28.
- 21 20. Qiu P, Caine E, Hou F, et al. The prevalence of distress and depression among women
22 in rural Sichuan province *PloS one* 2016;11(8):e0161097. doi: 10.1371/
23 journal.pone.0161097
- 24 21. Guangyuan Statistics Intranet. Statistical Communique on the 2012 Guangyuan
25 economic and social development. 2012
- 26 22. Statistic Bureau of Sichuan Province China. Sichuan Province Statistics Yearbook
27 2011
28 <http://www.sc.stats.gov.cn/tjcbw/tjnj/2011/chinese/dir/chinesemenu.htm2011>
29 1 [Available from:
30 <http://www.sc.stats.gov.cn/sctj/default.htm?status=NotExt&menu=20&sub=1,fa>
31 [lse](#) accessed 12.31.2012.
- 32 23. Koenig HG, Westlund RE, George LK, et al. Abbreviating the Duke Social Support
33 Index for use in chronically ill elderly individuals. *Psychosomatics*
34 1993;34(1):61-9. doi: 10.1016/s0033-3182(93)71928-3 [published Online First:
35 1993/01/01]
- 36 24. Zhang J, Conwell Y, Wiczorek WF, et al. Studying Chinese suicide with proxy-based
37 data: reliability and validity of the methodology and instruments in China. *The*
38 *Journal of nervous and mental disease* 2003;191(7):450.
- 39 25. Jia C, Zhang J. Psychometric characteristics of the Duke Social Support Index in a
40 young rural Chinese population. *Death studies* 2012;36(9):858-69.
- 41 26. Straus MA, Douglas EM. A short form of the Revised Conflict Tactics Scales, and
42 typologies for severity and mutuality. *Violence and Victims* 2004;19(5):507-20.

- 1 27. Zhang N, Gao Y, Wu R, et al. Validity and reliability of the short form of the revised
2 Conflict Tactics Scales in women suffering from domestic violence in rural areas.
3 *Chinese Mental Health Journal* 2014;28(5):381-84.
- 4 28. Kääriäinen M, Kanste O, Elo S, et al. Testing and verifying nursing theory by
5 confirmatory factor analysis. *Journal of Advanced Nursing* 2011;67(5):1163-72.
- 6 29. Meyers LS, Gamst G, Guarino AJ. Applied multivariate research: design and
7 interpretation: SAGE Publications, Incorporated 2006:539-68.
- 8 30. Krug EG, Mercy JA, Dahlberg LL, et al. The world report on violence and health. *The*
9 *lancet* 2002;360(9339):1083-88.
- 10 31. Muthén LK, Muthén BO. Mplus user's guide: The comprehensive modeling program
11 for applied researchers 2012.
- 12 32. Byrne BM. Structural equation modeling with Mplus: Basic concepts, applications,
13 and programming. New York: Routledge 2012.
- 14 33. Li Z, Chen Y, Wang X. A survey of domestic violence in rural China. *Chinese Journal of*
15 *Behavioral Medicine and Brain Science* 2003;12(2):228-30.
- 16 34. Tang CS-K, Lai BP-Y. A review of empirical literature on the prevalence and risk
17 markers of male-on-female intimate partner violence in contemporary China,
18 1987–2006. *Aggression and Violent Behavior* 2008;13(1):10-28.
- 19 35. Xue L, Pang S, Wu L, et al. Study on the influential factors of domestic violence against
20 married women in rural area of Tangshan. *Modern Preventive Medicine*
21 2007;34(17):3343-44.
- 22 36. Gao Y, Jacka T. Prevalence of intimate partner violence and its impact on women's
23 mental health in rural Western China: a study of a county in Ningxia. *Journal of*
24 *Peking University Health Sciences* 2012;44(3):379-86.
- 25 37. Taylor RL, Choy HYF. The illustrated encyclopedia of Confucianism: N-Z: The Rosen
26 Publishing Group 2005:496.
- 27 38. Jewkes R. Intimate partner violence: causes and prevention. *The Lancet*
28 2002;359(9315):1423-29.
- 29 39. Levinson D. Family violence in cross-cultural perspective. Thousand Oaks, CA, US:
30 Sage Publications, Inc 1989.
- 31 40. Mager AK. Gender and the making of a South African Bantustan: a social history of
32 the Ciskei, 1945-1959. 1999
- 33 41. Heise L, Ellsberg M, Gottemoeller M. Ending violence against women: population
34 reports Baltimore: Johns Hopkins University 1999.
- 35 42. Callaghan P, Morrissey J. Social support and health: a review. *Journal of Advanced*
36 *Nursing* 1993(18):203-10.
- 37 43. Coker AL, Smith PH, Thompson MP, et al. Social support protects against the negative
38 effects of partner violence on mental health. *Journal of Women's Health &*
39 *Gender-Based Medicine* 2002;11(5):465-76.
- 40 44. Harwell TS, Spence MR. Population surveillance for physical violence among adult
41 men and women, Montana 1998. *American Journal of Preventive Medicine*
42 2000;19(4):321-24.

- 1
2
3 1 45. World Health Organization. Preventing intimate partner and sexual violence against
4 2 women: taking action and generating evidence: World Health Organization 2010.
5 3 46. Romans S, Forte T, Cohen MM, et al. Who is most at risk for intimate partner violence?
6 4 A Canadian population-based study. *Journal of Interpersonal Violence*
7 5 2007;22(12):1495-514.
8 6 47. Vest JR, Catlin TK, Chen JJ, et al. Multistate analysis of factors associated with intimate
9 7 partner violence. *American Journal of Preventive Medicine* 2002;22(3):156-64.
10 8 48. Beydoun HA, Beydoun MA, Kaufman JS, et al. Intimate partner violence against adult
11 9 women and its association with major depressive disorder, depressive symptoms
12 10 and postpartum depression: a systematic review and meta-analysis. *Social*
13 11 *Science & Medicine* 2012;75(6):959-75. doi: 10.1016/j.socscimed.2012.04.025
14 12 49. Zhao F, Guo S, Wang L, et al. Investigation on the patterns and knowledge regarding
15 13 domestic violence among married women in rural areas of China. *Chinese Journal*
16 14 *of Epidemiology* 2006;27(8):664-68.
17 15 50. Chan KL. Study on child abuse and spouse battering: report on findings of household
18 16 survey : a consultancy study commissioned by the Social Welfare Department of
19 17 the Hong Kong SAR. Hong Kong: Department of Social Work and Social
20 18 Admission, the University of Hong Kong. Hong Kong, 2005.
21 19 51. Jewkes R, Levin J, Penn-Kekana L. Risk factors for domestic violence: findings from a
22 20 South African cross-sectional study. *Social Science & Medicine*
23 21 2002;55(9):1603-17.
24 22 52. Guo S, Wu J, Qu C, et al. Physical and sexual abuse of women before, during, and after
25 23 pregnancy. *International Journal of Gynecology & Obstetrics* 2004;84(3):281-86.
26 24 53. Gao G. An initial analysis of the effects of face and concern for other in Chinese
27 25 interpersonal communication. *International Journal of Intercultural Relations*
28 26 1998;22(4):467-82.
29 27 54. Pattie Yuk Yee LF. Hybridity in a guidance curriculum in Hong Kong. *International*
30 28 *Journal for the Advancement of Counselling* 2006;28(4):331-42.
31 29 55. Liu L, Xudong Z, Miller JK. Use of metaphors in Chinese family therapy: a qualitative
32 30 study. *Journal of Family Therapy* 2012
33 31 56. Sze YT, Hou J, Lan J, et al. Brief report: Profiling family therapy users of a therapy
34 32 center in Beijing. *The American Journal of Family Therapy* 2011;39(4):299-306.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Table 1.1 Demographic information of participants

Demographic Characters	n	IPV victim		Prevalence (%)	χ^2	p
		No	Yes			
Age	16-29	204	127	77	19.873	0.001
	30-39	225	159	66		
	40-49	468	320	146		
	50-59	364	276	88		
	60-69	188	138	50		
	70-	52	45	7	13.46	
Education	Never educated	497	375	122	11.378	0.023
	Primary school	623	442	181		
	Junior high school	283	183	100		
	High school	98	50	26		
	College and above	22	15	7		
Family annual income	0-9999 Yuan	265	194	71	3.237	0.663
	10000-19999 Yuan	371	263	108		
	20000-29999 Yuan	346	248	98		
	30000-39999 Yuan	213	152	61		
	40000-49999 Yuan	116	83	33		
	≥50000 Yuan	190	125	65	34.21	
Perceived family financial status*	Affluent	182	124	58	9.091	0.028
	Basic enough	788	582	206		
	Difficult	469	318	151		
	Very difficult	47	28	19		
Comparing with others*	Wealthier	81	62	19	1.363	0.506
	Same	858	606	252		
	Poorer	546	384	162		
Comparing with previous years*	Better	1119	810	309	6.095	0.047
	Same	228	147	81		
	Worse	137	94	43		

2 Note: "*" indicates missing data; "#" indicates using Fisher exact test; "IPV" is the abbreviation
 3 of intimate partner violence.

4

1 Table 1.2 The social support level of participants

Social support	Mean	IPV victim		F	p
		No	Yes		
Total score	37.33±5.13	37.77±4.74	36.28±5.86	26.289	0.000
Social interaction	7.81±1.70	7.82±1.70	7.81±1.68	0.002	0.962
Perceived social support	18.89±2.74	19.15±2.58	18.24±3.00	35.154	0.000
Instrumental social support	10.62±2.38	10.78±2.18	10.23±2.77	16.778	0.000
Family properties	7.54±2.97	7.50±2.93	7.63±3.05	0.604	0.437

2 Note: "IPV" is the abbreviation of intimate partner violence.

3

For peer review only

1 Table 2 IPV experiences of participants

IPV experiences	No (%)	Yes (%)
Ever experienced in the past 12 months	1065 (70.95%)	436 (29.05%)
Physical violence	1386 (92.34%)	115 (7.66%)
Psychological violence	1102 (73.42%)	399 (26.58%)
Sexual violence	1453 (96.80%)	48 (3.20%)

2 Note: "IPV" is the abbreviation of intimate partner violence.

3

For peer review only

1 Table 3 Factor loadings and path coefficients of the final model

Model	Dependent variables	Independent variables	Statistics results		
			Standardized coefficient	<i>p</i>	
Measurement model	OE	A3	0.61	0.00	
		A4	0.79	0.00	
		C1	0.84	0.00	
	PHV	C2	0.98	0.00	
		C3	0.45	0.00	
		C4	0.71	0.00	
	PSV	C5	0.76	0.00	
		C6	0.65	0.00	
		D1	0.55	0.00	
Structure model	SV	D2	0.42	0.00	
		D3	0.40	0.00	
		PHV	SS	-0.12	0.005
	PHV	OE	0.060	0.14	
		PSV	SS	-0.35	0.00
		OE	0.11	0.060	
	SV	SS	-0.12	0.021	
		OE	0.062	0.11	
		OE	A1	-0.18	0.00
Factor related	PSV	A2	0.30	0.00	
		SS	-0.068	0.18	
		A2	0.13	0.012	
	SV	OE	0.41	0.00	
		PHV	0.91	0.00	
		PHV	0.30	0.00	
PSV	PHV	0.46	0.00		
	C5	C1	0.31	0.00	
	D3	D2	0.41	0.00	

2 Note: OE=Objective economic status; SS=Social support; PHV=Physical violence; PSV=

3 Psychological violence; SV=Sexual violence; A1=Age; A2=Education; B1=Family annual income;

4 B2=Family properties; C1= Being pushed, shoved or slapped; C2=Being punched, kicked or

5 beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4= Being threatened to destroy

6 belongings; C5= Sex against will with physical force; C6= Sex against will without physical force;

7 D1=Social interaction; D2=Perceived social support; D3=Instrumental social support.

8

1 Table 4 Direct and indirect effects of the final model

Factors	Physical violence			Psychological violence			Sexual violence		
	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect	Direct effect	Indirect effect	Total effect
Age	—	0.005	0.005	—	0.030	0.030	—	0.005	0.005
Education	—	-0.011	-0.011	—	-0.056*	-0.056*	—	-0.011	-0.011
Objective economic status	0.060	-0.047*	0.013	0.11	-0.014*	-0.033	0.062	-0.047*	0.015
Social support	-0.12*	—	-0.12*	-0.35*	—	-0.35*	-0.12*	—	-0.12*

2 Note: "*" indicates $p < 0.05$.

3

For peer review only


Societal level	Community level	Interpersonal level	Personal level
Divorce regulations by government	Acceptance of traditional gender roles	Education disparity	Young age
Lack of legislation on IPV within marriage	High proportion of poverty	Male dominance in the family	Low socio-economic status/ income
Protective marriage law	High proportion of unemployment	Economic stress	Low education
Traditional gender norms and social norms supportive of violence	High proportion of female literacy	Low female contribution to household income	Mental disorder
	Low proportion of women with high level of autonomy	Number of children	Substance use
	Social isolation of nuclear families	Infidelity	Acceptance of violence and exposure to prior abuse
		Sexual jealousy	Child maltreatment
		Marital dissatisfaction	

2 Figure 1 The socio-ecological model of risk factors of IPV

peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1



2

3

Figure 2 The final confirmatory factor analysis model

4

Note: OE=Objective economic status; SS=Social support; SE= Subjective economic status; PHV=Physical violence; PSV= Psychological violence; SV=Sexual violence; A1=Age; A2=Education; A3=Family annual income; A3=Family properties; C1= Being pushed, shoved or slapped; C2=Being punched, kicked or beat-me-up; C3=Being insulted, swore, shouted, yelled at; C4= Being threatened to destroy belongings; C5= Sex against will with physical force; C6= Sex against will without physical force; D1=Social interaction; D2=Perceived social support; D3=Instrumental social support.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

Societal level	Community level	Interpersonal level	Personal level
Divorce regulations by government	Acceptance of traditional gender roles	Education disparity	Young age
Lack of legislation on IPV within marriage	High proportion of poverty	Male dominance in the family	Low socio-economic status/ income
Protective marriage law	High proportion of unemployment	Economic stress	Low education
Traditional gender norms and social norms supportive of violence	High proportion of female literacy	Low female contribution to household income	Mental disorder
	Low proportion of women with high level of autonomy	Number of children	Substance use
	Social isolation of nuclear families	Infidelity	Acceptance of violence and exposure to prior abuse
		Sexual jealousy	Child maltreatment
		Marital dissatisfaction	

Figure 1 The socio-ecological model of risk factors of IPV

303x121mm (300 x 300 DPI)




Figure 2 The final confirmatory factor analysis model

499x240mm (300 x 300 DPI)

Appendix 1 Latent and observation factors

Latent factors	Observed factors
	Age, A1
	Education, A2
Objective economic status, OE	Family annual income, A3
	Family properties, A4
Subjective economic status, SE	Perceived family financial status, B1
	Comparing with others, B2
	Comparing with previous years, B3
Physical violence, PHV	Being pushed, shoved or slapped, C1
	Being punched, kicked or beat-me-up, C2
Psychological violence, PSV	Being Insulted, swore, shouted, yelled at, C3
	Being threatened to destroy belongings, C4
Sexual violence, SV	Sex against will with physical force, C5
	Sex against will without physical force, C6
Social support, SS	Social interaction, D1
	Perceived social support, D2
	Instrumental social support, D3

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page number
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2,3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5
Objectives	3	State specific objectives, including any prespecified hypotheses	7,8
Methods			
Study design	4	Present key elements of study design early in the paper	8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	8
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	8,9
		<i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls	
		<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed	-
		<i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	10-12
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	10-12
Bias	9	Describe any efforts to address potential sources of bias	21
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	13,14
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	12-14
		(b) Describe any methods used to examine subgroups and interactions	No applicable
		(c) Explain how missing data were addressed	14
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed	8
		<i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed	
		<i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	
		(e) Describe any sensitivity analyses	No applicable

Continued on next page

Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	14
		(b) Give reasons for non-participation at each stage	21
		(c) Consider use of a flow diagram	-
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	14, 15
		(b) Indicate number of participants with missing data for each variable of interest	14
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	-
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	-
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	-
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	15,16
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	No applicable
		(b) Report category boundaries when continuous variables were categorized	14, 15
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	No applicable
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	No applicable
Discussion			
Key results	18	Summarise key results with reference to study objectives	17-21
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	21
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	17-21
Generalisability	21	Discuss the generalisability (external validity) of the study results	21, 22
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	22

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.