

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Young adult smokers' perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes: An online survey

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019662
Article Type:	Research
Date Submitted by the Author:	19-Sep-2017
Complete List of Authors:	Moodie, Crawford; University of Stirling, Institute for Social Marketing Hiscock, Rosemary; University of Bath, ; 1973 Thrasher, Jim; University of South Carolina Reid, Garth; NHS Health Scotland
Primary Subject Heading:	Smoking and tobacco
Secondary Subject Heading:	Public health
Keywords:	Smoking, Packaging, Inserts, Cigarettes

SCHOLARONE™
Manuscripts

Title: Young adult smokers' perceptions of cigarette pack inserts promoting cessation and
dissuasive cigarettes: An online survey

Authors: Crawford Moodie¹, Rosemary Hiscock², Jim Thrasher³, Garth Reid⁴

Affiliations:

¹ Institute for Social Marketing, Faculty of Health Sciences and Sport, University of Stirling,
Stirlingshire, Scotland

² Department for Health, University of Bath, England

³ Health Promotion, Education, and Behavior, Arnold School of Public Health, University of
South Carolina, Columbia, United States

⁴ Policy Evaluation, NHS Health Scotland, Edinburgh, Scotland

Corresponding author: Crawford Moodie, Institute for Social Marketing, Department of
Health Sciences and Sport, University of Stirling, Stirlingshire, Scotland FK9 4LA. Tel: +44
(0)1786 466456. Email: c.s.moodie@stir.ac.uk

Keywords: Smoking, Packaging, Inserts, Cigarettes

Word count: 4596

1
2
3 **Young adult smokers' perceptions of cigarette pack inserts promoting cessation and**
4 **dissuasive cigarettes: An online survey**
5
6
7
8

9 **ABSTRACT**

10
11 **Objectives:** To explore young adult smokers' perceptions of cigarette pack inserts promoting
12 cessation and cigarettes designed to be dissuasive.
13

14
15 **Design:** Cross-sectional online survey.
16

17
18 **Setting:** United Kingdom.
19

20
21 **Participants:** Of the 1970 young adult smokers recruited, the final sample was 1766
22 (89.6%); 50.3% were male and 71.6% white British. To meet the inclusion criteria
23 participants had to be 16-34 years old and smoke factory-made cigarettes.
24

25
26 **Primary and secondary outcome measures:** Salience of inserts, perceptions of inserts as
27 information provision, perceptions of inserts on quitting, support for inserts, and perceived
28 appeal, harm and trial of three cigarettes (a standard cigarette, a standard cigarette displaying
29 the warning 'Smoking kills' on the cigarette paper, and a green cigarette).
30

31
32 **Results:** Half the sample indicated that they would read inserts with three-fifths indicating
33 that they be a good way to provide information about quitting (61%). Just over half the
34 sample indicated that inserts would make them think more about quitting (53%), help if they
35 decided to quit (52%), are an effective way of encouraging smokers to quit (53%), and
36 supported having them in all packs (55%). Participants who smoked factory-made cigarettes
37 and other tobacco products (compared to exclusive factory-made cigarette smokers), had
38 made a quit attempt within the last six months (compared to those that had never made a quit
39 attempt), or were likely to make a successful quit attempt in the next six months (compared to
40 those unlikely to make a quit attempt in the next six months), were more likely to indicate
41 that inserts could assist with cessation. Multivariable logistic regression modelling suggested
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that the two dissuasive cigarettes were considered much less desirable (less appealing, more
4 harmful, less likely to be tried) than the standard cigarette.

5
6
7 **Conclusions:** Inserts and dissuasive cigarettes offer policy makers additional ways of using
8 the pack to reduce smoking.
9
10

11 12 13 **Strengths and limitations of this study**

- 14
15
16 ■ The main strength of this study is that it allows an insight into how young adult
17 smokers perceive two innovative tobacco control measures (pack inserts promoting
18 cessation and dissuasive cigarettes).
19
- 20
21
22 ■ The main limitations are that the study does not provide any insight into actual smoking
23 behaviour, the novelty of the stimuli and forced exposure to this, and the use of self-
24 selection.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

While packaging remains a key marketing driver for tobacco companies, more than 100 countries now require pictorial health warnings on cigarette packs,¹ which can limit pack appeal.² Some countries have gone even further by implementing plain (or standardised) packaging, which severely reduces the promotional power of the pack. The United Kingdom (UK) became the third country to fully implement standardised packaging in May 2017, following Australia in December 2012 and France in January 2017. In the UK all cigarette packs must be drab brown with pictorial warnings on 65% of the front and back of packs and additional health messages on 50% of the sides of the pack. Although these changes have reduced the ability of tobacco companies to use the pack to create favourable perceptions of the brand and of smoking, there is clearly more scope for using the packaging to dissuade consumers. Regulators and academics have typically focused on the exterior of the cigarette pack, with little consideration of how the pack interior, for instance pack inserts or cigarettes, could potentially be used to encourage smokers to think about their smoking behaviour. This is the focus of our study.

Tobacco companies have used the inside of the cigarette pack to communicate with consumers since the late 19th century, via cigarette cards, coupons and promotional inserts. Only in Canada are they required, by law, to include pack inserts with health messaging. Sixteen text-only inserts were required in packs between 2000 and 2012, with nine encouraging cessation and seven providing health risk information.³ These were replaced with eight new inserts, with coloured graphics and tips about quitting or the benefits of doing so, in 2012. Few studies have explored perceptions of pack inserts,⁴⁻⁸ with only two assessing smokers' perceptions of, and responses to, the inserts used in Canada.⁹⁻¹¹ In focus group research in Scotland,⁹ with smokers aged 16 and over who were shown seven of the inserts used in Canada, the general view was that they would capture attention and be read due to

1
2
3 their novelty and visibility when opening the pack. The positive messaging was liked and
4 thought to increase message engagement. The inserts were often preferred to the on-pack
5 warnings, although both were deemed necessary. Some participants suggested that inserts
6 could encourage them to stop smoking, and they were generally considered to have the
7 potential to alter the behaviour of younger people, would-be smokers and those wanting to
8 quit.⁹ In Canada, a longitudinal online survey with smokers aged 18 and over found that
9 between 26% and 31% at each wave reported having read pack inserts at least once in the
10 prior month; those intending to quit or having recently tried to do so were significantly more
11 likely to have read them.¹⁰ In addition, while reading warnings on the pack exterior decreased
12 over time, reading pack inserts increased over time, with more frequent reading
13 independently associated with self-efficacy to quit, quit attempts, and sustained quitting at
14 follow-up.¹¹

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29 The cigarette itself is also an important communications tool,^{12,13} which has long been
30 used by tobacco companies as a marketing device but has yet to be used by regulators to deter
31 smoking. As cigarettes are primarily responsible for tobacco related mortality and morbidity
32 and predicted to continue to dominate the global market for some time yet,¹⁴ research
33 exploring the potential impact of standardising the appearance of cigarettes to make them less
34 desirable is long overdue. Some recent research has examined consumer perceptions of
35 'dissuasive' cigarettes, including unattractively coloured cigarettes,^{15,16} cigarettes with the
36 warning 'Smoking kills' on the cigarette paper,^{17,18} and cigarettes displaying the 'minutes of
37 life lost due to smoking' on the cigarette paper.¹⁹ In each of these studies the dissuasive
38 cigarettes were generally viewed more negatively than regular cigarettes. For instance, a
39 qualitative study with young women smokers in New Zealand found that unattractively
40 coloured cigarettes, particularly green or brown coloured cigarettes, were perceived as more
41 harmful than other cigarettes, with it less likely that they or others their age would want to
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 use them.¹⁵ An in-home survey in the UK with 11-16 year olds, who were shown an image of
4 a cigarette stick displaying ‘Smoking kills’, found that 53% indicated that this would make
5 people want to give up smoking, 71% indicated that it would put people off starting to smoke,
6 and 85% supported having a warning on all cigarettes.¹⁸
7
8
9
10

11 In this study our objective was to explore, for the first time, young adult smokers’
12 perceptions of pack inserts and dissuasive cigarettes (a cigarette displaying the warning
13 ‘Smoking kills’ and a green coloured cigarette).
14
15
16
17
18
19

20 **METHODS**

21 **Design and sample**

22
23
24 An online survey was conducted in January-February 2016 with smokers aged 16-34 years
25 old in the UK; an online survey is a suitable approach for this age group given that 99% of
26 16-34 year olds in the UK are recent internet users.²⁰ The sample was recruited by online
27 market research company ‘Research Now’ (www.researchnow.com). The inclusion criteria
28 were that participants were factory-made cigarette smokers and aged 16-34 years. After
29 Research Now excluded those who had completed the survey in less than the minimum
30 completion time, which they had set prior to data collection commencing (n=193), and those
31 providing responses to open-ended questions that indicated that they had not taken the survey
32 seriously (n=11), the final sample was 1766 (89.6%). The final sample was 50.3% male, with
33 53.9% aged 25-34 years and 71.6% white British. Most participants smoked 10 or less
34 cigarettes per day, with 46.0% exclusive factory-made cigarette smokers (see Table 1 for
35 sample and smoking-related characteristics).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 Table 1 here
56
57
58
59
60

Procedure

An email invite was sent by Research Now to their online panel in the UK; Research Now is an established online market research company with their panel recruited from a range of internet sites, advertising and partnerships with other websites. Those eligible for inclusion were presented with an information page explaining the study aim (to explore what young adult smokers thought about cigarettes and pack inserts), and relevant ethical information (their right to withdraw at any time, assurances of confidentiality and anonymity, and contact details if they had any concerns). They were then presented with a consent page, with consent required for participation. Survey questions were presented in the same order for all participants, except the questions exploring perceptions of the three cigarettes (standard cigarette, warning cigarette, green cigarette), where the ordering was randomised; the ordering of the presentation of the three cigarettes (shown in Figure 1) was also randomised. There was no missing data as participants could only proceed to the next question if they had provided an answer to the previous question.

Figure 1 here

For each of the inserts questions participants were shown an image of one of four inserts, see Figure 2, chosen from the eight used in Canada as they were considered most relevant to our sample. The words 'Health Canada' were removed from the bottom of each insert to make them more relevant for participants in the UK. The median time for survey completion was 9 minutes 28 seconds. Participants received a nominal incentive for participation, as is common for online panels. The study received ethical approval from the School of Health Sciences Ethics Committee at the University of Stirling.

1
2
3
4
5 Figure 2 here
6
7
8

9 **Measures**
10

11
12
13 *Inserts: Salience and information provision*
14

15 Participants were asked 'If this type of insert was in your cigarette pack, do you think that
16 you would read it?' and 'If this type of insert was in your cigarette pack, do you think that
17 you would read it if you were interested in quitting?' They were also asked 'Do you think
18 that inserts would be a good way to provide information to smokers about quitting?'
19
20
21
22

23 Response options for each were 'Yes', 'No' and 'Not sure'.
24
25
26

27
28
29 *Inserts: Cessation*
30

31 Three questions assessed to what extent participants agreed or disagreed that inserts would
32 make them think about quitting, and help them quit: 'Do you agree or disagree that having
33 these types of inserts in every cigarette pack would make you think more about quitting?',
34
35
36
37
38
39
40
41
42
43

44 Response options for each were 'Strongly disagree', 'Disagree', 'Neither agree nor disagree',
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

50
51 *Inserts: Support*
52

53 A five-point semantic scale assessed support, with anchors 'All cigarette packs should have
54 inserts like this in them-No cigarette packs should have inserts like this in them'.
55
56
57
58
59
60

Cigarette design: Appeal, harm and trial

Seven-point semantic scales assessed appeal, harm and likely trial. Appeal was assessed via four scales, with anchors 'Attractive-Unattractive', 'Stylish-Not stylish', 'Not nice to be seen with-Nice to be seen with' and 'Not appealing to people my age-Appealing to people my age'. Harm was assessed via two scales, with anchors 'Looks harmful to health-Does not look harmful to health' and 'Makes me think about the dangers of smoking-Does not make me think about the dangers of smoking'. Likely trial was assessed via two scales, 'If a friend offered you each of these cigarettes, how likely would you be to try them?' and 'If someone your age who had never smoked before was going to try a cigarette, how likely do you think they would be to try each of these cigarettes?' Both scales assessing trial ranged from 'Not at all likely' to 'Very likely'.

Sociodemographic characteristics

Age, gender, ethnicity, educational attainment and economic status (based on chief income earner) were obtained. A count procedure was used to create a variable for low socioeconomic status (SES): low education (General Certificate of Secondary Education: GCSE or below) and/or low economic status (routine or manual occupation, long-term unemployed or long-term sick or disabled).

Smoking behaviour

Smoking status was assessed with 'Which of these best describes you?' with response options: 'I have never smoked', 'I used to smoke, but don't now', 'I smoke, but not every day', and 'I smoke every day'. Type of products used was assessed with 'What type(s) of tobacco products do you smoke?' with response options: 'Only factory-made (packet)

1
2
3 cigarettes', 'Factory-made and roll-your-own cigarettes', 'Factory-made cigarettes and other
4 tobacco products (e.g. cigars, shisha, etc)', 'Only roll-your-own cigarettes' and 'Only other
5 tobacco products (e.g. cigars, shisha, etc)'. The Heaviness of Smoking Index (HSI)²¹ was
6
7 used as a measure of dependence, based on daily consumption and time to first cigarette.
8
9

10 11 12 13 *Quitting and self-efficacy*

14
15 Participants were asked 'Have you ever made an attempt to quit smoking that lasted at least
16 24 hours?' (Yes within the last six months, Yes more than six months ago, I have never tried
17 to quit for more than 24 hours). They were also asked 'How likely are you to try to quit
18 smoking within the next six months?' (Not at all, A little, Moderately, Very, Extremely,
19 Don't know), with those responding 'Not at all', 'A little', 'Moderately' or 'Don't know'
20 classified as 'Unlikely to make a quit attempt in the next six months'. To measure quitting
21 self-efficacy, participants were asked 'If you decided to quit smoking in the next six months,
22 how sure are you that you would succeed?' (Not at all, A little, Moderately, Very, Extremely,
23 Don't know). Those who responded to the likelihood of quitting question with 'Very or
24 'Extremely' and to the quitting efficacy question with 'Not at all', 'A little', 'Moderately' or
25 'Don't know' were classified as 'unlikely to make a successful quit attempt in the next six
26 months'. Those who responded 'Very' or 'Extremely' to both questions were classified as
27 'likely to make a successful quit attempt in the next six months'.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **Analysis**

47
48 Data was analysed using Microsoft office Excel 2013, SPSS v22 and v23 and MIWin v2.33.²²
49
50 The insert variables were dichotomised into yes/agreement and no/disagreement/neutral/not
51 sure/don't know. The dichotomised insert variables were the outcomes of the logistic
52 regression models. The independent variables were gender, age, education, ethnicity,
53
54
55
56
57
58
59
60

1
2
3 dependence (tertiles of HSI), tobacco product(s) smoked, previous quit attempt lasting 24
4 hours, and likely efficacy of a quit attempt in the next six months. Percentages in agreement
5 were calculated. Age, gender and education (as a measure of SES) were entered into all
6 models to account for any sampling inadequacies. Other variables were entered where $p < 0.10$
7 in chi square tests.
8
9
10
11
12

13
14 The cigarette variables were assessed using seven-point semantic scales, with
15 percentages calculated for those indicating one of the three points nearest the undesirable
16 anchor (e.g. unattractive, not nice to be seen with, looks harmful to health). Differences
17 between the three cigarettes were tested using Cochran's Q and pairwise comparisons. A
18 factor analysis of the eight perception variables, collated for all three cigarettes, was
19 undertaken, with checks indicating that the data was suitable for factor analysis (Kaiser
20 Meyer Olkin=0.845, Bartlett's test of sphericity (approx. chi-square 18062.842, $df=276$,
21 $p < 0.001$), with no correlations between the variables > 0.9). The extraction method used was
22 Principal Axis Factoring and the criteria for extraction was eigenvalues > 1 . All eight variables
23 loaded on a single factor > 0.5 . High factor scores indicated that a cigarette was desirable and
24 low scores that it was undesirable. Visual inspection and the Kolmogorov-Smirnov test
25 indicated that the factor was non-normal (because responses for the dissuasive cigarettes
26 indicated they were undesirable generally) and attempts to normalise it using normit rankit
27 methods failed. Thus the factor was divided into tertiles and the tertile indicating undesirable
28 factor scores was compared with the other two tertiles. This was the outcome variable in
29 regression analysis.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 Multilevel logistic regression modelling, with second order PQL linearization, was
49 undertaken with cigarette type (at level one) clustered with individual participants (at level
50 two). All models included cigarette type as a fixed effect where the standard cigarette was
51 compared with the warning cigarette and green cigarette. Other fixed effects at the individual
52
53
54
55
56
57
58
59
60

(participant) level were sociodemographic and smoking-related characteristics. This main effects model tested which characteristics were associated with perceiving cigarettes as desirable. In order to understand which characteristics differentiated the desirability of the three types of cigarettes, interactions between cigarette type and each significant characteristic were tested. One interaction was found. Interacting variables were substituted by a cross classified variable (derived from cigarette type and the variable with which cigarette type significantly interacted). The reference category of the cross classified variable was varied in order to understand the interaction.

RESULTS

Perceptions of inserts

Half the sample indicated that they would read inserts, with approximately three-fifths indicating that they would read them if interested in quitting (60%), and that they would be a good way to provide information about quitting (61%). Just over half strongly agreed/agreed that inserts may make them think more about quitting (53%), help them if they decided to quit (52%), that they are an effective way of encouraging smokers to quit (53%), and that all cigarette packs should have inserts (55%), see Table 2.

Table 2 here

Sociodemographic differences in perceptions of inserts

Women were more likely than men to indicate that they would read inserts (aOR=1.24; 95%CI 1.02-1.50), and 25-34 year olds less likely than 16-19 year olds to think that they were a good way of providing information about quitting (aOR=0.76; 95%CI 0.60-0.98).

1
2
3 Compared with white British participants, white non-British (aOR=0.70; 95%CI 0.50-0.98)
4 and Asian (aOR=0.67; 95%CI 0.49-0.92) participants were less likely to suggest that they
5 would read inserts if trying to quit, white non-British (aOR=0.58; 95%CI 0.41-0.81) and
6
7 Black (aOR=0.61; 95%CI 0.38-0.98) participants were less likely to indicate that inserts
8 would make them think about quitting, and white non-British (aOR=0.62; 95%CI 0.44-0.87)
9 and Asian (aOR=0.70; 95%CI 0.51-0.96) participants were less likely to support having
10 inserts in all packs, see Table 3a.
11
12
13
14
15
16
17
18
19

20 **Smoking-related differences**

21
22 Compared to exclusive factory-made cigarette smokers, those who also smoked roll-your-
23 own cigarettes were more likely to indicate they would read inserts (aOR=1.35; 95%CI 1.09-
24 1.66), read them if trying to quit (aOR=1.61; 95%CI 1.30-2.00), that they would make them
25 think about quitting (aOR=1.31; 95%CI 1.06-1.62), help them if they decided to quit
26 (aOR=1.31; 95%CI 1.06-1.61), and that they would be an effective way of encouraging
27 smokers to quit (aOR=1.27; 95%CI 1.03-1.56). Compared to exclusive factory-made
28 cigarette smokers, those who also smoked other tobacco products (e.g. cigars, shisha) were
29 more likely to indicate they would read inserts if trying to quit (aOR=1.39; 95%CI 1.04-1.86)
30 and that inserts might help them if they decided to quit (aOR=1.34; 95%CI 1.01-1.78).
31
32
33
34
35
36
37
38
39
40
41

42 Participants who had made a quit attempt more than six months ago (aOR=1.30;
43 95%CI 1.00-1.69), or within the last six months (aOR=1.67; 95%CI 1.29-2.15), were more
44 likely to indicate that they would read inserts than those who had never made a quit attempt.
45 Those who had made a quit attempt in the last six months were also more likely than those
46 who had never made a quit attempt to indicate that inserts were a good way to provide
47 information about quitting (aOR=1.54; 95%CI 1.20-1.98), that they would read them if trying
48 to quit (aOR=1.51; 95%CI 1.17-1.94), make them think about quitting (aOR=1.46; 95%CI
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 1.14-1.88), help them if they decided to quit (aOR=1.35; 95%CI 1.05-1.73), and that they
4 would be an effective way of encouraging smokers to quit (aOR=1.33; 95%CI 1.04-1.71).
5
6

7 Compared to those likely to make a successful quit attempt in the next six months,
8 those unlikely to make a quit attempt in the next six months were less likely to indicate that
9 they would read inserts (aOR=0.58; 95%CI 0.44-0.75), read them if trying to quit
10 (aOR=0.74; 95%CI 0.55-0.99), that they would make them think about quitting (aOR 0.59
11 (0.45 to 0.78), help them if they decided to quit (aOR=0.51; 95%CI 0.38-0.67), that they
12 would be effective for smokers if they decided to quit (aOR=0.55; 95%CI 0.41-0.73), or
13 support them (aOR=0.56; 95%CI 0.42-0.74). Compared to those likely to make a successful
14 quit attempt in the next six months, those unlikely to make a successful quit attempt in the
15 next six months were more likely to read inserts if trying to quit (aOR=1.43; 95%CI 1.00-
16 2.06), thought that they were a good way to provide information to smokers about quitting
17 (aOR=1.46; 95%CI 1.02-2.08), and support them (aOR=1.43; 95%CI 1.00-2.04), see Table
18 3b.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 Table 3 here
36
37
38

39 **Perceptions of cigarette design**

40 With respect to harm, participants were less likely to think that the standard cigarette (SC)
41 (38.8%) looked harmful than the warning cigarette (WC) (69.1%) or green cigarette (GC)
42 (70.2%) ($p<0.001$), and that the SC (20.9%) made them think more about the dangers of
43 smoking than the WC (58.1%) or GC (53.5%) ($p<0.001$). Participants were also more likely
44 to indicate that the WC would make them think of the dangers of smoking than the GC
45 ($p=0.01$). In terms of appeal, participants were more likely to consider the SC (25.2%)
46 attractive than the WC (61.7%) or GC (68.7%) ($p<0.001$), and the SC (37.4%) as stylish than
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the WC (66.0%) or GC (69.4%) ($p<0.001$). The SC (19.8%) was also considered to be nicer
4 to be seen with than the WC (55.2%) or GC (60.2%) ($p<0.001$), and the SC (17.8%) was
5 viewed as not as appealing to people their age as the WC (51.5%) or GC (57.4) ($p<0.001$). In
6 terms of trial, 79.4% indicated that they would try a SC if offered by a friend (35.7% WC,
7 21.5% GC), and 70.1% indicated that a never smoker their age would be most likely to try a
8 SC (21.1% WC, 16.5% GC) (both $p<0.001$).

18 **Perceptions of cigarette desirability**

19
20 Main effects multivariable logistic regression modelling suggested that in comparison to the
21 SC, the WC (aOR=17.71; 95%CI 13.75-22.80) and GC (aOR=30.88; 95%CI 23.98-39.76)
22 were much more likely to be perceived as undesirable (i.e. less appealing, more harmful, less
23 likely to be tried). The model also indicated which smokers were more likely to rate the
24 cigarettes as undesirable: women were more likely than men (aOR=1.30; 95%CI 1.10-1.54),
25 and low SES more likely than those not low SES (aOR=1.26; 95%CI 1.06-1.50), to consider
26 all three cigarettes undesirable. Compared to exclusive factory-made cigarette smokers, those
27 who also smoked roll-your-own cigarettes (aOR=0.78; 95%CI 0.65-0.93) or other tobacco
28 products (aOR=0.73; 95%CI 0.56-0.93) were less likely to consider all three cigarettes
29 undesirable. Those not likely to make a quit attempt in the next six months were less likely
30 than those likely to make a quit attempt in the next six months (aOR=0.62; 95%CI 0.49-0.78)
31 to consider all three cigarettes undesirable.

32
33
34
35
36
37
38
39
40
41
42
43
44
45
46 Only one significant interaction, between cigarette type and SES, was found. Both
47 SES groups perceived the WC significantly more undesirable than the SC, and the GC
48 significantly more undesirable than the WC. Low SES were significantly more likely than
49 those not low SES to perceive the SC (aOR=17.71; 95%CI 13.75-22.80) and GC
50 (aOR=30.88; 95%CI 23.98-39.76) as undesirable; there was no difference for the WC
51
52
53
54
55
56
57
58
59
60

(aOR=0.99; 95%CI 0.78-1.25), see Figure 3.

Table 4 here

DISCUSSION

Our findings suggest that inserts highlighting the benefits of quitting or providing tips on how to do so may have the potential to encourage cessation, and dissuasive cigarettes may help to reduce the desirability of smoking. Greater attention to how the interior of the cigarette pack could be used to promote cessation appears warranted.

Health messages need to capture attention to be effective.²³ In this regard, at least half our sample indicated that they would read inserts (50%) and read them if interested in quitting (60%). In Canada, observational studies found that approximately a quarter of smokers reported reading them at least once within the last month,¹⁰ increasing to about one-third of smokers over two years of follow-up.¹¹ As in our study, smokers in Canada who had read/would read the inserts were more likely to be female, intend to quit or had recently tried to quit; in our study, they were also more likely to be white British, have moderate dependence, and use factory-made cigarettes and other tobacco products. Future research could explore why dual users (smokers of factory-made cigarettes and other tobacco products) were more likely to indicate that they would read inserts, but as inserts are typically only found in cigarette packs then for those who use other tobacco products they may be seen as more of a novelty and therefore more likely to capture attention.

Approximately three-fifths (61%) of smokers in our study thought that inserts were a good way to provide information about quitting to smokers, with only 25% disagreeing. In comparison, an earlier study in Canada, commissioned by Health Canada, found that 48% of smokers indicated that messaging on inserts was a good way to provide information to

1
2
3 smokers, with 47% disagreeing.⁵ Just over half our sample agreed/strongly agreed that inserts
4 may make them think more about quitting, help them if they decided to quit, and that they are
5 an effective way of encouraging smokers to quit, whereas in New Zealand only 34% of
6 smokers and recent quitters agreed/strongly agreed that inserts would be an effective way of
7 encouraging reduced consumption or quitting.⁶ There may be various reasons for the
8 differences between our findings and earlier research. For instance, when this earlier research
9 was conducted cigarette packs displayed text-only health warnings and it may be that having
10 pictorial warnings on packs, as is required in Scotland, may prompt smokers to look for
11 information on how to quit and the benefits of doing so. Insert design is also likely to be
12 relevant. Whereas the inserts used in earlier research were limited to text, the inserts used in
13 this study (which have been used in Canada since 2012) included coloured graphics, which
14 likely enhanced their impact. This would be consistent with the health communications and
15 warnings literature, which demonstrates the importance of supporting text with
16 pictorials.^{2,23,24} Future research exploring insert design (e.g. use of imagery, inclusion of
17 cessation resource information, length and framing of messages, etc) would be of value.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 More than half our sample supported the inclusion of inserts promoting cessation
36 inside every cigarette pack, with only a fifth opposing this. Within the European Union, the
37 recent Tobacco Products Directive (TPD)²⁵ does not require tobacco companies to include
38 health communication inserts in packs, but allows member states to introduce measures
39 beyond those specified. Among governmental representatives that responded to the
40 consultation on the revision of the TPD there was strong support for improving consumer
41 information via mandatory pictorial warnings, with those supportive arguing that additional
42 information, such as pack inserts, would help to deliver more accurate health information.²⁶ If
43 there is support for inserts among governmental representatives, and little opposition among
44 smokers (the group most likely to be resistant), they are clearly a viable option for regulators.

1
2
3 Tobacco industry journals describe the cigarette as an increasingly important
4 advertising medium for tobacco companies.¹² However, until recently, the public health focus
5 has been on the potential of regulating the contents of cigarettes to reduce palatability or
6 addictiveness,²⁷ with little consideration of the possibility of regulating the appearance of
7 cigarettes to reduce its importance as a promotional tool. We found that the two dissuasive
8 cigarettes were perceived as significantly more harmful and less appealing than the standard
9 cigarette, and less likely to encourage trial. The harm, appeal and trial items loaded onto a
10 single ‘undesirability’ factor, with the dissuasive cigarettes considered much more
11 undesirable than the standard cigarette. The findings are consistent with earlier research,
12 where cigarettes with the warning ‘Smoking kills’ were considered a constant reminder of the
13 associated harms and, partly due to the perceived discomfort of being observed by others
14 smoking a cigarette displaying this message, unappealing for smokers.^{8,16,17,18} Previous
15 studies have also found unattractively coloured cigarettes to be perceived as more harmful
16 than other cigarettes and also repellent, being a cigarette that young people did not think that
17 others their age would use.^{15,16,28,29} As with the inserts, the dissuasive cigarettes (and also the
18 standard cigarette) were considered more desirable among dual users than exclusive factory-
19 made cigarette smokers; again it is not clear why this was the case but further research with
20 dual users, or indeed those also using vaping devices (not assessed in this study), would be
21 fruitful.

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44 In terms of limitations, the cross-sectional design did not allow us to assess causality;
45 that inserts and dissuasive cigarettes are not available on the UK market prevents more robust
46 study designs such as longitudinal studies. Another potential limitation concerns the novelty
47 of the stimuli, which may have influenced responses, and forced exposure to the stimuli. In
48 addition, we only used four inserts, rather than the full set of eight used in Canada, which
49 includes inserts that be less relevant to our sample. While online surveys have been used for
50
51
52
53
54
55
56
57
58
59
60

1
2
3 previous research exploring cigarette packaging, inserts and dissuasive cigarettes,³⁰⁻³³ and are
4 a suitable survey mode for young adults, the use of an online panel and self-selection limits
5 the representativeness of our sample. In addition, the use of semantic differential scales can
6 be criticised because answers can be subject to various response biases, although we
7 attempted to diminish these through varying scale item direction and through our multivariate
8 modelling methodology.

9
10
11 It was argued, over two decades ago, that to offer greater protection to consumers
12 cigarettes should come in plain packs with messaging on both the pack exterior and interior.³⁴
13 This idea is a step closer in the UK, although there will still be no messaging on the pack
14 interior. That more than half of the participants in this study suggested that inserts may help
15 to promote cessation suggests that their inclusion in packs may be a meaningful supplement
16 to the on-pack warnings. Our findings suggest however that to offer the greatest protection to
17 consumers, it may be beneficial to supplement plain packaging and inserts with cigarettes
18 designed to be dissuasive. Unattractively coloured cigarettes would complement the
19 unattractively coloured packs, just as warnings on the cigarette would extend the warnings on
20 the cigarette pack. Both options are clearly viable.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40 **Contributors** CM designed the data collection tool and drafted and revised the paper. RH
41 analysed the data and drafted the Analysis and Results. JF and GR helped design the data
42 collection tool and commented on the paper. All authors read and approved the final
43 manuscript.

44
45
46
47
48
49
50
51
52
53
54 **Funding** This work was supported by Health Scotland. GR, who works for Health Scotland,
55 provided feedback on the survey and paper, but was not involved in the collection, analysis
56 and interpretation of the data.

57
58
59
60 **Competing interests** GR works for Health Scotland, who funded this study.

1
2
3 **Ethics approval** The study obtained ethics approval from the School of Health Sciences
4
5 Ethics Committee at the University of Stirling. Participants provided informed consent before
6
7 participating.
8

9 **Provenance and peer review** Not commissioned; externally peer reviewed.
10

11 **Data sharing statement** No additional data are available.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

REFERENCES

1. Canadian Cancer Society. *Cigarette package health warnings. International status report (Fifth edition)*. www.tobaccolabels.ca/wp/wp-content/uploads/2016/11/Cigarette-Package-Health-Warnings-International-Status-Report-English-CCS-Oct-2016.pdf (accessed 10 Sept 2017).
2. Hammond D. Health warning messages on tobacco products: a review. *Tob Control* 2011;20:327–37.
3. Mahood G. *Canada's tobacco package label or warning system: "telling the truth" about tobacco product risks*. Geneva: WHO, 2010.
4. Tandemar Research Inc. *Tobacco health warning messages, inserts and toxic constituents information study - Final Report*. Toronto: Tandemar Research Inc, 1992.
5. Environics. *Health warning messages on the flip/side and inserts of cigarette packaging. A survey of smokers. A report prepared for Health Canada*, 2000.
6. BRC Marketing and Social Research. *Smoking health warnings study*. Wellington: Ministry of Health, 2004. www.tobaccolabels.ca/wp/wp-content/uploads/2013/12/NZ-2004-Effectiveness-of-Different-Health-Warnings-in-Helping-People-Consider-Their-Smoking-Related-Behaviour-Government-Report.pdf (accessed 12 Aug 2017).
7. Gallopel-Morvan K, Moodie C, Hammond D, *et al*. Consumer understanding of cigarette emission labelling. *Eur J Pub Health* 2011;21:373-5.
8. Moodie C. Novel ways of using tobacco packaging to communicate health messages: Interviews with packaging and marketing experts. *Addict Res Theory* 2016;24:54-61.
9. Moodie C. Adult smokers' perceptions of cigarette pack inserts promoting cessation: A focus group study. *Tob Control* (in press).

10. Thrasher JT, Osman A, Abad EN, *et al.* The innovative use of cigarette package inserts to supplement pictorial health warnings: An evaluation of the Canadian policy. *Nicot Tob Res* 2015;17:870-5.
11. Thrasher JF, Swayampakala K, Cummings KM, *et al.* Cigarette package inserts can promote efficacy beliefs and sustained smoking cessation attempts: A longitudinal assessment of an innovative policy in Canada. *Prev Med* 2016;88:59-65.
12. Rossell S. Ready to roll. *Tob Report* 2017;2:44-5.
13. Smith KC, Washington C, Welding K, *et al.* Cigarette stick as valuable communicative real estate: a content analysis of cigarettes from 14 low-income and middle-income countries. *Tob Control* 2017;26:604-7.
14. Hedley D. Tobacco wars: A new hope - the changing future. *Tob J Int* 2015;3:31-4.
15. Hoek J, Robertson C. How do young adult female smokers interpret dissuasive cigarette cigarettes? *J Social Marketing* 2015;5:21-39.
16. Hoek J, Gendall P, Eckert C, *et al.* Dissuasive cigarette sticks: the next step in standardised ('plain') packaging? *Tob Control* 2016;15:699-705.
17. Moodie C, Purves R, McKell J, *et al.* Novel means of using cigarette packaging and cigarettes to communicate health risk and cessation messages: A qualitative study. *Internat J Mental Health Addiction* 2015;13:333-44.
18. Moodie C, Mackintosh AM, Gallopel-Morvan K, *et al.* Adolescents' perceptions of health warnings on cigarettes. *Nicot Tob Res* 2017;29:1232-7.
19. Hassan L, Shui E. No place to hide: two pilot studies assessing the effectiveness of adding a health warning to the cigarette cigarette. *Tob Control* 2015;24:e3-5.
20. ONS. *Internet users in the UK: 2017*.
www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2017 (accessed 15 Sept 2017).

- 1
2
3 21. Rasbash J, Browne W, Healy M, *et al.* *A user's guide to MLwiN, version 2.3*. Bristol:
4 Centre for multilevel modelling, University of Bristol, 2015.
5
6
7 22. Kozlowski LT, Porter CQ, Orleans T, *et al.* Predicting smoking cessation with self-
8 reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug Alcohol*
9 *Depend* 1994;34:211-6.
10
11
12
13 23. Wogalter MS, Conzola VC, Smith-Jackson TL. Research-based guidelines for
14 warning design and evaluation. *Applied Ergonomics* 2002;33:219-30.
15
16
17 24. Houts PD, Doak CC, Doak LG, *et al.* The role of pictures in improving health
18 communication: A review of research on attention, comprehension, recall, and
19 adherence. *Patient Educ Couns* 2006;61:173-90.
20
21
22
23 25. European Commission. Directive 2014/40/EU of the European parliament and of the
24 Council of 3 April 2014 on the approximation of the laws, regulations and
25 administrative provisions of the Member States concerning the manufacture,
26 presentation and sale of tobacco and related products and repealing Directive
27 2001/37/EC. *Official Journal of the European Union* 2014;L127:1-38.
28
29
30
31 26. European Commission. *Report on the public consultation on the possible revision of*
32 *the Tobacco Products Directive (2001/37/EC)*. Health and Consumers Directorate-
33 General – Directorate D – Health systems and products D4 – Substances of human
34 origin and tobacco control, 2011.
35
36 http://ec.europa.eu/health/tobacco/docs/consultation_report_en.pdf (accessed 10 Sept
37 2017).
38
39
40
41
42
43
44
45
46
47
48 27. Warner K. The national and international regulatory environment in tobacco control.
49 *Pub Health Res Practice* 2015;25:e2531527.
50
51
52
53 28. Ford A, Moodie C, Mackintosh AM, *et al.* Adolescent perceptions of cigarette
54 appearance. *Eur J Pub Health* 2014;24:464-8.
55
56
57
58
59
60

- 1
2
3 29. Moodie C, Ford A, Mackintosh AM, *et al.* Are all cigarettes just the same? Female's
4 perceptions of slim, coloured, aromatised and capsule cigarettes. *Health Educ Res*
5 2015;30:1-12.
6
7
8
9 30. Doxey J, Hammond D. Deadly in pink: The impact of female-oriented cigarette
10 packaging among young women. *Tob Control* 2011;20:353-60.
11
12 31. White CM, Hammond D, Thrasher JF, *et al.* The potential impact of plain packaging
13 of cigarette products among Brazilian young women: an experimental study. *BMC*
14 *Public Health* 2012;12:737.
15
16 32. Kotnowski K, Fong GT, Gallopel-Morvan K, *et al.* The impact of cigarette packaging
17 design among young females in Canada: findings from a discrete choice experiment.
18 *Nicot Tob Res* 2016;18:1348-56.
19
20 33. Hoek J, Gendall P, Maubach N, *et al.* Strong public support for plain packaging of
21 tobacco products. *Austr N Z J Pub Health* 2012;36:405-7.
22
23 34. Mahood G. Canada tobacco package warning system. *Tob Control* 1995;4:10-4.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Sample and smoking-related characteristics

<i>Characteristic</i>	<i>N</i>	<i>%</i>
Total	1766	100.0
Age group		
16-19	413	23.4
20-24	401	22.7
25-34	952	53.9
Gender		
Male	888	50.3
Female	878	49.7
Educational qualifications		
Other qualifications	1357	76.8
None or GCSE	409	23.2
Economic status		
Other status	1350	76.4
Routine or manual occupation, unemployed or long term sick	416	23.6
Socioeconomic status (SES)		
No indicators of low SES	1114	63.1
Low education and/or low SES	652	36.9
Ethnicity		
White British	1264	71.6
White non-British	162	9.2
Black (including mixed black and white)	79	4.5
Asian (including mixed Asian and white)	196	11.1
Other or not declared	65	3.7
Location		
England	1550	87.8
Scotland	109	6.2
Wales	73	4.1
Northern Ireland	34	1.9
Tobacco products used		
Only factory-made (packet) cigarettes	813	46.0
Factory-made and roll-your-own cigarettes	681	38.6
Factory-made cigarettes and other products (e.g. cigars, shisha)	272	15.4
Cigarettes per day		
10 or less	1272	72.0
11-20	433	24.5
21-30	46	2.6
31 or more	15	0.8
Time to first cigarette		
Within 5 minutes	263	14.9
6 to 30 minutes	570	32.3
31 to 60 minutes	315	17.8

<i>Characteristic</i>	<i>N</i>	<i>%</i>
After 60 minutes	618	35.0
Heaviness of Smoking Index (HSI)		
0 little dependence	601	34.0
1	257	14.6
2	418	23.7
3	293	16.6
4	156	8.8
5	28	1.6
6 high dependence	13	0.7
Dependence (Tertiles of HSI)		
Low-dependence	601	34.0
Mid-dependence	675	38.2
High-dependence	490	27.7
Made an attempt to quit smoking that lasted at least 24 hours?		
Yes, within the last six months	788	44.6
Yes, more than six months ago	552	31.3
No, I have never tried to quit smoking for more than 24 hours	426	24.1
How likely are you to try to quit smoking within the next six months?		
Not at all	198	11.2
A little	382	21.6
Moderately	508	28.8
Very	308	17.4
Extremely	272	15.4
Don't know	98	5.5
If you decided to quit smoking in the next six months, how sure are you that you would succeed?		
Not at all	147	8.3
A little	346	19.6
Moderately	612	34.7
Very	297	16.8
Extremely	241	13.6
Don't know	123	7.0
Quit approach		
Moderately or less likely to make quit attempt in next six months (unlikely to make a quit attempt in the next six months)	1186	67.2
Very or extremely likely to attempt but moderately or less likely to succeed (unlikely to make a successful quit attempt in the next six months)	304	17.2
Very or extremely likely to attempt and very or extremely likely to succeed (likely to make a successful quit attempt in the next six months)	276	15.6

Table 2: Perceptions of whether inserts would be read, are a good way to provide information, whether they would help smokers to think about quitting or quit, and support for them

	Yes %	No %	Not sure %
Would they be read	50	37	13
Would they be read if interested in quitting	60	25	15
Good way to provide information about quitting	61	25	14

	Agree %	Disagree %	Neither / Don't know %
Make you think more about quitting	53	18	29
Might help you if you decided to quit	52	19	29
Effective way of encouraging smokers to quit	53	17	30
All packs should have inserts	55	20	25

Table 3a: Logistic regression models exploring perceptions of inserts by sociodemographic characteristics (gender, age, education, ethnicity)^{1,2}

(n=1766)	Would read insert	Would read insert if trying to quit	Inserts make you think about quitting	Inserts might help you quit	Inserts a good way of providing information about quitting	Inserts are an effective way of encouraging smokers to quit	All packs should have inserts
Gender							
Male	1	1	1	1	1	1	1
Female	1.24 (1.02 to 1.50)	1.11 (0.91 to 1.35)	0.98 (0.81 to 1.19)	0.95 (0.79 to 1.15)	1.13 (0.93 to 1.37)	0.88 (0.73 to 1.07)	1.20 (0.99 to 1.46)
Age							
16-19	1	1	1	1	1	1	1
20-24	1.16 (0.87 to 1.54)	0.88 (0.66 to 1.18)	1.18 (0.89 to 1.56)	1.19 (0.89 to 1.58)	0.87 (0.65 to 1.16)	0.97 (0.73 to 1.28)	0.96 (0.72 to 1.29)
25-34	1.25 (0.97 to 1.60)	0.83 (0.65 to 1.07)	0.99 (0.78 to 1.26)	1.18 (0.92 to 1.50)	0.76 (0.60 to 0.98)	0.88 (0.69 to 1.12)	0.84 (0.65 to 1.07)
Education							
GCSEs (or equivalent) or none	1	1	1	1	1	1	1
More than GCSEs (or equivalent)	1.25 (0.99 to 1.58)	1.12 (0.89 to 1.42)	1.22 (0.97 to 1.54)	1.21 (0.97 to 1.52)	1.12 (0.89 to 1.40)	1.19 (0.95 to 1.50)	1.10 (0.87 to 1.40)
Ethnicity							
White British		1	1				1
White but not British		0.70 (0.50 to 0.98)	0.58 (0.41 to 0.81)				0.62 (0.44 to 0.87)
Black (inc mixed black & white)		0.92 (0.57 to 1.49)	0.61 (0.38 to 0.98)				0.99 (0.62 to 1.59)
Asian (inc mixed Asian & white)		0.67 (0.49 to 0.92)	1.19 (0.87 to 1.63)				0.70 (0.51 to 0.96)
other or not declared		0.84 (0.50 to 1.42)	1.06 (0.64 to 1.78)				1.08 (0.64 to 1.81)

¹Note smoking related characteristics (described in table 3b) were also entered into each model

²Blank cells indicate no significant relationship in bivariate analysis


Table 3b: Logistic regression models exploring perceptions of inserts by smoking related characteristics (dependence, tobacco products smoked, quit attempts, self-efficacy to quit)^{1,2}

(n=1766)	Would read insert	Would read insert if trying to quit	Inserts make you think about quitting	Inserts might help you quit	Inserts a good way of providing information about quitting	Inserts are an effective way of encouraging smokers to quit	All packs should have inserts
Dependence (tertiles of HSI)							
Lower dependence	1						1
Mid dependence	1.39 (1.11 to 1.76)						1.02 (0.80 to 1.29)
Higher dependence	1.22 (0.94 to 1.59)						0.86 (0.66 to 1.12)
Tobacco products smoked							
Only factory-made	1	1	1	1		1	
Factory-made and roll-your-own	1.35 (1.09 to 1.66)	1.61 (1.30 to 2.00)	1.31 (1.06 to 1.62)	1.31 (1.06 to 1.61)		1.27 (1.03 to 1.56)	
Factory-made cigarettes and other	1.20 (0.90 to 1.59)	1.39 (1.04 to 1.86)	1.22 (0.92 to 1.63)	1.34 (1.01 to 1.78)		1.20 (0.91 to 1.60)	
Quit attempt lasting at least 24 hours							
No	1	1	1	1	1	1	1
Yes, more than six months ago	1.30 (1.00 to 1.69)	1.12 (0.86 to 1.45)	1.20 (0.93 to 1.56)	1.05 (0.81 to 1.36)	1.16 (0.90 to 1.50)	1.07 (0.82 to 1.38)	0.78 (0.60 to 1.01)
Yes within the last six months	1.67 (1.29 to 2.15)	1.51 (1.17 to 1.94)	1.46 (1.14 to 1.88)	1.35 (1.05 to 1.73)	1.54 (1.20 to 1.98)	1.33 (1.04 to 1.71)	1.06 (0.82 to 1.37)
Efficacy of quit attempt in next 6 months							
Likely to quit	1	1	1	1	1	1	1
Likely to make unsuccessful attempt	1.01 (0.72 to 1.40)	1.43 (1.00 to 2.06)	0.97 (0.69 to 1.37)	0.92 (0.65 to 1.29)	1.46 (1.02 to 2.08)	1.10 (0.78 to 1.55)	1.43 (1.00 to 2.04)
Unlikely to make attempt	0.58 (0.44 to 0.75)	0.74 (0.55 to 0.99)	0.59 (0.45 to 0.78)	0.51 (0.38 to 0.67)	0.76 (0.57 to 1.01)	0.55 (0.41 to 0.73)	0.56 (0.42 to 0.74)

¹Note sociodemographic characteristics (described in table 3a) were also entered into each model

²Blank cells indicate no significant relationship in bivariate analysis

Table 4: Bar charts showing a combination of main effects and interactions between cigarette type and socioeconomic status on odds ratios for undesirability in multivariable models



¹ For the standard cigarette, participants categorised as ‘not low SES’ is the reference group, with an odds ratio of 1, thus they are not displayed



Figure 1: Pack inserts highlighting benefits of quitting or providing tips on how to do so

peer review only




Figure 2: Standard cigarette, warning cigarette and green cigarette

105x35mm (150 x 150 DPI)

peer review only

STROBE 2007 (v4) checklist of items to be included in reports of observational studies in epidemiology*
Checklist for cohort, case-control, and cross-sectional studies (combined)

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any pre-specified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	6
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7-10
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	10-12
Bias	9	Describe any efforts to address potential sources of bias	6
Study size	10	Explain how the study size was arrived at	6
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	10-12
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	10-12
		(b) Describe any methods used to examine subgroups and interactions	10-12
		(c) Explain how missing data were addressed	7
		(d) <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6, 28-29
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	25-26
		(b) Indicate number of participants with missing data for each variable of interest	NA
Outcome data	15*	<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	12-16
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	12-16
		(b) Report category boundaries when continuous variables were	28-29

		categorized	
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10-12
Discussion			
Key results	18	Summarise key results with reference to study objectives	16-18
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	18-19
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	16-18
Generalisability	21	Discuss the generalisability (external validity) of the study results	16-19
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	19

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes among young adult smokers in the United Kingdom: A cross-sectional online survey

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019662.R1
Article Type:	Research
Date Submitted by the Author:	23-Mar-2018
Complete List of Authors:	Moodie, Crawford; University of Stirling, Institute for Social Marketing Hiscock, Rosemary; University of Bath Thrasher, Jim; University of South Carolina Reid, Garth; NHS Health Scotland
Primary Subject Heading:	Smoking and tobacco
Secondary Subject Heading:	Public health
Keywords:	Smoking, Packaging, Inserts, Cigarettes

SCHOLARONE™
Manuscripts

Title: Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes among young adult smokers in the United Kingdom: A cross-sectional online survey

Authors: Crawford Moodie¹, Rosemary Hiscock², Jim Thrasher³, Garth Reid⁴

Affiliations:

¹ Institute for Social Marketing, Faculty of Health Sciences and Sport, University of Stirling, Stirlingshire, Scotland

² Department for Health, University of Bath, England

³ Health Promotion, Education, and Behavior, Arnold School of Public Health, University of South Carolina, Columbia, United States

⁴ Policy Evaluation, NHS Health Scotland, Edinburgh, Scotland

Corresponding author: Crawford Moodie, Institute for Social Marketing, Department of Health Sciences and Sport, University of Stirling, Stirlingshire, Scotland FK9 4LA. Tel: +44 (0)1786 466456. Email: c.s.moodie@stir.ac.uk

Keywords: Smoking, Packaging, Inserts, Cigarettes

Word count: 4899

1
2
3 **Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes**
4
5 **among young adult smokers: A cross-sectional online survey**
6
7

8
9 **ABSTRACT**

10
11 **Objectives:** To explore young adult smokers' perceptions of cigarette pack inserts promoting
12
13 cessation and cigarettes designed to be dissuasive.
14

15
16 **Design:** Cross-sectional online survey.
17

18
19 **Setting:** United Kingdom.

20
21 **Participants:** The final sample was 1766 young adult smokers, with 50.3% male and 71.6%
22
23 white British. To meet the inclusion criteria participants had to be 16-34 years old and smoke
24
25 factory-made cigarettes.

26
27 **Primary and secondary outcome measures:** Salience of inserts, perceptions of inserts as
28
29 information provision, perceptions of inserts on quitting, support for inserts, and perceived
30
31 appeal, harm and trial of three cigarettes (a standard cigarette, a standard cigarette displaying
32
33 the warning 'Smoking kills', and a green cigarette).
34

35
36 **Results:** Half the sample indicated that they would read inserts with three-fifths indicating
37
38 that they be a good way to provide information about quitting (61%). Just over half indicated
39
40 that inserts would make them think more about quitting (53%), help if they decided to quit
41
42 (52%), are an effective way of encouraging smokers to quit (53%), and supported having
43
44 them in all packs (55%). Participants who smoked factory-made cigarettes and other tobacco
45
46 products (compared to exclusive factory-made cigarette smokers), had made a quit attempt
47
48 within the last six months (compared to those that had never made a quit attempt), or were
49
50 likely to make a successful quit attempt in the next six months (compared to those unlikely to
51
52 make a quit attempt in the next six months), were more likely to indicate that inserts could
53
54 assist with cessation. Multivariable logistic regression modelling suggested that compared
55
56
57
58
59
60

1
2
3 with the standard cigarette, the cigarette with warning (adjusted Odds Ratio=17.71; 95%CI
4 13.75-22.80) and green cigarette (adjusted Odds Ratio=30.88; 95%CI 23.98-39.76) were
5
6
7 much less desirable (less appealing, more harmful, less likely to be tried).
8

9 **Conclusions:** Inserts and dissuasive cigarettes offer policy makers additional ways of using
10 the pack to reduce smoking.
11
12

13 **Strengths and limitations of this study**

- 14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
- 18 The main strength of this study is that it allows an insight into how young adult
19 smokers perceive two innovative tobacco control measures (pack inserts promoting
20 cessation and dissuasive cigarettes).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
- 18 Additional limitations include the novelty of the stimuli and forced exposure to this,
19 and the use of self-selection.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
4

INTRODUCTION

While packaging remains a key marketing driver for tobacco companies, more than 100 countries now require pictorial health warnings on cigarette packs,¹ which can limit pack appeal.² Some countries have gone even further by implementing plain (or standardised) packaging, which severely reduces the promotional power of the pack. The United Kingdom (UK) became the third country to fully implement standardised packaging in May 2017, following Australia in December 2012 and France in January 2017. In the UK all cigarette packs must be drab brown with pictorial warnings on 65% of the front and back of packs and additional health messages on 50% of the sides of the pack. Although these changes have reduced the ability of tobacco companies to use the pack to create favourable perceptions of the brand and of smoking, there is clearly more scope for using the packaging to dissuade consumers. Regulators and academics have typically focused on the exterior of the cigarette pack, with little consideration of how the pack interior, for instance pack inserts or cigarettes, which have long been used by tobacco companies to promote their brands, could potentially be used to encourage smokers to think about their smoking behaviour. This is the focus of our study.

Tobacco companies have used the inside of the cigarette pack to communicate with consumers since the late 19th century, via cigarette cards, coupons and promotional inserts. Only in Canada are they required, by law, to include pack inserts with health messaging. Sixteen text-only inserts were required in packs between 2000 and 2012, with nine encouraging cessation and seven providing health risk information.³ These were replaced with eight new inserts, with coloured graphics and positively framed messages about the benefits of quitting or tips on how to do so, in 2012. Few studies have explored perceptions of pack inserts,⁴⁻⁸ with only two assessing smokers' perceptions of, and responses to, the inserts used in Canada.⁹⁻¹¹ In focus group research in Scotland,⁹ with smokers aged 16 and

1
2
3 over who were shown seven of the inserts used in Canada, the general view was that they
4 would capture attention and be read due to their novelty and visibility when opening the pack.
5
6 Inserts were also thought to have a long lasting impact as they would be removed from the
7 pack and remain visible within the household or elsewhere, or as litter.⁹ The positive
8
9 messaging was liked and thought to increase message engagement. The inserts were often
10 preferred to the on-pack warnings, although both were deemed necessary. Some participants
11 suggested that inserts could encourage them to stop smoking, and they were generally
12 considered to have the potential to alter the behaviour of younger people, would-be smokers
13 and those wanting to quit.⁹ In Canada, a longitudinal online survey with smokers aged 18 and
14 over found that between 26% and 31% at each wave reported having read pack inserts at least
15 once in the prior month; those intending to quit or having recently tried to do so were
16 significantly more likely to have read them.¹⁰ In addition, while reading warnings on the pack
17 exterior decreased over time, reading pack inserts increased over time, with more frequent
18 reading independently associated with self-efficacy to quit, quit attempts, and sustained
19 quitting at follow-up.¹¹

20
21
22 The cigarette itself is also an important communications tool,^{12,13} which has long been
23 used by tobacco companies as a marketing device but has yet to be used by regulators to deter
24 smoking. As cigarettes are primarily responsible for tobacco related mortality and morbidity
25 and predicted to continue to dominate the global market for some time yet,¹⁴ research
26 exploring the potential impact of standardising the appearance of cigarettes to make them less
27 desirable is long overdue. Some recent research has examined consumer perceptions of
28 cigarettes that have been designed to be 'dissuasive', including unattractively coloured
29 cigarettes,^{15,16} cigarettes with the warning 'Smoking kills' on the cigarette paper,^{17,18} and
30 cigarettes displaying the 'minutes of life lost due to smoking' on the cigarette paper.¹⁹ In each
31 of these studies the dissuasive cigarettes were generally viewed more negatively than regular
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 cigarettes. For instance, a qualitative study with young women smokers in New Zealand
4 found that unattractively coloured cigarettes, particularly green or brown coloured cigarettes,
5 were perceived as more harmful than other cigarettes, with it less likely that they or others
6 their age would want to use them.¹⁵ An in-home survey in the UK with 11-16 year olds, who
7 were shown an image of a cigarette stick displaying ‘Smoking kills’, found that 53%
8 indicated that this would make people want to give up smoking, 71% indicated that it would
9 put people off starting to smoke, and 85% supported having a warning on all cigarettes.¹⁸

10
11 In this study our objective was to explore, for the first time, young adult smokers’
12 perceptions of pack inserts and dissuasive cigarettes (a cigarette displaying the warning
13 ‘Smoking kills’ and a green coloured cigarette).
14
15
16
17

18 **METHODS**

19 **Design and sample**

20 An online survey was conducted in January-February 2016 with smokers aged 16-34 years
21 old in the UK; an online survey is a suitable approach given that 99% of this age group in the
22 UK are recent internet users.²⁰ The sample (n=1970) was recruited by online market research
23 company ‘Research Now’ from their panel of over 400,000 people (www.researchnow.com).
24 After Research Now excluded those who had completed the survey in less than the minimum
25 completion time (n=193), which they had set prior to data collection commencing, and those
26 providing responses to open-ended questions that indicated that they had not taken the survey
27 seriously (n=11), the final sample was 1766 (89.6% of completed surveys). The final sample
28 was 50.3% male, with 53.9% aged 25-34 years and 71.6% white British. Most participants
29 smoked 10 or less cigarettes per day, with 46.0% exclusive factory-made cigarette smokers
30 (see Table 1 for sample and smoking-related characteristics).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 Table 1 here
6
7
8

9 **Procedure**

10 An email invite was sent by Research Now to their online panel in the UK. Research Now is
11 an established online market research company in the UK and elsewhere,²¹ with their panels
12 recruited from a wide range of sources, such as internet sites, advertising and partnerships
13 with other websites. Research Now, like other online panels, has details of their members'
14 demographics and other characteristics that are used to profile target samples. Response rate
15 details are not available when using this sampling methodology however as recording
16 contact, participation and refusal rates is not practical.²² For those that responded to the email
17 invite, they answered screening questions about their age, smoking status and types of
18 tobacco products used, with those that did not meet the inclusion criteria (factory-made
19 cigarette smokers aged 16-34 years) excluded.
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Those eligible for inclusion were presented with an information page explaining the
34 study aim (to explore what young adult smokers thought about cigarettes and pack inserts),
35 and relevant ethical information (their right to withdraw at any time, assurances of
36 confidentiality and anonymity, and contact details if they had any concerns or would like to
37 request a copy of the published findings). They were then presented with a consent page, with
38 consent required for participation. Survey questions were presented in the same order for all
39 participants, except the questions exploring perceptions of the three cigarettes (standard
40 cigarette, warning cigarette, green cigarette), where the ordering was randomised; the
41 ordering of the presentation of the three cigarettes (shown in Figure 1) was also randomised.
42 There was no missing data as participants could only proceed to the next question if they had
43 provided an answer to the previous question.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 Figure 1 here
6
7
8

9
10 Prior to the questions on inserts, participants were shown an image of a cigarette pack with an
11 insert shown in the front of the pack – as they typically appear in packs – alongside the text
12 ‘We have some questions on pack inserts, which can sometimes be found inside packs (see
13 image for example)’. For each question about inserts, participants were shown the question
14 and an image of one insert. Four different inserts were used in total, as shown in Figure 2,
15 with these chosen from the eight used in Canada as they were considered most relevant to our
16 sample. The words ‘Health Canada’ were removed from the bottom of each insert to make
17 them more relevant for participants in the UK. The median time for survey completion was 9
18 minutes 28 seconds. Participants received a nominal incentive (50 pence) for participation, as
19 is common for online panels. The study received ethical approval from the School of Health
20 Sciences Ethics Committee at the University of Stirling.
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 Figure 2 here
36
37
38

39 **Patient and public involvement**

40
41 There was no patient or public involvement in the development, design or conduct of this
42 study.
43
44
45

46 **Measures**

47
48
49
50
51
52 *Inserts: Salience and information provision*
53
54
55
56
57

1
2
3 Participants were asked ‘If this type of insert was in your cigarette pack, do you think that
4 you would read it?’ and ‘If this type of insert was in your cigarette pack, do you think that
5 you would read it if you were interested in quitting?’ They were also asked ‘Do you think
6 that inserts would be a good way to provide information to smokers about quitting?’⁵
7
8
9
10
11 Response options for each were ‘Yes’, ‘No’ and ‘Not sure’.
12
13

14 15 16 *Inserts: Cessation*

17
18 Three questions assessed to what extent participants agreed or disagreed that inserts would
19 make them think about quitting, and help them quit: ‘Do you agree or disagree that having
20 these types of inserts in every cigarette pack would make you think more about quitting?’,
21 ‘Do you agree or disagree that having these types of inserts in every cigarette pack might help
22 you if you decided to quit?’, and ‘Do you agree or disagree that having these types of inserts
23 inside every cigarette pack would be an effective way of helping smokers who want to quit?’⁶
24
25 Response options for each were ‘Strongly disagree’, ‘Disagree’, ‘Neither agree nor disagree’,
26 ‘Agree’, ‘Strongly agree’ and ‘Don’t know’.
27
28
29
30
31
32
33
34
35
36
37

38 *Inserts: Support*

39
40 A five-point semantic scale assessed support, with anchors ‘All cigarette packs should have
41 inserts like this in them-No cigarette packs should have inserts like this in them’.
42
43
44
45

46 *Cigarette design: Appeal, harm and trial*

47
48 Seven-point semantic scales assessed appeal, harm and likely trial. Appeal was assessed via
49 four scales, with anchors ‘Attractive-Unattractive’, ‘Stylish-Not stylish’, ‘Not nice to be seen
50 with-Nice to be seen with’ and ‘Not appealing to people my age-Appealing to people my
51 age’. Harm was assessed via two scales, with anchors ‘Looks harmful to health-Does not look
52
53
54
55
56
57
58
59
60

1
2
3 harmful to health' and 'Makes me think about the dangers of smoking-Does not make me
4 think about the dangers of smoking'. Likely trial was assessed via two scales, 'If a friend
5 offered you each of these cigarettes, how likely would you be to try them?' and 'If someone
6 your age who had never smoked before was going to try a cigarette, how likely do you think
7 they would be to try each of these cigarettes?' Both scales assessing trial ranged from 'Not at
8 all likely' to 'Very likely'.

15 *Sociodemographic characteristics*

16 Age, gender, ethnicity, educational attainment and economic status (based on chief income
17 earner) were obtained. A count procedure was used to create a variable for low
18 socioeconomic status (SES): low education (General Certificate of Secondary Education:
19 GCSE or below) and/or low economic status (routine or manual occupation, long-term
20 unemployed or long-term sick or disabled).

30 *Smoking behaviour*

31 Smoking status was assessed with 'Which of these best describes you?' with response
32 options: 'I have never smoked', 'I used to smoke, but don't now', 'I smoke, but not every
33 day', and 'I smoke every day'. Type of products used was assessed with 'What type(s) of
34 tobacco products do you smoke?' with response options: 'Only factory-made (packet)
35 cigarettes', 'Factory-made and roll-your-own cigarettes', 'Factory-made cigarettes and other
36 tobacco products (e.g. cigars, shisha, etc)', 'Only roll-your-own cigarettes' and 'Only other
37 tobacco products (e.g. cigars, shisha, etc)'. The Heaviness of Smoking Index (HSI)²³ was
38 used as a measure of dependence, based on daily consumption and time to first cigarette.
39
40
41
42
43
44
45
46
47
48
49
50

51 *Quitting and self-efficacy*

Participants were asked 'Have you ever made an attempt to quit smoking that lasted at least 24 hours?'²⁴ (Yes within the last six months, Yes more than six months ago, I have never tried to quit for more than 24 hours). They were also asked 'How likely are you to try to quit smoking within the next six months?'²⁵ (Not at all, A little, Moderately, Very, Extremely, Don't know), with those responding 'Not at all', 'A little', 'Moderately' or 'Don't know' classified as 'Unlikely to make a quit attempt in the next six months'. To measure quitting self-efficacy, participants were asked 'If you decided to quit smoking in the next six months, how sure are you that you would succeed?'²⁶ (Not at all, A little, Moderately, Very, Extremely, Don't know). Those who responded to the likelihood of quitting question with 'Very' or 'Extremely' and to the quitting efficacy question with 'Not at all', 'A little', 'Moderately' or 'Don't know' were classified as 'unlikely to make a successful quit attempt in the next six months'. Those who responded 'Very' or 'Extremely' to both questions were classified as 'likely to make a successful quit attempt in the next six months'.

Analysis

Data was analysed using Microsoft office Excel 2013, SPSS v22 and v23 and MLWin v2.33.²⁷ The insert variables were dichotomised into yes/agreement and no/disagreement/neutral/not sure/don't know. The dichotomised insert variables were the outcomes of the logistic regression models. The independent variables were gender, age, education, ethnicity, dependence (tertiles of HSI), tobacco product(s) smoked, previous quit attempt lasting at least 24 hours, and likely efficacy of a quit attempt in the next six months. Percentages in agreement were calculated. Age, gender and education (as a measure of SES) were entered into all models to account for any sampling inadequacies. Other variables were entered where $p < 0.10$ in chi square tests. The models were assessed for multicollinearity via comparison of standard errors²⁸ and none was found.

1
2
3 The cigarette variables were assessed using seven-point semantic scales, with
4 percentages calculated for those indicating one of the three points nearest the undesirable
5 anchor (e.g. unattractive, not nice to be seen with, looks harmful to health). Differences
6 between the three cigarettes were tested using Cochran's Q and pairwise comparisons. A
7 factor analysis of the eight perception variables, collated for all three cigarettes, was
8 undertaken, with checks indicating that the data was suitable for factor analysis (Kaiser
9 Meyer Olkin=0.845, Bartlett's test of sphericity (approx. chi-square 18062.842, df=276,
10 p<0.001), with no correlations between the variables >0.9). The extraction method used was
11 Principal Axis Factoring and the criteria for extraction was eigenvalues>1. All eight variables
12 loaded on a single factor with factor loadings that were >0.5. High factor scores indicated that
13 a cigarette was desirable and low scores that it was undesirable. Visual inspection and the
14 Kolmogorov-Smirnov test indicated that the factor was non-normal (because responses for
15 the dissuasive cigarettes indicated they were undesirable generally) and attempts to normalise
16 it using normit rankit methods failed. Thus the factor was divided into tertiles and the tertile
17 indicating undesirable factor scores was compared with the other two tertiles. This was the
18 outcome variable in regression analysis.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Multilevel logistic regression modelling, with second order PQL estimation,²⁹ was undertaken with cigarette type (at level one) clustered with individual participants (at level two). All models included cigarette type as a fixed effect, where the standard cigarette was compared with the warning cigarette and green cigarette. Other fixed effects at the individual (participant) level were sociodemographic and smoking-related characteristics. This main effects model tested which characteristics were associated with perceiving cigarettes as desirable. In order to understand which characteristics differentiated the desirability of the three types of cigarettes, interactions between cigarette type and each significant characteristic were tested. Only one interaction was found, between cigarette type and SES.

1
2
3 The interacting variables (cigarette type and SES) were substituted by a cross classified
4 variable which merged cigarette type and SES. This cross classified variable was split into
5 six categories: low SES standard cigarette, low SES warning cigarette, low SES green
6 cigarette, not low SES standard cigarette, not low SES warning cigarette, not low SES green
7 cigarette. To understand the interaction several models were run with the reference category
8 of the cross classified variable different each time.^{30,31}
9
10
11
12
13
14
15
16
17

18 RESULTS

19 Perceptions of inserts

20 Half the sample indicated that they would read inserts, with approximately three-fifths
21 indicating that they would read them if interested in quitting (60%), and that they would be a
22 good way to provide information about quitting (61%). Just over half strongly agreed/agreed
23 that inserts may make them think more about quitting (53%), help them if they decided to
24 quit (52%), that they are an effective way of encouraging smokers to quit (53%), and that all
25 cigarette packs should have inserts (55%), see Table 2.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Table 2 here

45 Sociodemographic differences in perceptions of inserts

46 Women were more likely than men to indicate that they would read inserts (aOR=1.24;
47 95%CI 1.02-1.50), and 25-34 year olds less likely than 16-19 year olds to think that they
48 were a good way of providing information about quitting (aOR=0.76; 95%CI 0.60-0.98).
49 Compared with white British participants, white non-British (aOR=0.70; 95%CI 0.50-0.98)
50 and Asian (aOR=0.67; 95%CI 0.49-0.92) participants were less likely to suggest that they
51
52
53
54
55
56
57
58
59
60

would read inserts if trying to quit, white non-British (aOR=0.58; 95%CI 0.41-0.81) and Black (aOR=0.61; 95%CI 0.38-0.98) participants were less likely to indicate that inserts would make them think about quitting, and white non-British (aOR=0.62; 95%CI 0.44-0.87) and Asian (aOR=0.70; 95%CI 0.51-0.96) participants were less likely to support having inserts in all packs, see Table 3a.

Smoking-related differences

Compared to exclusive factory-made cigarette smokers, those who also smoked roll-your-own cigarettes were more likely to indicate they would read inserts (aOR=1.35; 95%CI 1.09-1.66), read them if trying to quit (aOR=1.61; 95%CI 1.30-2.00), that they would make them think about quitting (aOR=1.31; 95%CI 1.06-1.62), help them if they decided to quit (aOR=1.31; 95%CI 1.06-1.61), and that they would be an effective way of encouraging smokers to quit (aOR=1.27; 95%CI 1.03-1.56). Compared to exclusive factory-made cigarette smokers, those who also smoked other tobacco products (e.g. cigars, shisha) were more likely to indicate they would read inserts if trying to quit (aOR=1.39; 95%CI 1.04-1.86) and that inserts might help them if they decided to quit (aOR=1.34; 95%CI 1.01-1.78).

Participants who had made a quit attempt more than six months ago (aOR=1.30; 95%CI 1.00-1.69), or within the last six months (aOR=1.67; 95%CI 1.29-2.15), were more likely to indicate that they would read inserts than those who had never made a quit attempt. Those who had made a quit attempt in the last six months were also more likely than those who had never made a quit attempt to indicate that inserts were a good way to provide information about quitting (aOR=1.54; 95%CI 1.20-1.98), that they would read them if trying to quit (aOR=1.51; 95%CI 1.17-1.94), make them think about quitting (aOR=1.46; 95%CI 1.14-1.88), help them if they decided to quit (aOR=1.35; 95%CI 1.05-1.73), and that they would be an effective way of encouraging smokers to quit (aOR=1.33; 95%CI 1.04-1.71).

1
2
3 Compared to those likely to make a successful quit attempt in the next six months,
4 those unlikely to make a quit attempt in the next six months were less likely to indicate that
5 they would read inserts (aOR=0.58; 95%CI 0.44-0.75), read them if trying to quit
6 (aOR=0.74; 95%CI 0.55-0.99), that they would make them think about quitting (aOR 0.59
7 (0.45 to 0.78), help them if they decided to quit (aOR=0.51; 95%CI 0.38-0.67), that they
8 would be effective for smokers if they decided to quit (aOR=0.55; 95%CI 0.41-0.73), or
9 support them (aOR=0.56; 95%CI 0.42-0.74). Compared to those likely to make a successful
10 quit attempt in the next six months, those unlikely to make a successful quit attempt in the
11 next six months were more likely to read inserts if trying to quit (aOR=1.43; 95%CI 1.00-
12 2.06), thought that they were a good way to provide information to smokers about quitting
13 (aOR=1.46; 95%CI 1.02-2.08), and support them (aOR=1.43; 95%CI 1.00-2.04), see Table
14 3b.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31 Table 3 here
32
33
34

35 **Perceptions of cigarette design**

36
37 With respect to harm, participants were less likely to think that the standard cigarette (SC)
38 looked harmful than the warning cigarette (WC) or green cigarette (GC) ($p<0.001$), and less
39 likely to think that the SC made them think more about the dangers of smoking than the WC
40 or GC ($p<0.001$). Participants were also less likely to indicate that the GC would make them
41 think of the dangers of smoking than the WC ($p=0.01$). In terms of appeal, participants were
42 more likely to consider the SC attractive, and stylish, than the WC or GC (both $p<0.001$). The
43 SC was also considered to be nicer to be seen with, and more appealing to people their age,
44 than the WC or GC (both $p<0.001$). In terms of trial, 79.4% indicated that they would try a
45 SC if offered by a friend (35.7% WC, 21.5% GC), and 70.1% indicated that a never smoker
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 their age would be most likely to try a SC (21.1% WC, 16.5% GC) (both $p < 0.001$), see Table
4
5 4.

6
7
8
9 Table 4 here

10 11 12 13 **Perceptions of cigarette desirability**

14
15 Main effects multivariable logistic regression modelling suggested that in comparison to the
16 SC, the WC (aOR=17.71; 95%CI 13.75-22.80) and GC (aOR=30.88; 95%CI 23.98-39.76)
17
18 were much more likely to be perceived as undesirable (i.e. less appealing, more harmful, less
19
20 likely to be tried). The model also indicated which smokers were more likely to rate the
21
22 cigarettes as undesirable: women were more likely than men (aOR=1.30; 95%CI 1.10-1.54),
23
24 and low SES more likely than those not low SES (aOR=1.26; 95%CI 1.06-1.50), to consider
25
26 all three cigarettes undesirable. Compared to exclusive factory-made cigarette smokers, those
27
28 who also smoked roll-your-own cigarettes (aOR=0.78; 95%CI 0.65-0.93) or other tobacco
29
30 products (aOR=0.73; 95%CI 0.56-0.93) were less likely to consider all three cigarettes
31
32 undesirable. Those not likely to make a quit attempt in the next six months were less likely
33
34 than those likely to make a quit attempt in the next six months (aOR=0.62; 95%CI 0.49-0.78)
35
36 to consider all three cigarettes undesirable.

37
38
39
40
41 Only one significant interaction, between cigarette type and SES, was found ($p < 0.05$).
42
43 Both SES groups perceived the WC significantly more undesirable than the SC, and the GC
44
45 significantly more undesirable than the WC. Low SES were significantly more likely than
46
47 those not low SES to perceive the SC (aOR=17.71; 95%CI 13.75-22.80) and GC
48
49 (aOR=30.88; 95%CI 23.98-39.76) as undesirable; there was no difference for the WC
50
51 (aOR=0.99; 95%CI 0.78-1.25).
52
53
54
55
56
57
58
59
60

DISCUSSION

Our findings suggest that inserts highlighting the benefits of quitting or providing tips on how to do so may have the potential to encourage cessation, and dissuasive cigarettes may help to reduce the desirability of smoking. Just as tobacco companies have used inserts and cigarette design to create interest in their products, our study suggests that greater attention to how these could be used to promote cessation appears warranted.

Health messages need to capture attention to be effective.³² In this regard, at least half our sample indicated that they would read inserts (50%) and read them if interested in quitting (60%). In Canada, an observational study found that approximately a quarter of smokers reported reading them at least once within the last month,¹⁰ increasing to about one-third of smokers over two years of follow-up.¹¹ As in our study, smokers in Canada who had read/would read the inserts were more likely to be female, intend to quit or had recently tried to quit; in our study, they were also more likely to be white British, have moderate dependence, and use factory-made cigarettes and other tobacco products. Future research could explore why dual users (smokers of factory-made cigarettes and other tobacco products) were more likely to indicate that they would read inserts, but as inserts are typically only found in cigarette packs then for those who use other tobacco products they may be seen as more of a novelty and therefore more likely to capture attention.

Approximately three-fifths (61%) of smokers in our study thought that inserts were a good way to provide information about quitting to smokers, with only 25% disagreeing. In comparison, an earlier study in Canada, commissioned by Health Canada, found that 48% of smokers indicated that messaging on inserts was a good way to provide information to smokers, with 47% disagreeing.⁵ Just over half our sample agreed/strongly agreed that inserts may make them think more about quitting, help them if they decided to quit, and that they are

1
2
3 an effective way of encouraging smokers to quit, whereas in New Zealand only 34% of
4 smokers and recent quitters agreed/strongly agreed that inserts would be an effective way of
5 encouraging reduced consumption or quitting.⁶ There may be various reasons for the
6 differences between our findings and earlier research. For instance, when this earlier research
7 was conducted cigarette packs displayed text-only health warnings and it may be that having
8 pictorial warnings on packs, as is required in Scotland, may prompt smokers to look for
9 information on how to quit and the benefits of doing so. Insert design is also likely to be
10 relevant. Whereas the inserts used in earlier research were limited to text, the inserts used in
11 this study (which have been used in Canada since 2012) included coloured graphics, which is
12 typical of promotional inserts used by tobacco companies and likely enhanced their impact.
13 This would be consistent with the health communications and warnings literature, which
14 demonstrates the importance of supporting text with pictorials.^{32,33} Future research exploring
15 insert design (e.g. use of imagery, inclusion of cessation resource information, length and
16 framing of messages, etc) would be of value.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 More than half our sample supported the inclusion of inserts promoting cessation
34 inside every cigarette pack, with only a fifth opposing this. Within the European Union, the
35 recent Tobacco Products Directive (TPD)³⁴ does not require tobacco companies to include
36 health communication inserts in packs, but allows member states to introduce measures
37 beyond those specified. Among governmental representatives that responded to the
38 consultation on the revision of the TPD there was strong support for improving consumer
39 information via mandatory pictorial warnings, with those supportive arguing that additional
40 information, such as pack inserts, would help to deliver more accurate health information.³⁵ If
41 there is support for inserts among governmental representatives, and little opposition among
42 smokers (the group most likely to be resistant), they are clearly a viable option for regulators.
43
44
45
46
47
48
49
50
51
52
53

54 Tobacco industry journals describe the cigarette as an increasingly important
55
56
57
58
59
60

1
2
3 advertising medium for tobacco companies.¹² However, until recently, the public health focus
4
5 has been on the potential of regulating the contents of cigarettes to reduce palatability or
6
7 addictiveness,³⁶ with little consideration of the possibility of regulating the appearance of
8
9 cigarettes to reduce its importance as a promotional tool. We found that the two dissuasive
10
11 cigarettes were perceived as significantly more harmful and less appealing than the standard
12
13 cigarette, and less likely to encourage trial. The harm, appeal and trial items loaded onto a
14
15 single ‘undesirability’ factor, with the dissuasive cigarettes considered much more
16
17 undesirable than the standard cigarette. The findings are consistent with earlier research,
18
19 where cigarettes with the warning ‘Smoking kills’ were considered a constant reminder of the
20
21 associated harms and, partly due to the perceived discomfort of being observed by others
22
23 smoking a cigarette displaying this message, unappealing for smokers.^{8,16,17,18} Previous
24
25 studies have also found unattractively coloured cigarettes to be perceived as more harmful
26
27 than other cigarettes and also repellent, being a cigarette that young people did not think that
28
29 others their age would use.^{15,16,37,38} As with the inserts, the dissuasive cigarettes (and also the
30
31 standard cigarette) were considered more desirable among dual users than exclusive factory-
32
33 made cigarette smokers; again it is not clear why this was the case but further research with
34
35 dual users, or indeed those also using vaping devices (not assessed in this study), would be
36
37 fruitful.
38
39
40

41
42 In terms of limitations, the cross-sectional design did not allow us to assess causality;
43
44 that inserts and dissuasive cigarettes are not available on the UK market prevents more robust
45
46 study designs such as longitudinal studies. Another potential limitation concerns the novelty
47
48 of the stimuli, which may have influenced responses, and forced exposure to the stimuli. In
49
50 addition, we only used four inserts, rather than the full set of eight used in Canada, which
51
52 includes inserts that be less relevant to our sample. While online surveys have been used for
53
54 previous research exploring cigarette packaging, inserts and dissuasive cigarettes,³⁹⁻⁴² and are
55
56
57
58
59
60

1
2
3 a suitable survey mode for young adults, the use of an online panel and self-selection limits
4 the representativeness of our sample. In addition, the use of semantic differential scales can
5 be criticised because answers can be subject to various response biases, although we
6 attempted to diminish these through varying scale item direction and through our multivariate
7 modelling methodology.
8
9
10
11
12

13
14 It was argued, over two decades ago, that to offer greater protection to consumers
15 cigarettes should come in plain packs with messaging on both the pack exterior and interior.⁴³
16 This idea is a step closer in the UK, although there will still be no messaging on the pack
17 interior. That more than half of the participants in this study suggested that inserts may help
18 to promote cessation suggests that their inclusion in packs may be a meaningful supplement
19 to the on-pack warnings. Our findings suggest however that to offer the greatest protection to
20 consumers, it may be beneficial to supplement plain packaging and inserts with cigarettes
21 designed to be dissuasive. Unattractively coloured cigarettes would complement the
22 unattractively coloured packs, just as warnings on the cigarette would extend the warnings on
23 the cigarette pack. Both options are clearly viable.
24
25
26
27
28
29
30
31
32
33
34

35
36
37 **Contributors** CM designed the data collection tool and drafted and revised the paper. RH
38 analysed the data and drafted the Analysis and Results. JT and GR helped design the data
39 collection tool and commented on the paper. All authors read and approved the final
40 manuscript.
41
42
43
44

45
46 **Funding** This work was supported by Health Scotland. GR, who works for Health Scotland,
47 provided feedback on the survey and paper, but was not involved in the collection, analysis
48 and interpretation of the data.
49
50
51

52 **Competing interests** GR works for Health Scotland, who funded this study.
53
54
55
56
57

1
2
3 **Ethics approval** The study obtained ethics approval from the School of Health Sciences
4
5 Ethics Committee at the University of Stirling. Participants provided informed consent before
6
7 participating.
8

9 **Provenance and peer review** Not commissioned; externally peer reviewed.
10

11 **Data sharing statement** No additional data are available.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

REFERENCES

1. Canadian Cancer Society. *Cigarette package health warnings. International status report (Fifth edition)*; 2016. www.tobaccolabels.ca/wp/wp-content/uploads/2016/11/Cigarette-Package-Health-Warnings-International-Status-Report-English-CCS-Oct-2016.pdf (accessed 10 Sept 2017).
2. Hammond D. Health warning messages on tobacco products: a review. *Tob Control* 2011;20:327–37.
3. Mahood G. *Canada's tobacco package label or warning system: "telling the truth" about tobacco product risks*. Geneva: WHO; 2010.
4. Tandemar Research Inc. *Tobacco health warning messages, inserts and toxic constituents information study - Final Report*. Toronto: Tandemar Research Inc, 1992.
5. Environics. *Health warning messages on the flip/side and inserts of cigarette packaging. A survey of smokers. A report prepared for Health Canada*; 2000.
6. BRC Marketing and Social Research. *Smoking health warnings study*. Wellington: Ministry of Health; 2004. www.tobaccolabels.ca/wp/wp-content/uploads/2013/12/NZ-2004-Effectiveness-of-Different-Health-Warnings-in-Helping-People-Consider-Their-Smoking-Related-Behaviour-Government-Report.pdf (accessed 12 Aug 2017).
7. Gallopel-Morvan K, Moodie C, Hammond D, *et al*. Consumer understanding of cigarette emission labelling. *Eur J Pub Health* 2011;21:373-5.
8. Moodie C. Novel ways of using tobacco packaging to communicate health messages: Interviews with packaging and marketing experts. *Addict Res Theory* 2016;24:54-61.
9. Moodie C. Adult smokers' perceptions of cigarette pack inserts promoting cessation: A focus group study. *Tob Control* 2018;27:72-7.

10. Thrasher JT, Osman A, Abad EN, *et al.* The innovative use of cigarette package inserts to supplement pictorial health warnings: An evaluation of the Canadian policy. *Nicot Tob Res* 2015;17:870-5.
11. Thrasher JF, Swayampakala K, Cummings KM, *et al.* Cigarette package inserts can promote efficacy beliefs and sustained smoking cessation attempts: A longitudinal assessment of an innovative policy in Canada. *Prev Med* 2016;88:59-65.
12. Rossell S. Ready to roll. *Tob Report* 2017;2:44-5.
13. Smith KC, Washington C, Welding K, *et al.* Cigarette stick as valuable communicative real estate: a content analysis of cigarettes from 14 low-income and middle-income countries. *Tob Control* 2017;26:604-7.
14. Hedley D. Tobacco wars: A new hope - the changing future. *Tob J Int* 2015;3:31-4.
15. Hoek J, Robertson C. How do young adult female smokers interpret dissuasive cigarettes? *J Social Marketing* 2015;5:21-39.
16. Hoek J, Gendall P, Eckert C, *et al.* Dissuasive cigarette sticks: the next step in standardised ('plain') packaging? *Tob Control* 2016;15:699-705.
17. Moodie C, Purves R, McKell J, *et al.* Novel means of using cigarette packaging and cigarettes to communicate health risk and cessation messages: A qualitative study. *Internat J Mental Health Addiction* 2015;13:333-44.
18. Moodie C, Mackintosh AM, Gallopel-Morvan K, *et al.* Adolescents' perceptions of health warnings on cigarettes. *Nicot Tob Res* 2017;29:1232-7.
19. Hassan L, Shui E. No place to hide: two pilot studies assessing the effectiveness of adding a health warning to the cigarette cigarette. *Tob Control* 2015;24:e3-5.
20. ONS. *Internet users in the UK: 2017*.
www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2017 (accessed 15 Sept 2017).

21. Hoek J, Gendall P, Eckert C, *et al.* Effects of brand variants on smokers' choice behaviours and risk perceptions. *Tob Control* 2016;25:160-5.
22. Moodie C, Sinclair L, Mackintosh AM, *et al.* How tobacco companies are perceived within the UK: An online panel. *Nicot Tob Res* 2016;18:1766-72.
23. Kozlowski LT, Porter CQ, Orleans T, *et al.* Predicting smoking cessation with self-reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug Alcohol Depend* 1994;34:211-6.
24. Etter JF, Sutton S. Assessing 'stage of change' in current and former smokers Trying to stop smoking: effects of perceived addiction, attributions for failure, and expectancy of success. *Addiction* 2002;97:1171-82.
25. Hammond D, Fong GT, McDonald PW, *et al.* Impact of the graphic Canadian warning labels on adult smoking behaviour. *Tob Control* 2003;12:391-5.
26. IARC. *IARC Handbooks of Cancer Prevention, Tobacco Control, Vol. 12: Methods for Evaluating Tobacco Control Policies*. Lyon, France: International Agency for Research on Cancer; 2008.
27. Rasbash J, Browne W, Healy M, *et al.* *A user's guide to MLwiN, version 2.3*. Bristol: Centre for multilevel modelling, University of Bristol; 2015.
28. Salmond C. *Fitting complex models using health survey data*; 2006.
www.otago.ac.nz/wellington/otago020178.pdf (accessed 15 Sept 2017).
29. Goldstein H, Rasbash J. Improved approximations for multilevel models with binary responses. *J Royal Statistical Society. Series A (Statistics in Society)* 1996;159:505-13.
30. Blakely T, Woodward A, Salmond C. Anonymous linkage of New Zealand mortality and Census data. *Australian NZ J Pub Health* 2000;24:92-5

- 1
2
3 31. Hiscock R, Murray S, Brose LS, *et al.* Behavioural therapy for smoking cessation:
4 The effectiveness of different intervention types for disadvantaged and affluent
5 smokers. *Addictive Behaviors* 2013;38:2787-96.
6
7
8
9 32. Wogalter MS, Conzola VC, Smith-Jackson TL. Research-based guidelines for
10 warning design and evaluation. *Applied Ergonomics* 2002;33:219-30.
11
12
13 33. Houts PD, Doak CC, Doak LG, *et al.* The role of pictures in improving health
14 communication: A review of research on attention, comprehension, recall, and
15 adherence. *Patient Educ Couns* 2006;61:173-90.
16
17
18
19 34. European Commission. Directive 2014/40/EU of the European parliament and of the
20 Council of 3 April 2014 on the approximation of the laws, regulations and
21 administrative provisions of the Member States concerning the manufacture,
22 presentation and sale of tobacco and related products and repealing Directive
23 2001/37/EC. *Official Journal of the European Union* 2014;L127:1-38.
24
25
26
27
28
29 35. European Commission. *Report on the public consultation on the possible revision of*
30 *the Tobacco Products Directive (2001/37/EC)*. Health and Consumers Directorate-
31 General – Directorate D – Health systems and products D4 – Substances of human
32 origin and tobacco control. 2011.
33 http://ec.europa.eu/health/tobacco/docs/consultation_report_en.pdf (accessed 10 Sept
34 2017).
35
36
37
38
39
40
41
42
43 36. Warner K. The national and international regulatory environment in tobacco control.
44 *Pub Health Res Practice* 2015;25:e2531527.
45
46
47
48 37. Ford A, Moodie C, Mackintosh AM, *et al.* Adolescent perceptions of cigarette
49 appearance. *Eur J Pub Health* 2014;24:464-8.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 38. Moodie C, Ford A, Mackintosh AM, *et al.* Are all cigarettes just the same? Female's
4 perceptions of slim, coloured, aromatised and capsule cigarettes. *Health Educ Res*
5 2015;30:1-12.
6
7
8
9 39. Doxey J, Hammond D. Deadly in pink: The impact of female-oriented cigarette
10 packaging among young women. *Tob Control* 2011;20:353-60.
11
12
13 40. White CM, Hammond D, Thrasher JF, *et al.* The potential impact of plain packaging
14 of cigarette products among Brazilian young women: an experimental study. *BMC*
15 *Public Health* 2012;12:737.
16
17
18 41. Kotnowski K, Fong GT, Gallopel-Morvan K, *et al.* The impact of cigarette packaging
19 design among young females in Canada: findings from a discrete choice experiment.
20 *Nicot Tob Res* 2016;18:1348-56.
21
22
23 42. Hoek J, Gendall P, Maubach N, *et al.* Strong public support for plain packaging of
24 tobacco products. *Austr N Z J Pub Health* 2012;36:405-7.
25
26
27 43. Mahood G. Canada tobacco package warning system. *Tob Control* 1995;4:10-4.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Sample and smoking-related characteristics

<i>Characteristic</i>	<i>N</i>	<i>%</i>
Total	1766	100.0
Age group		
16-19	413	23.4
20-24	401	22.7
25-34	952	53.9
Gender		
Male	888	50.3
Female	878	49.7
Educational qualifications		
Other qualifications	1357	76.8
None or GCSE	409	23.2
Economic status		
Other status	1350	76.4
Routine or manual occupation, unemployed or long term sick	416	23.6
Socioeconomic status (SES)		
No indicators of low SES	1114	63.1
Low education and/or low SES	652	36.9
Ethnicity		
White British	1264	71.6
White non-British	162	9.2
Black (including mixed black and white)	79	4.5
Asian (including mixed Asian and white)	196	11.1
Other or not declared	65	3.7
Location		
England	1550	87.8
Scotland	109	6.2
Wales	73	4.1
Northern Ireland	34	1.9
Tobacco products used		
Only factory-made (packet) cigarettes	813	46.0
Factory-made and roll-your-own cigarettes	681	38.6
Factory-made cigarettes and other products (e.g. cigars, shisha)	272	15.4
Cigarettes per day		
10 or less	1272	72.0
11-20	433	24.5
21-30	46	2.6
31 or more	15	0.8
Time to first cigarette		
Within 5 minutes	263	14.9
6 to 30 minutes	570	32.3
31 to 60 minutes	315	17.8

<i>Characteristic</i>	<i>N</i>	<i>%</i>
After 60 minutes	618	35.0
Heaviness of Smoking Index (HSI)		
0 little dependence	601	34.0
1	257	14.6
2	418	23.7
3	293	16.6
4	156	8.8
5	28	1.6
6 high dependence	13	0.7
Dependence (Tertiles of HSI)		
Low-dependence	601	34.0
Mid-dependence	675	38.2
High-dependence	490	27.7
Made an attempt to quit smoking that lasted at least 24 hours?		
Yes, within the last six months	788	44.6
Yes, more than six months ago	552	31.3
No, I have never tried to quit smoking for more than 24 hours	426	24.1
How likely are you to try to quit smoking within the next six months?		
Not at all	198	11.2
A little	382	21.6
Moderately	508	28.8
Very	308	17.4
Extremely	272	15.4
Don't know	98	5.5
If you decided to quit smoking in the next six months, how sure are you that you would succeed?		
Not at all	147	8.3
A little	346	19.6
Moderately	612	34.7
Very	297	16.8
Extremely	241	13.6
Don't know	123	7.0
Quit approach		
Moderately or less likely to make quit attempt in next six months (unlikely to make a quit attempt in the next six months)	1186	67.2
Very or extremely likely to attempt but moderately or less likely to succeed (unlikely to make a successful quit attempt in the next six months)	304	17.2
Very or extremely likely to attempt and very or extremely likely to succeed (likely to make a successful quit attempt in the next six months)	276	15.6

Table 2: Perceptions of whether inserts would be read, are a good way to provide information, whether they would help smokers to think about quitting or quit, and support for them

	Yes %	No %	Not sure %
Would they be read	50	37	13
Would they be read if interested in quitting	60	25	15
Good way to provide information about quitting	61	25	14

	Agree %	Disagree %	Neither / Don't know %
Make you think more about quitting	53	18	29
Might help you if you decided to quit	52	19	29
Effective way of encouraging smokers to quit	53	17	30
All packs should have inserts	55	20	25

Table 3a: Logistic regression models exploring perceptions of inserts by sociodemographic characteristics (gender, age, education, ethnicity)^{1,2}

(n=1766)	Would read insert	Would read insert if trying to quit	Inserts make you think about quitting	Inserts might help you quit	Inserts a good way of providing information about quitting	Inserts are an effective way of encouraging smokers to quit	All packs should have inserts
Gender							
Male	1	1	1	1	1	1	1
Female	1.24 (1.02 to 1.50)	1.11 (0.91 to 1.35)	0.98 (0.81 to 1.19)	0.95 (0.79 to 1.15)	1.13 (0.93 to 1.37)	0.88 (0.73 to 1.07)	1.20 (0.99 to 1.46)
Age							
16-19	1	1	1	1	1	1	1
20-24	1.16 (0.87 to 1.54)	0.88 (0.66 to 1.18)	1.18 (0.89 to 1.56)	1.19 (0.89 to 1.58)	0.87 (0.65 to 1.16)	0.97 (0.73 to 1.28)	0.96 (0.72 to 1.29)
25-34	1.25 (0.97 to 1.60)	0.83 (0.65 to 1.07)	0.99 (0.78 to 1.26)	1.18 (0.92 to 1.50)	0.76 (0.60 to 0.98)	0.88 (0.69 to 1.12)	0.84 (0.65 to 1.07)
Education							
GCSEs (or equivalent) or none	1	1	1	1	1	1	1
More than GCSEs (or equivalent)	1.25 (0.99 to 1.58)	1.12 (0.89 to 1.42)	1.22 (0.97 to 1.54)	1.21 (0.97 to 1.52)	1.12 (0.89 to 1.40)	1.19 (0.95 to 1.50)	1.10 (0.87 to 1.40)
Ethnicity							
White British		1	1				1
White but not British		0.70 (0.50 to 0.98)	0.58 (0.41 to 0.81)				0.62 (0.44 to 0.87)
Black (inc mixed black & white)		0.92 (0.57 to 1.49)	0.61 (0.38 to 0.98)				0.99 (0.62 to 1.59)
Asian (inc mixed Asian & white)		0.67 (0.49 to 0.92)	1.19 (0.87 to 1.63)				0.70 (0.51 to 0.96)
other or not declared		0.84 (0.50 to 1.42)	1.06 (0.64 to 1.78)				1.08 (0.64 to 1.81)

¹Note smoking related characteristics (described in table 3b) were also entered into each model

²Blank cells indicate no significant relationship in bivariate analysis

Table 3b: Logistic regression models exploring perceptions of inserts by smoking related characteristics (dependence, tobacco products smoked, quit attempts, self-efficacy to quit)^{1,2}

(n=1766)	Would read insert	Would read insert if trying to quit	Inserts make you think about quitting	Inserts might help you quit	Inserts a good way of providing information about quitting	Inserts are an effective way of encouraging smokers to quit	All packs should have inserts
Dependence (tertiles of HSI)							
Lower dependence	1						1
Mid dependence	1.39 (1.11 to 1.76)						1.02 (0.80 to 1.29)
Higher dependence	1.22 (0.94 to 1.59)						0.86 (0.66 to 1.12)
Tobacco products smoked							
Only factory-made	1	1	1	1		1	
Factory-made and roll-your-own	1.35 (1.09 to 1.66)	1.61 (1.30 to 2.00)	1.31 (1.06 to 1.62)	1.31 (1.06 to 1.61)		1.27 (1.03 to 1.56)	
Factory-made cigarettes and other	1.20 (0.90 to 1.59)	1.39 (1.04 to 1.86)	1.22 (0.92 to 1.63)	1.34 (1.01 to 1.78)		1.20 (0.91 to 1.60)	
Quit attempt lasting at least 24 hours							
No	1	1	1	1	1	1	1
Yes, more than six months ago	1.30 (1.00 to 1.69)	1.12 (0.86 to 1.45)	1.20 (0.93 to 1.56)	1.05 (0.81 to 1.36)	1.16 (0.90 to 1.50)	1.07 (0.82 to 1.38)	0.78 (0.60 to 1.01)
Yes within the last six months	1.67 (1.29 to 2.15)	1.51 (1.17 to 1.94)	1.46 (1.14 to 1.88)	1.35 (1.05 to 1.73)	1.54 (1.20 to 1.98)	1.33 (1.04 to 1.71)	1.06 (0.82 to 1.37)
Efficacy of quit attempt in next 6 months							
Likely to quit	1	1	1	1	1	1	1
Likely to make unsuccessful attempt	1.01 (0.72 to 1.40)	1.43 (1.00 to 2.06)	0.97 (0.69 to 1.37)	0.92 (0.65 to 1.29)	1.46 (1.02 to 2.08)	1.10 (0.78 to 1.55)	1.43 (1.00 to 2.04)
Unlikely to make attempt	0.58 (0.44 to 0.75)	0.74 (0.55 to 0.99)	0.59 (0.45 to 0.78)	0.51 (0.38 to 0.67)	0.76 (0.57 to 1.01)	0.55 (0.41 to 0.73)	0.56 (0.42 to 0.74)

¹Note sociodemographic characteristics (described in table 3a) were also entered into each model

²Blank cells indicate no significant relationship in bivariate analysis

Table 4: Perceptions of cigarette design (harm, appeal, trial)

	Standard cigarette % ¹	Cigarette with warning % ¹	Green cigarette % ¹
Harmful to health	38.8	69.1*	70.2*
Think of dangers	20.9	58.1* [#]	53.5*
Unattractive	25.2	61.7*	68.7*
Unstylish	37.4	66.0*	69.4*
Not nice to be seen with	19.8	55.2*	60.2*
Not appealing to people my age	17.8	51.5*	57.4*
Likely trial (personally)	79.4	35.7*	21.5*
Likely trial (for never smokers)	70.1	21.1*	16.5*

¹ Percentages shown indicate an answer within the three highest agreement categories on a seven point semantic scale.

* Significant difference in comparison to the standard cigarette (p<0.001)

Significant difference in comparison to the green cigarette (p<0.05)




Figure 1 Pack inserts highlighting the benefits of quitting or providing tips on how to do so

142x222mm (300 x 300 DPI)



Figure 2: Standard cigarette, warning cigarette and green cigarette

17x5mm (300 x 300 DPI)

For peer review only

STROBE 2007 (v4) checklist of items to be included in reports of observational studies in epidemiology*
Checklist for cohort, case-control, and cross-sectional studies (combined)

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any pre-specified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	6,7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8-11
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	10-12
Bias	9	Describe any efforts to address potential sources of bias	6
Study size	10	Explain how the study size was arrived at	6
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	10-12
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	11-12
		(b) Describe any methods used to examine subgroups and interactions	11-12
		(c) Explain how missing data were addressed	7
		(d) <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6,29-30
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	26-27
		(b) Indicate number of participants with missing data for each variable of interest	NA
Outcome data	15*	<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	13-16
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	13-16
		(b) Report category boundaries when continuous variables were	28-29

		categorized	
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10-12
Discussion			
Key results	18	Summarise key results with reference to study objectives	16-19
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	18
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	19-20
Generalisability	21	Discuss the generalisability (external validity) of the study results	17-19
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	20

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes among young adult smokers in the United Kingdom: A cross-sectional online survey

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-019662.R2
Article Type:	Research
Date Submitted by the Author:	19-Jun-2018
Complete List of Authors:	Moodie, Crawford; University of Stirling, Institute for Social Marketing Hiscock, Rosemary; University of Bath Thrasher, Jim; University of South Carolina Reid, Garth; NHS Health Scotland
Primary Subject Heading:	Smoking and tobacco
Secondary Subject Heading:	Public health
Keywords:	Smoking, Packaging, Inserts, Cigarettes

SCHOLARONE™
Manuscripts

Title: Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes among young adult smokers in the United Kingdom: A cross-sectional online survey

Authors: Crawford Moodie¹, Rosemary Hiscock², Jim Thrasher³, Garth Reid⁴

Affiliations:

¹ Institute for Social Marketing, Faculty of Health Sciences and Sport, University of Stirling, Stirlingshire, Scotland

² Department for Health, University of Bath, England

³ Health Promotion, Education, and Behavior, Arnold School of Public Health, University of South Carolina, Columbia, United States

⁴ Policy Evaluation, NHS Health Scotland, Edinburgh, Scotland

Corresponding author: Crawford Moodie, Institute for Social Marketing, Department of Health Sciences and Sport, University of Stirling, Stirlingshire, Scotland FK9 4LA. Tel: +44 (0)1786 466456. Email: c.s.moodie@stir.ac.uk

Keywords: Smoking, Packaging, Inserts, Cigarettes

Word count: 5045

1
2
3 **Perceptions of cigarette pack inserts promoting cessation and dissuasive cigarettes**
4 **among young adult smokers in the United Kingdom: A cross-sectional online survey**
5
6
7

8
9 **ABSTRACT**

10
11 **Objectives:** To explore young adult smokers' perceptions of cigarette pack inserts promoting
12 cessation and cigarettes designed to be dissuasive.
13

14
15 **Design:** Cross-sectional online survey.
16

17
18 **Setting:** United Kingdom.
19

20
21 **Participants:** The final sample was 1766 young adult smokers, with 50.3% male and 71.6%
22 white British. To meet the inclusion criteria participants had to be 16-34 years old and smoke
23 factory-made cigarettes.
24

25
26 **Primary and secondary outcome measures:** Salience of inserts, perceptions of inserts as
27 information provision, perceptions of inserts on quitting, support for inserts, and perceived
28 appeal, harm and trial of three cigarettes (a standard cigarette, a standard cigarette displaying
29 the warning 'Smoking kills', and a green cigarette).
30
31

32
33 **Results:** Half the sample indicated that they would read inserts with three-fifths indicating
34 that they be a good way to provide information about quitting (61%). Just over half indicated
35 that inserts would make them think more about quitting (53%), help if they decided to quit
36 (52%), are an effective way of encouraging smokers to quit (53%), and supported having
37 them in all packs (55%). Participants who smoked factory-made cigarettes and other tobacco
38 products (compared to exclusive factory-made cigarette smokers), had made a quit attempt
39 within the last six months (compared to those that had never made a quit attempt), or were
40 likely to make a successful quit attempt in the next six months (compared to those unlikely to
41 make a quit attempt in the next six months), were more likely to indicate that inserts could
42 assist with cessation. Multivariable logistic regression modelling suggested that compared
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 with the standard cigarette, the cigarette with warning (adjusted Odds Ratio=17.71; 95%CI
4 13.75-22.80) and green cigarette (adjusted Odds Ratio=30.88; 95%CI 23.98-39.76) were
5
6
7 much less desirable (less appealing, more harmful, less likely to be tried).
8

9 **Conclusions:** Inserts and dissuasive cigarettes offer policy makers additional ways of using
10 the pack to reduce smoking.
11
12

13 **Strengths and limitations of this study**

- 14
15
16
17
18 ▪ The main strength of this study is that it allows an insight into how young adult
19 smokers perceive two innovative tobacco control measures (pack inserts promoting
20 cessation and dissuasive cigarettes).
21
22
- 23
24 ▪ The main limitation of the study is that it does not provide any insight into actual
25 smoking behaviour.
26
27
- 28
29 ▪ Additional limitations include the novelty of the stimuli and forced exposure to this,
30 and the use of self-selection.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

While packaging remains a key marketing driver for tobacco companies, more than 100 countries now require pictorial health warnings on cigarette packs,¹ which can limit pack appeal.² Some countries have gone even further by implementing plain (or standardised) packaging, which severely reduces the promotional power of the pack. The United Kingdom (UK) became the third country to fully implement standardised packaging in May 2017, following Australia in December 2012 and France in January 2017. In the UK all cigarette packs must be drab brown with pictorial warnings on 65% of the front and back of packs and additional health messages on 50% of the sides of the pack. Although these changes have reduced the ability of tobacco companies to use the pack to create favourable perceptions of the brand and of smoking, there is clearly more scope for using the packaging to dissuade consumers. Regulators and academics have typically focused on the exterior of the cigarette pack, with little consideration of how the pack interior, for instance pack inserts or cigarettes, which have long been used by tobacco companies to promote their brands, could potentially be used to encourage smokers to think about their smoking behaviour. This is the focus of our study.

Tobacco companies have used the inside of the cigarette pack to communicate with consumers since the late 19th century, via cigarette cards, coupons and promotional inserts. Only in Canada are they required, by law, to include pack inserts with health messaging. Sixteen text-only inserts were required in packs between 2000 and 2012, with nine encouraging cessation and seven providing health risk information.³ These were replaced with eight new inserts, with coloured graphics and positively framed messages about the benefits of quitting or tips on how to do so, in 2012. Few studies have explored perceptions of pack inserts,⁴⁻⁸ with only two assessing smokers' perceptions of, and responses to, the inserts used in Canada.⁹⁻¹¹ In focus group research in Scotland,⁹ with smokers aged 16 and

1
2
3 over who were shown seven of the inserts used in Canada, the general view was that they
4 would capture attention and be read due to their novelty and visibility when opening the pack.
5
6 Inserts were also thought to have a long lasting impact as they would be removed from the
7 pack and remain visible within the household or elsewhere, or as litter.⁹ The positive
8
9 messaging was liked and thought to increase message engagement. The inserts were often
10 preferred to the on-pack warnings, although both were deemed necessary. Some participants
11 suggested that inserts could encourage them to stop smoking, and they were generally
12 considered to have the potential to alter the behaviour of younger people, would-be smokers
13 and those wanting to quit.⁹ In Canada, a longitudinal online survey with smokers aged 18 and
14 over found that between 26% and 31% at each wave reported having read pack inserts at least
15 once in the prior month; those intending to quit or having recently tried to do so were
16 significantly more likely to have read them.¹⁰ In addition, while reading warnings on the pack
17 exterior decreased over time, reading pack inserts increased over time, with more frequent
18 reading independently associated with self-efficacy to quit, quit attempts, and sustained
19 quitting at follow-up.¹¹

20
21
22 The cigarette itself is also an important communications tool,^{12,13} which has long been
23 used by tobacco companies as a marketing device but has yet to be used by regulators to deter
24 smoking. As cigarettes are primarily responsible for tobacco related mortality and morbidity
25 and predicted to continue to dominate the global market for some time yet,¹⁴ research
26 exploring the potential impact of standardising the appearance of cigarettes to make them less
27 desirable is long overdue. Some recent research has examined consumer perceptions of
28 cigarettes that have been designed to be 'dissuasive', including unattractively coloured
29 cigarettes,^{15,16} cigarettes with the warning 'Smoking kills' on the cigarette paper,^{17,18} and
30 cigarettes displaying the 'minutes of life lost due to smoking' on the cigarette paper.¹⁹ In each
31 of these studies the dissuasive cigarettes were generally viewed more negatively than regular
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 cigarettes. For instance, a qualitative study with young women smokers in New Zealand
4 found that unattractively coloured cigarettes, particularly green or brown coloured cigarettes,
5 were perceived as more harmful than other cigarettes, with it less likely that they or others
6 their age would want to use them.¹⁵ An in-home survey in the UK with 11-16 year olds, who
7 were shown an image of a cigarette stick displaying ‘Smoking kills’, found that 53%
8 indicated that this would make people want to give up smoking, 71% indicated that it would
9 put people off starting to smoke, and 85% supported having a warning on all cigarettes.¹⁸

10
11 In this study our objective was to explore, for the first time, young adult smokers’
12 perceptions of pack inserts and dissuasive cigarettes (a cigarette displaying the warning
13 ‘Smoking kills’ and a green coloured cigarette).
14
15
16
17

18 **METHODS**

19 **Design and sample**

20 An online survey was conducted in January-February 2016 with smokers aged 16-34 years
21 old in the UK; an online survey is a suitable approach given that 99% of this age group in the
22 UK are recent internet users.²⁰ The sample (n=1970) was recruited by online market research
23 company ‘Research Now’ from their panel of over 400,000 people (www.researchnow.com).
24 After Research Now excluded those who had completed the survey in less than the minimum
25 completion time (n=193), which they had set prior to data collection commencing, and those
26 providing responses to open-ended questions that indicated that they had not taken the survey
27 seriously (n=11), the final sample was 1766 (89.6% of completed surveys). The final sample
28 was 50.3% male, with 53.9% aged 25-34 years and 71.6% white British. Most participants
29 smoked 10 or less cigarettes per day, with 46.0% exclusive factory-made cigarette smokers
30 (see Table 1 for sample and smoking-related characteristics).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 here

Procedure

An email invite was sent by Research Now to their online panel in the UK. Research Now is an established online market research company in the UK and elsewhere,²¹ with their panels recruited from a wide range of sources, such as internet sites, advertising and partnerships with other websites. Research Now, like other online panels, has details of their members' demographics and other characteristics that are used to profile target samples. Response rate details are not available when using this sampling methodology however as recording contact, participation and refusal rates is not practical.²² For those that responded to the email invite, they answered screening questions about their age, smoking status and types of tobacco products used, with those that did not meet the inclusion criteria (factory-made cigarette smokers aged 16-34 years) excluded.

Those eligible for inclusion were presented with an information page explaining the study aim (to explore what young adult smokers thought about cigarettes and pack inserts), and relevant ethical information (their right to withdraw at any time, assurances of confidentiality and anonymity, and contact details if they had any concerns or would like to request a copy of the published findings). They were then presented with a consent page, with consent required for participation. Survey questions were presented in the same order for all participants, except the questions exploring perceptions of the three cigarette types (standard cigarette (SC), warning cigarette (WC), green cigarette (GC)), where the ordering was randomised; the ordering of the presentation of the three cigarettes (shown in Figure 1) was also randomised. There was no missing data as participants could only proceed to the next question if they had provided an answer to the previous question.

1
2
3
4
5 Figure 1 here
6
7
8

9
10 Prior to the questions on inserts, participants were shown an image of a cigarette pack with an
11 insert shown in the front of the pack – as they typically appear in packs – alongside the text
12 ‘We have some questions on pack inserts, which can sometimes be found inside packs (see
13 image for example)’. For each question about inserts, participants were shown the question
14 and an image of one insert. Four different inserts were used in total, as shown in Figure 2,
15 with these chosen from the eight used in Canada as they were considered most relevant to our
16 sample. The words ‘Health Canada’ were removed from the bottom of each insert to make
17 them more relevant for participants in the UK. The median time for survey completion was 9
18 minutes 28 seconds. Participants received a nominal incentive (50 pence) for participation, as
19 is common for online panels. The study received ethical approval from the School of Health
20 Sciences Ethics Committee at the University of Stirling.
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 Figure 2 here
36
37
38

39 **Patient and public involvement**

40
41 There was no patient or public involvement in the development, design or conduct of this
42 study.
43
44
45

46 **Measures**

47
48
49
50
51
52 *Inserts: Salience and information provision*
53
54
55
56
57

1
2
3 Participants were asked ‘If this type of insert was in your cigarette pack, do you think that
4 you would read it?’ and ‘If this type of insert was in your cigarette pack, do you think that
5 you would read it if you were interested in quitting?’ They were also asked ‘Do you think
6 that inserts would be a good way to provide information to smokers about quitting?’⁵
7
8
9
10
11 Response options for each were ‘Yes’, ‘No’ and ‘Not sure’.
12
13

14 15 16 *Inserts: Cessation*

17
18 Three questions assessed to what extent participants agreed or disagreed that inserts would
19 make them think about quitting, and help them quit: ‘Do you agree or disagree that having
20 these types of inserts in every cigarette pack would make you think more about quitting?’,
21 ‘Do you agree or disagree that having these types of inserts in every cigarette pack might help
22 you if you decided to quit?’, and ‘Do you agree or disagree that having these types of inserts
23 inside every cigarette pack would be an effective way of helping smokers who want to quit?’⁶
24
25 Response options for each were ‘Strongly disagree’, ‘Disagree’, ‘Neither agree nor disagree’,
26 ‘Agree’, ‘Strongly agree’ and ‘Don’t know’.
27
28
29
30
31
32
33
34
35
36
37

38 *Inserts: Support*

39
40 A five-point semantic scale assessed support, with anchors ‘All cigarette packs should have
41 inserts like this in them-No cigarette packs should have inserts like this in them’.
42
43
44
45

46 *Cigarette design: Appeal, harm and trial*

47
48 Seven-point semantic scales assessed appeal, harm and likely trial. Appeal was assessed via
49 four scales, with anchors ‘Attractive-Unattractive’, ‘Stylish-Not stylish’, ‘Not nice to be seen
50 with-Nice to be seen with’ and ‘Not appealing to people my age-Appealing to people my
51 age’. Harm was assessed via two scales, with anchors ‘Looks harmful to health-Does not look
52
53
54
55
56
57
58
59
60

1
2
3 harmful to health' and 'Makes me think about the dangers of smoking-Does not make me
4 think about the dangers of smoking'. Likely trial was assessed via two scales, 'If a friend
5 offered you each of these cigarettes, how likely would you be to try them?' and 'If someone
6 your age who had never smoked before was going to try a cigarette, how likely do you think
7 they would be to try each of these cigarettes?' Both scales assessing trial ranged from 'Not at
8 all likely' to 'Very likely'.

9
10
11
12
13
14
15
16 A factor analysis of the eight variables on appeal, harm and trial, collated for the three
17 cigarette types (SC, WC, GC), was undertaken. Checks indicated that the data was suitable
18 for factor analysis (Kaiser Meyer Olkin=0.845, Bartlett's test of sphericity (approx. chi-
19 square 18062.842, df=276, p<0.001), with no correlations between the variables >0.9). The
20 extraction method used was Principal Axis Factoring and the criteria for extraction was
21 eigenvalues>1. All eight variables loaded on a single factor with factor loadings that were
22 >0.5. High factor scores indicated that a cigarette was desirable and low scores that it was
23 undesirable. The factor was used as the outcome measure of cigarette desirability in the
24 regression analysis. Visual inspection and the Kolmogorov-Smirnov test indicated that the
25 factor was non-normal (because responses for the dissuasive cigarettes indicated they were
26 undesirable generally) and attempts to normalise it using normit rankit methods failed.
27 Therefore, the factor was divided into tertiles, with the tertile indicating undesirable factor
28 scores compared with the other two tertiles. This was the outcome variable in logistic
29 regression analysis.

30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 *Sociodemographic characteristics*

49
50 Age, gender, ethnicity, educational attainment and economic status (based on chief income
51 earner) were obtained. Preliminary analysis showed that education was associated with how
52 pack inserts were perceived, whereas both education and economic status were associated
53
54
55
56
57
58
59
60

1
2
3 with how cigarettes were perceived. As such, for the analysis of the cigarettes a count
4
5 procedure was used to create a variable for low socioeconomic status (SES): low education
6
7 (General Certificate of Secondary Education: GCSE or below) and/or low economic status
8
9 (routine or manual occupation, long-term unemployed or long-term sick or disabled).
10

11 12 13 *Smoking behaviour*

14
15 Smoking status was assessed with 'Which of these best describes you?' with response
16
17 options: 'I have never smoked', 'I used to smoke, but don't now', 'I smoke, but not every
18
19 day', and 'I smoke every day'. Type of products used was assessed with 'What type(s) of
20
21 tobacco products do you smoke?' with response options: 'Only factory-made (packet
22
23 cigarettes', 'Factory-made and roll-your-own cigarettes', 'Factory-made cigarettes and other
24
25 tobacco products (e.g. cigars, shisha, etc)', 'Only roll-your-own cigarettes' and 'Only other
26
27 tobacco products (e.g. cigars, shisha, etc)'. The Heaviness of Smoking Index (HSI)²³ was
28
29 used as a measure of dependence, based on daily consumption and time to first cigarette.
30
31
32
33
34

35 *Quitting and self-efficacy*

36
37 Participants were asked 'Have you ever made an attempt to quit smoking that lasted at least
38
39 24 hours?'²⁴ (Yes within the last six months, Yes more than six months ago, I have never
40
41 tried to quit for more than 24 hours). They were also asked 'How likely are you to try to quit
42
43 smoking within the next six months?'²⁵ (Not at all, A little, Moderately, Very, Extremely,
44
45 Don't know), with those responding 'Not at all', 'A little', 'Moderately' or 'Don't know'
46
47 classified as 'Unlikely to make a quit attempt in the next six months'. To measure quitting
48
49 self-efficacy, participants were asked 'If you decided to quit smoking in the next six months,
50
51 how sure are you that you would succeed?'²⁶ (Not at all, A little, Moderately, Very,
52
53 Extremely, Don't know). Those who responded to the likelihood of quitting question with
54
55
56
57
58
59
60

1
2
3 'Very or 'Extremely' and to the quitting efficacy question with 'Not at all', 'A little',
4
5 'Moderately' or 'Don't know' were classified as 'unlikely to make a successful quit attempt
6
7 in the next six months'. Those who responded 'Very' or 'Extremely' to both questions were
8
9 classified as 'likely to make a successful quit attempt in the next six months'.
10

11 12 13 **Analysis**

14
15 Data was analysed using Microsoft office Excel 2013, SPSS v22 and v23 and MLWin
16
17 v2.33.²⁷ The insert variables were dichotomised into yes/agreement and
18
19 no/disagreement/neutral/not sure/don't know. The dichotomised insert variables were the
20
21 outcomes of the logistic regression models. The independent variables were gender, age,
22
23 education, ethnicity, dependence (tertiles of HSI), tobacco product(s) smoked, previous quit
24
25 attempt lasting at least 24 hours, and likely efficacy of a quit attempt in the next six months.
26
27 Percentages in agreement were calculated. Age, gender and education (as a measure of SES)
28
29 were entered into all models to account for any sampling inadequacies. Other variables were
30
31 entered where $p < 0.10$ in chi square tests. The models were assessed for multicollinearity via
32
33 comparison of standard errors²⁸ and none was found.
34
35
36

37
38 For each of the eight seven-point semantic scales, the percentage of participants
39
40 choosing one of the three points nearest the undesirable anchor (e.g. unattractive, not nice to
41
42 be seen with, looks harmful to health) was calculated for each of the three cigarette types
43
44 (SC, WC, GC). Thus, 24 percentages were calculated. Differences between the three
45
46 cigarettes were tested using Cochran's Q and pairwise comparisons.
47

48
49 Multilevel logistic regression modelling of cigarette desirability, with second order
50
51 PQL estimation,²⁹ was undertaken with cigarette evaluations (participants' response to each
52
53 of the three cigarettes) clustered within individual participants. Therefore, cigarette
54
55 evaluations were level one cases and participants were entered at level two as a random
56
57

1
2
3 effect. All models included cigarette type as a fixed effect, where the standard cigarette was
4 compared with the warning cigarette and green cigarette. Other fixed effects at the individual
5 (participant) level were sociodemographic and smoking-related characteristics, which were
6 significantly associated with the outcome in multivariable models. This main effects model
7 tested which characteristics were associated with perceiving cigarettes as desirable. In order
8 to understand which characteristics differentiated the desirability of the three types of
9 cigarettes, interactions between cigarette type and each significant characteristic were tested.
10 Only one interaction was found, between cigarette type and SES. The interacting variables
11 (cigarette type and SES) were substituted by a cross classified variable which merged
12 cigarette type and SES. This cross classified variable was split into six categories: low SES
13 standard cigarette, low SES warning cigarette, low SES green cigarette, not low SES standard
14 cigarette, not low SES warning cigarette, not low SES green cigarette. To understand the
15 interaction five models were run with the reference category of the cross classified variable
16 different each time.^{30,31}

35 RESULTS

39 Perceptions of inserts

40 Half the sample indicated that they would read inserts, with approximately three-fifths
41 indicating that they would read them if interested in quitting (60%), and that they would be a
42 good way to provide information about quitting (61%). Just over half strongly agreed/agreed
43 that inserts may make them think more about quitting (53%), help them if they decided to
44 quit (52%), that they are an effective way of encouraging smokers to quit (53%), and that all
45 cigarette packs should have inserts (55%), see Table 2.

1
2
3 Table 2 here
4
5
6

7 **Sociodemographic differences in perceptions of inserts**

8
9 Women were more likely than men to indicate that they would read inserts (aOR=1.24;
10 95%CI 1.02-1.50), and 25-34 year olds less likely than 16-19 year olds to think that they
11 were a good way of providing information about quitting (aOR=0.76; 95%CI 0.60-0.98).
12
13 Compared with white British participants, white non-British (aOR=0.70; 95%CI 0.50-0.98)
14 and Asian (aOR=0.67; 95%CI 0.49-0.92) participants were less likely to suggest that they
15 would read inserts if trying to quit, white non-British (aOR=0.58; 95%CI 0.41-0.81) and
16 Black (aOR=0.61; 95%CI 0.38-0.98) participants were less likely to indicate that inserts
17 would make them think about quitting, and white non-British (aOR=0.62; 95%CI 0.44-0.87)
18 and Asian (aOR=0.70; 95%CI 0.51-0.96) participants were less likely to support having
19 inserts in all packs, see Table 3.
20
21
22
23
24
25
26
27
28
29
30
31
32

33 **Smoking-related differences**

34
35 Compared to exclusive factory-made cigarette smokers, those who also smoked roll-your-
36 own cigarettes were more likely to indicate they would read inserts (aOR=1.35; 95%CI 1.09-
37 1.66), read them if trying to quit (aOR=1.61; 95%CI 1.30-2.00), that they would make them
38 think about quitting (aOR=1.31; 95%CI 1.06-1.62), help them if they decided to quit
39 (aOR=1.31; 95%CI 1.06-1.61), and that they would be an effective way of encouraging
40 smokers to quit (aOR=1.27; 95%CI 1.03-1.56), see Table 3. Compared to exclusive factory-
41 made cigarette smokers, those who also smoked other tobacco products (e.g. cigars, shisha)
42 were more likely to indicate they would read inserts if trying to quit (aOR=1.39; 95%CI 1.04-
43 1.86) and that inserts might help them if they decided to quit (aOR=1.34; 95%CI 1.01-1.78).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Participants who had made a quit attempt more than six months ago (aOR=1.30; 95%CI 1.00-1.69), or within the last six months (aOR=1.67; 95%CI 1.29-2.15), were more likely to indicate that they would read inserts than those who had never made a quit attempt. Those who had made a quit attempt in the last six months were also more likely than those who had never made a quit attempt to indicate that inserts were a good way to provide information about quitting (aOR=1.54; 95%CI 1.20-1.98), that they would read them if trying to quit (aOR=1.51; 95%CI 1.17-1.94), make them think about quitting (aOR=1.46; 95%CI 1.14-1.88), help them if they decided to quit (aOR=1.35; 95%CI 1.05-1.73), and that they would be an effective way of encouraging smokers to quit (aOR=1.33; 95%CI 1.04-1.71).

Compared to those likely to make a successful quit attempt in the next six months, those unlikely to make a quit attempt in the next six months were less likely to indicate that they would read inserts (aOR=0.58; 95%CI 0.44-0.75), read them if trying to quit (aOR=0.74; 95%CI 0.55-0.99), that they would make them think about quitting (aOR 0.59; 95%CI 0.45-0.78), help them if they decided to quit (aOR=0.51; 95%CI 0.38-0.67), that they would be effective for smokers if they decided to quit (aOR=0.55; 95%CI 0.41-0.73), or support them (aOR=0.56; 95%CI 0.42-0.74). Compared to those likely to make a successful quit attempt in the next six months, those unlikely to make a successful quit attempt in the next six months were more likely to read inserts if trying to quit (aOR=1.43; 95%CI 1.00-2.06), thought that they were a good way to provide information to smokers about quitting (aOR=1.46; 95%CI 1.02-2.08), and support them (aOR=1.43; 95%CI 1.00-2.04).

Table 3 here

Perceptions of cigarette design

With respect to harm, participants were less likely to think that the standard cigarette (SC)

1
2
3 looked harmful than the warning cigarette (WC) or green cigarette (GC) ($p<0.001$), and less
4 likely to think that the SC made them think more about the dangers of smoking than the WC
5 or GC ($p<0.001$), see Table 4. Participants were also less likely to indicate that the GC would
6 make them think of the dangers of smoking than the WC ($p=0.01$). In terms of appeal,
7 participants were more likely to consider the SC attractive, and stylish, than the WC or GC
8 (both $p<0.001$). The SC was also considered to be nicer to be seen with, and more appealing
9 to people their age, than the WC or GC (both $p<0.001$). In terms of trial, whereas only 8.9%
10 indicated that they would be unlikely to try a SC if offered by a friend, this was 45.4% for the
11 WC and 66.5% for the GC (both $p<0.001$). Similarly, while only 14.8% indicated that a never
12 smoker their age would be unlikely to try a SC, this was 63.3% for the WC and 71.6% for the
13 GC (both $p<0.001$), see Table 4.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 Table 4 here
30
31
32

33 **Perceptions of cigarette desirability**

34
35 Main effects multivariable logistic regression modelling suggested that in comparison to the
36 SC, the WC (aOR=17.71; 95%CI 13.75-22.80) and GC (aOR=30.88; 95%CI 23.98-39.76)
37 were much more likely to be perceived as undesirable (i.e. less appealing, more harmful, less
38 likely to be tried). The model also indicated which smokers were more likely to rate the
39 cigarettes as undesirable: women were more likely than men (aOR=1.30; 95%CI 1.10-1.54),
40 and low SES more likely than those not low SES (aOR=1.26; 95%CI 1.06-1.50), to consider
41 all three cigarettes undesirable. Compared to exclusive factory-made cigarette smokers, those
42 who also smoked roll-your-own cigarettes (aOR=0.78; 95%CI 0.65-0.93) or other tobacco
43 products (aOR=0.73; 95%CI 0.56-0.93) were less likely to consider all three cigarettes
44 undesirable. Those not likely to make a quit attempt in the next six months were less likely
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 than those likely to make a quit attempt in the next six months (aOR=0.62; 95%CI 0.49-0.78)
4
5 to consider all three cigarettes undesirable.
6

7 Only one significant interaction, between cigarette type and SES, was found ($p<0.05$).
8
9 Both SES groups perceived the WC significantly more undesirable than the SC, and the GC
10 significantly more undesirable than the WC (see Table 5). Low SES participants were
11 significantly more likely than those not low SES to perceive the SC (aOR=1.89; 95%CI 1.18-
12 3.03) and GC (aOR=1.43; 95%CI 1.13-1.80) as undesirable; there was no difference for the
13 WC (aOR=0.99; 95%CI 0.78-1.25).
14
15
16
17
18
19
20
21

22 Table 5 here
23
24
25

26 **DISCUSSION**

27
28 Our findings suggest that inserts highlighting the benefits of quitting or providing tips on how
29 to do so may have the potential to encourage cessation, and dissuasive cigarettes may help to
30 reduce the desirability of smoking. Just as tobacco companies have used inserts and cigarette
31 design to create interest in their products, our study suggests that greater attention to how
32 these could be used to promote cessation appears warranted.
33
34
35
36
37
38

39 Health messages need to capture attention to be effective.³² In this regard, at least half
40 our sample indicated that they would read inserts (50%) and read them if interested in
41 quitting (60%). In Canada, an observational study found that approximately a quarter of
42 smokers reported reading them at least once within the last month,¹⁰ increasing to about one-
43 third of smokers over two years of follow-up.¹¹ Like the smokers in our study who indicated
44 that they would read the inserts, smokers in Canada who had read the inserts were more
45 likely to be female, intend to quit or had recently tried to quit; in our study, they were also
46 more likely to be white British, have moderate dependence, and use factory-made cigarettes
47
48
49
50
51
52
53
54
55
56
57

1
2
3 and other tobacco products. Future research could explore why dual users (smokers of
4 factory-made cigarettes and other tobacco products) were more likely to indicate that they
5 would read inserts, but as inserts are typically only found in cigarette packs then for those
6 who use other tobacco products they may be seen as more of a novelty and therefore more
7 likely to capture attention.
8
9
10
11
12

13
14 Approximately three-fifths (61%) of smokers in our study thought that inserts were a
15 good way to provide information about quitting to smokers, with only 25% disagreeing. In
16 comparison, an earlier study in Canada, commissioned by Health Canada, found that 48% of
17 smokers indicated that messaging on inserts was a good way to provide information to
18 smokers, with 47% disagreeing.⁵ Just over half our sample agreed/strongly agreed that inserts
19 may make them think more about quitting, help them if they decided to quit, and that they are
20 an effective way of encouraging smokers to quit, whereas in New Zealand only 34% of
21 smokers and recent quitters agreed/strongly agreed that inserts would be an effective way of
22 encouraging reduced consumption or quitting.⁶ There may be various reasons for the
23 differences between our findings and earlier research. For instance, when this earlier research
24 was conducted cigarette packs displayed text-only health warnings and it may be that having
25 pictorial warnings on packs, as is required in Scotland, may prompt smokers to look for
26 information on how to quit and the benefits of doing so. Insert design is also likely to be
27 relevant. Whereas the inserts used in earlier research were limited to text, the inserts used in
28 this study (which have been used in Canada since 2012) included coloured graphics, which is
29 typical of promotional inserts used by tobacco companies and likely enhanced their impact.
30
31 This would be consistent with the health communications and warnings literature, which
32 demonstrates the importance of supporting text with pictorials.^{32,33} Future research exploring
33 insert design (e.g. use of imagery, inclusion of cessation resource information, length and
34 framing of messages, etc) would be of value.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 More than half our sample supported the inclusion of inserts promoting cessation
4 inside every cigarette pack, with only a fifth opposing this. Within the European Union, the
5 Tobacco Products Directive (TPD)³⁴ does not require tobacco companies to include health
6 communication inserts in packs, but allows member states to introduce measures beyond
7 those specified. Among governmental representatives that responded to the consultation on
8 the revision of the TPD there was strong support for improving consumer information via
9 mandatory pictorial warnings, with those supportive arguing that additional information, such
10 as pack inserts, would help to deliver more accurate health information.³⁵ If there is support
11 for inserts among governmental representatives, and little opposition among smokers (the
12 group most likely to be resistant), they are clearly a viable option for regulators.
13
14
15
16
17
18
19
20
21
22
23

24 Tobacco industry journals describe the cigarette as an increasingly important
25 advertising medium for tobacco companies.¹² However, until recently, the public health focus
26 has been on the potential of regulating the contents of cigarettes to reduce palatability or
27 addictiveness,³⁶ with little consideration of the possibility of regulating the appearance of
28 cigarettes to reduce its importance as a promotional tool. We found that the two dissuasive
29 cigarettes were perceived as significantly more harmful and less appealing than the standard
30 cigarette, and less likely to encourage trial. The harm, appeal and trial items loaded onto a
31 single 'desirability' factor, with the dissuasive cigarettes considered much more undesirable
32 than the standard cigarette. The findings are consistent with earlier research, where cigarettes
33 with the warning 'Smoking kills' were considered a constant reminder of the associated
34 harms and, partly due to the perceived discomfort of being observed by others smoking a
35 cigarette displaying this message, unappealing for smokers.^{8,16,17,18} Previous studies have also
36 found unattractively coloured cigarettes to be perceived as more harmful than other cigarettes
37 and also repellent, being a cigarette that young people did not think that others their age
38 would use.^{15,16,37,38} As with the inserts, the dissuasive cigarettes (and also the standard
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 cigarette) were considered more desirable among dual users than exclusive factory-made
4
5 cigarette smokers; again it is not clear why this was the case but further research with dual
6
7 users, or indeed those also using vaping devices (not assessed in this study), would be
8
9 fruitful.

10
11 In terms of limitations, the cross-sectional design did not allow us to assess causality;
12
13 that inserts and dissuasive cigarettes are not available on the UK market prevents more robust
14
15 study designs such as longitudinal studies. Another potential limitation concerns the novelty
16
17 of the stimuli, which may have influenced responses, and forced exposure to the stimuli. In
18
19 addition, we only used four inserts, rather than the full set of eight used in Canada, which
20
21 includes inserts less relevant to our sample. While online surveys have been used for previous
22
23 research exploring cigarette packaging, inserts and dissuasive cigarettes,³⁹⁻⁴² and are a
24
25 suitable survey mode for young adults, the use of an online panel and self-selection limits the
26
27 representativeness of our sample. In addition, the use of semantic differential scales can be
28
29 criticised because answers can be subject to various response biases, although we attempted
30
31 to diminish these through varying scale item direction and through our multivariate modelling
32
33 methodology.

34
35
36
37 It was argued, over two decades ago, that to offer greater protection to consumers
38
39 cigarettes should come in plain packs with health messaging on both the pack exterior and
40
41 interior.⁴³ This idea is a step closer in the UK, although there will still be no messaging on the
42
43 pack interior. That more than half of the participants in this study suggested that inserts may
44
45 help to promote cessation suggests that their inclusion in packs may be a meaningful
46
47 supplement to the on-pack warnings. Our findings suggest however that to offer the greatest
48
49 protection to consumers, it may be beneficial to supplement plain packaging and inserts with
50
51 cigarettes designed to be dissuasive. Unattractively coloured cigarettes would complement
52
53
54
55
56
57
58
59
60

1
2
3 the unattractively coloured packs, just as warnings on the cigarette would extend the
4
5 warnings on the cigarette pack. Both options are clearly viable.
6
7

8
9 **Contributors** CM designed the data collection tool and drafted and revised the paper. RH
10
11 analysed the data and drafted the Analysis and Results. JT and GR helped design the data
12
13 collection tool and commented on the paper. All authors read and approved the final
14
15 manuscript.
16

17
18 **Funding** This work was supported by Health Scotland. GR, who works for Health Scotland,
19
20 provided feedback on the survey and paper, but was not involved in the collection, analysis
21
22 and interpretation of the data.
23

24 **Competing interests** GR works for Health Scotland, who funded this study.
25

26 **Ethics approval** The study obtained ethics approval from the School of Health Sciences
27
28 Ethics Committee at the University of Stirling. Participants provided informed consent before
29
30 participating.
31

32
33 **Provenance and peer review** Not commissioned; externally peer reviewed.
34

35 **Data sharing statement** No additional data are available.
36
37

38 39 **Figure Legends:** 40

41
42 Figure 1: Standard cigarette, warning cigarette and green cigarette
43

44
45 Figure 2: Pack inserts highlighting the benefits of quitting or providing tips on how to do so
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Canadian Cancer Society. *Cigarette package health warnings. International status report (Fifth edition)*; 2016. www.tobaccolabels.ca/wp/wp-content/uploads/2016/11/Cigarette-Package-Health-Warnings-International-Status-Report-English-CCS-Oct-2016.pdf (accessed 10 Sept 2017).
2. Hammond D. Health warning messages on tobacco products: a review. *Tob Control* 2011;20:327–37.
3. Mahood G. *Canada's tobacco package label or warning system: "telling the truth" about tobacco product risks*. Geneva: WHO; 2010.
4. Tandemar Research Inc. *Tobacco health warning messages, inserts and toxic constituents information study - Final Report*. Toronto: Tandemar Research Inc, 1992.
5. Environics. *Health warning messages on the flip/side and inserts of cigarette packaging. A survey of smokers. A report prepared for Health Canada*; 2000.
6. BRC Marketing and Social Research. *Smoking health warnings study*. Wellington: Ministry of Health; 2004. www.tobaccolabels.ca/wp/wp-content/uploads/2013/12/NZ-2004-Effectiveness-of-Different-Health-Warnings-in-Helping-People-Consider-Their-Smoking-Related-Behaviour-Government-Report.pdf (accessed 12 Aug 2017).
7. Gallopel-Morvan K, Moodie C, Hammond D, *et al*. Consumer understanding of cigarette emission labelling. *Eur J Pub Health* 2011;21:373-5.
8. Moodie C. Novel ways of using tobacco packaging to communicate health messages: Interviews with packaging and marketing experts. *Addict Res Theory* 2016;24:54-61.

- 1
2
3 9. Moodie C. Adult smokers' perceptions of cigarette pack inserts promoting cessation:
4 A focus group study. *Tob Control* 2018;27:72-7.
5
6
- 7 10. Thrasher JT, Osman A, Abad EN, *et al.* The innovative use of cigarette package
8 inserts to supplement pictorial health warnings: An evaluation of the Canadian policy.
9
10
11
12 *Nicot Tob Res* 2015;17:870-5.
- 13 11. Thrasher JF, Swayampakala K, Cummings KM, *et al.* Cigarette package inserts can
14 promote efficacy beliefs and sustained smoking cessation attempts: A longitudinal
15 assessment of an innovative policy in Canada. *Prev Med* 2016;88:59-65.
16
17
18
- 19 12. Rossell S. Ready to roll. *Tob Report* 2017;2:44-5.
20
21
- 22 13. Smith KC, Washington C, Welding K, *et al.* Cigarette stick as valuable
23 communicative real estate: a content analysis of cigarettes from 14 low-income and
24 middle-income countries. *Tob Control* 2017;26:604-7.
25
26
27
- 28 14. Hedley D. Tobacco wars: A new hope - the changing future. *Tob J Int* 2015;3:31-4.
29
- 30 15. Hoek J, Robertson C. How do young adult female smokers interpret dissuasive
31 cigarettes? *J Social Marketing* 2015;5:21-39.
32
33
- 34 16. Hoek J, Gendall P, Eckert C, *et al.* Dissuasive cigarette sticks: the next step in
35 standardised ('plain') packaging? *Tob Control* 2016;15:699-705.
36
37
38
- 39 17. Moodie C, Purves R, McKell J, *et al.* Novel means of using cigarette packaging and
40 cigarettes to communicate health risk and cessation messages: A qualitative study.
41
42
43
44 *Internat J Mental Health Addiction* 2015;13:333-44.
- 45 18. Moodie C, Mackintosh AM, Gallopel-Morvan K, *et al.* Adolescents' perceptions of
46 health warnings on cigarettes. *Nicot Tob Res* 2017;29:1232-7.
47
48
- 49 19. Hassan L, Shui E. No place to hide: two pilot studies assessing the effectiveness of
50 adding a health warning to the cigarette cigarette. *Tob Control* 2015;24:e3-5.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 20. ONS. *Internet users in the UK: 2017*.
4
5 [www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetuser](http://www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2017)
6
7 [s/2017](http://www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2017) (accessed 15 Sept 2017).
8
9 21. Hoek J, Gendall P, Eckert C, *et al*. Effects of brand variants on smokers' choice
10 behaviours and risk perceptions. *Tob Control* 2016;25:160-5.
11
12 22. Moodie C, Sinclair L, Mackintosh AM, *et al*. How tobacco companies are perceived
13 within the UK: An online panel. *Nicot Tob Res* 2016;18:1766-72.
14
15 23. Kozlowski LT, Porter CQ, Orleans T, *et al*. Predicting smoking cessation with self-
16 reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug Alcohol*
17 *Depend* 1994;34:211-6.
18
19 24. Etter JF, Sutton S. Assessing 'stage of change' in current and former smokers Trying
20 to stop smoking: effects of perceived addiction, attributions for failure, and
21 expectancy of success. *Addiction* 2002;97:1171-82.
22
23 25. Hammond D, Fong GT, McDonald PW, *et al*. Impact of the graphic Canadian
24 warning labels on adult smoking behaviour. *Tob Control* 2003;12:391-5.
25
26 26. IARC. *IARC Handbooks of Cancer Prevention, Tobacco Control, Vol. 12: Methods*
27 *for Evaluating Tobacco Control Policies*. Lyon, France: International Agency for
28 Research on Cancer; 2008.
29
30 27. Rasbash J, Browne W, Healy M, *et al*. *A user's guide to MLwiN, version 2.3*. Bristol:
31 Centre for multilevel modelling, University of Bristol; 2015.
32
33 28. Salmond C. *Fitting complex models using health survey data*; 2006.
34
35 www.otago.ac.nz/wellington/otago020178.pdf (accessed 15 Sept 2017).
36
37 29. Goldstein H, Rasbash J. Improved approximations for multilevel models with binary
38 responses. *J Royal Statistical Society. Series A (Statistics in Society)* 1996;159:505-
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 30. Blakely T, Woodward A, Salmond C. Anonymous linkage of New Zealand mortality
4 and Census data. *Australian NZ J Pub Health* 2000;24:92-5.
5
6
7 31. Hiscock R, Murray S, Brose LS, *et al.* Behavioural therapy for smoking cessation:
8 The effectiveness of different intervention types for disadvantaged and affluent
9 smokers. *Addict Behav* 2013;38:2787-96.
10
11
12
13 32. Wogalter MS, Conzola VC, Smith-Jackson TL. Research-based guidelines for
14 warning design and evaluation. *Applied Ergonomics* 2002;33:219-30.
15
16
17 33. Houts PD, Doak CC, Doak LG, *et al.* The role of pictures in improving health
18 communication: A review of research on attention, comprehension, recall, and
19 adherence. *Patient Educ Couns* 2006;61:173-90.
20
21
22
23 34. European Commission. Directive 2014/40/EU of the European parliament and of the
24 Council of 3 April 2014 on the approximation of the laws, regulations and
25 administrative provisions of the Member States concerning the manufacture,
26 presentation and sale of tobacco and related products and repealing Directive
27 2001/37/EC. *Official Journal of the European Union* 2014;L127:1-38.
28
29
30
31 35. European Commission. *Report on the public consultation on the possible revision of*
32 *the Tobacco Products Directive (2001/37/EC). Health and Consumers Directorate-*
33 *General – Directorate D – Health systems and products D4 – Substances of human*
34 *origin and tobacco control.* 2011.
35
36 http://ec.europa.eu/health/tobacco/docs/consultation_report_en.pdf (accessed 10 Sept
37 2017).
38
39
40
41
42
43
44
45
46
47
48 36. Warner K. The national and international regulatory environment in tobacco control.
49 *Pub Health Res Practice* 2015;25:e2531527.
50
51
52
53 37. Ford A, Moodie C, Mackintosh AM, *et al.* Adolescent perceptions of cigarette
54 appearance. *Eur J Pub Health* 2014;24:464-8.
55
56
57
58
59
60

- 1
2
3 38. Moodie C, Ford A, Mackintosh AM, *et al.* Are all cigarettes just the same? Female's
4 perceptions of slim, coloured, aromatised and capsule cigarettes. *Health Educ Res*
5 2015;30:1-12.
6
7
8
9 39. Doxey J, Hammond D. Deadly in pink: The impact of female-oriented cigarette
10 packaging among young women. *Tob Control* 2011;20:353-60.
11
12
13 40. White CM, Hammond D, Thrasher JF, *et al.* The potential impact of plain packaging
14 of cigarette products among Brazilian young women: an experimental study. *BMC*
15 *Public Health* 2012;12:737.
16
17
18 41. Kotnowski K, Fong GT, Gallopel-Morvan K, *et al.* The impact of cigarette packaging
19 design among young females in Canada: findings from a discrete choice experiment.
20 *Nicot Tob Res* 2016;18:1348-56.
21
22
23 42. Hoek J, Gendall P, Maubach N, *et al.* Strong public support for plain packaging of
24 tobacco products. *Austr N Z J Pub Health* 2012;36:405-7.
25
26
27 43. Mahood G. Canada tobacco package warning system. *Tob Control* 1995;4:10-4.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Sample and smoking-related characteristics

<i>Characteristic</i>	<i>N</i>	<i>%</i>
Total	1766	100.0
Age group		
16-19	413	23.4
20-24	401	22.7
25-34	952	53.9
Gender		
Male	888	50.3
Female	878	49.7
Educational qualifications		
Other qualifications	1357	76.8
None or GCSE	409	23.2
Economic status		
Other status	1350	76.4
Routine or manual occupation, unemployed or long term sick	416	23.6
Socioeconomic status (SES)		
No indicators of low SES	1114	63.1
Low education and/or low SES	652	36.9
Ethnicity		
White British	1264	71.6
White non-British	162	9.2
Black (including mixed black and white)	79	4.5
Asian (including mixed Asian and white)	196	11.1
Other or not declared	65	3.7
Location		
England	1550	87.8
Scotland	109	6.2
Wales	73	4.1
Northern Ireland	34	1.9
Tobacco products used		
Only factory-made (packet) cigarettes	813	46.0
Factory-made and roll-your-own cigarettes	681	38.6
Factory-made cigarettes and other products (e.g. cigars, shisha)	272	15.4
Cigarettes per day		
10 or less	1272	72.0
11-20	433	24.5
21-30	46	2.6
31 or more	15	0.8
Time to first cigarette		
Within 5 minutes	263	14.9
6 to 30 minutes	570	32.3

<i>Characteristic</i>	<i>N</i>	<i>%</i>
31 to 60 minutes	315	17.8
After 60 minutes	618	35.0
Heaviness of Smoking Index (HSI)		
0 little dependence	601	34.0
1	257	14.6
2	418	23.7
3	293	16.6
4	156	8.8
5	28	1.6
6 high dependence	13	0.7
Dependence (Tertiles of HSI)		
Low-dependence	601	34.0
Mid-dependence	675	38.2
High-dependence	490	27.7
Made an attempt to quit smoking that lasted at least 24 hours?		
Yes, within the last six months	788	44.6
Yes, more than six months ago	552	31.3
No, I have never tried to quit smoking for more than 24 hours	426	24.1
How likely are you to try to quit smoking within the next six months?		
Not at all	198	11.2
A little	382	21.6
Moderately	508	28.8
Very	308	17.4
Extremely	272	15.4
Don't know	98	5.5
If you decided to quit smoking in the next six months, how sure are you that you would succeed?		
Not at all	147	8.3
A little	346	19.6
Moderately	612	34.7
Very	297	16.8
Extremely	241	13.6
Don't know	123	7.0
Quit approach		
Moderately or less likely to make quit attempt in next six months (unlikely to make a quit attempt in the next six months)	1186	67.2
Very or extremely likely to attempt but moderately or less likely to succeed (unlikely to make a successful quit attempt in the next six months)	304	17.2
Very or extremely likely to attempt and very or extremely likely to succeed (likely to make a successful quit attempt in the next six months)	276	15.6

Table 2: Perceptions of whether inserts would be read, are a good way to provide information, whether they would help smokers to think about quitting or quit, and support for them

	Yes %	No %	Not sure %
Would they be read	50	37	13
Would they be read if interested in quitting	60	25	15
Good way to provide information about quitting	61	25	14

	Agree %	Disagree %	Neither / Don't know %
Make you think more about quitting	53	18	29
Might help you if you decided to quit	52	19	29
Effective way of encouraging smokers to quit	53	17	30
All packs should have inserts	55	20	25

Table 3: Logistic regression models exploring perceptions of inserts by sociodemographic and smoking related characteristics¹

(n=1766)	Would read insert	Would read insert if trying to quit	Inserts make you think about quitting	Inserts might help you quit	Inserts a good way of providing information about quitting	Inserts are an effective way of encouraging smokers to quit	All packs should have inserts
Gender (ref² = male)							
Female	1.24 (1.02 to 1.50)	1.11 (0.91 to 1.35)	0.98 (0.81 to 1.19)	0.95 (0.79 to 1.15)	1.13 (0.93 to 1.37)	0.88 (0.73 to 1.07)	1.20 (0.99 to 1.46)
Age (ref = 16-19)							
20-24	1.16 (0.87 to 1.54)	0.88 (0.66 to 1.18)	1.18 (0.89 to 1.56)	1.19 (0.89 to 1.58)	0.87 (0.65 to 1.16)	0.97 (0.73 to 1.28)	0.96 (0.72 to 1.29)
25-34	1.25 (0.97 to 1.60)	0.83 (0.65 to 1.07)	0.99 (0.78 to 1.26)	1.18 (0.92 to 1.50)	0.76 (0.60 to 0.98)	0.88 (0.69 to 1.12)	0.84 (0.65 to 1.07)
Education (ref = GCSEs (or equivalent) or none)							
More than GCSEs (or equivalent)	1.25 (0.99 to 1.58)	1.12 (0.89 to 1.42)	1.22 (0.97 to 1.54)	1.21 (0.97 to 1.52)	1.12 (0.89 to 1.40)	1.19 (0.95 to 1.50)	1.10 (0.87 to 1.40)
Ethnicity (ref = White British)							
White but not British		0.70 (0.50 to 0.98)	0.58 (0.41 to 0.81)				0.62 (0.44 to 0.87)
Black (inc mixed black & white)		0.92 (0.57 to 1.49)	0.61 (0.38 to 0.98)				0.99 (0.62 to 1.59)
Asian (inc mixed Asian & white)		0.67 (0.49 to 0.92)	1.19 (0.87 to 1.63)				0.70 (0.51 to 0.96)
other or not declared		0.84 (0.50 to 1.42)	1.06 (0.64 to 1.78)				1.08 (0.64 to 1.81)
Dependence (tertiles of HSI) (ref = lower dependence)							
Mid dependence	1.39 (1.11 to 1.76)						1.02 (0.80 to 1.29)
Higher dependence	1.22 (0.94 to 1.59)						0.86 (0.66 to 1.12)
Tobacco products smoked (ref = only factory-made cigarettes)							
Factory-made and roll-your-own	1.35 (1.09 to 1.66)	1.61 (1.30 to 2.00)	1.31 (1.06 to 1.62)	1.31 (1.06 to 1.61)		1.27 (1.03 to 1.56)	
Factory-made and other	1.20 (0.90 to 1.59)	1.39 (1.04 to 1.86)	1.22 (0.92 to 1.63)	1.34 (1.01 to 1.78)		1.20 (0.91 to 1.60)	
Quit attempt lasting at least 24 hours (ref = no)							
Yes, more than six months ago	1.30 (1.00 to 1.69)	1.12 (0.86 to 1.45)	1.20 (0.93 to 1.56)	1.05 (0.81 to 1.36)	1.16 (0.90 to 1.50)	1.07 (0.82 to 1.38)	0.78 (0.60 to 1.01)
Yes, within the last six months	1.67 (1.29 to 2.15)	1.51 (1.17 to 1.94)	1.46 (1.14 to 1.88)	1.35 (1.05 to 1.73)	1.54 (1.20 to 1.98)	1.33 (1.04 to 1.71)	1.06 (0.82 to 1.37)
Efficacy of quit attempt in next 6 months (ref = likely to quit)							
Likely to make unsuccessful attempt	1.01 (0.72 to 1.40)	1.43 (1.00 to 2.06)	0.97 (0.69 to 1.37)	0.92 (0.65 to 1.29)	1.46 (1.02 to 2.08)	1.10 (0.78 to 1.55)	1.43 (1.00 to 2.04)
Unlikely to make attempt	0.58 (0.44 to 0.75)	0.74 (0.55 to 0.99)	0.59 (0.45 to 0.78)	0.51 (0.38 to 0.67)	0.76 (0.57 to 1.01)	0.55 (0.41 to 0.73)	0.56 (0.42 to 0.74)

¹ Blank cells indicate no significant relationship in bivariate analysis; ² Odds ratios for reference categories are always 1.

Table 4: Perceptions of cigarette design (harm, appeal, trial)

	Standard cigarette (SC) %¹	Cigarette with warning (WC) %¹	Green Cigarette (GC) %¹
Harmful to health	38.8	69.1*	70.2*
Think of dangers	20.9	58.1*#	53.5*
Unattractive	25.2	61.7*	68.7*
Unstylish	37.4	66.0*	69.4*
Not nice to be seen with	19.8	55.2*	60.2*
Not appealing to people my age	17.8	51.5*	57.4*
Unlikely to try (personally)	8.9	45.4*	66.5*
Unlikely to try (for never smokers)	14.8	63.3*	71.6*

¹ Percentages shown indicate participants choosing one of the three points nearest the undesirable anchor on a seven-point semantic scale.

* Significant difference in comparison to the standard cigarette (p<0.001)

Significant difference in comparison to the green cigarette (p<0.05)

Table 5: Multilevel and multivariable modelling of perceiving cigarettes as undesirable (n=5298 cigarette evaluations, u=1766 participants)

		<i>Multivariable model</i>	<i>Multivariable model + cigarette*SES interaction</i>		
		<i>Odds ratio (95%CI)</i>	<i>Odds ratio (95%CI)</i>		
	cons	0.05 (0.04 to 0.07)	0.04 (0.03 to 0.06)		
Cigarette type (ref = standard cigarette ¹)	warning on cigarette	17.71 (13.75 to 22.80)	23.29 (16.40 to 33.08)		
	green cigarette	30.88 (23.98 to 39.76)	35.41 (24.93 to 50.29)		
Gender (ref = male)	Female	1.30 (1.10 to 1.54)	1.30 (1.10 to 1.55)		
SES (ref = higher SES)	low education AND/OR low economic status	1.26 (1.06 to 1.50)	1.89 (1.18 to 3.04)		
Ethnicity (ref = White British)	White but not British	0.96 (0.72 to 1.30)	0.96 (0.72 to 1.30)		
	Black (inc mixed black & white)	0.94 (0.62 to 1.42)	0.94 (0.62 to 1.42)		
	Asian (inc mixed Asian & white)	0.79 (0.60 to 1.05)	0.79 (0.60 to 1.05)		
	other or not declared	0.90 (0.58 to 1.42)	0.90 (0.57 to 1.42)		
Product category (ref = Factory-made only)	Factory-made and roll-your-own cigarettes	0.78 (0.65 to 0.90)	0.77 (0.64 to 0.93)		
	Factory-made cigarettes and other tobacco products (e.g. cigars, shisha, etc)	0.73 (0.56 to 0.93)	0.72 (0.56 to 0.93)		
Efficacy (ref = likely to quit)	Not likely to make a quit attempt in next six months	0.62 (0.49 to 0.78)	0.61 (0.49 to 0.78)		
	Likely to make unsuccessful attempt	1.05 (0.78 to 1.41)	1.05 (0.78 to 1.41)		
Interaction Cigarette type * SES (ref = SC*higher SES)	WC*low SES		0.52 (0.31 to 0.87)		
	GC*low SES		0.76 (0.46 to 1.26)		
	<i>Variation between participants (U(std err))</i>	<i>1.14(0.11)</i>	<i>1.14(0.11)</i>		
Models varying reference category of cross classified variable¹					
Reference Category:	SC not low SES	SC low SES	WC not low SES	WC low SES	GC not low SES
Cigarette type & SES					
SC: not low SES	1	0.53 (0.33 to 0.85)	0.04 (0.03 to 0.06)	0.04 (0.03 to 0.06)	0.03 (0.02 to 0.04)
SC: low SES	1.89 (1.18 to 3.03)	1	0.08 (0.06 to 0.12)	0.08 (0.06 to 0.12)	0.05 (0.04 to 0.08)
WC: not low SES	23.13 (16.28 to 32.85)	12.21 (8.48 to 17.58)	1	1.01 (0.80 to 1.28)	0.66 (0.55 to 0.79)
WC: low SES	22.83 (15.58 to 33.46)	12.05 (8.37 to 17.35)	0.99 (0.78 to 1.25)	1	0.65 (0.52 to 0.82)
GC: not low SES	35.09 (24.71 to 49.84)	18.52 (12.86 to 26.67)	1.52 (1.27 to 1.81)	1.54 (1.22 to 1.94)	1
GC: low SES	50.15 (34.29 to 73.35)	26.47 (18.35 to 38.19)	2.17 (1.72 to 2.74)	2.20 (1.74 to 2.77)	1.43 (1.13 to 1.80)

¹Variables included are those in the above models with the exception that cigarette type and SES are replaced with the cross classified variable

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 1: Standard cigarette, warning cigarette and green cigarette

17x5mm (300 x 300 DPI)

For peer review only




Figure 2: Pack inserts highlighting the benefits of quitting or providing tips on how to do so

142x222mm (300 x 300 DPI)

STROBE 2007 (v4) checklist of items to be included in reports of observational studies in epidemiology*
Checklist for cohort, case-control, and cross-sectional studies (combined)

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any pre-specified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	6,7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	8-12
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	9-12
Bias	9	Describe any efforts to address potential sources of bias	6
Study size	10	Explain how the study size was arrived at	6
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	10-13
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	12-13
		(b) Describe any methods used to examine subgroups and interactions	12-13
		(c) Explain how missing data were addressed	7
		(d) <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	NA
		(e) Describe any sensitivity analyses	NA
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	27-28
		(b) Indicate number of participants with missing data for each variable of interest	7
Outcome data	15*	<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	13-17
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	13-17
		(b) Report category boundaries when continuous variables were	30,32

		categorized	
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	12-13,29,31
Discussion			
Key results	18	Summarise key results with reference to study objectives	17-20
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	20
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	20
Generalisability	21	Discuss the generalisability (external validity) of the study results	17-20
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	21

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.