

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [editorial.bmjopen@bmj.com](mailto:editorial.bmjopen@bmj.com)

# BMJ Open

## Improving access to healthcare for women in low-income countries by addressing socio-economic determinants and health insurance: a generalized ordered logistic regression model

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023013
Article Type:	Research
Date Submitted by the Author:	19-Mar-2018
Complete List of Authors:	Bintabara, Deogratus; Tokyo Medical and Dental University, Global Health Entrepreneurship; The University of Dodoma, Public Health Nakamura, Keiko; Tokyo Medical and Dental University, Global Health Entrepreneurship Seino, Kaoruko; Tokyo Medical and Dental University, Global Health Entrepreneurship
Keywords:	HEALTH SERVICES ADMINISTRATION & MANAGEMENT, access to health care, women, Tanzania

SCHOLARONE™  
Manuscripts

Peer Review Only

1  
2  
3 1 **Improving access to healthcare for women in low-income countries by addressing socio-**  
4  
5 2 **economic determinants and health insurance: a generalized ordered logistic regression**  
6  
7 3 **model**  
8  
9

10 4 **Deogratius Bintabara<sup>1,2</sup>, Keiko Nakamura<sup>1</sup>, Kaoruko Seino<sup>1</sup>**  
11  
12

13 5 **Deogratius Bintabara, M.D.**  
14

15 6 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
16  
17 Graduate School of Tokyo Medical and Dental University,  
18  
19 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.  
20  
21  
22

23 9 <sup>2</sup>Department of Public Health, College of Health Sciences,  
24  
25 The University of Dodoma, P.O Box 259, Dodoma, Tanzania  
26  
27 Phone: (255) 767 966 114, Email: bintabaradeo@gmail.com  
28  
29  
30

31 12 **Keiko Nakamura, M.D., Ph.D., (Corresponding author)**  
32

33 13 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
34  
35 Graduate School of Tokyo Medical and Dental University,  
36  
37 Email: nakamura.ith@tmd.ac.jp  
38  
39  
40

41 16 **Kaoruko Seino, Ph.D.**  
42

43 17 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
44  
45 Graduate School of Tokyo Medical and Dental University,  
46  
47 Email: seino.ith@tmd.ac.jp  
48  
49  
50

51 20 **Keywords:** Health services administration and management, Access to healthcare, Women,  
52  
53 Tanzania  
54

55 22 **Word count:** 4830; Abstract 301  
56  
57  
58  
59  
60

## 1 ABSTRACT

2 **Objective:** This study aimed to use generalized ordered logistic regression model to explore the  
3 factors associated with problems in accessing healthcare among women in Tanzania as an  
4 example of low-income countries.

5 **Design:** Population-based cross-sectional survey

6 **Setting:** Nationwide representative data for women of reproductive age obtained from the 2015-  
7 2016 Tanzania Demographic Health Survey were analyzed.

8 **Primary outcome measures:** A composite variable called “problems in accessing healthcare”  
9 with five (0-4) categories was created based on the number of problems reported: getting  
10 permission to go to the doctor, getting money for advice or treatment, distance to a health facility  
11 and not wanting to go alone. Respondents who reported less or more of the problems were  
12 placed in lower or higher categories respectively.

13 **Results:** A total of 13,266 women aged 15-49 years, with a median age (IQR) of 27 (20–36)  
14 years were interviewed and included in this analysis. About two-thirds (65.53%) of the  
15 respondents reported at least one of the four major problems in accessing healthcare.  
16 Furthermore, after controlling for other variables included in the final model, women without  
17 any type of health insurance, those belonged to the poorest class of wealth index, those who had  
18 not attended any type of formal education, those who were not employed for cash, each year of  
19 increased age, and those who were divorced, separated or widowed were associated with greater  
20 problems in accessing healthcare.

1 **Conclusion:** This study found that a high proportion of women face problems in accessing  
2 healthcare in Tanzania. It also highlighted the influence of socio-economic determinants as well  
3 as the role of health insurance in accessing healthcare. The study recommends that, Ministry of  
4 Health, policy-makers, together with other agencies responsible for maternal health services,  
5 provide education for women regarding the importance of health insurance as a first-step in  
6 reducing problems associated with accessing healthcare.

### 7 **Strengths and limitations of this study**

- 8 ➤ This is the first study conducted in this region to explore the factors associated with  
9 problems in accessing healthcare for women using data obtained from a nationally  
10 representative sample with the application of a generalized ordered logistic regression  
11 model.
- 12 ➤ The use of a generalized ordered logistic regression model provided a significantly better  
13 fit to our data while at the same times being much more parsimonious; therefore, it  
14 clearly validated the factors associated with problems in accessing healthcare among  
15 women.
- 16 ➤ A statistical approach was used to adjust for the clustering effect and to weight the  
17 estimates to correct for non-responses and disproportionate sampling used during the  
18 design of the study.
- 19 ➤ The validity constraints of self-reported outcome and independent variables that cannot  
20 be externally validated may have resulted in misclassification bias.
- 21 ➤ As a cross-sectional study design was used, causality assumptions cannot be made;  
22 therefore, the results should be interpreted with caution.

## 1 INTRODUCTION

2 Despite the substantial decline in the global maternal mortality ratio (MMR), this victory  
3 cannot be celebrated in many low-income countries that continue to have high maternal death  
4 rates<sup>1,2</sup> which have been described as an “area of shameful failures of development.”<sup>3,4</sup> In  
5 Tanzania, the current statistics shows that the MMR has increased significantly by more than  
6 20% over the past 5 years,<sup>5,6</sup> despite an increased coverage of maternal health services.<sup>6</sup> The  
7 current high MMR (556 maternal deaths per 100,000 live-births) in Tanzania poses the question  
8 of whether the country can achieve the Sustainable Development Goals (SDGs) target of less  
9 than 140 maternal deaths per 100,000 live-births by the year 2030.<sup>7,8</sup> the majority of avoidable  
10 and unnecessary maternal deaths experienced in this region likely result from poor utilization of  
11 skilled maternal health services.<sup>9</sup>

12 Many social, cultural, geographical factors as well as education level and poverty  
13 reportedly play roles in the poor utilization of health services.<sup>10-12</sup> However, access to healthcare  
14 has been highlighted as the major barrier towards the utilization of maternal health services in  
15 low-income countries, especially in sub-Saharan Africa (SSA).<sup>9,13,14</sup> Access to healthcare can be  
16 broadly defined using four dimensions: availability, affordability, accessibility and  
17 acceptability,<sup>15</sup> but simply referred as the timely use of health services to achieve the desired  
18 health outcomes. Despite the fact that access to health care must be universal and guaranteed for  
19 all on an equitable basis,<sup>16</sup> women continue to face significant inequities in accessing and  
20 utilizing healthcare.<sup>17</sup>

21 Several problems have been mentioned as possible barriers for women in accessing  
22 healthcare,<sup>9</sup> and these can be grouped into two categories: the supply side, in which the facility

1 fails to provide good quality of healthcare and the demand side, in which the clients fail to utilize  
2 the available services because of their own personal reasons.<sup>15,18,19</sup> Despite the fact that it is  
3 important to understand both sides, the current study will focus on the demand side (women), as  
4 this side determines whether or not the healthcare services are used regardless of the presence of  
5 barriers.

6 Identifying the problems that women may experience in accessing healthcare, is  
7 important for addressing the barriers to maternal healthcare in many SSA countries including  
8 Tanzania.<sup>6,20</sup> This has led some studies to assess factors associated with access to healthcare.  
9 However, limited information is available regarding the type and characteristics of women who  
10 are collectively more likely to report having problems associated with access to healthcare in  
11 low-income countries such as Tanzania. Therefore, the current study explored the factors  
12 associated with problems in accessing health care among women in Tanzania.

### 13 **Generalized ordered logistic regression**

14 When the outcome variable has more than two categories that are ordered in nature, the  
15 most appropriate model is the one that can account for the ordering of multiple categories. The  
16 ordered logistic regression (ologit) model is the most commonly fitted model for this type of  
17 variable since it estimates the probability of the outcome belonging to a higher category rather  
18 than a less-than or equal to a given category. In the ologit model, the influence of each  
19 explanatory variable is presumed to be equal across the categories of outcome variable. This  
20 implies that the model provides the same odds ratio (OR) across the categories of outcome  
21 variable, thereby simplifying the interpretations.<sup>21,22</sup> However, the use of this OR across all  
22 categories is appropriate only when the proportional odds (parallel regression) assumption,  
23 which means the “equality of the log-odds across the different categories of the outcome

variable,” is met.<sup>23</sup> However, this assumption is often violated because it is very common for one or more of the coefficients or ORs to differ across the categories of outcome variable.<sup>24</sup> In such cases, it is advisable to use a non-ordinal model, such as multinomial logistic regression (mlogit). Unfortunately, such models are not only less parsimonious and difficult to interpret, compared with ologit models, but they also do not consider the ordinal nature of the variable.<sup>25</sup>

Generalized ordered logistic regression (gologit) established by Fu and later by William has been found to be an appropriate model for such cases<sup>24,26</sup> since it relaxes the proportional assumptions by allowing the effect of each explanatory variable to vary across different categories of outcome variable without modifying the data.<sup>27</sup> The gologit model can be written using the following formula (1):

$$P(Y_i > j) = \frac{\exp(\alpha_j + X_i \beta_j)}{1 + \exp(\alpha_j + X_i \beta_j)}, j = 1, 2, \dots, M - 1 \quad (1)$$

Where M is the number of categories of the ordinal dependent variable (Y), however, the logit model is a special case of the gologit model when M = 2. When M > 2, is equivalent to the series of binary logistic regressions, such as category 1 versus categories 2, 3, 4 and 5 (Y>1); categories 1 and 2 versus categories 3, 4, and 5 (Y>2); categories 1, 2, and 3 versus categories 4 and 5 (Y>3); and categories 1, 2, 3, and 4 versus category 5 (Y>4). Additionally, the ologit model is also a special case of the gologit model when the betas are the same for each value of j as shown in formula (2):

$$P(Y_i > j) = \frac{\exp(\alpha_j + X_i \beta)}{1 + \exp(\alpha_j + X_i \beta)}, j = 1, 2, \dots, M - 1 \quad (2)$$

When the betas change for some variables while for the other variables remain the same, the fitted model is described as being partial constrained, since it only allows the betas of the variables that met the proportional assumptions to be constrained while those not met the



1  
2  
3 1 aforementioned assumptions allowed to vary freely without constraint.<sup>28,29</sup> The gologit2  
4  
5 2 command in Stata is responsible for producing this type of model, and as shown in formula (3)  
6  
7  
8 3 below, the betas for X1 and X2 are constrained but the betas for X3 are not.

$$4 \quad P(Y_i > j) = \frac{\exp(\alpha_j + X1_i\beta1 + X2_i\beta2 + X3_i\beta3_j)}{1 + \exp(\alpha_j + X1_i\beta1 + X2_i\beta2 + X3_i\beta3_j)}, j = 1, 2, \dots, M - 1 \quad (3)$$

5  
6 To the best of our knowledge, there are no published articles that have used a gologit  
7  
8 model to explore factors associated with problems in accessing health care. Therefore the current  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

To the best of our knowledge, there are no published articles that have used a gologit model to explore factors associated with problems in accessing health care. Therefore the current study used gologit regression model to explore the factors associated with problems in accessing healthcare among women in Tanzania as an example of a low-income country.

## 1     **METHODS**

### 2     **Data sources**

3             The current study analyzed data from the 2015-2016 Tanzania Demographic Health  
4     Survey and Malaria Indicator Survey (TDHS-MIS). The survey was conducted by the National  
5     Bureau of Statistics (NBS) and the Office of Chief Government Statistician (OCGS), Zanzibar,  
6     in collaboration with the Ministry of Health, Community Development, Gender, Elderly, and  
7     Children (MoHCDGEC) of the Tanzania Mainland and the Ministry of Health (MoH), Zanzibar.

### 8     **Study design**

9             This study analyzed a nationwide population-based cross-sectional survey that used  
10    information obtained from interviewed women (15-49 years old) who were either residents or  
11    visitors in the household on the night before the survey.

### 12    **Sample size and sampling technique**

13            The 2015-2016 TDHS-MIS used a two stage cluster sampling technique. In the first stage,  
14    sample points (a total of 608 clusters) consisting of enumeration areas delineated for the 2012  
15    Tanzania Population and Housing Census were selected. In the second stage, households were  
16    selected systematically. A complete listing of households was established for all 608 selected  
17    clusters prior to the fieldwork. From this list, 22 households were then systematically selected  
18    from each cluster, yielding a representative probability sample of 13,376 households. Then, all  
19    eligible women and men between the ages of 15-49 years who were either residents or visitors in  
20    the household on the night before the survey were interviewed. Finally, a total of 13,266  
21    women and 3,514 men were interviewed.<sup>6</sup>

## 1 **Data collection and processing**

2 The 2015-2016 TDHS-MIS used four main types of questionnaires during data  
3 collection; however the current study used data collected by Woman's Questionnaire. After pre-  
4 testing of the questionnaires, the finalized and corrected version was used in the main survey  
5 data collection from August 22, 2015, through February 14, 2016. The data collection was  
6 performed by 64 female nurses who were trained and qualified the series of practical tests and  
7 examinations to be interviewers. Following the training, 16 teams were formed (3 for Zanzibar  
8 and 13 for Tanzania Mainland). Data entry was done concurrently with data collection in the  
9 field. After the paper questionnaires were completed, edited, and checked by both the field editor  
10 and the supervisor, the data was entered into a tablet equipped with a data entry programme.  
11 Data entry process included 100% double entry to minimize keying errors, and editing, were  
12 completed on March 21, 2016, while data cleaning and finalization were completed on April 22,  
13 2016.

## 14 **Measurement of variables**

15 **Outcome variable:** In this survey women were asked whether each of the following four  
16 factors was a problem in seeking medical advice or treatment when they were sick: getting  
17 permission to go to the doctor/health facility; getting money for advice, consultation or  
18 treatment; distance to the health facility; and not wanting to go alone. Then, a new composite  
19 variable called "problems in accessing healthcare" was created based on the number of problems  
20 reported with respondents who reported less or more were placed in lower or higher categories,  
21 respectively. The categories were assigned as 1, 2, 3, 4 and 5 for women who reported "no,"  
22 "one," "two," "three," and "four" problems to accessing healthcare, respectively. These

1 categories of a composite (outcome) variable were treated as ordinal numbers, with the  
2 assumption that conceptual differences between categories were exactly the same.

3 **Independent variables:** The current study included several independent variables that  
4 have been empirically and theoretically linked with the accessibility of healthcare among women.  
5 The respondents' ages were categorized into groups of "15-19," "20-34," and "35-49" years;  
6 marital status was grouped into "never married," "married/living together," and "divorced,  
7 separated or widowed." Education level was grouped into "none," "primary," "secondary," and  
8 "highest," (including college and all university level). Employment in the last 12 months was  
9 grouped into "not employed," "employed for cash" and "employed but paid in-kind." The area of  
10 residence was grouped into "urban" and "rural". Health insurance ownership was grouped as "no"  
11 for women who did not have any type of health insurance and "yes" for those who had any type  
12 of health insurance. The wealth index was computed based on household assets and housing  
13 characteristic. During the computation the households were given scores based on the number  
14 and kinds of consumer goods they owned, ranging from a television to a bicycle or car, plus  
15 housing characteristics, such as the source of drinking water, toilet facilities, and flooring  
16 materials. These scores were derived using a principal component analysis. National wealth  
17 quintiles were compiled by assigning the household score to each usual (de jure) household  
18 member, ranking each person in the household population by their score, and then dividing the  
19 distribution into five equal categories, each with 20% of the population, as "poorest, "poorer,"  
20 "middle," "richer", and "richest."<sup>6</sup>

21

## 1 Statistical approaches

2 In descriptive analyses, categorical variables were summarized using proportions and  
3 then presented in tables while quantitative variables were summarized using median and  
4 Interquartile Range (IQR).

5 **Model Fitting:** Since the outcome variable “problems in accessing healthcare” was  
6 ordinal in nature (a score based on the number of different reported problems), in which the  
7 order of its values corresponding to a hierarchy in meaning as in this study, therefore, the  
8 application of ordered logistic regression was recommended.<sup>30</sup> Stata 14 (StataCorp, College  
9 Texas) was used for the analysis in the current study. For all the analyses, the Stata survey set  
10 commands were used to adjust for the variability of clustering and all the estimates were  
11 weighted to correct for non-response and disproportionate sampling.

12 **Proportional ordered logistic model:** As the current study contained several factors to  
13 be considered in the models, a simple (binary) ordered logistics analysis was first performed to  
14 identify variables to be included in the multivariate models. Then, a proportional ordered logistic  
15 regression (ologit) model was used to assess the significance between outcome and independent  
16 variables. Although the ologit model provides results that are straightforward and easily to  
17 interpret the assumptions of a parallel-lines model (parallel regression) must be met for a correct  
18 interpretation. This assumption can be tested using the Brant test, which provides evidence of  
19 whether the assumption is violated.<sup>31</sup> If the Brant test provides a significant *P*-value (less than  
20 0.05), the assumption of parallel regression is violated; hence, the results from this model may  
21 lead to invalid interpretations.<sup>32</sup>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1           **Generalized ordered logistic model:** Since the Brant test command cannot work on  
2 weighted and svyset data like that used in the current analysis, a generalized ordered logistic  
3 (gologit) model was used instead, as this model provides results similar to the series of binary  
4 logistic regressions estimated using the Brant test.<sup>24</sup> The gologit compares higher categories to  
5 categories lower than or equal to the current category. Hence, positive coefficients indicate that  
6 higher values of independent variables make it more likely that the respondent will be classified  
7 in a higher category of the outcome variable (greater difficulties in assessing healthcare) than the  
8 current category. Conversely, negative coefficients indicate that higher values of the independent  
9 variable increase the likelihood of belonging to the current or lower category. However, the  
10 problem with the gologit model is that it frees all the variables from parallel-lines constraints,  
11 even though the assumption may be violated by one or a few of the variables.<sup>24</sup>

12           **Partially generalized ordered logistic model:** the gologit2 with “autofit” option  
13 command can be used to overcome the previous limitation of gologit model by fitting another  
14 model known as the partial proportional odds. This model allows some variables to be modeled  
15 with the proportional odds assumption while the parallel line constraint is relaxed for variables in  
16 which the assumption was not met. The model is less restrictive as it allows the coefficient of the  
17 variables to vary for the different categories that are compared. But, if this is not the case for all  
18 the variables, the model is called a partially constrained logistic model, which is model  
19 recommended for cases involving ordinal data.<sup>28,33</sup> However, the model does not appear to be  
20 parsimonious; therefore, an alternative (gamma) parameterization was performed to make the  
21 model more parsimonious and provide further understanding of the parallel regression  
22 assumptions. The model with gamma parameterization provides beta coefficients that have the  
23 similar coefficients for all pairs of categories of outcome variables. Also, the model provides

1  
2  
3 1 gamma coefficients that show the extent to which the parallel regression assumption is violated  
4  
5 2 by the variable. Hence, if the gamma coefficients for an independent variable are all equal to “0”,  
6  
7 3 then the parallel regression assumption is met for that variable; otherwise the assumption is  
8  
9 4 considered to be violated. If all the gamma coefficients are equal to “0” then the ologit model  
10  
11 5 will be obtained.  
12  
13

## 14 15 6 **Ethics statement**

16  
17  
18 7 This study was based on an analysis of existing public domain survey data sets that are  
19  
20 8 freely available online with all identifier information detached. The survey was approved by the  
21  
22 9 Ethics Committee of the ICF Macro at Calverton in the USA and by the National Institute of  
23  
24 10 Medical Research Ethics Committee in Tanzania. Informed consent was requested and obtained  
25  
26 11 from the participants prior to the interview.  
27  
28  
29

## 30 12 **Patient and Public Involvement Statement**

31  
32 13 Patient and public were not involved in the analysis of this study.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1 RESULTS

### 2 Respondents' characteristics

3 As shown in table 1, a total of 13,266 women between the ages 15-49 years were  
 4 interviewed and included in this analysis. The median age (IQR) of the respondents was 27 (20–  
 5 36) years. Overall, 8,210 respondents (61.89%) were living with their spouse at the time of the  
 6 interview. Only 183 (1.38%) had attained the highest level of education (college or university).  
 7 A total of 6,197 (46.71%) were employed for cash and 8,455 (63.73%) were living in a rural  
 8 residence, but only 1,200 (9.05%) reported having any kind of health insurance and 3,596  
 9 (27.11%) were categorized in the richest quintile. Almost half the respondents (49.49%),  
 10 reported that getting money for healthcare was the greatest problem in accessing healthcare.  
 11 Furthermore, about two-third (65.53%) of the respondents reported at least one of the four  
 12 problems in accessing healthcare.

13 **Table 1 Percent distribution of women between the ages 15-49 by selected background**  
 14 **characteristics, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	Frequency (n=weighted)	Percentage (%=weighted)
<b>Age (Median (IQR)=27, (20-36)</b>		
15-19	2,904	21.89
20-34	6,359	47.94
35-49	4,002	30.17
<b>Marital status</b>		
Never married	3,353	25.27
Married/living together	8,210	61.89
Divorced/separated/widowed	1,703	12.84
<b>Education</b>		
None	1,947	14.67.
Primary	8,211	61.90
Secondary	2,925	22.05



Highest	183	1.38
<b>Employed last 12 months</b>		
Not employed	3,033	22.86
Employed for cash	6,197	46.71
Employed but paid in-kind	4,036	30.43
<b>Residence</b>		
Urban	4,811	36.27
Rural	8,455	63.73
<b>Health insurance</b>		
No	12,066	90.95
Yes	1,200	9.05
<b>Wealth quintile</b>		
Lowest	2,246	16.93
Second	2,274	17.14
Middle	2,328	17.55
Fourth	2,822	21.27
Highest	3,596	27.11
<b>Type of Problems*</b>		
Getting money	6,565	49.49
Distance to facility	5,615	42.33
Not want to go alone	3,962	29.87
Getting permission	1,900	14.32
<b>No. of problems in accessing health</b>		
None	4,574	34.48
One problem	3,291	24.81
Two problems	2,547	19.20
Three problems	1,759	13.26
Four problems	1,095	8.25

Note: \*n and % do not add up to 13,266 and 100 %, respectively, because multiple responses were possible

### Proportional ordered logistic regression (ologit) model

The results from the final ologit model are shown in Table 2. In this model, the number of problems in accessing health care was significantly lower for women who had health insurance [POR=0.622, 95%CI: 0.531-0.728], a primary level of education; [POR=0.888,

1 95%CI: 0.796-0.992], a secondary level of education [POR=0.679, 95%CI: 0.580-0.796] a  
 2 highest level of education [POR= 0.506, 95%CI: 0.351-0.731], a middle wealth status  
 3 [POR=0.737, 95%CI; 0.640-0.848], a richer wealth status [POR=0.512, 95%CI: 0.433-0.604] a  
 4 richest wealth status [POR=0.342, 95%CI: 0.276-0.423], while the number of problems was  
 5 significantly higher for each year increase in age [POR=1.006, 95%CI: 1.001-1.011], for those  
 6 who were divorced, separated or widowed [POR=1.188, 95%CI: 1.025-1.377] and for those who  
 7 were employed but paid in kind [POR=1.221, 95%CI: 1.069-1.394].

8 **Table 2 Results of ologit model using problems in accessing health care as a response with**  
 9 **four categories, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	Coef	Std error	P-value	POR [95% CI]
<b>Health insurance (ref: No)</b>				
Yes	-.475	.081	0.000	0.622 [0.531-0.728]
<b>Residence (ref: Urban)</b>				
Rural	-.149	.0827	0.072	0.862 [0.732-1.013]
<b>Age (as continuous)</b>				
	.006	.002	0.015	1.006 [1.001-1.011]
<b>Marital status ref: Never married)</b>				
Married/living together	-.093	.061	0.127	0.911 [0.809-1.027]
Divorced/separated/widowed	.172	.075	0.022	1.188 [1.025-1.377]
<b>Education (ref: None)</b>				
Primary	-.118	.056	0.035	0.888 [0.796-0.992]
Secondary	-.387	.060	0.000	0.679 [0.580-0.796]
Highest	-.681	.187	0.000	0.506 [0.351-0.731]
<b>Wealth status (ref: Poorest)</b>				
Poorer	-.154	.080	0.055	0.858 [0.733-1.003]
Middle	-.306	.072	0.000	0.737 [0.640-0.848]
Richer	-.670	.085	0.000	0.512 [0.433-0.604]
Richest	-1.074	.109	0.000	0.342 [0.276-0.423]
<b>Employed last 12 months (ref: Not employed)</b>				
Employed for cash	-.014	.060	0.809	0.987 [0.877-1.108]
Employed but paid in kind	.200	.068	0.003	1.221 [1.069-1.394]

## 1 Gologit2 as separate binary logistic regression models

2 Table 3 shows the four separate binary logistic from the gologit2 model that were used to  
 3 assess the parallel regression assumption. The results indicate that the coefficients for all  
 4 categories of each of the independent variables were significant different except for age, wealth  
 5 status (richest), and marital status (divorced/separated/widowed). This means that age, wealth  
 6 and marital status failed to satisfy the parallel regression assumption hence the use of a  
 7 proportion odds ratio was not appropriate for these variables. Since the model frees all the  
 8 variables from parallel-lines constraints, we next used a partial gologit2 model.

9 **Table 3 Estimated coefficients from four binary regression variables of gologit for assessing**  
 10 **the parallel regression assumption, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	y>1	y>2	y>3	y>4
<b>Health insurance</b> (ref: No)				
Yes	-.487	-.484	-.400	-.221
<b>Residence</b> (ref: Urban)				
Rural	-.151	-.098	-.244	-.303
<b>Age</b> (as continuous)	.013	.004	.002	.003
<b>Marital status</b> ref: Never married)				
Married/living together	-.142	-.053	-.110	-.103
Divorced/separated/widowed	.332	.187	.035	.272
<b>Education</b> (ref: None)				
Primary	-.185	-.140	-.103	-.091
Secondary	-.438	-.351	-.444	-.212
Highest	-.736	-.500	-.835	-1.057
<b>Wealth status</b> (ref: Poorest)				
Poorer	-.170	-.239	-.124	.101
Middle	-.359	-.399	-.267	-.029
Richer	-.723	-.764	-.696	-.331
Richest	-1.233	-1.038	-.861	-.469

**Employed last 12 months** (ref: Not employed)

Employed for cash	-.076	-.003	-.076	.124
Employed but paid in kind	.223	.171	.223	.352

**Partial Gologit model with alternative gamma parameterization**

Table 4 shows the results of the partially constrained gologit model with alternative (gamma) parameterization for the outcome variable of problems in accessing healthcare. The results show an insignificant Wald test statistics, indicating that the model does not violate the proportional odds/parallel regression assumptions. However, constraints for parallel lines were not imposed for age, wealth status (richest) and marital status (divorced, separated or widowed). The remaining variables that met the parallel assumption can be interpolated in the same manner as for the ologit model as follows. The odds of reporting problems in accessing healthcare were 38% less among women who had any type of health insurance versus those who did not [POR = 0.622, 95%CI; 0.531-0.731]. Also, the odds of reporting problems in accessing healthcare were 12%, 32% and 48% less among women who attained primary [POR = 0.883, 95%CI; 0.788-0.990], secondary [POR = 0.683, 95%CI; 0.582-0.800], or the highest level of education [POR = 0.516, 95%CI; 0.360-0.741] respectively, versus those who reported not having attended any type of formal education. Additionally, the odds of reporting problems in accessing health care were 15%, 28% and 51% less among women who had a poorer [POR = 0.854, 95%CI; 0.726-1.006], middle [POR = . 0.725, 95%CI; 0.626-0.840], or richer class of wealth status [POR = 0.496, 95%CI; 0.417-0.590] respectively, versus those who were in the poorest class of wealth status. Furthermore, the odds of reporting problems in accessing healthcare were 22% higher among women who were employed but paid in-kind [POR = 1.220, 95%CI; 1.067-1.395], compared with those who were unemployed for the last 12 months before the survey.

1           The variables for which the constraints for parallel lines were not imposed were  
 2 interpreted as follows; the coefficients for age and marital status (divorced, separated or  
 3 widowed) were consistently positive, while those for wealth status (richest) were negative but  
 4 decreased across the cut-points. This means that for each year of increase in age and being  
 5 divorced, separated or widowed, women were more likely to report having a large number of  
 6 problems in accessing healthcare, with the greatest differences being that as the age increased  
 7 and for women who were divorced, separated or widowed, women were less likely to report  
 8 themselves as having few problems in accessing healthcare. Also, the women who were richest  
 9 tend to be less likely to report having many problems in accessing healthcare than the women  
 10 who were poorest, with the greatest differences being that the richest women were less likely to  
 11 report themselves as having many problems in accessing health care.

12 **Table 4 partially constrained gologit2 model with alternative gamma parameterization,**  
 13 **Tanzania DHS-MIS 2015-16 (N=13,266)**

	Variable	Coef	Std error	P-value	POR [95% CI]
<b>Beta</b>	<b>Health insurance</b> (ref: No)				
	Yes	-.474	.081	0.000	0.622 [0.531-0.731]
	<b>Residence</b> (ref: Urban)				
	Rural	-.153	.084	0.072	0.858 [0.728-1.012]
	<b>Age</b> (as continuous)	.011	.003	0.015	1.010 [1.001-1.017]
	<b>Marital status</b> ref: Never married)				
	Married/living together	-.104	.060	0.127	0.901 [0.801-1.014]
	Divorced/separated/widowed	.349	.096	0.022	1.418 [1.175-1.712]
	<b>Education</b> (ref: None)				
	Primary	-.124	.058	0.035	0.883 [0.788-0.990]
Secondary	-.382	.081	0.000	0.683 [0.582-0.800]	
Highest	-.661	.184	0.000	0.516 [0.360-0.741]	

	<b>Wealth status</b> (ref: Poorest)				
	Poorer	-.157	.083	0.055	0.854 [0.726-1.006]
	Middle	-.321	.075	0.000	0.725 [0.626-0.840]
	Richer	-.701	.088	0.000	0.496 [0.417-0.590]
	Richest	-1.234	.114	0.000	0.291 [0.233-0.364]
	<b>Employed last 12 months</b> (ref: Not employed)				
	Employed for cash	-.025	.059	0.809	0.975 [0.869-1.095]
	Employed but paid in-kind	.199	.068	0.003	1.220 [1.067-1.395]
<b>Gamma_2</b>	<b>Age</b>	-.007	.002	0.000	0.993 [0.989-0.998]
	<b>Wealth status</b> (Richest)	.246	.059	0.000	1.279 [1.140-1.435]
	<b>Marital status</b> (Divorced/separated/widowed)	-.206	.076	0.007	0.814 [0.701-0.945]
<b>Gamma_3</b>	<b>Age</b>	.008	.003	0.018	0.993 [0.986-0.999]
	<b>Wealth status</b> (Richest)	.415	.092	0.000	1.515 [1.265-1.814]
	<b>Marital status</b> (Divorced/separated/widowed)	-.288	.093	0.002	0.749 [0.625-0.899]
<b>Gamma_4</b>	<b>Age</b>	-.0131	.005	0.005	0.987 [0.978-0.996]
	<b>Wealth status</b> (Richest)	.671	.133	0.000	1.957 [1.508-2.540]
	<b>Marital status</b> (Divorced/separated /widowed)	-.570	.153	0.000	0.566 [0.419-0.764]

1 Note: Wald test of parallel lines assumption for the final model:  $F(33, 517) = 1.110$ ,  $P = 0.310$ .

2 An insignificant test statistic indicates that the final model does not violate the proportional odds/  
3 parallel lines assumption

4 \*POR= Proportional odds ratio

5

## 1 DISCUSSION

2 The aim of this study was to determine the factors associated with problems in accessing  
3 healthcare among women in Tanzania as an example of low-income country. To best of our  
4 knowledge, this is the first study in this region to explore the factors associated with problems in  
5 accessing healthcare using data obtained from a nationally representative sample with the  
6 application of generalized ordered logistic regression models. The study revealed that about two-  
7 thirds of women reported at least one of four major problems in accessing healthcare.  
8 Furthermore, after controlling for other variables included in the final model, women who did  
9 not have any type of health insurance, those who belonged to the poorest class of wealth index,  
10 those who did not have any type of formal education, those who were employed on a payment  
11 in-kind basis, each year of increased age and those who were divorced, separated or widowed  
12 were associated with greater problems in accessing healthcare.

13 The high proportion of women who reported problems (at least one problem) in accessing  
14 healthcare observed in this study is in agreement with the finding of a study performed in the  
15 Netherlands (69%) that assessed undocumented immigrant women.<sup>34</sup> Despite the Netherlands  
16 being a developed country and the differences in the methodological aspects between these two  
17 studies, the similarities observed might be due to the fact that the study population involved in  
18 the previous study in the Netherlands included only undocumented immigrants who did not have  
19 a residence permit giving them temporary residence. This study population is widely known to  
20 have many problems related in accessing healthcare,<sup>34-37</sup> compared with the general population.

21 Having health insurance is an essential element for timely access to healthcare and better  
22 health-related outcomes.<sup>38,39</sup> The current study found that less than one-tenth of the women

1 reported having any type of health insurance, furthermore, those women who had health  
2 insurance were found to be less likely to report having problems associated with access to  
3 healthcare. This result can be explained by the fact that having any type of health insurance  
4 makes someone not only more comfortable with receiving a wide range of service but also  
5 provides information regarding where and when to obtain healthcare without being afraid of the  
6 cost, which is usually covered by the insurance company. Similar findings have been reported in  
7 study conducted among Hispanics living in El Paso County, Texas.<sup>40</sup>

8 Money is critical to obtain health services such as medical treatment, and its absence may  
9 lead the greatest difficulty in accessing healthcare.<sup>41</sup> The current study found that about half of  
10 the women reported that money was a major problem in accessing healthcare. However, money  
11 and wealth are not synonymous: money can be used to obtain assets to build a household's  
12 wealth. This study found that women who were in the poorest class of the wealth index were  
13 more likely to report having many problems associated with access to healthcare, compared with  
14 women who were in the middle, richer and richest classes. Similar findings have been reported in  
15 a study conducted in Serbia<sup>42</sup> which found that respondents who were in the poorest class of the  
16 wealth index were less likely to access healthcare compared with those in middle, richer and  
17 richest. This finding may be explained by the fact that being in the poorest class requires  
18 individuals to spend their income on basic needs such as food; hence, healthcare costs are  
19 unlikely to be affordable.<sup>43</sup> Therefore, they are more likely to report having many problems in  
20 accessing healthcare.

21 Evidence from several studies shows that unemployment is associated with problems in  
22 accessing healthcare.<sup>44,45</sup> In contrast, to these previous studies and our expectations, the current  
23 study found an unclear relationship between employment status and access to healthcare. The



1 study revealed an insignificant association between unemployment and problems in accessing  
2 healthcare. The difference in findings might be due to differences in socio-cultural and economic  
3 determinants, since the previous studies were conducted in developed countries while the current  
4 study was conducted in a developing country. In developing countries, despite the fact that  
5 someone has employment there are a number of barriers that prevent women from accessing  
6 healthcare such as gender inequality, poor infrastructure, a lack of knowledge regarding  
7 maternal health services<sup>46,47</sup> and socio-cultural aspect such as poor perception toward young or  
8 male physicians.<sup>48</sup> Additionally, in poor resource settings, payment in-kind such as food, clothes,  
9 and other goods instead of cash is still practiced. In agreement with this conclusion, the current  
10 study found that women who were employed but paid in-kind were more likely to report having  
11 greater number of problems in accessing healthcare compared with unemployed women.

12 The current study applied the gologit2 command with an alternative parameterization so  
13 to allow the coefficients of variables that violated the parallel lines assumptions to vary among  
14 the categories of outcome variable. The variables age, wealth status (richest), and marital status  
15 (divorced, separated, or widowed) were found to vary for each category of outcome variable.  
16 However, the findings indicated that for each year of increase in age, women were more likely to  
17 report problems in accessing healthcare. This finding is in agreement to that from a study  
18 conducted in the United States, which found that an older age was associated with barriers in  
19 accessing healthcare.<sup>49</sup> Despite differences in environmental, geographical, cultural, economic,  
20 and study population involved in these studies: previous study included both men and women  
21 who were 65 years or older, while the current study included only women between ages of 15-49  
22 years older, still increase in age is associate with problems in accessing health. This might be due  
23 to that in age is more likely accompanied with decreased in working capability hence low

1 income, be retired and uninsured.<sup>50,51</sup> Also, women who were divorced, separated or widowed,  
2 were more likely to report difficulties in accessing healthcare than those who were never married.  
3 This finding can be explained by the fact that the women who were divorced, separated, or  
4 widowed were more likely to be older than those who had never married or were living with  
5 their partners. As mentioned earlier, older women were more likely to report having a larger  
6 number of problems associated with access to healthcare.

7 The current study subjected to some limitations such as misclassification bias. This might  
8 be introduced due to lack of external validation of self-reported information that could have  
9 affected the categorization of outcome variable. However, we reduce this effect by categorizing  
10 the outcome variable into five groups and used of gologit model that clearly validated the factors  
11 associated with problems in accessing healthcare among women in low-income countries. Also,  
12 being a cross-sectional in nature, the causality assumptions cannot be made; therefore, the results  
13 should be interpreted with caution.

## 14 CONCLUSION

15 Despite the SDG 3 and 5 emphasis on reducing the global maternal mortality ratio to less  
16 than 70 per 100 000 live births by the end of 2030 and to “leave no one behind” as a strategy to  
17 empower women, the current study revealed that a significantly high proportion of women have  
18 difficulty accessing healthcare in Tanzania. The present study also highlighted the influence of  
19 some socio-economic determinants as well as the role of health insurance in the ability of women  
20 to access healthcare. The study suggests that the Ministry of Health together with other agencies  
21 such as Non-Governmental Organizations (NGOs) responsible for maternal health services,  
22 should provide education about the importance of health insurance as a first-step in reducing

1 problems associated with accessing healthcare. Additionally, employment on a payment in-kind  
2 basis should be discouraged, as it seems to hinder women from accessing healthcare.

### 3 **Contributors**

4 DB originated the design of the study, performed statistical analysis, interpretation and  
5 drafted the manuscript. KN contributed to the design of the study and the interpretation advice of  
6 data. KS contributed to interpretation advice of data and drafted the manuscript. Both DB, KN &  
7 KS critically revised the draft manuscript. All authors read and approved the final manuscript.

### 8 **Acknowledgment**

9 We would like to acknowledge ICF International, Rockville, Maryland, USA, through  
10 DHS program for giving us permission to access the TDHS-MIS 2015-2016 dataset.

### 11 **Funding**

12 This research received no specific grant from any funding agency in the public,  
13 commercial or not-for-profit sectors.

### 14 **Competing interest**

15 None to declare.

### 16 **Provenance and peer review**

17 Not commissioned; externally peer reviewed.

### 18 **A data sharing statement**

19 The datasets used for the current analysis was generated from the original survey of  
20 Tanzania DHS-MIS datasets available from within the DHS program repository:  
21 <http://dhsprogram.com/data/available-datasets.cfm>

1  
2  
3 **1 REFERENCES**  
4  
5

- 6 2 1 Alkema L, Chou D, Hogan D, *et al.* Global, regional, and national levels and trends in  
7 maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a  
8 systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*  
9 2016;**387**:462–74.
- 10 6 2 World Health Organization (WHO). Trends in Maternal Mortality: 1990 to 2015:  
11 estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations  
12 Population Division. WHO; 2015. doi:ISBN 978 92 4 150363 1
- 13 9 3 Godefay H, Byass P, Graham WJ, *et al.* Risk Factors for Maternal Mortality in Rural  
14 Tigray, Northern Ethiopia: A Case-Control Study. *PLoS One* 2015;**10**:e0144975.
- 15 11 4 Lewis G. Maternal mortality in the developing world: why do mothers really die? *Obstet*  
16 *Med Med Pregnancy* 2008;**1**:2–6.
- 17 13 5 National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2011. Tanzania  
18 Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro;  
19 2011. [http://www.measuredhs.com/pubs/pdf/FR243/FR243\[24June2011\].pdf](http://www.measuredhs.com/pubs/pdf/FR243/FR243[24June2011].pdf). Accessed 4  
20 Dec 2017.
- 21 17 6 Ministry of Health, Community Development, Gender, Elderly and Children  
22 (MoHCDGEC) [Tanzania Mainland], Ministry of Health (MoH) [Zanzibar], National  
23 Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF.  
2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-  
MIS) 2015-16*. Dar es Salaam, Tanzania, and Rockville, Maryland, USA: MoHCDGEC,  
MoH, NBS, OCGS, and ICF; 2016. <https://dhsprogram.com/pubs/pdf/FR321/FR321.pdf>.  
Accessed 4 Dec 2017.

- 1  
2  
3 1 7 World Health Organization (WHO). Strategies toward ending preventable maternal  
4 mortality ( EPMM ). WHO; 2015. doi:ISBN 978 92 4 150848 3  
5  
6 2  
7  
8 3 8 Moran AC, Jolivet RR, Chou D, *et al.* A common monitoring framework for ending  
9 preventable maternal mortality, 2015–2030: phase I of a multi-step process. *BMC*  
10 *Pregnancy Childbirth* 2016;**16**:250.  
11  
12 5  
13  
14 6 9 Odetola TD. Health care utilization among rural women of child-bearing age: a nigerian  
15 experience. *Pan Afr Med J* 2015;**20**:1–7.  
16  
17 7  
18  
19 8 10 Babalola SO. Factors associated with use of maternal health services in Haiti: a multilevel  
20 analysis. *Rev Panam Salud Publica* 2014;**36**:1–9.  
21  
22 9  
23  
24 10 11 Ngomane S, Mulaudzi FM. Indigenous beliefs and practices that influence the delayed  
25 attendance of antenatal clinics by women in the Bohlabele district in Limpopo, South  
26 Africa. *Midwifery* 2012;**28**:30–8.  
27  
28 12  
29  
30 13 12 Ali HS, Abdalla A, Abdalla A. Understand Factors Influencing Accessibility of Pregnant  
31 Women to Antenatal Care Services Accessibility factors : Demographic characteristics of  
32 the study. *Heal Sci J* 2016;**10**:1–5.  
33  
34 14  
35  
36 15  
37  
38 16 13 Victora CG, Barros AJ, Axelson H, *et al.* How changes in coverage affect equity in  
39 maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of  
40 national surveys. *Lancet* 2012;**380**:1149–56.  
41  
42 18  
43  
44 19 14 Singh PK, Kumar C, Rai RK, *et al.* Factors associated with maternal healthcare services  
45 utilization in nine high focus states in India: a multilevel analysis based on 14 385  
46 communities in 292 districts. *Health Policy Plan* 2014;**29**:542–59.  
47  
48 20  
49  
50 21  
51 22 15 Donnell OO. Access to health care in developing countries : breaking down demand side  
52 barriers. *Cad Saude Publica* 2007;**23**:2820–34.  
53  
54 23  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 16 Kirby N. Access to healthcare services as a human right. *Med Law* 2010;**29**:487–96.
- 4  
5 2 17 Ganle JK, Parker M, Fitzpatrick R, *et al.* Inequities in accessibility to and utilisation of  
6  
7  
8 3 maternal health services in Ghana after user-fee exemption: a descriptive study. *Int J*  
9  
10 4  
11  
12 5 18 Exworthy M, Blane D, Marmot M. Tackling Health Inequalities in the United Kingdom:  
13  
14 6  
15 6 The Progress and Pitfalls of Policy. *Health Serv Res* 2003;**38**:1905–22.
- 16  
17 7 19 Kalu UI, M D-WO, Martha A, *et al.* The Socioeconomic Factors that Determine Women  
18  
19 8  
20 8 Utilization of Healthcare Services in Nigeria. *Int J Asian Soc Sci* 2017;**7**:359–66.
- 21  
22 9 20 Jacobs B, Ir P, Bigdeli M, *et al.* Addressing access barriers to health services: an  
23  
24 10  
25 10 analytical framework for selecting appropriate interventions in low-income Asian  
26  
27 11  
28 12 21 Whitehead J. Sample size calculations for ordered categorical data. *Stat Med*  
29  
30 13  
31 13 1993;**12**:2257–71.
- 32  
33 14 22 Lall R, Campbell MJ, Walters SJ, *et al.* A review of ordinal regression models applied on  
34  
35 15  
36 15 health-related quality of life assessments. *Stat Methods Med Res* 2002;**11**:49–67.
- 37  
38 16 23 Hosmer DW, Lemeshow S. *Applied Logistic Regression*. Hoboken, NJ, USA: John Wiley  
39  
40 17  
41 17 & Sons, Inc. 2000. doi:10.1002/0471722146
- 42  
43 18 24 Williams R. Generalized ordered logit/partial proportional odds models for ordinal  
44  
45 19  
46 19 dependent variables. *Stata J* 2006;**6**:58–82.
- 47  
48 20 25 Agga GE, Scott HM. Use of generalized ordered logistic regression for the analysis of  
49  
50 21  
51 21 multidrug resistance data. *Prev Vet Med* 2015;**121**:374–9.
- 52  
53 22 26 Fu VK. Estimating generalized ordered logit models. *Stata Tech Bull* :1998.  
54  
55 23  
56 23 <https://ideas.repec.org/a/tsj/stbull/y1999v8i44sg88.html#author-body>. Accessed 4 Dec

- 1  
2  
3 1 2017.  
4  
5 2 27 Liu X, Koirala H. Ordinal Regression Analysis: Using Generalized Ordinal Logistic  
6 Regression Models to Estimate Educational Data. *J Mod Appl Stat Methods* 2012;**11**:242–  
7  
8 3 54.  
9  
10 4  
11  
12 5 28 Michalaki P, Quddus MA, Pitfield D, *et al.* Exploring the factors affecting motorway  
13 accident severity in England using the generalised ordered logistic regression model. *J*  
14 *Safety Res* 2015;**55**:89–97.  
15  
16 6  
17 7  
18  
19 8 29 Williams R. Understanding and interpreting generalized ordered logit models. *J Math*  
20 *Sociol* 2016;**40**:7–20.  
21  
22 9  
23  
24 10 30 Quddus MA, Wang C, Ison SG. Road Traffic Congestion and Crash Severity:  
25 Econometric Analysis Using Ordered Response Models. *J Transp Eng* 2010;**136**:424–35.  
26  
27 11  
28 12 31 Brant R. Assessing Proportionality in the Proportional Odds Model for Ordinal Logistic  
29 Regression. *Biometrics* 1990;**46**:1171.  
30  
31 13  
32  
33 14 32 Ananth C V., Kleinbaum DG. Regression models for ordinal responses: A review of  
34 methods and applications. *Int J Epidemiol* 1997;**26**:1323–33.  
35  
36 15  
37  
38 16 33 Eluru N. Evaluating alternate discrete choice frameworks for modeling ordinal discrete  
39 variables. *Accid Anal Prev* 2013;**55**:1–11.  
40  
41 17  
42 18 34 Schoevers MA, Loeffen MJ, van den Muijsenbergh ME, *et al.* Health care utilisation and  
43 problems in accessing health care of female undocumented immigrants in the Netherlands.  
44  
45 19  
46  
47 20  
48  
49 21 35 Torres-Cantero AM, Miguel AG, Gallardo C, *et al.* Health care provision for illegal  
50 migrants: may health policy make a difference? *Eur J Public Health* 2007;**17**:483–5.  
51  
52 22  
53  
54 23 36 Nandi A, Galea S, Lopez G, *et al.* Access to and Use of Health Services Among  
55  
56  
57  
58  
59  
60



- 1  
2  
3 1 Undocumented Mexican Immigrants in a US Urban Area. *Am J Public Health*  
4  
5 2 2008;**98**:2011–20.  
6  
7  
8 3 37 Devillanova C. Social networks, information and health care utilization: Evidence from  
9  
10 4 undocumented immigrants in Milan. *J Health Econ* 2008;**27**:265–86.  
11  
12 5 38 Hsia J, Kemper E, Sofaer S, *et al*. Is Insurance a More Important Determinant of  
13  
14 6 Healthcare Access Than Perceived Health? Evidence from the Women’s Health Initiative.  
15  
16 7 *J Womens Health Gend Based Med* 2000;**9**:881–9.  
17  
18 8 39 Sommers BD, Gawande AA, Baicker K. Health Insurance Coverage and Health — What  
19  
20 9 the Recent Evidence Tells Us. *N Engl J Med* 2017;**377**:586–93.  
21  
22 10 40 Law J, VanDerslice J. Proximal and Distal Determinants of Access to Health Care among  
23  
24 11 Hispanics in El Paso County, Texas. *J Immigr Minor Heal* 2011;**13**:379–84.  
25  
26 12 41 Saha S. ‘More health for the money’: an analytical framework for access to health care  
27  
28 13 through microfinance and savings groups. *Community Dev J* 2014;**49**:618–30.  
29  
30 14 42 Jankovic J, Simic S, Marinkovic J. Inequalities that hurt: demographic, socio-economic  
31  
32 15 and health status inequalities in the utilization of health services in Serbia. *Eur J Public*  
33  
34 16 *Health* 2010;**20**:389–96.  
35  
36 17 43 Yaya S, Bishwajit G, Shah V. Wealth, education and urban–rural inequality and maternal  
37  
38 18 healthcare service usage in Malawi. *BMJ Glob Heal* 2016;**1**:e000085.  
39  
40 19 44 Driscoll AK, Bernstein AB. Health and access to care among employed and unemployed  
41  
42 20 adults: United States, 2009–2010. *NCHS Data Brief* 2012:1–8.  
43  
44 21 45 Maruthappu M, Watson RA, Watkins J, *et al*. Unemployment, public-sector healthcare  
45  
46 22 expenditure and colorectal cancer mortality in the European Union: 1990–2009. *Int J*  
47  
48 23 *Public Health* 2016;**61**:119–30.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 1 46 Bintabara D, Mpembeni RNM, Mohamed AA. Knowledge of obstetric danger signs  
4 among recently-delivered women in Chamwino district, Tanzania: a cross-sectional study.  
5  
6 2  
7  
8 3 *BMC Pregnancy Childbirth* 2017;**17**:276.  
9  
10 4 47 Tsawe M, Susuman A. Determinants of access to and use of maternal health care services  
11 in the Eastern Cape, South Africa: a quantitative and qualitative investigation. *BMC Res*  
12  
13 5  
14 6  
15 6  
16 7 48 Chiang C, Labeeb SA, Higuchi M, *et al.* Barriers to the use of basic health services among  
17 women in rural southern Egypt (Upper Egypt). *Nagoya J Med Sci* 2013;**75**:225–31.  
18  
19 8  
20 9 49 Fitzpatrick AL, Powe NR, Cooper LS, *et al.* Barriers to Health Care Access Among the  
21 Elderly and Who Perceives Them. *Am J Public Health* 2004;**94**:1788–94.  
22  
23 10  
24 11 50 Mold JW, Fryer GE, Thomas CH. Who Are the Uninsured Elderly in the United States? *J*  
25  
26 12  
27 12  
28 13 51 Hadley J, Waidmann T. Health Insurance and Health at Age 65: Implications for Medical  
29  
30  
31  
32  
33 14  
34  
35  
36 15  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Page 2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Page 5
Objectives	3	State specific objectives, including any prespecified hypotheses	Page 7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Page 8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 8-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Page 8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page 9-10
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 9-10
Bias	9	Describe any efforts to address potential sources of bias	Page 3, 8
Study size	10	Explain how the study size was arrived at	Page 8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page 10-12
		(b) Describe any methods used to examine subgroups and interactions	Page 10-12
		(c) Explain how missing data were addressed	N/A
		(d) If applicable, describe analytical methods taking account of sampling strategy	Page 10
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page 13
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page 14 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers of outcome events or summary measures	Page 14 and Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2,3,4
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Table 4
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 18-9, Table 4
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 24
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 24
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 24
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	N/A

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Improving access to healthcare for women in low-income countries by addressing socio-economic determinants and health insurance: a generalized ordered logistic regression model

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023013.R1
Article Type:	Research
Date Submitted by the Author:	11-May-2018
Complete List of Authors:	Bintabara, Deogratus; Tokyo Medical and Dental University, Global Health Entrepreneurship; The University of Dodoma, Public Health Nakamura, Keiko; Tokyo Medical and Dental University, Global Health Entrepreneurship Seino, Kaoruko; Tokyo Medical and Dental University, Global Health Entrepreneurship; Michigan University, Department of Environmental Health Sciences, School of Public Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Global health, Health policy, Health services research
Keywords:	HEALTH SERVICES ADMINISTRATION & MANAGEMENT, access to health care, women, Tanzania

SCHOLARONE™  
Manuscripts

1  
2  
3 1 **Improving access to healthcare for women in low-income countries by addressing socio-**  
4  
5 2 **economic determinants and health insurance: a generalized ordered logistic regression**  
6  
7 3 **model**  
8  
9

10 4 **Deogratius Bintabara<sup>1,2</sup>, Keiko Nakamura<sup>1</sup>, Kaoruko Seino<sup>1,3</sup>**  
11  
12

13 5 **Deogratius Bintabara, M.D.**  
14

15 6 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
16  
17 Graduate School of Tokyo Medical and Dental University,  
18  
19 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.  
20  
21

22 9 <sup>2</sup>Department of Public Health, College of Health Sciences,  
23  
24 The University of Dodoma, P.O Box 259, Dodoma, Tanzania  
25  
26 Phone: (255) 767 966 114, Email: bintabaradeo@gmail.com  
27  
28  
29

30 12 **Keiko Nakamura, M.D., Ph.D., (Corresponding author)**  
31

32 13 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
33  
34 Graduate School of Tokyo Medical and Dental University, Email: nakamura.ith@tmd.ac.jp  
35  
36

37 15 **Kaoruko Seino, Ph.D.**  
38

39 16 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
40  
41 Graduate School of Tokyo Medical and Dental University,  
42  
43 <sup>3</sup>Department of Environmental Health Sciences, School of Public Health,  
44  
45 University of Michigan, Ann Arbor, Michigan USA, Email: seino.ith@tmd.ac.jp  
46  
47  
48

49 20 **Keywords:** Health services administration and management, Access to healthcare, Women,  
50  
51 Tanzania  
52  
53

54 22 **Word count:** 6457; Abstract 277  
55  
56  
57  
58  
59  
60

## 1 ABSTRACT

2 **Objective:** This study aimed to explore the factors associated with accumulation of multiple  
3 problems in accessing healthcare among women in Tanzania as an example of a low-income  
4 country.

5 **Design:** Population-based cross-sectional survey

6 **Setting:** Nationwide representative data for women of reproductive age obtained from the 2015-  
7 2016 Tanzania Demographic Health Survey were analyzed.

8 **Primary outcome measures:** A composite variable called “problems in accessing healthcare”  
9 with five (0-4) categories were created based on the number of problems reported: getting  
10 permission to go to the doctor, getting money for advice or treatment, distance to a health facility  
11 and not wanting to go alone. Respondents who reported less or more of the problems were  
12 placed in lower or higher categories respectively.

13 **Results:** A total of 13,266 women aged 15-49 years, with a median age (IQR) of 27 (20–36)  
14 years were interviewed and included in this analysis. About two-thirds (65.53%) of the  
15 respondents reported at least one of the four major problems in accessing healthcare.  
16 Furthermore, after controlling for other variables included in the final model, women without  
17 any type of health insurance, those who belonged to the poorest class of wealth index, those who  
18 had not attended any type of formal education, those who were not employed for cash, each year  
19 of increased age, and those who were divorced, separated or widowed were associated with  
20 greater problems in accessing healthcare.

1           **Conclusion:** Despite available interventions, failure to address women's problems in a  
2 cumulative manner limits the efforts to eliminate problems in accessing healthcare for women in  
3 low-income countries such as Tanzania. The study recommends improving the uptake of health  
4 insurance and addressing socio-economic determinants as the first-step towards reducing  
5 women's problems associated with accessing healthcare.

### 6 **Strengths and limitations of this study**

- 7           ➤ This is the first study conducted in this region to explore the factors associated with  
8 problems in accessing healthcare for women using data obtained from a nationally  
9 representative sample with the application of a generalized ordered logistic regression  
10 model.
- 11           ➤ The use of a generalized ordered logistic regression model provided a significantly better  
12 fit to our data while at the same times is much more parsimonious; therefore, it clearly  
13 validated the factors associated with problems in accessing healthcare among women.
- 14           ➤ A statistical approach was used to adjust for the clustering effect and to weight the  
15 estimates to correct for non-responses and disproportionate sampling used during the  
16 design of the study.
- 17           ➤ The validity constraints of self-reported outcome and independent variables that cannot  
18 be externally validated may have resulted in misclassification bias.
- 19           ➤ As a cross-sectional study design was used, causality assumptions cannot be made;  
20 therefore, the results should be interpreted with caution.

## 1 INTRODUCTION

2 Despite the substantial decline in the global Maternal Mortality Ratio (MMR), this  
3 victory cannot be celebrated in many low-income countries that continue to have high maternal  
4 death rates<sup>1,2</sup> which have been described as being in an “area of shameful failures of  
5 development.”<sup>3,4</sup> In Tanzania, the current statistics show that the MMR has increased  
6 significantly by more than 20% over the past 5 years, despite the government efforts to  
7 strengthen its health system increasing the coverage of health facilities.<sup>5-7</sup> Through this, each  
8 ward has at least one dispensary and/or health centre, each district has at least one hospital, while  
9 each region has at least one referral hospital. Regardless of the differences in the level of these  
10 facilities in terms of functions, expertise, availability of services, and population coverage all of  
11 them are expected to provide basic maternal health services together with basic emergency  
12 obstetric care.<sup>7</sup> This resulted in an increased coverage of maternal health services such as  
13 antenatal care provided by a skilled provider (96 to 98%), facility delivery (50 to 63%), and  
14 births assisted by skilled providers (51 to 64%) between 2010 to 2016 respectively.<sup>5,6</sup> Therefore,  
15 reported high MMR (556 maternal deaths per 100,000 live-births) in Tanzania poses the question  
16 of whether the country can achieve the Sustainable Development Goals (SDGs) target of less  
17 than 140 maternal deaths per 100,000 live-births by the year 2030.<sup>8,9</sup> the majority of avoidable  
18 and unnecessary maternal deaths experienced in this region likely result from poor utilization of  
19 skilled maternal health services.<sup>10</sup>

20 Many social, cultural, geographical factors as well as education level and poverty  
21 reportedly play roles in the poor utilization of health services.<sup>11-13</sup> However, access to healthcare  
22 has been highlighted as the major barrier towards the utilization of maternal health services in  
23 low-income countries, especially in sub-Saharan Africa (SSA).<sup>10,14,15</sup> Access to healthcare can be



1 broadly defined using four dimensions: availability, affordability, accessibility and  
2 acceptability,<sup>16</sup> but simply to referred as the timely use of health services to achieve the desired  
3 health outcomes. Despite the fact that access to health care must be universal and guaranteed for  
4 all on an equitable basis,<sup>17</sup> women continue to face significant inequities in accessing and  
5 utilizing healthcare.<sup>18</sup>

6 Several problems have been mentioned as possible barriers for women in accessing  
7 healthcare,<sup>10</sup> and these can be grouped into two categories: the supply side, in which the facility  
8 fails to provide good quality of healthcare and the demand side, in which the clients fail to utilize  
9 the available services because of their own personal reasons.<sup>16,19,20</sup> Despite the fact that it is  
10 important to understand both sides, the current study will focus on the demand side (women), as  
11 this side determines whether or not the healthcare services are used regardless of the presence of  
12 barriers. Therefore, it is mainly based on providing insights into problems that women  
13 experienced in accessing healthcare for understanding the challenges embedded in interpersonal  
14 relations.

15 Based on the evidence from previous scholars many problems linked to women on the  
16 demand-side have been reported to be associated with access to healthcare. However, the  
17 following four; getting permission,<sup>21,22</sup> getting money,<sup>23</sup> distance to the health facility,<sup>24</sup> and not  
18 wanting to go alone (lack of spouse or family member escort)<sup>25,26</sup> were the major ones reported  
19 and discussed. Although it is uncertain whether women with multiple problems encounter greater  
20 difficulties in accessing healthcare, most of the previous studies assessed and discussed each of  
21 the four problems independently. Limited evidence of whether women with an accumulation of  
22 problems are more difficulty in accessing healthcare raised a need to create a composite variable  
23 that included all four problems to identify the women in the more disadvantaged group. A similar

1 approach has been used by researchers in another field to assess the severity of problems in  
2 accessing healthcare among individuals with disability in four African countries.<sup>27</sup> Moreover,  
3 evidence from recent studies determined that age, education, residence, health insurance  
4 ownership, wealth status, and occupation are strongly linked with access to healthcare.<sup>28–32</sup> The  
5 current study has also hypothesized that those factors have a relationship with the proposed  
6 composite outcome variable “problems in accessing healthcare.” Therefore, this study used a  
7 generalized ordered logistics regression model to explore the factors associated with  
8 accumulation of multiple problems in accessing healthcare among women in Tanzania as an  
9 example of a low-income country.

## 1 METHODS

### 2 Data sources

3 The current study analyzed data from the 2015-2016 Tanzania Demographic Health  
4 Survey and Malaria Indicator Survey (TDHS-MIS). The survey was conducted by the National  
5 Bureau of Statistics (NBS) and the Office of Chief Government Statistician (OCGS), Zanzibar,  
6 in collaboration with the Ministry of Health, Community Development, Gender, Elderly, and  
7 Children (MoHCDGEC) of the Tanzania Mainland and the Ministry of Health (MoH), Zanzibar.

### 8 Study design

9 This study analyzed a nationwide population-based cross-sectional survey that used  
10 information obtained from interviewed women (15-49 years old) who were either residents or  
11 visitors in the household on the night before the survey.

### 12 Sample size and sampling technique

13 The 2015-2016 TDHS-MIS used a two-stage cluster sampling technique in order to  
14 obtain a sample designed to provide nationally-representative results for all 30 Tanzania regions.  
15 In the first stage, sample points (a total of 608 clusters) consisting of enumeration areas  
16 delineated for the 2012 Tanzania Population and Housing Census were selected. In the second  
17 stage, households were selected systematically. A complete listing of households was established  
18 for all 608 selected clusters prior to the fieldwork. From this list, 22 households were then  
19 systematically selected from each cluster, yielding a representative probability sample of 13,376  
20 households. Then, all eligible women and men between the ages of 15-49 years who were either  
21 residents or visitors in the household on the night before the survey were interviewed. Finally, a  
22 total of 13,266 women and 3,514 men were interviewed.<sup>6</sup>

## 1 **Data collection and processing**

2           The 2015-2016 TDHS-MIS used four main types of questionnaires during data  
3 collection; however, the current study used data collected by the Woman's Questionnaire  
4 segment of the TDHS-MIS. After pre-testing of the questionnaires, the finalized and corrected  
5 version was used in the main survey data collected between August 22, 2015 and February 14,  
6 2016. The data collection was performed by 64 female nurses who were trained and qualified by  
7 a series of practical tests and examinations to be interviewers. Following the training, 16 teams  
8 were formed (3 for Zanzibar and 13 for Tanzania Mainland). The data entry was done  
9 concurrently with data collection in the field. After the paper questionnaires were completed,  
10 edited, and checked by both the field editor and the supervisor, the data was entered into a tablet  
11 equipped with a data entry programme. The data entry process included 100% double entry to  
12 minimize keying errors, and editing was completed on March 21, 2016, while data cleaning and  
13 finalization were completed on April 22, 2016.

## 14 **Measurement of variables**

15           **Outcome variable:** In this survey women were asked whether each of the following four  
16 factors was a problem in seeking medical advice or treatment when they were sick: getting  
17 permission to go to the doctor/health facility; getting money for advice, consultation or  
18 treatment; distance to the health facility; and not wanting to go alone. Then, a new composite  
19 variable called "problems in accessing healthcare" was created based on the number of problems  
20 reported with respondents who reported less or more were placed in lower or higher categories,  
21 respectively. The categories were assigned as 1, 2, 3, 4 and 5 for women who reported "no,"  
22 "one," "two," "three," and "four" problems to accessing healthcare, respectively. These

1 categories of a composite (outcome) variable were treated as ordinal numbers, with the  
2 assumption that conceptual differences between categories were exactly the same.

3 **Independent variables:** The current study included several independent variables that  
4 have been empirically and theoretically linked with the accessibility of healthcare among women.  
5 The respondents' ages were categorized into groups of "15-19," "20-34," and "35-49" years;  
6 marital status was grouped into "never married," "married/living together," and "divorced,  
7 separated or widowed." Education level was grouped into "none," "primary," "secondary," and  
8 "highest," (including college and all university level). Employment in the last 12 months was  
9 grouped into "not employed," "employed for cash" and "employed but paid in-kind." The area of  
10 residence was grouped into "urban" and "rural". Health insurance ownership was grouped as "no"  
11 for women who did not have any type of health insurance and "yes" for those who had any type  
12 of health insurance. The wealth index was computed based on household assets and housing  
13 characteristic. During the computation, the households were given scores based on the number  
14 and kinds of consumer goods they owned, ranging from a television to a bicycle or car, plus  
15 housing characteristics, such as the source of drinking water, toilet facilities, and flooring  
16 materials. These scores were derived using a principal component analysis. National wealth  
17 quintiles were compiled by assigning the household score to each usual (de jure) household  
18 member, ranking each person in the household population by their score, and then dividing the  
19 distribution into five equal categories, each with 20% of the population, as "poorest," "poorer,"  
20 "middle," "richer", and "richest."<sup>6</sup> The selection of these variables was based on studies that  
21 were conducted elsewhere.<sup>28-32</sup>

## 1 Statistical approaches

2 In descriptive analyses, categorical variables were summarized using proportions and  
3 then presented in tables while quantitative variables were summarized using the median and  
4 Interquartile Range (IQR).

5 **Model Fitting:** Since the outcome variable “problems in accessing healthcare” was  
6 ordinal in nature (a score based on the number of different reported problems), in which the  
7 order of its values corresponding to a hierarchy in meaning as in this study, therefore, the  
8 application of ordered logistic regression was recommended.<sup>33</sup> Stata 14 (StataCorp, College  
9 Texas) was used for the analysis in the current study. For all the analyses, the Stata survey set  
10 commands were used to adjust for the variability of clustering and all the estimates were  
11 weighted to correct for non-response and disproportionate sampling.

12 **Proportional ordered logistic model:** As the current study contained several factors to  
13 be considered in the models, a simple (binary) ordered logistics analysis was first performed to  
14 identify variables to be included in the multivariate models. Then, a proportional ordered logistic  
15 regression model was used to assess the significance of outcome and independent variables.  
16 Although the ordered logistic regression model provides results that are straightforward and easy  
17 to interpret the assumptions of a parallel-lines model (parallel regression) must be met for a  
18 correct interpretation.<sup>34–36</sup> This assumption can be tested using the Brant test, which provides  
19 evidence of whether the assumption is violated.<sup>37</sup> If the Brant test provides a significant *P*-value  
20 (less than 0.05), the assumption of parallel regression is violated; hence, the results from this  
21 model may lead to invalid interpretations.<sup>38</sup>

1  
2  
3 1       **Generalized ordered logistic regression model:** Since the Brant test command cannot  
4  
5 2 work on weighted and svyset data like that used in the current analysis, a generalized ordered  
6  
7 3 logistic regression model was used instead, as this model provides results similar to the series of  
8  
9 4 binary logistic regressions estimated using the Brant test.<sup>39</sup> The model compares higher  
10  
11 5 categories to categories lower than or equal to the current category.<sup>40</sup> Hence, positive coefficients  
12  
13 6 indicate that higher values of independent variables make it more likely that the respondent will  
14  
15 7 be classified in a higher category of the outcome variable (greater difficulties in assessing  
16  
17 8 healthcare) than the current category. Conversely, negative coefficients indicate that higher  
18  
19 9 values of the independent variable increase the likelihood of belonging to the current or lower  
20  
21 10 category. However, the problem with the generalized ordered logistic regression model is that it  
22  
23 11 frees all the variables from parallel-lines constraints, even though the assumption may be  
24  
25 12 violated by one or a few of the variables.<sup>39</sup> (see online supplementary table S1)

26  
27  
28  
29  
30  
31 13       **Partially generalized ordered logistic regression model:** When this model has the  
32  
33 14 “autofit” option command applied to it in Stata, it overcomes the previous limitation of the  
34  
35 15 generalized ordered logistic regression model by fitting another model known as the “partial  
36  
37 16 proportional odds.” This model allows some variables to be modeled with the proportional odds  
38  
39 17 assumption while the parallel line constraint is relaxed for variables in which the assumption was  
40  
41 18 not met. The model is less restrictive as it allows the coefficient of the variables to vary for the  
42  
43 19 different categories that are compared.<sup>41</sup> But if this is not the case for all the variables, the model  
44  
45 20 is called a “partially constrained logistic model,” which is a model recommended for cases  
46  
47 21 involving ordinal data.<sup>42,43</sup> However, the model does not appear to be parsimonious; therefore, an  
48  
49 22 alternative (gamma) parameterization was performed to make the model more parsimonious and  
50  
51 23 provide for further understanding of the parallel regression assumptions. The model with gamma  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 parameterization provides beta coefficients that have the similar coefficients for all pairs of  
2 categories of outcome variables. Also, the model provides gamma coefficients that show the  
3 extent to which the parallel regression assumption is violated by the variable. Hence, if the  
4 gamma coefficients for an independent variable are all equal to “0”, then the parallel regression  
5 assumption is met for that variable; otherwise, the assumption is considered to be violated. If all  
6 the gamma coefficients are equal to “0” then the ordered logistic regression model will be  
7 obtained, for more details (see online supplementary word document file S2).

### 8 **Ethics statement**

9 This study was based on an analysis of existing public domain survey data sets that are  
10 freely available online with all identifier information detached. The original TDHS-MIS  
11 protocols were reviewed equivalent by Institution Review Board (IRB) of ICF Macro at  
12 Calverton in the United States (U.S) and by the National Institute of Medical Research (NIMR)  
13 IRB in Tanzania. The ICF IRB ensures that the survey complied with the U.S. Department of  
14 Health and Human Services regulations for the protection of human subjects (45 CFR 46), while  
15 NIMR-IRB ensures that the surveys complied with laws and norms of Tanzania. Before  
16 interviews were performed, informed consent was requested and obtained from the participants.  
17 The participants were adequately informed about all relevant aspects of the survey, including its  
18 objective and interview procedures. All participants accepted to participate in the study signed an  
19 informed consent form.

### 20 **Patient and Public Involvement Statement**

21 Patient and public were not involved in the analysis of this study.



## 1 RESULTS

### 2 Respondents' characteristics

3 As shown in table 1, a total of 13,266 women between the ages 15-49 years were  
 4 interviewed and included in this analysis. The median age (IQR) of the respondents was 27 (20–  
 5 36) years. Overall, 8,210 respondents (61.89%) were living with their spouse at the time of the  
 6 interview. Only 183 (1.38%) had attained the highest level of education (college or university).  
 7 A total of 6,197 (46.71%) were employed for cash and 8,455 (63.73%) were living in a rural  
 8 residence, but only 1,200 (9.05%) reported having any kind of health insurance and 3,596  
 9 (27.11%) were categorized in the richest quintile. Almost half the respondents (49.49%),  
 10 reported that getting money for healthcare was the greatest problem in accessing healthcare.  
 11 Furthermore, about two-thirds (65.53%) of the respondents reported at least one of the four  
 12 problems in accessing healthcare.

13 **Table 1 Percent distribution of women between the ages 15-49 by selected background**  
 14 **characteristics, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	N (%) (Weighted)
<b>Age (Median (IQR)=27, (20-36)</b>	
15-19	2904 (21.89)
20-34	6360 (47.94)
35-49	4002 (30.17)
<b>Marital status</b>	
Never married	3353 (25.27)
Married/living together	8210 (61.89)
Divorced/separated/widowed	1703 (12.84)
<b>Education</b>	
None	1947 (14.67)
Primary	8211 (61.90)
Secondary	2925 (22.05)
Highest	183 (1.38)
<b>Employed last 12 months</b>	
Not employed	3033 (22.86)

Employed cor cash	6197 (46.71)
Employed but paid-in-kind	4036 (30.43)
<b>Residence</b>	
Urban	4811 (36.27)
Rural	8455 (63.73)
<b>Health insurance ownership</b>	
Yes	12066 (90.95)
No	1200 (9.05)
<b>Health quintile</b>	
Lowest	2246 (16.93)
Second	2274 (17.14)
Middle	2328 (17.55)
Fourth	2822 (21.27)
highest	3596 (27.11)
<b>Type of problems*</b>	
Getting money	6565 (49.49)
Distance to facility	5615 (42.33)
Not want to alone	3962 (29.87)
Getting permission	1900 (14.32)
<b>Number of problems in accessing healthcare</b>	
None	4574 (34.48)
One problem	3291 (24.81)
Two problems	2547 (19.20)
Three problems	1759 (13.26)
Four problems	1095 (8.25)

1 Note: \*n and % do not add up to 13,266 and 100 %, respectively, because multiple responses were  
 2 possible

### 3 Proportional ordered logistic regression model

4 The results from the final ordered logistic regression model are shown in Table 2. In this  
 5 model, for women who had any type of health insurance [POR = 0.622, 95%CI; 0.531-0.728],  
 6 the odds of being in higher categories of problems in accessing healthcare versus lower or equal  
 7 to a reference category were 0.622 times lower than those who did not. For each year increase in  
 8 age [POR = 1.006, 95%CI; 1.001-1.011], the odds of being in higher categories of problems in  
 9 accessing healthcare versus lower or equal to a reference category were 1.006 times greater. For  
 10 women who were divorced, separated, or widowed [POR = 1.188, 95%CI; 1.025-1.377], the

odds of being in higher categories of problems in accessing healthcare versus lower or equal to a reference category were 1.188 times greater than those who were never married. For women who attained primary [POR = 0.888, 95%CI; 0.796-0.992], secondary [POR = 0.679, 95%CI; 0.580-0.796], or the highest level of education [POR = 0.506, 95%CI; 0.351-0.731], the odds of being in higher categories of problems in accessing healthcare versus lower or equal to a reference category were 0.888, 0.679, and 0.506 times lower than those who reported not having attended any type of formal education respectively. For women who belonged to the poor [POR = 0.858, 95%CI; 0.733-1.003] middle [POR = 0.737, 95%CI; 0.640-0.848], richer [POR = 0.512, 95%CI; 0.433-0.604], or richest class of wealth status [POR = 0.342, 95%CI; 0.276-0.423], the odds of being in higher categories of problems in accessing healthcare versus lower or equal to a reference category were 0.858, 0.737, 0.512, and 0.342 times lower than those who were in the poorest class respectively.

**Table 2 Results of ordered logistic regression model using problems in accessing health care as a response with four categories, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	POR [95% CI]	P-value
<b>Health insurance</b> (ref: No)		
Yes	0.622 [0.531-0.728]	0.000
<b>Residence</b> (ref: Urban)		
Rural	0.862 [0.732-1.013]	0.072
<b>Age</b> (as continuous)	1.006 [1.001-1.011]	0.015
<b>Marital status</b> ref: Never married)		
Married/living together	0.911 [0.809-1.027]	0.127
Divorced/separated/widowed	1.188 [1.025-1.377]	0.022
<b>Education</b> (ref: None)		
Primary	0.888 [0.796-0.992]	0.035
Secondary	0.679 [0.580-0.796]	0.000
Highest	0.506 [0.351-0.731]	0.000
<b>Wealth status</b> (ref: Poorest)		
Poorer	0.858 [0.733-1.003]	0.055

Middle	0.737 [0.640-0.848]	0.000
Richer	0.512 [0.433-0.604]	0.000
Richest	0.342 [0.276-0.423]	0.000
<b>Employed last 12 months</b> (ref: Not employed)		
Employed for cash	0.987 [0.877-1.108]	0.809
Employed but paid in kind	1.221 [1.069-1.394]	0.003

## 1 The partial generalized ordered logistic regression model with alternative gamma 2 parameterization

3 Table 3 shows the results of the partially constrained generalized ordered logistic  
4 regression model with alternative (gamma) parameterization for the outcome variable of  
5 problems in accessing healthcare. The results show insignificant Wald test statistics, indicating  
6 that the model does not violate the proportional odds/parallel regression assumptions. However,  
7 constraints for parallel lines were not imposed for age, wealth status (richest) and marital status  
8 (divorced, separated or widowed). The remaining variables that met the parallel assumption can  
9 be interpolated in the same manner as for the ordered logistic regression model as follows. For  
10 women who had any type of health insurance [POR = 0.622, 95%CI; 0.531-0.731], the odds of  
11 being in higher categories of problems in accessing healthcare versus lower or equal to a  
12 reference category were 0.622 times lower than those who did not, given that other variables  
13 were held constant in the model. Also, for women who attained primary [POR = 0.883, 95%CI;  
14 0.788-0.990], secondary [POR = 0.683, 95%CI; 0.582-0.800], or the highest level of education  
15 [POR = 0.516, 95%CI; 0.360-0.741], the odds of being in higher categories of problems in  
16 accessing healthcare versus lower or equal to a reference category were 0.883, 0.683, and 0.516  
17 times lower than those who reported not having attended any type of formal education  
18 respectively, given that other variables were held constant in the model. Additionally, for women  
19 who belonged to middle [POR = 0.725, 95%CI; 0.626-0.840] or richer class of wealth status  
20 [POR = 0.496, 95%CI; 0.417-0.590), the odds of being in higher categories of problems in

1 accessing healthcare versus lower or equal to a reference category were 0.725 and 0.496 times  
 2 lower than those who were in the poorest class respectively given that other variables were held  
 3 constant in the model. Furthermore, for women who were employed but paid-in-kind [POR =  
 4 1.220, 95%CI; 1.067-1.395], the odds of being in higher categories of problems in accessing  
 5 healthcare versus lower or equal to a reference category were 1.22 times higher than those who  
 6 were unemployed for the last 12 months before the survey, given that the other variables were  
 7 held constant in the model.

8 The variables for which the constraints for parallel lines were not imposed were  
 9 interpreted as follows; the coefficients for age and marital status (divorced, separated or  
 10 widowed) were consistently positive, while those for wealth status (richest) were negative but  
 11 decreased across the cut-points. This means that for each year of increase in age and being  
 12 divorced, separated or widowed, women were more likely to report having a large number of  
 13 problems in accessing healthcare, with the greatest differences being that as the age increased  
 14 and for women who were divorced, separated or widowed, women were less likely to report  
 15 themselves as having few problems in accessing healthcare. Also, the women who were richest  
 16 tend to be less likely to report having many problems in accessing healthcare than the women  
 17 who were poorest, with the greatest differences being that the richest women were less likely to  
 18 report themselves as having many problems in accessing health care.

19 **Table 3 partially constrained generalized ordered logistic regression model with alternative**  
 20 **gamma parameterization, Tanzania DHS-MIS 2015-16 (N=13,266)**

	<b>Variable</b>	<b>POR [95% CI]</b>	<b>P-value</b>
<b>Beta</b>	<b>Health insurance</b> (ref: No)		
	Yes	0.622 [0.531-0.731]	0.000
	<b>Residence</b> (ref: Urban)		
	Rural	0.858 [0.728-1.012]	0.069

	<b>Age</b> (as continuous)	1.010 [1.001-1.017]	0.000
	<b>Marital status</b> (ref: Never married)		
	Married/living together	0.901 [0.801-1.014]	0.085
	Divorced/separated/widowed	1.418 [1.175-1.712]	0.000
	<b>Education</b> (ref: None)		
	Primary	0.883 [0.788-0.990]	0.033
	Secondary	0.683 [0.582-0.800]	0.000
	Highest	0.516 [0.360-0.741]	0.000
	<b>Wealth status</b> (ref: Poorest)		
	Poorer	0.854 [0.726-1.006]	0.059
	Middle	0.725 [0.626-0.840]	0.000
	Richer	0.496 [0.417-0.590]	0.000
	Richest	0.291 [0.233-0.364]	0.000
	<b>Employed last 12 months</b> (ref: Not employed)		
	Employed for cash	0.975 [0.869-1.095]	0.668
	Employed but paid in kind	1.220 [1.067-1.395]	0.004
Gamma_2	Age	0.993 [0.989-0.998]	0.000
	Wealth status (Richest)	1.279 [1.140-1.435]	0.000
	Marital status (Divorced/separated/widowed)	0.814 [0.701-0.945]	0.007
Gamma_3	Age	0.993 [0.986-0.999]	0.018
	Wealth status (Richest)	1.515 [1.265-1.814]	0.000
	Marital status (Divorced/separated/widowed)	0.749 [0.625-0.899]	0.002
Gamma_4	Age	0.987 [0.978-0.996]	0.005
	Wealth status (Richest)	1.957 [1.508-2.540]	0.000
	Marital status (Divorced/separated/widowed)	0.566 [0.419-0.764]	0.000

Note: Wald test of parallel lines assumption for the final model:  $F(33, 517) = 1.110$ ,  $P = 0.310$ .

An insignificant test statistic indicates that the final model does not violate the proportional odds/parallel lines assumption

\*POR= Proportional odds ratio

## 1 DISCUSSION

2 The aim of this study was to explore the factors associated with accumulation of multiple  
3 problems in accessing healthcare among women in Tanzania as an example of a low-income  
4 country. To best of our knowledge, this is the first study to explore the factors associated with  
5 problems in accessing healthcare by considering the accumulation of multiple problems someone  
6 has experienced. Furthermore, the study used a nationally representative sample from Tanzania  
7 with the application of generalized ordered logistic regression models, that provided the best  
8 models for ordinal data to validate the factors associated with problems in accessing healthcare.  
9 The study revealed that about two-thirds of women reported at least one of four major problems  
10 in accessing healthcare. Furthermore, after controlling for other variables included in the final  
11 model, women who did not have any type of health insurance, those who belonged to the poorest  
12 class of wealth index, those who did not have any type of formal education, those who were  
13 employed on a payment-in-kind basis, each year of increased age and those who were divorced,  
14 separated or widowed were associated with greater problems in accessing healthcare.

15 The high proportion of women who reportedly had problems (at least one problem) in  
16 accessing healthcare observed in this study is in agreement with the finding of a previous study  
17 performed in the Egypt.<sup>44</sup> These findings provide evidence that demand-side barriers such as  
18 cost of care, permission from their spouse, lack of someone to escort, and distance to the  
19 facility,<sup>45,46</sup> still ruin efforts of many African women in accessing healthcare. Because of  
20 cultural, social, and traditional perceptions in Africa that assumes maternal health is only a  
21 woman's responsibility, existing and new interventions should influence health service  
22 utilization to start at individual, households and community level as the one step towards  
23 eliminating demand-side barriers.<sup>23,47</sup>



1  
2  
3 1 Having health insurance is an essential element for timely access to healthcare and better  
4  
5 2 health-related outcomes.<sup>48,49</sup> Despite the availability of National Health Insurance Fund (NHIF)  
6  
7 3 in Tanzania since 2001 through Act No. 8 of 1999, more than 90% of women still are  
8  
9 4 uninsured.<sup>29,50</sup> The current study found that less than one-tenth of the women reported having  
10  
11 5 any type of health insurance. Moreover, those women who had health insurance were found to be  
12  
13 6 less likely to report having the multiple numbers of problems associated with access to  
14  
15 7 healthcare. This result can be explained by the fact that having any type of health insurance  
16  
17 8 makes someone not only more comfortable with receiving a wide range of service but also  
18  
19 9 provides information regarding where and when to obtain healthcare without being afraid of the  
20  
21 10 cost, which is usually covered by the insurance company. A similar finding has been reported in  
22  
23 11 a study conducted Ghana.<sup>28</sup> The similarity of the findings between these studies might be due to  
24  
25 12 both having used secondary data collected by a DHS program that applied similar methodology.  
26  
27 13 Furthermore, study participants in these two studies were from SSA. Therefore, both might have  
28  
29 14 similar socio-economic determinants.

30  
31  
32  
33  
34  
35  
36 15 Money is critical to obtain health services such as medical treatment, and its absence may  
37  
38 16 lead to the greatest difficulty in accessing healthcare.<sup>51</sup> The current study found that about half of  
39  
40 17 the women reported that money was a major problem in accessing healthcare. However, money  
41  
42 18 and wealth are not synonymous: money can be used to obtain assets to build a household's  
43  
44 19 wealth. This study found that women who were in the poorest class of the wealth index were  
45  
46 20 more likely to report having many problems associated with access to healthcare, compared with  
47  
48 21 women who were in the middle, richer and richest classes. Similar findings have been reported in  
49  
50 22 a study conducted in Serbia<sup>52</sup> which found that respondents who were in the poorest class of the  
51  
52 23 wealth index were less likely to access healthcare compared with those in middle, richer and  
53  
54  
55  
56  
57  
58  
59  
60



1 richest. This finding may be explained by the fact that being in the poorest class requires  
2 individuals to spend their income on basic needs such as food; hence, healthcare costs are  
3 unlikely to be affordable.<sup>53</sup> Therefore, they are more likely to report having many problems in  
4 accessing healthcare.

5 Evidence from several studies shows that unemployment is associated with problems in  
6 accessing healthcare.<sup>54,55</sup> In contrast to these previous studies and our expectations, the current  
7 study found an unclear relationship between employment status and access to healthcare. The  
8 study revealed an insignificant association between unemployment and problems in accessing  
9 healthcare. The difference in findings might be due to differences in socio-cultural and economic  
10 determinants since the previous studies were conducted in developed countries while the current  
11 study was conducted in a developing country. In developing countries, despite the fact that  
12 someone has employment there are a number of barriers that prevent women from accessing  
13 healthcare such as gender inequality, poor infrastructure, a lack of knowledge regarding  
14 maternal health services<sup>30,56</sup> and socio-cultural aspect such as poor perception toward young or  
15 male physicians.<sup>44</sup> Additionally, in poor resource settings, payment-in-kind such as food, clothes,  
16 and other goods instead of cash is still practiced. In agreement with this conclusion, the current  
17 study found that women who were employed but paid-in-kind were more likely to report having  
18 the greater number of problems in accessing healthcare compared with unemployed women.

19 The current study applied the generalized ordered logistics regression with an alternative  
20 parameterization so to allow the coefficients of variables that violated the parallel lines  
21 assumptions to vary among the categories of the outcome variable. The variables age, wealth  
22 status (richest), and marital status (divorced, separated, or widowed) were found to vary for each  
23 category of the outcome variable. However, the findings indicated that for each year of increase

1 in age, women were more likely to report problems in accessing healthcare. This finding is in  
2 agreement to that from the study conducted in the Ethiopia, which found that an older age was  
3 associated with problems in accessing maternal healthcare.<sup>57,58</sup> This might be due to that increase  
4 in age is more likely accompanied by a decrease in working capability (hence low income),  
5 being retired and being uninsured.<sup>59,60</sup> Also, women who were divorced, separated or widowed,  
6 were more likely to report difficulties in accessing healthcare than those who were never married.  
7 This finding can be explained by the fact that the women who were divorced, separated, or  
8 widowed were more likely to be older than those who had never married or were living with  
9 their partners. As mentioned earlier, older women were more likely to report having a larger  
10 number of problems associated with access to healthcare.

11 The current study was subjected to some limitations such as misclassification bias. This  
12 might be introduced due to lack of external validation of self-reported information that could  
13 affect the categorization of the outcome variable. However, we reduced this effect by  
14 categorizing the outcome variable into five groups and by the use of a generalized ordered  
15 logistics regression model that clearly validated the factors associated with problems in accessing  
16 healthcare among women in low-income countries. Also, being cross-sectional in nature, the  
17 causality assumptions cannot be made; therefore, the results should be interpreted with caution.

## 18 CONCLUSION

19 Despite several interventions that have been placed to increase women accessibility to  
20 healthcare, failure to address women's problems in a cumulative manner starting at individual,  
21 households and community levels limit the efforts to remove the demand-side barrier to  
22 accessing healthcare for women in low-income countries such as Tanzania. The study suggests  
23 that the Tanzanian government together with other agencies responsible for maternal health

1 services should emphasize that women be enrolled in any type of health insurance and that the  
2 addressing socio-economic determinants through health education interventions be the first-step  
3 forward to reducing problems associated with accessing healthcare.

#### 4 **Contributors**

5 DB originated the design of the study, performed statistical analysis, interpretation and  
6 drafted the manuscript. KN contributed to the design of the study and the interpretation advice of  
7 data. KS contributed to interpretation advice of data and drafted the manuscript. Both DB, KN &  
8 KS critically revised the draft manuscript. All authors read and approved the final manuscript.

#### 9 **Acknowledgment**

10 We would like to acknowledge ICF International, Rockville, Maryland, USA, through the  
11 DHS program for giving us permission to access the TDHS-MIS 2015-2016 dataset.

#### 12 **Funding**

13 This research received no specific grant from any funding agency in the public,  
14 commercial or not-for-profit sectors.

#### 15 **Competing interest**

16 None to declare.

#### 17 **Provenance and peer review**

18 Not commissioned; externally peer reviewed.

#### 19 **A data sharing statement**

1           The datasets used for the current analysis was generated from the original survey of  
2  
3  
4  
5           Tanzania DHS-MIS datasets available from within the DHS program repository:  
6  
7  
8           <http://dhsprogram.com/data/available-datasets.cfm>  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## 1 REFERENCES

- 2 1 Alkema L, Chou D, Hogan D, *et al.* Global, regional, and national levels and trends in  
3 maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a  
4 systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*  
5 2016;**387**:462–74. doi:10.1016/S0140-6736(15)00838-7
- 6 2 World Health Organization (WHO). Trends in Maternal Mortality: 1990 to 2015:  
7 estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations  
8 Population Division. 2015. doi:ISBN 978 92 4 150363 1
- 9 3 Godefay H, Byass P, Graham WJ, *et al.* Risk Factors for Maternal Mortality in Rural  
10 Tigray, Northern Ethiopia: A Case-Control Study. *PLoS One* 2015;**10**:e0144975.  
11 doi:10.1371/journal.pone.0144975
- 12 4 Lewis G. Maternal mortality in the developing world: why do mothers really die? *Obstet*  
13 *Med Med Pregnancy* 2008;**1**:2–6. doi:10.1258/om.2008.080019
- 14 5 National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2011. Tanzania  
15 Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro;  
16 2011. [http://www.measuredhs.com/pubs/pdf/FR243/FR243\[24June2011\].pdf](http://www.measuredhs.com/pubs/pdf/FR243/FR243[24June2011].pdf). Accessed 6  
17 May 2018.
- 18 6 Ministry of Health, Community Development, Gender, Elderly and Children  
19 (MoHCDGEC) [Tanzania Mainland], Ministry of Health (MoH) [Zanzibar], National  
20 Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF.  
21 2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-*  
22 *MIS) 2015-16*. Dar es Salaam, Tanzania, and Rockville, Maryland, USA: MoHCDGEC,  
23 MoH, NBS, OCGS, and ICF; 2016. <https://dhsprogram.com/pubs/pdf/FR321/FR321.pdf>.

- 1  
2  
3 1 Accessed 6 May 2018.  
4  
5 2 7 United Republic of Tanzania (URT) and World Health Organization (WHO). Health  
6  
7 Systems Profile in Tanzania. URT; 2004.  
8 3  
9  
10 4 <http://mbaralidc.go.tz/storage/app/media/uploaded-files/Health-tanzania.pdf> Accessed 6  
11  
12 5 May 2018.  
13  
14 6 8 World Health Organization (WHO). Strategies toward ending preventable maternal  
15  
16 mortality ( EPMM ). Published Online First: 2015. doi:ISBN 978 92 4 150848 3  
17 7  
18 8 9 Moran AC, Jolivet RR, Chou D, *et al.* A common monitoring framework for ending  
19  
20 preventable maternal mortality, 2015–2030: phase I of a multi-step process. *BMC*  
21  
22 *Pregnancy Childbirth* 2016;**16**:250. doi:10.1186/s12884-016-1035-4  
23  
24 10  
25 11 10 Odetola TD. Health care utilization among rural women of child-bearing age: a nigerian  
26  
27 experience. *Pan Afr Med J* 2015;**20**:1–7. doi:10.11604/pamj.2015.20.151.5845  
28  
29 12  
30 13 11 Babalola SO. Factors associated with use of maternal health services in Haiti: a multilevel  
31  
32 analysis. *Rev Panam Salud Publica* 2014;**36**:1–9.  
33  
34 14  
35 15 12 Ngomane S, Mulaudzi FM. Indigenous beliefs and practices that influence the delayed  
36  
37 attendance of antenatal clinics by women in the Bohlabele district in Limpopo, South  
38  
39 Africa. *Midwifery* 2012;**28**:30–8. doi:10.1016/j.midw.2010.11.002  
40  
41 17  
42 18 13 Ali HS, Abdalla A, Abdalla A. Understand Factors Influencing Accessibility of Pregnant  
43  
44 Women to Antenatal Care Services Accessibility factors : Demographic characteristics of  
45  
46 the study. *Heal Sci J* 2016;**10**:1–5. doi:10.4172/1791-809X.1000100507  
47  
48 20  
49 21 14 Victora CG, Barros AJ, Axelson H, *et al.* How changes in coverage affect equity in  
50  
51 maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of  
52  
53 national surveys. *Lancet* 2012;**380**:1149–56. doi:10.1016/S0140-6736(12)61427-5  
54  
55 23  
56  
57  
58  
59  
60

- 1  
2  
3 1 15 Singh PK, Kumar C, Rai RK, *et al.* Factors associated with maternal healthcare services  
4 utilization in nine high focus states in India: a multilevel analysis based on 14 385  
5  
6 2  
7 communities in 292 districts. *Health Policy Plan* 2014;**29**:542–59.  
8  
9 doi:10.1093/heapol/czt039  
10  
11 4  
12 5 16 Donnell OO. Access to health care in developing countries : breaking down demand side  
13 barriers. *Cad Saude Publica* 2007;**23**:2820–34. doi:S0102-311X2007001200003 [pii]  
14  
15 6  
16  
17 7 17 Kirby N. Access to healthcare services as a human right. *Med Law* 2010;**29**:487–96.  
18  
19 8 18 Ganle JK, Parker M, Fitzpatrick R, *et al.* Inequities in accessibility to and utilisation of  
20 maternal health services in Ghana after user-fee exemption: a descriptive study. *Int J*  
21  
22 9  
23 *Equity Health* 2014;**13**:89. doi:10.1186/s12939-014-0089-z  
24  
25 10  
26 11 19 Exworthy M, Blane D, Marmot M. Tackling Health Inequalities in the United Kingdom:  
27 The Progress and Pitfalls of Policy. *Health Serv Res* 2003;**38**:1905–22.  
28  
29 12  
30 doi:10.1111/j.1475-6773.2003.00208.x  
31  
32 13  
33 14 20 Kalu UI, M D-WO, Martha A, *et al.* The Socioeconomic Factors that Determine Women  
34 Utilization of Healthcare Services in Nigeria. *Int J Asian Soc Sci* 2017;**7**:359–66.  
35  
36 15  
37 doi:10.18488/journal.1/2017.7.5/1.5.359.366  
38  
39 16  
40 17 21 Danforth E, Kruk M, Rockers P, *et al.* Household Decision-making about Delivery in  
41 Health Facilities: Evidence from Tanzania. *J Heal Popul Nutr* 2009;**27**:696–703.  
42  
43 18  
44 doi:10.3329/jhpn.v27i5.3781  
45  
46 19  
47 20 22 Wilunda C, Scanagatta C, Putoto G, *et al.* Barriers to utilisation of antenatal care services  
48 in South Sudan: a qualitative study in Rumbek North County. *Reprod Health* 2017;**14**:65.  
49  
50 21  
51 doi:10.1186/s12978-017-0327-0  
52  
53 22  
54 23 23 Lowe M, Chen D-R, Huang S-L. Social and Cultural Factors Affecting Maternal Health in

- 1  
2  
3 1 Rural Gambia: An Exploratory Qualitative Study. *PLoS One* 2016;**11**:e0163653.  
4  
5 2 doi:10.1371/journal.pone.0163653  
6  
7  
8 3 24 Mbiza CR, Kazembe A, Simwaka A. Barriers to health-seeking practices during  
9  
10 4 pregnancy among adolescents in rural Blantyre, Malawi. *Afr J Midwifery Womens Health*  
11  
12 5 2014;**8**:59–65. doi:10.12968/ajmw.2014.8.2.59  
13  
14  
15 6 25 Kumbani L, Bjune G, Chirwa E, *et al.* Why some women fail to give birth at health  
16  
17 7 facilities: a qualitative study of women’s perceptions of perinatal care from rural Southern  
18  
19 8 Malawi. *Reprod Health* 2013;**10**:9. doi:10.1186/1742-4755-10-9  
20  
21  
22 9 26 Vidler M, Ramadurg U, Charantimath U, *et al.* Utilization of maternal health care services  
23  
24 10 and their determinants in Karnataka State, India. *Reprod Health* 2016;**13**:37.  
25  
26 11 doi:10.1186/s12978-016-0138-8  
27  
28  
29 12 27 Eide AH, Mannan H, Khogali M, *et al.* Perceived Barriers for Accessing Health Services  
30  
31 13 among Individuals with Disability in Four African Countries. *PLoS One*  
32  
33 14 2015;**10**:e0125915. doi:10.1371/journal.pone.0125915  
34  
35  
36 15 28 Browne JL, Kayode GA, Arhinful D, *et al.* Health insurance determines antenatal,  
37  
38 16 delivery and postnatal care utilisation: evidence from the Ghana Demographic and Health  
39  
40 17 Surveillance data. *BMJ Open* 2016;**6**:e008175. doi:10.1136/bmjopen-2015-008175  
41  
42  
43 18 29 Kibusi SM, Sunguya BF, Kimunai E, *et al.* Health insurance is important in improving  
44  
45 19 maternal health service utilization in Tanzania—analysis of the 2011/2012 Tanzania  
46  
47 20 HIV/AIDS and malaria indicator survey. *BMC Health Serv Res* 2018;**18**:112.  
48  
49 21 doi:10.1186/s12913-018-2924-1  
50  
51  
52 22 30 Tsawe M, Susuman A. Determinants of access to and use of maternal health care services  
53  
54 23 in the Eastern Cape, South Africa: a quantitative and qualitative investigation. *BMC Res*



- 1  
2  
3 1 *Notes* 2014;**7**:723. doi:10.1186/1756-0500-7-723  
4  
5 2 31 Mrisho M, Schellenberg JA, Mushi AK, *et al.* Factors affecting home delivery in rural  
6  
7 Tanzania. *Trop Med Int Heal* 2007;**12**:862–72. doi:10.1111/j.1365-3156.2007.01855.x  
8  
9 3  
10 4 32 Kyei-Nimakoh M, Carolan-Olah M, McCann T V. Access barriers to obstetric care at  
11  
12 health facilities in sub-Saharan Africa—a systematic review. *Syst Rev* 2017;**6**:110.  
13  
14 doi:10.1186/s13643-017-0503-x  
15  
16 6  
17 7 33 Quddus MA, Wang C, Ison SG. Road Traffic Congestion and Crash Severity:  
18  
19 Econometric Analysis Using Ordered Response Models. *J Transp Eng* 2010;**136**:424–35.  
20  
21 doi:10.1061/(ASCE)TE.1943-5436.0000044  
22  
23 9  
24 10 34 Whitehead J. Sample size calculations for ordered categorical data. *Stat Med*  
25  
26 1993;**12**:2257–71. doi:10.1002/sim.4780122404  
27  
28 12 35 Lall R, Campbell MJ, Walters SJ, *et al.* A review of ordinal regression models applied on  
29  
30 health-related quality of life assessments. *Stat Methods Med Res* 2002;**11**:49–67.  
31  
32 doi:10.1191/0962280202sm271ra  
33  
34 14  
35 15 36 Hosmer DW, Lemeshow S. *Applied Logistic Regression*. Hoboken, NJ, USA: : John  
36  
37 Wiley & Sons, Inc. 2000. doi:10.1002/0471722146  
38  
39 16  
40 17 37 Brant R. Assessing Proportionality in the Proportional Odds Model for Ordinal Logistic  
41  
42 Regression. *Biometrics* 1990;**46**:1171. doi:10.2307/2532457  
43  
44 18  
45 19 38 Ananth C V., Kleinbaum DG. Regression models for ordinal responses: A review of  
46  
47 20 methods and applications. *Int J Epidemiol* 1997;**26**:1323–33. doi:10.1093/ije/26.6.1323  
48  
49 21 39 Williams R. Generalized ordered logit/partial proportional odds models for ordinal  
50  
51 22 dependent variables. *Stata J* 2006;**6**:58–82. doi:st0097  
52  
53 23 40 Liu X, Koirala H. Ordinal Regression Analysis: Using Generalized Ordinal Logistic

- 1  
2  
3 1 Regression Models to Estimate Educational Data. *J Mod Appl Stat Methods* 2012;**11**:242–  
4  
5 2 54. doi:10.22237/jmasm/1335846000  
6  
7  
8 3 41 Williams R. Understanding and interpreting generalized ordered logit models. *J Math*  
9  
10 4 *Sociol* 2016;**40**:7–20. doi:10.1080/0022250X.2015.1112384  
11  
12 5 42 Eluru N. Evaluating alternate discrete choice frameworks for modeling ordinal discrete  
13  
14 6 variables. *Accid Anal Prev* 2013;**55**:1–11. doi:10.1016/j.aap.2013.02.012  
15  
16  
17 7 43 Michalaki P, Quddus MA, Pitfield D, *et al.* Exploring the factors affecting motorway  
18  
19 8 accident severity in England using the generalised ordered logistic regression model. *J*  
20  
21 9 *Safety Res* 2015;**55**:89–97. doi:10.1016/j.jsr.2015.09.004  
22  
23  
24 10 44 Chiang C, Labeeb SA, Higuchi M, *et al.* Barriers to the use of basic health services among  
25  
26 11 women in rural southern Egypt (Upper Egypt). *Nagoya J Med Sci* 2013;**75**:225–31.  
27  
28 12 45 Mselle LT, Kohi TW. Healthcare access and quality of birth care: narratives of women  
29  
30 13 living with obstetric fistula in rural Tanzania. *Reprod Health* 2016;**13**:87.  
31  
32 14 doi:10.1186/s12978-016-0189-x  
33  
34  
35 15 46 Shartz A, Long SK, Benatar S. Health Care Costs Are a Barrier to Care for Many  
36  
37 16 Women. Urban Inst. 2015.[http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf)  
38  
39 17 [to-Care-for-Many-Women.pdf](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf) (accessed 5 May 2018).  
40  
41  
42 18 47 Azuh D, Fayomi O, Ajayi, Lady. Socio-Cultural Factors of Gender Roles in Women's  
43  
44 19 Healthcare Utilization in Southwest Nigeria. *Open J Soc Sci* 2015;**3**:105–17.  
45  
46 20 doi:10.4236/jss.2015.34013  
47  
48  
49 21 48 Hsia J, Kemper E, Sofaer S, *et al.* Is Insurance a More Important Determinant of  
50  
51 22 Healthcare Access Than Perceived Health? Evidence from the Women's Health Initiative.  
52  
53 23 *J Womens Health Gen Based Med* 2000;**9**:881–9. doi:10.1089/152460900750020919  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 49 Sommers BD, Gawande AA, Baicker K. Health Insurance Coverage and Health — What  
4 the Recent Evidence Tells Us. *N Engl J Med* 2017;**377**:586–93.  
5  
6 2  
7  
8 3 doi:10.1056/NEJMs1706645  
9  
10 4 50 Marwa CW. Provision of National Health Insurance Fund Services to Its Members ; Pain  
11 or Gain ? *Unified J Sport Heal Sci* 2016;**2**:1–6.  
12  
13 5  
14 6 51 Saha S. ‘More health for the money’: an analytical framework for access to health care  
15 through microfinance and savings groups. *Community Dev J* 2014;**49**:618–30.  
16  
17 7  
18 8 doi:10.1093/cdj/bsu037  
19  
20 9 52 Jankovic J, Simic S, Marinkovic J. Inequalities that hurt: demographic, socio-economic  
21 and health status inequalities in the utilization of health services in Serbia. *Eur J Public*  
22  
23 10  
24 11 *Health* 2010;**20**:389–96. doi:10.1093/eurpub/ckp189  
25  
26 12 53 Yaya S, Bishwajit G, Shah V. Wealth, education and urban–rural inequality and maternal  
27 healthcare service usage in Malawi. *BMJ Glob Heal* 2016;**1**:e000085. doi:10.1136/bmjgh-  
28  
29 13 2016-000085  
30  
31 14  
32 15 54 Driscoll AK, Bernstein AB. Health and access to care among employed and unemployed  
33 adults: United States, 2009–2010. *NCHS Data Brief* 2012;**1**:1–8.  
34  
35 16  
36 17 55 Maruthappu M, Watson RA, Watkins J, *et al.* Unemployment, public-sector healthcare  
37 expenditure and colorectal cancer mortality in the European Union: 1990–2009. *Int J*  
38  
39 18  
40 19 *Public Health* 2016;**61**:119–30. doi:10.1007/s00038-015-0727-2  
41  
42 20 56 Bintabara D, Mpembeni RNM, Mohamed AA. Knowledge of obstetric danger signs  
43 among recently-delivered women in Chamwino district, Tanzania: a cross-sectional study.  
44  
45 21  
46 22 *BMC Pregnancy Childbirth* 2017;**17**:276. doi:10.1186/s12884-017-1469-3  
47  
48 23 57 Feyissa TR, Genemo GA. Determinants of Institutional Delivery among Childbearing Age  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Women in Western Ethiopia, 2013: Unmatched Case Control Study. *PLoS One*  
4  
5 2014;**9**:e97194. doi:10.1371/journal.pone.0097194  
6  
7  
8 3 58 Wilunda C, Quaglio G, Putoto G, *et al*. Determinants of utilisation of antenatal care and  
9  
10 4 skilled birth attendant at delivery in South West Shoa Zone, Ethiopia: a cross sectional  
11  
12 5 study. *Reprod Health* 2015;**12**:74. doi:10.1186/s12978-015-0067-y  
13  
14  
15 6 59 Mold JW, Fryer GE, Thomas CH. Who Are the Uninsured Elderly in the United States? *J*  
16  
17 7 *Am Geriatr Soc* 2004;**52**:601–6. doi:10.1111/j.1532-5415.2004.52169.x  
18  
19 8 60 Hadley J, Waidmann T. Health Insurance and Health at Age 65: Implications for Medical  
20  
21 9 Care Spending on New Medicare Beneficiaries. *Health Serv Res* 2006;**41**:429–51.  
22  
23 10 doi:10.1111/j.1475-6773.2005.00491.x  
24  
25  
26 11

**Supplementary Table S1: Estimated coefficients from four binary regression variables of generalized ordered logistics regressions for assessing the parallel regression assumption, Tanzania DHS-MIS 2015-16 (N=13,266)**

Variable	y>1	y>2	y>3	y>4
<b>Health insurance</b> (ref: No)				
Yes	-.487	-.484	-.400	-.221
<b>Residence</b> (ref: Urban)				
Rural	-.151	-.098	-.244	-.303
<b>Age</b> (as continuous)	.013	.004	.002	.003
<b>Marital status</b> ref: Never married)				
Married/living together	-.142	-.053	-.110	-.103
Divorced/separated/widowed	.332	.187	.035	.272
<b>Education</b> (ref: None)				
Primary	-.185	-.140	-.103	-.091
Secondary	-.438	-.351	-.444	-.212
Highest	-.736	-.500	-.835	-1.057
<b>Wealth status</b> (ref: Poorest)				
Poorer	-.170	-.239	-.124	.101
Middle	-.359	-.399	-.267	-.029
Richer	-.723	-.764	-.696	-.331
Richest	-1.233	-1.038	-.861	-.469
<b>Employed last 12 months</b> (ref: Not employed)				
Employed for cash	-.076	-.003	-.076	.124
Employed but paid in kind	.223	.171	.223	.352

Note: The Table presents four separate binary logistic from the generalized ordered logistics regressions model that were used to assess the parallel regression assumption. The results indicate that the coefficients for all categories of each of the independent variables were significant different except for age, wealth status (richest), and marital status (divorced/separated/widowed). This means that age, wealth and marital status failed to satisfy the parallel regression assumption hence the use of a proportion odds ratio was not appropriate for these variables. Since the model frees all the variables from parallel-lines constraints, the partial generalized ordered logistics model were the right model for this case.

1  
2  
3 **Supplementary document S2: Explain in details about the generalized logistic regression**  
4 **model that was used in the analysis of the current study.**  
5  
6

7  
8 **Generalized ordered logistic regression model**  
9

10 When the outcome variable has more than two categories that are ordered in nature, the  
11 most appropriate model is the one that can account for the ordering of multiple categories. The  
12 ordered logistic regression model is the most commonly fitted model for this type of variable  
13 since it estimates the probability of the outcome belonging to a higher category rather than a less-  
14 than or equal to a given category. In the ordered logistic regression model, the influence of each  
15 explanatory variable is presumed to be equal across the categories of outcome variable. This  
16 implies that the model provides the same odds ratio (OR) across the categories of outcome  
17 variable, thereby simplifying the interpretations. However, the use of this OR across all  
18 categories is appropriate only when the proportional odds (parallel regression) assumption,  
19 which means the “equality of the log-odds across the different categories of the outcome  
20 variable,” is met. Moreover, this assumption is often violated because it is very common for one  
21 or more of the coefficients or ORs to differ across the categories of outcome variable. In such  
22 cases, it is advisable to use a non-ordinal model, such as multinomial logistic regression.  
23 Unfortunately, such models are not only less parsimonious and difficult to interpret, compared  
24 with ordered logistic regression models, but also they do not consider the ordinal nature of the  
25 variable.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 Generalized ordered logistic regression established by Fu and later by William has been  
49 found to be an appropriate model for such cases, since it relaxes the proportional assumptions by  
50 allowing the effect of each explanatory variable to vary across different categories of outcome  
51 variable.  
52  
53  
54  
55  
56  
57  
58  
59  
60

variable without modifying the data. The generalized ordered logistic regression model can be written using the following formula (1):

$$P(Y_i > j) = \frac{\exp(\alpha_j + X_i \beta_j)}{1 + \exp(\alpha_j + X_i \beta_j)}, j = 1, 2, \dots, M - 1 \quad (1)$$

Where M is the number of categories of the ordinal dependent variable (Y), however, the logit model is a special case of the gologit model when M = 2. When M > 2, is equivalent to the series of binary logistic regressions, such as category 1 versus categories 2, 3, 4 and 5 (Y>1); categories 1 and 2 versus categories 3, 4, and 5 (Y>2); categories 1, 2, and 3 versus categories 4 and 5 (Y>3); and categories 1, 2, 3, and 4 versus category 5 (Y>4). Additionally, the ordered logistic regression model is also a special case of the generalized ordered logistic regression model when the betas are the same for each value of j as shown in formula (2):

$$P(Y_i > j) = \frac{\exp(\alpha_j + X_i \beta)}{1 + \exp(\alpha_j + X_i \beta)}, j = 1, 2, \dots, M - 1 \quad (2)$$

When the betas change for some variables while for the other variables remain the same, the fitted model is described as being partial constrained, since it only allows the betas of the variables that met the proportional assumptions to be constrained while those that have not met the aforementioned assumptions are allowed to vary freely without constraint. The gologit2 command in Stata is responsible for producing this type of model, and as shown in formula (3) below, the betas for X1 and X2 are constrained but the betas for X3 are not.

$$P(Y_i > j) = \frac{\exp(\alpha_j + X1_i \beta1 + X2_i \beta2 + X3_i \beta3_j)}{1 + \exp(\alpha_j + X1_i \beta1 + X2_i \beta2 + X3_i \beta3_j)}, j = 1, 2, \dots, M - 1 \quad (3)$$

**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Page 2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Page 5
Objectives	3	State specific objectives, including any prespecified hypotheses	Page 7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Page 8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 8-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Page 8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page 9-10
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 9-10
Bias	9	Describe any efforts to address potential sources of bias	Page 3, 8
Study size	10	Explain how the study size was arrived at	Page 8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page 10-12
		(b) Describe any methods used to examine subgroups and interactions	Page 10-12
		(c) Explain how missing data were addressed	N/A
		(d) If applicable, describe analytical methods taking account of sampling strategy	Page 10
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			



Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page 13
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page 14 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers of outcome events or summary measures	Page 14 and Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2,3,4
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Table 4
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 18-9, Table 4
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 24
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 24
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 24
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	N/A

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Improving access to health care for women in Tanzania by addressing socioeconomic determinants and health insurance

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023013.R2
Article Type:	Research
Date Submitted by the Author:	20-Jun-2018
Complete List of Authors:	Bintabara, Deogratus; Tokyo Medical and Dental University, Global Health Entrepreneurship; The University of Dodoma, Public Health Nakamura, Keiko; Tokyo Medical and Dental University, Global Health Entrepreneurship Seino, Kaoruko; Tokyo Medical and Dental University, Global Health Entrepreneurship; Michigan University, Department of Environmental Health Sciences, School of Public Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Global health, Health policy, Health services research
Keywords:	HEALTH SERVICES ADMINISTRATION & MANAGEMENT, access to health care, women, Tanzania

SCHOLARONE™  
Manuscripts

Only

1  
2  
3 1 **Improving access to health care for women in Tanzania by addressing socioeconomic**  
4  
5 2 **determinants and health insurance**  
6

7  
8 3 **Deogratus Bintabara<sup>1,2</sup>, Keiko Nakamura<sup>1</sup>, Kaoruko Seino<sup>1,3</sup>**  
9

10  
11 4 **Deogratus Bintabara, M.D.**

12  
13  
14 5 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

15  
16 6 Graduate School of Tokyo Medical and Dental University,

17  
18 7 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.

19  
20 8 <sup>2</sup>Department of Public Health, College of Health Sciences,

21  
22 9 The University of Dodoma, P.O Box 259, Dodoma, Tanzania

23  
24  
25 10 Phone: (255) 767 966 114, Email: bintabaradeo@gmail.com  
26  
27

28  
29 11 **Keiko Nakamura, M.D., Ph.D., (Corresponding author)**

30  
31 12 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

32  
33 13 Graduate School of Tokyo Medical and Dental University, Email: nakamura.ith@tmd.ac.jp  
34

35  
36 14 **Kaoruko Seino, Ph.D.**

37  
38 15 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

39  
40 16 Graduate School of Tokyo Medical and Dental University,

41  
42 17 <sup>3</sup>Department of Environmental Health Sciences, School of Public Health,

43  
44 18 University of Michigan, Ann Arbor, Michigan USA, Email: seino.ith@tmd.ac.jp  
45  
46  
47

48 19 **Keywords:** Health services administration and management, Access to healthcare, Women,  
49  
50 20 Tanzania

51  
52 21 **Word count:** 3649  
53  
54  
55  
56  
57

## 1 ABSTRACT

2 **Objective:** This study was performed to explore the factors associated with accumulation of  
3 multiple problems in accessing health care among women in Tanzania as an example of a low-  
4 income country.

5 **Design:** Population-based cross-sectional survey

6 **Setting:** Nationwide representative data for women of reproductive age obtained from the 2015  
7 – 2016 Tanzania Demographic Health Survey were analyzed.

8 **Primary outcome measures:** A composite variable, “problems in accessing health care,” with  
9 five (0 – 4) categories was created based on the number of problems reported: obtaining  
10 permission to go to the doctor, obtaining money to pay for advice or treatment, distance to a  
11 health facility, and not wanting to go alone. Respondents who reported fewer or more problems  
12 placed in lower and higher categories, respectively.

13 **Results:** A total of 13266 women aged 15 – 49 years, with a median age (IQR) of 27 (20 – 36)  
14 years were interviewed and included in the analysis. About two-third s (65.53%) of the  
15 respondents reported at least one of the four major problems in accessing health care.  
16 Furthermore, after controlling for other variables included in the final model, women without  
17 any type of health insurance, those belonging to the poorest class according to the wealth index,  
18 those who had not attended any type of formal education, those who were not employed for cash,  
19 each year of increasing age, and those who were divorced, separated, or widowed were  
20 associated with greater problems in accessing health care.

21

1           **Conclusion:** This study indicated the additive effects of barriers to health care in low-  
2 income countries such as Tanzania. Based on these results, improving uptake of health insurance  
3 and addressing social determinants of health are the first steps toward reducing women's  
4 problems associated with accessing health care.

### 5 **Strengths and limitations of this study**

- 6       ➤ This is the first study to show additive effects of problems in accessing health care and  
7       associated factors among women in Tanzania.
- 8       ➤ The study used a nationally representative sample with high response rate and robust  
9       sampling procedure.
- 10      ➤ As a cross-sectional study design was used, causality assumptions could not be made.  
11      Therefore, the results should be interpreted with caution.

## 1 INTRODUCTION

2 Despite the substantial decline in global maternal mortality ratio (MMR), low-income  
3 countries have not seen the same decline in MMR as those with higher incomes<sup>1,2</sup>; this situation  
4 is described as an “area of shameful failures of development.”<sup>3,4</sup> Current statistics show that the  
5 MMR has increased significantly by more than 20% over the past 5 years in Tanzania, despite  
6 the governmental efforts to strengthen the health system by increasing the coverage of health  
7 care facilities.<sup>5-7</sup> Each ward now has at least one dispensary and/or health center, each district  
8 has at least one hospital, while each region has at least one referral hospital. Regardless of  
9 differences in the levels of these facilities in terms of function, expertise, availability of services,  
10 and population coverage, all are expected to provide basic maternal health services together with  
11 basic emergency obstetric care.<sup>7</sup> This resulted in an increase in coverage of maternal health  
12 services, such as antenatal care by skilled providers (96% – 98%), delivery at a health care  
13 facility (50% – 63%), and births assisted by skilled providers (51% – 64%) between 2010 and  
14 2016.<sup>5,6</sup> Persistent high MMR (556 maternal deaths per 100000 live births) in Tanzania raises  
15 concerns reading whether the country can achieve the Sustainable Development Goals (SDGs)  
16 target of less than 140 maternal deaths per 100000 live births by 2030.<sup>8,9</sup> The majority of  
17 avoidable and unnecessary maternal deaths experienced in this region likely result from poor  
18 utilization of skilled maternal health services.<sup>10</sup>

19 Many social, cultural, and geographical factors as well as education level and poverty  
20 have been reported to play roles in the poor utilization of health services.<sup>11-13</sup> Access to health  
21 care has been highlighted as the major barrier toward the utilization of maternal health services  
22 in low-income countries, especially in sub-Saharan Africa (SSA).<sup>10,14,15</sup> Access to health care can  
23 be broadly defined based on availability, affordability, accessibility and acceptability,<sup>16</sup> but is

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2 Despite agreement that access to health care must be universal and guaranteed for all on an  
3 equitable basis,<sup>17</sup> women continue to face significant inequities in accessing and utilizing health  
4 care particularly in low-income countries.<sup>18</sup>

5 In relation to the problems experienced by women in accessing health care, the following  
6 four major problems have been addressed in previous studies: obtaining permission,<sup>19,20</sup>  
7 obtaining money,<sup>21</sup> distance to the health facility,<sup>22</sup> and not wanting to go alone (lack of spouse  
8 or family member escort).<sup>23,24</sup> Although it is unclear whether women with multiple problems  
9 encounter greater difficulties in accessing health care, most previous studies assessed and  
10 discussed each of these four problems independently. The limited evidence regarding whether  
11 women facing multiple problems have less access to health care suggested the need to create a  
12 composite variable that includes all four problems to identify groups of women at a greater  
13 disadvantage. A similar approach has been used to assess the severity of problems in accessing  
14 health care among individuals with disability in four African countries.<sup>25</sup> Moreover, recent  
15 studies indicated that age, education, residence, possession of health insurance, socioeconomic  
16 status, and occupation are strongly linked to access to health care.<sup>26–30</sup> The present study was  
17 performed to explore the factors associated with accumulation of multiple problems in accessing  
18 health care among women in Tanzania.

## 19 **METHODS**

### 20 **Data sources**

21 The present study used data from the 2015 – 2016 Tanzania Demographic Health Survey  
22 and Malaria Indicator Survey (TDHS-MIS) conducted by the National Bureau of Statistics

1 (NBS) and the Office of Chief Government Statistician (OCGS), Zanzibar, in collaboration with  
2 the Ministry of Health, Community Development, Gender, Elderly, and Children (MoHCDGEC)  
3 of the Tanzania Mainland and the Ministry of Health (MoH), Zanzibar. The technical support for  
4 the surveys was provided by ICF International under DHS program.

## 5 **Study design**

6 This study analyzed a nationwide population-based cross-sectional survey using  
7 information obtained by interviewing women (15 – 49 years old) who were either residents or  
8 visitors in the household on the night before the survey.

## 9 **Sample size and sampling technique**

10 The 2015 – 2016 TDHS-MIS used a two-stage cluster sampling technique to obtain a  
11 sample designed to provide nationally representative results according to all 30 regions of  
12 Tanzania. In the first stage, sample points (a total of 608 clusters) consisting of enumeration  
13 areas delineated for the 2012 Tanzania Population and Housing Census were selected. In the  
14 second stage, households were selected systematically. A complete listing of households was  
15 established for all 608 selected clusters prior to the fieldwork. From this list, 22 households were  
16 then systematically selected from each cluster, yielding a representative probability sample of  
17 13376 households. All eligible women in the selected households and men in subsample of one –  
18 third of selected households between the ages of 15 and 49 years who were either residents or  
19 visitors in the household on the night before the survey were then interviewed. Finally, a total of  
20 13266 women and 3514 men were interviewed.<sup>6</sup>

## 21 **Data collection and processing**

22 The 2015 – 2016 TDHS-MIS used four main types of questionnaires during data  
23 collection. However, only data collected with the Women's Questionnaire were used in the



1 present study. After pre-testing of the questionnaires, the finalized and corrected version was  
2 used in the main survey from August 22, 2015, through February 14, 2016. Data collection was  
3 performed by 64 female nurses who were trained and qualified to be interviewers through a  
4 series of practical tests and examinations. Following the training, 16 teams were formed (three  
5 for Zanzibar and 13 for Tanzania Mainland). Data entry was performed concurrently with data  
6 collection in the field. After the paper questionnaires were completed, edited, and checked by  
7 both the field editor and the supervisor, the data were entered into a tablet equipped with a data  
8 entry program. A 100% double entry data entry process was used to minimize keying errors, and  
9 editing was completed on March 21, 2016, while data cleaning and finalization were completed  
10 on April 22, 2016.

### 11 **Measurement of variables**

12 **Outcome variable:** In this survey women were asked whether each of the following four  
13 factors was a problem in seeking medical advice or treatment when they were ill: obtaining  
14 permission to go to the doctor/health facility; obtaining money to pay for advice, consultation or  
15 treatment; distance to the health facility; and not wanting to go alone. A new composite variable  
16 called “problems in accessing health care” was then created based on the number of problems  
17 reported, with respondents reporting fewer or more problems placed in lower and higher  
18 categories, respectively. The categories were assigned as 1, 2, 3, 4, and 5 for women who  
19 reported “no,” “one,” “two,” “three,” and “four” problems in accessing health care, respectively.  
20 These categories of a composite (outcome) variable were treated as ordinal numbers, with the  
21 assumption that conceptual differences between categories were identical.

22 **Independent variables:** Several independent variables that have been linked both  
23 empirically and theoretically with the accessibility of health care among women were included in

1 the present study. The respondents were divided into groups according to age as follows: 15 – 19,  
2 20 – 34, and 35 – 49 years. Women were grouped according to marital status as “never married,”  
3 “married/living together,” and “divorced, separated, or widowed.” Education level was classified  
4 as “none,” “primary,” “secondary,” and “higher” (including college and all university levels).  
5 Employment in the last 12 months was grouped into “not employed,” “employed for cash,” and  
6 “employed but paid-in-kind.” The area of residence was grouped into “urban” and “rural.”  
7 Possession of health insurance was grouped as “no” for women who did not have any type of  
8 health insurance and “yes” to those who had any type of health insurance. The wealth index was  
9 computed based on household assets and housing characteristics. During computation, the  
10 households were given scores based on the number and kinds of consumer goods they owned,  
11 ranging from a television to a bicycle or car, plus housing characteristics, such as the source of  
12 drinking water, toilet facilities, and flooring materials. These scores were derived using principal  
13 component analysis. National wealth quintiles were compiled by assigning the household score  
14 to each usual (de jure) household member, ranking each person in the household by their score,  
15 and then dividing the distribution into five equal categories, each with 20% of the population, as  
16 “poorest,” “poorer,” “middle,” “richer,” and “richest.”<sup>6</sup> The selection of these variables was  
17 based on previous studies.<sup>26–30</sup>

## 18 **Statistical approaches**

19 In descriptive analyses, categorical variables were summarized using proportions and  
20 then presented in tables, while quantitative variables were summarized using the median and  
21 interquartile range (IQR).

22 As the outcome variable “problems in accessing health care” was ordinal in nature (a  
23 score based on the number of different reported problems), in which the order of values

1 corresponded to a hierarchy in meaning as in this study, the application of ordered logistic  
2 regression was recommended.<sup>31</sup> Stata 14 (StataCorp, College Station, TX) was used for analysis  
3 in the present study. For all analyses, the Stata survey set commands were used to adjust for the  
4 variability of clustering and all the estimates were weighted to correct for non-responses and  
5 disproportionate sampling.

6 When assessing the associations between selected independent variables and the outcome  
7 variable, four models were tested for fit: a proportional ordered logistic model, a generalized  
8 ordered logistic regression model, and a generalized ordered logistic regression model with and  
9 without alternative parameterization. The generalized ordered logistic regression model with  
10 alternative parameterization was chosen as the best fit. This model allows some variables to be  
11 modeled with the proportional odds assumption while the parallel line constraint is relaxed for  
12 variables in which the assumption is not met. The model is less restrictive as it allows the  
13 coefficient of the variables to vary for the different categories that are compared.<sup>32</sup> The model  
14 provides gamma coefficients that show the extent to which the parallel regression assumption is  
15 violated by the variable.

## 16 **Ethics statement**

17 This study was based on an analysis of existing public domain survey data sets that are  
18 freely available online with all identifier information detached. The original TDHS-MIS  
19 protocols were reviewed by the Institution Review Board (IRB) of ICF Macro at Calverton in the  
20 USA and by the National Institute of Medical Research (NIMR) IRB in Tanzania. The ICF IRB  
21 ensured that the survey complied with the US Department of Health and Human Services  
22 regulations for the protection of human subjects (45 CFR 46), while NIMR-IRB ensured that the  
23 surveys complied with the laws and norms of Tanzania. The participants were adequately

1 informed about all relevant aspects of the survey, including its objective and interview  
 2 procedures. All participants accepted participating in the study signed informed consent prior to  
 3 the interviews.

#### 4 **Patient and Public Involvement Statement**

5 Patients and the public were not involved in the analysis of this study.

## 6 **RESULTS**

### 7 **Respondents' characteristics**

8 As shown in Table 1, a total of 13266 women between 15 and 49 years old were  
 9 interviewed and included in the analysis. The median age (IQR) of the respondents was 27 (20 –  
 10 36) years. About 62% of the respondents were living with their spouse at the time of the  
 11 interview. Less than 2% had attained the highest level of education (college or university).  
 12 Nearly two-thirds were employed but paid-in-kind and one-tenth reported having any type of  
 13 health insurance. Almost half of the respondents reported that obtaining money for health care  
 14 was the major problem in accessing health care. Furthermore, about two-thirds of the  
 15 respondents reported at least one of the four problems in accessing health care.

16 **Table 1 Percent distribution of women between the ages 15 – 49 years by selected**  
 17 **background characteristics, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

Variable	<i>n (%)</i> (Weighted)
<b>Age (Median (IQR)=27, (20-36)</b>	
15-19	2904 (21.89)
20-34	6360 (47.94)
35-49	4002 (30.17)
<b>Marital status</b>	
Never married	3353 (25.27)
Married/living together	8210 (61.89)
Divorced/separated/widowed	1703 (12.84)

**Education**

None	1947 (14.67)
Primary	8211 (61.90)
Secondary	2925 (22.05)
Highest	183 (1.38)

**Employed last 12 months**

Not employed	3033 (22.86)
Employed for cash	6197 (46.71)
Employed but paid-in-kind	4036 (30.43)

**Residence**

Urban	4811 (36.27)
Rural	8455 (63.73)

**Health insurance ownership**

Yes	12066 (90.95)
No	1200 (9.05)

**Health quintile**

Lowest	2246 (16.93)
Second	2274 (17.14)
Middle	2328 (17.55)
Fourth	2822 (21.27)
Highest	3596 (27.11)

**Types of problems\***

Obtaining money	6565 (49.49)
Distance to facility	5615 (42.33)
Not want to go alone	3962 (29.87)
Obtaining permission	1900 (14.32)

**Number of problems in accessing health care**

None	4574 (34.48)
One problem	3291 (24.81)
Two problems	2547 (19.20)
Three problems	1759 (13.26)
Four problems	1095 (8.25)

1 Note: \**n* and % do not add up to 13266 and 100%, respectively, because multiple responses were possible.

## 2 Partial generalized ordered logistic regression model with alternative gamma 3 parameterization

4 Table 2 shows the results of the partially constrained generalized ordered logistic  
5 regression model with alternative (gamma) parameterization for the outcome variable of  
6 problems in accessing health care. The results showed non-significant Wald test statistics,  
7 indicating that the model did not violate the proportional odds/parallel regression assumptions.

1  
2  
3 1 However, constraints for parallel lines were not imposed for age, wealth status (richest), and  
4  
5 2 marital status (divorced, separated, or widowed). The remaining variables that met the parallel  
6  
7 3 assumption can be interpolated in the same manner as for the ordered logistic regression model  
8  
9 4 as follows. The odds of being in higher categories of problems in accessing health care versus  
10  
11 5 lower or equal to a reference category were 0.622 times lower for women who had health  
12  
13 6 insurance than those who did not, given constant values for the other variables in the model.  
14  
15 7 Furthermore, the odds of being in higher categories of problems in accessing health care versus  
16  
17 8 less than or equal to a reference category were 1.22 times higher for women who were employed  
18  
19 9 but paid-in-kind than those who were unemployed for the last 12 months before the survey  
20  
21 10 keeping the other variables constant in the model.  
22  
23  
24  
25

26 11 The variables for which constraints for parallel lines were not imposed were interpreted  
27  
28 12 as follows: the coefficients for age and marital status (divorced, separated, or widowed) were  
29  
30 13 consistently positive, while those for wealth status (richest) were negative but decreased across  
31  
32 14 the cut-points. Therefore, for each year of increase in age and being divorced, separated, or  
33  
34 15 widowed, women were more likely to report having a larger number of problems in accessing  
35  
36 16 health care. The greatest differences were seen with increasing age and for those who were  
37  
38 17 divorced, separated, or widowed, women were less likely to report having few problems in  
39  
40 18 accessing health care. In addition, the women in the richest economic group tended to be less  
41  
42 19 likely to report having many problems in accessing health care than those in the poorest group,  
43  
44 20 with the greatest differences because the richest women were less likely to report having many  
45  
46 21 problems in accessing health care.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2 Generalized ordered logistic regression model with alternative gamma parameterization, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

	Variable	POR [95% CI]	P-value
<b>Beta</b>	<b>Health insurance</b> (ref: No)		
	Yes	0.622 [0.531 - 0.731]	0.000
	<b>Residence</b> (ref: Urban)		
	Rural	0.858 [0.728 - 1.012]	0.069
	<b>Age</b> (continuous)	1.010 [1.001 - 1.017]	0.000
	<b>Marital status</b> ref: Never married)		
	Married/living together	0.901 [0.801 - 1.014]	0.085
	Divorced/separated/widowed	1.418 [1.175 - 1.712]	0.000
	<b>Education</b> (ref: None)		
	Primary	0.883 [0.788 - 0.990]	0.033
	Secondary	0.683 [0.582 - 0.800]	0.000
	Highest	0.516 [0.360 - 0.741]	0.000
	<b>Wealth status</b> (ref: Poorest)		
	Poorer	0.854 [0.726 - 1.006]	0.059
	Middle	0.725 [0.626 - 0.840]	0.000
	Richer	0.496 [0.417 - 0.590]	0.000
	Richest	0.291 [0.233 - 0.364]	0.000
<b>Employed last 12 months</b> (ref: Not employed)			
Employed for cash	0.975 [0.869 - 1.095]	0.668	
Employed but paid in kind	1.220 [1.067 - 1.395]	0.004	
<b>Gamma_2</b>	Age	0.993 [0.989 - 0.998]	0.000
	Wealth status (Richest)	1.279 [1.140 - 1.435]	0.000
	Marital status (Divorced/separated/widowed)	0.814 [0.701 - 0.945]	0.007
<b>Gamma_3</b>	Age	0.993 [0.986 - 0.999]	0.018
	Wealth status (Richest)	1.515 [1.265 - 1.814]	0.000
	Marital status (Divorced/separated/widowed)	0.749 [0.625 - 0.899]	0.002
<b>Gamma_4</b>	Age	0.987 [0.978 - 0.996]	0.005
	Wealth status (Richest)	1.957 [1.508 - 2.540]	0.000
	Marital status (Divorced/separated/widowed)	0.566 [0.419 - 0.764]	0.000

Note: Wald test of parallel lines assumption for the final model:  $F(33, 517) = 1.110, P = 0.310$ .

A non-significant test statistic indicates that the final model does not violate the proportional odds/parallel lines assumption.

\*POR = Proportional odds ratio.



## 1 DISCUSSION

2 This study was performed to explore the factors associated with accumulation of multiple  
3 problems in accessing health care among women in Tanzania. To our knowledge, this is the first  
4 study to explore the factors associated with multiple problems in accessing health care.  
5 Furthermore, the study used a nationally representative sample from Tanzania with the  
6 application of generalized ordered logistic regression models, which provided the best models  
7 for ordinal data to validate the factors associated with problems in accessing health care. In the  
8 present study, about 65%, 40%, and 20% of women reported “one or more,” “two or more,” and  
9 “three or more” major problems in accessing health care, respectively. In addition, not having  
10 health insurance and low socioeconomic status as measured by wealth, education, and  
11 employment status were associated with accumulation of multiple problems in accessing health  
12 care.

13 The high proportion of women reporting problems in accessing health care in this study  
14 was consistent with the findings of a previous study performed in Egypt.<sup>33</sup> These findings  
15 provide evidence that demand-side barriers, such as cost of care, permission from their spouse,  
16 lack of someone to escort to a health care facility, and distance to the facility,<sup>34,35</sup> still prevent  
17 many African women from accessing health care. Due to the cultural, social, and traditional  
18 perceptions in Africa that maternal health is only the responsibility of women, existing and new  
19 interventions should influence health service utilization to begin at the individual, household, and  
20 community levels to eliminate such demand-side barriers.<sup>21,36</sup>

21 Having health insurance is an essential element for timely access to health care and better  
22 health-related outcomes.<sup>37,38</sup> Despite the availability of the National Health Insurance Fund



1 (NHIF) in Tanzania since 2001 through Act no. 8 of 1999, more than 90% of women are still  
2 uninsured.<sup>27,39</sup> Less than one-tenth of the women in the present study reported having any type of  
3 health insurance. Moreover, the women who had health insurance were less likely to report  
4 having multiple problems associated with access to health care. This may have been because  
5 having health insurance makes someone not only more comfortable with receiving a wide range  
6 of services but also ensures a wider choice regarding where and when to obtain health care  
7 without being afraid of the costs as they are covered by insurance. Similar findings were reported  
8 in a study conducted Ghana.<sup>26</sup> The similarity in the findings between these studies may have  
9 been because both used secondary data collected by a DHS program that applied a similar  
10 methodology. Furthermore, the participants in these two studies were from SSA, and therefore  
11 may have similar socioeconomic determinants.

12 The present study indicated a strong association between being in the poorest class of the  
13 wealth index and accumulation of multiple problems in accessing health care among women in  
14 Tanzania. This finding may be explained by the fact that being in the poorest class requires  
15 individuals to spend their income on basic needs, such as food, and health care costs may  
16 therefore be less likely to be affordable.<sup>40</sup> Such women are therefore more likely to report having  
17 many problems in accessing health care, as reported in other studies conducted in SSA.<sup>41-43</sup>

18 In contrast to our expectations, we found no significant association between  
19 unemployment and problems in accessing health care. However, this may have been because  
20 being employed is not enough to have full access to health care as there are other barriers  
21 preventing women from accessing health care, such as gender inequality, poor infrastructure, and  
22 lack of knowledge regarding maternal health services.<sup>28,33,44</sup> On the other hand, the results  
23 presented here indicated that women who are employed and receive wages in the form of

1 payment-in-kind, such as food, clothes, and other goods instead of cash, were likely to  
2 experience multiple problems in accessing health care.

3 The variables age, wealth status (richest), and marital status (divorced, separated, or  
4 widowed) were found to vary for each category of the outcome variable. The findings indicated  
5 that the accumulation of multiple problems in accessing health care was associated with older  
6 age and being divorced, separated, or widowed, consistent with the results of previous studies in  
7 other low-income countries.<sup>45,46</sup> Older age is more likely to be accompanied by decreased  
8 working capability, and hence low income, being retired, and uninsured.<sup>47</sup> As the risk of  
9 maternal complications increased with older age<sup>48,49</sup> and living without a spouse reduced the  
10 chance of having an escort to the health facility, efforts and support should be made to provide  
11 such women with access to health care.

12 This study had some limitations, including the risk of misclassification bias, which may  
13 have been introduced due to the lack of external validation of self-reported information that  
14 could have affected categorization of the outcome variable. However, we reduced this effect by  
15 categorizing the outcome variable into five groups and the use of a generalized ordered logistics  
16 regression model that clearly validated the factors associated with problems in accessing health  
17 care among women in low-income countries. In addition, causality assumptions could not be  
18 made due to the cross-sectional nature of the study, and therefore the results should be  
19 interpreted with caution.

20

## 1 **CONCLUSION**

2 The results of the present study provided evidence for additive effects of barriers to  
3 health care in low-income countries, such as Tanzania. Based on these results, improving access  
4 to health insurance and addressing social determinants of health represent the first steps toward  
5 reducing problems associated with accessing health care for women in low-income countries.

## 6 **Contributors**

7 DB originated the design of the study, performed statistical analysis, interpretation and  
8 drafted the manuscript. KN contributed to the design of the study and the interpretation advice of  
9 data. KS contributed to interpretation advice of data and drafted the manuscript. Both DB, KN &  
10 KS critically revised the draft manuscript. All authors read and approved the final manuscript.

## 11 **Acknowledgment**

12 We would like to acknowledge ICF International, Rockville, Maryland, USA, through  
13 DHS program for giving us permission to access the TDHS-MIS 2015-2016 dataset.

## 14 **Funding**

15 This research received no specific grant from any funding agency in the public,  
16 commercial or not-for-profit sectors.

## 17 **Competing interest**

18 None to declare.

## 19 **Provenance and peer review**

20 Not commissioned; externally peer reviewed.

## 21 **A data sharing statement**

1           The datasets used for the current analysis was generated from the original survey of  
2  
3  
4           1           Tanzania DHS-MIS datasets available from within the DHS program repository:  
5  
6           2  
7  
8           3           <http://dhsprogram.com/data/available-datasets.cfm>  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## 1 REFERENCES

- 2 1 Alkema L, Chou D, Hogan D, *et al.* Global, regional, and national levels and trends in  
3 maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a  
4 systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*  
5 2016;**387**:462–74. doi:10.1016/S0140-6736(15)00838-7
- 6 2 World Health Organization (WHO). Trends in Maternal Mortality: 1990 to 2015:  
7 estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations  
8 Population Division. 2015. doi:ISBN 978 92 4 150363 1
- 9 3 Godefay H, Byass P, Graham WJ, *et al.* Risk Factors for Maternal Mortality in Rural  
10 Tigray, Northern Ethiopia: A Case-Control Study. *PLoS One* 2015;**10**:e0144975.  
11 doi:10.1371/journal.pone.0144975
- 12 4 Lewis G. Maternal mortality in the developing world: why do mothers really die? *Obstet*  
13 *Med Med Pregnancy* 2008;**1**:2–6. doi:10.1258/om.2008.080019
- 14 5 National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2011. Tanzania  
15 Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro.  
16 [http://www.measuredhs.com/pubs/pdf/FR243/FR243\[24June2011\].pdf](http://www.measuredhs.com/pubs/pdf/FR243/FR243[24June2011].pdf) Accessed 6 May  
17 2018.
- 18 6 Ministry of Health, Community Development, Gender, Elderly and Children  
19 (MoHCDGEC) [Tanzania Mainland], Ministry of Health (MoH) [Zanzibar], National  
20 Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF.  
21 2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-*  
22 *MIS) 2015-16*. Dar es Salaam, Tanzania, and Rockville, Maryland, USA: MoHCDGEC,  
23 MoH, NBS, OCGS, and ICF; 2016. <https://dhsprogram.com/pubs/pdf/FR321/FR321.pdf>.

- 1  
2  
3 1 Accessed 6 May 2018.  
4  
5 2 7 United Republic of Tanzania (URT) and World Health Organization (WHO). Health  
6  
7 Systems Profile in Tanzania. URT; 2004.  
8 3  
9  
10 4 <http://mbaralidc.go.tz/storage/app/media/uploaded-files/Health-tanzania.pdf> Accessed 6  
11  
12 5 May 2018.  
13  
14 6 8 World Health Organization (WHO). Strategies toward ending preventable maternal  
15  
16 mortality ( EPMM ). Published Online First: 2015. doi:ISBN 978 92 4 150848 3  
17 7  
18 8 9 Moran AC, Jolivet RR, Chou D, *et al.* A common monitoring framework for ending  
19  
20 preventable maternal mortality, 2015–2030: phase I of a multi-step process. *BMC*  
21  
22 *Pregnancy Childbirth* 2016;**16**:250. doi:10.1186/s12884-016-1035-4  
23  
24 10  
25 11 10 Odetola TD. Health care utilization among rural women of child-bearing age: a nigerian  
26  
27 experience. *Pan Afr Med J* 2015;**20**:1–7. doi:10.11604/pamj.2015.20.151.5845  
28  
29 12  
30 13 11 Babalola SO. Factors associated with use of maternal health services in Haiti: a multilevel  
31  
32 analysis. *Rev Panam Salud Publica* 2014;**36**:1–9.  
33  
34 14  
35 15 12 Ngomane S, Mulaudzi FM. Indigenous beliefs and practices that influence the delayed  
36  
37 attendance of antenatal clinics by women in the Bohlabele district in Limpopo, South  
38  
39 Africa. *Midwifery* 2012;**28**:30–8. doi:10.1016/j.midw.2010.11.002  
40  
41 17  
42 18 13 Ali HS, Abdalla A, Abdalla A. Understand Factors Influencing Accessibility of Pregnant  
43  
44 Women to Antenatal Care Services Accessibility factors : Demographic characteristics of  
45  
46 the study. *Heal Sci J* 2016;**10**:1–5. doi:10.4172/1791-809X.1000100507  
47  
48 20  
49 21 14 Victora CG, Barros AJ, Axelson H, *et al.* How changes in coverage affect equity in  
50  
51 maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of  
52  
53 national surveys. *Lancet* 2012;**380**:1149–56. doi:10.1016/S0140-6736(12)61427-5  
54  
55 23  
56  
57  
58  
59  
60

- 1  
2  
3 1 15 Singh PK, Kumar C, Rai RK, *et al.* Factors associated with maternal healthcare services  
4 utilization in nine high focus states in India: a multilevel analysis based on 14 385  
5  
6 2  
7 communities in 292 districts. *Health Policy Plan* 2014;**29**:542–59.  
8  
9  
10 4  
11 doi:10.1093/heapol/czt039  
12  
13 5 16 Donnell OO. Access to health care in developing countries : breaking down demand side  
14  
15 6  
16 barriers. *Cad Saude Publica* 2007;**23**:2820–34. doi:S0102-311X2007001200003 [pii]  
17  
18 7 17 Kirby N. Access to healthcare services as a human right. *Med Law* 2010;**29**:487–96.  
19  
20 8 18 Ganle JK, Parker M, Fitzpatrick R, *et al.* Inequities in accessibility to and utilisation of  
21  
22 9  
23 maternal health services in Ghana after user-fee exemption: a descriptive study. *Int J*  
24  
25 10  
26 *Equity Health* 2014;**13**:89. doi:10.1186/s12939-014-0089-z  
27  
28 11 19 Danforth E, Kruk M, Rockers P, *et al.* Household Decision-making about Delivery in  
29  
30 12  
31 Health Facilities: Evidence from Tanzania. *J Heal Popul Nutr* 2009;**27**:696–703.  
32  
33 13  
34 doi:10.3329/jhpn.v27i5.3781  
35  
36 14 20 Wilunda C, Scanagatta C, Putoto G, *et al.* Barriers to utilisation of antenatal care services  
37  
38 15  
39 in South Sudan: a qualitative study in Rumbek North County. *Reprod Health* 2017;**14**:65.  
40  
41 16  
42 doi:10.1186/s12978-017-0327-0  
43  
44 17 21 Lowe M, Chen D-R, Huang S-L. Social and Cultural Factors Affecting Maternal Health in  
45  
46 18  
47 Rural Gambia: An Exploratory Qualitative Study. *PLoS One* 2016;**11**:e0163653.  
48  
49 19  
50 doi:10.1371/journal.pone.0163653  
51  
52 20 22 Mbiza CR, Kazembe A, Simwaka A. Barriers to health-seeking practices during  
53  
54 21  
55 pregnancy among adolescents in rural Blantyre, Malawi. *Afr J Midwifery Womens Health*  
56  
57 22  
58 2014;**8**:59–65. doi:10.12968/ajmw.2014.8.2.59  
59  
60 23 23 Kumbani L, Bjune G, Chirwa E, *et al.* Why some women fail to give birth at health

- 1 facilities: a qualitative study of women's perceptions of perinatal care from rural Southern  
2 Malawi. *Reprod Health* 2013;**10**:9. doi:10.1186/1742-4755-10-9
- 3 24 Vidler M, Ramadurg U, Charantimath U, *et al*. Utilization of maternal health care services  
4 and their determinants in Karnataka State, India. *Reprod Health* 2016;**13**:37.  
5 doi:10.1186/s12978-016-0138-8
- 6 25 Eide AH, Mannan H, Khogali M, *et al*. Perceived Barriers for Accessing Health Services  
7 among Individuals with Disability in Four African Countries. *PLoS One*  
8 2015;**10**:e0125915. doi:10.1371/journal.pone.0125915
- 9 26 Browne JL, Kayode GA, Arhinful D, *et al*. Health insurance determines antenatal,  
10 delivery and postnatal care utilisation: evidence from the Ghana Demographic and Health  
11 Surveillance data. *BMJ Open* 2016;**6**:e008175. doi:10.1136/bmjopen-2015-008175
- 12 27 Kibusi SM, Sunguya BF, Kimunai E, *et al*. Health insurance is important in improving  
13 maternal health service utilization in Tanzania—analysis of the 2011/2012 Tanzania  
14 HIV/AIDS and malaria indicator survey. *BMC Health Serv Res* 2018;**18**:112.  
15 doi:10.1186/s12913-018-2924-1
- 16 28 Tsawe M, Susuman A. Determinants of access to and use of maternal health care services  
17 in the Eastern Cape, South Africa: a quantitative and qualitative investigation. *BMC Res*  
18 *Notes* 2014;**7**:723. doi:10.1186/1756-0500-7-723
- 19 29 Mrisho M, Schellenberg JA, Mushi AK, *et al*. Factors affecting home delivery in rural  
20 Tanzania. *Trop Med Int Heal* 2007;**12**:862–72. doi:10.1111/j.1365-3156.2007.01855.x
- 21 30 Kyei-Nimakoh M, Carolan-Olah M, McCann T V. Access barriers to obstetric care at  
22 health facilities in sub-Saharan Africa—a systematic review. *Syst Rev* 2017;**6**:110.  
23 doi:10.1186/s13643-017-0503-x



- 1  
2  
3 1 31 Quddus MA, Wang C, Ison SG. Road Traffic Congestion and Crash Severity:  
4  
5 2 Econometric Analysis Using Ordered Response Models. *J Transp Eng* 2010;**136**:424–35.  
6  
7 doi:10.1061/(ASCE)TE.1943-5436.0000044  
8  
9  
10 4 32 Williams R. Understanding and interpreting generalized ordered logit models. *J Math*  
11  
12 5 *Sociol* 2016;**40**:7–20. doi:10.1080/0022250X.2015.1112384  
13  
14  
15 6 33 Chiang C, Labeeb SA, Higuchi M, *et al.* Barriers to the use of basic health services among  
16  
17 7 women in rural southern Egypt (Upper Egypt). *Nagoya J Med Sci* 2013;**75**:225–31.  
18  
19 8 34 Mselle LT, Kohi TW. Healthcare access and quality of birth care: narratives of women  
20  
21 9 living with obstetric fistula in rural Tanzania. *Reprod Health* 2016;**13**:87.  
22  
23 doi:10.1186/s12978-016-0189-x  
24  
25  
26 11 35 Shartz A, Long SK, Benatar S. Health Care Costs Are a Barrier to Care for Many  
27  
28 12 Women. Urban Inst. 2015.[http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf)  
29  
30 13 [to-Care-for-Many-Women.pdf](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf) (accessed 5 May 2018).  
31  
32  
33 14 36 Azuh D, Fayomi O, Ajayi, Lady. Socio-Cultural Factors of Gender Roles in Women's  
34  
35 15 Healthcare Utilization in Southwest Nigeria. *Open J Soc Sci* 2015;**3**:105–17.  
36  
37 doi:10.4236/jss.2015.34013  
38  
39  
40 17 37 Hsia J, Kemper E, Sofaer S, *et al.* Is Insurance a More Important Determinant of  
41  
42 18 Healthcare Access Than Perceived Health? Evidence from the Women's Health Initiative.  
43  
44 19 *J Womens Health Gend Based Med* 2000;**9**:881–9. doi:10.1089/152460900750020919  
45  
46  
47 20 38 Sommers BD, Gawande AA, Baicker K. Health Insurance Coverage and Health — What  
48  
49 21 the Recent Evidence Tells Us. *N Engl J Med* 2017;**377**:586–93.  
50  
51 22 doi:10.1056/NEJMsbl706645  
52  
53  
54 23 39 Marwa CW. Provision of National Health Insurance Fund Services to Its Members ; Pain  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 or Gain ? *Unified J Sport Heal Sci* 2016;**2**:1–6.  
4  
5 2 40 Yaya S, Bishwajit G, Shah V. Wealth, education and urban–rural inequality and maternal  
6  
7 healthcare service usage in Malawi. *BMJ Glob Heal* 2016;**1**:e000085. doi:10.1136/bmjgh-  
8  
9 2016-000085  
10  
11 4  
12 5 41 Shiferaw S, Spigt M, Godefrooij M, *et al.* Why do women prefer home births in Ethiopia?  
13  
14 *BMC Pregnancy Childbirth* 2013;**13**:5. doi:10.1186/1471-2393-13-5  
15  
16  
17 7 42 Lakew S, Tachbele E, Gelibo T. Predictors of skilled assistance seeking behavior to  
18  
19 pregnancy complications among women at southwest Ethiopia: a cross-sectional  
20  
21 community based study. *Reprod Health* 2015;**12**:109. doi:10.1186/s12978-015-0102-z  
22  
23  
24 10 43 Mwangome FK, Holding PA, Songola KM, *et al.* Barriers to hospital delivery in a rural  
25  
26 setting in coast province, Kenya: Community attitude and behaviours. *Rural Remote*  
27  
28 *Health* 2012;**12**. doi:22471588  
29  
30  
31 13 44 Bintabara D, Mpembeni RNM, Mohamed AA. Knowledge of obstetric danger signs  
32  
33 among recently-delivered women in Chamwino district, Tanzania: a cross-sectional study.  
34  
35 *BMC Pregnancy Childbirth* 2017;**17**:276. doi:10.1186/s12884-017-1469-3  
36  
37  
38 16 45 Tey N-P, Lai S. Correlates of and Barriers to the Utilization of Health Services for  
39  
40 Delivery in South Asia and Sub-Saharan Africa. *Sci World J* 2013;**2013**:1–11.  
41  
42 18 doi:10.1155/2013/423403  
43  
44  
45 19 46 Cheptum J, Gitonga M, Mutua E, *et al.* Barriers to Access and Utilization of Maternal and  
46  
47 Infant Health Services in Migori , Kenya. *Iiste* 2014;**4**:48–53.  
48  
49 21 47 Hadley J, Waidmann T. Health Insurance and Health at Age 65: Implications for Medical  
50  
51 Care Spending on New Medicare Beneficiaries. *Health Serv Res* 2006;**41**:429–51.  
52  
53 23 doi:10.1111/j.1475-6773.2005.00491.x  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 48 Ziadeh S, Yahaya A. Pregnancy outcome at age 40 and older. *Arch Gynecol Obstet*  
4  
5 2001;**265**:30–3. doi:10.1007/s004040000122  
6  
7  
8 3 49 Illah E, Mbaruku G, Masanja H, *et al.* Causes and risk factors for maternal mortality in  
9  
10 4 rural Tanzania - case of Rufiji Health and Demographic Surveillance Site (HDSS). *Afr J*  
11  
12 5 *Reprod Health* 2013;**17**:119–30.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Page 2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Page 5
Objectives	3	State specific objectives, including any prespecified hypotheses	Page 7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Page 8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 8-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Page 8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page 9-10
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 9-10
Bias	9	Describe any efforts to address potential sources of bias	Page 3, 8
Study size	10	Explain how the study size was arrived at	Page 8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page 10-12
		(b) Describe any methods used to examine subgroups and interactions	Page 10-12
		(c) Explain how missing data were addressed	N/A
		(d) If applicable, describe analytical methods taking account of sampling strategy	Page 10
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page 13
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page 14 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers of outcome events or summary measures	Page 14 and Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2,3,4
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Table 4
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 18-9, Table 4
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 24
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 24
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 24
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	N/A

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Improving access to health care for women in Tanzania by addressing socioeconomic determinants and health insurance

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023013.R3
Article Type:	Research
Date Submitted by the Author:	02-Aug-2018
Complete List of Authors:	Bintabara, Deogratus; Tokyo Medical and Dental University, Global Health Entrepreneurship; The University of Dodoma, Public Health Nakamura, Keiko; Tokyo Medical and Dental University, Global Health Entrepreneurship Seino, Kaoruko; Tokyo Medical and Dental University, Global Health Entrepreneurship; Michigan University, Department of Environmental Health Sciences, School of Public Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Global health, Health policy, Health services research
Keywords:	HEALTH SERVICES ADMINISTRATION & MANAGEMENT, access to health care, women, Tanzania

SCHOLARONE™  
Manuscripts

Only

1  
2  
3 1 **Improving access to health care for women in Tanzania by addressing socioeconomic**  
4  
5 2 **determinants and health insurance**  
6

7  
8 3 **Deogratus Bintabara<sup>1,2</sup>, Keiko Nakamura<sup>1</sup>, Kaoruko Seino<sup>1,3</sup>**  
9

10  
11 4 **Deogratus Bintabara, M.D.**

12  
13  
14 5 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

15  
16 6 Graduate School of Tokyo Medical and Dental University,

17  
18 7 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.

19  
20 8 <sup>2</sup>Department of Public Health, College of Health Sciences,

21  
22 9 The University of Dodoma, P.O Box 259, Dodoma, Tanzania

23  
24  
25 10 Phone: (255) 767 966 114, Email: bintabaradeo@gmail.com  
26  
27

28  
29 11 **Keiko Nakamura, M.D., Ph.D., (Corresponding author)**

30  
31 12 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

32  
33 13 Graduate School of Tokyo Medical and Dental University, Email: nakamura.ith@tmd.ac.jp  
34

35  
36 14 **Kaoruko Seino, Ph.D.**

37  
38 15 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,

39  
40 16 Graduate School of Tokyo Medical and Dental University,

41  
42 17 <sup>3</sup>Department of Environmental Health Sciences, School of Public Health,

43  
44 18 University of Michigan, Ann Arbor, Michigan USA, Email: seino.ith@tmd.ac.jp  
45  
46  
47

48 19 **Keywords:** Health services administration and management, Access to healthcare, Women,  
49  
50 20 Tanzania

51  
52 21 **Word count:** 3655  
53  
54  
55  
56  
57

## 1 ABSTRACT

2 **Objective:** This study was performed to explore the factors associated with accumulation of  
3 multiple problems in accessing health care among women in Tanzania as an example of a low-  
4 income country.

5 **Design:** Population-based cross-sectional survey

6 **Setting:** Nationwide representative data for women of reproductive age obtained from the 2015  
7 – 2016 Tanzania Demographic Health Survey were analyzed.

8 **Primary outcome measures:** A composite variable, “problems in accessing health care,” with  
9 five (0 – 4) categories was created based on the number of problems reported: obtaining  
10 permission to go to the doctor, obtaining money to pay for advice or treatment, distance to a  
11 health facility, and not wanting to go alone. Respondents who reported fewer or more problems  
12 placed in lower and higher categories, respectively.

13 **Results:** A total of 13266 women aged 15 – 49 years, with a median age (IQR) of 27 (20 – 36)  
14 years were interviewed and included in the analysis. About two-third s (65.53%) of the  
15 respondents reported at least one of the four major problems in accessing health care.  
16 Furthermore, after controlling for other variables included in the final model, women without  
17 any type of health insurance, those belonging to the poorest class according to the wealth index,  
18 those who had not attended any type of formal education, those who were not employed for cash,  
19 each year of increasing age, and those who were divorced, separated, or widowed were  
20 associated with greater problems in accessing health care.

21



1           **Conclusion:** This study indicated the additive effects of barriers to health care in low-  
2 income countries such as Tanzania. Based on these results, improving uptake of health insurance  
3 and addressing social determinants of health are the first steps toward reducing women's  
4 problems associated with accessing health care.

### 5 **Strengths and limitations of this study**

- 6       ➤ This is the first study to show additive effects of problems in accessing health care and  
7       associated factors among women in Tanzania.
- 8       ➤ The study used a nationally representative sample with high response rate and robust  
9       sampling procedure.
- 10      ➤ As a cross-sectional study design was used, causality assumptions could not be made.  
11      Therefore, the results should be interpreted with caution.

## 1 INTRODUCTION

2 Despite the substantial decline in global maternal mortality ratio (MMR), low-income  
3 countries have not seen the same decline in MMR as those with higher incomes.<sup>1,2</sup> This situation  
4 is described as an “area of shameful failures of development.”<sup>3,4</sup> Current statistics show that the  
5 MMR has increased significantly by more than 20% over the past 5 years in Tanzania, despite  
6 the governmental efforts to strengthen the health system by increasing the coverage of health  
7 care facilities.<sup>5-7</sup> Each ward now has at least one dispensary and/or health center, each district  
8 has at least one hospital, while each region has at least one referral hospital. Regardless of  
9 differences in the levels of these facilities in terms of function, expertise, availability of services,  
10 and population coverage, all are expected to provide basic maternal health services together with  
11 basic emergency obstetric care.<sup>7</sup> This resulted in an increase in coverage of maternal health  
12 services, such as antenatal care by skilled providers (96% – 98%), delivery at a health care  
13 facility (50% – 63%), and births assisted by skilled providers (51% – 64%) between 2010 and  
14 2016.<sup>5,6</sup> Persistent high MMR (556 maternal deaths per 100000 live births) in Tanzania raises  
15 concerns about whether the country can achieve the Sustainable Development Goals (SDGs)  
16 target of less than 140 maternal deaths per 100000 live births by 2030.<sup>8,9</sup> The majority of  
17 avoidable and unnecessary maternal deaths experienced in this region likely result from poor  
18 utilization of skilled maternal health services.<sup>10</sup>

19 Many social, cultural, and geographical factors as well as education level and poverty  
20 have been reported to play roles in the poor utilization of health services.<sup>11-13</sup> Access to health  
21 care has been highlighted as the major barrier toward the utilization of maternal health services  
22 in low-income countries, especially in sub-Saharan Africa (SSA).<sup>10,14,15</sup> Access to health care can  
23 be broadly defined based on availability, affordability, accessibility and acceptability,<sup>16</sup> but is

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2 Despite agreement that access to health care must be universal and guaranteed for all on an  
3 equitable basis,<sup>17</sup> women continue to face significant inequities in accessing and utilizing health  
4 care particularly in low-income countries.<sup>18</sup>

5 In relation to the problems experienced by women in accessing health care, the following  
6 four major problems have been addressed in previous studies: obtaining permission,<sup>19,20</sup>  
7 obtaining money,<sup>21</sup> distance to the health facility,<sup>22</sup> and not wanting to go alone (lack of spouse  
8 or family member escort).<sup>23,24</sup> Although it is unclear whether women with multiple problems  
9 encounter greater difficulties in accessing health care, most previous studies assessed and  
10 discussed each of these four problems independently. The limited evidence regarding whether  
11 women facing multiple problems have less access to health care suggested the need to create a  
12 composite variable that includes all four problems to identify groups of women at a greater  
13 disadvantage. A similar approach has been used to assess the severity of problems in accessing  
14 health care among individuals with disability in four African countries.<sup>25</sup> Moreover, recent  
15 studies indicated that age, education, residence, possession of health insurance, socioeconomic  
16 status, and occupation are strongly linked to access to health care.<sup>26–30</sup> The present study was  
17 performed to explore the factors associated with accumulation of multiple problems in accessing  
18 health care among women in Tanzania.

## 19 **METHODS**

### 20 **Data sources**

21 The present study used data from the 2015 – 2016 Tanzania Demographic Health Survey  
22 and Malaria Indicator Survey (TDHS-MIS) conducted by the National Bureau of Statistics

1 (NBS) and the Office of Chief Government Statistician (OCGS), Zanzibar, in collaboration with  
2 the Ministry of Health, Community Development, Gender, Elderly, and Children (MoHCDGEC)  
3 of the Tanzania Mainland and the Ministry of Health (MoH), Zanzibar. The technical support for  
4 the surveys was provided by ICF International under DHS program.

## 5 **Study design**

6 This study analyzed a nationwide population-based cross-sectional survey using  
7 information obtained by interviewing women (15 – 49 years old) who were either residents or  
8 visitors in the household on the night before the survey.

## 9 **Sample size and sampling technique**

10 The 2015 – 2016 TDHS-MIS used a two-stage cluster sampling technique to obtain a  
11 sample designed to provide nationally representative results according to all 30 regions of  
12 Tanzania. In the first stage, sample points (a total of 608 clusters) consisting of enumeration  
13 areas delineated for the 2012 Tanzania Population and Housing Census were selected. In the  
14 second stage, households were selected systematically. A complete listing of households was  
15 established for all 608 selected clusters prior to the fieldwork. From this list, 22 households were  
16 then systematically selected from each cluster, yielding a representative probability sample of  
17 13376 households. All eligible women in the selected households and men in subsample of one –  
18 third of selected households between the ages of 15 and 49 years who were either residents or  
19 visitors in the household on the night before the survey were then interviewed. Finally, a total of  
20 13266 women and 3514 men were interviewed.<sup>6</sup>

## 21 **Data collection and processing**

22 The 2015 – 2016 TDHS-MIS used four main types of questionnaires during data  
23 collection. However, only data collected with the Women's Questionnaire were used in the

1 present study. After pre-testing of the questionnaires, the finalized and corrected version was  
2 used in the main survey from August 22, 2015, through February 14, 2016. Data collection was  
3 performed by 64 female nurses who were trained and qualified to be interviewers through a  
4 series of practical tests and examinations. Following the training, 16 teams were formed (three  
5 for Zanzibar and 13 for Tanzania Mainland). Data entry was performed concurrently with data  
6 collection in the field. After the paper questionnaires were completed, edited, and checked by  
7 both the field editor and the supervisor, the data were entered into a tablet equipped with a data  
8 entry program. A 100% double entry data entry process was used to minimize keying errors, and  
9 editing was completed on March 21, 2016, while data cleaning and finalization were completed  
10 on April 22, 2016.

### 11 **Measurement of variables**

12 **Outcome variable:** In this survey women were asked whether each of the following four  
13 factors was a problem in seeking medical advice or treatment when they were ill: obtaining  
14 permission to go to the doctor/health facility; obtaining money to pay for advice, consultation or  
15 treatment; distance to the health facility; and not wanting to go alone. A new composite variable  
16 called “problems in accessing health care” was then created based on the number of problems  
17 reported, with respondents reporting fewer or more problems placed in lower and higher  
18 categories, respectively. The categories were assigned as 1, 2, 3, 4, and 5 for women who  
19 reported “no,” “one,” “two,” “three,” and “four” problems in accessing health care, respectively.  
20 These categories of a composite (outcome) variable were treated as ordinal numbers, with the  
21 assumption that conceptual differences between categories were identical.

22 **Independent variables:** Several independent variables that have been linked both  
23 empirically and theoretically with the accessibility of health care among women were included in

1 the present study. The respondents were divided into groups according to age as follows: 15 – 19,  
2 20 – 34, and 35 – 49 years. Women were grouped according to marital status as “never married,”  
3 “married/living together,” and “divorced, separated, or widowed.” Education level was classified  
4 as “none,” “primary,” “secondary,” and “higher” (including college and all university levels).  
5 Employment in the last 12 months was grouped into “not employed,” “employed for cash,” and  
6 “employed but paid-in-kind.” The area of residence was grouped into “urban” and “rural.”  
7 Possession of health insurance was grouped as “no” for women who did not have any type of  
8 health insurance and “yes” to those who had any type of health insurance. The wealth index was  
9 computed based on household assets and housing characteristics. During computation, the  
10 households were given scores based on the number and kinds of consumer goods they owned,  
11 ranging from a television to a bicycle or car, plus housing characteristics, such as the source of  
12 drinking water, toilet facilities, and flooring materials. These scores were derived using principal  
13 component analysis. National wealth quintiles were compiled by assigning the household score  
14 to each usual (de jure) household member, ranking each person in the household by their score,  
15 and then dividing the distribution into five equal categories, each with 20% of the population, as  
16 “poorest,” “poorer,” “middle,” “richer,” and “richest.”<sup>6</sup> The selection of these variables was  
17 based on previous studies.<sup>26–30</sup>

## 18 **Statistical approaches**

19 In descriptive analyses, categorical variables were summarized using proportions and  
20 then presented in tables, while quantitative variables were summarized using the median and  
21 interquartile range (IQR).

22 As the outcome variable “problems in accessing health care” was ordinal in nature (a  
23 score based on the number of different reported problems), in which the order of values

1 corresponded to a hierarchy in meaning as in this study, the application of ordered logistic  
2 regression was recommended.<sup>31</sup> Stata 14 (StataCorp, College Station, TX) was used for analysis  
3 in the present study. For all analyses, the Stata survey set commands were used to adjust for the  
4 variability of clustering and all the estimates were weighted to correct for non-responses and  
5 disproportionate sampling.

6 When assessing the associations between selected independent variables and the outcome  
7 variable, four models were tested for fit: a proportional ordered logistic model, a generalized  
8 ordered logistic regression model, and a generalized ordered logistic regression model with and  
9 without alternative parameterization. The generalized ordered logistic regression model with  
10 alternative parameterization was chosen as the best fit. This model allows some variables to be  
11 modeled with the proportional odds assumption while the parallel line constraint is relaxed for  
12 variables in which the assumption is not met. The model is less restrictive as it allows the  
13 coefficient of the variables to vary for the different categories that are compared.<sup>32</sup> The model  
14 provides gamma coefficients that show the extent to which the parallel regression assumption is  
15 violated by the variable.

## 16 **Ethics statement**

17 This study was based on an analysis of existing public domain survey data sets that are  
18 freely available online with all identifier information detached. The original TDHS-MIS  
19 protocols were reviewed by the Institution Review Board (IRB) of ICF Macro at Calverton in the  
20 USA and by the National Institute of Medical Research (NIMR) IRB in Tanzania. The ICF IRB  
21 ensured that the survey complied with the US Department of Health and Human Services  
22 regulations for the protection of human subjects (45 CFR 46), while NIMR-IRB ensured that the  
23 surveys complied with the laws and norms of Tanzania. The participants were adequately

1 informed about all relevant aspects of the survey, including its objective and interview  
 2 procedures. All participants accepted participating in the study signed informed consent prior to  
 3 the interviews.

#### 4 **Patient and Public Involvement Statement**

5 Patients and the public were not involved in the analysis of this study.

## 6 **RESULTS**

### 7 **Respondents' characteristics**

8 As shown in Table 1, a total of 13266 women between 15 and 49 years old were  
 9 interviewed and included in the analysis. The median age (IQR) of the respondents was 27 (20 –  
 10 36) years. About 62% of the respondents were living with their spouse at the time of the  
 11 interview. Less than 2% had attained the highest level of education (college or university).  
 12 Nearly two-thirds were employed but paid-in-kind and one-tenth reported having any type of  
 13 health insurance. Almost half of the respondents reported that obtaining money for health care  
 14 was the major problem in accessing health care. Furthermore, about two-thirds of the  
 15 respondents reported at least one of the four problems in accessing health care.

16 **Table 1 Percent distribution of women between the ages 15 – 49 years by selected**  
 17 **background characteristics, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

Variable	<i>n</i> (%) (Weighted)
<b>Age</b> (Median (IQR)=27, (20-36))	
15-19	2904 (21.89)
20-34	6360 (47.94)
35-49	4002 (30.17)
<b>Marital status</b>	
Never married	3353 (25.27)
Married/living together	8210 (61.89)
Divorced/separated/widowed	1703 (12.84)



<b>Education</b>		
None	1947	(14.67)
Primary	8211	(61.90)
Secondary	2925	(22.05)
Highest	183	(1.38)
<b>Employed last 12 months</b>		
Not employed	3033	(22.86)
Employed for cash	6197	(46.71)
Employed but paid-in-kind	4036	(30.43)
<b>Residence</b>		
Urban	4811	(36.27)
Rural	8455	(63.73)
<b>Health insurance ownership</b>		
Yes	12066	(90.95)
No	1200	(9.05)
<b>Health quintile</b>		
Lowest	2246	(16.93)
Second	2274	(17.14)
Middle	2328	(17.55)
Fourth	2822	(21.27)
Highest	3596	(27.11)
<b>Types of problems*</b>		
Obtaining money	6565	(49.49)
Distance to facility	5615	(42.33)
Not want to go alone	3962	(29.87)
Obtaining permission	1900	(14.32)
<b>Number of problems in accessing health care</b>		
None	4574	(34.48)
One problem	3291	(24.81)
Two problems	2547	(19.20)
Three problems	1759	(13.26)
Four problems	1095	(8.25)

1 Note: \*n and % do not add up to 13266 and 100%, respectively, because multiple responses were possible.

## 2 **Partial generalized ordered logistic regression model with alternative gamma** 3 **parameterization**

4 Table 2 shows the results of the partially constrained generalized ordered logistic  
5 regression model with alternative (gamma) parameterization for the outcome variable of  
6 problems in accessing health care. The results showed non-significant Wald test statistics,  
7 indicating that the model did not violate the proportional odds/parallel regression assumptions.

1  
2  
3 1 However, constraints for parallel lines were not imposed for age, wealth status (richest), and  
4  
5 2 marital status (divorced, separated, or widowed). The remaining variables that met the parallel  
6  
7 3 assumption can be interpolated in the same manner as for the ordered logistic regression model  
8  
9 4 as follows. The odds of being in higher categories of problems in accessing health care versus  
10  
11 5 lower or equal to a reference category were 0.622 [0.531 - 0.731] times lower for women who  
12  
13 6 had health insurance than those who did not, given constant values for the other variables in the  
14  
15 7 model. Furthermore, the odds of being in higher categories of problems in accessing health care  
16  
17 8 versus less than or equal to a reference category were 1.220 [1.067 - 1.395] times higher for  
18  
19 9 women who were employed but paid-in-kind than those who were unemployed for the last 12  
20  
21 10 months before the survey keeping the other variables constant in the model.  
22  
23  
24  
25

26 11 The variables for which constraints for parallel lines were not imposed were interpreted  
27  
28 12 as follows: the coefficients for age and marital status (divorced, separated, or widowed) were  
29  
30 13 consistently positive, while those for wealth status (richest) were negative but decreased across  
31  
32 14 the cut-points. Therefore, for each year of increase in age and being divorced, separated, or  
33  
34 15 widowed, women were more likely to report having a larger number of problems in accessing  
35  
36 16 health care. The greatest differences were seen with increasing age and for those who were  
37  
38 17 divorced, separated, or widowed, women were less likely to report having few problems in  
39  
40 18 accessing health care. In addition, the women in the richest economic group tended to be less  
41  
42 19 likely to report having many problems in accessing health care than those in the poorest group,  
43  
44 20 with the greatest differences because the richest women were less likely to report having many  
45  
46 21 problems in accessing health care.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 **Table 2 Generalized ordered logistic regression model with alternative gamma**  
 2 **parameterization, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

	<b>Variable</b>	<b>POR [95% CI]</b>	<b>P-value</b>
<b>Beta</b>	<b>Health insurance</b> (ref: No)		
	Yes	0.622 [0.531 - 0.731]	0.000
	<b>Residence</b> (ref: Urban)		
	Rural	0.858 [0.728 - 1.012]	0.069
	<b>Age</b> (continuous)	1.010 [1.001 - 1.017]	0.000
	<b>Marital status</b> ref: Never married)		
	Married/living together	0.901 [0.801 - 1.014]	0.085
	Divorced/separated/widowed	1.418 [1.175 - 1.712]	0.000
	<b>Education</b> (ref: None)		
	Primary	0.883 [0.788 - 0.990]	0.033
	Secondary	0.683 [0.582 - 0.800]	0.000
	Highest	0.516 [0.360 - 0.741]	0.000
	<b>Wealth status</b> (ref: Poorest)		
	Poorer	0.854 [0.726 - 1.006]	0.059
Middle	0.725 [0.626 - 0.840]	0.000	
Richer	0.496 [0.417 - 0.590]	0.000	
Richest	0.291 [0.233 - 0.364]	0.000	
<b>Employed last 12 months</b> (ref: Not employed)			
Employed for cash	0.975 [0.869 - 1.095]	0.668	
Employed but paid in kind	1.220 [1.067 - 1.395]	0.004	
<b>Gamma_2</b>	Age	0.993 [0.989 - 0.998]	0.000
	Wealth status (Richest)	1.279 [1.140 - 1.435]	0.000
	Marital status (Divorced/separated/widowed)	0.814 [0.701 - 0.945]	0.007
<b>Gamma_3</b>	Age	0.993 [0.986 - 0.999]	0.018
	Wealth status (Richest)	1.515 [1.265 - 1.814]	0.000
	Marital status (Divorced/separated/widowed)	0.749 [0.625 - 0.899]	0.002
<b>Gamma_4</b>	Age	0.987 [0.978 - 0.996]	0.005
	Wealth status (Richest)	1.957 [1.508 - 2.540]	0.000
	Marital status (Divorced/separated/widowed)	0.566 [0.419 - 0.764]	0.000

1 Note: Wald test of parallel lines assumption for the final model:  $F(33, 517) = 1.110, P = 0.310$ .

2 A non-significant test statistic indicates that the final model does not violate the proportional  
3 odds/parallel lines assumption.

4 \*POR = Proportional odds ratio.

## 5 DISCUSSION

6 This study was performed to explore the factors associated with accumulation of multiple  
7 problems in accessing health care among women in Tanzania. To our knowledge, this is the first  
8 study to explore the factors associated with multiple problems in accessing health care.  
9 Furthermore, the study used a nationally representative sample from Tanzania with the  
10 application of generalized ordered logistic regression models, which provided the best models  
11 for ordinal data to validate the factors associated with problems in accessing health care. In the  
12 present study, about 65%, 40%, and 20% of women reported “one or more,” “two or more,” and  
13 “three or more” major problems in accessing health care, respectively. In addition, not having  
14 health insurance and low socioeconomic status as measured by wealth, education, and  
15 employment status were associated with accumulation of multiple problems in accessing health  
16 care.

17 The high proportion of women reporting problems in accessing health care in this study  
18 was consistent with the findings of a previous study performed in Egypt.<sup>33</sup> These findings  
19 provide evidence that demand-side barriers, such as cost of care, permission from their spouse,  
20 lack of someone to escort to a health care facility, and distance to the facility,<sup>34,35</sup> still prevent  
21 many African women from accessing health care. Due to the cultural, social, and traditional  
22 perceptions in Africa that maternal health is only the responsibility of women, existing and new

1 interventions should influence health service utilization to begin at the individual, household, and  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 interventions should influence health service utilization to begin at the individual, household, and  
2 community levels to eliminate such demand-side barriers.<sup>21,36</sup>

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

3 Having health insurance is an essential element for timely access to health care and better  
4 health-related outcomes.<sup>37,38</sup> Despite the availability of the National Health Insurance Fund  
5 (NHIF) in Tanzania since 2001 through Act no. 8 of 1999, more than 90% of women are still  
6 uninsured.<sup>27,39</sup> Less than one-tenth of the women in the present study reported having any type of  
7 health insurance. Moreover, the women who had health insurance were less likely to report  
8 having multiple problems associated with access to health care. This may have been because  
9 having health insurance makes someone not only more comfortable with receiving a wide range  
10 of services but also ensures a wider choice regarding where and when to obtain health care  
11 without being afraid of the costs as they are covered by insurance. Similar findings were reported  
12 in a study conducted Ghana.<sup>26</sup> The similarity in the findings between these studies may have  
13 been because both used secondary data collected by a DHS program that applied a similar  
14 methodology. Furthermore, the participants in these two studies were from SSA, and therefore  
15 may have similar socioeconomic determinants.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

16 The present study indicated a strong association between being in the poorest class of the  
17 wealth index and accumulation of multiple problems in accessing health care among women in  
18 Tanzania. This finding may be explained by the fact that being in the poorest class requires  
19 individuals to spend their income on basic needs, such as food, and health care costs may  
20 therefore be less likely to be affordable.<sup>40</sup> Such women are therefore more likely to report having  
21 many problems in accessing health care, as reported in other studies conducted in SSA.<sup>41-43</sup>

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

22 In contrast to our expectations, we found no significant association between  
23 unemployment and problems in accessing health care. However, this may have been because

1 being employed is not enough to have full access to health care as there are other barriers  
2 preventing women from accessing health care, such as gender inequality, poor infrastructure, and  
3 lack of knowledge regarding maternal health services.<sup>28,33,44</sup> On the other hand, the results  
4 presented here indicated that women who are employed and receive wages in the form of  
5 payment-in-kind, such as food, clothes, and other goods instead of cash, were likely to  
6 experience multiple problems in accessing health care.

7 The variables age, wealth status (richest), and marital status (divorced, separated, or  
8 widowed) were found to vary for each category of the outcome variable. The findings indicated  
9 that the accumulation of multiple problems in accessing health care was associated with older  
10 age and being divorced, separated, or widowed, consistent with the results of previous studies in  
11 other low-income countries.<sup>45,46</sup> Older age is more likely to be accompanied by decreased  
12 working capability, and hence low income, being retired, and uninsured.<sup>47</sup> As the risk of  
13 maternal complications increased with older age<sup>48,49</sup> and living without a spouse reduced the  
14 chance of having an escort to the health facility, efforts and support should be made to provide  
15 such women with access to health care.

16 This study had some limitations, including the risk of misclassification bias, which may  
17 have been introduced due to the lack of external validation of self-reported information that  
18 could have affected categorization of the outcome variable. However, we reduced this effect by  
19 categorizing the outcome variable into five groups and the use of a generalized ordered logistics  
20 regression model that clearly validated the factors associated with problems in accessing health  
21 care among women in low-income countries. In addition, causality assumptions could not be  
22 made due to the cross-sectional nature of the study, and therefore the results should be  
23 interpreted with caution.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1

For peer review only

## 1 CONCLUSION

2 The results of the present study provided evidence for additive effects of barriers to  
3 health care in low-income countries, such as Tanzania. Based on these results, improving access  
4 to health insurance and addressing social determinants of health represent the first steps toward  
5 reducing problems associated with accessing health care for women in low-income countries.

## 6 Contributors

7 DB originated the design of the study, performed statistical analysis, interpretation and  
8 drafted the manuscript. KN contributed to the design of the study and the interpretation advice of  
9 data. KS contributed to interpretation advice of data and drafted the manuscript. Both DB, KN &  
10 KS critically revised the draft manuscript. All authors read and approved the final manuscript.

## 11 Acknowledgment

12 We would like to acknowledge ICF International, Rockville, Maryland, USA, through  
13 DHS program for giving us permission to access the TDHS-MIS 2015-2016 dataset.

## 14 Funding

15 This research received no specific grant from any funding agency in the public,  
16 commercial or not-for-profit sectors.

## 17 Competing interest

18 None to declare.

## 19 Provenance and peer review

20 Not commissioned; externally peer reviewed.

## 21 A data sharing statement



1           The datasets used for the current analysis was generated from the original survey of  
2  
3  
4           1           The datasets used for the current analysis was generated from the original survey of  
5  
6           2           Tanzania DHS-MIS datasets available from within the DHS program repository:  
7  
8           3           <http://dhsprogram.com/data/available-datasets.cfm>  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

1  
2  
3 **1 REFERENCES**  
4

- 5  
6 2 1 Alkema L, Chou D, Hogan D, *et al.* Global, regional, and national levels and trends in  
7  
8 3 maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a  
9  
10 4 systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*  
11  
12 5 2016;**387**:462–74. doi:10.1016/S0140-6736(15)00838-7  
13  
14  
15 6 2 World Health Organization (WHO). Trends in Maternal Mortality: 1990 to 2015:  
16  
17 7 estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations  
18  
19 8 Population Division. 2015. doi:ISBN 978 92 4 150363 1  
20  
21  
22 9 3 Godefay H, Byass P, Graham WJ, *et al.* Risk Factors for Maternal Mortality in Rural  
23  
24 10 Tigray, Northern Ethiopia: A Case-Control Study. *PLoS One* 2015;**10**:e0144975.  
25  
26 11 doi:10.1371/journal.pone.0144975  
27  
28  
29 12 4 Lewis G. Maternal mortality in the developing world: why do mothers really die? *Obstet*  
30  
31 13 *Med Med Pregnancy* 2008;**1**:2–6. doi:10.1258/om.2008.080019  
32  
33  
34 14 5 National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2011. Tanzania  
35  
36 15 Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro.  
37  
38 16 [http://www.measuredhs.com/pubs/pdf/FR243/FR243\[24June2011\].pdf](http://www.measuredhs.com/pubs/pdf/FR243/FR243[24June2011].pdf) Accessed 6 May  
39  
40 17 2018.  
41  
42  
43 18 6 Ministry of Health, Community Development, Gender, Elderly and Children  
44  
45 19 (MoHCDGEC) [Tanzania Mainland], Ministry of Health (MoH) [Zanzibar], National  
46  
47 20 Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF.  
48  
49 21 2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-*  
50  
51 22 *MIS) 2015-16*. Dar es Salaam, Tanzania, and Rockville, Maryland, USA: MoHCDGEC,  
52  
53 23 MoH, NBS, OCGS, and ICF; 2016. <https://dhsprogram.com/pubs/pdf/FR321/FR321.pdf>.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Accessed 6 May 2018.  
4  
5 2 7 United Republic of Tanzania (URT) and World Health Organization (WHO). Health  
6  
7 Systems Profile in Tanzania. URT; 2004.  
8 3  
9  
10 4 <http://mbaralidc.go.tz/storage/app/media/uploaded-files/Health-tanzania.pdf> Accessed 6  
11  
12 May 2018.  
13 5  
14 6 8 World Health Organization (WHO). Strategies toward ending preventable maternal  
15  
16 mortality ( EPMM ). Published Online First: 2015. doi:ISBN 978 92 4 150848 3  
17 7  
18 8 9 Moran AC, Jolivet RR, Chou D, *et al.* A common monitoring framework for ending  
19  
20 preventable maternal mortality, 2015–2030: phase I of a multi-step process. *BMC*  
21  
22 *Pregnancy Childbirth* 2016;**16**:250. doi:10.1186/s12884-016-1035-4  
23 10  
24 11 10 Odetola TD. Health care utilization among rural women of child-bearing age: a nigerian  
25  
26 experience. *Pan Afr Med J* 2015;**20**:1–7. doi:10.11604/pamj.2015.20.151.5845  
27 12  
28 13 11 Babalola SO. Factors associated with use of maternal health services in Haiti: a multilevel  
29  
30 analysis. *Rev Panam Salud Publica* 2014;**36**:1–9.  
31 14  
32 15 12 Ngomane S, Mulaudzi FM. Indigenous beliefs and practices that influence the delayed  
33  
34 attendance of antenatal clinics by women in the Bohlabele district in Limpopo, South  
35  
36 Africa. *Midwifery* 2012;**28**:30–8. doi:10.1016/j.midw.2010.11.002  
37 17  
38 18 13 Ali HS, Abdalla A, Abdalla A. Understand Factors Influencing Accessibility of Pregnant  
39  
40 Women to Antenatal Care Services Accessibility factors : Demographic characteristics of  
41  
42 the study. *Heal Sci J* 2016;**10**:1–5. doi:10.4172/1791-809X.1000100507  
43 19  
44 20 21 14 Victora CG, Barros AJ, Axelson H, *et al.* How changes in coverage affect equity in  
45  
46 maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of  
47  
48 national surveys. *Lancet* 2012;**380**:1149–56. doi:10.1016/S0140-6736(12)61427-5  
49 21  
50  
51 22  
52  
53 23  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 15 Singh PK, Kumar C, Rai RK, *et al.* Factors associated with maternal healthcare services  
4 utilization in nine high focus states in India: a multilevel analysis based on 14 385  
5  
6 2  
7 communities in 292 districts. *Health Policy Plan* 2014;**29**:542–59.  
8  
9 doi:10.1093/heapol/czt039  
10  
11 4  
12 5 16 Donnell OO. Access to health care in developing countries : breaking down demand side  
13 barriers. *Cad Saude Publica* 2007;**23**:2820–34. doi:S0102-311X2007001200003 [pii]  
14  
15 6  
16 7 17 Kirby N. Access to healthcare services as a human right. *Med Law* 2010;**29**:487–96.  
17  
18 8 18 Ganle JK, Parker M, Fitzpatrick R, *et al.* Inequities in accessibility to and utilisation of  
19 maternal health services in Ghana after user-fee exemption: a descriptive study. *Int J*  
20  
21  
22 9  
23  
24 10  
25  
26 11 19 Danforth E, Kruk M, Rockers P, *et al.* Household Decision-making about Delivery in  
27 Health Facilities: Evidence from Tanzania. *J Heal Popul Nutr* 2009;**27**:696–703.  
28  
29 12  
30  
31 13  
32  
33 14 20 Wilunda C, Scanagatta C, Putoto G, *et al.* Barriers to utilisation of antenatal care services  
34 in South Sudan: a qualitative study in Rumbek North County. *Reprod Health* 2017;**14**:65.  
35  
36 15  
37  
38 16  
39  
40 17 21 Lowe M, Chen D-R, Huang S-L. Social and Cultural Factors Affecting Maternal Health in  
41 Rural Gambia: An Exploratory Qualitative Study. *PLoS One* 2016;**11**:e0163653.  
42  
43 18  
44  
45 19  
46  
47 20 22 Mbiza CR, Kazembe A, Simwaka A. Barriers to health-seeking practices during  
48 pregnancy among adolescents in rural Blantyre, Malawi. *Afr J Midwifery Womens Health*  
49  
50 21  
51  
52 22  
53  
54 23 23 Kumbani L, Bjune G, Chirwa E, *et al.* Why some women fail to give birth at health

- 1 facilities: a qualitative study of women's perceptions of perinatal care from rural Southern  
2  
3 1 facilities: a qualitative study of women's perceptions of perinatal care from rural Southern  
4 Malawi. *Reprod Health* 2013;**10**:9. doi:10.1186/1742-4755-10-9  
5  
6 2  
7  
8 3 24 Vidler M, Ramadurg U, Charantimath U, *et al*. Utilization of maternal health care services  
9 and their determinants in Karnataka State, India. *Reprod Health* 2016;**13**:37.  
10  
11  
12 5  
13  
14  
15 6 25 Eide AH, Mannan H, Khogali M, *et al*. Perceived Barriers for Accessing Health Services  
16 among Individuals with Disability in Four African Countries. *PLoS One*  
17  
18  
19 8  
20  
21  
22 9 26 Browne JL, Kayode GA, Arhinful D, *et al*. Health insurance determines antenatal,  
23  
24 10  
25  
26 11  
27  
28 12 27 Kibusi SM, Sunguya BF, Kimunai E, *et al*. Health insurance is important in improving  
29  
30  
31 13  
32  
33 14  
34  
35 15  
36  
37  
38 16 28 Tsawe M, Susuman A. Determinants of access to and use of maternal health care services  
39  
40 17  
41  
42 18  
43  
44 19 29 Mrisho M, Schellenberg JA, Mushi AK, *et al*. Factors affecting home delivery in rural  
45  
46  
47 20  
48  
49 21 30 Kyei-Nimakoh M, Carolan-Olah M, McCann T V. Access barriers to obstetric care at  
50  
51 22  
52  
53  
54 23  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 31 Quddus MA, Wang C, Ison SG. Road Traffic Congestion and Crash Severity:  
4  
5 Econometric Analysis Using Ordered Response Models. *J Transp Eng* 2010;**136**:424–35.  
6  
7 doi:10.1061/(ASCE)TE.1943-5436.0000044  
8  
9  
10 4 32 Williams R. Understanding and interpreting generalized ordered logit models. *J Math*  
11  
12 *Sociol* 2016;**40**:7–20. doi:10.1080/0022250X.2015.1112384  
13  
14  
15 6 33 Chiang C, Labeeb SA, Higuchi M, *et al.* Barriers to the use of basic health services among  
16  
17 women in rural southern Egypt (Upper Egypt). *Nagoya J Med Sci* 2013;**75**:225–31.  
18  
19 8 34 Mselle LT, Kohi TW. Healthcare access and quality of birth care: narratives of women  
20  
21 living with obstetric fistula in rural Tanzania. *Reprod Health* 2016;**13**:87.  
22  
23 doi:10.1186/s12978-016-0189-x  
24  
25  
26 11 35 Shartz A, Long SK, Benatar S. Health Care Costs Are a Barrier to Care for Many  
27  
28 Women. Urban Inst. 2015.[http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf)  
29  
30 [to-Care-for-Many-Women.pdf](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf) (accessed 5 May 2018).  
31  
32  
33 14 36 Azuh D, Fayomi O, Ajayi, Lady. Socio-Cultural Factors of Gender Roles in Women's  
34  
35 Healthcare Utilization in Southwest Nigeria. *Open J Soc Sci* 2015;**3**:105–17.  
36  
37 doi:10.4236/jss.2015.34013  
38  
39  
40 17 37 Hsia J, Kemper E, Sofaer S, *et al.* Is Insurance a More Important Determinant of  
41  
42 Healthcare Access Than Perceived Health? Evidence from the Women's Health Initiative.  
43  
44 *J Womens Health Gend Based Med* 2000;**9**:881–9. doi:10.1089/152460900750020919  
45  
46  
47 20 38 Sommers BD, Gawande AA, Baicker K. Health Insurance Coverage and Health — What  
48  
49 the Recent Evidence Tells Us. *N Engl J Med* 2017;**377**:586–93.  
50  
51 doi:10.1056/NEJMsbl706645  
52  
53  
54 23 39 Marwa CW. Provision of National Health Insurance Fund Services to Its Members ; Pain  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 or Gain ? *Unified J Sport Heal Sci* 2016;**2**:1–6.  
4  
5 2 40 Yaya S, Bishwajit G, Shah V. Wealth, education and urban–rural inequality and maternal  
6  
7 healthcare service usage in Malawi. *BMJ Glob Heal* 2016;**1**:e000085. doi:10.1136/bmjgh-  
8  
9 2016-000085  
10  
11 4  
12 5 41 Shiferaw S, Spigt M, Godefrooij M, *et al.* Why do women prefer home births in Ethiopia?  
13  
14 *BMC Pregnancy Childbirth* 2013;**13**:5. doi:10.1186/1471-2393-13-5  
15  
16 6  
17 7 42 Lakew S, Tachbele E, Gelibo T. Predictors of skilled assistance seeking behavior to  
18  
19 pregnancy complications among women at southwest Ethiopia: a cross-sectional  
20  
21 community based study. *Reprod Health* 2015;**12**:109. doi:10.1186/s12978-015-0102-z  
22  
23 9  
24 10 43 Mwangome FK, Holding PA, Songola KM, *et al.* Barriers to hospital delivery in a rural  
25  
26 setting in coast province, Kenya: Community attitude and behaviours. *Rural Remote*  
27  
28 *Health* 2012;**12**. doi:22471588  
29  
30 12  
31 13 44 Bintabara D, Mpembeni RNM, Mohamed AA. Knowledge of obstetric danger signs  
32  
33 among recently-delivered women in Chamwino district, Tanzania: a cross-sectional study.  
34  
35 *BMC Pregnancy Childbirth* 2017;**17**:276. doi:10.1186/s12884-017-1469-3  
36  
37 15  
38 16 45 Tey N-P, Lai S. Correlates of and Barriers to the Utilization of Health Services for  
39  
40 Delivery in South Asia and Sub-Saharan Africa. *Sci World J* 2013;**2013**:1–11.  
41  
42 18  
43 doi:10.1155/2013/423403  
44  
45 19 46 Cheptum J, Gitonga M, Mutua E, *et al.* Barriers to Access and Utilization of Maternal and  
46  
47 Infant Health Services in Migori , Kenya. *Iiste* 2014;**4**:48–53.  
48  
49 21 47 Hadley J, Waidmann T. Health Insurance and Health at Age 65: Implications for Medical  
50  
51 Care Spending on New Medicare Beneficiaries. *Health Serv Res* 2006;**41**:429–51.  
52  
53 22  
54 23 doi:10.1111/j.1475-6773.2005.00491.x  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 48 Ziadeh S, Yahaya A. Pregnancy outcome at age 40 and older. *Arch Gynecol Obstet*  
4  
5 2001;**265**:30–3. doi:10.1007/s004040000122  
6  
7  
8 3 49 Illah E, Mbaruku G, Masanja H, *et al.* Causes and risk factors for maternal mortality in  
9  
10 4 rural Tanzania - case of Rufiji Health and Demographic Surveillance Site (HDSS). *Afr J*  
11  
12 5 *Reprod Health* 2013;**17**:119–30.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only



**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
<b>Title and abstract</b>	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	Page 2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Page 5
Objectives	3	State specific objectives, including any prespecified hypotheses	Page 7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Page 8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 8-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Page 8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page 9-10
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 9-10
Bias	9	Describe any efforts to address potential sources of bias	Page 3, 8
Study size	10	Explain how the study size was arrived at	Page 8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page 10-12
		(b) Describe any methods used to examine subgroups and interactions	Page 10-12
		(c) Explain how missing data were addressed	N/A
		(d) If applicable, describe analytical methods taking account of sampling strategy	Page 10
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page 13
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page 14 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers of outcome events or summary measures	Page 14 and Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2,3,4
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Table 4
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 18-9, Table 4
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 24
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 24
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 24
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	N/A

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Improving access to health care for women in Tanzania by addressing socioeconomic determinants and health insurance: a population-based cross-sectional survey

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023013.R4
Article Type:	Research
Date Submitted by the Author:	08-Aug-2018
Complete List of Authors:	Bintabara, Deogratus; Tokyo Medical and Dental University, Global Health Entrepreneurship; The University of Dodoma, Public Health Nakamura, Keiko; Tokyo Medical and Dental University, Global Health Entrepreneurship Seino, Kaoruko; Tokyo Medical and Dental University, Global Health Entrepreneurship; Michigan University, Department of Environmental Health Sciences, School of Public Health
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Global health, Health policy, Health services research
Keywords:	HEALTH SERVICES ADMINISTRATION & MANAGEMENT, access to health care, women, Tanzania

SCHOLARONE™  
Manuscripts

Only

1  
2  
3 1 **Improving access to health care for women in Tanzania by addressing socioeconomic**  
4  
5 2 **determinants and health insurance: a population-based cross-sectional survey**  
6  
7

8 3 **Deogratus Bintabara<sup>1,2</sup>, Keiko Nakamura<sup>1</sup>, Kaoruko Seino<sup>1,3</sup>**  
9  
10

11 4 **Deogratus Bintabara, M.D.**

12 5 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
13

14 6 Graduate School of Tokyo Medical and Dental University,  
15

16 7 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.  
17

18 8 <sup>2</sup>Department of Public Health, College of Health Sciences,  
19

20 9 The University of Dodoma, P.O Box 259, Dodoma, Tanzania  
21

22 10 Phone: (255) 767 966 114, Email: bintabaradeo@gmail.com  
23  
24  
25

26 11 **Keiko Nakamura, M.D., Ph.D., (Corresponding author)**

27 12 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
28

29 13 Graduate School of Tokyo Medical and Dental University, Email: nakamura.ith@tmd.ac.jp  
30  
31

32 14 **Kaoruko Seino, Ph.D.**

33 15 <sup>1</sup>Department of Global Health Entrepreneurship, Division of Public Health,  
34

35 16 Graduate School of Tokyo Medical and Dental University,  
36

37 17 <sup>3</sup>Department of Environmental Health Sciences, School of Public Health,  
38

39 18 University of Michigan, Ann Arbor, Michigan USA, Email: seino.ith@tmd.ac.jp  
40  
41

42 19 **Keywords:** Health services administration and management, Access to healthcare, Women,  
43  
44 20 Tanzania  
45

46 21 **Word count:** 3655  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1 ABSTRACT

2 **Objective:** This study was performed to explore the factors associated with accumulation of  
3 multiple problems in accessing health care among women in Tanzania as an example of a low-  
4 income country.

5 **Design:** Population-based cross-sectional survey

6 **Setting:** Nationwide representative data for women of reproductive age obtained from the 2015  
7 – 2016 Tanzania Demographic Health Survey were analyzed.

8 **Primary outcome measures:** A composite variable, “problems in accessing health care,” with  
9 five (0 – 4) categories was created based on the number of problems reported: obtaining  
10 permission to go to the doctor, obtaining money to pay for advice or treatment, distance to a  
11 health facility, and not wanting to go alone. Respondents who reported fewer or more problems  
12 placed in lower and higher categories, respectively.

13 **Results:** A total of 13266 women aged 15 – 49 years, with a median age (IQR) of 27 (20 – 36)  
14 years were interviewed and included in the analysis. About two-thirds (65.53%) of the  
15 respondents reported at least one of the four major problems in accessing health care.  
16 Furthermore, after controlling for other variables included in the final model, women without  
17 any type of health insurance, those belonging to the poorest class according to the wealth index,  
18 those who had not attended any type of formal education, those who were not employed for cash,  
19 each year of increasing age, and those who were divorced, separated, or widowed were  
20 associated with greater problems in accessing health care.

21

1           **Conclusion:** This study indicated the additive effects of barriers to health care in low-  
2 income countries such as Tanzania. Based on these results, improving uptake of health insurance  
3 and addressing social determinants of health are the first steps toward reducing women's  
4 problems associated with accessing health care.

### 5 **Strengths and limitations of this study**

- 6       ➤ This is the first study to show additive effects of problems in accessing health care and  
7       associated factors among women in Tanzania.
- 8       ➤ The study used a nationally representative sample with high response rate and robust  
9       sampling procedure.
- 10      ➤ As a cross-sectional study design was used, causality assumptions could not be made.  
11      Therefore, the results should be interpreted with caution.

## 1 INTRODUCTION

2 Despite the substantial decline in global maternal mortality ratio (MMR), low-income  
3 countries have not seen the same decline in MMR as those with higher incomes.<sup>1,2</sup> This situation  
4 is described as an “area of shameful failures of development.”<sup>3,4</sup> Current statistics show that the  
5 MMR has increased significantly by more than 20% over the past 5 years in Tanzania, despite  
6 the governmental efforts to strengthen the health system by increasing the coverage of health  
7 care facilities.<sup>5-7</sup> Each ward now has at least one dispensary and/or health center, each district  
8 has at least one hospital, while each region has at least one referral hospital. Regardless of  
9 differences in the levels of these facilities in terms of function, expertise, availability of services,  
10 and population coverage, all are expected to provide basic maternal health services together with  
11 basic emergency obstetric care.<sup>7</sup> This resulted in an increase in coverage of maternal health  
12 services, such as antenatal care by skilled providers (96% – 98%), delivery at a health care  
13 facility (50% – 63%), and births assisted by skilled providers (51% – 64%) between 2010 and  
14 2016.<sup>5,6</sup> Persistent high MMR (556 maternal deaths per 100000 live births) in Tanzania raises  
15 concerns about whether the country can achieve the Sustainable Development Goals (SDGs)  
16 target of less than 140 maternal deaths per 100000 live births by 2030.<sup>8,9</sup> The majority of  
17 avoidable and unnecessary maternal deaths experienced in this region likely result from poor  
18 utilization of skilled maternal health services.<sup>10</sup>

19 Many social, cultural, and geographical factors as well as education level and poverty  
20 have been reported to play roles in the poor utilization of health services.<sup>11-13</sup> Access to health  
21 care has been highlighted as the major barrier toward the utilization of maternal health services  
22 in low-income countries, especially in sub-Saharan Africa (SSA).<sup>10,14,15</sup> Access to health care can  
23 be broadly defined based on availability, affordability, accessibility and acceptability,<sup>16</sup> but is

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 simply referred to as the timely use of health services to achieve the desired health outcomes.  
2 Despite agreement that access to health care must be universal and guaranteed for all on an  
3 equitable basis,<sup>17</sup> women continue to face significant inequities in accessing and utilizing health  
4 care particularly in low-income countries.<sup>18</sup>

5 In relation to the problems experienced by women in accessing health care, the following  
6 four major problems have been addressed in previous studies: obtaining permission,<sup>19,20</sup>  
7 obtaining money,<sup>21</sup> distance to the health facility,<sup>22</sup> and not wanting to go alone (lack of spouse  
8 or family member escort).<sup>23,24</sup> Although it is unclear whether women with multiple problems  
9 encounter greater difficulties in accessing health care, most previous studies assessed and  
10 discussed each of these four problems independently. The limited evidence regarding whether  
11 women facing multiple problems have less access to health care suggested the need to create a  
12 composite variable that includes all four problems to identify groups of women at a greater  
13 disadvantage. A similar approach has been used to assess the severity of problems in accessing  
14 health care among individuals with disability in four African countries.<sup>25</sup> Moreover, recent  
15 studies indicated that age, education, residence, possession of health insurance, socioeconomic  
16 status, and occupation are strongly linked to access to health care.<sup>26–30</sup> The present study was  
17 performed to explore the factors associated with accumulation of multiple problems in accessing  
18 health care among women in Tanzania.

## 19 **METHODS**

### 20 **Data sources**

21 The present study used data from the 2015 – 2016 Tanzania Demographic Health Survey  
22 and Malaria Indicator Survey (TDHS-MIS) conducted by the National Bureau of Statistics



1 (NBS) and the Office of Chief Government Statistician (OCGS), Zanzibar, in collaboration with  
2 the Ministry of Health, Community Development, Gender, Elderly, and Children (MoHCDGEC)  
3 of the Tanzania Mainland and the Ministry of Health (MoH), Zanzibar. The technical support for  
4 the surveys was provided by ICF International under DHS program.

## 5 **Study design**

6 This study analyzed a nationwide population-based cross-sectional survey using  
7 information obtained by interviewing women (15 – 49 years old) who were either residents or  
8 visitors in the household on the night before the survey.

## 9 **Sample size and sampling technique**

10 The 2015 – 2016 TDHS-MIS used a two-stage cluster sampling technique to obtain a  
11 sample designed to provide nationally representative results according to all 30 regions of  
12 Tanzania. In the first stage, sample points (a total of 608 clusters) consisting of enumeration  
13 areas delineated for the 2012 Tanzania Population and Housing Census were selected. In the  
14 second stage, households were selected systematically. A complete listing of households was  
15 established for all 608 selected clusters prior to the fieldwork. From this list, 22 households were  
16 then systematically selected from each cluster, yielding a representative probability sample of  
17 13376 households. All eligible women in the selected households and men in subsample of one –  
18 third of selected households between the ages of 15 and 49 years who were either residents or  
19 visitors in the household on the night before the survey were then interviewed. Finally, a total of  
20 13266 women and 3514 men were interviewed.<sup>6</sup>

## 21 **Data collection and processing**

22 The 2015 – 2016 TDHS-MIS used four main types of questionnaires during data  
23 collection. However, only data collected with the Women's Questionnaire were used in the

1 present study. After pre-testing of the questionnaires, the finalized and corrected version was  
2 used in the main survey from August 22, 2015, through February 14, 2016. Data collection was  
3 performed by 64 female nurses who were trained and qualified to be interviewers through a  
4 series of practical tests and examinations. Following the training, 16 teams were formed (three  
5 for Zanzibar and 13 for Tanzania Mainland). Data entry was performed concurrently with data  
6 collection in the field. After the paper questionnaires were completed, edited, and checked by  
7 both the field editor and the supervisor, the data were entered into a tablet equipped with a data  
8 entry program. A 100% double entry data entry process was used to minimize keying errors, and  
9 editing was completed on March 21, 2016, while data cleaning and finalization were completed  
10 on April 22, 2016.

### 11 **Measurement of variables**

12 **Outcome variable:** In this survey women were asked whether each of the following four  
13 factors was a problem in seeking medical advice or treatment when they were ill: obtaining  
14 permission to go to the doctor/health facility; obtaining money to pay for advice, consultation or  
15 treatment; distance to the health facility; and not wanting to go alone. A new composite variable  
16 called “problems in accessing health care” was then created based on the number of problems  
17 reported, with respondents reporting fewer or more problems placed in lower and higher  
18 categories, respectively. The categories were assigned as 1, 2, 3, 4, and 5 for women who  
19 reported “no,” “one,” “two,” “three,” and “four” problems in accessing health care, respectively.  
20 These categories of a composite (outcome) variable were treated as ordinal numbers, with the  
21 assumption that conceptual differences between categories were identical.

22 **Independent variables:** Several independent variables that have been linked both  
23 empirically and theoretically with the accessibility of health care among women were included in

1 the present study. The respondents were divided into groups according to age as follows: 15 – 19,  
2 20 – 34, and 35 – 49 years. Women were grouped according to marital status as “never married,”  
3 “married/living together,” and “divorced, separated, or widowed.” Education level was classified  
4 as “none,” “primary,” “secondary,” and “higher” (including college and all university levels).  
5 Employment in the last 12 months was grouped into “not employed,” “employed for cash,” and  
6 “employed but paid-in-kind.” The area of residence was grouped into “urban” and “rural.”  
7 Possession of health insurance was grouped as “no” for women who did not have any type of  
8 health insurance and “yes” to those who had any type of health insurance. The wealth index was  
9 computed based on household assets and housing characteristics. During computation, the  
10 households were given scores based on the number and kinds of consumer goods they owned,  
11 ranging from a television to a bicycle or car, plus housing characteristics, such as the source of  
12 drinking water, toilet facilities, and flooring materials. These scores were derived using principal  
13 component analysis. National wealth quintiles were compiled by assigning the household score  
14 to each usual (de jure) household member, ranking each person in the household by their score,  
15 and then dividing the distribution into five equal categories, each with 20% of the population, as  
16 “poorest,” “poorer,” “middle,” “richer,” and “richest.”<sup>6</sup> The selection of these variables was  
17 based on previous studies.<sup>26–30</sup>

## 18 **Statistical approaches**

19 In descriptive analyses, categorical variables were summarized using proportions and  
20 then presented in tables, while quantitative variables were summarized using the median and  
21 interquartile range (IQR).

22 As the outcome variable “problems in accessing health care” was ordinal in nature (a  
23 score based on the number of different reported problems), in which the order of values

1 corresponded to a hierarchy in meaning as in this study, the application of ordered logistic  
2 regression was recommended.<sup>31</sup> Stata 14 (StataCorp, College Station, TX) was used for analysis  
3 in the present study. For all analyses, the Stata survey set commands were used to adjust for the  
4 variability of clustering and all the estimates were weighted to correct for non-responses and  
5 disproportionate sampling.

6 When assessing the associations between selected independent variables and the outcome  
7 variable, four models were tested for fit: a proportional ordered logistic model, a generalized  
8 ordered logistic regression model, and a generalized ordered logistic regression model with and  
9 without alternative parameterization. The generalized ordered logistic regression model with  
10 alternative parameterization was chosen as the best fit. This model allows some variables to be  
11 modeled with the proportional odds assumption while the parallel line constraint is relaxed for  
12 variables in which the assumption is not met. The model is less restrictive as it allows the  
13 coefficient of the variables to vary for the different categories that are compared.<sup>32</sup> The model  
14 provides gamma coefficients that show the extent to which the parallel regression assumption is  
15 violated by the variable.

## 16 **Ethics statement**

17 This study was based on an analysis of existing public domain survey data sets that are  
18 freely available online with all identifier information detached. The original TDHS-MIS  
19 protocols were reviewed by the Institution Review Board (IRB) of ICF Macro at Calverton in the  
20 USA and by the National Institute of Medical Research (NIMR) IRB in Tanzania. The ICF IRB  
21 ensured that the survey complied with the US Department of Health and Human Services  
22 regulations for the protection of human subjects (45 CFR 46), while NIMR-IRB ensured that the  
23 surveys complied with the laws and norms of Tanzania. The participants were adequately

1 informed about all relevant aspects of the survey, including its objective and interview  
 2 procedures. All participants accepted participating in the study signed informed consent prior to  
 3 the interviews.

#### 4 **Patient and Public Involvement Statement**

5 Patients and the public were not involved in the analysis of this study.

## 6 **RESULTS**

### 7 **Respondents' characteristics**

8 As shown in Table 1, a total of 13266 women between 15 and 49 years old were  
 9 interviewed and included in the analysis. The median age (IQR) of the respondents was 27 (20 –  
 10 36) years. About 62% of the respondents were living with their spouse at the time of the  
 11 interview. Less than 2% had attained the highest level of education (college or university).  
 12 Nearly two-thirds were employed but paid-in-kind and one-tenth reported having any type of  
 13 health insurance. Almost half of the respondents reported that obtaining money for health care  
 14 was the major problem in accessing health care. Furthermore, about two-thirds of the  
 15 respondents reported at least one of the four problems in accessing health care.

16 **Table 1 Percent distribution of women between the ages 15 – 49 years by selected**  
 17 **background characteristics, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

Variable	<i>n</i> (%) (Weighted)
<b>Age</b> (Median (IQR)=27, (20-36))	
15-19	2904 (21.89)
20-34	6360 (47.94)
35-49	4002 (30.17)
<b>Marital status</b>	
Never married	3353 (25.27)
Married/living together	8210 (61.89)
Divorced/separated/widowed	1703 (12.84)

<b>Education</b>		
None	1947	(14.67)
Primary	8211	(61.90)
Secondary	2925	(22.05)
Highest	183	(1.38)
<b>Employed last 12 months</b>		
Not employed	3033	(22.86)
Employed for cash	6197	(46.71)
Employed but paid-in-kind	4036	(30.43)
<b>Residence</b>		
Urban	4811	(36.27)
Rural	8455	(63.73)
<b>Health insurance ownership</b>		
Yes	12066	(90.95)
No	1200	(9.05)
<b>Health quintile</b>		
Lowest	2246	(16.93)
Second	2274	(17.14)
Middle	2328	(17.55)
Fourth	2822	(21.27)
Highest	3596	(27.11)
<b>Types of problems*</b>		
Obtaining money	6565	(49.49)
Distance to facility	5615	(42.33)
Not want to go alone	3962	(29.87)
Obtaining permission	1900	(14.32)
<b>Number of problems in accessing health care</b>		
None	4574	(34.48)
One problem	3291	(24.81)
Two problems	2547	(19.20)
Three problems	1759	(13.26)
Four problems	1095	(8.25)

1 Note: \*n and % do not add up to 13266 and 100%, respectively, because multiple responses were possible.

## 2 Partial generalized ordered logistic regression model with alternative gamma 3 parameterization

4 Table 2 shows the results of the partially constrained generalized ordered logistic  
5 regression model with alternative (gamma) parameterization for the outcome variable of  
6 problems in accessing health care. The results showed non-significant Wald test statistics,  
7 indicating that the model did not violate the proportional odds/parallel regression assumptions.

1  
2  
3 1 However, constraints for parallel lines were not imposed for age, wealth status (richest), and  
4  
5 2 marital status (divorced, separated, or widowed). The remaining variables that met the parallel  
6  
7 3 assumption can be interpolated in the same manner as for the ordered logistic regression model  
8  
9 4 as follows. The odds of being in higher categories of problems in accessing health care versus  
10  
11 5 lower or equal to a reference category were 0.622 [0.531 - 0.731] times lower for women who  
12  
13 6 had health insurance than those who did not, given constant values for the other variables in the  
14  
15 7 model. Furthermore, the odds of being in higher categories of problems in accessing health care  
16  
17 8 versus less than or equal to a reference category were 1.220 [1.067 - 1.395] times higher for  
18  
19 9 women who were employed but paid-in-kind than those who were unemployed for the last 12  
20  
21 10 months before the survey keeping the other variables constant in the model.  
22  
23  
24  
25

26 11 The variables for which constraints for parallel lines were not imposed were interpreted  
27  
28 12 as follows: the coefficients for age and marital status (divorced, separated, or widowed) were  
29  
30 13 consistently positive, while those for wealth status (richest) were negative but decreased across  
31  
32 14 the cut-points. Therefore, for each year of increase in age and being divorced, separated, or  
33  
34 15 widowed, women were more likely to report having a larger number of problems in accessing  
35  
36 16 health care. The greatest differences were seen with increasing age and for those who were  
37  
38 17 divorced, separated, or widowed, women were less likely to report having few problems in  
39  
40 18 accessing health care. In addition, the women in the richest economic group tended to be less  
41  
42 19 likely to report having many problems in accessing health care than those in the poorest group,  
43  
44 20 with the greatest differences because the richest women were less likely to report having many  
45  
46 21 problems in accessing health care.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Table 2 Generalized ordered logistic regression model with alternative gamma parameterization, Tanzania DHS-MIS 2015 – 2016 (n = 13266)**

	<b>Variable</b>	<b>POR [95% CI]</b>	<b>P-value</b>
<b>Beta</b>	<b>Health insurance</b> (ref: No)		
	Yes	0.622 [0.531 - 0.731]	0.000
	<b>Residence</b> (ref: Urban)		
	Rural	0.858 [0.728 - 1.012]	0.069
	<b>Age</b> (continuous)	1.010 [1.001 - 1.017]	0.000
	<b>Marital status</b> ref: Never married)		
	Married/living together	0.901 [0.801 - 1.014]	0.085
	Divorced/separated/widowed	1.418 [1.175 - 1.712]	0.000
	<b>Education</b> (ref: None)		
	Primary	0.883 [0.788 - 0.990]	0.033
	Secondary	0.683 [0.582 - 0.800]	0.000
	Highest	0.516 [0.360 - 0.741]	0.000
	<b>Wealth status</b> (ref: Poorest)		
	Poorer	0.854 [0.726 - 1.006]	0.059
	Middle	0.725 [0.626 - 0.840]	0.000
Richer	0.496 [0.417 - 0.590]	0.000	
Richest	0.291 [0.233 - 0.364]	0.000	
<b>Employed last 12 months</b> (ref: Not employed)			
Employed for cash	0.975 [0.869 - 1.095]	0.668	
Employed but paid in kind	1.220 [1.067 - 1.395]	0.004	
<b>Gamma_2</b>	Age	0.993 [0.989 - 0.998]	0.000
	Wealth status (Richest)	1.279 [1.140 - 1.435]	0.000
	Marital status (Divorced/separated/widowed)	0.814 [0.701 - 0.945]	0.007
<b>Gamma_3</b>	Age	0.993 [0.986 - 0.999]	0.018
	Wealth status (Richest)	1.515 [1.265 - 1.814]	0.000
	Marital status (Divorced/separated/widowed)	0.749 [0.625 - 0.899]	0.002
<b>Gamma_4</b>	Age	0.987 [0.978 - 0.996]	0.005
	Wealth status (Richest)	1.957 [1.508 - 2.540]	0.000
	Marital status (Divorced/separated/widowed)	0.566 [0.419 - 0.764]	0.000



1 Note: Wald test of parallel lines assumption for the final model:  $F(33, 517) = 1.110, P = 0.310$ .

2 A non-significant test statistic indicates that the final model does not violate the proportional  
3 odds/parallel lines assumption.

4 \*POR = Proportional odds ratio.

## 5 DISCUSSION

6 This study was performed to explore the factors associated with accumulation of multiple  
7 problems in accessing health care among women in Tanzania. To our knowledge, this is the first  
8 study to explore the factors associated with multiple problems in accessing health care.  
9 Furthermore, the study used a nationally representative sample from Tanzania with the  
10 application of generalized ordered logistic regression models, which provided the best models  
11 for ordinal data to validate the factors associated with problems in accessing health care. In the  
12 present study, about 65%, 40%, and 20% of women reported “one or more,” “two or more,” and  
13 “three or more” major problems in accessing health care, respectively. In addition, not having  
14 health insurance and low socioeconomic status as measured by wealth, education, and  
15 employment status were associated with accumulation of multiple problems in accessing health  
16 care.

17 The high proportion of women reporting problems in accessing health care in this study  
18 was consistent with the findings of a previous study performed in Egypt.<sup>33</sup> These findings  
19 provide evidence that demand-side barriers, such as cost of care, permission from their spouse,  
20 lack of someone to escort to a health care facility, and distance to the facility,<sup>34,35</sup> still prevent  
21 many African women from accessing health care. Due to the cultural, social, and traditional  
22 perceptions in Africa that maternal health is only the responsibility of women, existing and new

1 interventions should influence health service utilization to begin at the individual, household, and  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 interventions should influence health service utilization to begin at the individual, household, and  
2 community levels to eliminate such demand-side barriers.<sup>21,36</sup>

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

3 Having health insurance is an essential element for timely access to health care and better  
4 health-related outcomes.<sup>37,38</sup> Despite the availability of the National Health Insurance Fund  
5 (NHIF) in Tanzania since 2001 through Act no. 8 of 1999, more than 90% of women are still  
6 uninsured.<sup>27,39</sup> Less than one-tenth of the women in the present study reported having any type of  
7 health insurance. Moreover, the women who had health insurance were less likely to report  
8 having multiple problems associated with access to health care. This may have been because  
9 having health insurance makes someone not only more comfortable with receiving a wide range  
10 of services but also ensures a wider choice regarding where and when to obtain health care  
11 without being afraid of the costs as they are covered by insurance. Similar findings were reported  
12 in a study conducted Ghana.<sup>26</sup> The similarity in the findings between these studies may have  
13 been because both used secondary data collected by a DHS program that applied a similar  
14 methodology. Furthermore, the participants in these two studies were from SSA, and therefore  
15 may have similar socioeconomic determinants.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

16 The present study indicated a strong association between being in the poorest class of the  
17 wealth index and accumulation of multiple problems in accessing health care among women in  
18 Tanzania. This finding may be explained by the fact that being in the poorest class requires  
19 individuals to spend their income on basic needs, such as food, and health care costs may  
20 therefore be less likely to be affordable.<sup>40</sup> Such women are therefore more likely to report having  
21 many problems in accessing health care, as reported in other studies conducted in SSA.<sup>41-43</sup>

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

22 In contrast to our expectations, we found no significant association between  
23 unemployment and problems in accessing health care. However, this may have been because

1 being employed is not enough to have full access to health care as there are other barriers  
2 preventing women from accessing health care, such as gender inequality, poor infrastructure, and  
3 lack of knowledge regarding maternal health services.<sup>28,33,44</sup> On the other hand, the results  
4 presented here indicated that women who are employed and receive wages in the form of  
5 payment-in-kind, such as food, clothes, and other goods instead of cash, were likely to  
6 experience multiple problems in accessing health care.

7 The variables age, wealth status (richest), and marital status (divorced, separated, or  
8 widowed) were found to vary for each category of the outcome variable. The findings indicated  
9 that the accumulation of multiple problems in accessing health care was associated with older  
10 age and being divorced, separated, or widowed, consistent with the results of previous studies in  
11 other low-income countries.<sup>45,46</sup> Older age is more likely to be accompanied by decreased  
12 working capability, and hence low income, being retired, and uninsured.<sup>47</sup> As the risk of  
13 maternal complications increased with older age<sup>48,49</sup> and living without a spouse reduced the  
14 chance of having an escort to the health facility, efforts and support should be made to provide  
15 such women with access to health care.

16 This study had some limitations, including the risk of misclassification bias, which may  
17 have been introduced due to the lack of external validation of self-reported information that  
18 could have affected categorization of the outcome variable. However, we reduced this effect by  
19 categorizing the outcome variable into five groups and the use of a generalized ordered logistics  
20 regression model that clearly validated the factors associated with problems in accessing health  
21 care among women in low-income countries. In addition, causality assumptions could not be  
22 made due to the cross-sectional nature of the study, and therefore the results should be  
23 interpreted with caution.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1

For peer review only

## 1 **CONCLUSION**

2 The results of the present study provided evidence for additive effects of barriers to  
3 health care in low-income countries, such as Tanzania. Based on these results, improving access  
4 to health insurance and addressing social determinants of health represent the first steps toward  
5 reducing problems associated with accessing health care for women in low-income countries.

## 6 **Contributors**

7 DB originated the design of the study, performed statistical analysis, interpretation and  
8 drafted the manuscript. KN contributed to the design of the study and the interpretation advice of  
9 data. KS contributed to interpretation advice of data and drafted the manuscript. Both DB, KN &  
10 KS critically revised the draft manuscript. All authors read and approved the final manuscript.

## 11 **Acknowledgment**

12 We would like to acknowledge ICF International, Rockville, Maryland, USA, through  
13 DHS program for giving us permission to access the TDHS-MIS 2015-2016 dataset.

## 14 **Funding**

15 This research received no specific grant from any funding agency in the public,  
16 commercial or not-for-profit sectors.

## 17 **Competing interest**

18 None to declare.

## 19 **Provenance and peer review**

20 Not commissioned; externally peer reviewed.

## 21 **A data sharing statement**

1  
2  
3 1 The datasets used for the current analysis was generated from the original survey of  
4  
5 2 Tanzania DHS-MIS datasets available from within the DHS program repository:  
6  
7 3 <http://dhsprogram.com/data/available-datasets.cfm>  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

1  
2  
3 **1 REFERENCES**  
4

- 5  
6 2 1 Alkema L, Chou D, Hogan D, *et al.* Global, regional, and national levels and trends in  
7  
8 3 maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a  
9  
10 4 systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*  
11  
12 5 2016;**387**:462–74. doi:10.1016/S0140-6736(15)00838-7  
13  
14  
15 6 2 World Health Organization (WHO). Trends in Maternal Mortality: 1990 to 2015:  
16  
17 7 estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations  
18  
19 8 Population Division. 2015. doi:ISBN 978 92 4 150363 1  
20  
21  
22 9 3 Godefay H, Byass P, Graham WJ, *et al.* Risk Factors for Maternal Mortality in Rural  
23  
24 10 Tigray, Northern Ethiopia: A Case-Control Study. *PLoS One* 2015;**10**:e0144975.  
25  
26 11 doi:10.1371/journal.pone.0144975  
27  
28  
29 12 4 Lewis G. Maternal mortality in the developing world: why do mothers really die? *Obstet*  
30  
31 13 *Med Med Pregnancy* 2008;**1**:2–6. doi:10.1258/om.2008.080019  
32  
33  
34 14 5 National Bureau of Statistics (NBS) [Tanzania] and ICF Macro. 2011. Tanzania  
35  
36 15 Demographic and Health Survey 2010. Dar es Salaam, Tanzania: NBS and ICF Macro.  
37  
38 16 [http://www.measuredhs.com/pubs/pdf/FR243/FR243\[24June2011\].pdf](http://www.measuredhs.com/pubs/pdf/FR243/FR243[24June2011].pdf) Accessed 6 May  
39  
40 17 2018.  
41  
42  
43 18 6 Ministry of Health, Community Development, Gender, Elderly and Children  
44  
45 19 (MoHCDGEC) [Tanzania Mainland], Ministry of Health (MoH) [Zanzibar], National  
46  
47 20 Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF.  
48  
49 21 2016. *Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-*  
50  
51 22 *MIS) 2015-16*. Dar es Salaam, Tanzania, and Rockville, Maryland, USA: MoHCDGEC,  
52  
53 23 MoH, NBS, OCGS, and ICF; 2016. <https://dhsprogram.com/pubs/pdf/FR321/FR321.pdf>.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 Accessed 6 May 2018.  
4  
5 2 7 United Republic of Tanzania (URT) and World Health Organization (WHO). Health  
6  
7 Systems Profile in Tanzania. URT; 2004.  
8 3  
9  
10 4 <http://mbaralidc.go.tz/storage/app/media/uploaded-files/Health-tanzania.pdf> Accessed 6  
11  
12 May 2018.  
13 5  
14 6 8 World Health Organization (WHO). Strategies toward ending preventable maternal  
15  
16 mortality ( EPMM ). Published Online First: 2015. doi:ISBN 978 92 4 150848 3  
17 7  
18 8 9 Moran AC, Jolivet RR, Chou D, *et al.* A common monitoring framework for ending  
19  
20 preventable maternal mortality, 2015–2030: phase I of a multi-step process. *BMC*  
21  
22 *Pregnancy Childbirth* 2016;**16**:250. doi:10.1186/s12884-016-1035-4  
23 10  
24 11 10 Odetola TD. Health care utilization among rural women of child-bearing age: a nigerian  
25  
26 experience. *Pan Afr Med J* 2015;**20**:1–7. doi:10.11604/pamj.2015.20.151.5845  
27 12  
28 13 11 Babalola SO. Factors associated with use of maternal health services in Haiti: a multilevel  
29  
30 analysis. *Rev Panam Salud Publica* 2014;**36**:1–9.  
31 14  
32 15 12 Ngomane S, Mulaudzi FM. Indigenous beliefs and practices that influence the delayed  
33  
34 attendance of antenatal clinics by women in the Bohlabele district in Limpopo, South  
35  
36 Africa. *Midwifery* 2012;**28**:30–8. doi:10.1016/j.midw.2010.11.002  
37 17  
38 18 13 Ali HS, Abdalla A, Abdalla A. Understand Factors Influencing Accessibility of Pregnant  
39  
40 Women to Antenatal Care Services Accessibility factors : Demographic characteristics of  
41  
42 the study. *Heal Sci J* 2016;**10**:1–5. doi:10.4172/1791-809X.1000100507  
43 19  
44 20 21 14 Victora CG, Barros AJ, Axelson H, *et al.* How changes in coverage affect equity in  
45  
46 maternal and child health interventions in 35 Countdown to 2015 countries: an analysis of  
47  
48 national surveys. *Lancet* 2012;**380**:1149–56. doi:10.1016/S0140-6736(12)61427-5  
49 21  
50 22  
51 23  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 1 15 Singh PK, Kumar C, Rai RK, *et al.* Factors associated with maternal healthcare services  
4 utilization in nine high focus states in India: a multilevel analysis based on 14 385  
5  
6 2  
7 communities in 292 districts. *Health Policy Plan* 2014;**29**:542–59.  
8  
9 doi:10.1093/heapol/czt039  
10  
11 4  
12 5 16 Donnell OO. Access to health care in developing countries : breaking down demand side  
13 barriers. *Cad Saude Publica* 2007;**23**:2820–34. doi:S0102-311X2007001200003 [pii]  
14  
15 6  
16 7 17 Kirby N. Access to healthcare services as a human right. *Med Law* 2010;**29**:487–96.  
17  
18 8 18 Ganle JK, Parker M, Fitzpatrick R, *et al.* Inequities in accessibility to and utilisation of  
19 maternal health services in Ghana after user-fee exemption: a descriptive study. *Int J*  
20  
21  
22 9  
23  
24 10  
25  
26 11 19 Danforth E, Kruk M, Rockers P, *et al.* Household Decision-making about Delivery in  
27 Health Facilities: Evidence from Tanzania. *J Heal Popul Nutr* 2009;**27**:696–703.  
28  
29 12  
30  
31 13  
32  
33 14 20 Wilunda C, Scanagatta C, Putoto G, *et al.* Barriers to utilisation of antenatal care services  
34 in South Sudan: a qualitative study in Rumbek North County. *Reprod Health* 2017;**14**:65.  
35  
36 15  
37  
38 16  
39  
40 17 21 Lowe M, Chen D-R, Huang S-L. Social and Cultural Factors Affecting Maternal Health in  
41 Rural Gambia: An Exploratory Qualitative Study. *PLoS One* 2016;**11**:e0163653.  
42  
43 18  
44  
45 19  
46  
47 20 22 Mbiza CR, Kazembe A, Simwaka A. Barriers to health-seeking practices during  
48 pregnancy among adolescents in rural Blantyre, Malawi. *Afr J Midwifery Womens Health*  
49  
50 21  
51  
52 22  
53  
54 23 23 Kumbani L, Bjune G, Chirwa E, *et al.* Why some women fail to give birth at health

- 1 facilities: a qualitative study of women's perceptions of perinatal care from rural Southern  
2  
3 1 facilities: a qualitative study of women's perceptions of perinatal care from rural Southern  
4 Malawi. *Reprod Health* 2013;**10**:9. doi:10.1186/1742-4755-10-9  
5  
6 2  
7  
8 3 24 Vidler M, Ramadurg U, Charantimath U, *et al*. Utilization of maternal health care services  
9 and their determinants in Karnataka State, India. *Reprod Health* 2016;**13**:37.  
10  
11  
12 5  
13  
14  
15 6 25 Eide AH, Mannan H, Khogali M, *et al*. Perceived Barriers for Accessing Health Services  
16 among Individuals with Disability in Four African Countries. *PLoS One*  
17  
18  
19 8  
20 2015;**10**:e0125915. doi:10.1371/journal.pone.0125915  
21  
22 9 26 Browne JL, Kayode GA, Arhinful D, *et al*. Health insurance determines antenatal,  
23 delivery and postnatal care utilisation: evidence from the Ghana Demographic and Health  
24  
25  
26 11  
27 Surveillance data. *BMJ Open* 2016;**6**:e008175. doi:10.1136/bmjopen-2015-008175  
28  
29 12 27 Kibusi SM, Sunguya BF, Kimunai E, *et al*. Health insurance is important in improving  
30 maternal health service utilization in Tanzania—analysis of the 2011/2012 Tanzania  
31  
32  
33 14  
34 HIV/AIDS and malaria indicator survey. *BMC Health Serv Res* 2018;**18**:112.  
35  
36 15  
37 doi:10.1186/s12913-018-2924-1  
38  
39 16 28 Tsawe M, Susuman A. Determinants of access to and use of maternal health care services  
40 in the Eastern Cape, South Africa: a quantitative and qualitative investigation. *BMC Res*  
41  
42  
43 18  
44 Notes 2014;**7**:723. doi:10.1186/1756-0500-7-723  
45  
46 19 29 Mrisho M, Schellenberg JA, Mushi AK, *et al*. Factors affecting home delivery in rural  
47 Tanzania. *Trop Med Int Heal* 2007;**12**:862–72. doi:10.1111/j.1365-3156.2007.01855.x  
48  
49 21 30 Kyei-Nimakoh M, Carolan-Olah M, McCann T V. Access barriers to obstetric care at  
50 health facilities in sub-Saharan Africa—a systematic review. *Syst Rev* 2017;**6**:110.  
51  
52  
53 23  
54 doi:10.1186/s13643-017-0503-x  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 31 Quddus MA, Wang C, Ison SG. Road Traffic Congestion and Crash Severity:  
4  
5 2 Econometric Analysis Using Ordered Response Models. *J Transp Eng* 2010;**136**:424–35.  
6  
7 3 doi:10.1061/(ASCE)TE.1943-5436.0000044  
8  
9  
10 4 32 Williams R. Understanding and interpreting generalized ordered logit models. *J Math*  
11  
12 5 *Sociol* 2016;**40**:7–20. doi:10.1080/0022250X.2015.1112384  
13  
14 6 33 Chiang C, Labeeb SA, Higuchi M, *et al.* Barriers to the use of basic health services among  
15  
16 7 women in rural southern Egypt (Upper Egypt). *Nagoya J Med Sci* 2013;**75**:225–31.  
17  
18 8 34 Mselle LT, Kohi TW. Healthcare access and quality of birth care: narratives of women  
19  
20 9 living with obstetric fistula in rural Tanzania. *Reprod Health* 2016;**13**:87.  
21  
22 10 doi:10.1186/s12978-016-0189-x  
23  
24 11 35 Shartz A, Long SK, Benatar S. Health Care Costs Are a Barrier to Care for Many  
25  
26 12 Women. Urban Inst. 2015.[http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf)  
27  
28 13 [to-Care-for-Many-Women.pdf](http://hrms.urban.org/briefs/Health-Care-Costs-Are-a-Barrier-to-Care-for-Many-Women.pdf) (accessed 5 May 2018).  
29  
30  
31 14 36 Azuh D, Fayomi O, Ajayi, Lady. Socio-Cultural Factors of Gender Roles in Women's  
32  
33 15 Healthcare Utilization in Southwest Nigeria. *Open J Soc Sci* 2015;**3**:105–17.  
34  
35 16 doi:10.4236/jss.2015.34013  
36  
37 17 37 Hsia J, Kemper E, Sofaer S, *et al.* Is Insurance a More Important Determinant of  
38  
39 18 Healthcare Access Than Perceived Health? Evidence from the Women's Health Initiative.  
40  
41 19 *J Womens Health Gend Based Med* 2000;**9**:881–9. doi:10.1089/152460900750020919  
42  
43 20 38 Sommers BD, Gawande AA, Baicker K. Health Insurance Coverage and Health — What  
44  
45 21 the Recent Evidence Tells Us. *N Engl J Med* 2017;**377**:586–93.  
46  
47 22 doi:10.1056/NEJMsbl706645  
48  
49 23 39 Marwa CW. Provision of National Health Insurance Fund Services to Its Members ; Pain  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 or Gain ? *Unified J Sport Heal Sci* 2016;**2**:1–6.  
4  
5 2 40 Yaya S, Bishwajit G, Shah V. Wealth, education and urban–rural inequality and maternal  
6  
7 healthcare service usage in Malawi. *BMJ Glob Heal* 2016;**1**:e000085. doi:10.1136/bmjgh-  
8  
9 2016-000085  
10  
11 4  
12 5 41 Shiferaw S, Spigt M, Godefrooij M, *et al.* Why do women prefer home births in Ethiopia?  
13  
14 *BMC Pregnancy Childbirth* 2013;**13**:5. doi:10.1186/1471-2393-13-5  
15  
16 6  
17 7 42 Lakew S, Tachbele E, Gelibo T. Predictors of skilled assistance seeking behavior to  
18  
19 pregnancy complications among women at southwest Ethiopia: a cross-sectional  
20  
21 community based study. *Reprod Health* 2015;**12**:109. doi:10.1186/s12978-015-0102-z  
22  
23 9  
24 10 43 Mwangome FK, Holding PA, Songola KM, *et al.* Barriers to hospital delivery in a rural  
25  
26 setting in coast province, Kenya: Community attitude and behaviours. *Rural Remote*  
27  
28 *Health* 2012;**12**. doi:22471588  
29  
30 12  
31 13 44 Bintabara D, Mpembeni RNM, Mohamed AA. Knowledge of obstetric danger signs  
32  
33 among recently-delivered women in Chamwino district, Tanzania: a cross-sectional study.  
34  
35 *BMC Pregnancy Childbirth* 2017;**17**:276. doi:10.1186/s12884-017-1469-3  
36  
37 15  
38 16 45 Tey N-P, Lai S. Correlates of and Barriers to the Utilization of Health Services for  
39  
40 Delivery in South Asia and Sub-Saharan Africa. *Sci World J* 2013;**2013**:1–11.  
41  
42 18  
43 doi:10.1155/2013/423403  
44  
45 19 46 Cheptum J, Gitonga M, Mutua E, *et al.* Barriers to Access and Utilization of Maternal and  
46  
47 Infant Health Services in Migori , Kenya. *Iiste* 2014;**4**:48–53.  
48  
49 21 47 Hadley J, Waidmann T. Health Insurance and Health at Age 65: Implications for Medical  
50  
51 Care Spending on New Medicare Beneficiaries. *Health Serv Res* 2006;**41**:429–51.  
52  
53 22  
54 23 doi:10.1111/j.1475-6773.2005.00491.x  
55  
56  
57  
58  
59  
60

- 1  
2  
3 1 48 Ziadeh S, Yahaya A. Pregnancy outcome at age 40 and older. *Arch Gynecol Obstet*  
4  
5 2001;**265**:30–3. doi:10.1007/s004040000122  
6  
7  
8 3 49 Illah E, Mbaruku G, Masanja H, *et al.* Causes and risk factors for maternal mortality in  
9  
10 4 rural Tanzania - case of Rufiji Health and Demographic Surveillance Site (HDSS). *Afr J*  
11  
12 5 *Reprod Health* 2013;**17**:119–30.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
<b>Title and abstract</b>	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	Page 2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Page 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Page 5
Objectives	3	State specific objectives, including any prespecified hypotheses	Page 7
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Page 8
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 8-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Page 8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Page 9-10
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Page 9-10
Bias	9	Describe any efforts to address potential sources of bias	Page 3, 8
Study size	10	Explain how the study size was arrived at	Page 8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Page 10-12
		(b) Describe any methods used to examine subgroups and interactions	Page 10-12
		(c) Explain how missing data were addressed	N/A
		(d) If applicable, describe analytical methods taking account of sampling strategy	Page 10
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Page 13
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Page 14 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers of outcome events or summary measures	Page 14 and Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2,3,4
		(b) Report category boundaries when continuous variables were categorized	Table 1
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Table 4
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 18-9, Table 4
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 24
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 24
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 24
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	N/A

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).