


BMJ Open Prevalence and correlates of smoking and nicotine dependence: results of a nationwide cross-sectional survey among Singapore residents

Shazana Shahwan ¹, Edimansyah Abdin,¹ Saleha Shafie,¹ Sherilyn Chang,¹ Rajeswari Sambasivam,¹ Yunjue Zhang,¹ Janhavi Ajit Vaingankar,¹ Yik Ying Teo,² Derrick Heng,³ Siow Ann Chong,¹ Mythily Subramaniam¹

To cite: Shahwan S, Abdin E, Shafie S, *et al.* Prevalence and correlates of smoking and nicotine dependence: results of a nationwide cross-sectional survey among Singapore residents. *BMJ Open* 2019;**9**:e032198. doi:10.1136/bmjopen-2019-032198

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-032198>).

Received 07 June 2019
Revised 23 August 2019
Accepted 19 September 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Research Division, Institute of Mental Health, Singapore, Singapore

²Saw Swee Hock School of Public Health, National University of Singapore, Singapore

³Epidemiology & Disease Control Division, Ministry of Health, Government of Singapore, Singapore, Singapore

Correspondence to
Ms Shazana Shahwan;
Shazana_MOHAMED_@SHAHWAN@imh.com.sg

ABSTRACT

Background Since the Singapore Mental Health Study in 2010 which reported a 16.0% prevalence rate for current smokers and 4.5% for nicotine dependence, new anti-smoking strategies have been implemented. The aim of this study was to compare smoking trends from the 2010 study with the second Singapore Mental Health Study in 2016 (SMHS 2016).

Methods A survey of 6126 individuals aged 18 years and above randomly selected among Singapore residents was conducted using the same methodology as the 2010 study. The measures used in this analysis were sociodemographic questions, the Composite International Diagnostic Interview which assessed for psychiatric disorders, the Fagerstrom Test for Nicotine Dependence and a list of chronic physical conditions that were prevalent in Singapore. Logistic regression analyses were used to test for associations between smoking/nicotine-dependence and other measures.

Results In the SMHS 2016, 16.1% were current smokers and 3.3% were nicotine-dependent. As compared with non-smokers, current smokers were more likely to be younger, male gender, of ethnic minority and had lower/vocational education level. Younger age, male gender, lower/vocational education and psychiatric disorders (major depression, bipolar disorder and alcohol use disorders) predicted nicotine dependence. No associations were found between nicotine dependence and any of the chronic conditions.

Conclusion The prevalence of current smokers in the population has plateaued while that of nicotine dependence has decreased from 2010. However, the study did not investigate the use of e-cigarettes. Inequalities in smoking and nicotine dependence continue to pervade the population particularly among those of ethnic minority, lower/vocational education and the mentally ill.

INTRODUCTION

Cigarette smoking is a leading cause of preventable death worldwide.¹ Smoking-related diseases contribute significantly to the global rise in incidence of non-communicable diseases in both developed and developing countries.² Smoking ranks high among

Strengths and limitations of this study

- The study reports the prevalence and correlates of smoking and nicotine dependence based on a large representative sample of the Singapore population.
- The same methodology and instruments employed in the 2010 study was applied allowing direct comparisons with the earlier study.
- The study did not include individuals below 18 years of age and did not assess for alternative forms of smoking such as e-cigarettes that are gaining popularity.
- Reliance on self-report could lead to an underestimation of the true prevalence of nicotine dependence and associations with mental and medical conditions.

public health problems in the world, with an estimated 7.4–9.7 million tobacco-attributable deaths by 2030.³

In 2003, the World Health Assembly adopted the Framework Convention on Tobacco Control (FCTC) to take steps to reduce both the supply of and demand for tobacco products. This treaty is now ratified by 181 countries.² To help those countries fulfil their commitment to the FCTC, the WHO disseminated recommendations consisting of six strategies: monitor tobacco use; protect people from tobacco smoke; offer help to quit tobacco use; warn about the dangers of tobacco; enforce bans on tobacco advertising, promotion and sponsorship; and raise taxes on tobacco. Steady decline in smoking prevalence rates has been witnessed in countries such as New Zealand,⁴ Turkey and Sweden⁵ since the implementation of FCTC strategies.

However, these changes have not occurred uniformly across all population groups. An upward trend in smoking debut in early adolescence was found in a European study.⁶

Disparities in smoking prevalence in underprivileged populations are also widening. In the USA, for example, less than 20% of those at or above the poverty level smoke compared with 30% of those below the poverty level. Other socioeconomic measures associated with inequalities in smoking include education, income and neighbourhood deprivation.⁷ The higher prevalence of smoking in individuals from lower socioeconomic status (SES) groups is the single most important cause of socioeconomic differences in mortality.⁸

Another subpopulation with exceptionally high rates of smoking internationally is individuals with mental illness. Individuals with mental illness smoke at rates approximately twice that of adults without mental disorders.⁹ Smoking is believed to account for the majority of excess mortality among individuals with serious mental illness.¹⁰ Life expectancy among people with severe mental illness is 25 years less than that among the general population.¹¹ Monitoring trends in the population and between subgroups aid countries in taking necessary corrections or new actions for tobacco control.

Singapore is a city-state located in Southeast Asia with a multi-ethnic population and was among the first 40 countries to ratify the FCTC. It exercises stringent smoking policies and extensive regulations on the demand and supply of tobacco.¹² In the nationwide Singapore Mental Health Study conducted in 2010 (SMHS 2010), Picco and colleagues¹³ reported local smoking prevalence rates of 16%. Smokers were more likely to be of younger age, male gender, Malay ethnicity and have lower education. Prevalence of nicotine dependence was higher in those with alcohol abuse and those experiencing chronic pain. Singapore aims to lower smoking prevalence rates to 12% by 2020 through a multipronged strategy composed of preventing initiation among the youth, public education and specific programmes for target groups, and providing more support and access to smoking cessation programmes.¹⁴

The purpose of this study was to compare and contrast smoking trends from the 2010 study with the second Singapore Mental Health Study which began in 2016 (SMHS 2016).¹⁵ As stricter anti-smoking laws (e.g., raising minimum smoking age to 21 years, expanding smoke-free zones) and new campaigns were launched after 2010, we hypothesise that there will be a decline in the prevalence of smoking and nicotine dependence in the SMHS 2016. This study also examined sociodemographic risk factors of smoking and nicotine dependence as well as the association of nicotine dependence with lifetime psychiatric and physical disorders.

METHODS

Participants and procedure

The SMHS 2016 was conducted between 2016 and 2018 following the same procedures as the SMHS 2010.¹⁶ This population-based, cross-sectional study included Singapore citizens and permanent residents aged 18 years

and above living in Singapore. The sampling frame was based on a national population registry of all citizens and permanent residents in Singapore, and is updated regularly. Individuals were randomly selected using a disproportionate stratified sampling design with 16 strata defined according to ethnicity (Chinese, Malay, Indian, Others) and age groups (18–34, 35–49, 50–64, 65 and above). Residents aged 65 and above, Malays, and Indians were over sampled to ensure that an adequate sample size would be achieved to improve the reliability of estimates for the subgroup analysis. We requested 15 907 records of Singapore residents. 11 100 records were eventually released in eight different batches. About 20% of these were ineligible cases (e.g., ineligible language, incorrect address) which were excluded from the response rate calculation. In all, 6126 respondents were interviewed, giving a response rate of 69.5%. Data on household structure were not collated and not accounted for in the analysis.

An invitation letter was sent to each respondent followed by a personal home visit by a trained interviewer to obtain his/her agreement to participate in the survey. Trained interviewers from a survey research company conducted face-to-face interviews with those who agreed to participate in the study. The questionnaires were available in English, Chinese and Malay. Residents who were incapable of doing an interview due to severe physical/mental conditions, language barriers; were living outside the country, institutionalised/hospitalised and those who were not contactable due to incomplete/incorrect address were excluded from the survey. Consent was obtained from all participants prior to commencement of any study procedure. Parental consent was also obtained for minors aged 18–20 years.

MEASURES

Sociodemographic information

Data on gender, age groups (18–34, 35–49, 50–64 and ≥65 years), ethnicity (Chinese, Malay, Indian and Others), marital status (single, married, divorced/separated or widowed), educational level (primary and below, secondary, vocational institute, pre-university/junior college, diploma and university), employment status (employed, unemployed and economically inactive) and household income were collected.

Psychiatric disorders

The WHO World Mental Health Composite International Diagnostic Interview is a structured instrument used to generate diagnoses of Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV) disorders using established algorithms with organic exclusion criteria and hierarchical rules. Modules on depression, mania, generalised anxiety disorder, obsessive compulsive disorder (OCD) and alcohol use were included in the survey.

Smoking and nicotine dependence

Information on smoking was collected through a question that asked participants whether they were current

smokers, ex-smokers or non-smokers who never smoked before. The 6-item Fagerstrom Test for Nicotine Dependence was used to assess physical dependence on tobacco smoking. Scores of 4 or less are classified as low dependence while scores of 8 to 10, as very high dependence. We categorised those with scores five and above as dependence as defined by previous studies,¹⁷ including our previous study¹³ to ensure consistency for comparison.

Chronic medical conditions

Respondents were asked to report whether 'a doctor ever told you that you have any of the following...'. This was followed by a list of 18 chronic medical conditions that are prevalent in Singapore in the form of a checklist. These disorders were reclassified into the following nine types of physical disorders: hypertension, hyperlipidemia, diabetes, asthma, chronic pain, cardiovascular diseases, ulcers, thyroid problems and cancer.

Statistical analysis

All estimates were weighted to adjust for over-sampling and post-stratified for age and ethnicity distributions between the survey sample and the Singapore resident population. Mean and SD were calculated for continuous variables, and frequencies and percentages for categorical variables. The sociodemographic characteristics were compared among the groups and tested for significant differences using χ^2 tests. This was followed by multiple logistic regression and multinomial logistic regression analyses to explore the sociodemographic correlates of nicotine dependence, and current or ex-smoking status. Gender-specific analyses were also conducted to compare the prevalence rates between 2010 and 2016 as well to explore sociodemographic correlates of nicotine dependence, current and ex-smoking status. Statistical significance was evaluated at the <0.05 level using two-sided tests. All statistical analyses were carried out using the Statistical Analysis Software System V.9.

Patient and public involvement

This research was done without patient involvement. Patients were not invited to comment on the study design and were not consulted to develop patient relevant outcomes or interpret the results. Patients were not invited to contribute to the writing or editing of this document for readability or accuracy.

RESULTS

Prevalence of smoking and nicotine dependence

Table 1 shows the sociodemographic characteristics of the sample classified by smoker, ex-smoker and non-smoker status. Table 2 summarises the prevalence rates in 2010 and 2016. Among the population, 16.1% were current smokers, 10.5% were ex-smokers, while 3.3% had nicotine dependence. The prevalence of smokers among men was 27.1% and among women it was 5.3%. Online supplementary table 1 provides further information of

the prevalence of current smokers, ex-smokers and nicotine dependence by age group and gender.

Sociodemographic correlates of nicotine dependence

Table 3 shows the sociodemographic correlates of current smokers and ex-smokers. As compared with non-smokers, those who were current smokers were more likely to be of Malay, Indian or Other ethnicity (vs Chinese), male gender, divorced/separated (compared with married), have lower education level (i.e., primary or secondary education) or vocational qualifications (i.e., polytechnic or technical education) (compared with university degree). Those in the older age groups (≥ 50 years compared with 18–34 years), economically inactive (compared with employed) and with a monthly household income of more than SGD 10,000 (compared with a monthly household income of SGD 2000) were less likely to be a current smoker. Compared with non-smokers, ex-smokers, similarly, were of Malay or other ethnicity, male gender, divorced/separated and had lower/vocational education level. Online supplementary table 2a,b present the sociodemographic correlates of current and ex-smokers of men and women, respectively. The results are consistent for both sexes with exception to income, where the highest income group (SGD 10,000 and above) was associated with lower odds of smoking in men, whereas the lower income group (SGD 2000–3999) was associated with lower odds of smoking in women. The relationship between marital status and smoking however was inconsistent with no association found for men and a lower odds of current smoking associated with being widowed in women (although being divorced/separated was associated with higher odds of being a current or ex-smoker for the overall sample).

Multiple logistic regression showed (table 4) that men had seven times higher risk of nicotine dependence than women. Furthermore, nicotine dependence was significantly higher in those with lower or vocational educational qualifications (compared with university education). Older age (≥ 50 years, compared with 18–34 years), being economically inactive (compared with employed) and monthly household income of SGD 4000–5999 (compared with less than SGD 2000) was associated with lower risk of nicotine dependence. Online supplementary table 3a,b present the sociodemographic correlates of nicotine dependence in men and women, respectively. The results for both sexes are consistent with that of the overall sample.

Relationship between nicotine dependence and psychiatric and physical disorders

Those with nicotine dependence were significantly more likely to have major depressive disorder (MDD), bipolar disorder, alcohol abuse and alcohol dependence. Gender difference was observed for OCD where women with nicotine dependence were six times more likely to have OCD than those without nicotine dependence but this

**Table 1** Sociodemographic profile of the sample by smoking status

Sociodemographic characteristics	N	Smoking status			Total (n=6107)	
		Smoker (n=1176)	Ex-smoker (n=750)	Non-smoker (n=4181)		
		Weighted percentage				
Age group (years) (Mean=45.2)	18–34	1707	35.1	23.7	30.5	30.4
	35–49	1496	30.8	29.9	29.2	29.6
	50–64	1626	27.1	28.4	26.6	26.9
	65+	1297	7.0	18.0	13.7	13.1
Gender	Female	3058	16.5	24.3	61.7	50.4
	Male	3068	83.5	75.7	38.3	49.6
Ethnicity	Chinese	1782	63.7	71.1	79.0	75.7
	Malay	1990	23.9	16.3	9.4	12.5
	Indian	1844	9.8	8.3	8.5	8.7
	Others	510	2.6	4.3	3.1	3.1
Marital status	Never married	1544	35.5	22.6	31.3	31
	Married	3843	54.7	66.8	59.8	59.8
	Divorced/separated	343	7.9	7.8	4.2	5.2
	Widowed	396	2.0	2.8	4.7	4.1
Education	Primary and below	1187	21.8	20.8	14.4	16.3
	Secondary	1648	29.6	25.3	21.2	23
	Pre-U/JC	304	3.2	4.6	6.8	6
	Vocational/ITE	508	15.9	8.8	3.8	6.3
	Diploma	1024	18.8	18.2	19.2	19
	University	1455	10.7	22.3	34.6	29.4
Employment	Employed	4055	83.4	73.5	69.3	72
	Economically inactive*	1716	9.3	22.0	25.8	22.7
	Unemployed	354	7.3	4.5	4.8	5.3
Household income (SGD/month)	Below 2000	1147	21.0	20.8	14.8	16.5
	2000–3999	1331	27.2	18.1	18.7	20
	4000–5999	1113	23.0	21.2	21.0	21.4
	6000–9999	1003	19.4	20.0	22.7	21.8
	10 000 and above	861	9.5	19.9	22.8	20.3

*Includes homemakers, students and retirees/pensioners.

ITE, Institute of Technical Education; JC, Junior College; Pre-U, Pre-University; SGD, Singapore dollars.

Table 2 Prevalence of Current smokers, Ex-smokers and Nicotine dependence in SMHS 2010 and SMHS 2016

	2010	2016	P value
Current smokers	16.0	16.1	n.s.
Male	27.0	27.1	n.s.
Female	5.6	5.3	n.s.
Ex-smokers	10.8	10.5	n.s.
Nicotine dependence	4.5	3.3	0.007

χ^2 analysis.

n.s., not statistically significant; SMHS, Singapore Mental Health Study.

association was not observed in men. These results are summarised in [table 5](#).

No associations were found between nicotine dependence and any of the chronic conditions.

DISCUSSION

The prevalence of smoking in the general population remained at 16% from our 2010 national survey.¹³ Prevalence rates in men and women, likewise, remained at about 27% in men and about 5% in women indicating a plateau in smoking prevalence. The sharpest decline occurred between the 1980s and the 2000s¹⁸ with local rates hovering around 15% in the past 10 years.¹³ However,

Table 3 Sociodemographic correlates of Current smokers and Ex-smokers

		Current smoker vs non-smoker			Ex-smoker vs non-smoker		
		OR	95% CI	P value	OR	95% CI	P value
Age group	18–34	Ref			Ref		
	35–49	0.9	(0.6 to 1.3)	0.48	1.1	(0.7 to 1.6)	0.84
	50–64	0.4	(0.2 to 0.6)	<0.0001	0.7	(0.5 to 1.2)	0.18
	65 and above	0.2	(0.1 to 0.4)	<0.0001	0.8	(0.5 to 1.4)	0.43
Gender	Male	8.8	(6.5 to 11.8)	<0.0001	5.5	(4.0 to 7.5)	<0.0001
	Female	Ref			Ref		
Ethnicity	Chinese	Ref			Ref		
	Malay	2.4	(1.9 to 3.0)	<0.0001	1.9	(1.5 to 2.5)	<0.0001
	Indian	1.4	(1.1 to 1.8)	0.004	1.1	(0.9 to 1.5)	0.32
	Others	1.7	(1.2 to 2.6)	0.008	2.0	(1.3 to 2.8)	0.001
Marital	Single	Ref			Ref		
	Married	0.9	(0.6 to 1.2)	0.42	1.3	(0.8 to 2.0)	0.27
	Divorced/separated	1.9	(1.1 to 3.5)	0.04	2.5	(1.3 to 4.8)	0.008
	Widowed	0.8	(0.3 to 1.8)	0.57	0.7	(0.3 to 1.8)	0.52
Education	Primary	13.8	(7.9 to 24.1)	<0.0001	3.8	(2.3 to 6.3)	<0.0001
	Secondary	7.3	(4.5 to 12.0)	<0.0001	2.6	(1.7 to 4.1)	<0.0001
	Pre-U/JC	1.7	(0.8 to 3.8)	0.19	1.5	(0.7 to 3.0)	0.26
	Vocational/ITE	7.8	(4.7 to 13.0)	<0.0001	3.1	(1.8 to 5.3)	<0.0001
	Diploma	3.1	(2.0 to 4.8)	<0.0001	1.8	(1.2 to 2.8)	0.008
	University	Ref			Ref		
Employment	Employed	Ref			Ref		
	Economically inactive	0.3	(0.2 to 0.5)	<0.0001	0.9	(0.6 to 1.3)	0.46
	Unemployed	0.9	(0.5 to 1.5)	0.65	0.9	(0.5 to 1.6)	0.8
Household income (SGD/month)	Below 2000	Ref			Ref		
	2000–3999	0.8	(0.6 to 1.2)	0.23	0.7	(0.5 to 1.1)	0.1
	4000–5999	0.8	(0.5 to 1.1)	0.16	0.8	(0.5 to 1.3)	0.43
	6000–9999	0.9	(0.6 to 1.3)	0.45	0.9	(0.5 to 1.4)	0.54
	10 000 and above	0.6	(0.3 to 1.0)	0.04	1.0	(0.6 to 1.6)	0.88

Multinomial logistic regression analysis controlled for potential confounders including age, ethnicity, marital status, education, employment and household income.

ITE, Institute of Technical Education; JC, junior college; Pre-U, Pre-University; SGD, Singapore dollars.

a desirable shift in nicotine dependence from 4.5% in 2010 to 3.3% in 2016 was observed. With one of the lowest smoking prevalence rates in the world, Singapore's challenge is to go beyond these rates to achieve the target set at 12% by 2020.¹² Novel endgame solutions such as prohibiting the sales of tobacco to citizens born after year 2000 and using plain packaging have been proposed.¹⁹

It was noteworthy that as many as one in four adult men are current smokers accounting for the vast majority of smokers in the country, with men being seven times more likely than women to be nicotine-dependent. More recently, Subramaniam and colleagues²⁰ through focus group discussions with Singaporean youths identified multiple personal (e.g., coping), social (e.g., for networking) and familial influences (e.g., early exposure) on young adults' smoking behaviours which provide

actionable information for further anti-smoking initiatives. Factors such as traditional values, normative gender expectations and economic independence have been purported for the wide margin of difference between the sexes.²¹

Not surprisingly, smoking and nicotine dependence groups were over-represented in those with lower or vocational qualifications and less likely to be associated with higher income. Marques-Vidal and colleagues²² suggested that those with higher levels of education are more responsive to social initiatives to cut down smoking and anti-smoking messages or have more contact with exemplary role models. Despite the rising cigarette prices/taxes with the average cost of a pack of 20 cigarettes priced at US\$9.66 (SGD\$13.31),²³ higher levels of smoking and nicotine dependence were observed among

**Table 4** Sociodemographic correlates of nicotine dependence

		OR	95% CI	P value
Age group	18–34	Ref		
	35–49	0.6	(0.3 to 1.1)	0.34
	50–64	0.3	(0.2 to 0.6)	0.001
	65 and above	0.3	(0.1 to 0.7)	0.007
Gender	Male	6.9	(3.6 to 13.2)	<0.0001
	Female	Ref		
Ethnicity	Chinese	Ref		
	Malay	1.5	(0.9 to 2.3)	0.1
	Indian	1.3	(0.8 to 2.0)	0.31
	Others	1.4	(0.6 to 3.2)	0.41
Marital	Single	Ref		
	Married	0.8	(0.5 to 1.5)	0.5
	Divorced/separated	0.8	(0.3 to 1.8)	0.57
	Widowed	1.7	(0.4 to 6.7)	0.43
Education	Primary	37.0	(8.3 to 165.1)	<0.0001
	Secondary	23.3	(5.8 to 92.6)	<0.0001
	Pre-U/JC	2.1	(0.4 to 9.8)	0.359
	Vocational/ITE	16.0	(4.0 to 63.8)	<0.0001
	Diploma	10.0	(2.7 to 37.0)	<0.0001
Employment	University	Ref		
	Employed	Ref		
	Economically inactive	0.1	(0.1 to 0.3)	<0.0001
Household income (SGD/month)	Unemployed	0.6	(0.3 to 1.4)	0.26
	Below 2000	Ref		
	2000–3900	0.8	(0.5 to 1.5)	0.56
	4000–5999	0.4	(0.2 to 0.8)	0.009
	6000–9999	0.6	(0.3 to 1.4)	0.24
10 000 and above	0.6	(0.2 to 1.4)	0.21	

ITE, Institute of Technical Education; JC, Junior College; Pre-U, Pre-University; SGD, Singapore dollars.

those with the lowest income levels suggesting alternative strategies are needed to reduce morbidity and mortality due to smoking for this group.

Two other sociodemographic factors that were associated with current smoking prevalence were age and ethnic minority status (Malay and Others ethnic groups). An age effect was observed with the prevalence of smoking being higher in the younger age groups despite the combined efforts of raising the minimum age for smoking, increasing cigarette prices and smoking prevention and cessation programmes in institutes of higher learning in the recent years. Shahwan and colleagues³⁴ through focus group discussions with youths identified various elements that were deemed to be efficacious in anti-smoking campaigns (e.g., positive tone, low-fear visual images, low ‘controlling’ language) which may be translated into continued efforts towards further reducing smoking rates in youths. With regard to ethnicity, almost all Malays in

Singapore are Muslims. While drinking alcohol is clearly forbidden in Islam, smoking is deemed by many Muslims as acceptable. Ethnic differences may also represent residual confounding by socioeconomic influences that have not been adequately controlled using our proxy measures.

The correlates of smoking and nicotine dependence identified in this study (i.e., the association between smoking and younger age, male gender, ethnic minority, lower/vocational education) are similar to the findings of the 2010 study, representing the stability and persistence of these factors.

The relationship between being economically inactive and smoking is less clear. While unemployment is defined as being out of work and actively seeking work, economic inactivity exists when a person is without any form of employment and is not actively seeking work. The majority of individuals in the economic inactivity group

Table 5 Prevalence and OR of other lifetime psychiatric disorders in people with nicotine dependence

Lifetime psychiatric disorders	Sample	%	SE	OR	95% CI	P value
MDD	Total	14.8	0.4	3.0	(1.7 to 5.5)	<0.0001
	Male	14.9	3.9	3.3	(1.7 to 6.5)	<0.0001
	Female	13.6	5.7	1.6	(0.6 to 4.4)	0.359
Dysthymia	Total	0.1	0.0	0.4	(0.1 to 3.5)	0.420
	Male	0.1	0.1	0.5	(0.06 to 4.9)	0.590
	Female	0.0	0.0	.	.	.
Bipolar disorder	Total	5.2	0.0	3.7	(1.6 to 8.8)	0.003
	Male	5.1	2.2	4.0	(1.5 to 10.8)	0.006
	Female	5.8	3.5	2.9	(0.7 to 11.2)	0.131
Generalised anxiety disorder	Total	1.6	0.0	1.0	(0.4 to 2.2)	0.960
	Male	1.3	0.6	0.7	(0.3 to 1.9)	0.540
	Female	4.9	3.1	2.5	(0.6 to 10.2)	0.210
OCD	Total	5.1	0.0	1.5	(0.6 to 3.7)	0.340
	Male	2.5	0.8	0.7	(0.4 to 1.6)	0.429
	Female	26.4	13.9	6.2	(1.5 to 24.6)	0.010
Alcohol abuse	Total	25.8	0.0	6.7	(4.0 to 11.3)	<0.0001
	Male	27.2	4.7	6.7	(3.9 to 11.4)	<0.0001
	Female	14.2	11.2	5.9	(0.9 to 37.2)	0.060
Alcohol dependence	Total	3.0	0.0	4.0	(1.4 to 11.5)	0.009
	Male	2.7	1.2	3.2	(1.0 to 10.0)	0.040
	Female	5.3	3.3	25.7	(5.8 to 113.8)	<0.0001

Multiple logistic regression analyses in total sample and by gender specification, adjusted for age. MDD, major depressive disorder; OCD, obsessive compulsive disorder; SE, standard error.

consist of housewives, retirees and students. As such, we speculate that there were several protective factors against smoking for this group such as higher education, spousal support, increasing health concerns with advancing age and the desire to improve longevity and quality of life.

Nicotine dependence was significantly associated with alcohol abuse as well as alcohol dependence. Nicotine-dependent individuals were about seven times more likely to abuse alcohol and four times more likely to be dependent on alcohol than those who were non-nicotine dependent. Psychosocial factors, such as risk-prone personality traits, greater opportunities and inclinations to drink, have been widely accepted as reasons for the well-documented link between smoking and alcoholism.²⁵ However, these psychosocial factors may not completely account for the association between smoking and alcohol problems. Some authors have speculated that the progression from the use of alcohol and tobacco to abuse may be facilitated by effects of early-stage use on central reward circuitry.²⁶

An association between nicotine dependence and MDD and bipolar disorder, which was not apparent in 2010, emerged in this study. The proportion of individuals with nicotine dependence who had MDD and bipolar disorder increased from 7% to 14% and 1% to 5%, respectively. The prevalence of nicotine dependence in patients with mood disorders has been reported to range from 50% to 70% compared with 25% in the general population in other studies.²⁷ The comorbidity between nicotine dependence

and mood disorders may be explained in at least two ways. First, various studies have demonstrated shared genetic and environmental influences. Second, it could also be argued that (i) depression increases the risks of smoking (i.e., through self-medication) or (ii) smoking increases the risk of depression.²⁸ Given that the proportion of MDD and bipolar disorder among individuals with nicotine dependence increased from 2010 to 2016, further exploration of this relationship is warranted.

A gender difference was found in the association between nicotine dependence and OCD where 26.4% of women with nicotine dependence had OCD compared with 2.5% for men. We identified three other studies that have similarly found an association between smoking and OCD only in women.^{29–31} However, our finding differs from the vast majority of clinical studies that have shown that patients with OCD are less likely to smoke compared with the general population.^{32–33} As suggested by Wu and colleagues, this may have to do with differences between clinical and community samples and further research is needed to shed light on the association between nicotine dependence and OCD in men and women.

Current population-level tobacco control interventions may be less effective for those with mental illness. Health promotion campaigns and smoking policies that use stigma (e.g., the peril that smokers bring to the rest of the population) as the main motivating factor for giving up smoking may contribute to social isolation among those

with psychiatric disorders.³⁴ Thus, these efforts are more likely to perpetuate smoking inequalities than remove them. Cook and colleagues³⁵ found that individuals receiving mental health treatment are not only less likely to smoke but are more likely to quit, suggesting that the mental health facility is a promising setting to promote smoking cessation in this group.

The study had several limitations. We did not include individuals below 18 years of age and did not assess other forms of smoking such as use of e-cigarettes which is gaining popularity despite its ban in Singapore. Second, we relied on self-report which could lead to an underestimation of true prevalence of nicotine dependence and associations with mental and medical conditions. Third, as this was a cross-sectional study, we are unable to determine causality. Fourth, although we achieved a fair response rate of 69.5%, there were sociodemographic differences between respondent and non-respondent groups. Respondents were more likely to be in the younger age group (ie, 18–35 years compared with 35–49 years; OR=0.65, $p<0.0001$; 50–64 years; OR=0.68, $p<0.0001$; 65+years; OR=0.82, $p=0.005$) and of Malay or Indian ethnicity (compared with Chinese; OR=1.87, $p<0.0001$ and OR=1.91, $p<0.0001$, respectively). This could lead to obscuring true prevalence as mental health determinants differ between responders and non-responders.³⁶ In order to minimise the impact of this bias, non-response weighting was used to statistically adjust for these differences. The strengths of this study include the large sample size, the use of structured, well-validated instruments and a methodology similar to the 2010 study that allows for a direct comparison between these two time points.

Acknowledgements We would like to thank the interviewers who collected the data for the study.

Contributors SAC, MS and JAV were involved in the planning of the study and provided advice and guidance throughout the conduct study, and manuscript writing. ShS, SaS, SC, RS and YZ were involved in the conduct of the study. EA conducted the statistical analyses. DH and YYT provided expert opinion. All authors reviewed the final document.

Funding The study was funded by the Ministry of Health Singapore and Temasek Foundation Innovates.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Shazana Shahwan <http://orcid.org/0000-0003-4584-0751>

REFERENCES

- Méndez D, Alshanqeety O, Warner KE. The potential impact of smoking control policies on future global smoking trends. *Tob Control* 2013;22:46–51.
- WHO. Who | parties to the who framework convention on tobacco control, 2017. Available: https://www.who.int/fctc/signatories_parties/en/
- Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006;3:e442.
- Ng M, Freeman MK, Fleming TD, et al. Smoking prevalence and cigarette consumption in 187 countries, 1980–2012. *JAMA* 2014;311.
- Bilano V, Gilmour S, Moffiet T, et al. Global trends and projections for tobacco use, 1990–2025: an analysis of smoking indicators from the who comprehensive information systems for tobacco control. *The Lancet* 2015;385:966–76.
- Marcon A, Pesce G, Calciano L, et al. Trends in smoking initiation in Europe over 40 years: a retrospective cohort study. *PLoS One* 2018;13:e0201881.
- Hill S, Amos A, Clifford D, et al. Impact of tobacco control interventions on socioeconomic inequalities in smoking: review of the evidence. *Tob Control* 2014;23:e89–97.
- Stringhini S, et al. Association of socioeconomic position with health behaviors and mortality. *JAMA* 2010;303.
- Lasser K, Boyd JW, Woolhandler S, et al. Smoking and mental illness: a population-based prevalence study. *JAMA* 2000;284:2606–10.
- Campion J, Checinski K, Nurse J, et al. Smoking by people with mental illness and benefits of smoke-free mental health services. *Adv psychiatr. treat* 2008;14:217–28.
- Kessler RC, Chiu WT, Demler O, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National comorbidity survey replication. *Arch Gen Psychiatry* 2005;62:617.
- Amul GGH, Pangestu Pang T. Progress in tobacco control in Singapore: lessons and challenges in the implementation of the framework convention on tobacco control. *Asia Pac Policy Stud* 2018;5:102–21.
- Picco L, Subramaniam M, Abdin E, et al. Smoking and nicotine dependence in Singapore: findings from a cross-sectional epidemiological study. *Ann Acad Med Singapore* 2012;41:325–34.
- Health Promotion Board. Health promotion board unveils two-pronged strategy to step up tobacco control efforts on world no tobacco day, 2014–4. Available: <https://www.hpb.gov.sg/article/health-promotion-board-unveils-two-pronged-strategy-to-step-up-tobacco-control-efforts-on-world-no-tobacco-day>
- Subramaniam M, Abdin E, Vaingankar JA, et al. Tracking the mental health of a nation: prevalence and correlates of mental disorders in the second Singapore mental health study. *Epidemiol Psychiatr Sci* 2019;32:1–10.
- Subramaniam M, Vaingankar J, Heng D, et al. The Singapore mental health study: an overview of the methodology. *Int J Methods Psychiatr Res* 2012;21:149–57.
- Manimunda SP, Benegal V, Sugunan AP, et al. Tobacco use and nicotine dependency in a cross-sectional representative sample of 18,018 individuals in Andaman and Nicobar Islands, India. *BMC Public Health* 2012;12:515.
- Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010. *The Lancet* 2012;380:2224–60.
- Chia KS. TODAYonline | Singapore should aim to be smoke-free, not just smoke-lite. TODAYonline, 2017. Available: <https://www.todayonline.com/commentary/singapore-should-aim-be-smoke-free-not-just-smoke-lite> [Accessed 7 Mar 2019].
- Subramaniam M, Shahwan S, Fauziana R, et al. Perspectives on smoking initiation and maintenance: a qualitative exploration among Singapore youth. *Int J Environ Res Public Health* 2015;12:8956–70.
- Pierce JP, White VM, Emery SL. What public health strategies are needed to reduce smoking initiation? *Tob Control* 2012;21:258–64.
- Marques-Vidal P, Cerveira J, Paccaud F, et al. Smoking trends in Switzerland, 1992–2007: a time for optimism? *J Epidemiol Community Health* 2011;65:281–6.
- Lim J-L. How much does smoking cost in Singapore? 2018. Available: <https://www.imoney.sg/articles/smoking-cost-singapore/> [Accessed 8 Mar 2019].
- Shahwan S, Fauziana R, Satghare P, et al. Qualitative study of Singaporean youths' perception of antismoking campaigns: what works and what does not. *Tob Control* 2016;25:e101–6.
- Dinn WM, Aycicegi A, Harris CL. Cigarette smoking in a student sample: neurocognitive and clinical correlates. *Addict Behav* 2004;29:107–26.
- Doyon WM, Dong Y, Ostroumov A, et al. Nicotine decreases ethanol-induced dopamine signaling and increases self-administration via stress hormones. *Neuron* 2013;79:530–40.

- 27 Glassman AH, Helzer JE, Covey LS, *et al.* Smoking, smoking cessation, and major depression. *JAMA* 1990;264:1546–9.
- 28 Munafò MR, Hitsman B, Rende R, *et al.* Effects of progression to cigarette smoking on depressed mood in adolescents: evidence from the National longitudinal study of adolescent health. *Addiction* 2008;103:162–71.
- 29 Grabe HJ, Meyer C, Hapke U, *et al.* Lifetime-comorbidity of obsessive-compulsive disorder and subclinical obsessive-compulsive disorder in northern Germany. *Eur Arch Psychiatry Clin Neurosci* 2001;251:130–5.
- 30 Wu P, Goodwin RD, Fuller C, *et al.* The relationship between anxiety disorders and substance use among adolescents in the community: specificity and gender differences. *J Youth Adolesc* 2010;39:177–88.
- 31 Dell’Osso B, Nicolini H, Lanzagorta N, *et al.* Cigarette smoking in patients with obsessive compulsive disorder: a report from the International College of obsessive compulsive spectrum disorders (ICOCS). *CNS Spectr* 2015;20:469–73.
- 32 Bejerot S, von Knorring L, Ekselius L. Personality traits and smoking in patients with obsessive-compulsive disorder. *Eur Psychiatry* 2000;15:395–401.
- 33 Abramovitch A, Pizzagalli DA, Geller DA, *et al.* Cigarette smoking in obsessive-compulsive disorder and unaffected parents of OCD patients. *European Psychiatry* 2015;30:137–44.
- 34 Farrimond HR, Joffe H. Pollution, peril and poverty: a British study of the stigmatization of smokers. *J Community Appl Soc Psychol* 2006;16:481–91.
- 35 Cook BL, Wayne GF, Kafali EN, *et al.* Trends in smoking among adults with mental illness and association between mental health treatment and smoking cessation. *JAMA* 2014;311.
- 36 de Winter AF, Oldehinkel AJ, Veenstra R, *et al.* Evaluation of non-response bias in mental health determinants and outcomes in a large sample of pre-adolescents. *Eur J Epidemiol* 2005;20:173–81.