

Tobacco consumption and secondhand smoke exposure in vehicles: a cross-sectional study

Ariadna Curto,^{1,2} Jose M Martínez-Sánchez,^{1,3,4} Esteve Fernández^{1,3,4}

To cite: Curto A, Martínez-Sánchez JM, E Fernández. Tobacco consumption and secondhand smoke exposure in vehicles: a cross-sectional study. *BMJ Open* 2011;1:e000418. doi:10.1136/bmjopen-2011-000418

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://bmjopen.bmj.com>).

Received 27 September 2011
Accepted 31 October 2011

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

¹Tobacco Control Unit, Cancer Control and Prevention Programme, Institut Català d'Oncologia-ICO, L'Hospitalet de Llobregat, Barcelona, Spain

²Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Barcelona, Spain

³Cancer Control and Prevention Group, Institut d'Investigació Biomèdica de Bellvitge-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

⁴Department of Clinical Sciences, Universitat de Barcelona, Campus de Bellvitge, L'Hospitalet del Llobregat, Barcelona, Spain

Correspondence to

Dr Jose M Martínez-Sánchez; jmmartinez@iconcologia.net

ABSTRACT

Objectives: To estimate the prevalence of tobacco consumption and secondhand smoke (SHS) exposure in private cars, commercial vehicles and taxis in the city of Barcelona in Spain.

Design setting and participants: We carried out an observational cross-sectional study in 2011. We selected a systematic sample of 2442 private cars, commercial vehicles and taxis on 40 public roads regulated by traffic lights in all 10 districts of Barcelona. We calculated the prevalence rates and 95% CIs of smoking and SHS exposure in cars, and the corresponding ORs adjusting for the potential confounding variables.

Results: The prevalence of tobacco consumption was 5.5% (95% CI 4.6% to 6.4%) and was greater for commercial vehicles (9.8%; 95% CI 7.1% to 12.5%). The prevalence of SHS exposure was 5.2% (95% CI 3.8% to 6.6%) and 2.2% (95% CI 0.5% to 3.9%) of passengers under 14 years of age were exposed to SHS in vehicles.

Conclusions: This study highlights the need to promote public health measures aimed at reducing tobacco consumption in vehicles, especially in the presence of children, as well as enforcement of the current Spanish law against smoking in commercial vehicles and taxis.

INTRODUCTION

Secondhand smoke (SHS) contains a complex mixture of pollutants that includes toxic and irritant compounds, as well as carcinogens.^{1,2} SHS has been classified by the International Agency for Research on Cancer (IARC) as a type I carcinogen in humans.² In 2004 it was estimated that 1% of global mortality (603 000 deaths) was attributable to SHS exposure.³ Specifically, in Spain, it was estimated that between 1228 and 3237 deaths from lung cancer and myocardial infarction were attributable to SHS exposure in 2002.⁴

Although there is a dose–response effect and no level of SHS exposure can be

ARTICLE SUMMARY

Article focus

- Tobacco consumption and secondhand smoke (SHS) exposure in vehicles should be targeted because of the high concentrations of SHS due to the confined environment.
- Tobacco consumption while driving can also increase the risk of traffic accidents due to driver distraction.
- Few studies have used direct observation to examine the consumption of tobacco and SHS exposure in vehicles.

Key messages

- Reducing smoking in vehicles in the presence of children must be prioritised.
- Smoking laws should be enforced in commercial vehicles and taxis.
- Smoking in vehicles should be recognised by legislation as a distraction.

Strengths and limitations of this study

- The main limitation of this study is the inherent observer bias, especially as regards variables such as age, where physical appearance can result in misclassification by the observer.
- Obtaining a truly random and representative sample of vehicles in circulation in a city is difficult.
- This observational study avoids the information bias generated by the use of self-reported questionnaires.
- A pilot study found that direct observational studies are useful for monitoring smoking by motor vehicle drivers.

considered safe,⁵ the intensity of the exposure depends on the length of exposure and the concentration of SHS in the air. Therefore, because they are confined environments, tobacco consumption and SHS exposure in vehicles deserves special attention as SHS concentrations can be much higher than in the home, workplace or leisure settings such as discos and bars.^{6,7} Moreover, concentrations of PM_{2.5} (particulate matter with a diameter below 2.5 µm) in

vehicles where people are smoking^{6–10} can exceed the limits recommended by the US Environmental Protection Agency.¹¹

Few studies have examined the consumption of tobacco and SHS exposure in vehicles through direct observation^{12–13}; to our knowledge, only one cross-sectional study using questionnaires has been carried out in Spain.¹⁴ The objective of this study was to estimate the prevalence of tobacco consumption and SHS exposure in private cars, commercial vehicles and taxis in the city of Barcelona.

METHODS

Sampling, sample size and site selection

A cross-sectional study was conducted using direct observation of the occupants of private cars, commercial vehicles and taxis in the city of Barcelona. The fieldwork was carried out in April and May 2011. For each of the 10 districts of Barcelona, we randomly selected two public roads of the 15 with the greatest traffic flow rates and two public roads of the five with the greatest density of nurseries and primary and secondary schools to ensure the presence of children in vehicles. We consulted the 2011 'Araña de Tráfico' of the Mobility Services Division of Barcelona City Council when choosing the public roads.¹⁵ We selected a total of 40 public roads (four per district, two with greater traffic flow rates and two with greater school density). For each of the public roads, a traffic light was selected as the observation point where a trained observer directly observed the vehicles and recorded the variables of interest on a specially designed data collection sheet.

The theoretical sample size was 2401 vehicles assuming an expected prevalence of 50% with a 95% confidence interval (CI) and a precision of 2%. Two previous studies in Italy and Spain had prevalence rates of 10% and 20%, respectively.^{12–14} However, we assumed a 50% prevalence to maximise statistical power in order to stratify the results by type of vehicle. The final sample size was 2442 vehicles. The number of observations was distributed in proportion to the rates of traffic flow on each public road. The observations were carried out when traffic lights were set at red for vehicles. We systematically selected the first two vehicles in the lane adjacent to the observer. We excluded adjacent lanes which were restricted to buses, taxis and bicycles. The study included any private car, commercial vehicle (tourist related, adapted mixed vehicles and vans carrying a company logo or slogan) or taxi. We excluded buses, coaches, trucks, motorcycles, bicycles and other public service vehicles (ambulances, police cars, driving school cars, etc). We also excluded cars with tinted windows that did not allow the occupants to be observed. When a vehicle did not meet the inclusion criteria, we observed the next vehicle in line. The observations were carried out continuously for an average duration of 1 h from 8:00 to 11:00 h and from 17:00 to 19:00 h, Monday through Friday.

Study variables

We defined the variables for drivers, passengers and vehicles. The driver variables studied were: tobacco consumption (yes, no), approximate age (18–34, 35–64, and ≥ 65 years) and sex (male, female). Passenger variables were: total number of passengers, number of passengers who smoked and number of passengers under 14 years of age. Vehicle variables were: type of vehicle (private car, commercial vehicle, taxi), number of open windows, and whether the driver's window was open (yes, no). We also collected contextual variables (day of the week, district, time, weather and number of lanes on the road).

The two main variables, tobacco consumption by drivers and passengers, were defined as the presence of any burning tobacco product in the hand or mouth of the driver or a passenger. From these variables, two new variables were created: total tobacco consumption (driver and/or passenger) and SHS exposure, that is, if an occupant (driver or passenger) was exposed to SHS. The district variable was categorised into three groups according to the socioeconomic status of the district in question.¹⁶

We conducted a pilot study¹⁷ before the fieldwork to evaluate the feasibility of the observations and standardise the data collection sheet, as well as to analyse the degree of agreement between two observers in obtaining information. This pilot study demonstrated the feasibility of the direct observation method and perfect inter-observer concordance for monitoring the consumption of tobacco and number of passengers under 14 years of age (simple inter-observer agreement of 100% and κ coefficient=1.0). However, lower inter-observer agreement was found regarding the driver's age (simple inter-observer agreement of 94.3% and κ coefficient=0.865).¹⁷

Statistical analysis

We carried out a descriptive analysis of tobacco consumption and SHS exposure stratified by driver, passengers and vehicle variables, and by context. We calculated the prevalence of smoking and SHS exposure and their 95% CIs.¹⁸ We performed a χ^2 test to compare the prevalence rates. We fitted a logistic regression model to obtain the adjusted OR and 95% CI of smoking in cars by drivers. All analyses were performed using the statistical package SPSS V.15.

As personal data or biological samples were not used, approval was not required by the Ethics Committee of Bellvitge University Hospital.

RESULTS

We observed 2442 vehicles, of which 71.1% were private cars, 19.7% commercial vehicles and 9.2% taxis. More than half of the observations (53.9%) were conducted on public roads with more than two lanes. The majority (77.8%) of drivers were men and the most common age range was 36–64 years (69.6%). There were no passengers in 62.6% of the vehicles observed. There was

a passenger under 14 years of age in 29.7% of the vehicles with passengers (11.1% of all vehicles). All windows were closed in 53.4% of vehicles observed.

Table 1 shows the prevalence of smoking and total SHS exposure of the driver and/or passenger according to type of vehicle. The prevalence of tobacco consumption in vehicles (private, commercial and taxis) was 5.5% (95% CI 4.6% to 6.4%). The prevalence of tobacco consumption was higher in commercial vehicles (9.8%, 95% CI 7.1% to 12.5%). The prevalence of tobacco consumption among drivers was 4.7% (95% CI 3.9% to 5.5%) and 2.4% (95% CI 1.4% to 3.4%) among passengers; this difference was maintained according to vehicle type (table 1).

The overall prevalence of SHS exposure was 5.2% (95% CI 3.8% to 6.6%). Commercial vehicle occupants were the most exposed to SHS (12.7%, 95% CI 7.1% to 18.3%), while in taxis there was no observed tobacco consumption or occupant exposure to SHS. Passengers were slightly more likely to be exposed to SHS than drivers (3.0% vs 2.2%; $p=0.283$). This difference was threefold higher in commercial vehicles (9.7% vs 3.0%; $p=0.024$). The prevalence of passengers under 14 years old exposed to SHS in vehicles was 2.2% (95% CI 0.5% to 3.9%) (table 1).

Table 2 shows the prevalence of driver tobacco consumption according to vehicle type, driver and vehicle variables, and context. The prevalence of smoking in drivers was almost twofold higher in men than in women (5.2% vs 2.9%; $p=0.031$). There was a statistically significant downward trend in the consumption of tobacco according to the age of the driver, although this trend was not statistically significant in commercial vehicles or taxis. Although consumption of tobacco by drivers was higher in districts with more socioeconomic deprivation (5.2%, 95% CI 3.8% to 6.6%), this difference was not statistically significant. The overall prevalence of tobacco consumption was higher in the morning than in the afternoon (5.3% vs 3.8%; $p=0.097$), especially in commercial vehicles

(11.6% vs 4.1%; $p=0.006$). The prevalence of tobacco consumption among drivers was greater on cloudy than sunny days, regardless of vehicle type (table 2).

In bivariate logistic regression models, we confirmed a stronger association with driver tobacco consumption when the driver was a man (OR 1.79, 95% CI 1.05 to 3.07), was 18–34 years old (OR 9.61, 95% CI 1.31 to 70.53), was driving a commercial vehicle (OR 2.48, 95% CI 1.66 to 3.69) and had the window open (OR 10.50, 95% CI 5.86 to 18.82) (table 3). After adjusting a saturated model with all potential confounders, this association was statistically significant only for the driver's window being open (OR 11.05, 95% CI 6.08 to 20.09), mornings (OR 1.83, 95% CI 1.20 to 2.79) and cloudy weather (OR 1.69, 95% CI 1.11 to 2.57) (table 3).

DISCUSSION

This is the first study in Spain to estimate the prevalence of tobacco consumption and SHS exposure in vehicles through direct observation and shows that both are high in vehicles in Barcelona, especially commercial vehicles.

The prevalence of tobacco consumption in vehicles observed in our study is similar to that demonstrated in studies in Italy¹² and New Zealand,¹³ where vehicles were also directly observed. However, SHS exposure in vehicles in our study (5.2%) is much lower than that seen in the New Zealand study (23.7%). This could be related to social and other variations such as the different population sizes between the areas observed in Wellington in New Zealand (<410 000 inhabitants) and in Barcelona in Spain (about 1.6 millions inhabitants) and also the different years in which the studies were carried out.

The prevalence rates obtained in studies using questionnaires^{14–19} are considerably higher than those estimated in studies using direct observations as in the present study. One reason could be the inability of an observational study to monitor the entire car journey, during which the driver can smoke at any time or smoke more than one cigarette, although the driver's consumption of tobacco during the entire journey can

Table 1 Prevalence (%) and 95% confidence interval (CI) of tobacco consumption and secondhand smoke (SHS) exposure in vehicles in the city of Barcelona, Spain (2011)

	Total		Private		Commercial		Taxi	
	n (%)	(95% CI)	n (%)	(95% CI)	n (%)	(95% CI)	n (%)	(95% CI)
Tobacco consumption								
Driver and/or passenger	2442 (5.5)	(4.6 to 6.4)	1736 (4.7)	(3.7 to 5.7)	482 (9.8)	(7.1 to 12.5)	224 (2.2)	(0.3 to 4.1)
Driver	2442 (4.7)	(3.9 to 5.5)	1736 (3.8)	(2.9 to 4.7)	482 (8.9)	(6.4 to 11.4)	224 (2.2)	(0.3 to 4.1)
Passenger	913 (2.4)	(1.4 to 3.4)	685 (2.6)	(1.4 to 3.8)	134 (3.0)	(0.1 to 5.9)	94 (0.0)	(0.0 to 3.9)
SHS exposure								
Any occupant	913 (5.2)	(3.8 to 6.6)	685 (4.3)	(2.8 to 5.8)	134 (12.7)	(7.1 to 18.3)	94 (0.0)	(0.0 to 3.9)
Driver	913 (2.2)	(1.2 to 3.2)	685 (2.3)	(1.2 to 3.4)	134 (3.0)	(0.1 to 5.9)	94 (0.0)	(0.0 to 3.9)
Passenger	913 (3.0)	(1.9 to 4.1)	685 (2.0)	(1.0 to 3.0)	134 (9.7)	(4.7 to 14.7)	94 (0.0)	(0.0 to 3.9)
Passenger <14 years old	271 (2.2)	(0.5 to 3.9)	253 (2.0)	(0.3 to 3.7)	12 (8.3)	(1.5 to 35.4)	6 (0.0)	(0.0 to 39.0)

Table 2 Prevalence (%) and 95% confidence interval (CI) of driver's tobacco consumption by vehicle type, driver and vehicle variables and context in the city of Barcelona, Spain

	Total (n = 2442)		Private (n = 1736)		Commercial (n = 482)		Taxi (n = 224)	
	% (95% CI)	p Value	% (95% CI)	p Value	% (95% CI)	p Value	% (95% CI)	p Value
Sex								
Male	5.2 (4.2 to 6.2)	0.031	4.2 (3.1 to 5.3)	0.223	9.2 (6.5 to 11.9)	0.351	2.3 (0.3 to 4.3)	0.708
Female	2.9 (1.5 to 4.3)		2.9 (1.4 to 4.4)		3.8 (0.7 to 18.9)		0 (0.0 to 39.0)	
Age								
18–34 years	6.6 (4.6 to 8.6)	0.002	6.1 (3.9 to 8.3)	0.001	8.4 (3.6 to 13.2)	0.945	5.3 (0.9 to 24.6)	0.345
35–64 years	4.3 (3.3 to 5.3)		3.2 (2.2 to 4.2)		9.4 (6.3 to 12.5)		2.0 (0.1 to 3.9)	
≥65 years	0.7 (0.1 to 4.0)		0.8 (0.1 to 4.4)		0 (0.0 to 27.8)		0 (0.0 to 65.8)	
Socioeconomic status*		0.628		0.277		0.585		0.268
Low	5.2 (3.8 to 6.6)		4.7 (3.1 to 6.3)		7.4 (3.7 to 11.1)		3.9 (0.1 to 7.7)	
Medium	4.3 (3.0 to 5.6)		3.1 (1.8 to 4.4)		9.3 (5.3 to 13.3)		1.2 (0.2 to 6.6)	
High	4.5 (2.7 to 6.3)		3.4 (1.5 to 5.3)		11.1 (4.6 to 17.6)		0 (0.0 to 8.8)	
Driver's open window		<0.001		<0.001		0.001		0.089
Yes	9.3 (7.6 to 11.0)		9.1 (6.9 to 11.3)		12.3 (8.6 to 16.0)		3.5 (0.5 to 6.5)	
No	1.0 (0.5 to 1.5)		0.6 (0.1 to 1.1)		3.3 (0.7 to 5.9)		0 (0.0 to 4.5)	
Time		0.097		0.947		0.006		0.150
Morning (8:00–11:00 h)	5.3 (4.1 to 6.5)		3.8 (2.6 to 5.0)		11.6 (8.0 to 15.2)		0.9 (0.2 to 4.7)	
Afternoon (17:00–19:00 h)	3.8 (2.6 to 5.0)		3.8 (2.4 to 5.2)		4.1 (1.1 to 7.1)		3.7 (0.1 to 7.3)	
Weather		0.230		0.506		0.855		0.004
Sunny	4.3 (3.3 to 5.3)		3.6 (2.6 to 4.6)		9.1 (5.9 to 12.3)		0.6 (0.1 to 3.3)	
Cloudy	5.4 (3.8 to 7.0)		4.3 (2.5 to 6.1)		8.6 (4.3 to 12.9)		7.1 (0.4 to 13.8)	

*According to socioeconomic status of district.

Table 3 Crude and adjusted odds ratio (OR) and 95% confidence interval (CI) of smoking in cars by drivers

	cOR (95% CI)	p Value	aOR (95% CI)	p Value
Sex				
Male	1.79 (1.05 to 3.07)	0.033	1.46 (0.83 to 2.59)	0.193
Female	1		1	
Age				
18–34 years old	9.61 (1.31 to 70.53)	0.026	6.85 (0.92 to 51.18)	0.061
35–64 years old	6.11 (0.84 to 44.30)	0.073	5.12 (0.69 to 37.81)	0.109
≥65 years old	1		1	
Vehicle type				
Private	1		1	
Commercial	2.48 (1.66 to 3.69)	<0.001	1.36 (0.89 to 2.10)	0.158
Taxi	0.58 (0.23 to 1.45)	0.242	0.34 (0.13 to 0.86)	0.023
Socioeconomic status*				
Low	1.15 (0.69 to 1.91)	0.605	1.18 (0.69 to 2.00)	0.548
Medium	0.93 (0.55 to 1.58)	0.801	0.98 (0.57 to 1.68)	0.930
High	1		1	
Driver's open window				
Yes	10.50 (5.86 to 18.82)	<0.001	11.05 (6.08 to 20.09)	<0.001
No	1		1	
Time				
Morning (8:00–11:00 h)	1.40 (0.94 to 2.08)	0.099	1.83 (1.20 to 2.79)	0.005
Afternoon (17:00–19:00 h)	1		1	
Weather				
Sunny	1		1	
Cloudy	1.27 (0.86 to 1.89)	0.231	1.69 (1.11 to 2.57)	0.014

*According to socioeconomic status of district.

cOR, crude OR; aOR, adjusted OR derived from a logistic regression model adjusted for driver's sex, age and window, vehicle type, district, time and weather.

be obtained in classical cross-sectional studies using questionnaires. This limitation could lead to an underestimation of prevalence as occurs in studies in bars, which have reported an almost threefold higher number of smokers identified using questionnaires compared with the number found through direct observations.²⁰ This aspect should be addressed in future studies.

On the other hand, tobacco consumption in our study was higher when any window in the vehicle was open, as in previous studies.¹³ This may be due to the belief by smokers that the adverse health effects of tobacco consumption in vehicles are minimised when a window is open.²¹ However, although levels of PM_{2.5} are lower when the vehicle is ventilated, the levels are still unhealthy.^{6 9 10 22}

The prevalence of SHS exposure in children was high compared to that observed in the Italian study (0.9%).¹² This difference could be attributed to the fact that in our study there were schools on half of the public roads selected. This fact should be taken into account since children are inevitably more vulnerable to the effects of SHS exposure.²³ Indeed, a study conducted in Ireland²⁴ found an increased likelihood of developing respiratory and allergic symptoms in children aged 13–14 years old exposed to SHS in vehicles, with wheezing as the most obvious symptom. Therefore, it is clear public health policies in Spain need to focus on reducing tobacco consumption in private cars if children are present. Moreover, a study conducted through telephone surveys

in four countries (Australia, Canada, UK and USA) concluded that the majority of smokers (more than 60%) would ban smoking in vehicles if children are present,²⁵ with higher approval by smokers with less tobacco dependence and with young children (<5 years old) versus those without children. This support for restricting smoking in vehicles is also found in studies among adolescents.^{26 27}

The observed prevalence of tobacco consumption in commercial vehicles and taxis is especially high considering that under Law 42/2010²⁸ smoking in commercial vehicles and taxis in Spain is banned as they are places of work and public places. Furthermore, since our study found a very high prevalence of tobacco consumption and SHS exposure in commercial vehicles, as seen in Italy,¹² the current legislation should be more strictly enforced.

On the other hand, tobacco consumption in taxis was never observed in the presence of passengers. Although the passengers of taxis were not directly exposed to SHS, recent studies^{29 30} have highlighted the possible effects of exposure to SHS particles deposited on vehicle seats, as dust or in the air. The involuntary inhalation, ingestion or skin absorption of particles is known as thirdhand smoke exposure.²⁹ While there is still insufficient evidence to assess the health hazards from thirdhand smoke, further studies should investigate the potential for exposure to thirdhand smoke and its health effects.

Tobacco consumption while driving is also a distraction that increases the risk of traffic accidents.^{31–33} It is estimated that drivers who smoke are 1.5 times more likely to have an accident than non-smoking drivers.³⁴ Distraction caused by smoking is associated with, among others, decreased dexterity controlling the steering wheel and reduced attention to the road (when the cigarette is being lit or put out and as a result of eye irritation caused by the carbon monoxide present in SHS), in addition to the lower perception of risk that smokers have at the wheel versus non-smokers.³⁴ Indeed, a survey conducted by the Spanish National Road Safety Observatory of the General Directorate of Traffic³⁵ in 2005 revealed that 83.8% of interviewed drivers thought that tobacco consumption was a source of distraction while driving. In addition, 76.8% thought smoking was a fairly or very dangerous behaviour while driving. Given the evidence of the health hazards of smoking while driving and the broad general perception of risk by the population, smoking should be treated the same way as other distractions recognised by Spanish law, such as the use of mobile phones or other manually operated devices, for instance GPS systems.³⁶

The main limitation of this study derives from the inherent observer bias, especially regarding variables such as age, as people can look younger or older than they actually are. However, the pilot study¹⁷ carried out by two independent observers showed almost perfect agreement in observations. We cannot disregard selection bias, since the choice of public roads with a higher density of schools could have resulted in the number of children present in vehicles being over-represented. Other potential limitations of the study are that field-work was conducted only in spring and at a particular time of day. However, the Spanish National Health Interview Surveys conducted in several waves during a single year have not shown seasonality in tobacco consumption (prevalence rates by waves for several years are available at <http://www.ine.es>). Moreover, similar prevalence rates of smoking in vehicles have been reported in autumn,¹² which also suggests no seasonal differences. Finally, we do not know whether the time-window used for sampling in our study is representative of the entire 24 h in a day. While there are fewer vehicles during the night, we do not know if those driving during our sampling time smoke less or more than drivers during the rest of the day or night.

We have used logistic regression as a measure of association in a cross-sectional study. However, the OR only overestimates the prevalence ratio when it is above 20%,³⁷ which is not the case for our data. Also noteworthy is the difficulty of obtaining a truly random and representative sample of vehicles in circulation in a city. Our sampling approach, designed to be representative of all 10 districts of Barcelona, was designed to minimise this limitation. A strength of an observational study like this compared with studies based on the use of questionnaires is the lack of information bias inherent in self-reported data.

In conclusion, this study provides an estimation of smoking in some types of vehicles and highlights the need for public health measures aimed at reducing tobacco consumption in private cars, especially in the presence of children, as well as the enforcement of measures to control smoking in commercial vehicles and taxis.

Acknowledgements We acknowledge the collaboration of Ángel López Rodríguez for providing 'la Araña de Tráfico' of the city of Barcelona.

Funding This work was funded by the Instituto de Salud Carlos III, Government of Spain (RTICC RD06/0020/0089) and the Ministry of Universities and Research, Government of Catalonia (grant 2009SGR192).

Competing interests None.

Contributors All authors contributed to the design of the study. AC collected the data, prepared the database and analyzed the data. All authors revised the results and contributed to the interpretation of results. AC drafted the manuscript, which was critically revised by JMMS and EF. All authors approved the final version.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data available.

REFERENCES

1. US Environmental Protection Agency. *Respiratory Health Effects of Passive Smoking: Lung Cancer and Other Disorders*. Washington, DC: Environmental Protection Agency, 1992:6–90.
2. International Agency for Research on Cancer. *Monographs on the Evaluation of Carcinogenic Risks to Humans (Vol 83). Tobacco Smoke and Involuntary Smoking*. Lyon: International Agency for Research on Cancer, 2002.
3. Öberg M, Jaakkola MS, Woodward A, *et al*. Worldwide burden of disease from exposure to secondhand smoke: a retrospective analysis of data from 192 countries. *Lancet* 2011;377:139–46.
4. López MJ, Pérez-Ríos M, Schiaffino A, *et al*. Mortality attributable to passive smoking in Spain, 2002. *Tob Control* 2007;16:373–7.
5. Dept. of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion Office on Smoking and Health. *The Health Consequences of Smoking: a report of the Surgeon General*. Washington, DC: Surgeon General, 2004.
6. Jones MR, Navas-Acién A, Yuan J, *et al*. Secondhand tobacco smoke concentrations in motor vehicles: a pilot study. *Tob Control* 2009;18:399–404.
7. Vardavas CI, Linardakis M, Kafatos AG. Environmental tobacco smoke exposure in motor vehicles: a preliminary study. *Tob Control* 2006;15:415.
8. Rees VW, Connolly GN. Measuring air quality to protect children from secondhand smoke in cars. *Am J Prev Med* 2006;31:363–8.
9. Sendzik T, Fong GT, Travers MJ, *et al*. An experimental investigation of tobacco smoke pollution in cars. *Nicotine Tob Res* 2009;11:627–34.
10. Ott W, Klepeis N, Switzer P. Air change rates of motor vehicles and in-vehicle pollutant concentrations from secondhand smoke. *J Expo Sci Environ Epidemiol* 2008;18:312–25.
11. US Environmental Protection Agency. *National Ambient Air Quality Standards For Particulate Matter; Final Rule (Report Num. 71)*. Washington, DC: Environmental Protection Agency, 2006.
12. Sbrogiò L, Frison G, Tagliapietra L, *et al*. [Observed smoking in car: results of a study of the Regional Health Prevention Service of Veneto, Northern Italy] (In Italian). *Epidemiol Prev* 2010;34:43–7.
13. Martin J, George R, Andrews K, *et al*. Observed smoking in cars: a method and differences by socioeconomic area. *Tob Control* 2006;15:409–11.
14. Lushchenkova O, Fernández E, López MJ, *et al*. [Secondhand smoke exposure in Spanish adult non-smokers following the introduction of an anti-smoking law] (In Spanish). *Rev Esp Cardiol* 2008;61:687–94.
15. *Direcció de Serveis de Mobilitat, Ajuntament de Barcelona. Dades Bàsiques de Mobilitat*. 2011. http://w3.bcn.es/XMLServeis/XMLHomeLinkPI/0,4022,173198596_257817017_1,00.html (accessed 19 Jun 2011).
16. Domínguez-Berjón MF, Borrell C. [Mortality and socioeconomic deprivation in the census tracts and districts of Barcelona] (In Spanish). *Gac Sanit* 2005;19:363–9.

17. Martínez-Sánchez JM, Curto A, Fernández E. [Agreement between two observers in the measurement of tobacco consumption, use of safety belt and cell phones in vehicles] (In Spanish). *Gac Sanit*. Published Online first: 27 October 2011. doi:10.1016/j.gaceta.2011.07.011.
18. Wilson EB. Probable inference, the law of succession, and statistical inference. *J Am Stat Assoc* 1927;22:209–12.
19. Hitchman SC, Fong GT, Borland R, et al. Predictors of smoking in cars with nonsmokers: findings from the 2007 wave of the International Tobacco Control Four Country Survey. *Nicotine Tob Res* 2010;12:374–80.
20. Repace JL. Exposure analysis. In: Ott WR, Steinemann AC, Wallace LA, eds. *Exposure to Secondhand Smoke*. NY: Taylor & Francis Group, 2007:201–35.
21. Kegler MC, Escoffery C, Butler S. A qualitative study on establishing and enforcing smoking rules in family cars. *Nicotine Tob Res* 2008;10:493–7.
22. Invernizzi G, Ruprecht AA, Mazza R, et al. [Smoking in car: monitoring pollution of particulate matter, of organic volatile compounds and of carbon monoxide. The effect of opening the driver's window] (In Italian). *Epidemiol Prev* 2010;34:35–42.
23. Bearer CF. How are children different from adults? *Environ Health Perspect* 1995;103(Suppl 6):7–12.
24. Kabir Z, Manning PJ, Holohan J, et al. Secondhand smoke exposure in cars and respiratory health effects in children. *Eur Respir J* 2009;34:629–33.
25. Hitchman SC, Fong GT, Zanna MP, et al. Support and correlates of support for banning smoking in cars with children: findings from the ITC Four Country Survey. *Eur J Public Health* 2011;21:360–5.
26. Leatherdale ST, Smith P, Ahmed R. Youth exposure to smoking in the home and in cars: how often does it happen and what do youth think about it? *Tob Control* 2008;17:86–92.
27. Leatherdale ST, Ahmed R. Second-hand smoke exposure in homes and in cars among Canadian youth: current prevalence, beliefs about exposure, and changes between 2004 and 2006. *Cancer Causes Control* 2009;20:855–65.
28. Boletín Oficial del Estado. Ley 42/2010 por la que se modifica la Ley 28/2005 de medidas sanitarias frente al tabaquismo y reguladora de la venta, el suministro, el consumo y la publicidad de los productos del tabaco. <http://www.boe.es/boe/dias/2010/12/31/pdfs/BOE-A-2010-20138.pdf> (accessed 21 Jun 2011).
29. Matt GE, Quintana PJ, Destailats H, et al. Thirdhand tobacco smoke: emerging evidence and arguments for a multidisciplinary research agenda. *Environ Health Perspect* 2011;119:1218–26.
30. Matt GE, Quintana PJ, Hovell MF, et al. Residual tobacco smoke pollution in used cars for sale: air, dust, and surfaces. *Nicotine Tob Res* 2008;10:1467–75.
31. Sacks JJ, Nelson DE. Smoking and injuries: an overview. *Prev Med* 1994;23:515–20.
32. Wen CP, Tsai SP, Cheng TY, et al. Excess injury mortality among smokers: a neglected tobacco hazard. *Tob Control* 2005;14(Suppl 1):i28–32.
33. Buñuel JM, Córdoba R, del Castillo M, et al. Smoking and nonfatal traffic accidents. *Atención Primaria* 2003;31:349–53.
34. Brison RJ. Risk of automobile accidents in cigarette smokers. *Can J Public Health* 1990;81:102–6.
35. Soria J. Si conduce, mejor no fume. *Tráfico y Seguridad Vial* 2005;172:38–9.
36. Boletín Oficial del Estado. Ley 17/2005, de 19 de julio, por la que se regula el permiso y la licencia de conducción por puntos y se modifica el texto articulado de la ley sobre tráfico, circulación de vehículos a motor y seguridad vial. <http://www.boe.es/boe/dias/2005/07/20/pdfs/A25781-25793.pdf> (accessed 21 Jun 2011).
37. Schiaffino A, Rodríguez M, Pasarín MI, et al. [Odds ratio or prevalence ratio? Their use in cross-sectional studies]. (In Spanish). *Gac Sanit* 2003;17:70–4.

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found
Introduction		
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported
Objectives	3	State specific objectives, including any prespecified hypotheses
Methods		
Study design	4	Present key elements of study design early in the paper
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group
Bias	9	Describe any efforts to address potential sources of bias
Study size	10	Explain how the study size was arrived at
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses
Results		
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest
Outcome data	15*	Report numbers of outcome events or summary measures
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses

Discussion		
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.