

BMJ Open Attitudes and willingness toward out-of-hospital cardiopulmonary resuscitation: a questionnaire study among the public trained online in China

Yi Jiang ^{1,2}, Bangsheng Wu,^{1,2} Long Long,^{1,2} Jiaying Li,^{1,2} Xiaoqing Jin¹

To cite: Jiang Y, Wu B, Long L, *et al.* Attitudes and willingness toward out-of-hospital cardiopulmonary resuscitation: a questionnaire study among the public trained online in China. *BMJ Open* 2020;**10**:e038712. doi:10.1136/bmjopen-2020-038712

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-038712>).

Received 23 March 2020
Revised 25 August 2020
Accepted 01 September 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Emergency Center, Zhongnan Hospital of Wuhan University, 169 Donghu Road, Wuhan 430071, China

²The Second Clinical School, Wuhan University, Wuhan 430071, China

Correspondence to

Dr Xiaoqing Jin;
redjin@whu.edu.cn

ABSTRACT

Objectives The incidence of bystander cardiopulmonary resuscitation (CPR) is low in China. CPR training could improve public attitudes and willingness, but at present, the attitudes of the public after online training are unclear. This study investigated individual attitudes towards CPR, the willingness to perform it in emergencies along with the main obstacles and the overall effects of online training.

Design Questionnaires were distributed to investigate the public attitudes and willingness towards performing bystander CPR.

Setting Questionnaires were accessible after the online course 'First Aid'.

Participants 1888 students who attended 'First Aid' from December 2019 to 1 January 2020 and then completed the questionnaire voluntarily.

Results The majority understood CPR (96.7%) and displayed a willingness to learn (98.4%) and to disseminate CPR knowledge (82.0%). Characteristics associated with more positive attitudes included women, the 26–35-year olds and those in medical-related occupations ($p < 0.05$). Only 34.8% had CPR training before. Most people would willingly perform CPR on a close family member. Compared with the standard CPR (S-CPR), the public preferred chest compression-only CPR (CO-CPR) ($p < 0.01$). The top three obstacles to performing CO-CPR were lack of confidence (26.7%), fear of harming the victim (23.4%) and causing legal trouble (20.7%), while regarding S-CPR, fear of disease transmission (22.9%) ranked second. Women, those in poor health and in medical-related occupations, were more likely to perform CPR ($p < 0.05$). The confidence to perform CPR was improved remarkably after online training ($p < 0.05$).

Conclusions The overwhelming majority of respondents showed positive attitudes and willingness towards CPR. In some cases, there is still reluctance, especially towards S-CPR. Obstacles arise mainly due to lack of confidence in administering CPR, while online CPR training can markedly improve it. Therefore, we should focus on disseminating CPR knowledge, targeting those who are less willing to perform CPR and helping overcome their obstacles by online training.

Strengths and limitations of this study

- We conducted a questionnaire study to investigate the public attitudes and willingness towards cardiopulmonary resuscitation (CPR) after online training.
- Our questionnaire advanced the understanding of the influencing factors and obstacles of performing CPR.
- We provided a three-level ordinal variable in evaluating willingness and used ordinal regression to analyse the correlation between demographic characteristics and willingness to perform CPR.
- Our online training and questionnaire focused on theoretical knowledge, rather than on the practical skills of CPR.
- This survey was released through the internet only to the public that attended our online course, who usually paid more attention to CPR, which may lead to more optimistic results.

BACKGROUND

Out-of-hospital cardiac arrest (OHCA) remains a major cause of mortality worldwide. The mean incidence worldwide of OHCA is estimated to be 55/100 000 person-years.¹ In China, according to the national cardiovascular disease centre statistics, about 550 000 cardiac arrests happen every year, while the survival rates in OHCA are less than 1%. The sudden death of a public figure in 2019 pushed this topic to the forefront again, with many more online hits for 'cardiopulmonary resuscitation' (CPR).² Early initiation of CPR improves survival rates from OHCA,³ but the average rate for bystanders performing CPR in eight large and medium-sized cities in China is just 4.5%,⁴ while in the USA and in Europe, it is around 40.2% and 47.4%, respectively.^{5,6}

CPR training is a pivotal factor in improving the performance rate of CPR and has been

developed around the world according to the guidelines of the American Heart Association (AHA).⁷ Unfortunately, the training rate in China is less than 26%, while there is legislation mandating CPR training in some schools in the USA and in several European countries.^{4,8,9} In addition, due to the limitations of training location, instructor and equipment, live CPR training is not as accessible as it is online.¹⁰ Thus, in order to disseminate first-aid knowledge including CPR to a wider public, we developed 'First Aid', an online course. It has been stated that trained people will be more likely to perform CPR without much hesitation,¹¹ yet we have little data about the effects of online training in China. Consequently, the aim of this study was to explore attitudes and willingness towards CPR among the public in China who had taken online CPR training as well as to evaluate the effects of this training.

MATERIALS AND METHODS

Study design

The data for this survey were acquired via questionnaires distributed to the public through the internet from December 2019 to 1 January 2020, from those who had taken our online training (<http://www.icourses.cn/cuoc/>). The survey was carried out after the public had finished the online CPR training, and the link to the questionnaire was attached at the end of the course. Before filling the questionnaire, all participants were informed about the aim of the study and were assured that participation was voluntary and results would be anonymous. And the refusal to participate would be without any consequences. Only persons attached to the research team had access to the data.

Questionnaires

The research team developed the survey questions and precoded answer options after reviewing the literature as well as after consulting two Emergency Medicine Specialists in our hospital. The questionnaire consisted of four parts with 54 questions (see online supplemental file 1). The first part asked for personal demographic characteristics (age, gender, education level, occupation, health condition, family members with diseases, whether witnessed OHCA). The second section contained questions regarding the attitudes and willingness towards CPR, asking whether they knew the meaning of CPR, their willingness to learn and disseminate CPR knowledge as well as their motives, opinions on developing CPR training courses in university, the importance of chest compression (CC) and mouth-to-mouth ventilation (MMV) in cardiac arrest, their history of training in CPR, for those with none, their reasons for not receiving CPR training. The next section assessed the participants' willingness to perform CPR in eight hypothetical scenarios involving cardiac arrest that we programmed according to a previous study (see online supplemental table S1).¹² Additionally, respondents were asked to select the reason

if they were reluctant to perform either compression-only CPR (CO-CPR) or standard CPR (S-CPR). Finally, to quantify the confidence level before and after our online CPR training, we used scores from '1' (completely not confident) to '10' (very confident), in order to evaluate the overall effects of online CPR training.

Data analysis

Data were analysed by IBM SPSS V.20.0 (IBM Corporation, New York, USA). Some questions permitted the respondents to give more than one answer, resulting sometimes in percentages that add to more than 100%. Qualitative variables were reported as absolute frequencies and proportions, while quantitative variables were presented as the median. The difference between groups was analysed with χ^2 tests or Fisher exact tests. It produced a three-level ordinal variable, with a zero indicating not willing to perform CPR in any scenario, and two indicating being likely to perform in all scenarios. Ordinal regression was used to analyse the correlation between demographic characteristics and willingness to perform CPR. Variables with p values <0.1 at the univariate level were incorporated into the model. The level of significant difference was set at p<0.05.

Patient and public involvement

Public were not involved in the development of the methodology for the current study. However, academic discussion with two Emergency Medicine Specialists in our hospital have jointly promoted the design of the questionnaire. There are currently no plans to involve public in dissemination. Study objectives, benefits and mandatory information were explained to the public and the participants were voluntary to fill the questionnaire after taking online training.

RESULTS

Respondent demographic characteristics

We received a total of 1899 questionnaires, of which 1888 were valid, and the effective rate was 99.4%. Respondents were composed of 1091 women (57.8%) and 797 men (42.2%). Most respondents (65.4%) were between 18 and 25 years old, with bachelor's degree (67.4%) and in non-medical-related occupations (78.5%). 5.3% estimated their own health as poor, while 9.1% (n=171) confirmed their family members had life-threatening diseases. Only 139 of the respondents (7.4%) claimed they had witnessed someone collapse (table 1).

Attitudes towards CPR

Most of the respondents had positive attitudes towards CPR. 96.7% understood CPR. 98.4% and 82.0% of the respondents were willing to learn and to disseminate CPR knowledge, respectively. A great majority wanted to study CPR to avoid unnecessary death (87.6%) as well as to help others (86.4%). More participants (98.8%) attached great significance to timely CC in cardiac arrest, in

Table 1 Demographic characteristics of the respondents (n=1888)

Characteristics	Sample size, N (%)
Gender	
Male	797 (42.2)
Female	1091 (57.8)
Age, years	
<18	159 (8.4)
18–25	1234 (65.4)
26–35	314 (16.6)
>35	181 (9.6)
Education level	
<Associate's degree	395 (20.9)
Bachelor's degree	1273 (67.4)
Master's degree or above	220 (11.7)
Occupation	
Medical related	407 (21.5)
Non-medical related	1481 (778.5)
Health condition	
Good	1041 (55.1)
General	748 (39.6)
Poor	99 (5.3)
Family members have life-threatening diseases	
Yes	171 (9.1)
No	1717 (90.9)
Have you ever encountered someone in need of CPR performance?	
Yes	139 (7.4)
No	1749 (92.6)

CPR, cardiopulmonary resuscitation.

contrast with 81.9% to MMV. Moreover, 1213 participants (64.2%) had never received CPR training before, among which 50.9% did not know where to take the training, 37.6% claimed there would be a cost and 16.2% indicated that they had no time owing to a busy schedule (see online supplemental table S2). Women, younger people between 26 and 35 years old, and those in medical-related occupations were more positive towards CPR ($p<0.05$). The training rate differed by age and occupations. Those whose family members had life-threatening diseases had a higher rate of previous training ($p<0.05$) (table 2).

Willingness to perform CPR in different scenarios

Across all eight hypothetical scenarios of OHCA, people were more likely to perform CO-CPR instead of S-CPR ($p<0.01$). Most people reported a higher likelihood of performing CPR on a family member, a relative or a friend. It showed that when the victim was a close family member, the rate of willingness to perform CO-CPR was 98.8%, and S-CPR was 97.0%, whereas the public was least likely to perform CPR on an elderly person, with

the rates 84.0% for CO-CPR and 69.5% for S-CPR. (see figure 1 and online supplemental table S3). Analysing the effects of demographic characteristics on willingness by ordinal regression, we found statistical differences in gender, in occupation and in health condition groups. Women were more willing to perform CO-CPR than men. Those in medical-related occupations, in poor health and those who had had the experience of witnessing someone collapse had stronger motivation ($p<0.05$) (see figure 2 and online supplemental table S4).

Reasons for not performing CO-CPR or S-CPR

Lack of confidence (26.6%), fear of harming the victim (23.4%) and fear of causing legal trouble for themselves (20.7%) were the top three reasons for not performing CO-CPR. There existed a significant difference between men and women concerning motive in all eight scenarios ($p<0.05$). As shown in table 3, when analysing the major barriers to performing CO-CPR, male respondents mostly reported fear of causing legal trouble (23.8%), whereas female respondents mainly reported lack of confidence (29.0%). Regarding S-CPR, the major obstacle was lack of confidence (28.7%), which varied by gender ($p<0.05$). The next two reasons were fear of disease transmission (22.9%) and fear of harming the victim (21.5%) (see table 3 and online supplemental table S5).

The effects of online CPR training

We investigated the confidence to perform CPR in emergencies both pre and postonline training, assigning a '1' to the least confidence and '10' to the greatest confidence. The mean score was 1.89 points before, in comparison with 4.47 points after it, a difference of about 2.4 times. Of note was that participants scoring 1 point dropped as a proportion of the whole from 56.0% to 1.3%, accompanied by respondents scoring 6 points increasing from 2.8% to 22.1%. Although no one got more than 6 points, it showed their confidence improved markedly after online CPR training ($p<0.05$) (figure 3).

DISCUSSION

Positive attitudes towards CPR among the online-trained public

Early effective CPR plays a vital role in saving patients with OHCA, promoting their survival rate. However, the worldwide rate of bystander CPR is below 50% on average and is below 6% in China.^{4,13} In this study, we determined that the majority of the Chinese public understood the meaning of CPR after online training. Their desire to learn CPR (98.4%) stemmed from positive attitudes towards avoiding unnecessary death and helping others. Compared with the past, public attitudes have improved significantly after online training.¹⁴ Our result showed that 7.4% respondents had witnessed someone collapse, higher than in the study of Huang *et al* (4.3%).¹⁵ This may be because the tragedy of OHCA now draws more attention from people in this country. Women seem to



Table 2 Attitudes towards CPR and previous training experience in relation to demographic characteristics

	Gender		Age				Education level				Family members with life-threatening diseases				Occupation		Ever trained in CPR		
	Total	Male	Female	<18	18-25	26-35	>35	Associate's degree		Bachelor's degree		Master's degree or above		Yes	No	Medical-related	Non-medical-related	Yes	No
								<35	35-40	41-45	46-50	51+	Yes						
All respondents (1888)	1888	797	1055	159	1234	314	241	395	1273	220	171	1717	407	1481	657	1231			
Ever trained in CPR (%)	34.8	36.1	33.8	27	36.8	33.4	30.4*	31.1	36.6	30.9	37.3	31.4†	62.2	27.3†	N/A	N/A			
Understanding what is CPR (%)	96.7	96.7	96.7	95	97.3	96.8	93.9	93.9	97.4	97.7†	97.4	95.7†	99	96.1†	99.1	95.4†			
Would like to learn CPR knowledge (%)	98.4	97.2	99.2†	94.3	98.6	100	97.2†	96.5	99.1	97.3†	99.1	97.4	98.5	98.3†	98.3	98.4			
Would like to disseminate CPR knowledge (%)	82	79.2	84.0†	69.8	82.2	85.7	85.1†	73.7	83.4	89.1	83.4	80.0†	87.2	80.6†	86.9	79.4†			

*p<0.05.
†p<0.01, the χ^2 test were between each of the groups respectively.
CPR, cardiopulmonary resuscitation.




Figure 1 Proportions (%) of public being willing to perform chest compression-only cardiopulmonary resuscitation (CO-CPR) and standard CPR (S-CPR) in different scenarios. ‘CO-CPR’ means CO-CPR will be applied to the victim and so on. The difference was statistically significant between each group ($p < 0.01$).

attach more importance to CPR,¹⁶ which may be they had inadequate knowledge and skills of CPR. Because young people are good at obtaining wider information including the serious consequences of OHCA, they can recognise the importance of CPR earlier, which may be why 26–35-year olds were more eager to learn CPR. In this regard, many training courses are aimed at the young community, and while the elderly may have been trained before, their confidence and skills have eroded with time since their last training. This highlights an ongoing need for CPR refresher training.^{17 18} People in non-medical-related occupations attach less importance to CPR, which could be attributable to the fact that CPR training is not recommended as a compulsory curriculum, except in medical schools in China. Moreover, previous training experience does help improve their attitudes. Taken together, CPR training can be enhanced and directed towards specific groups, including the elderly and the non-medical-related public, in the future.




Figure 2 Association between the public demographic characteristics and willingness to perform chest compression-only cardiopulmonary resuscitation (CO-CPR) and standard CPR (S-CPR). OR and 95% CIs were estimated by ordinal regression.

Table 3 The major reasons for not performing CO-CPR or S-CPR

Reasons	CO-CPR (%)			S-CPR (%)		
	All	Male	Female	All	Male	Female
Lack of confidence	26.6	23.3	29.0†	28.7	24.8	31.5†
Fear of harming the victim	23.4	20.9	25.3†	21.5	17.6	24.4†
Fear of causing trouble (legal trouble)	20.7	23.8	18.4†	20.6	23.9	18.2†
Fear of disease transmission	7.7	9.0	6.8†	22.9	22.9	22.9

* $p < 0.05$.

†** $p < 0.01$, the χ^2 test were between the gender groups.

CO-CPR, chest compression-only CPR; CPR, cardiopulmonary resuscitation; S-CPR, standard CPR.

Willingness and factors influencing in performing CO-CPR and S-CPR

In all eight hypothetical scenarios, people were reluctant to perform S-CPR compared with CO-CPR, which was in accordance with previous studies.^{13 19} MMV was more intimate so the reluctance is stronger, and the relationship with the victim influences their willingness.^{16 20 21} In contrast with the past, the public’s overall willingness has increased, with the updating guidelines and evolving society.^{22 23} Our results suggest that the public in poor health or with sick family members present more positive attitudes and willingness towards CPR. We believe that this may be due to compassion and empathy, though there were some differences in previous studies.^{15 16 21} Given that first-aid courses are not compulsory except in medical schools in China, those in medical-related occupations view CPR more positively. Overall, the positive attitudes shown in this study should encourage us to disseminate CPR knowledge to a larger population, and CPR training can be enhanced and directed towards specific groups.




Figure 3 Proportions (%) of scored confidence to perform cardiopulmonary resuscitation (CPR) on emergency pre-CPR and post-CPR online training. Scores from ‘1’ (completely not confident) to ‘10’ (very confident) quantified the confidence level. The confidence change was statistically significant ($p < 0.05$).



Reluctance to perform CO-CPR and S-CPR

Our survey indicated that the top three reasons for being unwilling to perform CO-CPR were lack of confidence, fear of harming the victim and causing legal trouble. Men mostly chose 'fear of causing trouble (legal trouble)' as their major obstacle, whereas women mainly had a lack of confidence. Maybe men are more concerned with self-protection, while women exhibit more sympathy in emergencies.²² Solid confidence builds on adequate theoretical knowledge and skills, otherwise CPR will carry more risk.^{21 24} In addition, fear of causing legal trouble, which refers to the absence or imperfection of relevant legislation, also contributes to the performing reluctance.¹⁵ This suggests that the government implements legislation to protect performers and provides training to build confidence towards bystander CPR. As for S-CPR, the main obstacle was lack of confidence (28.7%) as well, especially among women (31.5%). The second difficulty, fear of disease transmission (22.9%), covered a large percentage (94%) in Sweden²⁵ and 58.9% in South Africa,²⁶ but in Crimea only about 2.1%–9.7% of people worried about disease transmission.²¹ These differences may stem from cultural and regional epidemiological differences such as the high HIV prevalence in South Africa.²⁷ However, the '2010 Cardiopulmonary Resuscitation (CPR) and the American Heart Association Cardiovascular Emergency Guidelines' allows non-medical professionals to perform CO-CPR, to encourage a wider public to participate in first aid.²⁸ Some studies have indicated that performing CO-CPR can still improve the patients with cardiac arrest survival.^{29–31} We encourage the public to actively engage in CO-CPR first when confronting OHCA, to improve the rate of survival. It is reported that specially modified curriculum, targeting preidentified obstacles to providing CPR, can increase public willingness to attempt it.³² In light of these barriers towards performing CPR, our training will emphasise the importance of CPR, and will target the different genders, in order to achieve a greater overcoming of the obstacles.

The necessity of online CPR training promotion in China

CPR training has a positive impact on the attitudes towards performing CPR, which improves the survival rate of OHCA.^{33 34} In our study, 34.8% respondents had previous CPR training, which was much more than the study of Chen *et al* in 2015 (8%).¹⁴ Although the average rate of CPR training (26%) in China has improved during these years,³⁵ it is still lower in contrast with 53% in Crimea,²¹ 54.7% in the USA³⁶ and 55.7% in Australia.³⁷ The primary reasons for not receiving CPR training were not knowing where to study and no free training locations, which were similar to those noted in a previous investigation.²⁶ Since 2015, AHA has recommended online training as resources for teaching and learning. Students take video-based courses before simulation training, with instructors providing feedback to standardise their performing.³⁸ It is noteworthy that public confidence in performing CPR improved 2.4-fold after our online training, which is consistent with the study of Cason and Stiller.¹⁰ In recent years, online training is an a feasible way for the public to gain first-aid knowledge, and according

to several studies has brought a similar quality to the learning of CPR techniques.^{39–41} Therefore, as a nation, we should call for full use of online platforms, to provide quality and free CPR training for the wider public.

Limitations

Our study has several limitations. First, this survey was released through the internet to the public who had attended our online CPR training. These respondents usually paid more attention to CPR, thus leading to more optimistic results. Still, it does show the overall benefit of CPR online training. Second, our online training and questionnaire focused on theoretical knowledge, rather than on the practical skills of CPR. Future investigations should explore the practical skills more and then fully evaluate the effects of online CPR training. Finally yet importantly, the majority of respondents who were motivated to learn CPR were 18–25 years old. The results of our study might therefore be more positive than in reality, and each limitation warrants further research.

CONCLUSIONS

The public has overwhelmingly positive attitudes and willingness to attempt CPR after online training. There are many factors influencing their attitudes and willingness, including gender, age and occupation. Nevertheless, the public still show a reluctance to some extent to perform CO-CPR or S-CPR, especially the latter. Obstacles to CPR include lack of confidence, fear of harming victim, fear of causing legal trouble or fear of disease transmission. Notably, online CPR training leads to a remarkable improvement in the public's confidence, confirming its beneficial effects and advantages when compared with the effects thus far of offline training. By taking advantage of online CPR training, it can be more convenient for the public to get free and flexible training. Considering the low bystander CPR rate in China and our study results, we should emphasise the importance of CPR and focus on overcoming the psychological and operational obstacles in performing CPR.

Acknowledgements The authors would like to thank all the participants who made this study possible. They are grateful to two Emergency Medicine Specialists in Zhongnan Hospital for providing valuable comments on the questionnaire design.

Contributors YJ was the first author. She analysed the data, interpreted the data and was a major contributor in writing the manuscript. BW participated in formulation of the figure, in data processing and in revising the manuscript. LL took part in the questionnaire design and modified the manuscript. JL interpreted the data and modified the manuscript. XJ designed the study, collected data and gave critical appraisal of the manuscript. All authors read and approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the ethical review committee of Zhongnan Hospital of Wuhan University, Number: 2019048.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. No additional data are available

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Yi Jiang <http://orcid.org/0000-0002-3531-0637>

REFERENCES

- Berdowski J, Berg RA, Tijssen JGP, *et al*. Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation* 2010;81:1479–87.
- National Health Commission of the People's Republic of China. *The transcript of National Health Commission's regular press conference on December 26, 2019*, 2019.
- Bakke HK, Wisborg T. We need to include bystander first aid in trauma research. *Scand J Trauma Resusc Emerg Med* 2017;25:32.
- Xu F, Zhang Y, Chen Y. Cardiopulmonary resuscitation training in China: current situation and future development. *JAMA Cardiol* 2017;2:469–70.
- Hock OME, SS D, Amatullah D, *et al*. Outcomes for out-of-hospital cardiac arrests across 7 countries in Asia: the pan Asian resuscitation outcomes study (PAROS). *Resuscitation* 2015;96.
- Gräsner J-T, Lefering R, Koster RW, *et al*. EuReCa ONE-27 nations, one Europe, one registry: a prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation* 2016;105:188–95.
- Potts J, Lynch B, Jerry P. The American heart association CPR Anytime program: the potential impact of highly accessible training in cardiopulmonary resuscitation. *J Cardiopulm Rehabil* 2006;26:346–54.
- Malta Hansen C, Zinckernagel L, Ersbøll AK, *et al*. Cardiopulmonary Resuscitation Training in Schools Following 8 Years of Mandating Legislation in Denmark: A Nationwide Survey. *J Am Heart Assoc* 2017;6.
- Semeraro F, Wingen S, Schroeder DC, *et al*. Kids save lives implementation in Europe: a survey through the ERC research net. *Resuscitation* 2016;107:e7–9.
- Cason CL, Stiller J. Performance outcomes of an online first aid and CPR course for laypersons. *Health Educ J* 2011;70:458–67.
- Kappus RM, McCullough G. The feasibility of a novel method of bystander CPR training: a pilot study. *Am J Emerg Med* 2020;38:594–7.
- Taniguchi T, Sato K, Fujita T, *et al*. Attitudes to bystander cardiopulmonary resuscitation in Japan in 2010. *Circ J* 2012;76:1130–5.
- AM-H H, Mizubuti GB, Wan S. Improving the success rate of chest compression-only CPR by untrained Bystanders in adult out-of-hospital cardiac arrest: maintaining airway patency may be the way forward. *Anesth Analg* 2018;126.
- Chen Z-Q, Zhao Y, Lu Z-H, *et al*. Awareness and attitudes of Chinese students towards cardiopulmonary resuscitation. *Emerg Med J* 2010;27:907–10.
- Huang Q, Hu C, Mao J. Are Chinese students willing to learn and perform bystander cardiopulmonary resuscitation? *J Emerg Med* 2016;51:712–20.
- Pei-Chuan Huang E, Chiang W-C, Hsieh M-J, *et al*. Public knowledge, attitudes and willingness regarding bystander cardiopulmonary resuscitation: a nationwide survey in Taiwan. *J Formos Med Assoc* 2019;118:572–81.
- Krammel M, Schnaubelt S, Weidenauer D, *et al*. Gender and age-specific aspects of awareness and knowledge in basic life support. *PLoS One* 2018;13:e0198918.
- Christenson J, Nafziger S, Compton S, *et al*. The effect of time on CPR and automated external defibrillator skills in the public access defibrillation trial. *Resuscitation* 2007;74:52–62.
- Becker T, Maciel C, Gul S, *et al*. Bystander CPR: public perception and barriers. *Critical Care Medicine* 2019;47.
- Becker TK, Gul SS, Cohen SA, *et al*. Public perception towards bystander cardiopulmonary resuscitation. *Emerg Med J* 2019;36:660–5.
- Birkun A, Kosova Y. Social attitude and willingness to attend cardiopulmonary resuscitation training and perform resuscitation in the Crimea. *World J Emerg Med* 2018;9:237–48.
- Lu C, Jin Y, Meng F, *et al*. An exploration of attitudes toward bystander cardiopulmonary resuscitation in university students in Tianjin, China: a survey. *Int Emerg Nurs* 2016;24:28–34.
- Moon S, Ryou HW, Ahn JY, *et al*. A 5-year change of knowledge and willingness by sampled respondents to perform bystander cardiopulmonary resuscitation in a metropolitan city. *PLoS One* 2019;14:e0211804.
- Dobbie F, MacKintosh AM, Clegg G, *et al*. Attitudes towards bystander cardiopulmonary resuscitation: results from a cross-sectional general population survey. *PLoS One* 2018;13:e0193391.
- Axelsson A, Thorén A, Holmberg S, *et al*. Attitudes of trained Swedish lay rescuers toward CPR performance in an emergency. A survey of 1012 recently trained CPR rescuers. *Resuscitation* 2000;44:27–36.
- Ojifinni K, Motara F, Laher AE. Knowledge, attitudes and perceptions regarding basic life support among teachers in training. *Cureus* 2019;11:e6302.
- Global Aids Update. Ending AIDS: progress towards 90-90-90 targets 2017. Available: https://www.unaids.org/sites/default/files/media_asset/Global_AIDS_update_2017_en.pdf [Accessed 1 Jul 2020].
- Association AH. 2010 American heart association guidelines for cardiopulmonary resuscitation and emergency CardiovascularCare science, 2010. Available: http://circ.ahajournals.org/content/vol122/18_suppl_3/
- Riva G, Ringh M, Jonsson M, *et al*. Survival in out-of-hospital cardiac arrest after standard cardiopulmonary resuscitation or chest compressions only before arrival of emergency medical services. *Circulation* 2019;139:2600–9.
- Rea TD, Fahrenbruch C, Culley L, *et al*. CPR with chest compression alone or with rescue breathing. *N Engl J Med Overseas Ed* 2010;363:423–33.
- Iwami T, Kitamura T, Kiyohara K, *et al*. Dissemination of chest Compression-only cardiopulmonary resuscitation and survival after out-of-hospital cardiac arrest. *Circulation* 2015;132:415–22.
- Ross EM, Redman TT, Mapp JG, *et al*. Stop the bleed: the effect of hemorrhage control education on laypersons' willingness to respond during a traumatic medical emergency. *Prehosp Disaster Med* 2018;33:127–32.
- Fratta KA, Bouland AJ, Vesselinov R, *et al*. Evaluating barriers to community CPR education in Baltimore, Maryland, USA. *Am J Emerg Med* 2019.
- Bouland AJ, Halliday MH, Comer AC, *et al*. Evaluating barriers to bystander CPR among laypersons before and after Compression-only CPR training. *Prehosp Emerg Care* 2017;21:662–9.
- Chen M, Wang Y, Li X, *et al*. Public knowledge and attitudes towards bystander cardiopulmonary resuscitation in China. *Biomed Res Int* 2017;2017:3250485.
- Hawkes CA, Brown TP, Booth S, *et al*. Attitudes to cardiopulmonary resuscitation and defibrillator use: a survey of UK adults in 2017. *J Am Heart Assoc* 2019;8:e008267.
- Cartledge S, Saxton D, Finn J, *et al*. Australia's awareness of cardiac arrest and rates of CPR training: results from the Heart Foundation's HeartWatch survey. *BMJ Open* 2020;10:e033722.
- Bhanji F, Donoghue AJ, Wolff MS, *et al*. Part 14: education: 2015 American heart association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2015;132:S561–73.
- Celik S. A media comparison study on first aid instruction. *Health Educ J* 2013;72:95–101.
- Madou T, Iserbytt P, Tom M. Mastery versus self-directed blended learning in basic life support: a randomised controlled trial. *Acta Cardiol* 2019;1–7.
- Lynch BC, Einspruch EL. Brief, self-directed CPR training for lay responders is as effective as traditional instruction. *Circulation* 2004;110:414–14.

Dear friends,

In order to have a deeper understanding of public knowledge and attitudes toward cardiopulmonary resuscitation (CPR), as well as willingness to perform CPR under different scenarios, we hereby conduct this research and hope to get your support and cooperation. Your answers are only for the statistical analysis of the researcher. There is no correct or incorrect answer to each question, and the questions do not involve personal privacy, nor any impact on your work and life. We promise to keep all information confidential. Therefore, you are supposed to answer carefully and truthfully in accordance with the instructions, or make such judgment as you think appropriate. The truthfulness of your answers is very important to the accuracy of our research. Thank you for your cooperation! **(If you don't want to fill in the form, just quit.)**

Demographic characteristics

1. Gender:

Male Female

2. Age(years):

<18 18-25 26-35 36-45 56-55 ≥56

3. Education level

Primary school Junior high school High school

Associate's degree Bachelor's degree Master's degree or above

4. Occupation:

Medical-related Non-medical-related

5. Health condition:

Very poor Poor General Good

6. Do you have any family member with life-threatening diseases?

Yes No

7. Have you ever encountered someone in need of cardiopulmonary resuscitation (CPR) performance out of hospital?

Yes No

Part.1 Knowledge and attitudes toward CPR.

Instruction: Please choose the answer you think is most suitable according to the degree of the following description and your actual situation.

1. Do you understand what is cardiopulmonary resuscitation (CPR)?

Yes No

2. Would you like to learn CPR knowledge?

Yes No Do not know/Haven't thought about it

3. Reasons for learning CPR (multiple-choice):

A family history of cardiac arrest

Avoiding unnecessary death

Helping others

Other

4. Would you like to disseminate CPR knowledge?

Yes No Don't know/Haven't thought about it

5. What is your opinion on developing CPR training courses in university?

It should be compulsory for all students

It should be optional for all students

It should be compulsory for medical-related students

Do not support CPR training courses in university

6. How important do you think chest compression is for cardiac arrest patients?

Uncertain Not important Not very important Very important

7. How important do you think mouth-to-mouth ventilation is for patients suffered cardiac arrest?

Uncertain Not important Not very important Very important

8. Have you ever received any training in CPR?

Yes No

9. The reasons that you have not received CPR training (multiple-choice):

- Not paying attention to or thinking about it
- Want but do not know where to learn
- No place for free training
- Too busy with work and study to learn
- Want to learn, but do not want to take examination after training

10. Your confidence in performing CPR in the event of an emergency out of hospital is (1 for "not confident at all", 10 for "very confident") :

pre-online CPR training: 1 2 3 4 5 6 7 8 9 10

post-online CPR training: 1 2 3 4 5 6 7 8 9 10

Part.2 Willingness to perform CPR in eight hypothetical scenarios

Instruction: In the following questionnaire, different scenarios are set respectively. Please check the answer you think is the most suitable according to the degree of the following description and your actual situation.

A complete male stranger, who is around 18-50 years old, has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR to him?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of chest compression
- Lack of confidence
- Unwilling to touch strangers

- Nervousness and fear
- Fear of harming the victim
- None of my business
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation (Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch strangers
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- The stranger being male
- Feel sick
- Other

A complete female stranger, who is around 18-50 years old, has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR to her?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to her?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of chest compression

Lack of confidence

Unwilling to touch strangers

Nervousness and fear

Fear of harming the victim

None of my business

Other

4. The reason you don't want to perform mouth-to-mouth ventilation

(Multiple-choice):

Fear of causing trouble (legal trouble)

Fear of disease transmission

Poor knowledge of mouth-to-mouth ventilation

Lack of confidence

Unwilling to touch strangers

Nervousness and fear

Fear of harming the victim

None of my business

Feel embarrassed

The stranger being female

Feel sick

Other

A complete old stranger (regardless of gender) has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR to him/her?

Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him/her?

Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

Fear of causing trouble (legal trouble)

Fear of disease transmission

- Poor knowledge of chest compression
- Lack of confidence
- Unwilling to touch strangers
- Nervousness and fear
- Fear of harming the victim
- None of my business
- The stranger being old
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation

(Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch strangers
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- The stranger being old
- Feel sick
- Other

A close family member (parents/sisters/brothers) has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR to him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of disease transmission
- Poor knowledge of chest compression
- Lack of confidence
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation (Multiple-choice):

- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch others
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- Feel sick
- Other

A general relative has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR for him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of disease transmission
- Poor knowledge of chest compression

- Lack of confidence
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation (Multiple-choice):

- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch others
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- Feel sick
- Other

A friend has suffered cardiac arrest and needs immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR for him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of chest compression
- Lack of confidence
- Unwilling to touch others

- Nervousness and fear
- Fear of harming the victim
- None of my business
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation

(Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch others
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- Feel sick
- Other

At the scene of the traffic accident, a person, badly injured and bleeding with wounds, is in need of immediate CPR. You are the first witness without any other helpers.

1. Will you perform chest compression-only CPR for him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

2. Will you perform standard CPR (including chest compression and mouth-to-mouth ventilation) to him/her?

- Absolutely yes Maybe yes Maybe no Absolutely no

3. The reason you don't want to perform CPR (Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of chest compression
- Lack of confidence

- Unwilling to touch strangers
- Nervousness and fear
- Fear of harming the victim
- None of my business
- The stranger being injured and bleeding
- Other

4. The reason you don't want to perform mouth-to-mouth ventilation

(Multiple-choice):

- Fear of causing trouble (legal trouble)
- Fear of disease transmission
- Poor knowledge of mouth-to-mouth ventilation
- Lack of confidence
- Unwilling to touch strangers
- Nervousness and fear
- Fear of harming the victim
- None of my business
- Feel embarrassed
- The stranger being injured and bleeding
- Feel sick
- Other

Table S1 Lists of different scenarios.

Description of different scenarios.
(1) Male Stranger A strange man who is between the ages of 18 and 50 has sudden cardiac arrest and needs an immediate CPR. You are the first eyewitness and there are no other people on the spot to help.
(2) Female Stranger A strange woman who is between the ages of 18 and 50 has sudden cardiac arrest and needs an immediate CPR. You are the first eyewitness and there are no other people on the spot to help.
(3) A Child An unknown child (regardless of gender) has cardiac arrest and requires an immediate CPR. You are the first eyewitness and there is no other person to help.
(4) An Elderly Person An unknown elderly person (regardless of gender) has cardiac arrest and requires immediate CPR. You are the first eyewitness and there is no other person to help.
(5) A Close Family Member Your close family member (parent, sibling, etc.) has cardiac arrest and requires immediate CPR. You are the first eyewitness and no one else can help.
(6) A Relative A relative (cousin, uncle, aunt, etc.) has cardiac arrest and requires immediate CPR. You are the first eyewitness and there is no other person to help.
(7) A Friend Your friend has cardiac arrest and requires immediate CPR. You are the first eyewitness and there is no other person to help.
(8) A Trauma Victim At the scene of a traffic accident, there is a trauma victim with serious injury and bleeding, and you need to perform CPR immediately. You are the first eyewitness and there is no other people on the scene to help.

CPR: Cardiopulmonary resuscitation

Table S2 Respondents' attitudes toward cardiopulmonary resuscitation (CPR) (n=1888).

Responses	N (%)
Understanding what is CPR	
Yes	1826 (96.7)
No	62 (3.3)
Would like to learn CPR knowledge	
Yes	1857 (98.4)
No	31 (1.6)
Reasons for learning CPR (multiple-choice)	
A family history of cardiac arrest	512 (27.1)
Avoiding unnecessary death	1654 (87.6)
Helping others	1631 (86.4)
Others	386 (20.4)
Would like to disseminate CPR knowledge	
Yes	1548(82.0)
No	48(2.5)
Uncertain	292(15.5)
Opinions on developing CPR training courses in university	
Compulsory courses for all the students	1263(66.9)
Optional courses for all the students	509 (27.0)
Only for medical-related students	109 (5.8)
Negative answer	7 (0.3)
How important do you think CC is for cardiac arrest	
Uncertain	17 (0.9)
Not important	1 (0.0)
Not very important	7 (0.3)
Very important	1863 (98.8)
How important do you think MMV is for cardiac arrest	
Uncertain	92 (4.9)
Not important	15 (0.8)
Not very important	235 (12.4)
Very important	1546(81.9)
Ever trained in CPR	
Yes	657 (34.8)
No	1231 (65.2)
Reasons for not attending CPR training (multiple-choice)	
Do not know where the training is	961 (50.9)
Cost	709 (37.6)
Lack of time	305 (16.2)
Others	387 (20.5)

CPR = cardiopulmonary resuscitation; CC: Chest compressions; MMV: Mouth-to-mouth ventilation

Table S3 Willingness to perform CO-CPR and S-CPR in different scenarios.

Scenarios	CO-CPR (N, %)		S-CPR (N, %)		P value
	Yes	No	Yes	No	
Male Stranger	1740 (92.2)	148 (7.8)	1349 (71.5)	539 (28.6)	<0.01
Female Stranger	1741 (92.2)	147 (7.8)	1476 (78.2)	412 (21.8)	
A Child	1740 (92.2)	148 (7.8)	1618 (85.7)	270 (14.3)	
An Elderly Person	1585 (84.0)	303 (16.1)	1312 (69.5)	576 (30.5)	
A Close Family Member	1865 (98.8)	23 (1.2)	1831 (97.0)	57 (3.0)	
A Relative	1829 (96.9)	59 (3.1)	1696 (89.8)	192 (10.2)	
A Friend	1845 (97.7)	43 (2.3)	1751 (92.7)	137 (7.3)	
A Trauma Victim	1676 (88.8)	212 (11.2)	1452 (76.9)	436 (23.1)	
Total	1477 (78.2)	411 (21.8)	1155 (61.2)	733 (38.8)	

CO-CPR: chest compression-only CPR; S-CPR: standard CPR.

The χ^2 test were between the CO-CPR and S-CPR groups.

Table S4 Predictive factors associated with performing CO-CPR and S-CPR using ordinal regression.

Respondent Variables	Reference group	CO-CPR		S-CPR	
		Odds Ratio(95%CI)	P value	Odds Ratio(95%CI)	P value
Gender (Male)	Female	0.687(-0.153-0.858)	0.001	1.039(0.860-1.254)	0.694
Occupation (Medical-related)	Non-medical-related	1.315(0.573-1.774)	0.073	1.510(1.191-1.914)	0.001
Health condition (Good) (General)	Poor	0.520(0.331-0.819)	0.005	0.645(0.426-0.977)	0.039
		0.897(0.712-1.130)	0.354	0.802(0.661-0.972)	0.025
Ever encountered someone in need of CPR performance (No)	Yes	0.473(0.283-0.791)	0.004	N/A	N/A
Ever trained in CPR (No)	Yes	0.820(0.640-1.05)	0.118	N/A	N/A

CPR: Cardiopulmonary resuscitation; CO-CPR: chest compression-only CPR; S-CPR: standard CPR.

Table S5 Reasons for not performing CO-CPR or S-CPR in different scenarios.

Reasons	CO-CPR (%)			S-CPR (%)		
	All	Male	Female	All	Male	Female
Lack of confidence	26.6	23.3	29.0**	28.7	24.8	31.5**
Fear of harming the victim	23.4	20.9	25.3**	21.5	17.6	24.4**
Fear of causing trouble (legal trouble)	20.7	23.8	18.4**	20.6	23.9	18.2**
Nervousness and fear	16.7	15.1	17.9**	20.0	16.3	22.8**
Poor knowledge of CPR	12.7	13.8	11.9**	17.6	17.8	17.4
Depending on victims' situation	11.9	12.9	11.1*	13.5	13.2	13.8
Fear of disease transmission	7.7	9.0	6.8**	22.9	22.9	22.9
Unwilling to touch strangers	5.2	5.9	4.7**	10.7	8.3	12.5**
Others	3.8	4.7	3.1**	4.9	5.4	4.5**
None of my business	0.7	1.1	0.5**	1.0	1.4	0.7**

CPR: Cardiopulmonary resuscitation; CO-CPR: chest compression-only CPR; S-CPR: standard CPR.

*P<0.05, **P<0.01, the χ^2 test were between the gender groups.