BMJ Open Trends and differences in sexual practices and sexually transmitted infections in men who have sex with men only (MSMO) and men who have sex with men and women (MSMW): a repeated cross-sectional study in Melbourne, Australia

Mario Martín-Sánchez ^{1,2}, Richard Case,¹ Christopher Fairley,^{1,3} Jane S Hocking,⁴ Catriona Bradshaw,^{1,3} Jason Ong ^{1,3}, Marcus Y Chen,^{1,3} Eric P F Chow ^{1,3,4}

To cite: Martín-Sánchez M, Case R, Fairley C, *et al.* Trends and differences in sexual practices and sexually transmitted infections in men who have sex with men only (MSMO) and men who have sex with men and women (MSMW): a repeated cross-sectional study in Melbourne, Australia. *BMJ Open* 2020;**10**:e037608. doi:10.1136/bmjopen-2020-037608

► Prepublication history and additional material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-037608>).

Received 09 February 2020
Revised 19 October 2020
Accepted 29 October 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Associate Professor Eric P F Chow;
eric.chow@monash.edu and
Dr Mario Martín-Sánchez;
mmartin@hku.hk

ABSTRACT

Objectives In the 2010s, there has been an increase in sexually transmitted infections (STI) in men who have sex with men (MSM) in Australia, and since 2015 also in urban heterosexuals. Men who have sex with both men and women (MSMW) have characteristics that may differ from both men who have sex with men only (MSMO) and heterosexual men. We aimed to compare the sexual practices and the trends in HIV/STI positivity between MSMO and MSMW.

Design Repeated cross-sectional study.

Setting A sexual health centre in Melbourne, Australia.

Participants MSM aged 18 years and above who attended the Melbourne Sexual Health Centre for the first time between 2011 and 2018. This includes 12 795 MSMO and 1979 MSMW.

Primary outcome measures Demographic characteristics, sexual practices and HIV/STI positivity.

Results Compared with MSMW, MSMO were more likely to practice anal sex and to have condomless receptive anal sex with casual male partners, and less likely to have a current regular relationship. Over the 8-year period, there was an increase in condomless receptive anal sex with casual male partners for both groups (MSMO: from 46.2% to 63.3%, $p_{\text{trend}} < 0.001$; MSMW: from 41.3% to 57.9%, $p_{\text{trend}} = 0.011$). Syphilis positivity increased in MSMO (from 5.5% to 7.9%, $p_{\text{trend}} = 0.012$) and MSMW (from 0.9% to 6.4%, $p_{\text{trend}} = 0.004$) and HIV remained stable. Gonorrhoea increased among MSMO from 2011 to 2014 (from 6.7% to 9.6%, $p_{\text{trend}} = 0.002$), and remained stable from 2015 to 2018. MSMO had higher odds of testing positive for gonorrhoea (adjusted OR (aOR) 1.36, 95% CI 1.13 to 1.64), chlamydia (aOR 1.39, 95% CI 1.16 to 1.67), syphilis (aOR 1.74, 95% CI 1.37 to 2.22) and HIV (aOR 4.60, 95% CI 2.43 to 8.70) than MSMW.

Conclusions MSMW have overall lower condomless sex and lower HIV/STI positivity. In the last years, changes in sexual practices in MSM have affected both MSMW and MSMO leading to an increased STI risk.

Strengths and limitations of this study

- We compared sexual practices and HIV/sexually transmitted infections positivity in a high number of men who have sex with men only (N=12 795) and men who have sex with men and women (N=1979) over an 8-year period.
- We provide a comprehensive description of sexual practices distinguishing between condom use with regular or casual sexual partners, and in receptive or insertive anal sex.
- The study was conducted in one urban major sexual health clinic and it is possible that our findings may not be generalisable to other settings.
- We were unable to analyse the statistical significance of the temporal trends during the whole study period for gonorrhoea and chlamydia due to a change in the diagnostic test in 2015.

INTRODUCTION

Despite considerable public health efforts and biomedical advances,¹ the rate of sexually transmitted infections (STIs) is increasing at an alarming rate particularly among gay, bisexual and other men who have sex with men (MSM) worldwide.² Among the Australian population, the notification rate, defined as the number of new cases reported to the national infectious diseases surveillance system, increased an 80% for gonorrhoea (from 65.5 to 118.0 per 100,000), a 14% for chlamydia (from 364.5 to 416.8 per 100,000) and has more than doubled for syphilis (from 7.8 to 18.3 per 100,000) between 2013 and 2017.³ In males, most of the early rises

in gonorrhoea and syphilis were accounted for by rises in MSM. However, in the late 2010s, there has been a dramatic increase in both infections in the Australian heterosexual population to levels not seen since the 1980s.⁴ In stark contrast to the rise in the notification rate of gonorrhoea and syphilis, there has been an 11% decline in the HIV notification rate from 4.5 per 100 000 in 2013 to 4.0 in 2017 among MSM in Australia, with most of the reduction occurring between 2016 and 2017 after the introduction of pre-exposure prophylaxis (PrEP) in late 2016.^{3 5–7}

Past studies have shown that sexual practices among MSM have changed over time (eg, an increase in condomless anal sex).^{8–12} However, there have been very limited studies examining whether there are any differences in sexual practices between men who have sex with men only (MSMO) and men who have sex with both men and women (MSMW or bisexual MSM).^{13 14} Therefore, this study aimed to compare the characteristics and sexual practices, and describe the trends in HIV/STI positivity in MSMO and MSMW attending a large sexual health clinic in Melbourne between 2011 and 2018. Additionally, we aimed to explore differences within MSMW depending on the most frequent gender of their sexual partners.

METHODS

We conducted a repeated cross-sectional analysis of retrospective data of MSM attending the Melbourne Sexual Health Centre (MSHC), Australia, between 2011 and 2018. MSHC is a public clinic that offers a range of free clinical services regarding sexual health. MSHC is the largest sexual health clinic in Victoria and provided more than 50 000 clinical consultations in 2019, approximately 40% of clients are MSM.⁵ On arrival, all clients are invited to complete a questionnaire via computer-assisted self-interviewing (CASI), which collects client's demographic characteristics (eg, age, country of birth, Aboriginal and Torres Strait Islander status), sexual practices (ie, gender of partners, type of partners (regular or casual), number of partners, condom use and anal sex practice (receptive anal sex (RAS) and insertive anal sex (IAS))), sex work status and injecting drug use practice. Data on sexual practices and injecting drug use were measured in the past 12 months, and sex work status was defined as clients who had ever worked in the sex industry during lifetime. Clients are allowed to decline to answer any questions they may wish to.

Clients who were male, aged 18 years and above, reported having sex with another man in the past 12 months, and attended MSHC for the first time between 2011 and 2018 were included in this analysis. We only included data from the client's first visit to MSHC to avoid any bias including men with repeated visits as they might be at a higher risk of HIV/STI with different sexual practices. We categorised MSM into two main categories based on their self-reported sexual practices: (1) MSMO and (2) MSMW. We further categorised MSMW into two

groups: (1) MSM who reported more female partners than male partners in the past 12 months (MSMW-W) and (2) MSM who reported more male partners than female partners in the past 12 months (MSMW-M). We compared the characteristics between the MSMW-M and MSMW-W. Men with an equal number of male and female partners in the past 12 months (N=336) were excluded in the comparisons among MSMW subgroups but they were included in the overall MSMW category.

We also extracted the HIV/STI testing results on the day. STI testing results included syphilis (serologically confirmed by rapid plasma regain test, *Treponema pallidum* enzyme immunoassay and *Treponema pallidum* particle agglutination assay and including all syphilis stages), gonorrhoea (stratified by anatomical site: anorectal, urethral and oropharyngeal) and chlamydia (stratified by anatomical site: anorectal and urethral). For HIV, we analysed separately incident cases (diagnosed on the day of the first visit using screening assay followed by a confirmatory Western Blot assay) and MSM with a previous HIV diagnosis visiting MSHC for the first time. There was a major change in gonorrhoea and chlamydia testing at our clinic.¹⁵ Prior to 2015, gonorrhoea was diagnosed by using culture and chlamydia was diagnosed by nucleic acid amplification test using the BD ProbeTec Strand Displacement Amplification Assay (Becton, Dickinson and Co, Sparks, Maryland, USA). From March 2015 onwards, testing for both gonorrhoea and chlamydia was performed using the Aptima Combo 2 Transcription-Mediated Amplification Assay (AC2) (Hologic Gen-Probe, San Diego, California, USA). HIV and syphilis testing methods did not change during the study period.

Before July 2015, screening for urethral gonorrhoea was only performed in MSM with urethral symptoms while screening for urethral chlamydia was performed in all MSM as per the Australian guidelines.¹⁶ However, since July 2015, all MSM were screened for both urethral gonorrhoea and chlamydia regardless of the presence of symptoms as per our clinic policy.¹⁷ Therefore, the positivity for urethral gonorrhoea was calculated as the number of men tested positive divided by the number of men who were tested for urethral chlamydia to avoid the bias of testing urethral gonorrhoea only among symptomatic MSM until July 2015, and this approach was also used elsewhere.¹⁸ In addition, routine screening for oropharyngeal chlamydia among all MSM at MSHC were introduced in April 2017; before then, only MSM who reported as a contact of infection were tested. Routine screening for HIV, syphilis, oropharyngeal and anorectal gonorrhoea, and anorectal chlamydia were conducted among all MSM and the screening guidelines did not change over the study period at MSHC.

χ^2 test was used to compare the demographic characteristics and sexual practices between risk groups. We examined the annual trends of sexual practices and HIV/STI positivity for MSMO and MSMW using χ^2 trend test. Annual trends for condomless IAS and RAS with causal male partners were examined separately. We reported any

condomless sex (ie, IAS and/or RAS) with regular male partners for the trend analysis due to the small number of participants who reported having regular male partners. For chlamydia and gonorrhoea positivity, we calculated the annual trends for 2011–2014 and 2015–2018 separately due to the change of the diagnostic assays.¹⁵ Univariable and multivariable logistic regression was used to examine the association between sexual practice (MSMW or MSMO) and HIV/STI positivity. We assessed HIV and each STI separately, and this included (1) gonorrhoea at any anatomical site, (2) chlamydia at any anatomical site, (3) syphilis, (4) new HIV diagnosis on the day of the first visit and (5) previous HIV diagnosis, as the dependent variables. Therefore, five different logistic regression models were conducted, and the independent variables included sexual practice (being MSMW or MSMO and using MSMW as reference), the year of the visit and all potential confounding factors (ie, variables with $p < 0.20$ in the univariable analyses) in the multivariable analysis. Missing data were presented as 'no information'. We repeated the same procedure using sexual practice categorised in MSMO, MSMW-M and MSMW-W as the independent variable with MSMW-W as the reference group. We reported the crude odds ratio (OR) and adjusted OR, and their respective 95% confidence interval (CI). The 0.05 level was used for statistical significance in all the analysis. All statistical analyses were conducted using SPSS V.25.

Patient and public involvement

Patients were not directly involved in this study; only data gathered retrospectively and coming from electronic health records was used.

RESULTS

Demographic characteristics

There were 18851 MSM attending MSHC for the first time between 2011 and 2018. We excluded 1909 MSM (10.1%) who had had sexual contact with another man but declined to report the number of male sexual

partners, 1630 MSM (8.6%) who reported no male sexual partner in the past 12 months at the current visit but had reported male partners at subsequent visit at the clinic and 538 (2.9%) duplicate records from the same individual on the same day. The remaining 14774 men were included in this data analysis, and the number of men increased from 1215 in 2011 to 2468 in 2018 but the proportion of MSMO and MSMW remained stable over the period (table 1). Overall, there were 12795 MSMO (86.6%) and 1979 MSMW (13.4%). Among 1979 MSMW, 804 (40.6%) were MSMW-W, 839 (42.4%) MSMW-M and 336 (17.0%) had an equal number of male and female partners in the past 12 months.

Among the 14774 MSM, the median age was 27 years (interquartile range [IQR]: 23–33) and MSMW were slightly older than MSMO (median age: 29 vs 27; $p < 0.001$) (table 2). The proportion of men born in Australia was higher in MSMW than in MSMO (55.3% vs 48.5%, $p < 0.001$). The proportion of injecting drug use in the past 12 months was higher in MSMW than MSMO (2.0% vs 1.6%, $p = 0.032$; table 2); but injecting drug use was similar between MSMW-W and MSMW-M (2.5% vs 2.1%, $p = 0.777$; online supplemental table S1).

Sexual practices

The proportion of men who reported sex overseas in the past 12 months was comparable between MSMW and MSMO (37.1% vs 34.5%; $p = 0.060$); and more MSMW-W (43.0%) reported sex overseas than MSMW-M (34.6%; $p = 0.002$) (online supplemental table S1). Less than half (40.1%) of MSMO had a male regular partner. Among MSMW, 8.5% had a regular male partner, 33.5% a regular female partner, and 4.6% had regular male and female partners (table 2).

The median number sexual partners (regardless gender of the partners) in the past 12 months, was higher among MSMW (7, IQR: 5–12 in MSMW-M; and 6, IQR: 4–11 in MSMW-W) than MSMO (5, IQR: 3–10; $p < 0.001$) (online supplemental table S1). The proportion of men with ≥ 5 sexual partners increased significantly in MSMW

Table 1 Number of men who have sex with men attending the Melbourne Sexual Health Centre between 2011 and 2018, stratified by men who have sex with men only (MSMO) and men who have sex with men and women (MSMW)

	2011	2012	2013	2014	2015	2016	2017	2018	Total
MSMO	1047 (86.2)	1282 (87.7)	1422 (85.5)	1433 (85.9)	1658 (86.5)	1872 (86.7)	1938 (87.1)	2143 (86.8)	12795 (86.6)
MSMW	168 (13.8)	180 (12.3)	241 (14.5)	235 (14.1)	258 (13.5)	286 (13.3)	286 (12.9)	325 (13.2)	1979 (13.4)
MSMW-W	70 (41.7)	70 (38.9)	98 (40.7)	108 (46.0)	91 (35.3)	114 (39.9)	116 (40.6)	137 (42.2)	804 (40.6)
MSMW-M	64 (38.1)	78 (43.3)	96 (39.8)	90 (38.3)	127 (49.2)	117 (40.9)	124 (43.4)	143 (44.0)	839 (42.4)
MSMW(=)	34 (20.2)	32 (17.8)	47 (19.5)	37 (15.7)	40 (15.5)	55 (19.2)	46 (16.1)	45 (13.8)	336 (17.0)
Total	1215	1462	1663	1668	1916	2158	2224	2468	14774

Data are presented as n (%) where n is the number of men in each category. The proportion of MSMO and MSMW was calculated using the total number of men each year as the denominator. The proportion of MSMW-W, MSMW-M and MSMW(=) was calculated using the total number of MSMW each year as the denominator. MSMW(=), men who have sex with men and women but reported an equal number of male and female partners in the past 12 months; MSMW-M, men who have sex with men and women and had more male than female sexual partners in the past 12 months; MSMW-W, men who have sex with men and women and had more female than male sexual partners in the past 12 months.

Table 2 Demographic characteristics and sexual practices among men who have sex with men attending a sexual health centre in Melbourne, 2011–2018, stratified by men who have sex with men only (MSMO) and men who have sex with men and women (MSMW)

	Men who have sex with men only (N=12 795)	Men who have sex with men and women (N=1979)	P value
	n (%)	n (%)	
Age (years), median (IQR)	27 (23–33)	29 (24–38)	<0.001
Country of birth			<0.001
Australia	6208 (48.5)	1094 (55.3)	
Overseas	6110 (47.8)	790 (39.9)	
No information	477 (3.7)	95 (4.8)	
Aboriginal and Torres Strait Islander Status			0.224
Indigenous origin	150 (1.2)	17 (0.9)	
Non-indigenous origin	11 595 (90.6)	1783 (90.1)	
No information	1050 (8.2)	179 (9.0)	
Sex overseas in the past 12 months			0.060
Yes	4415 (34.5)	735 (37.1)	
No	7571 (59.2)	1131 (57.2)	
No information	809 (6.3)	113 (5.7)	
Sex work during lifetime			0.574
Yes	71 (0.6)	13 (0.7)	
No	12 724 (99.4)	1966 (99.3)	
Injected drug use in the past 12 months			0.032
Yes	211 (1.6)	40 (2.0)	
No	12 363 (96.6)	1890 (95.5)	
No information	221 (1.7)	49 (2.5)	
No of female and male sexual partners in the past 12 months, median (IQR)	5 (3–10)	6 (4–11)	<0.001
No of male sexual partners in the past 12 months, median (IQR)	5 (3–10)	2 (1–5)	<0.001
No of female sexual partners in the past 12 months, median (IQR)	NA	2 (1–5)	–
Regular sexual partner			–
No regular partner	7513 (58.7)	1053 (53.2)	
Regular male partner	5132 (40.1)	169 (8.5)	
Regular female partner	NA	663 (33.5)	
Regular female and male partner	NA	91 (4.6)	
No information	150 (1.2)	3 (0.2)	
Anal sexual practice with regular male partners†			0.118
Receptive only	435 (8.5)	15 (5.8)	
Insertive only	465 (9.1)	16 (6.2)	
Receptive and insertive	3921 (76.4)	214 (82.3)	
No anal sex	234 (4.6)	9 (3.5)	
No information	77 (1.5)	6 (2.3)	
Condom use with regular male partners in the past 12 months for receptive anal sex*‡			0.005
Always	1241 (28.5)	85 (37.1)	
Not always	3115 (71.5)	144 (62.9)	
Condom use with regular male partners in the past 12 months for insertive anal sex§*			0.010
Always	1254 (28.6)	84 (36.5)	
Not always	3132 (71.4)	146 (63.5)	
Vaginal or anal sex with regular female partner*¶			–
Yes	NA	729 (96.7)	

Continued

Table 2 Continued

	Men who have sex with men only (N=12 795)	Men who have sex with men and women (N=1979)	P value
	n (%)	n (%)	
No	NA	10 (1.3)	
No information	NA	15 (2.0)	
Condom use with regular female partners in the past 12 months*, **			–
Always	NA	132 (18.1)	
Not always	NA	597 (81.9)	
Casual sexual partner			–
No casual partner	806 (6.3)	11 (0.6)	
Casual male partner(s)	11 969 (93.5)	194 (9.8)	
Casual female partner(s)	NA	30 (1.5)	
Casual female and male partner(s)	NA	1744 (88.1)	
No information	20 (0.2)	0 (0)	
Anal sexual practice with casual male partners			<0.001
Receptive only	1262 (9.9)	131 (6.6)	
Insertive only	1344 (10.5)	282 (14.2)	
Receptive and insertive	7835 (61.2)	1060 (53.6)	
No anal sex	1692 (13.2)	422 (21.3)	
No information	662 (5.2)	84 (4.2)	
Condom use with casual male partners in the past 12 months for receptive anal sex††*			0.012
Always	4207 (46.2)	597 (50.1)	
Not always	4890 (53.8)	594 (49.9)	
Condom use with casual male partners in the past 12 months for insertive anal sex*‡‡			0.223
Always	4276 (46.6)	649 (48.4)	
Not always	4903 (53.4)	693 (51.6)	
Vaginal or anal sex with casual female partner			–
Yes	NA	1686 (85.2)	
No	NA	56 (2.8)	
No information	NA	237 (12.0)	
Condom use with casual female partners in the past 12 months§§*			–
Always	NA	548 (32.5)	
Not always	NA	1138 (67.5)	

*‘Not always’ was defined as men who sometimes, usually or never used a condom with their sexual partners in the past 12 months (with female partners for vaginal and/or anal sex, with male partners for anal sex).

†Only including MSM with regular male partners or regular male and female partners.

‡Only including MSM that referred receptive only or receptive and insertive anal sex with regular male partners.

§Only including MSM that referred insertive only or receptive and insertive anal sex with regular male partners.

¶Only including MSM with regular female partners or regular male and female partners.

**Only including MSM that referred vaginal and/or anal sex with regular female partner.

††Only including MSM that referred receptive only or receptive and insertive anal sex with casual male partners.

‡‡Only including MSM that referred insertive only or receptive and insertive anal sex with casual male partners.

§§Only including MSM that referred vaginal or anal sex with casual female partners.

MSM, men who have sex with men; NA, not applicable.

(from 63.1% in 2011 to 70.2% in 2018, $p_{\text{trend}}=0.014$), as well as in MSMO (from 50.8% to 56.4%; $p_{\text{trend}}=0.002$) (online supplemental tables S2 and S3). In addition, the proportion of MSMW with ≥ 5 female sexual partners increased significantly from 23.2% in 2011 to 29.5% in 2018 ($p_{\text{trend}}=0.039$) (online supplemental table S3).

Anal sex practices with casual male partners varied between MSMO and MSMW. The proportion of MSMO having RAS only with casual male partners was higher (9.9%) than in MSMW (6.6%); however, the proportion of MSMO having IAS only with casual male partners was lower (10.5%) than in MSMW (14.2%; $p<0.001$). Condomless anal sex with casual male partners in the

Figure 1 Proportion of condomless anal and/or vaginal sex in the past 12 months with casual partners between 2011 and 2018, stratified by men who have sex with men only (MSMO) and men who have sex with men and women (MSMW). MSM reporting condomless anal and/or vaginal sex includes MSM who sometimes, usually or never used a condom with their sexual partners in the past 12 months.

past 12 months was more common in MSMO than in MSMW for RAS (53.8% *vs* 49.9%; $p=0.012$) but there was no significant difference in IAS. Two-thirds (67.5%) of MSMW had condomless vaginal and/or anal sex with female casual partners in the past 12 months (table 2). Among MSMO, condomless RAS with casual male partners in the past 12 months increased from 46.2% to 63.3% ($p_{\text{trend}} < 0.001$) and IAS from 45.8% to 64.2% ($p_{\text{trend}} < 0.001$) from 2011 to 2018 (figure 1 and online supplemental table S2). Similarly, among MSMW, there was an increase from 41.3% to 57.9% in condomless RAS ($p_{\text{trend}} = 0.011$), the proportion of condomless IAS with casual male partners did not change during the study period ($p_{\text{trend}} = 0.087$), and the proportion of condomless sex with casual female partners remained high and only increased marginally from 65.4% to 71.0% ($p_{\text{trend}} = 0.056$) (figure 1 and online supplemental table S3). MSMW-M compared with MSMW-W showed higher condomless sex with casual male partners (55.0% *vs* 46.1% for RAS; $p=0.004$ and 57.0% *versus* 47.1% for IAS; $p=0.001$) and lower condomless sex with casual female partners (60.9% *vs* 75.6%; $p < 0.001$) (online supplemental table S1).

Regarding regular male partners, both condomless RAS and condomless IAS were more commonly reported in MSMO (RAS: 71.5%; IAS: 71.4%) than in MSMW (RAS 62.9%; IAS: 63.5%; $p=0.005$; $p=0.010$) (table 1) and condomless anal sex regardless of IAS or RAS, increased from 69.2% to 78.4% in MSMO ($p_{\text{trend}} < 0.001$) but not in MSMW ($p_{\text{trend}} = 0.435$) (figure 1 and online supplemental tables S2, S3).

HIV/STI positivity

Overall, MSMO had a higher HIV, syphilis, gonorrhoea and chlamydia positivity than MSMW. MSMO had higher extragenital gonorrhoea and chlamydia positivity than MSMW; however, the positivity for urethral gonorrhoea and urethral chlamydia did not differ between the two groups (table 3). In the period 2011–2014, there was an increase in oropharyngeal and anorectal and oropharyngeal gonorrhoea among MSMO, while chlamydia remained stable. There was no increase in gonorrhoea and chlamydia positivity in the period 2015–2018 in either group (online supplemental tables S4 and S5). Among MSMW, MSMW-M had higher extragenital gonorrhoea and chlamydia positivity but similar urethral gonorrhoea and chlamydia, HIV and syphilis compared with MSMW-W (online supplemental table S6).

There was a 43.5% increase in syphilis positivity from 5.5% (43/778) in 2011 to 7.9% (152/1917) in 2018 ($p_{\text{trend}} = 0.012$) in MSMO (online supplemental table S4) and more than a sixfold increase in syphilis positivity from 0.9% (1/112) in 2011 to 6.4% (19/299) in 2018 ($p_{\text{trend}} = 0.004$) in MSMW, although the number of infections remained low in this group (figure 2 and online supplemental table S5). HIV positivity did not change in either group between 2011 and 2018 (figure 2 and online supplemental tables S4, S5).

After adjusting for other potential confounding factors, MSMO had higher odds of testing positive for gonorrhoea at any anatomical site (aOR 1.36, 95% CI 1.13 to 1.64), chlamydia at any anatomical site (aOR 1.39; 95% CI 1.16

Table 3 Positivity for gonorrhoea, Chlamydia, syphilis and HIV among men who have sex with men attending a sexual health centre in Melbourne, 2011–2018, stratified by men who have sex with men only (MSMO) and men who have sex with men and women (MSMW)

HIV/STI	Men who have sex with men only (N=12795)			Men who have sex with men and women (N=1979)			P value
	No of men tested, N	No of positive, n	Positivity, % (95% CI)	No of men tested, N	No of positive, n	Positivity, % (95% CI)	
Gonorrhoea							
Oropharyngeal	11 810	682	5.8 (5.4 to 6.2)	1711	68	4.0 (3.1 to 5.0)	0.002
Urethral*	11 699	390	3.3 (3.0 to 3.7)	1885	52	2.8 (2.1 to 3.6)	0.192
Anorectal	11 080	745	6.7 (6.3 to 7.2)	1404	58	4.1 (3.2 to 5.3)	<0.001
Any anatomical site†	11 951	1321	11.1 (10.5 to 11.6)	1829	140	7.7 (6.5 to 9.0)	<0.001
Chlamydia							
Urethral	11 699	410	3.5 (3.2 to 3.9)	1885	73	3.9 (3.0 to 4.8)	0.423
Anorectal	11 043	989	9.0 (8.4 to 9.5)	1404	83	5.9 (4.7 to 7.3)	<0.001
Any anatomical site†	11 891	1272	10.7 (10.1 to 11.3)	1907	141	7.4 (6.3 to 8.7)	<0.001
Syphilis	11 317	797	7.0 (6.6 to 7.5)	1773	81	4.6 (3.6 to 5.6)	<0.001
HIV							
New diagnosis‡	11 102	262	2.4 (2.1 to 2.7)	1774	10	0.6 (0.3 to 1.0)	<0.001
Previous diagnosis	12 795	316	2.5 (2.2 to 2.8)	1979	7	0.4 (0.1 to 0.7)	<0.001

*Before 2015, urethral gonorrhoea testing was only performed among symptomatic MSM with urethral symptoms or self-reported as contact of infection. The number of people tested corresponds to the number of people tested for urethral chlamydia.

†Any anatomical site for gonorrhoea includes urethral, anorectal and oropharyngeal; while any site for chlamydia includes urethral and anorectal.

‡Only including HIV cases diagnosed on the day of the first visit.

CI, confidence interval ; STI, sexually transmitted infection.

Figure 2 Positivity for syphilis and HIV between 2011 and 2018, stratified by men who have sex with men only (MSMO) and men who have sex with men and women (MSMW).

Table 4 Association between sexual practice and gonorrhoea, Chlamydia, syphilis and HIV positivity among men who have sex with men only (MSMO) and men who have sex with men and women (MSMW) attending a sexual health centre in Melbourne, 2011–2018

HIV/STI	OR (95% CI)	P value	aOR (95% CI)*	P value
Gonorrhoea (any anatomical site)†				
MSMO	1.50 (1.25 to 1.80)	<0.001	1.36 (1.13 to 1.64)	0.001
MSMW	1		1	ref
Chlamydia (any anatomical site)†				
MSMO	1.50 (1.25 to 1.80)	<0.001	1.39 (1.16 to 1.67)	<0.001
MSMW	1		1	ref
Syphilis				
MSMO	1.58 (1.25 to 2.00)	<0.001	1.74 (1.37 to 2.22)	<0.001
MSMW	1		1	ref
HIV (new diagnosis)‡				
MSMO	4.26 (2.26 to 8.03)	<0.001	4.60 (2.43 to 8.70)	<0.001
MSMW	1		1	ref
HIV (previous diagnosis)				
MSMO	7.13 (3.37 to 15.11)	<0.001	9.08 (4.26 to 19.37)	<0.001
MSMW	1	ref	1	ref

*Adjusted by age, country of birth, sex overseas, injected drug use in the past 12 months, regular partner, number of male sexual partners in the past 12 months, condom use with casual male partners in the past 12 months and year of visit.

†Any anatomical site for gonorrhoea includes urethral, anorectal and oropharyngeal; while any site for chlamydia includes urethral and anorectal.

‡Only including HIV cases diagnosed on the day of the first visit.

STI, sexually transmitted infection; aOR, adjusted odds ratio; OR, odds ratio; MSMO, men who have sex with men only; MSMW, men who have sex with men and women; CI, confidence interval.

to 1.67), syphilis (aOR 1.74, 95% CI 1.37 to 2.22) and HIV (aOR 4.60, 95% CI 2.43 to 8.70) compared with MSMW (table 4). Additionally, MSMO had higher odds of testing positive for gonorrhoea at any anatomical site (aOR 1.58; 95% CI 1.15 to 2.18), chlamydia at any anatomical site (aOR 1.47; 95% CI 1.09 to 1.99), syphilis (aOR 2.11, 95% CI 1.36 to 3.27) and HIV (aOR 5.49, 95% CI 1.74 to 17.30) compared with MSMW-W in the adjusted analysis; however, there was no significant difference in HIV/STI positivity between MSMW-M and MSMW-W nor MSMW-M and MSMO (table 5).

DISCUSSION

In this study of 12795 MSM only (MSMO) and 1979 MSMW in Melbourne, Australia, we found significant changes in sexual practices among MSMW between 2011 and 2018, a period in which rates of STIs had been increasing in both MSM and the heterosexual population. Specifically, we found that condom use with casual female partners remained low during the study period and that condomless anal sex with casual male partners had increased in MSMW, and these results echo the rise in syphilis positivity over the same period. While some sexual risk practices and STI positivity were generally lower in MSMW than in MSMO, their rising rates could indicate more transmission from MSMW to their female

partners than had occurred previously. Clarifying this issue and what factors are causing the rising rates of STIs is likely to contribute significantly to the design of STI control programmes.

Several studies have assessed HIV and sexual practices in MSMW, but their results are conflicting. A meta-analysis published in 2014 concluded that MSMW had a lower HIV prevalence (16.9%) compared with MSMO (33.3%) but the prevalence is higher than men who have sex with women only (3.5%), and this is consistent with our findings.¹⁹ The authors also found that MSMW (15.9%) were less likely to engage in condomless RAS compared with MSMO (35.0%).¹⁹ Other factors such as drug and alcohol use, other risky practices (eg, group sex, paying for sex), frequent HIV testing and the use of biomedical interventions (eg, pre-exposure prophylaxis and post-exposure prophylaxis) are also associated with HIV acquisition,^{20–22} and very limited studies examining these factors between MSMO and MSMW. However, some other studies conducted in different settings, such as China, India and the USA, have shown the risk of HIV among MSMW is similar or even at a higher risk compared with MSMO.^{13 23–25}

Our data suggest that MSMO have a higher positivity for syphilis, chlamydia and gonorrhoea than MSMW; however, the comparison of STI positivity between MSMO

Table 5 Association between sexual practice and gonorrhoea, Chlamydia, syphilis and HIV positivity among men who have sex with men only (MSMO) and men who have sex with men and women (MSMW) attending a sexual health centre in Melbourne, 2011–2018

HIV/STI	OR (95% CI)	P value	aOR (95% CI)*	P value
Gonorrhoea (any anatomical site)†				
MSMO	1.96 (1.45 to 2.67)	<0.001	1.58 (1.15 to 2.18)	0.005
MSMW-M	1.91 (1.31 to 2.78)	0.001	1.41 (0.94 to 2.12)	0.093
MSMW-W	1	ref	1	ref
Chlamydia (any anatomical site)†				
MSMO	1.87 (1.39 to 2.50)	<0.001	1.47 (1.09 to 1.99)	0.013
MSMW-M	1.63 (1.12 to 2.38)	0.010	1.19 (0.80 to 1.77)	0.394
MSMW-W	1	ref	1	ref
Syphilis				
MSMO	2.18 (1.45 to 3.27)	<0.001	2.11 (1.36 to 3.27)	0.001
MSMW-M	1.48 (0.88 to 2.49)	0.142	1.08 (0.61 to 1.93)	0.791
MSMW-W	1	ref	1	ref
HIV (new diagnosis)‡				
MSMO	6.01 (1.92 to 18.80)	0.002	5.49 (1.74 to 17.30)	0.004
MSMW-M	2.38 (0.61 to 9.24)	0.210	1.81 (0.43 to 7.63)	0.422
MSMW-W	1	ref	1	ref
HIV (previous diagnosis)				
MSMO	21.22 (2.98 to 151.30)	0.002	17.91 (2.50 to 128.43)	0.004
MSMW-M	6.30 (0.76 to 52.45)	0.089	4.10 (0.47 to 35.42)	0.200
MSMW-W	1	ref	1	ref

*Adjusted by age, country of birth, sex overseas, injected drug use in the past 12 months, regular partner, number of male sexual partners in the past 12 months, condom use with casual male partners in the past 12 months and year of visit.

†Any anatomical site for gonorrhoea includes urethral, anorectal and oropharyngeal, while any site for chlamydia includes urethral and anorectal.

‡Only including HIV cases diagnosed on the day of the first visit.

aOR, adjusted odds ratio; CI, confidence interval; MSMO, men who have sex with men only; MSMW-M, men who have sex with men and women and had more male than female sexual partner in the past 12 months; MSMW-W, men who have sex with men and women and had more female than male sexual partner in the past 12 months; OR, odds ratio; STI, sexually transmitted infection.

and MSMW varies across published studies. For example, Davis *et al* conducted a study among Chinese MSM showing no differences between MSMW and MSMO in the lifetime prevalence of syphilis, gonorrhoea, chlamydia and genital warts but a higher prevalence of genital herpes in MSMW (6.6%) compared with MSMO (0.4%).²⁵ In the USA, MSMW had a 57% increased odds of having an STI compared with heterosexuals, but no differences were found between MSMO and heterosexuals or MSMO and MSMW.²⁴ Similarly, an Indian study has shown that there were no differences in any STI positivity (ie, either syphilis, gonorrhoea or chlamydia) between MSMO and MSMW.²³ Furthermore, MSMW are also less likely to ever have an HIV/STI test compared with MSMO,^{23 26 27} which is consistent with an Australian qualitative study suggesting MSMW have a poor sexual health knowledge compared with MSMO.²⁸ Furthermore, the rise in antimicrobial resistance in STI,²⁹ particularly gonorrhoea, is of particular concern. Transmission of antimicrobial

resistance across risk populations (eg, between MSMW and female) may have occurred.^{30 31}

Our data also show that MSMW are less likely to have RAS with male partners and less likely to have condomless RAS. These findings may explain the reasons why MSMW have a lower positivity for anorectal gonorrhoea and chlamydia than MSMO in our study. Importantly, we did not find any significant differences in urethral gonorrhoea and chlamydia between MSMO and MSMW despite differences in anal sex and condom use. This may be because some urethral chlamydial infections in MSMW are acquired from women and urethral gonorrhoea is commonly acquired from no anal sexual activity such as oral sex.^{32 33} The disparities of HIV and syphilis positivity between MSMO and MSMW were more pronounced compared with gonorrhoea and chlamydia, and there was a striking increase in syphilis positivity in both groups.

The differential sexual practices and HIV/STI positivity among MSMW^{34 35} may also be due to other health issues

such as mental health and substance use, that also place them as an especially vulnerable population.^{36 37} Bisexual men may face unique stressors, such as specific prejudices against them, pressures or negative attitudes from both heterosexual and lesbian, gay, bisexual, and transgender (LGBT) individuals, which can have a negative impact on their health.^{37 38} These factors may also render bisexual men less likely to discuss their sexual health and disclose their sexual orientation with their general practitioners¹⁴ and, hence not receive the appropriate sexual healthcare and management. An LGBT-friendly and non-judgemental approach is essential to build up the trust between clinicians and patients. Furthermore, the current health promotion and prevention programmes mainly targeting the MSM population as a whole. It is possible that some MSMW may not be engaged in the LGBT community and hence they may not be targeted with the current programmes. Future public health campaigns and prevention programmes should also tailor for MSMW via different channels.

Our study, including a high number of MSM over an 8-year period, provides relevant and updated information to understand changes in sexual practices and STI epidemiology occurring in Australia and across the world. We also provide detailed information regarding condom use, differencing between both casual or regular partners and RAS or IAS. Nevertheless, there are several limitations in the study that must be considered. First, data were self-reported by MSHC clients which may influence the results regarding sexual practices, including the number and gender of the sexual partners due to social desirability or recall bias. However, the use of a CASI to collect sexual history at the clinic has shown to improve the accuracy and reduce biased information. Besides that, data on sexual orientation were not collected and the categorisation of MSMO and MSMW was based on self-reported sexual practice rather than sexual identity. This means that MSMW in this study might not define themselves as bisexual men. Second, this study was conducted in one urban major sexual health clinic. It is possible that MSM attending a sexual health clinic are more likely to be at increased sexual risk and hence our findings may not be generalisable to the whole MSM population in Australia or in other settings. Third, other sexual practices that may increase the risk of HIV/STI (eg, chemsex, group sex, rimming or saliva use as a lubricant)^{39–42} were not routinely collected in the clinic and hence not included in the study. Fourth, we only included individuals who attended the clinic for the first time. Sexual practices might change with age and further longitudinal cohort studies examining the changes in sexual practices among individuals would be required. Lastly, we were unable to examine the temporal trends for gonorrhoea and chlamydia over the 8-year period because we changed the diagnostic test for gonorrhoea and chlamydia at our clinic in 2015.

CONCLUSION

In our study population, MSMW were a heterogeneous group in which sexual practices and STI positivity varied between MSMW with mainly female partners and MSMW with mainly male partners. Compared with MSMO, MSMW were less likely to engage in condomless sex and had a lower HIV/STI positivity. From 2011 to 2018, changes in the sexual practices in MSM have affected both MSMW and MSMO leading to an increased risk of STI in both subgroups. Further studies also including heterosexual men and women are needed to better understand the recent changes in the STI epidemiology.

Author affiliations

¹Melbourne Sexual Health Centre, Alfred Health, Melbourne, Victoria, Australia

²Preventive Medicine and Public Health Training Unit, Parc de Salut Mar - Pompeu Fabra University - Agència de Salut Pública de Barcelona (PSMar-UPF-ASPB), Barcelona, Spain

³Central Clinical School, Monash University, Melbourne, Victoria, Australia

⁴Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia

Twitter Mario Martín-Sánchez @mmsanchz

Acknowledgements The authors would like to acknowledge Afrizal Afrizal for his assistance with data extraction at the Melbourne Sexual Health Centre.

Contributors EPFC and CF planned, conceived and designed the study. RC conducted some preliminary analyses. MM-S performed the literature review, conducted the statistical analysis and wrote the first draft of the manuscript. EPFC provided statistical advice and oversaw the study. RC, CF, JSH, CB, JO, MYC and EPFC assisted with data interpretation. MM-S, RC, CF, JSH, CB, JO, MYC and EPFC critically revised it for important intellectual content and approved the final version of the manuscript.

Funding EPFC is supported by an Australian National Health and Medical Research Council (NHMRC) Emerging Leadership Investigator Grant (GNT1172873). CKF is supported by an Australian NHMRC Leadership Investigator Grants (GNT1172900).

Disclaimer The funder had no role in the study design, collection of data, writing, or decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the Alfred Hospital Ethics Committee, Melbourne, Australia (project number 83/18).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Mario Martín-Sánchez <http://orcid.org/0000-0002-2221-5331>

Jason Ong <http://orcid.org/0000-0001-5784-7403>

Eric P F Chow <http://orcid.org/0000-0003-1766-0657>

REFERENCES

- World Health Organization. *Report on global sexually transmitted infection surveillance, 2018*. Geneva: World Health Organization, 2018.
- Kirkcaldy RD, Weston E, Segurado AC, *et al*. Epidemiology of gonorrhoea: a global perspective. *Sex Health* 2019;16:401.
- Kirby Institute. Annual surveillance report 2018 HIV, viral hepatitis and sexually transmissible infections in Australia, 2018. Available: https://kirby.unsw.edu.au/sites/default/files/kirby/report/KI_Annual-Surveillance-Report-2018_0.pdf
- Jasek E, Chow EP, Ong JJ, *et al*. Sexually transmitted infections in Melbourne, Australia from 1918 to 2016: nearly a century of data. *Commun Dis Intell Q Rep* 2017;41:E212–22.
- Needleman R, Chow EPF, Towns JM, *et al*. Access to sexual health services after the rapid roll out of the launch of pre-exposure prophylaxis for HIV in Melbourne, Australia: a retrospective cross-sectional analysis. *Sex Health* 2018;15:528–32.
- Ryan KE, Mak A, Stooze M, *et al*. Protocol for an HIV pre-exposure prophylaxis (PrEP) population level intervention study in Victoria Australia: the PrEPX study. *Front Public Health* 2018;6:1–11.
- Chow EPF, Medland NA, Denham I, *et al*. Decline in new HIV diagnoses among MSM in Melbourne. *Lancet HIV* 2018;5:e479–81.
- Zhang Kudon H, Mulatu MS, Song W, *et al*. Trends in Condomless sex among MSM who participated in CDC-Funded HIV Risk-Reduction interventions in the United States, 2012–2017. *J Public Health Manag Pract* 2020:01143.
- Hess KL, Crepez N, Rose C, *et al*. Trends in sexual behavior among men who have sex with men (MSM) in high-income countries, 1990–2013: a systematic review. *AIDS Behav* 2017;21:2811–34.
- Oldenburg CE, Nunn AS, Montgomery M, *et al*. Behavioral changes following uptake of HIV pre-exposure prophylaxis among men who have sex with men in a clinical setting. *AIDS Behav* 2018;22:1075–9.
- Chow EPF, Grulich AE, Fairley CK. Epidemiology and prevention of sexually transmitted infections in men who have sex with men at risk of HIV. *Lancet HIV* 2019;6:e396–405.
- Holt M, Lea T, Mao L, *et al*. Community-Level changes in condom use and uptake of HIV pre-exposure prophylaxis by gay and bisexual men in Melbourne and Sydney, Australia: results of repeated behavioural surveillance in 2013–17. *Lancet HIV* 2018;5:e448–56.
- Jeffries WL. Beyond the bisexual bridge: sexual health among U.S. men who have sex with men and women. *Am J Prev Med* 2014;47:320–9.
- Friedman MR, Stall R, Silvestre AJ, *et al*. Stuck in the middle: longitudinal HIV-related health disparities among men who have sex with men and women. *J Acquir Immune Defic Syndr* 2014;66:213–20.
- Cornelisse VJ, Chow EPF, Huffam S, *et al*. Increased detection of pharyngeal and rectal gonorrhea in men who have sex with men after transition from culture to nucleic acid amplification testing. *Sex Transm Dis* 2017;44:114–7.
- Templeton D, Read P, Varma R, *et al*. Australian sexually transmitted infection & HIV testing guidelines 2014 for asymptomatic men who have sex with men, 2014: 217–29. http://stipu.nsw.gov.au/wp-content/uploads/STIGMA_Testing_Guidelines_Final_v5.pdf
- Ong JJ, Fethers K, Howden BP, *et al*. Asymptomatic and symptomatic urethral gonorrhoea in men who have sex with men attending a sexual health service. *Clin Microbiol Infect* 2017;23:555–9.
- Chow EPF, Tomnay J, Fehler G, *et al*. Substantial increases in Chlamydia and gonorrhea positivity unexplained by changes in individual-level sexual behaviors among men who have sex with men in an Australian sexual health service from 2007 to 2013. *Sex Transm Dis* 2015;42:81–7.
- Friedman MR, Wei C, Klem ML, Lou KM, *et al*. HIV infection and sexual risk among men who have sex with men and women (MSMW): a systematic review and meta-analysis. *PLoS One* 2014;9:e87139.
- Chou R, Evans C, Hoveman A, *et al*. Preexposure prophylaxis for the prevention of HIV infection: evidence report and systematic review for the US preventive services Task force. *JAMA* 2019;321:2214–30.
- Woelf SE, Maisto SA. Alcohol use and risk of HIV infection among men who have sex with men. *AIDS Behav* 2009;13:757–82.
- Hutchinson AB, Farnham PG, Sansom SL, *et al*. Cost-Effectiveness of frequent HIV testing of high-risk populations in the United States. *J Acquir Immune Defic Syndr* 2016;71:323–30.
- Ramakrishnan L, Ramanathan S, Chakrapani V, *et al*. Comparison of sexual risk, HIV/STI prevalence and intervention exposure among men who have sex with men and women (MSMW) and men who have sex with men only (MSMO) in India: implications for HIV prevention. *AIDS Behav* 2015;19:2255–69.
- Dyer TV, Khan MR, Sandoval M, *et al*. Drug use and sexual HIV transmission risk among men who have sex with men and women (MSMW), men who have sex with men only (MSMO), and men who have sex with women only (MSWO) and the female partners of MSMW and MSWO: a network perspective. *AIDS Behav* 2017;21:3590–8.
- Davis A, Best J, Luo J, *et al*. Differences in risk behaviours, HIV/STI testing and HIV/STI prevalence between men who have sex with men and men who have sex with both men and women in China. *Int J STD AIDS* 2016;27:840–9.
- Jeffries WL. HIV testing among bisexual men in the United States. *AIDS Educ Prev* 2010;22:356–70.
- Mirandola M, Gios L, Sherriff N, *et al*. Socio-Demographic characteristics, sexual and Test-Seeking behaviours amongst men who have sex with both men and women: results from a Bio-behavioural survey in 13 European cities. *AIDS Behav* 2017;21:3013–25.
- Smardon T, Vaisey A, Chow EPF, *et al*. Sexual practices and healthcare use of men who have sex with men only and men who have sex with men and women – a qualitative comparison. *Australas Sex Heal Conf Jt with 2019 Australas HIV&AIDS Conf* 2019;2019.
- Williamson DA, Fairley CK, Howden BP, *et al*. Trends and Risk Factors for Antimicrobial-Resistant *Neisseria gonorrhoeae*, Melbourne, Australia, 2017 to 2018. *Antimicrob Agents Chemother* 2019;63:e01221–19.
- Williamson DA, Chow EPF, Gorrie CL, *et al*. Bridging of *Neisseria gonorrhoeae* lineages across sexual networks in the HIV pre-exposure prophylaxis era. *Nat Commun* 2019;10:1–10.
- Kwong JC, Chow EPF, Stevens K, *et al*. Whole-Genome sequencing reveals transmission of gonococcal antibiotic resistance among men who have sex with men: an observational study. *Sex Transm Infect* 2018;94:151–7.
- Barbee LA, Khosropour CM, Dombrowski JC, *et al*. An estimate of the proportion of symptomatic gonococcal, chlamydial and non-gonococcal non-chlamydial urethritis attributable to oral sex among men who have sex with men: a case-control study. *Sex Transm Infect* 2016;92:155–60.
- Nash JL, Hocking JS, Read TRH, *et al*. Contribution of sexual practices (other than anal sex) to bacterial sexually transmitted infection transmission in men who have sex with men: a cross-sectional analysis using electronic health records. *Sex Transm Infect* 2014;90:55–7.
- Chow EPF, Wilson DP, Zhang L. Estimating HIV incidence among female partners of bisexual men in China. *Int J Infect Dis* 2012;16:e312–20.
- Ong JJ, Landika A, Fairley CK, *et al*. Characteristics, sexual practices and sexually transmissible infections diagnoses of men who have sex with men and use non-occupational HIV post-exposure prophylaxis in Victoria, Australia. *Sex Health* 2016;13:555–9.
- Gonzales G, Henning-Smith C, Henning C. Health disparities by sexual orientation: results and implications from the behavioral risk factor surveillance system. *J Community Health* 2017;42:1163–72.
- Feinstein BA, Health GM, Bisexuality DC. Minority stress, and health. *Curr Sex Heal Rep* 2018;9:42–9.
- Marti-Pastor M, Ferrer M, Alonso J, *et al*. Association of enacted stigma with depressive symptoms among gay and bisexual men who have sex with men: Baltimore, 2011 and 2014. *LGBT Health* 2020;7:47–59.
- Chow EP, Fairley CK. The role of saliva in gonorrhoea and Chlamydia transmission to extragenital sites among men who have sex with men: new insights into transmission. *J Int AIDS Soc* 2019;22 Suppl 6:e25354.
- Maxwell S, Shahmanesh M, Gafos M. Chemsex behaviours among men who have sex with men: a systematic review of the literature. *Int J Drug Policy* 2019;63:74–89.
- van den Boom W, Davidovich U, Heuker J, *et al*. Is group sex a higher-risk setting for HIV and other sexually transmitted infections compared with Dyadic sex among men who have sex with men? *Sex Transm Dis* 2016;43:99–104.
- Phillips TR, Fairley CK, Bradshaw CS, *et al*. Group sex among men who have sex with men in the era of PrEP: a cross-sectional study. *J Acquir Immune Defic Syndr* 2020. doi:10.1097/QAI.0000000000002550. [Epub ahead of print: 19 Oct 2020].