

# BMJ Open Prevalence of limited health literacy and its associated factors in patients with type 2 diabetes mellitus in Perak, Malaysia: a cross-sectional study

Adina Abdullah <sup>1</sup>, Chirk Jenn Ng <sup>1</sup>, Su May Liew,<sup>1</sup> Subashini Ambigapathy,<sup>2</sup> Paranthaman V,<sup>3</sup> Karuthan Chinna<sup>4</sup>

**To cite:** Abdullah A, Ng CJ, Liew SM, *et al*. Prevalence of limited health literacy and its associated factors in patients with type 2 diabetes mellitus in Perak, Malaysia: a cross-sectional study. *BMJ Open* 2020;**10**:e039864. doi:10.1136/bmjopen-2020-039864

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-039864>).

Received 29 April 2020  
Revised 02 September 2020  
Accepted 13 October 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Primary Care Medicine, University Malaya Primary Care Research Group (UMPCRG), Faculty of Medicine, University Malaya, Kuala Lumpur, Malaysia

<sup>2</sup>Health Clinic Buntong, Ministry of Health Malaysia, Ipoh, Malaysia

<sup>3</sup>Health Clinic Greentown, Ministry of Health Malaysia, Ipoh, Malaysia

<sup>4</sup>School of Medicine, Faculty of Health and Medical Sciences, Taylor's University, Kuala Lumpur, Malaysia

**Correspondence to**  
Dr Adina Abdullah;  
[adina@ummc.edu.my](mailto:adina@ummc.edu.my)

## ABSTRACT

**Objective** Limited health literacy in patients with type 2 diabetes mellitus (T2DM) led to poorer diabetes knowledge, less medication adherence and increased healthcare cost. The purpose of this paper was to report the prevalence of limited health literacy in patients with T2DM and to identify factors that are associated with it. **Design** A cross-sectional study was conducted from January to March 2018; data on patients' sociodemographic characteristics, diabetes knowledge, perceived social support and health literacy level were collected. Health literacy level was measured using the European Health Literacy Survey Questionnaire (HLS-EU-Q47). **Setting** Patients were recruited from four primary care clinics in Perak, Malaysia. **Participants** Adult patients diagnosed with T2DM who attended the study clinics during the study period. **Primary outcome variable** Patients with HLS-EU-Q47 General Index of  $\leq 33$  points were classified as having limited health literacy.

**Results** The prevalence of limited health literacy was 65.3% (n=279). In bivariate analysis, patients' ethnicity (p=0.04), highest education level (p<0.001), monthly income (p=0.003), having health insurance (p=0.007), English language fluency (p<0.001), Malay language fluency (p=0.021), attending diabetes education sessions (p<0.001), perceived social support (p<0.001) and diabetes knowledge (p=0.019) were factors associated with limited health literacy. In logistic regression, not being fluent in English was associated with limited health literacy (OR=2.36, 95% CI 1.30 to 4.30) whereas having high perceived social support (OR=0.52, 95% CI 0.40 to 0.69) and having attended diabetes education sessions (OR=0.42, 95% CI 0.27 to 0.68) were associated with adequate health literacy.

**Conclusion** The prevalence of limited health literacy is high among patients with T2DM in Perak, Malaysia. Strategies to improve health literacy in these patients must consider the influences of English fluency, attendance at diabetes education sessions and social support, and may need to adopt a universal approach to addressing limited health literacy.

## BACKGROUND

Health literacy is defined as the degree to which individuals can access, understand,

## Strengths and limitations of this study

- This is the first study to measure health literacy using the European Health Literacy Survey Questionnaire in patients with type 2 diabetes mellitus in Malaysia.
- The study questionnaire was made available in three languages: Bahasa Malaysia, English and Mandarin to ensure inclusion of patients from various ethnicities.
- The study was conducted in primary care clinics from urban and rural settings.
- This study was conducted in only one state in Malaysia which limits its generalisability.
- The cross-sectional design of the study means that we could not determine causality and temporal associations between patients' health literacy levels and associated factors.

appraise and apply health information to make informed health decisions. It is closely related to literacy and entails the individual's knowledge, motivation and competencies.<sup>1</sup> It encompasses the abilities and skills required by patients to deal with the complexity and demands of the health condition or systems where patients are increasingly expected to be partners in care. Recent research showed that health literacy skills are important for patients with type 2 diabetes mellitus (T2DM). When patients are diagnosed with T2DM, they needed to understand new terminologies and learn self-care concepts quickly in order to make healthcare decisions related to treatment and to perform daily self-management activities for optimal diabetes control.<sup>2</sup> Studies showed that patients with T2DM and limited health literacy struggled to understand print instructions, health advisories or warnings.<sup>3</sup> They have poorer diabetes knowledge,<sup>4,5</sup> are less adherent to medication<sup>6</sup> and spend more money on their health.<sup>7</sup> Furthermore, limited health literacy

led to patients having poorer patient–doctor communication and participating less in shared decision-making.<sup>8</sup> Studies evaluating health literacy-targeted patient education interventions in patients with T2DM showed improvement of glycaemic control.<sup>9 10</sup>

Previous studies measured health literacy levels in patients with T2DM and showed the prevalence of limited health literacy to be high; it ranges from 60% to 85%.<sup>11</sup> Many of these studies were conducted in high-income countries in North America and Europe. However, nearly 79% of patients with T2DM live in low-income to middle-income countries, like Malaysia.<sup>12</sup> Diabetes is a major health problem in Malaysia. In 2015, the prevalence of diabetes in the entire country of Malaysia was 17.5%, which was a 15% increase from the prevalence in 2011.<sup>13</sup> The prevalence of diabetes risk factors is also a concern; the prevalence of abdominal obesity ranges from 55.6% to 57.4%<sup>14 15</sup> with the prevalence of gestational diabetes reaching 8.66% in 2010.<sup>16</sup> High prevalence of these risk factors means that diabetes will continue to be a major health problem in Malaysia for years to come. In order to face this problem, the Ministry of Health, Malaysia, had come up with new policies, strategic plans and clinical practice guidelines (CPGs) on management of patients with T2DM. These guidelines include strategies on screening, prevention and use of new efficacious drugs. Despite this effort, up to 30% of Malaysian patients with T2DM did not achieve the target glycaemic control of haemoglobin A1c (HbA1c) >6.5% as defined by the Malaysian CPG on management of T2DM;<sup>17</sup> this led to a high proportion of patients having microvascular (75%) and macrovascular (29%) diabetes complications.<sup>18 19</sup>

Based on previous studies on health literacy and T2DM, health literacy may be a new target for intervention for patients with T2DM in Malaysia.<sup>20–22</sup> Therefore, this study aimed to measure the prevalence of limited health literacy in patients with T2DM and to identify factors that are associated with limited health literacy. Findings in this study could help to identify new targets for interventions in order to improve management of patients with T2DM in Malaysia.

## METHODS

### Study design and settings

This cross-sectional study was conducted at four primary care clinics in Perak, Malaysia. Perak is the second largest state in Peninsular Malaysia. The state of Perak was chosen because it has a diabetes prevalence similar to Malaysia's prevalence (11.9%, 95% CI 9.9% to 14.3%) and a population with the highest prevalence of adequate health literacy in Malaysia (19.5%, 95% CI 13.2% to 27.9%).<sup>13</sup> The latter is important to ensure the sample is from a population with a range of health literacy levels. Four primary care clinics were purposively identified from 17 clinics in Kinta District: Klinik Kesihatan Jelapang, Klinik Kesihatan Buntong, Klinik Kesihatan Manjoi and Klinik

Kesihatan Menglembu. These clinics were selected to ensure ethnicity and socioeconomic representations.


### Participant selection

Eligible participants were adult men or women with a known diagnosis of T2DM. The diagnosis was confirmed by participants' medical records. The exclusion criteria were those who were unable to communicate in English, Bahasa Malaysia or Mandarin; had dementia as determined by an Elderly Cognitive Assessment Questionnaire Score of 5 points or less; and who were determined to be too ill based on vital sign assessments or interactions with the research assistants administering the questionnaire.

### Data collection

All participants who fulfilled the inclusion and exclusion criteria and attended the study clinics for diabetes follow-up from January to March 2018 were recruited. This study used a self-designed questionnaire to capture participants' sociodemographic data: age, gender, ethnicity, marital status, highest educational attainment, language fluency, employment status, monthly income and health insurance coverage. Language fluency in both English and Malaysian national language, Bahasa Malaysia, was assessed using a 4-point Likert Scale. Participants self-reported their ability to speak, read and write in the said language. Higher scores on the scale indicate better language fluency. In addition, health literacy level, diabetes knowledge and perceived social support were measured using validated tools. Data on disease outcomes (diabetes control and blood pressure control) were gathered from participants' medical records. This questionnaire was made available in three languages: English, Bahasa Malaysia and Mandarin and participants chose the language they were most familiar with. The questionnaire was administered with the help of trained research assistants and supervised by the principal investigator.

A pretest of the questionnaire was conducted. Nine face-to-face interviews were conducted with five Malay, two Chinese and two Indian participants. During this process, the participants and the researcher answered the questionnaire together. The participants were asked to voice their thoughts on the questionnaire, particularly with regards to the clarity of the questions and the ease of answering them. The comments from these participants were used to improve the readability and clarity of the questionnaire. Minimal changes were made to the questions included in the questionnaire, including the removal of a question regarding self-rated health status, which the pretest participants had difficulty understanding. In the actual study only one<sup>1</sup> participant did not complete all the measurements for the European Health Literacy Survey Questionnaire (HLS-EU-Q47). The participants took on average 15 mins to complete the questionnaire with assistance from the trained research assistants.



**Figure 1** The conceptual framework for the study measuring factors associated with limited health literacy.

### Sample size calculation

The sample size was calculated based on the formula by Peduzzi *et al.*<sup>23</sup> Therefore, the required minimal sample size for logistic regression was 331. This number was inflated 20% to account for the non-responder rate giving a final sample size of 397.<sup>24</sup>

### Conceptual model of the study

In order to identify possible associated factors of health literacy, we reviewed previous literature and conducted an expert review. A conceptual model of factors that had been found to be associated with health literacy level is presented in figure 1. This study focused on measuring the personal factors that may be associated with limited health literacy in patients with T2DM.

### Measures

#### Health literacy level

Health literacy level was determined using the General Health Literacy Index of HLS-EU-Q47. It contains 47 items that are phrased as direct questions, and the participants have to indicate on a 4-point Likert Scale how easy it is to find, understand, judge and apply health information. The original English questionnaire was validated in English with a good internal consistency of 0.97.<sup>25</sup> The Malay and Mandarin translations were validated in a five Asian country survey with both having a Cronbach's  $\alpha$  of 0.96.<sup>26</sup> The General Health Literacy Index was calculated using the formula:  $Index = (mean - 1) * (50/3)$ . The General Health Literacy Index has a minimum score of 0 and a maximum of 50. Participants were classified as having limited health literacy if their General Health Literacy Index was  $\leq 33$  points.<sup>27</sup>

#### Diabetes knowledge

The participants' diabetes knowledge was measured using the Michigan Diabetes Knowledge Test (MDKT).<sup>28</sup> The MDKT consists of 14 multiple-choice questions, each with one correct answer. The knowledge score is determined by giving 1 point for each correct answer and 0 for a wrong answer or no response. The total knowledge scores range from 0 to 14 with higher scores indicating higher levels of diabetes knowledge. The original version was in English (Cronbach's  $\alpha=0.70$ ); the Bahasa Malaysian

translation was also found to be valid and reliable (Cronbach's  $\alpha=0.702$ ).<sup>29</sup>

### Perceived social support

Perceived social support was measured using the Multi-dimensional Scale of Perceived Social Support (MSPSS), which measured perceptions of support from three sources: family, friends and significant others.<sup>30</sup> The MSPSS is self-administered and contains 12 items, with 4 items for each subscale. Higher scores reflect higher perceived support. The original English MSPSS has excellent internal consistency, with a Cronbach's  $\alpha$  of 0.91 for the total scale. Cronbach's  $\alpha$  for the translated Malay and validated versions was 0.89.<sup>31</sup>

### Disease outcomes

Diabetes-related measurements included the duration of diabetes diagnosis, attendance at diabetes education sessions as well as glycaemic and blood pressure controls. Glycaemic control was assessed using the mean of the three most recent HbA1c values over the last 1 year taken from the patients' medical records. The blood pressure reading used to determine blood pressure control was also the mean of the three most recent blood pressure readings in the patients' medical records over the last 1 year.

### Data analysis

All statistical analyses were conducted using SPSS V.21.0 (IBM SPSS Statistics for Windows, Armonk, New York, USA). First, we reported the prevalence of limited health literacy and the descriptive statistics analysis of the sample and measured variables. Then we computed the bivariate relationships between measured variables and health literacy groups (limited or adequate health literacy) using the  $\chi^2$  test for categorical data and the independent t-test or one-way analyses of variance for continuous parametrical data. Then, a binary logistic regression was performed to identify the factors associated with limited health literacy. Variables that had a p value  $<0.25$  in the univariate analysis were included in the logistic regression. The resulting ORs were reported with 95% CIs and the p values. The results were checked for influential cases and multicollinearity. Regression diagnostics showed no influential cases. Furthermore, there was no multicollinearity among the variables. The statistical significance level for the variables inputted in the final model was set at  $p<0.05$ .

### Patient and public involvement

There were no funds or time allocated for patient and public involvement so we were unable to involve patients. We have invited patients to help us develop our dissemination strategy.

### RESULTS

A total of 428 from 450 patients with T2DM approached, agreed to be involved in the study (response rate=95.1%). The mean General Health Literacy Index Score was 31.86

**Table 1** Participants' characteristics

Characteristics		N (%)	Mean±SD
Age (years)			58.1±10.6
Gender	Female	258 (60.3)	
Ethnicity	Malay	185 (43.8)	
	Chinese	96 (22.7)	
	Indian	141 (33.4)	
Marital status	Single/divorced/separated/widowed	99 (23.2)	
	Married	328 (76.8)	
Highest educational attainment	Primary school	133 (31.2)	
	Junior high school	126 (29.6)	
	High school	113 (26.5)	
	University/masters/PhD	54 (12.7)	
Employment status	Yes	152 (35.9)	
	No, never worked	123 (29.1)	
	No, stop working or retired	148 (35.0)	
Monthly income	Less than minimum wage	243 (56.8)	
Health insurance ownership	No	330 (77.6)	
English fluency	Not fluent	275 (64.3)	
Bahasa Malaysia fluency	Not fluent	87 (20.3)	
Diabetes duration (years)			5.70±3.48
Attended diabetes education session(s)	Yes	199 (46.5)	
Perceived social support (MSPSS Score)			5.23±0.92
Glycaemic control	Have not achieved target	295 (68.9)	
Blood pressure control	Have not achieved target	130 (30.4)	
Diabetes knowledge test (MDKT Score)			7.70±2.31

MDKT, michigan diabetes knowledge test; MSPSS, multidimensional scale of perceived social support.

out of 50 (SD=7.04). The prevalence of limited health literacy in patients with T2DM was 65.3% (n=279). The majority of participants answered the questionnaire in Bahasa Malaysia (n=282, 65.8%) followed by Mandarin (n=82, 19.2%) and English (n=64, 15.5%). One patient has more than 30% of data missing, thus was not analysed. Final number of patients included in the analysis was 427. The mean age of the study population was 58.1 years (SD=10.6), and 60.3% (n=258) were female, 43.8% were Malay, 33.4% were Indian and 22.7% were Chinese. The mean duration of diabetes was 5.7 years (SD=3.48), 46.5% (n=199) had attended the diabetes education sessions, 68.9% (n=295) had not achieved target glucose control, conversely 30.4% (n=130) had not achieved target blood pressure control. The mean score for perceived social support was 5.23 (SD=0.92) and the mean diabetes knowledge was 7.70 (SD=2.31) (table 1).

All variables with a value of  $p < 0.25$  in the univariate analysis were included in the multivariate analysis. This analysis was performed using a binary logistic regression (table 2). The results of the likelihood ratio test indicated that perceived social support, attendance at diabetes education sessions in the clinic and English fluency were

statistically significant factors associated with limited health literacy ( $\chi^2(15)=89.06$ ,  $p < 0.001$ ). Those who were not fluent in English (OR=2.36, 95% CI 1.30 to 4.30) were twice as likely to present with limited health literacy. Patients with better perceived social support (OR=0.52, 95% CI 0.40 to 0.69) and those who attended diabetes education sessions at the clinic (OR=0.42, 95% CI 0.27 to 0.68) were more likely to have adequate health literacy than those who did not.

### Consent to participate

All participants provided written informed consent prior to participating in the study.

### DISCUSSION

In this study, we found that more than two-thirds of the patients with T2DM attending primary care clinics in Perak, Malaysia have limited health literacy. Malaysia seems to be facing the same challenges as other countries with regards to limited health literacy in patients with T2DM. However, the prevalence in Malaysia is much higher than in several high-income countries such as the

**Table 2** Factors associated with limited health literacy (n=427)

Variables	Adequate HL		P value (univariate)	aOR	95% CI	P value (multivariate)
	n (%)	Limited HL n (%)				
Age (mean±SD)	57.17±10.39	58.63±10.72	0.179‡	1.00	0.97–1.02	0.757
Gender						
Male	59 (39.9)	110 (39.4)	0.930			
Female	89 (60.1)	169 (60.6)				
Ethnicity						
Malay	72 (49.7)	113 (40.9)	0.040§	0.82	0.45–1.49	0.506
Chinese	23 (15.9)	73 (26.4)		1.08	0.50–2.34	0.837
Indian	50 (34.5)	90 (32.6)		Reference		
Marital status						
Single/divorced/separated/widowed	33 (22.4)	66 (23.7)	0.779			
Married	114 (77.6)	213 (76.3)				
Highest education						
Primary school	29 (19.6)	103 (37.2)	<0.001§	1.91	0.68–5.35	0.217
Junior high school	33 (22.3)	93 (33.6)		2.00	0.83–4.84	0.124
High school	53 (35.8)	60 (21.7)		1.41	0.64–3.07	0.394
University/Masters/PhD	33 (22.3)	21 (7.6)		Reference		
Employment status						
Yes	59 (40.4)	92 (33.3)	0.088‡	0.85	0.43–1.71	0.652
No, never worked	33 (22.6)	90 (32.6)		1.16	0.60–2.22	0.66
No, stopped working or retired	54 (37.0)	94 (34.1)		Reference		
Monthly income						
Less than RM1000	70 (47.3)	173 (62)	0.003‡	0.77	0.41–1.43	0.401
Equal to or more than RM1000	78 (52.7)	106 (38)		Reference		
Health Insurance						
No	103 (70.1)	226 (81.6)	0.007§	1.34	0.75–2.40	0.319
Yes	44 (29.9)	51 (18.4)		Reference		
English fluency						
Not fluent	69 (46.6)	205 (73.5)	<0.001§	2.36	1.30–4.30	0.005§
Fluent	79 (53.4)	74 (26.5)		Reference		
Bahasa Malaysia fluency						
Not fluent	21 (14.2)	66 (23.7)	0.021§	1.30	0.60–2.83	0.51
Fluent	127 (85.8)	213 (76.3)		Reference		
Diabetes duration (years)	5.60±3.14	5.75±3.65	0.686			
Attended diabetes education session(s)						
Yes	91 (61.5)	108 (38.7)	<0.001§	0.42	0.27–0.68	<0.001§
No	57 (38.5)	171 (61.3)		Reference		
Perceived Social Support (MSPSS Score) (mean±SD)	5.58±0.93	5.05±0.86	<0.001*	0.52	0.40–0.69	<0.001§
Glycaemic control*						
Have not achieved target	97 (65.5)	197 (70.6)	0.282			
Achieved target	51 (34.5)	82 (29.4)				
Blood pressure control†						
Have not achieved target	49 (33.1)	81 (29)	0.384			

Continued



Table 2 Continued

Variables	Adequate HL	Limited HL	P value	aOR	95% CI	P value
	n (%)	n (%)	(univariate)			(multivariate)
Achieved target	99 (66.9)	198 (71)				
Diabetes knowledge test (MDKT Score) (mean±SD)	8.06±2.11	7.51±2.39	0.019§	0.99	0.89–1.10	0.819

\*HbA1c ≤ 6.5% or FBS ≤ 7.0.

†Blood pressure ≤ 135/75 mm Hg.

‡p<0.25.

§p<0.05.

aOR, adjusted OR; HbA1c, haemoglobin A1c; MDKT, michigan diabetes knowledge test; MSPSS, multidimensional scale of perceived social support.

USA (30%), Canada (15.6%)<sup>32</sup> and countries in Europe, such as Switzerland (8.7%)<sup>33</sup> and the Netherlands (9.7%).<sup>34</sup> There had not been a comprehensive study to explain the reason for these discrepancies. However, use of different health literacy measurement tools may have contributed to the differences in prevalence.<sup>11</sup> Findings from our study suggested that sociodemographic characteristics and healthcare support may also play an important role in determining the prevalence level.

Patients with less perceived social support, not fluent in English and, those who did not attend diabetes education sessions were more likely to have limited health literacy. Patients with better perceived social support from friends, family and significant others were more likely to have adequate health literacy. Studies have shown that people with a larger social network are more likely to use interpersonal communication to seek health information than are people with a smaller social network.<sup>35</sup> Qualitative studies suggest that people with a long-term health condition often draw on the health literacy skills of others in their social network<sup>36</sup> and that people with high health literacy tend to pass on health information to others.<sup>37</sup> This finding highlights the importance of social support in the management of patients with T2DM. Family members and significant others should play a more central role in empowering patients with T2DM.

Limited health literacy is also associated with not attending diabetes self-management education (DSME) sessions delivered at the clinics. DSME is an integral part of diabetes management, and many guidelines recommend that all patients with T2DM receive DSME at diagnosis and as needed thereafter.<sup>38</sup> Diabetes education sessions that tailor education to patients with low health literacy levels have been shown to improve patient knowledge, activation, self-efficacy, diabetes distress and self-care behaviours.<sup>39</sup> Currently, DSME is delivered by trained diabetes nurse educators. However, there is a significant lack of diabetes educators in Malaysia's healthcare system. Since 2004, the training courses for diabetes nurse educators by the Ministry of Health have produced approximately 900 trained diabetes nurse educators to practise in a total of 1195 public hospitals and primary

care clinics. Although DSME is expected as part of the standard care for all patients with T2DM, in reality, only approximately 4%–7% of patients have received it.<sup>40 41</sup> In our study, only 50% of patients surveyed attended DSME sessions at their clinics. Interventions are needed to increase patients' participation in DSME, and a review of the DSME programme to include training for health literacy may be a game changer in diabetes management.

This study identified three factors associated with limited health literacy that need to be considered when planning for interventions. In contrast to other studies, this study found the association between health literacy level and education level only at the univariate analysis level. In the final regression model, education level was not significantly associated with health literacy; instead, patients' fluency in English displayed a significant association. Since the multicollinearity analysis was not significant, there were no confounders for this relationship. Limited health literacy is less likely in patients who are fluent in English. A previous study showed that patients who could understand health information better were more likely to have higher health literacy levels.<sup>3</sup> It is thought that language influences the amount of information available to patients especially online. Also, in Malaysia, health information and patients' interactions with healthcare providers are delivered mostly in English. Since most patients received their information from their healthcare providers, limitations in the English language may limit their understanding and appraisal. There may also be other explanations for this observation. Therefore, we plan to embark on a qualitative study to further explore the reasons for this observation.

The strengths of our study is that it had expanded the understanding of health literacy levels in patients with T2DM. Previously, we conducted a systematic review and meta-analysis of prevalence of limited health literacy in the same population of patients and found a lack of data from low-income to middle-income countries. Most prevalence data came from studies conducted in the USA and Europe.<sup>42</sup> This study is only the second study done in Malaysia and the first to measure multidimensional aspects of health literacy. The other study showed

85.8% of patients have limited health literacy but used the Newest Vital Signs (NVS) to measure health literacy level.<sup>24</sup> The NVS only measures the numeracy component of health literacy<sup>36</sup> and has been shown to have administration issues in certain populations in Malaysia.<sup>43</sup> In contrast, the HLS-EU-Q47 was well validated in the Malaysian population and was available in three languages commonly used by the multiethnic population of Malaysia.<sup>26</sup> The use of HLS-EU-Q47 to measure HL in patients with T2DM is limited. So far only one other study in Spain had used this tool and reported 81.5% of patients with T2DM had limited health literacy.<sup>44</sup> This study also added to the data on health literacy in Malaysia. Health literacy is a new research topic in Malaysia that is gaining importance. The health literacy levels of the adult population in Malaysia was first measured in 2015 and the survey found only 6.6% of Malaysian adults had adequate health literacy.<sup>13</sup> Our study highlighted the need for further studies on limited health literacy as the problem is also found in patient populations and not just in the general population.

However, this study is not without limitations. Data for this study were collected only in Perak, which may not be representative of all states in Malaysia. Nevertheless, it was not our aim to generalise the findings to the entire country; rather, this study will form a basis to call for a countrywide measure of the health literacy level in patients with T2DM. Since the health literacy level and the associated factors were measured cross-sectionally, causal relationships could not be determined. Patients with type 2 DM who attended DSME or who have better social support may already have adequate health literacy level despite these factors rather than as a result of these factors. A qualitative study is planned to follow-up on these findings to fully understand the influence and impact of these factors on patients' health literacy level.

## CONCLUSIONS

In summary, this study found limited health literacy to be prevalent in patients with T2DM. This means that most patients with T2DM attending primary care clinics are unable to fully use their abilities to find, understand, appraise and apply information to optimally manage their disease. Strategies to improve health literacy are required, and must consider issues related to limited English fluency, attendance to diabetes education sessions and social support.

**Acknowledgements** The authors thank the study subjects for participating in this study. The authors also thank Dr James Gnanasigamani, Dr Azlin Yusma bt Md Yusup, Dr S Shailaja Siva Sankaran, Mastura bt Awang, Uma A/P Muthu, Sattieswary A/P Govindasamy and Massarah Kee bt Shaharuddin Kee for their support in data collection and participant recruitment. The authors specially thank Dr Alison A. Jackson for proofreading the manuscript.

**Contributors** CJN and SML contributed to the conceptualisation of the study, study design and the writing of the manuscript. SA and PV contributed to the data collection and data analysis. KC contributed to the study design and data analysis. AA contributed to the conceptualisation of the study, study design, data collection,

data analysis and the writing of the manuscript. All authors read and approved the final version of the manuscript.

**Funding** This work was supported by the University Malaya Research Grant, Special Grant, grant no. BKS047-2017.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. The deidentified participant data are available upon reasonable request to the corresponding author. She could be contacted at adina@ummc.edu.my.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Adina Abdullah <http://orcid.org/0000-0003-1545-7110>

Chirk Jenn Ng <http://orcid.org/0000-0002-8320-1603>

## REFERENCES

- Sørensen K, Van den Broucke S, Fullam J, *et al.* Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012;12:80.
- Selden CR, Zorn M, Ratzan SC, *et al.* *National library of medicine current bibliographies in medicine: health literacy*. Bethesda, MD: National Institutes of Health, 2000.
- Chen G-D, Huang C-N, Yang Y-S, *et al.* Patient perception of understanding health education and instructions has Moderating effect on glycemic control. *BMC Public Health* 2014;14:683.
- Powell CK, Hill EG, Clancy DE. The relationship between health literacy and diabetes knowledge and readiness to take health actions. *Diabetes Educ* 2007;33:144–51.
- Caruso R, Magon A, Baroni I, *et al.* Health literacy in type 2 diabetes patients: a systematic review of systematic reviews. *Acta Diabetol* 2018;55:1–12.
- Thurston MM, Bourg CA, Phillips BB, *et al.* Impact of health literacy level on aspects of medication nonadherence reported by underserved patients with type 2 diabetes. *Diabetes Technol Ther* 2015;17:187–93.
- Haun JN, Patel NR, French DD, *et al.* Association between health literacy and medical care costs in an integrated healthcare system: a national population based study. *BMC Health Serv Res* 2015;15:249.
- DeWalt DA, Boone RS, Pignone MP. Literacy and its relationship with self-efficacy, trust, and participation in medical decision making. *Am J Health Behav* 2007;31 Suppl 1:27–35.
- Rothman RL, Dewalt DA, Malone R. Influence of patient literacy on the effectiveness of a primary care-based diabetes disease management program. *JAMA* 2004;292:1711–6.
- Gerber BS, Brodsky IG, Lawless KA, *et al.* Implementation and evaluation of a low-literacy diabetes education computer multimedia application. *Diabetes Care* 2005;28:1574–80.
- Abdullah A, Liew SM, Salim H, *et al.* Prevalence of limited health literacy among patients with type 2 diabetes mellitus: a systematic review. *PLoS One* 2019;14:e0216402–16.
- Ogurtsova K, da Rocha Fernandes JD, Huang Y, *et al.* IDF diabetes atlas: global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res Clin Pract* 2017;128:40–50.
- Institute for Public Health (IPH). *National health and morbidity survey 2015 (NHMS 2015) vol. II: non-communicable diseases, risk factors & other health problems*, 2015. ISBN: 978-983-2387-23-7.
- Zaki M, Robaayah Z, Chan SP, *et al.* Malaysia shape of the nation (MySoN): a primary care based study of abdominal obesity in Malaysia. *Med J Malaysia* 2010;65 Suppl A:143–9.
- Mohamad WNW, Ismail Aal-S, Khir ASM, *et al.* Prevalence of metabolic syndrome and its risk factors in adult Malaysians: results of a nationwide survey. *Diabetes Res Clin Pract* 2012;96:91–7.
- Jeganathan R, Karalasingam SD, eds. *2nd report of national obstetrics registry 2010*. Putrajaya: National Obstetrics Registry and the Clinical Research Centre, 2013.

- 17 CPG Secretariat Ministry of Health Malaysia. *Clinical practice guidelines for the management of type 2 diabetes mellitus*. 5th edition. Kuala Lumpur: Joint Publication of the Ministry of Health Malaysia, Academy of Medicine Malaysia, Malaysian Endocrine & Metabolic Society and Diabetes Malaysia, 2015: 140.
- 18 Mafauzy M, Hussein Z, Chan SP. The status of diabetes control in Malaysia: results of DiabCare 2008. *Med J Malaysia* 2011;66:175–81.
- 19 Bak Leong G, Loke Meng O, Yam Ngo L. *21st report of the Malaysian dialysis & transplant registry 2013*. Kuala Lumpur: The National Renal Registry Malaysia, 2014.
- 20 Rothman R, Malone R, Bryant B, et al. The relationship between literacy and glycemic control in a diabetes disease-management program. *Diabetes Educ* 2004;30:263–73.
- 21 Kim S, Love F, Quistberg DA, et al. Association of health literacy with self-management behavior in patients with diabetes. *Diabetes Care* 2004;27:2980–2.
- 22 Sarkar U, Karter AJ, Liu JY, et al. Hypoglycemia is more common among type 2 diabetes patients with limited health literacy: the diabetes study of northern California (distance). *J Gen Intern Med* 2010;25:962–8.
- 23 Peduzzi P, Concato J, Kemper E, et al. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996;49:1373–9.
- 24 Azreena E, Suriani I, Juni MH, et al. Factors associated with health literacy among type 2 diabetes mellitus patients attending a government health clinic, 2016. *Int J Public Heal Clin Sci* 2016;3:50–64.
- 25 Sørensen K, Van den Broucke S, Pelikan JM, et al. Measuring health literacy in populations: illuminating the design and development process of the European health literacy survey questionnaire (HLS-EU-Q). *BMC Public Health* 2013;13.
- 26 Duong TV, Aringazina A, Baisunova G, et al. Measuring health literacy in Asia: validation of the HLS-EU-Q47 survey tool in six Asian countries. *J Epidemiol* 2017;27:80–6.
- 27 Sørensen K, Pelikan JM, Röthlin F, et al. Health literacy in Europe: comparative results of the European health literacy survey (HLS-EU). *Eur J Public Health* 2015;25:1053–8.
- 28 Fitzgerald JT, Funnell MM, Hess GE, et al. The reliability and validity of a brief diabetes knowledge test. *Diabetes Care* 1998;21:706–10.
- 29 Al-Qazaz HK, Hassali MA, Shafie AA, et al. The 14-item Michigan diabetes knowledge test: translation and validation study of the Malaysian version. *Pract Diab Int* 2010;27:238–41.
- 30 Zimet GD, Dahlem NW, Zimet SG, et al. The multidimensional scale of perceived social support. *J Pers Assess* 1988;52:30–41.
- 31 Ng CG, Amer Siddiq AN, Aida SA, et al. Validation of the Malay version of the multidimensional scale of perceived social support (MSPSS-M) among a group of medical students in faculty of medicine, University Malaya. *Asian J Psychiatr* 2010;3:3–6.
- 32 Al Sayah F, Majumdar SR, Johnson JA. Association of inadequate health literacy with health outcomes in patients with type 2 diabetes and depression: secondary analysis of a controlled trial. *Can J Diabetes* 2015;39:259–65.
- 33 Mantwill S, Schulz PJ. Low health literacy associated with higher medication costs in patients with type 2 diabetes mellitus: evidence from matched survey and health insurance data. *Patient Educ Couns* 2015;98:1625–30.
- 34 van der Heide I, Wang J, Droomers M, et al. The relationship between health, education, and health literacy: results from the Dutch adult literacy and life skills survey. *J Health Commun* 2013;18 Suppl 1:172–84.
- 35 Askelson NM, Campo S, Carter KD. Completely isolated? health information seeking among social isolates. *Health Educ Behav* 2011;38:116–22.
- 36 Edwards M, Wood F, Davies M, et al. 'Distributed health literacy': longitudinal qualitative analysis of the roles of health literacy mediators and social networks of people living with a long-term health condition. *Health Expect* 2015;18:1180–93.
- 37 Ellis J, Mullan J, Worsley A, et al. The role of health literacy and social networks in arthritis patients' health information-seeking behavior: a qualitative study. *Int J Family Med* 2012;2012:1–6.
- 38 American Diabetes Association. Standards of medical care in diabetes D 2017. *Diabetes Care* 2017;35.
- 39 Wallace AS, Seligman HK, Davis TC, et al. Literacy-appropriate educational materials and brief counseling improve diabetes self-management. *Patient Educ Couns* 2009;75:328–33.
- 40 Duncan I, Birkmeyer C, Coughlin S, et al. Assessing the value of diabetes education. *Diabetes Educ* 2009;35:752–60.
- 41 Li R, Shrestha SS, Lipman R, et al. Diabetes self-management education and training among privately insured persons with newly diagnosed diabetes — United States, 2011–2012. *Morb Mortal Wkly Rep* 2014;63:1045–9.
- 42 Abdullah A, Liew SM, Salim H, et al. Prevalence of limited health literacy among patients with type 2 diabetes mellitus: a systematic review. *PLoS One* 2019;14:e0216402.
- 43 Norrafizah J, Asiah M, Suraiya S, et al. Assessment of health literacy among people in a rural area in Malaysia using newest vital signs assessment. *BJESBS* 2016;16:1–7.
- 44 Fernández-Silva MJ, Alonso-González A, González-Pérez E. Alfabetización en salud en pacientes Con diabetes tipo 2: un Estudio transversal Con El cuestionario HLS-EU-Q47. *Med Fam Semer* 2018;1:30–6.