

Table S3: Percentage of patients with a comorbidity in the year before the start of their wound. Those with diabetes are a subset of those with endocrinological comorbidities.

	Abscess	Burn	Diabetic foot ulcer	Leg ulcer (arterial)	Leg ulcer (mixed)	Leg ulcer (unspecified)	Leg ulcer (venous)	Open wound	Pressure ulcer	Surgical wound	Trauma	Unspecified	All wounds
Cardiovascular	36%	35%	70%	67%	83%	45%	78%	44%	53%	47%	38%	51%	53%
Dermatological	48%	52%	60%	44%	63%	50%	41%	51%	40%	51%	54%	45%	49%
Endocrinological	52%	56%	99%	78%	97%	18%	52%	69%	60%	63%	55%	57%	60%
Diabetes	50%	52%	100%	78%	97%	13%	37%	67%	60%	59%	51%	52%	57%
Gastroenterological	29%	32%	42%	67%	33%	29%	19%	34%	47%	37%	31%	35%	33%
Genito-urinary	21%	24%	28%	44%	33%	18%	19%	25%	27%	28%	24%	24%	24%
Immunological	7%	11%	13%	11%	17%	3%	11%	10%	7%	9%	8%	9%	9%
Malnutrition	0%	2%	2%	0%	0%	5%	4%	1%	13%	1%	2%	3%	3%
Musculoskeletal	41%	56%	62%	67%	63%	61%	59%	49%	73%	57%	49%	56%	56%
Neurological	16%	21%	24%	11%	27%	24%	33%	18%	67%	22%	17%	17%	24%
Oncological	14%	16%	25%	11%	20%	16%	7%	14%	27%	18%	15%	19%	16%
Ophthalmological	8%	15%	20%	0%	27%	13%	4%	11%	33%	13%	11%	10%	12%
Psychiatric	44%	40%	39%	33%	53%	47%	15%	35%	33%	35%	32%	35%	35%
Renal	8%	5%	30%	22%	33%	13%	30%	12%	40%	14%	9%	16%	18%
Respiratory	38%	44%	45%	56%	43%	42%	33%	36%	33%	38%	36%	37%	38%
Mean number of comorbidities per patient	3.7	4.1	5.7	6.0	5.7	3.9	4.0	4.1	5.6	4.3	3.8	4.2	4.1