

**Supplementary Table 1** Treatment duration and dosing

	<b>Regorafenib (N=100)</b>
Time on treatment, months <sup>a</sup>	
Mean (SD)	3.9 (4.1)
Median (range)	2.5 (0.1–20.6)
Time on treatment >18 weeks, n (%) <sup>a</sup>	29 (29)
Number of cycles completed	
Mean (SD)	4.4 (4.3)
Median (range)	3 (1–22)
Completed ≥5 cycles, n (%)	29 (29)
Actual daily dose <sup>b</sup>	
Mean (SD), mg	150.8 (15.1)
Median (range), mg	160.0 (101.5–160.0)
<160 mg, n (%)	34 (34)
160 mg, n (%)	66 (66)
Percent of planned dose, % <sup>c</sup>	
Mean (SD)	87.3 (16.4)
Patients with treatment modifications, n (%)	
Any	72 (72)
Dose reduction	33 (33)
Treatment interruption/delay	64 (64)
Dose re-escalation <sup>d</sup>	7 (7)

<sup>a</sup>Includes treatment interruptions/delays.

<sup>b</sup>Includes only days on which a dose was received (i.e. excludes treatment interruptions/delays).

<sup>c</sup>Planned dose describes the intended initial dose according to the protocol.

<sup>d</sup>Permitted at the discretion of the treating physician, up to a maximum of 160 mg/day.

SD=standard deviation