Correction: *The premature closure of ROMPA clinical trial: mortality reduction in septic shock by plasma adsorption*

Giménez-Esparza C, Portillo-Requena C, Colomina-Climent F, *et al.* The premature closure of ROMPA clinical trial: mortality reduction in septic shock by plasma adsorption. *BMJ Open* 2019;9:e030139. doi: 10.1136/bmjopen-2019-030139.

This article was previously published with an error.

The following statement in the discussion section is incorrect: 'In addition, as in our study, the COMPACT 2 group used a randomisation system based on prognostic scores,⁶ which means that the groups will not be similar until the end of recruitment (which is reason to introduce the propensity score in our results).' The COMPACT 2 group used a blocked randomization schedule (randomly permuting blocks of four and six), with stratification according to site and the presence of septic shock at admission.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2020;**10**:e030139corr1. doi:10.1136/bmjopen-2019-030139corr1