Correction: recorded poor insight as a predictor of service use outcomes: cohort study of patients with first-episode psychosis in a large mental healthcare database

Ramu N, Kolliakou A, Sanyal J, *et al.* Recorded poor insight as a predictor of service use outcomes: cohort study of patients with first-episode psychosis in a large mental healthcare database. *BMJ Open* 2019;9:e028929. doi: 10.1136/bmjopen-2019-028929

This article has been corrected since it was published.

The authors have corrected the coding error (in the variable measuring insight, the main exposure) and have re-run the analyses. Therefore, 'Results', 'Conclusions' in 'Abstract' section; 'Patients', 'Unadjusted and adjusted main outcomes' in 'Results' section; 'Discussion' and 'Conclusion' have been updated. All the tables have been revised and updated.

However, the main findings of the paper remain unaltered.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2020;**10**:e028929corr1. doi:10.1136/bmjopen-2019-028929corr1