BMJ Open Cardiac arrest as a reportable condition: a cohort study of the first 6 years of the Norwegian out-of-hospital cardiac arrest registry

Ingvild Beathe Myrhaugen Tjelmeland ^{1,2}, Kristin Alm-Kruse,^{3,4} Lars-Jøran Andersson,⁵ Ståle Bratland,⁵ Arne-Ketil Hafstad,⁶ Bjørn Haug,⁷ Jørund Langørgen,⁸ Alf Inge Larsen,^{9,10} Thomas Werner Lindner,¹¹ Jan Erik Nilsen,¹ Theresa M Olasveengen,^{4,12} Eldar Soreide,^{13,14} Eirik Skogvoll,^{15,16} Jo Kramer-Johansen^{1,4}

To cite: Tjelmeland IBM, Alm-Kruse K, Andersson L-J, *et al.* Cardiac arrest as a reportable condition: a cohort study of the first 6 years of the Norwegian out-of-hospital cardiac arrest registry. *BMJ Open* 2020;**10**:e038133. doi:10.1136/bmjopen-2020-038133

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-038133>).

Received 28 February 2020
Revised 23 April 2020
Accepted 12 May 2020

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Ingvild Beathe Myrhaugen Tjelmeland; ingvild@nako.no

ABSTRACT

Objectives The Norwegian Cardiac Arrest Registry (NorCAR) was established in 2013 when cardiac arrest became a mandatory reportable condition. The aim of this cohort study is to describe how the world's first mandatory, population-based cardiac arrest registry evolved during its first 6 years.

Setting Norway has a total population of 5.3 million inhabitants with a population density that varies considerably. All residents are assigned a unique identifier number, giving nationally approved registries access to information about all births and deaths in the country. Data in the registry are entered by data processors; public employees with close links to the emergency medical services. All data processors undergo a standardised training and meet for yearly retraining and updates.

Participants All events of cardiac arrest where bystanders or healthcare professionals have started cardiopulmonary resuscitation or performed defibrillation are included into the NorCAR.

Primary and secondary outcome measures Since the establishment of the registry, the number of reporting health trusts, the number of reported events and the corresponding population at risk were followed year by year. Outcome is measured as changes in inclusion rate, incidence per 100 000 inhabitants and survival to 30 days after cardiac arrest.

Results In total, 14 849 cases were registered over 6 years, between 2013 and 2018. The number of health trusts reporting rose steadily from 2013. Within 3 years, all trusts reported to the registry with an increasing number of events reported; going from 1101 to 3400 per year. The prevalence of bystander cardiopulmonary resuscitation increased slightly, but the population incidence of survival did not change.

Conclusion Declaring cardiac arrest as a reportable condition and close follow-up of all reporting areas is essential when building a national registry.

INTRODUCTION AND BACKGROUND

Response, treatment and outcome after out-of-hospital cardiac arrest (OHCA) is an

Strengths and limitations of this study

- Cohort study of the world's first mandatory cardiac arrest registry presenting establishment and the first results.
- Includes almost 15 000 cases of cardiac arrest.
- Merging information from a lot of different ambulance services uncovers challenges related to the understanding of the different data definitions.
- The Norwegian registry uses the national unique personal identifier number, and generalisability of the method might be limited.

internationally recognised quality indicator for the emergency medical service (EMS).¹ The links in the chain-of-survival highlight both the time-sensitive nature of interventions and the concept that high quality in all stages of treatment is necessary to improve patient outcome.²

A population-based cardiac arrest registry allows systematic assessment of the various components and aspects of resuscitation care; early recognition of cardiac arrest, bystander cardiopulmonary resuscitation (CPR), handling of emergency calls, dispatch assisted CPR (DA-CPR), ambulance dispatch and care, and hospital treatment. Registry data may facilitate an understanding of how each element in a system affects survival and identify potential for improvement.³

The reported population incidence of OHCA in which CPR is initiated by a bystander or the EMS varies worldwide. A recent European study reports an incidence between 28 and 91 per 100 000 inhabitants.⁴ The Australasian Resuscitation Outcomes Consortium (Aus-ROC) report similar estimates between 35 and 54 per 100 000 inhabitants.⁵ The outcome after OHCA varies

between and within countries. Sweden, establishing a voluntary registry already in 1990, reports the incidence of survival after OHCA between 3.7 and 10.4 per 100 000 inhabitants,⁶ similar to prior publications from single centres in Norway.^{7,8}

In 2013, Norwegian Cardiac Arrest Registry (NorCAR) became part of the Norwegian Cardiovascular Disease Registry (NCVDR) run by Norwegian Institute of Public Health and reporting to the registry became mandatory.

The aim of this paper is to describe how the inclusion rate, the incidence and estimated survival developed in this world's first mandatory, population-based cardiac arrest register over its first 6 years.

METHODS

Setting

Norway has 5.3 million inhabitants with a population density that varies considerably between urban and rural areas (Oslo 223/km²; Finnmark 1.6/km²). The Norwegian single-payer public insurance covers all aspects of healthcare. Specialist healthcare is organised in 4 regional and 19 local Health Trusts. For medical emergencies, there is one dedicated medical emergency telephone number, 113, that terminates at 16 local Emergency Medical Communications Centres (EMCCs).

EMCCs are staffed with nurses and ambulance personnel, and dispatch the EMS. The EMS response may include ambulance, physician staffed helicopters/rapid response car, ambulance boats as well as first responders from the fire brigade and out-of-office-hours general practitioners, all dependent on the location of the arrest and the available resources. Telephone-guided first aid, including DA-CPR, has been an integrated EMCC routine since 1995. DA-CPR is provided to bystanders when the patient is unconscious and not breathing normally.

Data management and governance

All residents in Norway are assigned an 11-digit unique personal identifier number. This ID is a common identifier across all public registries, including health systems and can be used by approved national registries to confirm data about birth, name, address and date of death. Provisional ID can be generated for temporary residents or when ID is unknown.

Registration of diseases and procedures related to cardiovascular conditions in NCVDR is regulated to allow the collection, analysis and use of sensitive data.⁹ Reporting to NorCAR is mandatory as it is part of NCVDR, and there is no need for consent; rather all healthcare personnel and facilities have a duty to report. Through the online reporting system, the strictest level of patient privacy and data integrity is maintained in accordance with best practice and regulations.

Entry into NorCAR is made by data processors, often part-time nurses or paramedics with other tasks related to cardiac arrest education, organisation or quality improvement. All data processors are employed by the

local Health Trusts. To ensure equal understanding of data definitions, data processors undergo standardised training and meet for yearly retraining and updates.

NorCAR is governed by a national advisory board representing all Regional Health Trusts, the Norwegian Resuscitation Council, the Norwegian Society of Cardiology, the Norwegian Myocardial Infarction Registry and the National Advisory Unit of Prehospital Emergency Medicine. The board also includes a data processor and a patient representative.

Patient and public involvement

The user representative involved in this project is part of NorCARs steering committee. He represents the patient organisation National Association of Heart and Lung Disease that has 54 000 members across Norway suffering from cardiovascular diseases, including myocardial infarction, heart failure and stroke. He provides a channel for communication to the patient population as well as to the boards of health trusts through his network of fellow user representatives. He has been involved in the development of this research project.

Event registration

All patients that are unresponsive without normal breathing, where bystander, first responders, EMS or other healthcare professionals initiate CPR or defibrillation, are included in the registry. DA-CPR is defined as treatment alongside basic or advanced CPR or defibrillation. Patients are included irrespective of having a confirmed cardiac arrest on EMS arrival or not. Patients who do not receive any CPR are not included, neither are newborns in need of resuscitation at birth.

Data sources

Registration of an event often starts in the EMCC when DA-CPR is initiated. After the event the attending ambulance personnel fill in a dedicated form and transfer the information to the data processor. The local data processors collect information on the event and the patient, checks data quality and collect additional information from the defibrillator and hospital records when applicable. To ensure complete local coverage, data processors routinely check for missing cases in the EMCC databases, air-ambulance patient report forms, hospital electronic records and the Norwegian Patient Registry. Several logical checks have been implemented in the electronic report form, preventing typing errors or non-logical entries and finally the national registry staff performs a second quality control of all submitted data.

Variables

The dataset in NorCAR adheres to the Utstein dataset and definitions,¹⁰ with some minor modifications to adjust for local conditions, and include information on the event, treatment and outcome. A full description of our dataset can be found in the online supplementary appendix.

Statistical methods

When calculating bystander CPR rates, all EMS witnessed cardiac arrests are excluded from the analysis. When calculating the incidence of ROSC and 30-day survival, only patients treated by EMS, or patients who have been successfully resuscitated using an Automatic External Defibrillator (AED) before EMS arrival, are included.

Descriptive measures are provided as mean with SD or median with range, as appropriate according to the distribution of data. For regions reporting part of a year, we calculate population incidence by dividing number of patients by the corresponding fraction of the person-years for that region. Yearly trends in the number of reported events, the population incidence and the incidence of survivors with 95% CIs were investigated using Poisson regression with year as predictor variable. Yearly trends in the prevalence of bystander CPR were investigated with logistic regression with year as predictor variable. A $p < 0.05$ was considered as statistically significant. The software R¹¹ was used for the statistical analyses.

Data protection

NorCAR is a national person-identifiable health registry that does not require consent from the registered individual or next of kin.⁹

RESULTS

Regional and national development since inception

A total of 14849 OHCA events were registered in NorCAR from 2013 to 2018. The number of participating health trusts rose steadily, and by May 2016 data collection and reporting was established in all of Norway (figure 1).

The reported number of OHCA per year increased from 1101 in 2013 to 3400 in 2018 (figure 2, $p < 0.01$). The estimated population incidence also increased from 46 (95% CI 45 to 50) to 64 (62 to 66) per 100 000 inhabitants ($p < 0.01$). The incidence of OHCA treated by ambulance increased from 44 to 53 per 100 000 inhabitants ($p < 0.01$). (table 1)

Utstein comparator group

The Utstein comparator group of 2014¹⁰ is defined as patients experiencing witnessed cardiac arrest where the first documented rhythm is shockable. In addition, we

Figure 1 Map of Norway showing health trusts that report data to the National registry. Green: data collected the entire year, yellow: data collected for parts of the year, red: did not report.

Figure 2 All patients included in the registry. Left axis and bars: number of out-of-hospital cardiac arrests treated by ambulance, by year of registration. Patients with EMS-observed collapse are identified as light grey. Right axis and lines: population incidence of ambulance treated arrests (blue), and population incidence of 30-day survival (green); per 100 000 inhabitants. EMS, emergency medical service.

have included witnessed cardiac arrest-patients successfully resuscitated before EMS arrival to our Utstein comparator group. These patients have been defibrillated before EMS personnel arrive on scene and have spontaneous circulation when EMS arrives (shown as orange in figure 3).

The incidence of the Utstein comparator group has varied between 7.4 and 8.9 per 100 000 inhabitants, with a survival incidence in this group of 3.5 per 100 000 inhabitants (42%). The incidence of ambulance witnessed cardiac arrests with shockable first rhythm has been about 2 per 100 000 inhabitants with a survival incidence of 1.2 per 100 000 inhabitants (63%). The survivors in the Utstein comparator group and ambulance witnessed arrests constitute more than 70% of the total number of 30-day survivors (figure 3).

Patient demography and outcome

Median age was 65 (IQR 55–79) years and 67% were male. In 80% of the cases, CPR was started before EMS arrival (table 1). Among the 13 003 ambulance treated patients, 15 per 100 000 inhabitants (30%) achieved return of spontaneous circulation (ROSC) that lasted long enough for hospital admission or for transferral to the next treating prehospital unit (sustained ROSC).

Likewise, 30-day survival was 7.1 per 100 000 inhabitants (14%) of the ambulance treated patients, and did not change over the period investigated ($p = 0.98$). If we include patients defibrillated before EMS arrival, the overall survival is 7.4 per 100 000 inhabitants (15%).

DISCUSSION

The main findings from the first 6 years of the world's first mandatory cardiac arrest registry was an initial increase in the reported incidence of OHCA, levelling off after a few years. We observed a stable incidence of survival at 30 days as well as a bystander CPR rate of 80% during

Table 1 OHCA cases that receive attempted resuscitation included in the Norwegian cardiac arrest registry from 2013 to 2018

	2013	%	Inct	%	2014	%	Inct	%	2015	%	Inct	%	2016	%	Inct	%	2017	%	Inct	%	2018	%	Inct	%	All cases	%	Inct	%
Population covered	2370729		46		3395193		43		4793741		53		5220143		61		5295619		60		5312343		64		26387768		56	
No of patients	1101				1444				2549				3168				3187				3400				14849			
Age (median, 25th and 75th percentiles)	64 (54, 79)				65 (56, 79)				65 (55, 80)				64 (54, 78)				65 (54, 79)				65 (55, 79)				65 (55, 79)			
Male gender	749	68	32	972	67	29	1744	68	36	2100	66	40	2135	67	40	2222	65	42	9922	67	38				9922	67	38	
Place of cardiac arrest home	695	63	29	897	62	26	1555	61	32	1884	59	36	1986	62	38	2129	63	40	9146	62	35				9146	62	35	
Witnessed by bystander	605	55	26	751	52	22	1312	51	27	1605	51	31	1656	52	31	1619	48	30	7548	51	29				7548	51	29	
Witnessed by EMS	155	14	6.5	162	11	4.8	329	13	6.9	389	12	7.5	401	13	7.6	356	10	6.7	1792	12	6.8				1792	12	6.8	
Presumed cardiac cause	882	80	37	1040	72	31	1752	69	37	2186	69	42	2183	68	41	2256	66	42	10299	69	39				10299	69	39	
No minus EMS witnessed	946				1280				2220				2779				2786			13055								
Bystander CPR	719	76			1004	78			1691	76			2227	80			2277	82		10402	80							
EMS treated	1038		44		1354		40		2320		48		2735		52		2731		52	13003		49						
Initial shockable rhythm	287	28	12	352	26	10	572	25	12	634	23	12	643	24	12	691	24	13	3179	24	12				3179	24	12	
Sustained ROSC	379	37	16	446	33	13	734	32	15	810	30	16	762	28	14	767	27	14	3898	30	15				3898	30	15	
30-day survival	190	18	8	182	13	5.4	361	16	7.5	375	14	7.2	368	13	6.9	402	14	7.6	1878	14	7.1				1878	14	7.1	
EMS treated + ROSC before EMS arrival	1043		44		1362		40		2331		49		2753		53		2751		52	13086		50						
30-day survival	195	19	8.2	187	14	5.5	371	16	7.7	392	14	7.5	382	14	7.2	422	15	7.9	1949	15	7.4				1949	15	7.4	
Utstein comparator group*	210		8.9		250		7.4		403		8.4		463		8.9		446		8.4	2230		8.5						
ROSC	138	66	5.8	158	63	4.7	236	59	4.9	259	56	5	260	58	4.9	255	56	4.8	1306	59	4.9				1306	59	4.9	
30-day survival	104	50	4.4	99	40	2.9	172	43	3.6	187	40	3.6	186	42	3.5	181	40	3.4	929	42	3.5				929	42	3.5	
EMS witnessed with shockable rhythm	49		2.1		48		1.4		84		1.8		106		2		122		2.3	519		2						
ROSC	37	76	1.6	36	75	1.1	62	74	1.3	76	72	1.5	86	70	1.6	80	73	1.5	377	73	1.4				377	73	1.4	
30-day survival	35	71	1.5	32	67	0.9	48	57	1	65	61	1.2	73	60	1.4	75	68	1.4	328	63	1.2				328	63	1.2	
Resuscitated before EMS arrival (n)	5				8				11				18				20		83							83		
30-day survival	5				5				10				17				14		71							71		

*Utstein comparator group including patients defibrillated before EMS arrival, and presenting with ROSC as first registered rhythm by EMS.

Incidence of cardiac arrest per 100 000 inhabitants

CPR, cardiopulmonary resuscitation; EMS, emergency medical services; OHCA, out-of-hospital cardiac arrest.

Figure 3 The Utstein comparator group (witnessed collapse and shockable first rhythm). Left axis and bars: number of cardiac arrest patients by year of registration. Patients with EMS-observed collapse are identified as light grey. Patients successfully resuscitated with AED before EMS arrival are orange. Right axis and lines: population incidence of the total Utstein comparator group (including EMS witnessed cases and those resuscitated before EMS arrival) (blue line), and population incidence of 30-day survival in the total group (green); per 100 000 inhabitants. AED, Automatic External Defibrillator; EMS, emergency medical service.

the whole observation period. The likely explanation for the increasing number of events is a more complete registration.

The reported rate of bystander CPR of 80% is among the highest ever published.⁴ It is not fully understood why Norway has such high bystander CPR rate, but DA-CPR being implemented in all EMCCs already in 1995, a culture of willingness to help as well as repeated CPR training in school and adult life gives our inhabitants knowledge of and a willingness to perform CPR. The historical background also includes the first ever resuscitation training manikin being ‘born’ in Norway in 1960.¹² The current public strategy to increase bystander CPR involves collaborations between the Norwegian Health Directorate, non-governmental organisations and EMCCs.¹³ A study of the term ‘Bystander CPR’¹⁴ and an article from Aus-ROC¹⁵ both underline the importance of clear definitions of what constitutes a ‘bystander’. In the Norwegian registry, we have not seen any big changes in the rate of bystander CPR over the years, indication that the interpretation has not changed, and that the high bystander rates are representative for our nation.

The initial increases seen in total numbers and ambulance treated cardiac arrests per population at risk, has levelled off. We believe that this is due to an initial under-reporting of patients that were not treated by ambulance because they had clear signs of death, or the patient did not survive the event and was not transported to hospital. The stable survival rate at around 7 per 100 000 inhabitants, despite an increase in overall reported incidence, supports this. Reports from Denmark¹⁶ and Sweden¹⁷ indicate that survivors might also be under-reported to national registries, and a validation process of inclusions would further strengthen the registry.

As the number of included patients increase, the proportion of survivors decrease, again stressing the importance of reporting results as incidence rates in addition to percentages. Consistent method for reporting the number of cardiac arrests, compared with the number of inhabitants in the region covered, makes it possible to compare results across regions and between countries as suggested by Becker *et al*.¹⁸ Comparison of results can also be done for subgroups of patients, such as the Utstein Comparator group (witnessed and shockable first rhythm). As this group is more homogeneous in cause and treatment, percentages may be appropriate. In the Norwegian registry, ROSC rate and survival to 30 days in the Utstein comparator group were 59% and 42%, respectively. The ROSC rate is similar to the overall ROSC rate for this group in the EuReCa-two study, but the survival to 30 days is higher than reported both in the European study and in the Aus-ROC epistudy.^{4,5}

Some patients do not receive EMS treatment though bystanders have started CPR before ambulance arrival (12%, table 1). The reasons for EMS personnel not to start treatment can be signs of death, do-not-resuscitate-orders, further resuscitation is considered futile due to circumstances or comorbidities, or that the patient presents a pulse and signs of life on EMS arrival. Patients who have received bystander CPR, but have a pulse when EMS arrives, cannot be considered confirmed cardiac arrests. To be consistent and conservative, we do not include these patients when calculating survival with one exception; the patient has received a shock from an AED before EMS arrival. The number of reported patients in this group has increased from 5 in 2013 to more than 20, adding up to 83 patients over 6 years. The increase in the reported numbers could be due to a better reporting combined with increasing availability of AEDs in public places, use of first responders and the implementation of an AED-registry directly available in the maps in the EMCC.¹³ It is, however, important to acknowledge that there are a lot of barriers to an increased use of AED and one such barrier is that the majority of cardiac arrests occurring at home (62%).

Limitations of the study

Collecting data from different local registries has resulted in some challenges related to the understanding of the different data definitions in the Utstein template. Translating the dataset and distributing it to all EMS personnel takes time, meaning that differences between Health Trusts the first years of the registry might be related to different interpretation of the definitions just as well as real differences. We hope that the continuous work from the registry, including standardised training and yearly meetings has contributed to a common interpretation. It is also unknown if the focus on treatment of OHCA has changed registration practices or if there is a changing culture.¹⁸

Missing information in the registry is handled by returning the registration to the relevant health trust,

who then correct the data. Missing information that cannot be retrieved is handled by excluding the patient from the relevant analysis. It is, however, a challenge that by the time the national registry returns the form to the data processor, the cardiac arrest happened more than 3 months ago. Retrieving information more than 3 months after the event is difficult, and recall bias is substantial.

CONCLUSION

Having cardiac arrest as a reportable condition and closely following up all submitted registry data is essential when building a national registry. Defining cardiac arrest as a reportable condition made it possible to include all regions of Norway within a 4-year period, and increased the number of reported cardiac arrests cases to an incidence of 64 per 100 000 inhabitants. Survival to 30 days stayed stable with an incidence of 7.4 per 100 000 inhabitants.

Author affiliations

¹Division of Prehospital Services, Oslo University Hospital, Oslo, Norway

²Institute for Emergency Medicine, University Hospital Schleswig Holstein, Kiel, Germany

³Division of Emergencies and Critical Care, Oslo University Hospital, Oslo, Norway

⁴Institute of Clinical Medicine, University of Oslo, Oslo, Norway

⁵Clinic of Emergency Medicine, University Hospital of North Norway, Tromsø, Norway

⁶National Association for Heart and Lung Disease, Jessheim, Norway

⁷Department of Cardiology, Akershus University Hospital, Lorenskog, Norway

⁸Department of Heart Disease, Haukeland University Hospital, Bergen, Norway

⁹Department of Cardiology, Stavanger University Hospital, Stavanger, Norway

¹⁰Department of Clinical Science, University of Bergen, Bergen, Norway

¹¹Regional Competency Center for Emergency Medicine (RAKOS), Stavanger University Hospital, Stavanger, Norway

¹²Department of Research and Development and Department of Anesthesiology, Oslo University Hospital, Oslo, Norway

¹³Critical Care and Anaesthesiology Research Group, Stavanger University Hospital, Stavanger, Norway

¹⁴Department of Clinical Medicine, University of Bergen, Bergen, Hordaland, Norway

¹⁵Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway

¹⁶Clinic of Anaesthesia and Intensive Care, St. Olav University Hospital, Trondheim, Norway

Acknowledgements The authors wish to thank all the local data processors and all former members of the steering committee; Petter Andreas Steen, Mads Gilbert, Kaare Harald Bønå, Are Helseth and Per Halvorsen.

Contributor IBMT: contributed in the conception, planning, design, acquisition of data, analysis and interpretation of data, first draft of article, revision of draft and approval of final manuscript. KA-K: contributed in the design, acquisition of data, analysis and interpretation of data, revision of draft and approval of final manuscript. L-JA: contributed in the conception, planning, design, revision of draft and approval of final manuscript. SB: contributed in the conception, planning, revision of draft and approval of final manuscript. A-KH: contributed in the design, revision of draft and approval of final manuscript. BH: contributed in the conception, planning, design, revision of draft and approval of final manuscript. JL: contributed in the conception, planning, design, revision of draft and approval of final manuscript. ALL: contributed in the conception, planning, design, revision of draft and approval of final manuscript. TWL: contributed in the design, revision of draft and approval of final manuscript. JEN: contributed in the conception, planning, design, revision of draft and approval of final manuscript. TMO: contributed in the design, revision of draft and approval of final manuscript. EIS: contributed in the conception, planning, design, revision of draft and approval of final manuscript. EIS: contributed in the conception, planning, design, analysis and interpretation of data, revision of draft and approval of final manuscript. JK-J: contributed in the

conception, planning, design, acquisition of data, analysis and interpretation of data, first draft of article, revision of draft and approval of final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Map disclaimer The depiction of boundaries on this map does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. This map is provided without any warranty of any kind, either express or implied.

Competing interests There are no competing interests to report for the submitted work. TMO reports grants from Zoll Foundation and grants from Laerdal foundation, outside the submitted work. TWL reports paid and unpaid consultant work for Laerdal Medical and the Norwegian Air Ambulance Foundation outside the submitted work.

Patient consent for publication Not required.

Ethics approval This study was approved by Regional Ethical Committee (REK 60984) and data handling was done in accordance with recommendations from the local Data Protection Officer at Oslo University Hospital (20/01070).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. The data that support the findings of this study are available from the Norwegian Cardiac Arrest Registry through an application to the Norwegian Institute of Public Health. Restrictions apply to the availability of these data, as only aggregated data can be shared unless there is an approval from an ethical committee and the steering committee of the registry.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Ingvild Beathe Myrhaugen Tjelmeland <http://orcid.org/0000-0003-0362-1008>

REFERENCES

- 1 Meaney PA, Bobrow BJ, Mancini ME, *et al*. Cardiopulmonary resuscitation quality: [corrected] improving cardiac resuscitation outcomes both inside and outside the hospital: a consensus statement from the American Heart Association. *Circulation* 2013;128:417–35.
- 2 Hazinski MF, Nolan JP, Billi JE, *et al*. Part 1: Executive summary: 2010 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Circulation* 2010;122:S250–75.
- 3 Global Resuscitation Alliance. Secondary global resuscitation alliance 2017. Available: <https://www.globalresuscitationalliance.org> [Accessed 21 Apr 2020].
- 4 Gräsner J-T, Whent J, Herlitz J, *et al*. Survival after out-of-hospital cardiac arrest in Europe - Results of the EuReCa TWO study. *Resuscitation* 2020;148:218–26.
- 5 Beck B, Bray J, Cameron P, *et al*. Regional variation in the characteristics, incidence and outcomes of out-of-hospital cardiac arrest in Australia and New Zealand: results from the Aus-ROC Epistery. *Resuscitation* 2018;126:49–57.
- 6 Årsrapport HJ. *Yearly report of 2017 from the Swedish cardiac arrest registry*. 2018. Svenska Hjärt-Lungräddningsregistret, 2017.
- 7 Lindner TW, Søreide E, Nilsen OB, *et al*. Good outcome in every fourth resuscitation attempt is achievable—an Utstein template report from the Stavanger region. *Resuscitation* 2011;82:1508–13.
- 8 Nilsen JML, Haug B, Rasmussen JR, Clinic of Emergency Medicine, University Hosp of North Norway, Tromsø, Norway. Doubled survival from out-of-hospital cardiac arrest in a rural community in North-Norway following implementation of an aggressive chest pain protocol with early prehospital thrombolysis for STEMI. *Circulation* 2011.
- 9 Helse- og omsorgsdepartementet. *Forskrift Om innsamling OG behandling AV helseopplysninger I Nasjonalt register over hjerte- OG karlidelser (personal health data filing system act)*. Oslo: Helse- og omsorgsdepartementet, 2014.

- 10 Perkins GD, Jacobs IG, Nadkarni VM, *et al.* Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein resuscitation registry templates for out-of-hospital cardiac arrest: a statement for healthcare professionals from a task force of the International liaison Committee on resuscitation (American heart association, European resuscitation Council, Australian and New Zealand Council on resuscitation, heart and stroke Foundation of Canada, InterAmerican heart Foundation, resuscitation Council of southern Africa, resuscitation Council of Asia); and the American heart association emergency cardiovascular care Committee and the Council on cardiopulmonary, critical care, perioperative and resuscitation. *Resuscitation* 2015;96:328–40.
- 11 R Foundation for Statistical Computing. A language and environment for statistical computing 2017.
- 12 Tjomsland N, Baskett P, Laerdal AS, Asmund S, Laerdal. *Resuscitation* 2002;53:115–9.
- 13 The Directorate of Health. *Saving lives together*. helsedirektoratet, 2018.
- 14 Maurer H, Masterson S, Tjelmeland IB, *et al.* When is a bystander not a bystander any more? a European survey. *Resuscitation* 2019;136:78–84.
- 15 Beck B, Bray JE, Smith K, *et al.* Description of the ambulance services participating in the Aus-ROC Australian and New Zealand out-of-hospital cardiac arrest Epistry. *Emerg Med Australas* 2016;28:673–83.
- 16 Ringgren KB, Christensen HC, Schønau L. *Dansk hjertestopregister 2018 Danish cardiac arrest registry*, 2019.
- 17 Strömsöe A, Svensson L, Axelsson Å B, *et al.* Validity of reported data in the Swedish cardiac arrest register in selected parts in Sweden. *Resuscitation* 2013;84:952–6.
- 18 Becker LB, Smith DW, Rhodes KV. Incidence of cardiac arrest: a neglected factor in evaluating survival rates. *Ann Emerg Med* 1993;22:86–91.