Supplemental material

BMJ Publishing Group Limited (BMJ) disclaims all liability and responsibility arising from any reliance placed on this supplemental material which has been supplied by the author(s)

BMJ Open

Supplementary Table. Socio-demographic and clinical characteristics within sub-groups

Socio-demographic and clinical characteristics within sub-groups (percentages)

Socio-demo	Centre 1 Observational (N=70)	Centre 1 Experimental (N=100)	Centre 2 Experimental (N=100)	Centre 3 Experimental (N=107)	Test	*p*-Value
Refusal rate	19%	19%	14%	20%	$\chi^2 = 1.78$	$p = 0.62$
Employment Status					$\chi^2 = 2.64$	$p = 0.85$
Employed	41%	40%	46%	46%		
Non-employed	24%	28%	29%	27%		
Retired	34%	32%	25%	27%		
Marital Status					$\chi^2 = 2.93$	$p = 0.40$
Married/Cohabitant	63%	70%	73%	64%		
Single (Widowed/Divorced)	37%	30%	27%	36%		
Education					$\chi^2 = 4.39$	$p = 0.22$
Primary/Secondary School	43%	54%	48%	40%		
High School/University	57%	46%	52%	60%		
Age (mean ± standard deviation)	58 ± 9.7	58 ± 10.5	57 ± 9.5	55 ± 10.7	$F = 1.50$	$p = 0.22$
Therapy§						
Chemotherapy prescribed	27%	36%	35%	35%	$\chi^2 = 1.66$	$p = 0.64$
Hormonal therapy prescribed	67%	76%	84%	74%	$\chi^2 = 6.96$	$p = 0.07$
Radiotherapy prescribed	66%	62%	60%	77%	$\chi^2 = 7.54$	$p = 0.06$
Immunotherapy prescribed	9%	8%	4%	7%	$\chi^2 = 1.91$	$p = 0.59$
Cancer stage ***					**$\chi^2 = 21.81$**	**$p = 0.01$**
Stage 0	38%	20%	31%	21%		
Stage I	45%	55%	36%	44%		
Stage II	8%	17%	22%	31%		
Stage III	9%	7%	10%	5%		

Significant parameters are written in bold
§ more than a therapy could be prescribed to each patient
*** 22 missing info