

Supplement

Table S1: Comparison of Indian population and study sample after applying weights

Age group	Proportion of 25 to 90 year old population	
	India (Census 2011)	Survey
25-29	16.9	16.9
30-64	72.8	72.5
65-90	10.3	10.6

Table S2: Differential impact of financial difficulties on health care service provision and health and wellbeing at different severities of lockdown

Outcomes	Lockdown	
	Most severe	Other
	OR (95%CI)	OR (95%CI)
Non-availability	4.32 (2.21, 8.43)	3.31 (2.23, 4.92)
Non-affordability	4.06 (1.80, 9.17)	7.17 (5.06, 10.15)
Non-accessibility	11.81 (6.24, 22.33)	1.82 (1.34, 2.46)
Inadequacy	7.00 (3.03, 16.18)	4.48 (3.07, 6.54)
Inappropriate	10.33 (2.95, 36.21)	3.26 (2.15, 4.94)
Negative Patient experience	1.06 (0.12, 9.18)	6.65 (3.37, 13.14)
Discontinuity of care	15.22 (7.58, 30.57)	2.94 (2.08, 4.15)
Poor Health	1.29 (0.69, 2.41)	2.69 (1.91, 3.79)
WHO-5 Screening for depression	4.69 (1.72, 12.77)	4.54 (3.31, 6.24)
Emotional loneliness	0.77 (0.35, 1.68)	2.21 (1.52, 3.20)
Social loneliness	2.20 (0.86, 5.64)	2.22 (1.47, 3.36)
Loneliness emotional and/or social	4.00 (1.36, 11.73)	2.18 (1.24, 3.82)