

Figure S1: Flow of study participants in the first follow-up of the Canadian Longitudinal Study on Aging (CLSA)




Figure S2: Percentage of adults who indicated their last used e-cigarette contained nicotine by demographic factors and cigarette smoking status

Table S1: Comparison of participants with acceptable grade on spirometry and those with missing spirometry data

	Included in the analysis n (%)	Not included in the analysis* n (%)	p-value
Age			<.0001
45-54	3569 (17.5)	503 (12.2)	
55-64	7132 (34.9)	1160 (28.2)	
65-74	5990 (29.3)	1192 (28.9)	
75-85	3727 (18.3)	1266 (30.7)	
Sex			<.0001
Male	9769 (47.8)	2249 (54.6)	
Female	10649 (52.2)	1872 (45.4)	
Education			<.0001
No post-secondary education	868 (4.3)	282 (6.8)	
Diploma/certificate below bachelor's	1766 (8.7)	438 (10.7)	
Bachelor's degree	1423 (7.0)	349 (8.5)	
Above bachelor's degree	16334 (80.1)	3043 (74.0)	
Annual household income			<.0001
< \$20,000	740 (3.8)	274 (7.3)	
\$20,000 - <50,000	3850 (20.0)	978 (26.2)	
\$50,000 - <100,000	7001 (36.33)	1335 (35.8)	
\$100,000 - <150,000	3995 (20.7)	631 (16.9)	
≥ \$150,000	3685 (19.1)	516 (13.8)	
Cigarette smoking			<.0001
Never	9929 (48.9)	1967 (48.1)	
0-14 pack-years	6418 (31.6)	1208 (29.5)	
≥ 15 pack-years	3954 (19.5)	919 (22.4)	
E-cigarette ever use			0.0569
Yes	927 (4.5)	215 (5.2)	
No	19489 (95.5)	3899 (94.8)	

*Participants with contraindications are excluded. Proportions in the table are unweighted estimates.