BMJ Open Common elements of service delivery models that optimise quality of life and health service use among older people with advanced progressive conditions: a tertiary systematic review

Joanne Bayly ^{1,2}, Anna E Bone,¹ Clare Ellis-Smith,¹ India Tunnard,¹ Shuja Yaqub,¹ Deokhee Yi ¹, Kennedy B Nkhoma ³, Amelia Cook,¹ Sarah Combes,^{3,4,5} Sabrina Bajwah,¹ Richard Harding ¹, Caroline Nicholson,^{4,5} Charles Normand,^{1,6} Shalini Ahuja,⁷ Pamela Turrillas,¹ Yoshiyuki Kizawa,⁸ Tatsuya Morita,⁹ Nanako Nishiyama ¹⁰, Satoru Tsuneto,¹¹ Paul Ong,¹² Irene J Higginson,¹ Catherine J Evans ^{1,13}, Matthew Maddocks ¹

To cite: Bayly J, Bone AE, Ellis-Smith C, *et al.* Common elements of service delivery models that optimise quality of life and health service use among older people with advanced progressive conditions: a tertiary systematic review. *BMJ Open* 2021;**11**:e048417. doi:10.1136/bmjopen-2020-048417

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-048417>).

CJE and MM are joint senior authors.

Received 04 January 2021
Accepted 05 November 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Matthew Maddocks;
matthew.maddocks@kcl.ac.uk

ABSTRACT

Introduction Health and social care services worldwide need to support ageing populations to live well with advanced progressive conditions while adapting to functional decline and finitude. We aimed to identify and map common elements of effective geriatric and palliative care services and consider their scalability and generalisability to high, middle and low-income countries.

Methods Tertiary systematic review (Cochrane Database of Systematic Reviews, CINAHL, Embase, January 2000–October 2019) of studies in geriatric or palliative care that demonstrated improved quality of life and/or health service use outcomes among older people with advanced progressive conditions. Using frameworks for health system analysis, service elements were identified. We used a staged, iterative process to develop a ‘common components’ logic model and consulted experts in geriatric or palliative care from high, middle and low-income countries on its scalability.

Results 78 studies (59 geriatric and 19 palliative) spanning all WHO regions were included. Data were available from 17 739 participants. Nearly half the studies recruited patients with heart failure (n=36) and one-third recruited patients with mixed diagnoses (n=26). Common service elements (≥80% of studies) included collaborative working, ongoing assessment, active patient participation, patient/family education and patient self-management. Effective services incorporated patient engagement, patient goal-driven care and the centrality of patient needs. Stakeholders (n=20) emphasised that wider implementation of such services would require access to skilled, multidisciplinary teams with sufficient resource to meet patients’ needs. Identified barriers to scalability included the political and societal will to invest in and prioritise palliative and geriatric care for older people, alongside geographical and socioeconomic factors.

Conclusion Our logic model combines elements of effective services to achieve optimal quality of life and health service use among older people with advanced

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ We draw on and synthesise a diverse evidence base of geriatric and palliative care for older people with progressive advanced conditions across the globe.
- ⇒ The review was conducted by a multidisciplinary and international group representing broad methodological expertise and perspectives.
- ⇒ Our common components logic model is a recombination of effective service elements. However, we were unable to assert how outcomes may be influenced by different combinations of components and their interactions.
- ⇒ Our stakeholder consultation identified significant barriers to scalability where country health budgets cannot meet the growing population need, and where multidisciplinary care is not available.

progressive conditions. The model transcends current best practice in geriatric and palliative care and applies across the care continuum, from prevention of functional decline to end-of-life care.

PROSPERO registration number CRD42020150252.

INTRODUCTION

Globally, more people are living into old age¹ with the largest proportional increase occurring in those 80 years and above.^{2,3} By 2050, 80% of older people will live in low-income and middle-income countries (LMICs).⁴ The concomitant risks of multimorbidity and/or frailty⁵ mean more people experience a trajectory of prolonged and uncertain functional decline. Health and social care needs and their impact on physical functioning are more heterogeneous¹ in older populations,

shaped by multiple interacting factors related to the individual and their environment. These population changes bring new societal challenges related to health and social care policy, spending, workforce and security, regardless of the developmental context.

The WHO Member States' commitment to achieve Universal Health Coverage (UHC) by 2030 provides an opportunity to plan health and social care delivery for the future. Palliative care has recently been included as an essential service that is fundamental to achieving UHC.⁶ While prevention remains a priority across the health continuum, a shift in health systems is needed to balance disease-modifying interventions with services where improving quality of life is the main goal of care. In older people with advanced (incurable) and progressive diseases, health systems must align to support the dual priorities of living well while adapting to a gradual decline in function. Access to appropriate care and support is recognised as a basic human right,⁷ yet access varies according to socioeconomic and geographical variables.^{8,9} Budget constraints require maximum value from the resources used to improve outcomes.¹⁰ The importance of integrated working across services is consistently advocated in global guidance on health service provision for advanced disease¹¹ and older people.¹²

Our previous meta-review outlined two service delivery models for older people towards the end of life; 'integrated geriatric care' and 'integrated palliative care'.¹² Both showed potential to improve quality of life and patterns of health service use, but with differing emphasis on either function or symptoms and concerns. Our findings underscored the imperative of access to services based on the likelihood of benefit and integration of care using comprehensive assessment, case management and/or collaborative working.¹² However, use of systematic reviews as the unit of analysis prevented a detailed description of service model elements, and suppressed the heterogeneity across the primary studies.

This review aimed to detail service delivery models that optimise quality of life and health services use for older people aged 60 years and over with advanced progressive health conditions. We defined 'advanced' to include disease stage, people described as in their last 1 or 2 years of life or people accessing a service typically used in advanced disease stage, such as nursing home or palliative care. Our objectives were to: (1) identify and map common elements of effective service delivery models within primary studies; (2) outline the similarities and differences across models of geriatric care or palliative care and (3) consider the scalability of effective models, attending to implementation and economic requirements.

METHOD

Study design

This review builds on our previous meta-review, where the methods are described in detail.¹² Here, we conducted a

tertiary review of individual empirical studies ('primary studies') from the meta-review.¹² This was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis.¹³ We then used logic modelling¹⁴ and a stakeholder consultation to support the analysis and interpretation¹⁵ of findings. This study was registered on PROSPERO prior to data extraction.

Patient and public involvement

Patients and members of the public were not involved in the design, conduct, reporting or dissemination of this research.

Search strategy

For the purposes of this tertiary review, in October 2019 we updated our original meta-review search to identify systematic reviews that included a meta-analysis that demonstrated overall effectiveness on at least one outcome for quality of life (including symptom burden and function) and/or health service use outcome. The systematic review eligibility criteria and search terms are reported in online supplemental materials 1 and 2. From the eligible systematic reviews, we identified primary studies with evidence of effect on our selected outcomes of quality of life and/or health service use. Inclusion criteria for primary studies comprised: (1) experimental study design; (2) contributed data to meta-analysis and (3) reported a point estimate of effect in the same direction as the meta-analysis. One reviewer (JB) evaluated all systematic reviews and primary studies for eligibility and a second (MM, AB or CE-S) double-screened studies, with inconsistencies resolved by consensus. Duplicate primary studies were identified and removed.

Data extraction

Data on study population, outcomes and context were extracted. Service delivery models were classified as either integrated geriatric or palliative care. Data identification and extraction was informed by a framework for health-care systems analysis, the checklist CATWOE (customers, actors, transformation processes, world view, owner, environmental constraints).^{16,17} For each CATWOE domain, (eg, customers, actors), a list of service elements was identified. Service elements were categorised as present, absent or unclear by two individuals (from JB, AB, CE-S, SY, DY, NK, SB, CE, MM) and reviewed as a team. Identification of the elements for each CATWOE domain was, informed by the Template for Intervention Description and Replication checklist for complex health service interventions¹⁸ and prior studies on geriatric,¹⁹ integrated,²⁰ transitional²¹ and palliative care²² online supplemental material 3 details the elements for each CATWOE domain.

Quality appraisal

The methodological quality of systematic reviews and primary studies was appraised using the A MeaSurement Tool to Assess systematic Reviews (AMSTAR) tool²³

and Cochrane Risk of Bias Tool, respectively.²⁴ We used the quality appraisal in the systematic reviews when the Cochrane Risk of Bias Tool was used, otherwise assessment was by two researchers (JB and IT). We did not exclude studies from analysis based on quality.

Development of logic model

We used a staged and iterative approach following Rohwer *et al*'s guidance on logic models for complex health interventions¹⁴ incorporating analysis of extracted data followed by a stakeholder consultation.

The frequency and proportion of service elements^{16 17} was summarised overall and separately for integrated geriatric and palliative care models. The proportion was calculated using studies where the element was categorised as present or absent. We mapped service elements present in $\geq 50\%$ of integrated geriatric and/or palliative care studies by CATWOE domain to existing logic model templates¹⁴ (see online supplemental material 4). To compare the presence of service elements between integrated geriatric and palliative care models we conducted χ^2 tests (or Fisher's exact tests where counts were low).

We appraised the potential for the common components of effective interventions to be generalised and scalable, defined as the ability 'to be expanded under real world conditions to reach a greater proportion of the eligible population while retaining effectiveness'.²⁵

We shared an interim logic model and consulted a purposive sample of healthcare researchers, clinical-academics

and clinicians from high, middle and low-income countries with expertise in either geriatric or palliative care, hospital or community based. We used the Context and Implementation of Complex Interventions Framework (CICI) to develop a response form with free-text open questions on the barriers and facilitators to providing the elements of care, for their respective country and healthcare settings.²⁶ The CICI framework domains informed the identification and collation of the narrative responses on the context and implementation considerations. We developed the logic model by synthesising the findings from the tertiary review and the stakeholder consultation, using an iterative process of team discussion and consensus.¹⁴

RESULTS

Study retrieval

Ten systematic reviews met eligibility, seven from the meta-review^{27–33} and three from the updated search.^{34–36}

The reviews reported 180 potentially eligible studies, of which 47 were duplicates. Of the 133 remaining studies, 78 met eligibility (figure 1).

Characteristics of included studies

Of the 78 included studies, 59 were categorised as integrated geriatric care and 19 as integrated palliative care (table 1 and online supplemental material 5). All WHO regions were represented, though studies were

Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart for selection of primary studies.

Table 1 Summary characteristics of included studies N=78

Variable		Frequency n (%)		
		All n=78	Geriatric n=59	Palliative n=19
WHO region	Americas	46 (59)	36 (61)	10 (53)
	Europe	22 (28)	16 (27)	6 (32)
	Southeast Asia	3 (4)	2 (3)	1 (5)
	West Pacific	6 (8)	5 (8)	1 (5)
	Africa	1 (1)	1 (2)	0
Country income status	High	75 (96)	17 (29)	58
	Upper—middle	2 (3)	1 (2)	1
	Lower—middle	1 (1)	0	1
	Low	0	0	0
Population by main diagnosis	Heart failure	36 (46)	32 (54)	4 (21)
	No main diagnosis	23 (29)	23 (39)	0
	Cancer	14 (18)	2 (3)	12 (63)
	Single	4 (5)	1 (2)	3 (16)
	Mixed	10 (13)	1 (2)	9 (47)
	Heart failure +diabetes	1 (1)	1 (2)	0
	Heart failure +depression	1 (1)	1 (2)	0
	Multiple sclerosis	1 (1)	0	1 (5)
	Mixed diagnosis (COPD, cancer, HF, ILD, MND)	1 (1)	0	1 (5)
	HIV infection	1 (1)	0	1 (5)
Population by referral criteria	People with heart failure	38 (49)	34 (58)	4 (21)
	People with acute episode of illness	17 (22)	17 (29)	0
	People with advanced cancer	13 (17)	2 (3)	11(58)
	Older people (varied age ranges)	6 (8)	6 (10)	0
	People with HIV	1 (1)	0	1 (5)
	People with multiple sclerosis	1 (1)	0	1 (5)
	Advanced mixed diagnoses	1 (1)	0	1 (5)
	People with cancer commencing chemotherapy	1 (1)	1 (2)	1 (5)
Health organisation funding	State funded health organisation	35 (45)	26 (44)	9 (47)
	For profit health organisation	37 (47)	28 (47)	9 (47)
	Non-profit health organisation	6 (8)	5 (8)	1 (5)
Care setting	Mixed settings	29 (37)	20 (34)	9 (47)
	Hospital inpatients and home	6 (8)	6 (10)	0
	Hospital inpatients and outpatients	5 (6)	5 (8)	0
	Hospital outpatients and home	10 (13)	4 (7)	6 (32)
	Hospital inpatients, outpatients and home	7 (9)	4 (7)	3 (16)
	Hospital emergency room and home	1 (1)	1 (2)	0
	Home	16 (21)	13 (22)	3 (16)
	Hospital outpatients	15 (19)	9 (15)	6 (32)
	Hospital inpatients	13 (17)	12 (20)	1 (5)
	Community settings	3 (4)	3 (5)	0

COPD, chronic obstructive pulmonary disease; HF, heart failure; ILD, interstitial lung disease; MND, motor neuron disease.

predominantly from the North American region of the Americas (n=46) or Europe (n=22), with fewer from Western Pacific (n=6), Southeast Asia (n=3) and only a

single study from Africa. Most studies were from high-income countries (n=75). The number of study participants ranged from 20 to 1632, with data available from

17739 participants overall. Nearly half of all studies recruited patients with heart failure (n=36) and one-third recruited patients with mixed diagnoses (n=26). Palliative care studies most often recruited by cancer diagnosis (n=12). Study interventions were delivered across multiple care settings (n=31), in participants' homes (n=15) or in hospital (outpatients n=14; inpatients n=12) (table 1).

Quality appraisal

The 10 systematic reviews were assessed as of moderate quality (online supplemental material 6). Primary studies were assessed as low-to-moderate risk of bias overall (online supplemental material 6). Where high risk of bias was found, this most frequently related to challenges of blinding participants and personnel leading to possible performance and detection bias. Risk of bias tended to be lower for palliative care compared with geriatric care studies (online supplemental material 7).

Service delivery elements

Services most frequently used collaborative working and case management to support integrated working between professionals (table 2). Patient/family education was present in all studies. Other common elements, present in ≥80% of studies were ongoing assessment, active patient participation and evidence of patient engagement in their care. The least common elements overall were: bereavement support; 24-hour home visits or access to physicians; links to residential hospice facilities and joint provision of care across health and social care services. No studies reported delivering interventions in residential care/nursing homes or use of volunteers.

Comparing between integrated geriatric and palliative care, palliative care interventions had a higher frequency of end-of-life expertise and training, professional psychosocial support, spiritual support and physician home visits. In contrast geriatric care interventions more often featured early rehabilitation assessment and self-management, though the differences were not statistically significant (table 2).

Service delivery agents

All interventions were delivered by qualified health-care professionals (staff who have received nationally recognised and regulated training and education), most often working in multidisciplinary teams. Over 90% of studies involved trained medical and nursing clinicians and 59% involved members of the wider healthcare team, including physiotherapists, occupational therapists and social workers. Geriatric care services were delivered by physicians from geriatrics, cardiology and general practice, whereas palliative care services involved physicians from cardiology, neurology, respiratory medicine, oncology, psychiatry, primary care and palliative medicine. While involvement of physiotherapists was reported across all studies, fewer occupational therapists and

dietitians were reported in those from palliative care. No studies reported the involvement of volunteers (table 3).

Service outcomes including costs

Forty-five studies (58%) were included based on an effect on quality of life alone. Fifty-seven studies (73%) used a disease or population specific measure to quantify quality of life (online supplemental material 5) and five studies (6%) employed the EuroQoL 5-Dimension (EQ-5D). Thirty-three studies (42%) reported utilisation of acute care services (eg, hospital admission, readmission after discharge) or community care services and 20 studies (26%) calculated costs of health services utilisation. Only a minority (n=12%/15%) demonstrated an effect on both quality of life and health service use, all of which were geriatric care studies. No study used costs and EQ-5D to generate information required for health economic decision-making (table 4).

Common components logic model

The interim logic model highlighted key elements present in the majority (<80%) of included studies.

Elements more common in integrated palliative care compared with geriatric care studies were; professional psychosocial support, advance care planning, caregiver engagement, joint decision-making and expert consultation with other providers. Elements more common in geriatric care studies included a social worker or dietitian as a delivery agent and care planning organised around the service, delivering the same intervention to all patients but with individual tailoring (figure 2).

Elements more common in geriatric care studies included a social worker or dietitian as a delivery agent and care planning organised around the service with the same intervention being delivered to all patients with individual tailoring (figure 2).

Stakeholder perspectives on scalability

The context and implementation considerations identified from the stakeholder responses were incorporated into the logic model (figure 2). Stakeholders (n=20) contributed views from high-income countries (n=12) (UK, Japan, Taiwan, Portugal and Chile) and LMICs (n=8) (Uganda, Malawi, South Africa, Ghana, Zimbabwe, China, India and Bangladesh) contributed views. They described increasing patient complexity with rapid population ageing and the associated rise in multimorbidity, frailty and dementia. This raised particular challenges in LMICs where health services have historically focused on prevention and management of infectious diseases and where there has been a recent increased burden of non-communicable disease. Specialist services being based in major city hospitals were described as a barrier to providing care to rural populations. Recruiting, training and retaining skilled staff to work in rural areas and having a multidisciplinary team including allied health professionals and specialist doctors and nurses was considered infeasible for many rural areas.

Table 2 Service delivery model elements N=78

	All n (%)	Geriatric n (%)	Palliative n (%)	Sig†
<i>Method of supporting integrated working</i>				
Collaborative working	64 (82)	46 (78)	18 (95)	0.17*
Case management	61 (78)	46 (78)	15 (79)	1.00*
Comprehensive assessment	51 (65)	36 (68)	15 (79)	0.36
<i>Actors-workforce</i>				
Professional education	76 (100)	58 (100)	18 (100)	1.00
MDT care	54 (72)	42 (73)	12 (71)	1.00*
Rehabilitation expertise training	34 (50)	27 (50)	7 (50)	1.00
End-of-life expertise training	18 (25)	1 (2)	17 (90)	<0.001*
<i>Transformation-service model elements/components</i>				
Patient family education	60 (100)	49 (100)	11 (100)	0.93
Medication review	51 (80)	40 (77)	11 (92)	0.43*
Self-management	48 (80)	41 (84)	7 (64)	0.21*
Systematic risk screening	47 (69)	37 (70)	10 (67)	1.00*
Contact with GP or attending doctor	46 (68)	33 (65)	13 (77)	0.37
Practical support	41 (68)	34 (69)	7 (64)	0.73*
Medical intervention	52 (67)	39 (66)	13 (68)	0.85
Individualised MDT plan	40 (61)	29 (59)	11 (65)	0.69
Complex/medication management	37 (58)	30 (59)	7 (54)	0.75
Discharge planning	36 (52)	29 (55)	7 (44)	0.44
Professional psychosocial support	38 (51)	26 (44)	12 (80)	0.01
Team case rounds	25 (40)	18 (37)	7 (50)	0.37
Early rehab assessment	25 (38)	21 (40)	4 (29)	0.54
Advanced care planning	23 (30)	9 (16)	14 (78)	<0.001
Emergency response plan	15 (21)	12 (22)	3 (20)	1.00*
Spiritual support	13 (18)	2 (3)	11 (79)	<0.001*
Bereavement support	4 (5)	0 (0)	4 (25)	0.002*
<i>Transformation-mode of delivery</i>				
Ongoing assessment	66 (87)	50 (86)	16 (89)	1.00*
Face-to-face and telephone	41 (53)	31 (53)	10 (53)	0.10
Face-to-face interaction	31 (40)	23 (39)	8 (42)	0.81
Access to inpatient beds	21 (30)	18 (32)	3 (21)	0.53*
Physician home visits	11 (15)	4 (7)	7 (37)	0.04*
24-hour physician access	6 (10)	5 (11)	1 (7)	1.00*
Telephone only	5 (6)	4 (7)	1 (5)	1.00*
24-hour home visits	1 (1)	1 (2)	0 (0)	1.00*
Online only	1 (1)	1 (2)	0 (0)	0.10*
<i>Transformation-operational tools and guidance to support practice</i>				
Standard comprehensive assessment	38 (59)	26 (55)	12 (71)	0.27
<i>Worldview-methods of integrated working</i>				
Link to hospital	57 (78)	41 (72)	16 (100)	0.02*
Expert consult with other providers	40 (58)	24 (45)	16 (100)	<0.001
Link between community services	31 (50)	22 (45)	9 (69)	0.12
Joint provision-health and social care	7 (10)	4 (7)	3 (20)	0.16*
Link to residential hospice	5 (7)	1 (2)	4 (27)	0.005*

Continued

Table 2 Continued

	All n (%)	Geriatric n (%)	Palliative n (%)	Sig†
<i>Worldview-conceptual model</i>				
Patient engagement	71 (99)	53 (98)	18 (100)	1.00*
Active patient participation	67 (99)	50 (98)	17 (100)	1.00*
Centrality of patient needs	64 (91)	46 (89)	18 (100)	0.33*
Patient goal driven care	56 (81)	40 (77)	16 (94)	0.16*
Ongoing/continuous care	46 (67)	33 (62)	13 (81)	0.16
Joint decision-making	38 (69)	25 (61)	13 (93)	0.04*
Service driven care planning	38 (54)	34 (65)	4 (21)	0.001*
Needs and benefit-driven care planning	33 (46)	18 (35)	15 (79)	0.001
Caregiver engagement	32 (55)	22 (50)	10 (71)	0.16

*Fisher's exact test.
†Sig=significance for difference in presence of service delivery element between geriatric and palliative care studies.
GP, General Practitioner; MDT, Multidisciplinary Team.

Stakeholders from LMICs considered that overall health budgets were inadequate to meet the population need, and multidisciplinary care was considered unaffordable. The voluntary sector was often seen as important to augment publicly funded services. In some contexts, continuity of care is impeded when individually funded services compete for resources rather than collaborate. There are challenges to multidisciplinary working in systems where health workers receive payment directly from patients, as this was considered a financial disincentive to making referrals for expert consultation. Social deprivation was cited as an important barrier to accessing care, especially in health systems with out-of-pocket expenses or private insurance.

Stakeholders described how cultural norms influence care provision. Death denying attitudes in some cultures influence uptake of palliative care services. Some countries have limited recognition or respect for the specialties of palliative care and geriatric care. The role of the family and the health system to provide care was identified to vary across countries influenced by cultural beliefs such as filial piety, gender-related norms and changing intergenerational family structures. Acknowledging faith and religion were identified as factors supporting the delivery of individualised care aligned with spiritual needs in hospice and nursing homes.

Increasing education levels and internet access were identified as factors that are changing patient and family participation in joint decision-making. Finally, stakeholders recognised an increasing political will to invest in services for older people supported by a growing public and research agenda and established regulatory frameworks. However, this did not always equate to increases in funding. A lack of policies and clinical governance for specialist palliative and geriatric care was reported as a problem, like legal restrictions on opioid prescribing limiting effective medication management of pain.

DISCUSSION

We used rigorous methods to detail service delivery models that optimise quality of life and health service use outcomes among older people with advanced progressive conditions. Effective services commonly used collaborative working between professionals and specialties, comprehensive and ongoing assessment, patient/family education and active patient participation. Aligned to this, effective services consistently incorporated patient engagement, patient goal-driven care and the centrality of patient needs in care delivery. Our logic model encompasses a breadth of elements that aim to 'protect' (discharge planning and falls prevention programmes), 'reactivate' (disease management, self-management and exercise programmes), 'compensate' (symptom management, support with capabilities for activities of daily living) and 'support' (enhancing social assets and provision of home care). Such practices may together support older people to maintain intrinsic capacity and functional ability³⁷ and to compress functional decline across the life course.^{38 39} This broad focus, together with consideration of social factors, extends health and social care beyond provision at the point of decline to meet the dual priorities of living well while adapting to a gradual functional decline.¹

This review has several strengths. It was conducted by a large multidisciplinary team with a range of methodological expertise and representation from many regions of the world. We synthesised a diverse literature with studies across different patient populations and needs across the trajectory of advanced disease. We used recognised frameworks to categorise studies, extract data and consult with stakeholders in order to develop a visual logic model applicable to different international settings. There are some limitations to consider. Data on study context is limited to country, country income status and the system for funding healthcare. Further information to support

Table 3 Service delivery model agents

Delivery agent	All n (%)	Geriatric n (%)	Palliative n (%)	Sig†
<i>Physicians</i>				
Geriatrician	14 (18)	14 (24)	0 (0)	0.02
Cardiologist	15 (19)	12 (20)	3 (16)	1.0
Palliative care physician	12 (15)	0 (0)	12 (63)	<0.001*
Neurologist	1 (1)	0 (0)	1 (5)	0.24*
Respiratory physician	1 (1)	0 (0)	1 (5)	0.24*
Oncologist	4 (5)	0 (0)	4 (21)	0.001*
Psychiatrist	2 (3)	0 (0)	2 (11)	0.06*
Physician	18 (23)	17 (29)	1 (5)	0.06*
Primary care doctor (GP)	5 (6)	4 (7)	1 (5)	0.55*
Physician assistant	2 (3)	2 (3)	0 (0)	0.43*
<i>Nurses</i>				
Nurse	24 (31)	22 (37)	2 (11)	0.28
Advanced nurse practitioner	13 (17)	8 (14)	5 (26)	0.17*
Specialist cardiac nurse	12 (15)	10 (17)	2 (11)	0.40*
Primary care nurse	9 (8)	8 (14)	1 (5)	0.30*
Specialist geriatric nurse	6 (8)	6 (10)	0 (0)	0.18*
Case manager	5 (6)	3 (5)	2 (11)	0.35*
Specialist palliative nurse	4 (5)	1 (2)	3 (16)	0.43*
Specialist rehabilitation nurse	1 (1)	1 (2)	0 (0)	0.76*
Specialist HIV nurse	1 (1)	0 (0)	1 (5)	0.24*
Oncology nurse	1 (1)	0 (0)	1 (5)	0.24*
<i>Allied health professionals</i>				
Physiotherapist	23 (29)	17 (29)	6 (32)	0.85
Occupational therapist	14 (18)	12 (20)	2 (11)	0.28*
Dietitian	16 (21)	14 (24)	2 (11)	0.18*
Psychologist	9 (15)	6 (10)	3 (16)	0.38*
Pharmacologist/pharmacist	7 (9)	7 (12)	0 (0)	0.13*
Chaplain	4 (5)	1 (2)	3 (16)	0.43*
Audiologist	1 (1)	1 (2)	0 (0)	0.76*
Speech and language therapist	1 (1)	1 (2)	0 (0)	0.76*
<i>Social care</i>				
Social worker	21 (27)	17 (29)	4 (21)	0.51
Home care service manager	3 (4)	3 (5)	0 (0)	0.43*
Social assistant	4 (1)	3 (5)	1 (5)	0.68*
<i>Other professionals</i>				
Unspecified wider 'MDT'	11 (14)	9 (15)	2 (11)	0.47*
Exercise instructor	2 (3)	2 (3)	0 (0)	0.57*

*Fisher's exact test.

†Significance for difference in presence of service delivery element between geriatric and palliative care studies.

an evaluation of how interventions could be scaled and implemented would be valuable. Stakeholders identified limited applicability for some service elements, including multidisciplinary care, within low-income countries where health budgets cannot meet the growing

population need. Change beyond the health system, into education and health promotion, may be required to implement services that meet the challenge of rising incidence in diseases of ageing.⁴⁰ As in other reviews of complex interventions in this population,¹⁶ we were

Table 4 Number of studies reporting quality of life and health services use outcomes

		Health service use			Subtotal
		None	More than 1	1+and costs	
Quality of life	None	0	6	15	21
	More than 1	40	7	5	52
	1+ and EQ-5D	4	0	0	5
	Subtotal	45	13	20	78

EQ-5D, EuroQoL 5-Dimension.

unable to discern the specific mechanisms of action that make each component effective. In part this was linked to our data extraction framework. For example, we did not extract data on how interventions provided care across care boundaries during care transitions, yet elements including ongoing assessment and links between community services indicate this may have been occurring.

Our findings build on previous reviews. Bainbridge *et al*²⁹ found that ‘linkages with hospital,’ ‘multiprofessional teams’ and ‘end of life care expertise and training’ were critical to the delivery of end-of-life home care. In a review of integrated care for older people, Briggs *et al*²⁰ found that multidisciplinary teams, comprehensive assessment and case management were most frequently reported. We

show the importance of a capable workforce that works collaboratively across disciplinary boundaries, to provide comprehensive and ongoing multidimensional assessment. This model of care requires active patient engagement, participation and self-management with tailored care centred on the needs of individuals.⁴¹ It allows for a shared understanding between the person(when able and/or the family) and the team providing their care, facilitating joint decision-making that addresses their priorities in their context.⁴²

We also provide new insights into the range of health and social care providers associated with effective services in this population. Services were frequently delivered by multidisciplinary teams of health and

Figure 2 Common components logic model detailing effective service delivery models for older people with advanced progressive conditions .

social care professionals with formal training in core skills of comprehensive assessment, communication and symptom management. These teams can support people to self-manage a progressive condition and help people close to them to provide care. Investment in training and education is required to achieve greater coverage and ensure the skills base keeps up with the needs of this growing population. Uncoupling skills from professional roles and working towards a generalist skills set may be most beneficial. However, this should ideally be accompanied by access to specialists for ongoing support and supervision. Volunteers may provide additional support that supplements or enhances usual health and social care provision.^{43 44} The absence of volunteers in studies probably reflects the fact that most were conducted in high-income countries.

Service elements that we consider relevant to the target population but not brought forward to our logic model include joint provision across health and social care and early rehabilitation assessment. Neglecting social care can have a considerable negative effect on quality of life for older people, their family and friends and lead to increased patient and carer morbidity and mortality.⁴⁵ Integrated care should follow older people as they transition from acute to community care.⁴⁶ However workforce issues continue to influence the integration of health and social care delivery⁴⁷ as highlighted in our stakeholder consultation. Early rehabilitation assessment was detailed in only 40% and 19% of geriatric and palliative care studies, respectively. Given that maintaining independence, normality and social participation are high priorities for older people towards the end of life,⁴⁸ this was a surprising finding. It may relate to a focus on physical symptoms arising from advanced disease rather than functional needs, and the presumption that decline is an inevitability of disease progression.⁴⁹ The increasing prominence of rehabilitation in palliative care to challenge this misconception is therefore timely.^{50 51}

The logic model is a recombination of different services and we were unable to assert how effectiveness may be influenced by different combinations of components and their interactions. Consequently, the model remains untested as a whole.⁵² However the model can inform health and social care policy and support the conceptual and organisational development of services. We recommend that the clinical and cost-effectiveness of interventions, underpinned by our proposed model, should be tested in older people with multimorbidity based on need, rather than diagnostic condition, over longer trajectories and across care boundaries. Implications for policy are presented in [box 1](#).

CONCLUSION

Our logic model brings together common elements of interventions found to optimise quality of life and health service use among older people with advanced progressive conditions. These included collaborative working

Box 1 Implications for policy

- ⇒ Configure services for the whole trajectory of chronic progressive conditions up until the end of life and move away from a focus on acute episodes of care.
- ⇒ Plan and deliver education to drive provision of a capable workforce. A broad range of professional education courses and training in core skills of geriatric and palliative care, including comprehensive assessment, communication and symptom management specific to individual need is required.
- ⇒ Incentivise interdisciplinary and collaborative working between professional disciplines and across health and social care settings, to optimise high-quality individualised service provision and care co-ordination. This integrated care, when aligned to need rather than diagnostic condition, will increase the reach and impact of services and promote equitable access.
- ⇒ Enable robust evaluation by embedding routine outcome measurement in health and social care settings. These should include measures of intrinsic capacity, functional ability, symptom experience and quality of life. Measures should capture the changes in health and social well-being that are associated with the provision of high quality individualised care across the care continuum from protect to support and end-of-life care.

between professionals and specialties, ongoing assessment, active patient participation, patient/family education and patient self-management, while effective service delivery approaches consistently incorporated patient engagement, patient goal-driven care and the centrality of patient needs.

These elements transcend best practices in geriatric care and palliative care to optimise patient outcomes across the continuum, from prevention of functional decline to end-of-life care. The logic model serves as a useful resource for health systems looking to strengthen their response to population ageing. It can guide provision of health and social care that is aligned to the needs of this rapidly growing population. Such care should allow older people across the globe to live fully, with minimal suffering, and to die with dignity.

Author affiliations

¹Cicely Saunders Institute for Palliative Care, Policy and Rehabilitation, King's College London, London, UK

²St Barnabas Hospice, Worthing, UK

³Florence Nightingale Faculty of Nursing Midwifery and Palliative Care, King's College London, London, UK

⁴St Christopher's Hospice, London, UK

⁵University of Surrey Faculty of Health and Medical Sciences, Guildford, UK

⁶Centre for Health Policy and Management, The University of Dublin Trinity College, Dublin, Ireland

⁷Health Service and Population Research Department, King's College London Institute of Psychiatry Psychology and Neuroscience, London, UK

⁸Department of Palliative Medicine, Kobe University, Kobe, Japan

⁹Palliative and Supportive Care Division, Seirei Mikatahara Hospital, Hamamatsu, Japan

¹⁰Graduate School of Comprehensive Rehabilitation, Osaka Prefecture University, Habikino, Japan

¹¹Department of Human Health Sciences, Department of Palliative Medicine, Kyoto University Hospital, Kyoto, Japan

¹²WHO Centre for Health Development (WKC), Kobe, Japan

¹³Sussex Community NHS Foundation Trust, Brighton, UK

Twitter Kennedy B Nkhoma @KennedyNkhoma6, Richard Harding @RHardingCSL, Irene J Higginson @IJ_Higginson, Catherine J Evans @CatherineJanee1 and Matthew Maddocks @MTMaddocks

Acknowledgements We thank Olivia Dix for proofreading the manuscript and Hamid Benalia for helping with the production of the figures.

Contributors JB, AB, CE-S, RH, CNO, SA, PT, YM, TK, NN, ST, PO, IH, CE and MM conceived and designed the study. JB, AB, CE-S, IT, SY, DY, KBN, AC, SC, SB, CE and MM extracted data. JB, AB, CE-S, SY, AC, SC, CNI, CNO, RH, KBN, CE and MM analysed data. JB, AB, AC, SC, CNI, PO, CE and MM drafted the manuscript. All authors critically revised the draft and approved the final manuscript. MM is responsible for the overall content as the guarantor.

Funding This research was supported by the World Health Organization Centre for Health Development (WHO Kobe Centre – WKC: K19002), the Dunhill Medical Trust (grant number RPGF1906\177) and the National Institute of Health Research Applied Research Collaboration South London (NIHRARC South London NIHR200152) at King's College Hospital NHS Foundation Trust. AB is supported by the Dunhill Medical Trust (number RPGF1906\177) and Cicely Saunders International. SC is funded by a Health Education England/NIHR Clinical Doctoral Research Fellowship (ICA-CDRF-2017-03-012) and CNI by a Health Education England/NIHR Senior Clinical Lectureship (ICA-SCL-2018-04-ST2-001). IH is an NIHR Senior Investigator Emeritus. CE is funded by a Health Education England/NIHR Senior Clinical Lectureship (ICA-SCL-2015-01-001) and MM is funded by an NIHR Career Development Fellowship (CDF-2017-10-009). This publication presents independent research supported by the National Institute for Health Research (NIHR). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, NIHR or the Department of Health and Social Care.

Competing interests PO reported that he was an employee of the funding sponsor, the WHO, and was involved in the extraction, analysis and interpretation of data. All other authors have no competing interests.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Extracted data is available on request from the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Joanne Bayly <http://orcid.org/0000-0001-9478-8932>
Deokhee Yi <http://orcid.org/0000-0003-4894-1689>
Kennedy B Nkhoma <http://orcid.org/0000-0002-2991-8160>
Richard Harding <http://orcid.org/0000-0001-9653-8689>
Nanako Nishiyama <http://orcid.org/0000-0002-7580-4009>
Catherine J Evans <http://orcid.org/0000-0003-0034-7402>
Matthew Maddocks <http://orcid.org/0000-0002-0189-0952>

REFERENCES

- WHO. *World report on aging and health*. Geneva: World Health Organisation, 2015.
- Evans CJ, Ho Y, Daveson BA, *et al*. Place and cause of death in centenarians: a population-based observational study in England, 2001 to 2010. *PLoS Med* 2014;11: :e1001653.
- Nations, U. World population prospects: the 2015 revision, key findings and advance tables. Rep No. ESA/P/WP. 241 2015.
- WHO. Ageing and health, 2018. Available: <http://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
- Clegg A, Young J, Iliffe S, *et al*. Frailty in elderly people. *The Lancet* 2013;381:752–62.
- World, H., Organisation. Universal health coverage (UHC), 2020. Available: [https://www.who.int/news-room/fact-sheets/detail/universal-health-coverage-\(uhc\)](https://www.who.int/news-room/fact-sheets/detail/universal-health-coverage-(uhc))
- Ahmedzai SH, Costa A, Blengini C, *et al*. A new international framework for palliative care. *Eur J Cancer* 2004;40:2192–200.
- Walshe C, Todd C, Caress A, *et al*. Patterns of access to community palliative care services: a literature review. *J Pain Symptom Manage* 2009;37:884–912.
- Cohen LL. Racial/Ethnic disparities in hospice care: a systematic review. *J Palliat Med* 2008;11:763–8.
- Porter ME. What is value in health care? *N Engl J Med* 2010;363:2477–81.
- WPCA and who, global atlas of palliative care at the end of life. 2014, world palliative care alliance: London www.who.int/nmh_global_atlas_of_palliative_care.pdf (genethique.org)
- Evans CJ, Ison L, Ellis-Smith C, *et al*. Service delivery models to maximize quality of life for older people at the end of life: a rapid review. *Milbank Q* 2019;97:113–75.
- Moher D, Shamseer L, Clarke M, *et al*. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- Rohwer A, Booth A, Pfadenhauer L. Guidance on the use of logic models in health technology assessments of complex interventions. *International Journal of Technology Assessment in Health Care* 2016 core.ac.uk
- Popay J, Roberts H, Sowden A. Guidance on the conduct of narrative synthesis in systematic reviews: a product from the ESRC methods programme 2006 Guidance on the conduct of narrative synthesis in systematic reviews — Teesside University's Research Portal
- Brereton L, Clark J, Ingleton C, *et al*. What do we know about different models of providing palliative care? Findings from a systematic review of reviews. *Palliat Med* 2017;31:781–97.
- Checkland P, Tsouvalis C. *Reflecting on SSM: the link between root definitions and conceptual models*. *Systems Research and Behavioral Science* 1997;14:153–68.
- Hoffmann TC, Glasziou PP, Boutron I, *et al*. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:g1687.
- Fox MT, Sidani S, Persaud M, *et al*. Acute care for elders components of acute geriatric unit care: systematic descriptive review. *J Am Geriatr Soc* 2013;61:939–46.
- Briggs AM, Valentijn PP, Thiagarajan JA, *et al*. Elements of integrated care approaches for older people: a review of reviews. *BMJ Open* 2018;8:e021194.
- Naylor MD, Shaid EC, Carpenter D, *et al*. Components of comprehensive and effective transitional care. *J Am Geriatr Soc* 2017;65:1119–25.
- Bainbridge D, Seow H, Sussman J. Common components of efficacious in-home end-of-life care programs: a review of systematic reviews. *J Am Geriatr Soc* 2016;64:632–9.
- Shea BJ, Grimshaw JM, Wells GA, *et al*. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10.
- Higgins JPT, Altman DG, Gøtzsche PC, *et al*. The Cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- Milat AJ, King L, Bauman AE, *et al*. The concept of scalability: increasing the scale and potential adoption of health promotion interventions into policy and practice. *Health Promot Int* 2013;28:285–98.
- Pfadenhauer LM, Gerhardus A, Mozygemba K, *et al*. Making sense of complexity in context and implementation: the context and implementation of complex interventions (CICI) framework. *Implement Sci* 2017;12:21.
- De Coninck L, Bekkering GE, Bouckaert L, *et al*. Home- and community-based occupational therapy improves functioning in frail older people: a systematic review. *J Am Geriatr Soc* 2017;65:1863–9.
- Ekdahl AW, Sjöstrand F, Ehrenberg A, *et al*. Frailty and comprehensive geriatric assessment organized as CGA-ward or CGA-consult for older adult patients in the acute care setting: a systematic review and meta-analysis. *Eur Geriatr Med* 2015;6:523–40.
- Haun MW, Estel S, Rücker G, *et al*. Early palliative care for adults with advanced cancer. *Cochrane Database Syst Rev* 2017;6:CD011129.

- 30 Kavalieratos D, Corbelli J, Zhang D, *et al.* Association between palliative care and patient and caregiver outcomes: a systematic review and meta-analysis. *JAMA* 2016;316:2104–14.
- 31 McAlister FA, Stewart S, Ferrua S, *et al.* Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol* 2004;44:810–9.
- 32 Phillips CO, Wright SM, Kern DE, *et al.* Comprehensive discharge planning with postdischarge support for older patients with congestive heart failure: a meta-analysis. *JAMA* 2004;291:1358–67.
- 33 Fox MT, Persaud M, Maimets I, *et al.* Effectiveness of acute geriatric unit care using acute care for elders components: a systematic review and meta-analysis. *J Am Geriatr Soc* 2012;60:2237–45.
- 34 Kassianos AP, Ioannou M, Koutsantoni M, *et al.* The impact of specialized palliative care on cancer patients' health-related quality of life: a systematic review and meta-analysis. *Support Care Cancer* 2018;26:61–79.
- 35 Cui X, Dong W, Zheng H, *et al.* Collaborative care intervention for patients with chronic heart failure: a systematic review and meta-analysis. *Medicine* 2019;98:e14867.
- 36 Fulton JJ, LeBlanc TW, Cutson TM, *et al.* Integrated outpatient palliative care for patients with advanced cancer: a systematic review and meta-analysis. *Palliat Med* 2019;33:123–34.
- 37 Sezgin D OCR, Liew A, O' Donovan M. On behalf of work package 7 partners, *Intermediate care interventions for older adults*. Health Programme of the European Union, NHS Lanarkshire 2019.
- 38 Gore PG, Kingston A, Johnson GR, *et al.* New horizons in the compression of functional decline. *Age Ageing* 2018;47: :764–8.
- 39 Organization, W.H. *Integrated care for older people: guidelines on community-level interventions to manage declines in intrinsic capacity*. Geneva: Organization, W.H, 2017.
- 40 Sleeman KE, de Brito M, Etkind S, *et al.* The escalating global burden of serious health-related suffering: projections to 2060 by world regions, age groups, and health conditions. *Lancet Glob Health* 2019;7:e883–92.
- 41 Evans CJ, Bone AE, Yi D, *et al.* Community-Based short-term integrated palliative and supportive care reduces symptom distress for older people with chronic noncancer conditions compared with usual care: a randomised controlled single-blind mixed method trial. *Int J Nurs Stud* 2021;120:103978.
- 42 Kodner DL, Spreeuwenberg C. Integrated care: meaning, logic, applications, and implications--a discussion paper. *Int J Integr Care* 2002;2:e12.
- 43 Comas-Herrera A, Fernandez J-L, Hancock R, *et al.* COVID-19: implications for the support of people with social care needs in England. *J Aging Soc Policy* 2020;32:365–72.
- 44 Walshe C, Dodd S, Hill M, *et al.* How effective are volunteers at supporting people in their last year of life? A pragmatic randomised wait-list trial in palliative care (ELSA). *BMC Med* 2016;14:203.
- 45 Humphries R, Thorlby R, Holder H, *et al.* *Social care for older people: home truths*. London: King's Fund, 2016.
- 46 Robinson, L. and G.O.f.S. *Present and future configuration of health and social care services to enhance robustness in older age*. London: Foresight, Government Office for Science, 2015.
- 47 Colombo Fet al. *Help wanted? Providing and paying for long-term care OECD health policy studies*. Paris: OECD Publishing, 2011.
- 48 Tiberini R, Turner K, Talbot-Rice H. *Rehabilitation in palliative care, in textbook of palliative care*. Cham: Springer, 2018: 1–29.
- 49 Nicholson C, Davies JM, George R, *et al.* What are the main palliative care symptoms and concerns of older people with multimorbidity?—a comparative cross-sectional study using routinely collected Phase of Illness, Australia-modified Karnofsky performance status and integrated palliative care outcome scale data. *Ann Palliat Med* 2018;7:S164–75.
- 50 Kitzman DW, Whellan DJ, Duncan P, *et al.* Physical rehabilitation for older patients hospitalized for heart failure. *N Engl J Med* 2021;385:203–16.
- 51 Nottelmann L, Groenvold M, Vejlgaard TB, *et al.* Early, integrated palliative rehabilitation improves quality of life of patients with newly diagnosed advanced cancer: the Pal-Rehab randomized controlled trial. *Palliat Med* 2021;35:1344–55.
- 52 Glasziou PP, Chalmers I, Green S, *et al.* Intervention synthesis: a missing link between a systematic review and practical treatment(s). *PLoS Med* 2014;11:e1001690.