BMJ Open Early-term birth and its association with universal two-child policy: a national cross-sectional study in China

Jie Zhang,^{1,2} Gareth J Williams,³ Guanghua Wang,⁴ Jingjing Chen,^{1,2} Mengyu Zhang,^{1,2} Wenchong Du,⁵ Jing Zhu,⁶ Jun Zhang,⁶ Jing Hua^{1,2}

To cite: Zhang J, Williams GJ, Wang G, *et al.* Early-term birth and its association with universal two-child policy: a national cross-sectional study in China. *BMJ Open* 2021;**11**:e054959. doi:10.1136/bmjopen-2021-054959

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-054959>).

JZ, GJW and GW contributed equally.

Received 30 June 2021

Accepted 15 November 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Professor Jing Hua;
Jinghua@tongji.edu.cn

ABSTRACT

Objective To describe the epidemiology of early-term birth (ETB) at the national level in China, and explore the association and mediating factors between ETB and policy between universal two-child policy and ETB, so as to explain the potential reason for such a relationship and provide evidence for future ETB interventions in the era of the new birth control policy.

Design Cross-sectional study.

Participants The cross-sectional study used data from China Labour and Delivery Survey between 2015 and 2016. A total of 75 132 survey data collected from 89 hospitals in 25 provinces were included in the analysis. We further explored the association between the universal two-child policy and ETB.

Results The weighted incidence of ETB was 30.1 per 100 all births (95% CI 30.06% to 30.14%) or 29.88 per 100 live births (95% CI 29.97% to 30.05%) between 2015 and 2016 in China. There was an association between the universal two-child policy and ETB (relative risk, RR 1.19, 95% CI 1.15 to 1.23), which was not mediated by maternal age (RR 1.17, 95% CI 1.13 to 1.22), previous uterine scars (RR 1.18, 95% CI 1.14 to 1.22), parity (RR 1.19, 95% CI 1.15 to 1.24) and other measured conditions (each $p < 0.05$). Stratified analysis showed that the association between universal two-child policy and ETB were the strongest in multiparous young women or women without previous uterine scars (each $p < 0.05$), and disappeared in all women of advanced maternal age (each $p > 0.05$).

Conclusion The incidence of ETB was high in China when compared with most of reported countries, and there might be a link between two-child policy and ETB. Obstetric practice such as selective induced labour and caesarean section should be revised with ETB risks in mind, when ETB is more likely to happen under the universal two-child policy. Preventing ETB should not be neglected in multiparous young women or those without previous uterine scars under the new policy.

INTRODUCTION

Early-term birth (ETB) is defined as infants delivered at a gestation age between 37^{0/7} and 38^{6/7} weeks¹, which accounts for 15%–31% of all deliveries, and far exceeds the number of preterm births.² Preterm infants are often at a higher risk of morbidity and mortality compared with full-term infants,³ and recent

Strengths and limitations of this study

- The strength of this study is first report the rates of early-term birth (ETB) before and after the implementation of the universal two-child policy.
- The most important population for further intervention of ETB under the new policy was presented using a stratified analysis.
- The evidence provided by our study was reliable because it based a large sample in China at national level.
- Missing and excluded information appeared not to substantially affect the results in our study using a sensitivity analysis.
- We do not have a complete profile of mediating factors such as smoking, physical activity, eating habits and economic status.

reports also suggested that ETB is associated with significantly higher health risks, including higher neonatal, postnatal, and infant mortality and morbidity rates when compared with full-term infants (>38 gestational weeks)⁴; ETB can also have a long-term impact on children's cognition and behaviour.⁵ However, the epidemiology of ETB is scarce in China despite a large number of studies on preterm birth.^{6–12}

Since October 2015, China's one-child policy has been replaced with a universal two-child policy (which allows a family to have two children). However, the relationship between universal two-child policy and newborn outcomes births is still unclear. Studies have reported that the change of the birth control policy might result in nationwide changes to a range of health outcomes, such as a more equal sex ratio, reduced neonatal mortality,¹³ and increased risk of birth defects.¹⁴ However, the new two-child policy might also be associated with outcomes related to birth term.

One of the potential consequences which is associated with the two-child policy might be an increased rate of ETB, which can be influenced by a range of factors, for example,

pregnancies in mothers of advanced maternal age (≥ 34 years) have increased since the implementation of the universal two-child policy,^{14–17} and there is an increased risk of pregnancy complications in women of advanced maternal age.^{14 18} Complications during pregnancy, such as placental ischaemia and diabetes mellitus, increase the risk of ETB.¹⁹ Additionally, a higher primary caesarean section (C-Section) rate in China means that a more pregnant women who have had a previous birth have uterine scars.^{20 21} To avoid rupturing the uterus, early elective birth is more likely to be a decision for women with previous scars^{3 22 23}; the uterus scar site is more susceptible to tearing and rupture when subjected to pressure during labour. Therefore, with those possible influencing factors, a higher incidence of ETB might be observed after the implementation of the universal two-child policy.

Furthermore, studies showed that the relationship of universal two-child policy with obstetric issues can vary across different maternal characteristics.^{13 16 24} For example, it has been reported that the associations between the two-child policy and sex ratios differed by maternal age, parity, and history of C-section.¹³ It has also been found that the association of two-child policy with caesarean birth rate varies between nulliparous and multiparous women.²⁴ Moreover, the risk of childbearing policy on incidence of birth defects has been found to be higher in women aged 30–34 years old when compared with their younger counterparts.¹⁴

In the current study, we aimed to describe the epidemiology of ETB in China by using a nationwide dataset from the China Labour and Delivery Survey (CLDS). Additionally, we analysed the association between the incidence of ETB and the two-child policy in terms of different maternal characteristics (such as maternal age, previous uterine scars and parity). We also explored the factors that might mediate the association between ETB and policy so as to explain such a relationship and offer evidence for future ETB interventions in the era of the new birth control policy.

METHODS

Sample

Data for this national cross-sectional study were acquired from the dataset of the CLDS, which is a population-based multicentre study that describes the incidence, risk factors and possible underlying causes of ETB in China. We extracted the data between 2015 and 2016 because the one-child policy has been replaced with a two-child policy since October 2015. Participation was solicited through a national obstetric conference hosted by Shanghai First Maternity and Infant Hospital, Tongji University School of Medicine.

We used the same methodology as the WHO Global Survey of Maternal and Perinatal Health and the WHO Multi-Country Survey of Maternal and Newborn Health to perform data collection.^{25 26} The hospitals that expressed an interest in taking part in the research were

asked to provide basic information about the hospitals, and only those with 1000 or more deliveries per year (having sufficient obstetrical faculty) were eligible for being included in this study. To ensure that there was a representative sample of ETB, we selected the hospitals with 1000 or more deliveries per year which covered most Chinese provinces (25 out of 34). With reference with the WHO methodology,^{25 26} for hospitals with an annual rate of more than 6000 deliveries per year, we divided the 1-year period into 13 time-sections and randomly selected 6 weeks of records, whereas for hospitals with an annual delivery rate of less than 6000 per year, ten weeks were randomly selected within a 1-year period, divided into five time sections. Within each selected week, all births delivered at 24 or more completed weeks of gestation or weighing 500 grams or more at birth were eligible. Medical records were retrieved and reviewed, and information was extracted by trained research nurses. The study design, organisation and implementation have also been reported in another study.²⁷ Using the WHO methodology allows for comparability between these results and the studies of other countries.

A data coordination centre was established to take charge of establishing, managing and maintaining the database and website, coordination among hospitals and investigators' training. This study did not involve individual informed consent because the centre collected anonymous clinical information only. All information was kept confidential, and only the cooperators of our research could access the database when their application was approved by the data coordination centre. Additionally, the centre was also responsible for data logic checking, assessing of study protocol and other quality control measures of the data.

A total of 75 132 birth records were initially included in the analysis from 89 hospitals in China between 1 March 2015 and 31 December 2016. Among these birth records, a total of 73 567 live birth were included for calculating the incidence of ETB at a national level. In order to further analyse the association of ETB with the implementation of the universal two-child policy, we chose the women who were pregnant after October 2015 (the universal two-child policy was announced) and gave birth to children after July 2016 (the effective period: 9 months after the announcement of the universal two-child policy) according to previous studies.¹⁵ The women who gave birth to a child before October 2015 (before the announcement of the universal two-child policy) were included as controls. A total of 28 966 live singleton full-term birth were included in the final analysis (figure 1).

Patient and public involvement

Patients or the public were not involved in the design, conduct, reporting or dissemination plans of our research.

Exposure, outcomes and other covariates

ETB, which was defined as newborn delivery occurring between 37^{0/7} to 38^{6/7} weeks of gestation, was the main

Figure 1 Flow chart of study population. Flow chart of the study inclusion and exclusion criteria for review of total participant samples and how the final number of the cohort was established. ETB, early-term birth.

outcome of our study. Gestational age was determined by obstetrical providers with the use of standard criteria that took into consideration the clinical history and the results of the earliest ultrasound examination.^{28 29} If the gestational age as calculated from an early ultrasound is contradictory to the one calculated directly from the last menstrual period, it is the age calculated from the early ultrasound that is used for the rest of the pregnancy.^{30 31}

We included the maternal age, previous uterine scar, parity and precursors as mediating factors (which refer to the factors that could mediate the relationship between the independent and dependent variables so as to explain the reason for such a relationship to exist). Precursors for delivery were defined using the following four categories: spontaneous labour, premature rupture of membranes (PROM), indicated and elective induction or C-Section, which has also been used to assess the reasons for late preterm birth according to previous study.³² Spontaneous labour was defined as women with the spontaneous start of labour and with or without complications or rupture of membranes during pregnancy. PROM was defined where a woman presented with PROM and underwent induction labour or prelabour C-Section, but exclusive of spontaneous labour. Indicated deliveries included women without spontaneous labour or PROM who underwent induction labour or prelabour C-Section for maternal, fetal or obstetric complications.^{33 34} We also categorised as indicated any delivery where induction of labour or

indication for prelabour C-Section was not recorded, but the pregnancy had other complications such as hyperthyroidism or hypothyroidism, and hypertensive disease or gestational diabetes. Elective induction or C-Section included elective inductions or C-Section as identified by the site with no other obstetrical, fetal or maternal conditions, and there was no reason for induction or C-Section provided, or no other obstetrical, fetal or maternal conditions.^{35 36}

Additionally, it has been reported that the pre-pregnancy body mass index (BMI) is related with gestational ages.^{37 38} We included it as a potential confounder when considering the association between two-child policy and ETB. We used maternal prepregnancy weight and height to calculate BMI, which was grouped into three categories: underweight ($<18.5 \text{ kg/m}^2$), normal weight ($18.5\text{--}24.9 \text{ kg/m}^2$) and overweight or obesity ($\geq 25 \text{ kg/m}^2$) according to the WHO BMI classification.³⁹ Maternal height was presented in quartiles (Q1: 100–160 cm, Q2: 161–165 cm, Q3: 166–168 cm, Q4: 169–190 cm).

Statistical analysis

To better represent the epidemiology of ETB in China, we calculated weight for each birth in the survey by using the number of deliveries in each province from the 2016 China Statistical Yearbook, compiled by the National Bureau of Statistics of China (<http://www.stats.gov.cn/tjsj/ndsj/2016/indexch.htm>). We calculated a 95% CI of the incidence of Early term with the Normal approximation. We analysed the difference between the precursors of delivery in ETB before and after the implementation of the universal two-child policy using the partitions of the χ^2 method.

We presented the maternal and newborn's characteristics in the study population by the status of birth control policy and by the different gestational ages (online supplemental file 1). These characteristics were considered as potential confounders (covariates) when we analysed the association between two-child policy and ETB. A multivariable regression model was used to analyse the associations between universal two-child policy and ETB when adjusting for or not adjusting for the maternal and newborn characteristics (as confounders).

We carried out a stratified analysis according to maternal age, parity, and previous uterine scar to explore potential disparities in the association of ETB with a universal two-child policy. To further clarify the association, we carried out analyses to detect the joint effect of each age group by subgroup of parity or previous uterine scar. We added the interaction term (maternal age $\times$ parity or maternal age $\times$ uterine scars) to the model. When an interaction is confirmed (with statistical significance), strata-specific estimates are reported.

Additionally, we did a series of sensitivity analyses by including all subjects and excluding women who had gestational week <37 , with Gemellary pregnancy and stillbirth or newborn death, late-term and post-term birth,

missing of covariates on the association between two-child policy and ETB.

Relative risk (RR) and 95% CI was estimated using a multilevel modified Poisson regression approach.⁴⁰ For controlling the unmeasured confounding factors from different hospitals, we used a multilevel model (considering the hospital as the cluster) in our study. The application of Poisson regression on multinomial data has been shown to overestimate the RR.^{40 41} We rectified this using a robust error variance procedure known as sandwich estimation, in our modified Poisson regression⁴⁰ using PROC GENMOD in SAS V.9.4 software. A *p* value of <0.05 was considered statistically significant.

RESULTS

Among all births (*n*=75 132) and live births (*n*=73 567), the unweighted incidence of ETB was 31.88% per 100 all births (95% CI 31.55% to 32.21%) and 31.72% per 100 live births (95% CI 31.38% to 32.06%), respectively. After weighting adjustment, the weighted incidence of ETB was 30.1% per 100 all births (95% CI 30.06% to 30.14%) and 29.88% per 100 live births (95% CI 29.97% to 30.05%) in China, respectively. Of 75 132 birth, 41.75% of all weighted births occurred with spontaneous labour, followed by indicated (41.32%), PROM (13.83%) and elective induction or C-Section (3.10%). Of 73 567 live births, 41.75% occurred with spontaneous labour, followed by indicated (41.32%), PROM (14.09%) and elective induction or C-Section (2.45%). There were substantial variations of ETB according to maternal age, previous uterine scars and parity (table 1).

Of 28 966 women, 7566 women were included in the exposure group (after the implementation of two-child policy) and 21 400 women (before the implementation of two-child policy) were included in the control group. There was a statistically significant difference in most of the maternal and newborn characteristics between the exposure and control groups, and between ETB and completely full-term birth (*p*<0.05 online supplemental eTable 1), so these were included as mediating or confounding factor in our study. In table 2, the results showed there were associations between universal two-child policy and ETB when not adjusting for (RR 1.19, 95% CI 1.15 to 1.23, *p*<0.001) or adjusting for maternal age (RR 1.17, 95% CI 1.13 to 1.22, *p*<0.001), previous uterine scars (RR 1.18, 95% CI 1.14 to 1.22, *p*<0.001), parity (RR 1.19, 95% CI 1.15 to 1.24, *p*<0.001) and all above three conditions (RR 1.17, 95% CI 1.13 to 1.21, *p*<0.001). The statistical significance remained when precursors for delivery (RR 1.18, 95% CI 1.14 to 1.23, *p*<0.001) and all measured characteristics (RR 1.19, 95% CI 1.14 to 1.22, *p*<0.001) were added to the adjustment.

In the stratified analysis, we found that the association of universal two-child policy with ETB varied by different ages, previous uterine scars and parity (table 3). The association between universal two-child policy and ETB differed significantly by maternal age and uterine scars (*p*_{interaction}=0.0026). We analysed the association when stratified by age × previous uterine scars (figure 2). The association of universal two-child policy with ETB was statistically significant in women younger than 30 years of age with previous uterine scars (adjusted RR=1.19,

Table 1 Incidence of early-term birth by maternal age, previous uterine scar, parity and precursors for delivery (*n*=73 567)

Characteristics	Total	Unweighted		Weighted	
		Incidence (%)	95% CI	Incidence (%)	95% CI
Maternal age					
<30	45 982	25.88	25.56 to 26.20	24.81	24.50 to 25.12
30–34	19 141	31.87	31.53 to 32.21	30.67	30.34 to 31.00
≥35	8 444	35.14	34.80 to 35.48	34.38	34.04 to 34.72
Previous uterine scars					
Yes	10 681	42.84	42.48 to 43.20	40.25	39.90 to 40.60
No	62 886	26.07	25.75 to 26.39	24.81	24.50 to 25.12
Parity					
Nulliparous	41 302	24.96	24.65 to 25.27	23.27	22.96 to 23.58
Multiparous	32 265	33.04	32.70 to 33.38	30.84	30.51 to 31.17
Precursors for delivery					
Spontaneous labour	33 958	24.82	24.51 to 25.13	23.94	23.63 to 24.25
PROM	8 616	33.57	33.23 to 33.91	34.58	34.24 to 34.92
Indicated*	28 815	31.67	31.33 to 32.01	29.79	29.46 to 30.12
Elective induction or C-Section	2 178	23.92	23.61 to 24.23	21.53	21.23 to 21.83

*‘Indicated’ precursors refer to women with maternal, fetal or obstetric complications.

C-Section, caesarean section; PROM, premature rupture of membranes.

Table 2 Association between universal two-child policy and early-term birth when adjusting for maternal and newborn's characteristics (n=28 966)

Adjusting variables	Adjusted RR (95% CI)	P value
Crude*	1.19 (1.15 to 1.23)	<0.0001
Maternal age	1.17 (1.13 to 1.22)	<0.0001
Previous uterine scars	1.18 (1.14 to 1.22)	<0.0001
Parity	1.19 (1.15 to 1.24)	<0.0001
Maternal age +previous uterine scars+parity	1.17 (1.13 to 1.21)	<0.0001
Maternal age +previous uterine scars+parity+precursors for delivery	1.18 (1.14 to 1.23)	<0.0001
Maternal age +previous uterine scars+parity+precursors for delivery +the other maternal and newborn characteristics†	1.18 (1.14 to 1.22)	<0.0001

*Not adjusted for any variables.

†The other maternal and newborn characteristics were shown in online supplemental eTable 1.
RR, relative risk.

$p<0.05$) and without previous uterine scars (adjusted RR=1.25, $p<0.05$). The association remained among those aged from 30 to 34 without previous uterine scars (adjusted RR=1.19, $p<0.05$), however, disappeared in those with previous uterine scars ($p>0.05$). The association disappeared for all women aged more than 34 years with and without previous uterine scars (each $p>0.05$). The crude RR and 95% CI of RR were shown in figure 2.

We found the associations between universal two-child policy and ETB differed significantly by maternal age and

parity ($p_{\text{interaction}}=0.0155$). Therefore, we ran further analyses for strata-specific estimates by age $\times$ parity (figure 1). The universal two-child policy was positively associated with ETB among nulliparous and multiparous women younger than 30 years old (adjusted RR were 1.23 and 1.31, respectively, $p<0.05$), nulliparous and multiparous women aged 30–40 years (adjusted RR were 1.14 and 1.16, respectively, each $p<0.05$), but the association disappeared among nulliparous and multiparous women older than 34 years when adjusting for maternal and newborn characteristics (each $p>0.05$). The crude RR and 95% CI of RR were shown in figure 3.

Sensitivity analyses, in which we did not exclude any variables or only excluded preterm births, gemellary pregnancy, stillbirth and neonatal death, and missing covariates yielded similar results when compared with our main analysis (each $p>0.05$). The finding changed with marginal significance, in contrast to our main analysis, when we excluded late-term and post-term birth from the overall subjects ($p=0.05$, online supplemental eFigure 1).

DISCUSSION

To our knowledge, the current study is the first to describe the incidence of ETB in China at a national level. We observed an incidence of about 30% of ETB between 2015 and 2016 in China and an increased risk of ETB after the announcement of the universal two-child policy. However, the association between ETB and the universal two-child policy was not mediated by maternal age, previous uterine scars, parity and other measured characteristics. Furthermore, the stratified analysis showed that the effects of the universal two-child policy on ETB was strongest in multiparous young women or those without previous uterine scars, and disappeared in all women

Table 3 Association between universal two-child policy and early-term birth stratified by maternal age, previous uterine scars and parity(n=28 966)

Adjusted variables	Crude RR* (95% CI)	P value	Adjusted RR† (95% CI)	P value
Maternal age				
<30	1.22 (1.16 to 1.28)	<0.0001	1.24 (1.18 to 1.30)	<0.0001
30–34	1.15 (1.08 to 1.22)	<0.0001	1.14 (1.07 to 1.22)	<0.0001
≥35	1.06 (0.97 to 1.15)	0.1816	1.05 (0.97 to 1.15)	0.2241
Previous uterine scars				
No	1.20 (1.16 to 1.25)	<0.0001	1.21 (1.16 to 1.26)	<0.0001
Yes	1.08 (1.02 to 1.15)	0.0104	1.08 (1.01 to 1.15)	0.0263
Parity				
Nulliparous	1.18 (1.12 to 1.24)	<0.0001	1.19 (1.13 to 1.26)	<0.0001
Multiparous	1.20 (1.14 to 1.25)	<0.0001	1.17 (1.12 to 1.23)	<0.0001

*Not adjusted for any variables.

†Adjusted for maternal sociodemographic status, maternal health conditions and newborn's characteristic.

‡Adjusted for maternal sociodemographic status.

§Adjusted for maternal health conditions.

¶Adjusted for newborn's characteristics.

RR, relative risk.

Figure 2 Association between universal two-child policy and early-term birth (ETB) in women without previous uterine scars when not adjusting other variables (A), without previous uterine scars when adjusting other variables (B), with previous uterine scars when not adjusting other variables (C), with previous uterine scars when adjusting other variables (D), by maternal age group (n=28 966). The association between universal two-child policy and ETB when stratified by age × previous uterine scars. RR, relative risk. * $p < 0.05$, ** $p < 0.01$

of advanced maternal age (irrespective of parity and previous uterine scars).

The incidence of ETB in different countries is varied. For example, a population study reported that the average incidence of ETB was 25.3% among six high-income countries.⁴² In that study, a lower incidence was reported in Denmark, Finland, Norway, and Sweden (all below 20%), and a higher incidence was reported in the USA and Canada (higher than 20%). In other countries, the reported incidence of ETB is fairly consistent. For example, a multicentred study from 44 hospitals in Portugal reported an incidence of ETB of 27%.⁴³ Another national study in Brazil reported an incidence of ETB of 29.8% in singleton deliveries.⁴⁴ The incidence of ETB in China of about 30% between 2015 and 2016 can be therefore considered as similar to Portugal and Brazil, but higher than most of reported countries. The variations of ETB across different maternal age, parity and women with or without previous uterine scars in China were consistent with previous studies.^{19 45–49}

Of the reported ETB cases in the current study, about 3% were due to having an elective delivery. It has been reported that the ETB rates in the USA decreased from 33.0% in 2006 to 21.1% in 2014 among births with clinician-initiated obstetric intervention, and from 29.7% in 2006 to 27.1% in 2014 among births without clinician-initiated obstetric intervention.⁴² These studies^{42 50 51} found that reductions in elective obstetric intervention

at ETB may reflect the success of interdisciplinary health-care teams that focus on decreasing elective births before 39 weeks.^{52–54} Therefore, our result suggests that the ETB rate in China may decrease if obstetric intervention reduces the number of unnecessary elective deliveries.

Owing to the timing of the initiation of the universal two-child policy across the research period, we explored ETB rate changes after the implementation of the universal two-child policy, so as to provide evidence for future ETB interventions in the era of the new policy. Our study reported that the incidence of ETB increased after the implementation of the universal two-child policy, however, the association stayed statistically significant after mediated by maternal age, previous uterine scar, parity and the other maternal and newborn factors in table 1. The results suggest that the two-child policy might bring other risk factors that were not included in our study. A recent study suggested that under the newly relaxed universal two-child policy, women may experience higher risk of mental health difficulties during their second pregnancy,⁵⁵ and poorer mental health may lead to a higher chance of ETB. Further research is necessary to explore the association between ETB and maternal and newborn outcomes, to reveal more details of the underlying mechanisms of ETB, and inform new interventions for pregnant women.

We found that there was an association between China's two-child policy and ETB which differed across

Figure 3 Association between universal two-child policy and early-term birth (ETB) in nulliparous women when not adjusting other variables (A), in nulliparous women when adjusting other variables (B), in multiparous women when not adjusting other variables (C), in multiparous women when adjusting other variables (D), by maternal age group (n=28 966). The diagram analysed the association between universal two-child policy and ETB when stratified by age × parity. RR, relative risk. * $p < 0.05$, ** $p < 0.01$

subjects with different maternal characteristics, especially, maternal age. One likely explanation for why the association was not present in women of advanced maternal age (but remained in women with younger age), is a ‘ceiling effect’. Women of advanced maternal age are at risk of higher rates of ETB as they are more vulnerable to maternal complications and adverse birth outcomes. Obstetric complications such as placental abruption,⁵⁶ malpresentation and postpartum haemorrhage^{45 46 57 58} are higher in women of advanced maternal age. Unexplained stillbirths increase with advancing maternal age and with increasing gestational age in both nulliparous and multiparous women. Therefore, the induction of labour or prelabour C-Section in women of advanced maternal age is widely practiced as an intervention perceived to reduce the risk of adverse perinatal outcomes.⁴⁶ Those factors may reduce the difference in ETB rates in women of advanced maternal age before and after the implementation of the universal two-child policy.

We also found that the effect of the universal two-child policy on ETB was the strongest in young women without previous uterine scars, or who are multiparous. It has been reported that early elective birth before 39 weeks is more likely to occur in women with previous scars when obstetricians do not appreciate the risk of morbidity of infants before 39 weeks and prefer to maintain autonomy in determining the timing of delivery.³ Therefore, the probability of early elective birth in women with previous

uterine scars may moderate the difference in ETB before and after the implementation of the universal two-child policy. Additionally, the effects of the policy on ETB increased more significantly in multiparous young women than their nulliparous counterparts. One possible explanation is that first-time mothers are more likely to have a spontaneous vaginal delivery, instead of an elective birth week, so that they are able to reduce the risk of potential health complications that might arise from a second pregnancy,⁵⁹ thus reducing the difference in ETB rate in nulliparous women before and after the implementation of the universal two-child policy.

Strengths and limitations

With the large sample in our study, we first reported the incidence of ETB at a national level and found an association between universal two-child policy and ETB. However, there were several limitations to our study. First, we do not have a complete profile of factors that may influence the association between universal two-child policy and ETB. For example, we have comparatively little information on maternal behaviour, such as smoking, physical activity and eating habits, and were unable to measure the influence of psychological disorders (eg, depression, anxiety) on ETB in this study. Furthermore, future studies including sociological variables such as maternal economy, marital status, income, education and occupation could help further explain these findings.

We will explore these potential factors in our subsequent study. Second, we excluded the stillbirth (or newborn death), preterm birth, gemellary pregnancy, late-term and post-term birth, missing covariates in our study, and thus our findings may not be generalised to these populations. However, in the sensitivity analysis of our study, it did not appear to substantially affect the results.

CONCLUSION

The incidence of ETB was high in China when compared with most of reported countries and practice patterns should be adjusted to respond to ETB, when ETB was found to be more likely to happen under the universal two-child policy. The underlying mechanisms have yet to be elucidated; the mediating effects of maternal age, parity and previous uterine scars on the association between universal two-child policy and ETB need to be explored further, in a more diverse dataset, with a more comprehensive range of information for potential confounding factors.

Author affiliations

¹Department of Women and Children's Health Care, School of Medicine, Tongji University, Shanghai, China

²Shanghai Key Laboratory of Maternal Fetal Medicine, Shanghai First Maternity and Infant Hospital, School of Medicine, Tongji University, Shanghai, China

³School of Social Sciences, Nottingham Trent University, Nottingham, UK

⁴Department of Obstetrics and Gynecology, Tongren Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

⁵Department of Psychology, Nottingham Trent University, Nottingham, UK

⁶Ministry of Education-Shanghai Key Laboratory of Children's Environmental Health, Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Contributors JH conceptualized and designed the study, reviewed and critically revised the manuscript. JZ, GW and WD drafted the initial manuscript. JH, JZ, JC and WD conducted the data analysis. JC, JZ, JZ and MZ established the cohort and provided intellectual inputs to the manuscript. GJW were responsible for and supervising data collection and critical revision of the manuscript for important intellectual content. All authors reviewed and approved the final version of the manuscript.

Funding This study was partly supported by National Natural Science Foundation of China (81673179), the Science and Technology Commission of Shanghai Municipality (18140903100, 19140903100), Shanghai Municipal Health Commission (2020YJZX0213), Clinical Research Plan of Shanghai Hospital Development Center (SHDC2020CR1047B-003).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the Ethics Committees of Xinhua Hospital (XHEC-C2015-006) and WHO (A65899). Since only anonymous clinical information were collected, no individual informed consent was obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Jing Hua <http://orcid.org/0000-0001-7613-0926>

REFERENCES

1. Spong CY. Defining "term" pregnancy: recommendations from the defining "term" pregnancy workgroup. *JAMA* 2013;309:2445–6.
2. Mally PV, Agathis NT, Bailey SM. Early term infants are at increased risk of requiring neonatal intensive care. *World J Pediatr* 2016;12:76–81.
3. Craighead DV. Early term birth: understanding the health risks to infants. *Nurs Womens Health* 2012;16:136–45.
4. Reddy UM, Bettgowda VR, Dias T, et al. Term pregnancy: a period of heterogeneous risk for infant mortality. *Obstet Gynecol* 2011;117:1279–87.
5. Nielsen TM, Pedersen MV, Miliidou I. Long-Term cognition and behavior in children born at early term gestation: a systematic review. *Acta Obstet Gynecol Scand* 2019;98:1227–34.
6. Chen C, Zhang JW, Xia HW, et al. Preterm birth in China between 2015 and 2016. *Am J Public Health* 2019;109:1597–604.
7. Zou L, Wang X, Ruan Y, et al. Preterm birth and neonatal mortality in China in 2011. *Int J Gynaecol Obstet* 2014;127:243–7.
8. Xue Q, Shen F, Gao Y, et al. An analysis of the medical indications for preterm birth in an obstetrics and gynaecology teaching hospital in Shanghai, China. *Midwifery* 2016;35:17–21.
9. Zhang X, Zhou M, Chen L, et al. Risk factors for preterm birth: a case-control study in rural area of Western China. *Int J Clin Exp Med* 2015;8:4527–32.
10. Lu L, Qu Y, Tang J, et al. Risk factors associated with late preterm births in the underdeveloped region of China: a cohort study and systematic review. *Taiwan J Obstet Gynecol* 2015;54:647–53.
11. Liu L, Liu J-meng, Liu Y-hui, et al. [Prevalence of preterm birth among singletons in 10 counties (cities) of China, 1993–2005]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2007;28:1051–4.
12. Newnham JP, Sahota DS, Zhang CY, et al. Preterm birth rates in Chinese women in China, Hong Kong and Australia - the price of Westernisation. *Aust N Z J Obstet Gynaecol* 2011;51:426–31.
13. Fan SL, Xiao CN, Zhang YK, et al. How does the two-child policy affect the sex ratio at birth in China? A cross-sectional study. *BMC Public Health* 2020;20:789.
14. Zhang X, Chen L, Wang X, et al. Changes in maternal age and prevalence of congenital anomalies during the enactment of China's universal two-child policy (2013–2017) in Zhejiang Province, China: an observational study. *PLoS Med* 2020;17:e1003047.
15. Li H-T, Xue M, Hellerstein S, et al. Association of China's universal two child policy with changes in births and birth related health factors: national, descriptive comparative study. *BMJ* 2019;366:14680.
16. Teng X, Shane MI, Pan S. The changing situation about maternal age, risk factors and pregnancy outcomes after the two-child policy: a retrospective cohort study. *Ann Palliat Med* 2020;9:824–34.
17. Liu J, Song L, Qiu J, et al. Reducing maternal mortality in China in the era of the two-child policy. *BMJ Glob Health* 2020;5:e002157.
18. Jolly M, Sebire N, Harris J, et al. The risks associated with pregnancy in women aged 35 years or older. *Hum Reprod* 2000;15:2433–7.
19. Brown HK, Speechley KN, Macnab J, et al. Biological determinants of spontaneous late preterm and early term birth: a retrospective cohort study. *BJOG* 2015;122:491–9.
20. Feng XL, Wang Y, An L, et al. Cesarean section in the People's Republic of China: current perspectives. *Int J Womens Health* 2014;6:59–74.
21. Liu X, Landon MB, Cheng W, et al. Cesarean delivery on maternal Request in China: what are the risks and benefits? *Am J Obstet Gynecol* 2015;212:817 e1–9.
22. Kayem G, Raiffort C, Legardeur H. Specific particularities of uterine scars and their impact on the risk of uterine rupture in case of trial of labor. *J Gynecol Obstet Biol Reprod* 2012;41:753–71.
23. Kiran TSU, Chui YK, Bethel J, et al. Is gestational age an independent variable affecting uterine scar rupture rates? *Eur J Obstet Gynecol Reprod Biol* 2006;126:68–71.
24. Yan J, Wang L, Yang Y. The trend of caesarean birth rate changes in China after 'universal two-child policy' era: a population-based study in 2013–2018. *BMC Med* 2020;18:249.

- 25 Shah A, Faundes A, Machoki M'Imunya, *et al.* Methodological considerations in implementing the who global survey for monitoring maternal and perinatal health. *Bull World Health Organ* 2008;86:126–31.
- 26 Souza JP, Gülmezoglu AM, Vogel J, *et al.* Moving beyond essential interventions for reduction of maternal mortality (the who multicountry survey on maternal and newborn health): a cross-sectional study. *Lancet* 2013;381:1747–55.
- 27 Chen C, Zhang JW, Xia HW, *et al.* Preterm birth in China between 2015 and 2016. *Am J Public Health* 2019;109:1597–604.
- 28 Cooper S, Somerset D. Determination of gestational age by ultrasound. *J Obstet Gynaecol Can* 2016;38:337.
- 29 Bisahnyui P, Nkfusai CN, Bede F, *et al.* Comparative study of clinical methods versus ultrasound methods for accurate gestational age determination in different trimesters of pregnancy, Ndop district Hospital, North West region, Cameroon. *Pan Afr Med J* 2020;37:4.
- 30 Andersen HF, Johnson TR, Barclay ML, *et al.* Gestational age assessment. I. analysis of individual clinical observations. *Am J Obstet Gynecol* 1981;139:173–7.
- 31 Hittner HM, Gorman WA, Rudolph AJ. Examination of the anterior vascular capsule of the lens: II. assessment of gestational age in infants small for gestational age. *J Pediatr Ophthalmol Strabismus* 1981;18:52–4.
- 32 Laughon SK, Reddy UM, Sun L, *et al.* Precursors for late preterm birth in singleton gestations. *Obstet Gynecol* 2010;116:1047–55.
- 33 Krispin E. Management of premature rupture of membranes at term: the need to correct a recurring mistake in articles, chapters, and recommendations of professional organizations. *Am J Obstet Gynecol* 2017;217:661 e1–61.
- 34 Jiang H, Lu C, Zhou J, *et al.* Cesarean section and pregnancy outcomes of preterm premature rupture of membranes under different fertility policies in China. *Transl Pediatr* 2021;10:973–83.
- 35 Coulm B, Blondel B, Alexander S, *et al.* Elective induction of labour and maternal Request: a national population-based study. *BJOG* 2016;123:2191–7.
- 36 Entringer AP, Pinto M, Gomes MAdSM. Cost-Effectiveness analysis of natural birth and elective C-section in supplemental health. *Rev Saude Publica* 2018;52:91.
- 37 Sharifzadeh F, Kashanian M, Jouhari S, *et al.* Relationship between pre-pregnancy maternal BMI with spontaneous preterm delivery and birth weight. *J Obstet Gynaecol* 2015;35:354–7.
- 38 Torloni MR, Betrán AP, Daher S, *et al.* Maternal BMI and preterm birth: a systematic review of the literature with meta-analysis. *J Matern Fetal Neonatal Med* 2009;22:957–70.
- 39 WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157–63.
- 40 Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
- 41 Cao Q, Wu Z, Sun Y, *et al.* [Application of negative binomial regression and modified Poisson regression in the research of risk factors for injury frequency]. *Wei Sheng Yan Jiu* 2011;40:702–4.
- 42 Richards JL, Kramer MS, Deb-Rinker P, *et al.* Temporal trends in late preterm and early term birth rates in 6 high-income countries in North America and Europe and association with clinician-initiated obstetric interventions. *JAMA* 2016;316:410–9.
- 43 Barros JG, Clode N, Graça LM. Prevalence of late preterm and early term birth in Portugal. *Acta Med Port* 2016;29:249–53.
- 44 Barros FC, Rabello Neto Ddel, Villar J, *et al.* Caesarean sections and the prevalence of preterm and early-term births in Brazil: secondary analyses of national birth registration. *BMJ Open* 2018;8:e021538.
- 45 Carolan M, Davey M-A, Biro MA, *et al.* Older maternal age and intervention in labor: a population-based study comparing older and younger first-time mothers in Victoria, Australia. *Birth* 2011;38:24–9.
- 46 Carolan M, Frankowska D. Advanced maternal age and adverse perinatal outcome: a review of the evidence. *Midwifery* 2011;27:793–801.
- 47 Murta EFC, Nomellini RS. Is repeated caesarean section a consequence of elective caesarean section? *Lancet* 2004;364:649–50.
- 48 Mohammed A-BF, Bayo AI, Abu-Jubara MF. Timing of elective repeated cesarean delivery in patients with previous two or more cesarean section. *J Matern Fetal Neonatal Med* 2013;26:10–12.
- 49 Hutcheon JA, Joseph KS, Kinniburgh B, *et al.* Maternal, care provider, and institutional-level risk factors for early term elective repeat cesarean delivery: a population-based cohort study. *Matern Child Health J* 2014;18:22–8.
- 50 Little SE, Robinson JN, Puopolo KM, *et al.* The effect of obstetric practice change to reduce early term delivery on perinatal outcome. *J Perinatol* 2014;34:176–80.
- 51 Osterman MJ, Martin JA. Recent declines in induction of labor by gestational age. *NCHS Data Brief* 2014;155:1–8.
- 52 Trembath AN, Iams JD, Walsh M. Quality initiatives related to moderately preterm, late preterm, and early term births. *Clin Perinatol* 2013;40:777–89.
- 53 Oshiro BT, Kowalewski L, Sappenfield W, *et al.* A multistate quality improvement program to decrease elective deliveries before 39 weeks of gestation. *Obstet Gynecol* 2013;121:1025–31.
- 54 Berrien K, Deventer J, French A, *et al.* The perinatal quality collaborative of North Carolina's 39 weeks project: a quality improvement program to decrease elective deliveries before 39 weeks of gestation. *N C Med J* 2014;75:169–76.
- 55 Lu L, Duan Z, Wang Y, *et al.* Mental health outcomes among Chinese prenatal and postpartum women after the implementation of universal two-child policy. *J Affect Disord* 2020;264:187–92.
- 56 Fretts RC, Schmittiel J, McLean FH, *et al.* Increased maternal age and the risk of fetal death. *N Engl J Med* 1995;333:953–7.
- 57 Hung TH. Advanced maternal age and adverse perinatal outcome: a call for investigations on Asian women. *Taiwan J Obstet Gynecol* 2008;47:257–8.
- 58 Jacobsson B, Ladfors L, Milsom I. Advanced maternal age and adverse perinatal outcome. *Obstetrics & Gynecology* 2004;104:727–33.
- 59 Zhao J, Shan N, Yang X, *et al.* Effect of second child intent on delivery mode after Chinese two child policy implementation: a cross sectional and prospective observational study of nulliparous women in Chongqing. *BMJ Open* 2017;7:e018823.